Experimental evolution of virulence and associated traits in a *Drosophila melanogaster* –

***Wolbachia* symbiosis**

David Monnin*, Natacha Kremer, Caroline Michaud, Manon Villa, Hélène Henri, Emmanuel

Desouhant, Fabrice Vavre

Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive

UMR 5558, F-69622 Villeurbanne, France

* Corresponding author: david.monnin@hotmail.fr (School of Biological and Chemical Sciences,

Queen Mary University of London, Mile End Road, London E1 4NS, UK)

Abstract

Evolutionary theory predicts that vertically transmitted symbionts are selected for low

virulence, as their fitness is directly correlated to that of their host. In contrast with this prediction,

the *Wolbachia* strain *w*MelPop drastically reduces its *Drosophila melanogaster* host lifespan at

high rearing temperatures. It is generally assumed that this feature is maintained because the

*D. melanogaster–w*MelPop symbiosis is usually not exposed to environmental conditions in which

the symbiont is virulent. To test this hypothesis, we submitted *w*MelPop-infected *D. melanogaster*

lines to 17 generations of experimental evolution at a high temperature, while enforcing late

reproduction. The fly survival was measured at different time points, as well as two traits that have

been proposed to be causally responsible for *w*MelPop virulence: its relative density and the mean

number of octomom copies present in its genome. We hypothesised that these conditions would

select for a reduced *w*MelPop virulence, a reduced *w*MelPop density, and a reduced octomom copy

number. Our results indicate that density, octomom copy number and virulence are correlated.

However, contrary to our expectations, we could not detect any reduction in virulence during the

course of evolution. We discuss the significance of our results with respect to the evolutionary

causes of *w*MelPop virulence and propose that intra-host selection could explain this conundrum.

Introduction

Symbionts live inside (endosymbionts) or on (ectosymbionts) bigger organisms referred to

as hosts. As a result, their evolutionary success crucially depends on their ability to colonize, or be

transmitted to, new hosts. Deleterious symbionts, also known as parasites, exploit their hosts to

maximize their transmission, but in doing so usually face a trade-off. Indeed, the reproductive rate

of a parasite is likely to be positively correlated to both its instantaneous transmission rate and its

virulence (the reduction in fitness it incurs to its host). Increased virulence is in turn associated,

notably through increased host mortality, with a reduction in the opportunities for transmission

(Anderson & May 1982, Ewald 1983, see Alizon et al. 2009 for a review and defence of this

hypothesis).

Theoretical and experimental works suggest that the mode of transmission – which ranges

from fully vertical (from parents to offspring) to fully horizontal (contagion) – is an important

factor shaping parasite virulence. The fitness of vertically transmitted symbionts being tightly

correlated to that of their hosts, they are expected to be selected for low virulence (Ewald 1983).

This reasoning, just like the transmission/virulence trade-off hypothesis, rests solely on the

inter-host level of selection, ignoring the potential effects of intra-host selection. Indeed, the most

competitive symbionts within a host may replicate faster and be more virulent than what would be

optimal from the standpoint of inter-host selection (Alizon et al. 2013). However, the causal link

from vertical transmission to low virulence is well supported by the experimental evolution of

parasites following the enforcement of different transmission modes (Bull et al. 1991, Turner et al.

1998, Messenger et al. 1999, Stewart et al. 2005). Accordingly, the most widespread

vertically-transmitted symbiont in insects, the intracellular bacterium *Wolbachia*, is often found to

be either avirulent (Hoffmann et al. 1994, Giordano et al. 1995, Bourtzis et al. 1996, Poinsot &

Merçot 1997, Hoffmann et al. 1998) or slightly virulent (Hoffmann et al. 1990, Turelli & Hoffmann

1995, Clancy & Hoffmann 1997).

The most striking exception to this pattern is *w*MelPop, a *Wolbachia* strain hosted by

Drosophila melanogaster that drastically reduces its host lifespan at high rearing temperatures

(Min & Benzer 1997). This exceptional virulence raises the question of its causes, both proximal

and ultimate. Min & Benzer (1997) suggested that the proximal cause of the virulence of *w*MelPop

is its over-replication, which was further supported by subsequent studies (McGraw et al. 2002,

Strunov et al. 2013). Furthermore, the repetition of an eight-gene region ("octomom"), has recently

been proposed as the genomic basis of *w*MelPop high density and virulence (Chrostek & Teixeira

2015). However, both the link between octomom copy number and density and the link between

density and virulence have been called into question (Rohrscheib et al. 2016). The ultimate cause

of *w*MelPop virulence has been assumed to be non-adaptive: as this *Wolbachia* strain is not known

to occur in nature, and as flies are typically reared at temperature lower than those at which

*w*MelPop is virulent and reproduce early in the lab, there is no selective pressure for reduced

virulence (Reynolds et al. 2003). If this explanation is correct, raising *w*MelPop-infected flies at a

high temperature, while enforcing late reproduction, should select for: (i) a reduced *w*MelPop

virulence, (ii) a reduced *w*MelPop density (under the assumption that *w*MelPop virulence at high

temperature is due to its high density), and (iii) a reduced number of octomom copies (under the

assumption that *w*MelPop virulence at high temperature is due to its high octomom copy number).

In the present study, we used experimental evolution over 17 generations to test these

hypotheses, raising *w*MelPop infected flies at 29°C, and enforcing reproduction eight days after

emergence. In addition, in one experimental condition, we fed the flies with paraquat, as it was

previously shown to increase the survival of *w*MelPop-infected flies (while having no effect on the

survival of uninfected flies) and to reduce the symbiotic density (Monnin et al. 2016). We

hypothesised that this treatment would reduce or cancel the selective pressure for reduced

virulence, density, and octomom copy number, resulting in higher values for these parameters,

compared to the control condition. Our results contribute to the ongoing debate concerning the

relationships between *w*MelPop octomom copy number, density, and virulence (Rohrscheib et al.

2016, Chrostek & Teixeira 2017, Rohrscheib et al. 2017).

Material & methods

<u>Biological material</u>

We used 17 isofemale lines of *w*MelPop-infected *D. melanogaster*[w1118]. These lines,

originated from flies provided by Scott O'Neill (Monash University, Australia), were founded

approximately six months before the start of the experiment, and then maintained under controlled

rearing conditions at 18°C (12 LD cycle), a temperature at which *w*MelPop is expected to be

avirulent (Reynolds et al. (2003) found no effect of the infection on mortality rate at 19°C and no

virulence of wMelPop was ever detected at a lower temperature). The experiment, by contrast, was

conducted at 29°C (12 LD cycle), a temperature at which wMelPop is known to be virulent (Min

& Benzer 1997, Strunov et al. 2013, Monnin et al. 2016). The flies were fed, and laid eggs on,

standard drosophila medium (David & Clavel 1965), both before and during the experiment.

Experimental evolution procedures

Emerging flies from the 17 isofemale lines were mixed to constitute the founding generation

G0 and were allowed to lay eggs. Twenty experimental lines were then constituted, each originating

from 100 randomly-picked eggs that were deposited in vials containing 1.5 g of *Drosophila*

medium. The experimental lines thus constituted were split into 10 "paraquat lines", in which

larvae developed on drosophila medium supplemented with 150 µL of a 10 mM paraquat solution

(as described in Monnin et al. 2016), and 10 "control lines", in which the paraquat was replaced by

150 µL of water. The G1 flies emerging from each vial were then allowed to lay their eggs on

unsupplemented medium for eight days. For each experimental line, a subset (n=100) of the eggs

laid on the eighth day following emergence was deposited in new vial similar to those in which the

parents developed – *i.e.*, paraquat-supplemented medium for the paraquat lines and water-

supplemented medium for the control lines. The protocol was repeated for the emerging G2 flies,

and again until G17. The evolution of wMelPop virulence, density and mean octomom copy

number was measured as follow: in each line, eggs laid seven days after emergence were collected

at G2, G3, G9, G12 and G17 and allowed to develop on unsupplemented medium. Then, emerging

females were used either to assess their survival, or their *Wolbachia* density and mean octomom

copy number (fig. 1).

Figure 1. Summary diagram of the experimental evolution experiment, showing the generations (G0-G17) at which the different measurements were performed.

Though we chose – based on preliminary observations – the eight days period to enforce

late reproduction while reducing the risk of line extinction, several lines nevertheless went extinct

before the end of the experiment, as a result of lack of eggs. Three control lines and two paraquat

lines went extinct between G3 and G9, four control lines and one paraquat line between G9 and

G12, and one control line and two paraquat lines between G12 and G17.

Survival assays

Fly survival was measured, as a proxy for *w*MelPop virulence, at G3 and G9. At emergence,

15 females from each experimental line were dispatched into three vials containing

sugar-supplemented agar (10% sugar). The surviving flies were then counted every day. The line

extinction rate was calculated as another proxy for *w*MelPop virulence, both before and after the

last survival measurement (G9).

DNA extraction

The *w*MelPop density and mean octomom copy number were measured at G2, G9, G12 and

G17. For each generation and line, four of the emerging females were transferred on

sugar-supplemented agar (10% sugar), to be collected four days later and then kept at -80°C.

Because of technical issues during DNA extraction or qPCR, the final number of replicates varies

from 2 to 4 (mean number > 3.8). The DNA extractions were performed using the 96 wells Biobasic

EZ-10 BS4372 kit. The entire flies were mechanically crushed for 30 s at 25 Hz using a 5-mm

stainless steel bead in a TissueLyser (Qiagen). Three hundred µL of ACL solution and 20 µL of

16 g.L^{-1} proteinase K were then added to the samples. Following an overnight incubation at 56°C,

300 µL of AB solution were added. The samples were then transferred on the plate from the kit,

and the purification was performed following the instructions of the manufacturer. The DNA was

eluted in 100 µL of elution buffer and stored at -20°C.

Quantitative PCR

Both relative *w*MelPop density and mean octomom copy number were measured in each

DNA extract by quantitative PCR, conforming to the Minimum Information for Publication of

Quantitative real-time PCR Experiments (MIQE) guidelines (Bustin et al. 2009). The relative

*w*MelPop density was quantified by amplifying two monocopy genes: one *Wolbachia* gene

(*wd0505*) and one *D. melanogaster* gene (*rp49*). The mean octomom copy number was quantified

by amplifying two *Wolbachia* genes: one that is part of the octomom region (*wd0513*), and one that

is not (*wd0505*). The primers used to amplify *rp49* were as follow: *rp49dd*-F: 5'-CTG-CCC-ACC-

GGA-TTC-AAG-3', *rp49dd*-R: 5'-CGA-TCT-CGC-CGC-AGT-AAA-C-3'. The primers used to

amplify the *wd0505* and *wd0513* genes were published in Chrostek et al. (2013). The primers were

synthetized by Eurogentec (Seraing, Belgium). qPCR mixes consisted of 5 μL of Sso Advanced

SYBR Green Supermix (Bio-Rad, Hercules, California, USA), 1 μL of water, 0.5 μL of each primer

(10 mM), and 3 μL of DNA samples diluted at 1/10. The reaction conditions for amplification were

95°C for 3 min, followed by 40 cycles of 95°C for 10 s, 59°C for 10 s and 68°C for 15 s. A melting

curve was recorded from 70°C to 95°C to ensure the specific amplification of the transcript.

Amplification and detection of DNA by quantitative PCR were performed with CFX96 instrument

(Bio-Rad). Standard curves were plotted using seven dilutions (10^2 to 10^7 copies) of a previously

amplified PCR product that had been purified using the kit Nucleospin extract II (Machery-Nagel).

Primer sets exhibited mean efficiencies of 93.2% ± 1.2 for *rp49dd*, 91.0% ± 0.7 for *wd0505,* and

92.7% ± 0.6 for *wd0513*. Two technical replicates per biological replicate were used for the

determination of DNA starting quantities. As the deviations between these duplicates were below

0.5 cycles, the mean Cp values were calculated. As PCR efficiency were close to 100%, we

calculated the relative *w*MelPop density and the mean octomom copy number using the following

formulas: $2^{Cp\_rp49-Cp\_wd0505}$ and $2^{Cp\_wd0505-Cp\_wd0513}$, respectively.

<u>Relationship between mean octomom copy number and the age of the host</u>

To assess the relationship between the mean octomom copy number and the age of the host,

we performed qPCR as described above on samples obtained in the course of a previously

published experiment (Monnin et al. 2016). Briefly, *w*MelPop-infected *D. melanogaster*[w1118] were

raised until emergence on standard drosophila medium supplemented with water (control),

paraquat (as described above). Emerging females were either immediately collected and kept at

-80°C (1 fly per treatment) or transferred on sugar-supplemented agar (10% sugar) to be collected

at days 7, 14 and 21 of adulthood (four flies per day and treatment). DNA extractions were

performed as described above.

<u>Statistical analyses</u>

We used the R software (version 3.3.2) for all analyses (R Core Team 2016).

We tested for evolution in survival between G3 and G9 using a mixed generalized linear

model with a gamma distribution. Generation (G3 or G9) and treatment (control or paraquat) were

treated in the model as fixed factors, while the line and vial factors were included as random effects.

The evolution of density and mean octomom copy number was tested using mixed linear

models, with the generation included as a fixed quantitative variable, the treatment as a fixed

qualitative variable, and the line factor included as a random effect. Normality and

homoscedasticity were checked graphically for both models.

The relationship between *Wolbachia* density and survival at G9 was tested by fitting a linear

model with median longevity as the response variable, mean density as a quantitative explanatory

variable, and the treatment (control or paraquat) as an explanatory factor.

The relationship between density and mean octomom copy number was tested by fitting a

mixed linear model with density as the response variable, mean octomom copy number as a fixed

quantitative explanatory variable, and generation, line and treatment as random effects.

The relationship between age and mean octomom copy number was tested by fitting a linear

model with mean octomom copy number as the response variable, age as a quantitative explanatory

variable and treatment as a qualitative explanatory variable.

Results & discussion

<u>Test of the predictions</u>

We followed the evolution of *w*MelPop virulence, density, and octomom copy number at a high temperature while enforcing late reproduction. Our first prediction was that during the experimental evolution, *w*MelPop virulence would have decreased in the control lines, and less so, or not at all, in the paraquat lines. To test this, we compared fly survival at G3 and G9.

Survival data at G3 confirmed that mortality began before the eighth day of the adult life of the flies (fig. 2a), suggesting that the late reproduction we enforced should have induced a selective pressure for decreased *w*MelPop virulence, at least in the control lines. In the paraquat lines, we expected a lesser reduction, or no reduction at all, in virulence.

Neither the interaction between generation and treatment (χ^2_1=3.23, p=0.07) nor the treatment effect (χ^2_1=0.38, p=0.54) was statistically significant, suggesting that contrary to our prediction, the paraquat treatment did not modify the evolution of virulence, as compared to the control condition. Furthermore, there was no effect of the generation factor, indicating that virulence did not decrease between G3 and G9 (χ^2_1=0.98, p=0.32; fig. 2b; median longevity at G3 in the control lines: 11 days; at G9 in the control lines: 10.5 days; at G3 in the paraquat lines: 10 days; at G9 in the paraquat lines: 11 days). As the survival was not assessed after G9, *w*MelPop virulence possibly decreased subsequently. As a decreased virulence would presumably lead to a decreased probability of extinction, we tested this hypothesis by comparing the rate of line extinction per generation before and after G9. In the control condition, it was 0.038 before G9 (3 out of 10 lines went extinct at some point during these eight generations), and 0.079 after G9 (5 out of 7 lines went extinct at some point during these nine generations), suggesting that no decrease

in virulence occurred during the experimental evolution, even after G9. Likewise, in the paraquat

condition, the rate of line extinction per generation was 0.025 before G9 (2 out of 10 lines went

extinct at some point during these eight generations), and 0.042 after G9 (3 out of 8 lines went

extinct at some point during these nine generations).

Figure 2. Evolution of fly survival. (a) Survival of female flies in the control condition at G3. Dotted lines represent
the 95 % confidence interval. The red vertical line shows the age (8 days) at which reproduction was enforced during
the experimental evolution. (b) Mean longevity (± standard error) in all lines in the paraquat (orange) and control
(black) conditions, at generations 3 and 9.

The discrepancy between the first prediction and the results invalidate the next two

predictions, as the expected reduction in *Wolbachia* density was supposed to causally explain the

expected reduction in virulence, and the expected reduction in mean octomom copy number was

supposed to causally explain the expected reduction in density. Accordingly, we observed neither

a decrease in density nor in mean octomom copy number during the experimental evolution. On

the contrary, both density ($\chi^2_1=27.84$, p<0.001; fig. 3a) and mean octomom copy number

($\chi^2_1=16.95$, p<0.001; fig. 3b) increased (mean density in the control lines at G2: 11.73; G9: 26.33;

G12: 21.05; G17: 20.50; in the paraquat lines at G2: 13.05; G9: 12.74; G12: 18.59; G17: 33.23;

mean octomom copy number in the control lines at G2: 2.50; G9: 3.45; G12: 2.66; G17: 3.43; in

the paraquat lines at G2: 2.84; G9: 2.67; G12: 4.27; G17: 3.97). Neither the interactions between

generation and treatment (density: χ^2_1=0.52, p=0.47; octomom copy number: χ^2_1=1.21, p=0.27) nor

the treatment effects (density: χ^2_1=0.44, p=0.51; octomom copy number: χ^2_1=0.44, p=0.51) were

statistically significant.

Figure 3. Evolution of *Wolbachia* density and octomom copy number. (a) Mean *Wolbachia* relative density (± standard
error) in all lines in the paraquat (orange) and control (black) conditions, at generations 2, 9, 12 and 17. (b) Mean
octomom copy number (± standard error) in all lines in the paraquat (orange) and control (black) conditions, at
generations 2, 9, 12 and 17.

As some lines went extinct before the end of the experiment, our statistical analyses may

have been biased if those extinctions were not random. However, this is unlikely to explain why

our predictions were not vindicated. Indeed, although a reduction in virulence could have been

concealed by the preferential extinction of lines exhibiting low virulence, this is the opposite of

what we would expect.

Potential explanations for the falsification of the predictions

The falsification of our hypotheses is especially puzzling, as our results show that both

density and octomom copy number evolved, suggesting that it is not by lack of time or heritable

variability that the *D. melanogaster–w*MelPop interaction did not evolve in the expected ways.

Among our starting assumptions was that the virulence of *w*MelPop is due to its high

density, which in turn is due to its high octomom copy number. It is therefore noteworthy that both

assumptions were recently called into question (Rohrscheib et al. 2016; see also Chroskek &

Teixeira 2017, Rohrscheib et al. 2017). If, as claimed by Rohrscheib et al. (2016), the virulence of

*w*MelPop is independent of its density and octomom copy number, our second and third predictions

do not hold. Although this would not explain why virulence did not decrease during the

experimental evolution, it would remove our puzzlement regarding the increase in *Wolbachia*

density and octomom copy number. However, our results do not support Rohrscheib et al. (2016)

findings. First, *w*MelPop density and fly survival were negatively related at G9, the only generation

at which both survival and density data were collected ($F_{1,18}=5.25$, p=0.04; fig. 4a). Second,

*w*MelPop density and mean octomom copy number were found to be positively related ($\chi^2_1=116.31$,

p<0.001; fig. 4b).

Figure 4. (a) Correlation between *Wolbachia* density and fly survival. Each circle represents a line in the paraquat (orange) or control (black) condition, at G9. The black line represents the regression of median longevity on mean density (Data for both conditions are pooled, as the treatment effect was found not to be statistically significant). Statistics from Pearson's correlation test are indicated. (b) Correlation between *Wolbachia* density and octomom copy number. Each symbol represents an individual fly, from the paraquat (orange) or control (black) condition, at G2 (squares), G9 (circles), G12 (triangles), or G17 (crosses). The black line represents the regression of density on octomom copy number. Statistics from Pearson's correlation test, with all conditions pooled together, are indicated.

It is therefore more likely that our assessment of the evolutionary forces at play, rather than our assumptions regarding the proximal causes of *w*MelPop virulence, was mistaken. The increase, over time, in both density and octomom copy number appear to be consistent with two non-mutually exclusive hypotheses. First, this pattern could be explained by the relaxation of an unknown selective pressure maintaining a low octomom copy number in flies kept at relatively low temperatures without enforcement of late reproduction, perhaps amplified by a mutational bias favouring a high octomom copy number. As each new generation of experimental evolution was started with only 100 eggs, genetic drift could have been increased by the experimental evolution, possibly limiting the efficiency of natural selection. However, it remains difficult to see why the selective pressure in favour of a low octomom copy number should be weaker in the experimental evolution setting than in the maintenance conditions, as our current knowledge of *w*MelPop

suggests that the opposite is true: in the absence of *w*MelPop-induced mortality in the maintenance

conditions, the octomom copy number is not expected to be under selection.

A second explanation for the observed pattern is that it results from directional selection in

favour of high density, high octomom copy number variants, more or less constrained, in each line,

by the amount of available heritable variation. Other things being equal, flies harbouring low

density, low octomom copy number should be longer-lived than their counterparts infected by high

density, high octomom copy number variants. As a result, inter-host selection is expected to favour

the former over the latter. However, intra-host selection is also likely to affect the evolution of

virulence (Alizon et al. 2013). Indeed, one can expect rapidly-replicating *w*MelPop variants to

reach a higher frequency than their slowly-replicating counterparts, and therefore be more likely to

be transmitted to their host offspring. Crucially, the competitive advantage of rapidly-replicating

variants will increase with host age, as it is positively correlated with the number of *w*MelPop

generations. 'Late reproduction' may therefore allow intra-host selection to be more powerful than

inter-host selection. This is especially likely given that survival was still high at eight days (fig.

2a), suggesting that inter-host selection may have been relatively weak. Chrostek & Teixeira (2018)

provide evidence for such intra-host selection by showing that the mean octomom copy number

increases as flies age and reach a plateau by day eight of adulthood. Our own results confirm the

increase ($F_{1,24}=25.64$, $p<0.001$), but no plateau was observed (fig. 5).

Figure 5. Relationship between the age of the fly and the octomom copy number of *w*MelPop. Data pooled from two treatments (paraquat and control), as neither the interaction between age and treatment ($F_{1,22}=2.79$, p=0.11) nor the treatment effect ($F_{1,23}=0.14$, p=0.72) were significant.

307
308
309

More research is needed to determine the evolutionary consequences of this intra-host

dynamic. Rapid evolution of octomom copy number in *w*MelPop suggests that intra-host selection

could play an important role in the evolution of vertically transmitted symbiont and may contribute

to explain the persistence of relatively virulent *Wolbachia* strains in nature.

Taken together, our results show that *Wolbachia* traits associated with virulence can evolve

rapidly following a switch in selection regime. These evolutionary changes challenge common

assumptions on the evolution of virulence in vertically transmitted symbionts, as traits positively

correlated with virulence were found to evolve. This could indicate that the intra-host level of

selection plays a significant, yet underexplored, role in shaping associations with vertically

transmitted symbionts.

Data availability

Raw data and scripts are available online: https://doi.org/10.5281/zenodo.3778214

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article.

Acknowledgments

We thank Nicole Lara for providing the *Drosophila* media. This work was supported by the Agence Nationale de la Recherche: 'ImmunSymbArt' ANR-2010-BLAN-170101 and 'RESIST' ANR-16-CE02-0013-01. This work was performed within the framework of the LABEX ECOFECT (ANR-11-LABX-0048) of Université de Lyon, within the programme 'Investissements d'Avenir' (ANR-11-IDEX-0007).

References

Alizon S, de Roode JC & Michalakis Y (2013) Multiple infections and the evolution of virulence. *Ecol. Lett.* **16:** 556–567

Alizon S, Hurford A, Mideo N & Van Baalen M (2009) Virulence evolution and the trade-off hypothesis: History, current state of affairs and the future. *J. Evol. Biol.* **22:** 245–259

Anderson RM & May RM (1982) Coevolution of hosts and parasites. *Parasitology* **85:** 411

Bourtzis K, Nirgianaki A, Markakis G & Savakis C (1996) *Wolbachia* infection and cytoplasmic

incompatibility in *Drosophila* species. *Genetics* **144:** 1063–73

Bull JJ, Molineux IJ & Rice WR (1991) Selection of Benevolence in a Host-Parasite System.

Evolution **45:** 875–882

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl

MW, Shipley GL & Vandesompele J (2009). The MIQE guidelines: minimum information

for publication of quantitative real-time PCR experiments. *Clinical chemistry* **55**, 611-622.

Chrostek E, Marialva MSP, Esteves SS, Weinert L a, Martinez J, Jiggins FM & Teixeira L (2013)

Wolbachia variants induce differential protection to viruses in *Drosophila melanogaster*: a

phenotypic and phylogenomic analysis. *PLoS Genet.* **9:** e1003896

Chrostek E & Teixeira L (2015) Mutualism Breakdown by Amplification of *Wolbachia* Genes.

PLOS Biol. **13:** e1002065

Chrostek E & Teixeira L (2017) Comment on Rohrscheib et al. 2016 'Intensity of mutualism

breakdown is determined by temperature not amplification of *Wolbachia* genes'. *PLoS*

Pathog. **13:** e1006540

Chrostek E & Teixeira L (2018) Within host selection for faster replicating bacterial symbionts.

PLoS ONE **13**: e0191530.

Clancy DJ & Hoffmann AA (1997) Behavior of *Wolbachia* endosymbionts from *Drosophila*

simulans in *Drosophila serrata*, a novel host. *Am. Nat.* **149:** 975–988

David JR & Clavel MF (1965) Interaction entre le génotype et le milieu d'élevage. Conséquences

sur les caractéristiques du développement de la Drosophile. *Bull. Biol. Fr. Belg.* **99:** 369–378

Ewald PW (1983) Host-parasites relations, vectors, and the evolution of disease severity. *Annu.*

Rev. Ecol. Evol. Syst. **14:** 465–485

Giordano R, O'Neill SL & Robertson HM (1995) *Wolbachia* infections and the expression of

cytoplasmic incompatibility in *Drosophila sechellia* and *D. mauritiana*. *Genetics* **140:** 1307–

1317

Hoffmann AA, Clancy DJ & Merton E (1994) Cytoplasmic incompatibility in Australian

populations of *Drosophila melanogaster*. *Genetics* **136:** 993–999

Hoffmann AA, Turelli M & Harshman LG (1990) Factors affecting the distribution of cytoplasmic

incompatibility in *Drosophila simulans*. *Genetics* **126:** 933–948

Hoffmann AA, Hercus M & Dagher H (1998) Population dynamics of the *Wolbachia* infection

causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* **148:** 221–231

McGraw E a, Merritt DJ, Droller JN & O'Neill SL (2002) *Wolbachia* density and virulence

attenuation after transfer into a novel host. *Proc. Natl. Acad. Sci. U. S. A.* **99:** 2918–23

Messenger SL, Molineux IJ & Bull JJ (1999) Virulence evolution in a virus obeys a trade-off. *Proc.*

R. Soc. B Biol. Sci. **266:** 397–404

Min KT & Benzer S (1997) *Wolbachia*, normally a symbiont of Drosophila, can be virulent,

causing degeneration and early death. *Proc. Natl. Acad. Sci. U. S. A.* **94:** 10792–6

Monnin D, Kremer N, Berny C, Henri H, Dumet A, Voituron Y, Desouhant E & Vavre F (2016)

Influence of oxidative homeostasis on bacterial density and cost of infection in *Drosophila–*

Wolbachia symbioses. *J. Evol. Biol.* **29:** 1211–1222

Poinsot D & Merçot H (2014) *Wolbachia* infection in *Drosophila simulans*: does the female host

bear a physiological cost? *Evolution* **51:** 180–186

R Core Team (2016). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reynolds KT, Thomson LJ & Hoffmann AA (2003) The Effects of Host Age, Host Nuclear

Background and Temperature on Phenotypic Effects of the Virulent *Wolbachia* Strain

popcorn in *Drosophila melanogaster*. *Genetics* **1034:** 1027–1034

Rohrscheib CE, Frentiu FD, Horn E, Ritchie FK, van Swinderen B, Weible MW, O'Neill SL &

Brownlie JC (2017) Response to: Comment on Rohrscheib et al. 2016 'Intensity of mutualism

breakdown is determined by temperature not amplification of *Wolbachia* genes'. *PLOS*

Pathog. **13:** e1006521

Rohrscheib CE, Frentiu FD, Horn E, Ritchie FK, van Swinderen B, Weible MW, O'Neill SL &

Brownlie JC (2016) Intensity of Mutualism Breakdown Is Determined by Temperature Not

Amplification of *Wolbachia* Genes. *PLoS Pathog.* **12:** 1–15

Stewart AD, Logsdon Jr. JM & Kelley SE (2005) An empirical study of the evolution of virulence

under both horizontal and vertical transmission. *Evolution* **59:** 730–739

Strunov A, Kiseleva E & Gottlieb Y (2013) Spatial and temporal distribution of pathogenic

Wolbachia strain *w*MelPop in *Drosophila melanogaster* central nervous system under

different temperature conditions. *J. Invertebr. Pathol.* **114:** 22–30

Turelli M & Hoffmann AA (1995) Cytoplasmic Incompatibility in *Drosophila simulans*: Dynamics

and Parameter Estimates from Natural Populations. *Genetics* **140:** 1319–1338

Turner P, Cooper V & Lenski R (1998) Tradeoff between horizontal and vertical modes of

transmission in bacterial plasmids. *Evolution* **52:** 315–329