Experimental evolution of virulence and associated traits in a *Drosophila melanogaster* –

***Wolbachia* symbiosis**

David Monnin*, Natacha Kremer, Caroline Michaud, Manon Villa, Hélène Henri, Emmanuel

Desouhant, Fabrice Vavre

Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive

UMR 5558, F-69622 Villeurbanne, France

* Corresponding author: david.monnin@hotmail.fr (School of Biological and Chemical Sciences,

Queen Mary University of London, Mile End Road, London E1 4NS, UK)

Abstract

Evolutionary theory predicts that vertically transmitted symbionts are selected for low

virulence, as their fitness is directly correlated to that of their host. In contrast with this prediction,

the *Wolbachia* strain *w*MelPop drastically reduces its *Drosophila melanogaster* host lifespan at

high rearing temperatures. It is generally assumed that this feature is maintained because the

D. melanogaster–*w*MelPop symbiosis is usually not exposed to environmental conditions in which

the symbiont is virulent. To test this hypothesis, we submitted *w*MelPop-infected *D. melanogaster*

lines to 17 generations of experimental evolution at a high temperature, while enforcing late

reproduction by artificial selection. The fly survival was measured at different time points, as well

as two traits that have been proposed to be causally responsible for *w*MelPop virulence: its relative

density and the mean number of copies of octomom, an 8-genes region of the *Wolbachia* genome.

We hypothesised that these conditions (high temperature and late reproduction) would select for a

reduced *w*MelPop virulence, a reduced *w*MelPop density, and a reduced octomom copy number.

Our results indicate that density, octomom copy number and virulence are correlated to each other.

However, contrary to our expectations, we could not detect any reduction in virulence during the

course of evolution. We discuss the significance of our results with respect to the evolutionary

causes of *w*MelPop virulence.

Introduction

Symbionts live inside (endosymbionts) or on (ectosymbionts) bigger organisms referred to

as hosts. As a result, their evolutionary success crucially depends on their ability to colonize, or be

transmitted to, new hosts. Deleterious symbionts, also known as parasites, exploit their hosts in a

way that maximizes their transmission, but usually face a trade-off between instantaneous

transmission rate and opportunities for transmission. Indeed, the reproductive rate of a parasite is

likely to be positively correlated to both its instantaneous transmission rate and its virulence (the

reduction in fitness it incurs to its host). Increased virulence is in turn associated, notably through

increased host mortality, with a reduction in the opportunities for transmission (Anderson & May

1982, Ewald 1983, see Alizon et al. 2009 for a review and defence of this hypothesis).

Theoretical and experimental works suggest that the mode of transmission – which ranges

from fully vertical (from parents to offspring) to fully horizontal (contagion) – is an important

factor shaping parasite virulence. The fitness of vertically transmitted symbionts being tightly

correlated to that of their hosts, they are expected to be selected for low virulence (Ewald 1983).

This reasoning, just like the transmission/virulence trade-off hypothesis, rests solely on the

inter-host level of selection (*i.e.*, selection between symbionts infecting different hosts), ignoring

the potential effects of intra-host selection (*i.e.*, selection between symbionts infecting the same

host). Indeed, the most competitive symbionts within a host may replicate faster and be more

virulent than what would be optimal from the standpoint of inter-host selection (Alizon et al. 2013).

However, the causal link from vertical transmission to low virulence is well supported by the

experimental evolution of parasites following the enforcement of different transmission modes

(Bull et al. 1991, Turner et al. 1998, Messenger et al. 1999, Stewart et al. 2005). Accordingly, the

most widespread vertically-transmitted symbiont in insects, the intracellular bacterium *Wolbachia*,

is often found to be either avirulent (Hoffmann et al. 1994, Giordano et al. 1995, Bourtzis et al.

1996, Poinsot & Merçot 1997, Hoffmann et al. 1998) or slightly virulent (Hoffmann et al. 1990,

Turelli & Hoffmann 1995, Clancy & Hoffmann 1997).

The most striking exception to this pattern is *w*MelPop, a *Wolbachia* strain hosted by

Drosophila melanogaster that drastically reduces its host lifespan at high rearing temperatures

(Min & Benzer 1997). This exceptional virulence raises the question of its causes, both proximal

and ultimate. Min & Benzer (1997) suggested that the proximal cause of the virulence of *w*MelPop

is its over-replication, which was further supported by subsequent studies (McGraw et al. 2002,

Strunov et al. 2013). Furthermore, the repetition of an eight-gene region ("octomom"), has recently

been proposed as the genomic basis of *w*MelPop high density and virulence (Chrostek & Teixeira

2015). Octomom includes genes encoding proteins potentially involved in DNA replication, repair,

recombination, transposition or transcription (Chrostek et al. 2003) and was found to be correlated

with *w*MelPop density and virulence (Chrostek & Teixeira 2015). However, both the link between

octomom copy number and density and the link between density and virulence have been called

into question. Rohrscheib et al. (2016) indeed found that differences in survival between flies

reared at 24°C and 29°C cannot be explained by differences in bacterial density. Similarly,

comparing flies of the same age but reared at different temperatures, or of different ages but reared

at the same temperature, they found no correlation between octomom copy number and either

density or virulence. However, to exclude the possibility that octomom copy number has an effect

on density, and density on virulence, the effect of these variables should be assessed independently

of temperature and age. The ultimate cause of *w*MelPop virulence has been assumed to be

non-adaptive: as this *Wolbachia* strain is not known to occur in nature but may have originated in

the lab, and as flies are typically reared at temperature lower than those at which *w*MelPop is

virulent and reproduce early in the lab, there is no selective pressure for reduced virulence

(Reynolds et al. 2003). It was indeed found that *w*MelPop reduces fly survival at 25°C, but not at

19°C (Reynolds et al. 2003). If this explanation is correct, raising *w*MelPop-infected flies at a high

temperature, while enforcing late reproduction, should select for: (i) a reduced *w*MelPop virulence,

(ii) a reduced *w*MelPop density (under the assumption that *w*MelPop virulence at high temperature

is due to its high density), and (iii) a reduced number of octomom copies (under the assumption

that *w*MelPop virulence at high temperature is due to its high octomom copy number).

In the present study, we used experimental evolution in conjunction with artificial selection

over 17 generations to test these hypotheses, raising *w*MelPop infected flies at 29°C, and enforcing

reproduction eight days after emergence. In addition, in one experimental condition, we fed the

flies with paraquat (1,1′dimethyl-4-4′bipyridynium dichloride), a pro-oxidant compound and

herbicide that was previously shown to increase the survival of *w*MelPop-infected flies (while

having no effect on the survival of uninfected flies) and to reduce the symbiotic density (Monnin

et al. 2016). We hypothesised that this treatment would reduce or cancel the selective pressure for

reduced virulence, density, and octomom copy number, resulting in higher values for these

parameters, compared to the control condition. The results of these experiments were interpreted

in light of our assessment of the relationship between octomom copy number and the age of the

host. Despite not being known to occur in nature, the *D. melanogaster-w*MelPop association can

help to understand the real-life dynamics of host-*Wolbachia* associations, especially in their early

stages. Indeed, one can expect that the low virulence typically observed in *Wolbachia* symbioses

is the outcome of a relatively long process of coevolution between the symbiotic partners. Newly

formed associations, by contrast, might be more likely to exhibit levels of virulence similar to what

is observed in *D. melanogaster-w*MelPop associations. Furthermore, our results contribute to the

ongoing debate concerning the relationships between *w*MelPop octomom copy number, density,

and virulence (Rohrscheib et al. 2016, Chrostek & Teixeira 2017, Rohrscheib et al. 2017).

Material & methods

<u>Biological material</u>

We used 17 isofemale lines of *w*MelPop-infected *D. melanogaster*[w1118]. These lines,

originated from flies provided by Scott O'Neill (Monash University, Australia), were founded

approximately six months before the start of the experiment, and then maintained under controlled

rearing conditions at 18°C (12 LD cycle), a temperature at which *w*MelPop is expected to be

avirulent (Reynolds et al. (2003) found no effect of the infection on mortality rate at 19°C and no

virulence of *w*MelPop was ever detected at a lower temperature). The flies were maintained on

standard drosophila medium (David & Clavel 1965).

Experimental evolution procedures

The experiment was conducted at 29°C (12 LD cycle), a temperature at which wMelPop is known to be virulent (Min & Benzer 1997, Strunov et al. 2013, Monnin et al. 2016). Emerging flies from the 17 isofemale lines were mixed to constitute the founding generation G0 and were allowed to lay eggs. Twenty experimental lines were then constituted, each originating from 100 randomly-picked eggs that were deposited in vials containing 1.5 g of *Drosophila* medium. The experimental lines thus constituted were split into 10 "paraquat lines", in which larvae developed on drosophila medium supplemented with 150 µL of a 10 mM paraquat (1,1′dimethyl-4-4′bipyridynium dichloride) solution (as described in Monnin et al. 2016), and 10 "water lines", in which the paraquat was replaced by 150 µL of water. The G1 flies emerging from each vial were then allowed to lay their eggs on unsupplemented medium for eight days (the medium was not supplemented with paraquat or water, as this would have been a confounding factor for the subsequent measurements). The medium was changed regularly, including at the end of day seven, and the eggs already laid on it were therefore removed. For each experimental line, a subset (n=100) of the eggs laid on the eighth day following emergence was deposited in new vial similar to those in which the parents developed – *i.e.*, paraquat-supplemented medium for the paraquat lines and water-supplemented medium for the water lines. The protocol was repeated for the emerging G2 flies, and again until G17. The evolution of wMelPop virulence, density and mean octomom copy number was measured as follow: in each line, eggs laid seven days after emergence were collected at G2, G3, G9, G12 and G17 and allowed to develop on unsupplemented medium. Then, emerging females were used either to assess their survival, or their *Wolbachia* density and mean octomom copy number (fig. 1). These traits were not measured at every single generation, as

we do not expect the changes from one generation to the next to be so great as to justify measuring

all traits at each generation.

Figure 1. Summary diagram of the experimental evolution experiment, showing the generations (G0-G17) at which
the different measurements were performed.

Though we chose – based on preliminary observations – the eight days period to enforce

late reproduction while reducing the risk of line extinction, several lines nevertheless went extinct

before the end of the experiment, as a result of lack of egg laying. Three control lines and two

paraquat lines went extinct between G3 and G9, four control lines and one paraquat line between

G9 and G12, and one control line and two paraquat lines between G12 and G17.

Survival assays

Fly survival was measured, as a proxy for *w*MelPop virulence, at G3 and G9. At emergence,

15 females from each experimental line were dispatched into three vials containing

sugar-supplemented agar (10% sugar). The surviving flies were then counted every day. The line

extinction rate – the ratio of lines that went extinct over the total number of lines – was calculated

for both the water and the paraquat lines, both during the 8 generations prior to G9 and during the

9 subsequent generations. It serves as another proxy for wMelPop virulence. We used Fisher's

exact tests to assess differences in extinction rates.

DNA extraction

The wMelPop density and mean octomom copy number were measured at G2, G9, G12 and

G17. For each generation and line, four of the emerging females were transferred on

sugar-supplemented agar (10% sugar), to be collected four days later and then kept at -80°C.

Because of technical issues during DNA extraction or qPCR, the final number of replicates varies

from 2 to 4 (mean number > 3.8). The DNA extractions were performed using the 96 wells Biobasic

EZ-10 BS4372 kit. The entire flies were mechanically crushed for 30 s at 25 Hz using a 5-mm

stainless steel bead in a TissueLyser (Qiagen). Three hundred µL of ACL solution and 20 µL of

16 g.L^{-1} proteinase K were then added to the samples. Following an overnight incubation at 56°C,

300 µL of AB solution was added. The samples were then transferred on the plate from the kit, and

the purification was performed following the instructions of the manufacturer. The DNA was eluted

in 100 µL of elution buffer and stored at -20°C.

Quantitative PCR

Both relative wMelPop density and mean octomom copy number were measured in each

DNA extract by quantitative PCR, conforming to the Minimum Information for Publication of

Quantitative real-time PCR Experiments (MIQE) guidelines (Bustin et al. 2009). The relative

wMelPop density was quantified by amplifying two monocopy genes: one *Wolbachia* gene

(*wd0505*) and one *D. melanogaster* gene (*rp49*). The mean octomom copy number was quantified

by amplifying two *Wolbachia* genes: one that is part of the octomom region (*wd0513*), and one that

is not (*wd0505*). The primers used to amplify *rp49* were as follow: *rp49dd*-F: 5'-CTG-CCC-ACC-

GGA-TTC-AAG-3', *rp49dd*-R: 5'-CGA-TCT-CGC-CGC-AGT-AAA-C-3'. The primers used to

amplify the *wd0505* and *wd0513* genes were published in Chrostek et al. (2013). The primers were

synthetized by Eurogentec (Seraing, Belgium). qPCR mixes consisted of 5 µL of Sso Advanced

SYBR Green Supermix (Bio-Rad, Hercules, California, USA), 1 µL of water, 0.5 µL of each primer

(10 mM), and 3 µL of DNA samples diluted at 1/10. The reaction conditions for amplification were

95°C for 3 min, followed by 40 cycles of 95°C for 10 s, 59°C for 10 s and 68°C for 15 s. A melting

curve was recorded from 70°C to 95°C to ensure the specific amplification of the transcript.

Amplification and detection of DNA by quantitative PCR were performed with CFX96 instrument

(Bio-Rad). Standard curves were plotted using seven dilutions (10^2 to 10^7 copies) of a previously

amplified PCR product that had been purified using the kit Nucleospin extract II (Machery-Nagel).

Primer sets exhibited mean efficiencies of 93.2% $\pm$ 1.2 for *rp49dd*, 91.0% $\pm$ 0.7 for *wd0505,* and

92.7% $\pm$ 0.6 for *wd0513*. Two technical replicates per biological replicate were used for the

determination of DNA starting quantities. As the deviations between these duplicates were below

0.5 cycles, the mean Cp values were calculated. As PCR efficiency were close to 100%, we

calculated the relative *w*MelPop density and the mean octomom copy number using the following

formulas: $2^{\text{Cp\_}rp49-\text{Cp\_}wd0505}$ and $2^{\text{Cp\_}wd0505-\text{Cp\_}wd0513}$, respectively.

Relationship between mean octomom copy number and the age of the host

To evaluate the possibility that intra-host selection is involved in the evolution of *w*MelPop

octomom copy number, we assessed the relationship between the mean octomom copy number and

the age of the host. To do so, we performed qPCR as described above on samples obtained in the

course of a previously published experiment (Monnin et al. 2016). Briefly, *w*MelPop-infected

D. melanogaster[w1118] were raised until emergence on standard drosophila medium supplemented

with water or paraquat (as described above). Emerging females were either immediately collected

and kept at -80°C (1 fly per treatment) or transferred on sugar-supplemented agar (10% sugar) to

be collected at days 7, 14 and 21 of adulthood (four flies per day and per treatment). DNA

extractions were performed as described above.

<u>Statistical analyses</u>

We used the R software (version 3.3.2) for all analyses (R Core Team 2016).

We tested for evolution in survival between G3 and G9 using a mixed generalized linear

model with a gamma distribution (in the absence of censored data, a survival model is not

necessary, and the gamma distribution is suitable for continuous positive data). Individual

longevity was used as the response variable. Generation (G3 or G9) and treatment (water or

paraquat) were treated in the model as fixed factors, while the line and vial factors were included

as random effects.

The evolution of density and mean octomom copy number was tested using mixed linear

models, with the generation included as a fixed quantitative variable, the treatment as a fixed

qualitative variable, and the line factor included as a random effect. Normality and

homoscedasticity were checked graphically for both models.

The relationship between *Wolbachia* density and survival at G9 was tested by fitting a linear

model with median longevity as the response variable, mean density as a quantitative explanatory

variable, and the treatment (water or paraquat) as an explanatory factor.

220 The relationship between density and mean octomom copy number was tested by fitting a

221 mixed linear model with density as the response variable, mean octomom copy number as a fixed

222 quantitative explanatory variable, and generation, line and treatment as random effects.

223 The relationship between age and mean octomom copy number was tested by fitting a linear

224 model with mean octomom copy number as the response variable, age as a quantitative explanatory

225 variable and treatment as a qualitative explanatory variable.

Results & discussion

Test of the predictions

229 We followed the evolution of wMelPop virulence, density, and octomom copy number at a

230 high temperature while enforcing late reproduction. Our first prediction was that during the

231 experimental evolution, wMelPop virulence would have decreased in the water lines, and less so,

232 or not at all, in the paraquat lines. To test this, we compared fly survival at G3 and G9. We can be

233 confident that fly mortality is induced by wMelPop, as the strong reduction of host longevity

234 induced by wMelPop has been established previously. A study performed in our lab (Monnin et al.

235 2016), using the same *D. melanogaster* line and the same survival protocol, found a large mean

236 difference in longevity (4.4±0.66 days) between wMelPop-infected and uninfected individuals.

237 Survival data at G3 confirmed that mortality began before the eighth day of the adult life of

238 the flies (fig. 2a), suggesting that the late reproduction we enforced should have induced a selective

239 pressure for decreased wMelPop virulence, at least in the water lines. In the paraquat lines, we

240 expected a lesser reduction, or no reduction at all, in virulence.

Neither the interaction between generation and treatment (χ^2_1=3.23, p=0.07) nor the

treatment effect (χ^2_1=0.38, p=0.54) was statistically significant, suggesting that the paraquat

treatment did not modify the evolution of survival, as compared to the water condition.

Furthermore, there was no effect of the generation factor, indicating that survival did not increase

between G3 and G9 (χ^2_1=0.98, p=0.32; fig. 2b; median longevity at G3 in the control lines: 11 days;

at G9 in the control lines: 10.5 days; at G3 in the paraquat lines: 10 days; at G9 in the paraquat

lines: 11 days). As the survival was last assessed at G9, we cannot directly test whether it evolved

between G9 and G17. Given that an increased survival would presumably lead to a decreased

probability of extinction, we compared the rates of line extinction per generation before and after

G9. In the water condition, it was 0.038 before G9 (3 out of 10 lines went extinct at some point

during these eight generations), and 0.079 after G9 (5 out of 7 lines went extinct at some point

during these nine generations), showing that no increase in extinction rate occurred during the

experimental evolution, even after G9. Likewise, in the paraquat condition, the rate of line

extinction per generation was 0.025 before G9 (2 out of 10 lines went extinct at some point during

these eight generations), and 0.042 after G9 (3 out of 8 lines went extinct at some point during

these nine generations). Differences in extinction rate between the early and late generations were

not significant (Fisher's exact tests, p=0.15 in the water lines, p=0.61 in the paraquat lines, p=0.16

overall).

Taken together, these survival and extinction rates suggest the contrary to our prediction,

the virulence of *w*MelPop did not decrease during the experiment and was not differentially

affected by the treatment (paraquat or water).

Figure 2. Evolution of fly survival. (a) Survival of female flies in the water condition at G3. Dotted lines represent the 95 % confidence interval. The red vertical line shows the age (8 days) at which reproduction was enforced during the experimental evolution. The corresponding figure for the paraquat condition is provided as supplementary material (b) Mean longevity (± standard error) in all lines in the paraquat (orange) and water (black) conditions, at generations 3 and 9.

The discrepancy between the first prediction and the results invalidates the next two predictions, as the expected reduction in *Wolbachia* density was supposed to causally explain the expected reduction in virulence, and the expected reduction in mean octomom copy number was supposed to causally explain the expected reduction in density. Accordingly, we observed neither a decrease in density nor in mean octomom copy number during the experimental evolution. On the contrary, both density (χ^2_1=27.84, p<0.001; fig. 3a) and mean octomom copy number (χ^2_1=16.95, p<0.001; fig. 3b) increased (mean density in the water lines at G2: 11.73; G9: 26.33; G12: 21.05; G17: 20.50; in the paraquat lines at G2: 13.05; G9: 12.74; G12: 18.59; G17: 33.23; mean octomom copy number in the water lines at G2: 2.50; G9: 3.45; G12: 2.66; G17: 3.43; in the paraquat lines at G2: 2.84; G9: 2.67; G12: 4.27; G17: 3.97). Neither the interactions between generation and treatment (density: χ^2_1=0.52, p=0.47; octomom copy number: χ^2_1=1.21, p=0.27) nor

the treatment effects (density: $\chi^2_1=0.44$, p=0.51; octomom copy number: $\chi^2_1=0.44$, p=0.51) were

statistically significant.

Figure 3. Evolution of *Wolbachia* density and octomom copy number. (a) Mean *Wolbachia* relative density (± standard error) in all lines in the paraquat (orange) and water (black) conditions, at generations 2, 9, 12 and 17. (b) Mean octomom copy number (± standard error) in all lines in the paraquat (orange) and control (black) conditions, at generations 2, 9, 12 and 17.

Potential explanations for the falsification of the predictions

The falsification of our predictions is especially puzzling, as our results show that both

density and octomom copy number evolved, suggesting that it is not by lack of time or heritable

variability that the *D. melanogaster–w*MelPop interaction did not evolve in the expected ways.

As some lines went extinct before the end of the experiment, our statistical analyses may

have been biased if those extinctions were not random. However, it is unlikely to explain why our

predictions were not vindicated. Indeed, although a reduction in virulence could have been

concealed by the preferential extinction of lines exhibiting low virulence, this pattern is the

opposite of what we would expect. Furthermore, the observed increase in relative *Wolbachia*

density and mean octomom copy number is not likely to be due to a mere increase in genetic

similarity: at G2, no measured fly had a relative density superior to 40, contrary to 21% of the G17

flies (the maximum being 143); similarly, no flies at G2 had a mean octomom copy number over

6, contrary to 18% of the flies measured at G17 (the maximum being 12). Finally, we did not vary

the experimental temperature – a factor recently shown to impact on the evolution of

D. melanogaster-Wolbachia associations by Mazzucco et al. (2020) – and used only one host

genetic background (*D. melanogaster*[w1118]) and one *Wolbachia* strain (*w*MelPop). We cannot

exclude the possibility that different conditions would have led to different coevolutionary

outcomes.

Among our starting assumptions was that the virulence of *w*MelPop is due to its high

density, which in turn is due to its high octomom copy number. It is therefore noteworthy that both

assumptions were recently called into question (Rohrscheib et al. 2016; see also Chroskek &

Teixeira 2017, Rohrscheib et al. 2017). If, as claimed by Rohrscheib et al. (2016), the virulence of

*w*MelPop is independent of its density and octomom copy number, our second and third predictions

do not hold. Although this would not explain why virulence did not decrease during the

experimental evolution, it would remove our puzzlement regarding the increase in *Wolbachia*

density and octomom copy number. However, our results do not support Rohrscheib et al. (2016)

findings. First, *w*MelPop density and fly survival were negatively related at G9, the only generation

at which both survival and density data were collected ($F_{1,18}=5.25$, $p=0.04$; fig. 4a). Second,

*w*MelPop density and mean octomom copy number were found to be positively related ($\chi^2_1=116.31$,

$p<0.001$; fig. 4b).

Figure 4. (a) Correlation between *Wolbachia* density and fly survival. Each circle represents a line in the paraquat (orange) or water (black) condition, at G9. The black line represents the regression of median longevity on mean density (data for both conditions are pooled, as the treatment effect was found not to be statistically significant). Statistics from Pearson's correlation test are indicated. (b) Correlation between *Wolbachia* density and octomom copy number. Each symbol represents an individual fly, from the paraquat (orange) or water (black) condition, at G2 (squares), G9 (circles), G12 (triangles), or G17 (crosses). The black line represents the regression of density on octomom copy number. Statistics from Pearson's correlation test, with all conditions pooled together, are indicated.

It is therefore more likely that our assessment of the evolutionary forces at play, rather than our assumptions regarding the proximal causes of *w*MelPop virulence, was mistaken. The increase, over time, in both density and octomom copy number, appears to be consistent with two non-mutually exclusive hypotheses. First, this pattern could be explained by the relaxation of an unknown selective pressure maintaining a low octomom copy number in flies kept at relatively low temperatures without enforcement of late reproduction, perhaps amplified by a mutational bias favouring a high octomom copy number. As each new generation of experimental evolution was started with only 100 eggs, genetic drift could have been increased by the experimental evolution/selection protocol, possibly limiting the efficiency of natural selection. However, it remains difficult to see why the selective pressure in favour of a low octomom copy number should be weaker in the experimental evolution setting than in the maintenance conditions, as our current

knowledge of *w*MelPop suggests that the opposite is true: in the absence of *w*MelPop-induced

mortality in the maintenance conditions, the octomom copy number is not expected to be under

selection.

A second explanation for the observed pattern is that it results from directional selection in

favour of high density, high octomom copy number variants, more or less constrained, in each line,

by the amount of available heritable variation. Other things being equal, flies harbouring low

density, low octomom copy number should be longer-lived and therefore have more opportunities

to transmit their symbionts to their offspring than their counterparts infected by high density, high

octomom copy number variants. As a result, inter-host selection is expected to favour *w*Melpop

variants that do not reach densities so high that they compromise the fitness of their host. However,

intra-host selection is also likely to affect the evolution of virulence (Alizon et al. 2013). Indeed,

one can expect high density, high octomom copy number variants to replicate more rapidly than

(and therefore outnumber) their low density, low octomom copy number counterparts, which will

make them more likely to be transmitted to their host offspring. Crucially, the competitive

advantage of rapidly-replicating variants will increase with host age, as it is positively correlated

with the number of *w*MelPop generations. (At 29°C, the doubling time of *w*MelPop was estimated

by Duarte et al. (preprint) to vary from 0.88 to 1.38 days.) 'Late reproduction' may therefore allow

intra-host selection to be more powerful than inter-host selection. This is especially likely given

that survival was still high at eight days (fig. 2a), suggesting that inter-host selection may have

been relatively weak. Chrostek & Teixeira (2018) provide evidence for such intra-host selection

by showing that the mean octomom copy number increases as flies age and reach a plateau by day

eight of adulthood. Our own results confirm the increase ($F_{1,24}=25.64$, $p<0.001$), but no plateau

was observed (fig. 5). These results show that intra-host variability can exist, and therefore allow

for intra-host selection, despite the bottleneck experienced by vertically transmitted symbionts

(estimated to range from 850 to 8000 cells in aphid-*Buchnera* symbioses (Mira & Moran 2002)).

Figure 5. Relationship between the age of the fly and the octomom copy number of *w*MelPop. Data pooled from two
treatments (paraquat and water), as neither the interaction between age and treatment ($F_{1,22}=2.79$, p=0.11) nor the
treatment effect ($F_{1,23}=0.14$, p=0.72) were significant.

More research is needed to determine the evolutionary consequences of this intra-host

dynamic. Rapid evolution of octomom copy number in *w*MelPop suggests that intra-host selection

could play an important role in the evolution of vertically transmitted symbiont and may contribute

to explain the persistence of relatively virulent *Wolbachia* strains in nature.

Taken together, our results show that *Wolbachia* traits associated with virulence can evolve

rapidly following a switch in selection regime. These evolutionary changes challenge common

assumptions on the evolution of virulence in vertically transmitted symbionts, as traits positively

correlated with virulence were found to evolve. This could indicate that the intra-host level of

selection plays a significant, yet underexplored, role in shaping associations with vertically

transmitted symbionts.

Data availability

Raw data, scripts and additional figure are available online:

https://doi.org/10.5281/zenodo.4065517

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content

of this article. NK & FV are PCIEvolBiol recommenders.

Acknowledgments

We thank Nicole Lara for providing the *Drosophila* media. This work was supported by the Agence

Nationale de la Recherche: 'ImmunSymbArt' ANR-2010-BLAN-170101 and 'RESIST' ANR-16-

CE02-0013-01. This work was performed within the framework of the LABEX ECOFECT (ANR-

11-LABX-0048) of Université de Lyon, within the programme 'Investissements d'Avenir' (ANR-

11-IDEX-0007).

References

Alizon S, de Roode JC & Michalakis Y (2013) Multiple infections and the evolution of virulence.

Ecol. Lett. **16:** 556–567

Alizon S, Hurford A, Mideo N & Van Baalen M (2009) Virulence evolution and the trade-off

hypothesis: History, current state of affairs and the future. *J. Evol. Biol.* **22:** 245–259

Anderson RM & May RM (1982) Coevolution of hosts and parasites. *Parasitology* **85:** 411

Bourtzis K, Nirgianaki A, Markakis G & Savakis C (1996) *Wolbachia* infection and cytoplasmic

incompatibility in *Drosophila* species. *Genetics* **144:** 1063–73

Bull JJ, Molineux IJ & Rice WR (1991) Selection of Benevolence in a Host-Parasite System.

Evolution **45:** 875–882

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl

MW, Shipley GL & Vandesompele J (2009). The MIQE guidelines: minimum information

for publication of quantitative real-time PCR experiments. *Clinical chemistry* **55**, 611-622.

Chrostek E, Marialva MSP, Esteves SS, Weinert L a, Martinez J, Jiggins FM & Teixeira L (2013)

Wolbachia variants induce differential protection to viruses in *Drosophila melanogaster*: a

phenotypic and phylogenomic analysis. *PLoS Genet.* **9:** e1003896

Chrostek E & Teixeira L (2015) Mutualism Breakdown by Amplification of *Wolbachia* Genes.

PLOS Biol. **13:** e1002065

Chrostek E & Teixeira L (2017) Comment on Rohrscheib et al. 2016 'Intensity of mutualism

breakdown is determined by temperature not amplification of *Wolbachia* genes'. *PLoS*

Pathog. **13:** e1006540

Chrostek E & Teixeira L (2018) Within host selection for faster replicating bacterial symbionts. *PLoS ONE* **13**: e0191530.

Clancy DJ & Hoffmann AA (1997) Behavior of *Wolbachia* endosymbionts from *Drosophila simulans* in *Drosophila serrata*, a novel host. *Am. Nat.* **149:** 975–988

David JR & Clavel MF (1965) Interaction entre le génotype et le milieu d'élevage. Conséquences sur les caractéristiques du développement de la Drosophile. *Bull. Biol. Fr. Belg.* **99:** 369–378

Duarte EH, Carvalho A, Lopez-Madrigal S & Teixeira L (2020) Regulation of *Wolbachia* proliferation by the amplification and deletion of an addictive genomic island. *bioRxiv* (https://doi.org/10.1101/2020.09.08.288217)

Ewald PW (1983) Host-parasites relations, vectors, and the evolution of disease severity. *Annu. Rev. Ecol. Evol. Syst.* **14:** 465–485

Giordano R, O'Neill SL & Robertson HM (1995) *Wolbachia* infections and the expression of cytoplasmic incompatibility in *Drosophila sechellia* and *D. mauritiana*. *Genetics* **140:** 1307–1317

Hoffmann AA, Clancy DJ & Merton E (1994) Cytoplasmic incompatibility in Australian populations of *Drosophila melanogaster*. *Genetics* **136:** 993–999

Hoffmann AA, Turelli M & Harshman LG (1990) Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* **126:** 933–948

Hoffmann AA, Hercus M & Dagher H (1998) Population dynamics of the *Wolbachia* infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* **148:** 221–231

Mazzucco R, Nolte V, Vijayan T & Schlötterer C (2020) Long-Term Dynamics Among *Wolbachia*

Strains During Thermal Adaptation of Their *Drosophila melanogaster* Hosts. *Frontiers in*

Genetics **11**: 482.

McGraw E a, Merritt DJ, Droller JN & O'Neill SL (2002) *Wolbachia* density and virulence

attenuation after transfer into a novel host. *Proc. Natl. Acad. Sci. U. S. A.* **99:** 2918–23

Messenger SL, Molineux IJ & Bull JJ (1999) Virulence evolution in a virus obeys a trade-off. *Proc.*

R. Soc. B Biol. Sci. **266:** 397–404

Min KT & Benzer S (1997) *Wolbachia*, normally a symbiont of Drosophila, can be virulent,

causing degeneration and early death. *Proc. Natl. Acad. Sci. U. S. A.* **94:** 10792–6

Mira A & Moran NA (2002) Estimating population size and transmission bottlenecks in maternally

transmitted endosymbiotic bacteria. *Microbial Ecology* **44:** 137-143.

Monnin D, Kremer N, Berny C, Henri H, Dumet A, Voituron Y, Desouhant E & Vavre F (2016)

Influence of oxidative homeostasis on bacterial density and cost of infection in *Drosophila–*

Wolbachia symbioses. *J. Evol. Biol.* **29:** 1211–1222

Poinsot D & Merçot H (2014) *Wolbachia* infection in *Drosophila simulans*: does the female host

bear a physiological cost? *Evolution* **51:** 180–186

R Core Team (2016). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reynolds KT, Thomson LJ & Hoffmann AA (2003) The Effects of Host Age, Host Nuclear

Background and Temperature on Phenotypic Effects of the Virulent *Wolbachia* Strain

popcorn in *Drosophila melanogaster*. *Genetics* **1034:** 1027–1034

Rohrscheib CE, Frentiu FD, Horn E, Ritchie FK, van Swinderen B, Weible MW, O'Neill SL &

Brownlie JC (2017) Response to: Comment on Rohrscheib et al. 2016 'Intensity of mutualism

breakdown is determined by temperature not amplification of *Wolbachia* genes'. *PLOS*

Pathog. **13:** e1006521

Rohrscheib CE, Frentiu FD, Horn E, Ritchie FK, van Swinderen B, Weible MW, O'Neill SL &

Brownlie JC (2016) Intensity of Mutualism Breakdown Is Determined by Temperature Not

Amplification of *Wolbachia* Genes. *PLoS Pathog.* **12:** 1–15

Stewart AD, Logsdon Jr. JM & Kelley SE (2005) An empirical study of the evolution of virulence

under both horizontal and vertical transmission. *Evolution* **59:** 730–739

Strunov A, Kiseleva E & Gottlieb Y (2013) Spatial and temporal distribution of pathogenic

Wolbachia strain *w*MelPop in *Drosophila melanogaster* central nervous system under

different temperature conditions. *J. Invertebr. Pathol.* **114:** 22–30

Turelli M & Hoffmann AA (1995) Cytoplasmic Incompatibility in *Drosophila simulans*: Dynamics

and Parameter Estimates from Natural Populations. *Genetics* **140:** 1319–1338

Turner P, Cooper V & Lenski R (1998) Tradeoff between horizontal and vertical modes of

transmission in bacterial plasmids. *Evolution* **52:** 315–329