# Coevolution of song and egg coloration: multimodal mating signals?

Silu Wang
University of California, Berkeley
Email: siluwang.evo@gmail.com

**Abstract**
The divergence of reproductive traits frequently underpins the evolution of reproductive
isolation. One of the most enduring puzzles on this subject concerns the variability in egg
coloration among species of tinamou (Tinamidae) —a group of birds endemic to neotropics
(Cabot 1992). Specifically, some tinamous lay glossy and colorful eggs while others lay less
colorful eggs. Here I tested the hypothesis that tinamou egg coloration is a mating signal and its
diversification was driven by reinforcement. For most tinamou species, the male guard the nest
that is sequentially visited and laid eggs in by multiple females. The colorations of the existing
eggs in the nest could signal mate quality and species identities to the upcoming females to the
nest, preventing costly hybridization, thus were selected to diverge among species. If so, egg
colors should coevolve with the known mating signals as the tinamou lineages diverged. The
tinamou songs are important mating signals and are highly divergent among species. I found that
the egg luminance was significantly associated with the first principle component of the song
variables among 31 tinamou species (after correcting for phylogenetic signal). Egg color and
songs could be multimodal mating signals that are divergently selected as different tinamou
species diverged. Mating signal evolution could be opportunistic and even exploit post-mating
trait as premating signals.

26
**Introduction**
Reproductive trait divergence is crucial for speciation because these traits frequently
underpin barriers to gene flow (Koski and Ashman 2016; Pfennig 2016). A very puzzling
reproductive trait divergence exists in tinamou egg coloration. Tinamiformes (common name:
tinamou) is the most species-rich order of Palaeognathae, containing 48 extant species (Figure 1)
(Cabot 1992; Davies 2002). Tinamous are mostly dull in plumage, but various species of
tinamous lay brightly colored eggs ranging from brilliant magenta, to pink, purple, turquoise, and
olive green (Figure 1) (Cabot 1992). Despite its evolutionary importance, tinamou egg color
divergence remains a mystery due to the cryptic and elusive plumage and behavior of the birds
(Cabot 1992; Davies 2002).
Two main hypotheses have been made about evolutionary mechanism of egg color
divergence: (1) aposematism hypothesis (Swynnerton 1916) states that eggs are brightly colored
to warn predators of their distastefulness; (2) mating signaling hypothesis, which predicts that
egg color are employed for mating recognition and attraction (Weeks 1973; Brennan 2010). The
aposematic function of egg coloration was thought to be unlikely, because egg predators are
nocturnal mammals or reptiles that prioritize chemical cues over visual cues (Skutch 1966; Cabot
1992; Brennan 2010). The mating signal hypothesis is plausible considering the ecology and
mating systems of tinamou species. In most tinamou species, males collect and incubate eggs
laid by multiple females (Cabot 1992; Davies 2002). An initial female is attracted by distinctive
songs to the nest guarded a male. If mating occurs, the female lays eggs in the nest and leaves
before other females come and lay more eggs that are all incubated by the same male (Cabot
1992; Davies 2002). The colors of the existing eggs in the nest could signal intra-specific mate
identity to prevent costly hybridization. Such alternative "mating channel" might be especially
favored when the plumage is adapted for camouflage (Cabot 1992; Davies 2002). If egg colors of
different tinamou species is adapted for mate recognition, reinforcement (Dobzhansky 1937;
Mayr 1942; Liou and Price 1994; Servedio 2000) could have driven egg color divergence among
different tinamou species.

Here I test the mating signal hypothesis of tinamou egg color divergence. If tinamou egg
colors are employed as a species-specific mate recognition signal, it should coevolve with other
mate-recognition signals as the tinamou lineages diverged. Songs of tinamous are known to be
important and highly divergent mating signals among species (Cabot 1992; Bertelli and Tubaro
2002; Laverde-R and Cadena 2014). In addition to songs, multimodality of mating signals is
selected for efficient mate-recognition in complex environment (Partan and Marler 1999; Rowe
1999; Secondi et al. 2015; Halfwerk et al. 2019). When glamorous plumage is costly due to
increased predation risk, alternative signal modality such as egg coloration could be favored.
Since the birds themselves (instead of the eggs) tend to attract nest predation (Brennan 2010),
egg coloration could be employed to fulfill signal multimodality, bypassing the evolutionary
constraints for plumage. To investigate this possibility, here I test whether egg coloration
coevolved with songs as tinamou species diverged.
66
67                                                **Methods**
To test coevolution of egg color and song in tinamous, I tested the association of songs
and egg colors among 32 tinamou species in R (Hothorn et al. 2008). Tinamou egg coloration
data was extracted from existing tinamou egg color analysis (Schläpfer 2017), which quantified
coloration of the eggs from 32 tinamou species (Figure 1). Since tinamou egg colors are known
to decay over the course museum storage, tinamou nest photos are used to best represent the
functional egg color (Schläpfer 2017). Briefly, for each species, the egg coloration reflected in
hue, chroma, and luminance in the CIELAB color space was quantified (Schläpfer 2017).
Song data was acquired from a previous study (Bertelli and Tubaro 2002), in which four
song variables from 40 tinamou species were quantified: maximum frequency (Hz), minimum
frequency (Hz), emphasized frequency (frequency of the note with highest amplitude in the song,
Hz), and bandwidth (the difference between maximum and minimum frequency, Hz). Because
these variables are correlated (Figure S1), I used principle component analysis (PCA) to generate
PC1 that captured 81.2% of the variation in song data. Four song variables were shifted to center
around zero and scaled to unit variance preceding PCA.
I tested whether egg hue, chroma, and luminance is respectively associated with song
PC1 among tinamous, accounting for the potential confounding phylogenetic signals. I used the
Tinamidae phylogenetic tree inferred with both molecular (1143 bp mitochondrial and 1145 bp
nuclear) and morphological characters (237 characters) (Bertelli and Porzecanski 2004), in which
the polytomy was resolved with additional morphological and life history traits (Bertelli 2017). I
first tested if there is phylogenetic signal (Pagel 1999) in each of the egg color variable with
phytools (Revell 2012). If no significant phylogenetic signal was identified, I ran permutation
test of independence between song PC1 and the egg color variable with *coin* package. If there is
significant ($p < 0.05$) phylogenetic signal, I first calculated independent contrasts of egg colors
and song PC1 with Ape (Paradis and Schliep 2019), then executed independence test. To correct
for multiple hypotheses testing, I conducted False Discovery Rate correction (Benjamini and
Hochberg 1995) to correct for three tests.
94
95                                                **Results**
96      I found a significant association between tinamou songs and egg luminance. Song PC1
97   effectively represents variation among the song variables in tinamous (Figure 2A). Significant
98   phylogenetic signal was observed in egg hue (K = 0.43, $p = 0.049$) and chroma (K= 0.96, $p =$
99   0.001), but not in luminance (K = 0.37, $p = 0.17$). The tinamou song PC1 was significantly

associated with egg luminance ($Z = -2.42$, $p = 0.015$, $p_{corrected} = 0.045$; Figure 2B), but not with egg hue ($Z = 0.34$, $p = 0.72$, $p_{corrected} = 0.75$), nor with egg chroma ($Z = -0.32$, $p = 0.75$, $p_{corrected} = 0.75$).

**Figure 1 Phylogeny of tinamous and variability in egg colour** The Tinamidae phylogeny was inferred with molecular and morphological data (Bertelli and Porzecanski 2004).

**Figure 2 Principal components analysis of tinamou song characteristics and their association with egg colour** (A) a biplot of the song PCA with blue vectors representing projected song variables (labeled in red). (B) There is significant association between egg luminance and song PC1 (permutation test of independence, $Z = -2.42$, $R^2 = 0.20$, $p = 0.015$, $p_{corrected} = 0.045$). The color of dots represent the RGB color of the eggs of each species.

116
## Discussion
117
The coevolution of tinamou song and egg coloration supports the hypothesis that tinamou
egg coloration is an alternative mating signal (Weeks 1973; Brennan 2010). Egg colors and song
could be divergently selected as multimodal mate-recognition signals among tinamou species
with similar appearance. When the plumage modality is constrained by anti-predation adaptation,
egg coloration could be opportunistically adopted to fulfill mating signal multimodality. This
study sheds light on the evolution of multimodal sexual signals that bypasses natural selection
for plumage camouflage in the most species-rich order of Palaeognathae.
Egg coloration could be both a pre- and post-mating signals in tinamous. In many other
bird species, egg color is known to be post-mating signals indicating female quality and to
influence male incubation and promiscuity (Soler et al. 2005; English 2009). In tinamous, egg
coloration can also function as premating signal as well, received by females. For most of the
tinamou species, males guard and incubate the eggs laid by multiple females (Cabot 1992). The
coloration of the existing eggs in the nest could be a mating signal received by the upcoming
females to the nest. As premating signals, egg colors could reflect mate species identity (Sætre et
al. 1997; Servedio and Noor 2003; Secondi et al. 2015) and mate quality, stimulating female
mate choice copying (Dugatkin 1992; Gibson and Höglund 1992).
Then why do tinamous adopt egg colors as mating signals in addition to songs? Songs
that are important mating signals in tinamous (Bertelli and Tubaro 2002) and are involved in
duetting between mating partners (Boesman et al. 2018). However, such acoustic modality is
usually insufficient for mate communication in complex environment, thus signal redundancy or
multimodality is further selected for mate communication to alleviate mate-searching effort
and/or hybridization (Partan and Marler 1999; Rowe 1999; Uy et al. 2008; Secondi et al. 2015).
Plumage patterning and coloration are frequently adopted as inter-specific and intra-specific
mating signals at a finer spatial scale in birds (Uy et al. 2008, 2009; Seddon et al. 2013).
However, most tinamou species exhibited cryptic plumage as an adaption for anti-predatory
camouflage (Cabot 1992; Davies 2002). Sexual selection for plumage elaboration would
compromise the adaptation for camouflage favored by natural selection. Egg coloration can be an
alternative modality of mate signaling enrichment towards mate-searching refinement, bypassing
the conflict between natural and sexual selection.
Reinforcement may have driven tinamou egg color and song divergence among
sympatric or parapatric tinamou species. Many closely-related tinamou species demonstrate
historical and/or contemporary range overlap (Cabot 1992), suggesting potential sympatry in the
course of their speciation history. Ecological and/or intrinsic incompatibility among diverged
lineages leads to reduced hybrid fitness, which in turn select for premating signal divergence to
avoid costly hybridization (Dobzhansky 1940; Mayr 1942; Liou and Price 1994; Servedio 2000).
Mating signal multimodality is needed to ensure mate recognition in complex heterospecific
environment (Uy et al. 2008; Secondi et al. 2015). The divergence of tinamou song and egg
coloration could jointly reduce heterospecific reproductive efforts among sympatric tinamou
species. Closely related tinamou species that are currently in allopatry could still harbor
footprints of historical character displacement of egg colors formed at historical sympatry. For
example, the three closely related species *Crypturellus erythropus*, *C. notivagus*, *C. variegatus*
demonstrate distinct egg coloration, which may have been driven by reinforcement at historical
sympatry. The historical geography and ecology around the speciation event is most relevant to
recreate the potential egg color displacement. However, it is challenging to retrieve such

information, because the splits among these three tinamou species have long passed (~34%
sequence divergence).
The genetic underpinning of such song and egg color association is unclear. The simplest
genetic mechanism underlying such association is pleiotropy (one gene affecting multiple traits)
(Fisher 1930; Williams 1957). For example, the pleiotropic *foraging* gene underpins the
association among social behavior and lifehistory traits in natural populations (de Belle et al.
1989; Mery et al. 2007; Wang and Sokolowski 2017). There might be an omnipotent pleiotropic
gene linking egg coloration, songs, among many other traits. If so, a wide array of traits
underpinned by such pleiotropy are expected to coevolve. However, the association between
song and egg color of tinamous is fairly specific: only luminance (but not hue or chroma) of the
egg color is associated with tinamou songs. Such specificity in song-egg-color association is
consistent with mating signal multimodality in which specifically signals were coupled among
modalities (Gilliard 1956; Partan and Marler 1999; Hebets and Papaj 2005). Various signal
modality can be underpinned by different genes and selected to coevolve as multimodal signals.
Future investigation of genetic underpinnings of tinamou song and egg colors would further shed
light on this interesting co-divergence. In sum, the results presented herein are concordant with
the hypothesis that egg coloration in tinamous serves as mating signals either for mate attraction
or mate species-recognition.
180
181

**Acknowledgement**
Thank Dahong Chen, Patricia Brennan, Daniel Hanley, Julia Clarke, and Jonathan
Rolland for helpful discussions. Thank Ken Thompson, Dahong Chen, and Jonathan Rolland for
comments on the manuscript.

**Reference**
Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B 57:289–300.
Bertelli, S. 2017. Advances on tinamou phylogeny: an assembled cladistic study of the volant
palaeognathous birds. Cladistics 33:351–374.
Bertelli, S., and A. L. Porzecanski. 2004. Tinamou (tinamidae) systematics: a preliminary
combined analysis of morphology and molecules. Ornitol. Neotrop. 15:293–299.
Bertelli, S., and P. L. Tubaro. 2002. Body mass and habitat correlates of song structure in a
primitive group of birds. Biol. J. Linn. Soc. 77:423–430.
Boesman, P., O. Claessens, T. V. V. Costa, V. Pelletier, J. Ingels, and A. Renaudier. 2018. Songs
of Rusty Tinamou Crypturellus brevirostris and duetting in Crypturellus species. Bull. Br.
Ornithol. Club 138:69–78.
Brennan, P. L. R. 2010. Clutch predation in great tinamous Tinamus major and implications for
the evolution of egg color. J. Avian Biol. 41:419–426.
Cabot, J. 1992. Tinamiformes. Pp. 112–144 *in* D. Hoyo, A. Elliot, and J. Sargatal, eds.
Handbook of birds of the world. Barcelona.
Davies, S. J. J. F. 2002. Ratities and tinamous. Tinamidae, Rheidae, Dromaiidae, Casuariidae,
Apterygidae, Struthionidae. Oxford Univ. Press, New York.
de Belle, J. S., A. J. Hilliker, and M. B. Sokolowski. 1989. Genetic localization of foraging (for):
a major gene for larval behavior in Drosophila melanogaster. Genetics 123:157–164.
Dobzhansky, T. 1937. Genetics and the Origin of Species. Columbia University Press, New

York.

Dobzhansky, T. 1940. Speciation as a Stage in Evolutionary Divergence. Am. Nat. 74:312–321.

Dugatkin, L. A. 1992. Sexual Selection and Imitation: Females Copy the Mate Choice of Others.
Am. Nat. 139:1384–1389.

English, P. 2009. Why is a robin's egg blue? Exploring the evolution of egg colour in birds.
Queen's University.

Fisher, R. A. 1930. The genetical theory of natural selection. Oxford Univ. Press, New York,
NY.

Gibson, R., and J. Höglund. 1992. Copying sexual selection. Trends Ecol. Evol. 7:229–232.

Gilliard, E. T. 1956. Bower Ornamentation versus Plumage Characters in Bower-Birds. Auk
73:450–451.

Halfwerk, W., J. Varkevisser, R. Simon, E. Mendoza, C. Scharff, and K. Riebel. 2019. Toward
testing for multimodal perception of mating signals. Front. Ecol. Evol. 7:1–7.

Hebets, E. A., and D. R. Papaj. 2005. Complex signal function: Developing a framework of
testable hypotheses.

Hothorn, T., K. Hornik, M. A. van De Wiel, and A. Zeileis. 2008. Implementing a Class of
Permutation Tests COIN Package. J. Stat. Softw. 28:1–23.

Koski, M. H., and T. L. Ashman. 2016. Reproductive character displacement and environmental
filtering shape floral variation between sympatric sister taxa. Evolution (N. Y). 70:2616–
2622.

Laverde-R., O., and C. D. Cadena. 2014. Taxonomy and conservation: A tale of two tinamou
species groups (Tinamidae, Crypturellus). J. Avian Biol. 45:484–492.

Liou, L. W., and T. D. Price. 1994. Speciation by Reinforcement of Premating Isolation.
Evolution (N. Y). 48:1451–1459.

Mayr, E. 1942. Systematics and the origin of species. Columbia University Press, New York.

Mery, F., A. T. Belay, A. K. C. So, M. B. Sokolowski, and T. J. Kawecki. 2007. Natural
polymorphism affecting learning and memory in Drosophila. Proc. Natl. Acad. Sci. U. S. A.
104:13051–13055.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–884.

Paradis, E., and K. Schliep. 2019. Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics 35:526–528.

Partan, S., and P. Marler. 1999. Communication goes multimodal. Science (80-. ). 283:1272–
1273.

Pfennig, K. S. 2016. Reinforcement as an initiator of population divergence and speciation. Curr.
Zool. 62:145–154.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other
things). Methods Ecol. Evol. 3:217–223.

Rowe, C. 1999. Receiver psychology and the evolution of multicomponent signals. Anim.
Behav., doi: 10.1006/anbe.1999.1242.

Sætre, G. P., T. Moum, S. Bureš, M. Král, M. Adamjan, and J. Moreno. 1997. A sexually
selected character displacement in flycatchers reinforces premating isolation. Nature
387:589–592.

Schläpfer, K. 2017. The colorful eggs of the tinamous.

Secondi, J., G. Rodgers, F. Bayle, S. Sourice, and M. Théry. 2015. Mate preference, species
recognition and multimodal communication in heterogeneous environments. Evol. Ecol.
29:217–227.

Seddon, N., C. A. Botero, J. A. Tobias, P. O. Dunn, H. E. A. MacGregor, D. R. Rubenstein, J. A.
C. Uy, J. T. Weir, L. A. Whittingham, and R. J. Safran. 2013. Sexual selection accelerates
signal evolution during speciation in birds. Proc. R. Soc. B Biol. Sci. 280:20131065.
Servedio, M. R. 2000. Reinforcement and the genetics of nonrandom mating. Evolution (N. Y).
54:21–29.
Servedio, M. R., and M. A. Noor. 2003. The role of reinforcement in speciation: Theory and
data. Annu. Rev. Ecol. Syst. 34:339–364.
Skutch, A. F. 1966. A breeding bird census and nesting success in Central America. Ibis (Lond.
1859). 108:1–16.
Soler, J. J., J. Moreno, J. M. Avilés, and A. P. Møller. 2005. Blue and green egg-color intensity
is associated with parental effort and mating system in passerines: support for the sexual
selection hypothesis. Evolution (N. Y). 59:636–644.
Swynnerton, C. F. M. 1916. On the Coloration of the Mouths and Eggs of Birds: On the
Coloration of Eggs. Ibis (Lond. 1859). 58:529–606.
Uy, J. A. C., R. G. Moyle, and C. E. Filardi. 2008. Plumage and song differences mediate species
recognition between incipient flycatcher species of the Solomon Islands. Evolution (N. Y).
63:153–164.
Uy, J. A. C., R. G. Moyle, C. E. Filardi, and Z. A. Cheviron. 2009. Difference in Plumage Color
Used in Species Recognition between Incipient Species Is Linked to a Single Amino Acid
Substitution in the Melanocortin-1 Receptor. Am. Nat. 174:244–254.
Wang, S., and M. B. Sokolowski. 2017. Aggressive behaviours, food deprivation and the
foraging gene. R. Soc. open Sci. 4:170042.
Weeks, S. . 1973. The behavior of the red-winged tinamou (Rhynchotus rufescens). Zoologica
58:13–40.
Williams, G. C. 1957. Pleiotropy, Natural Selection, and the Evolution of Senescence. Evolution
(N. Y). 11:398–411.
280

281  **Supplementary information**

282
283 **Figure S1 Scatterplots showing correlations among the four song variables**.