Title

Flood-induced interspecific interactions in spring-fed tributary as an ecosystem function

of heterogeneous river networks

Authors

Masaru Sakai[1] · Ryoshiro Wakiya[2] · Gosuke Hoshi[3]

Affiliations

[1] Fukushima Regional Collaborative Research Center, National Institute for

Environmental Studies, Japan, 10-2 Fukasaku, Miharu, Tamura District, Fukushima

963-7700 Japan

[2] Atmosphere and Ocean Research Institute, the University of Tokyo, 5-1-5

Kashiwanoha, Kashiwa, Chiba 277-8564 Japan

[3] Graduate School of Science and Engineering, Chuo University, 1-13-27 Kasuga,

Bunkyo-ku, Tokyo 112-8551 Japan

✉ Masaru Sakai

sakai.masaru@nies.go.jp

ORCID iD: 0000-0001-5361-0978

Abstract

Understanding the migratory community dynamics of river networks is important for

maintaining lotic system integrity. River animals migrate to their preferred habitats in

spatiotemporally heterogeneous river environments. Spring-fed habitats are uniquely

characterized by stable temperature and flow regimes, which create suitable spawning

habitats for the chum salmon *Oncorhynchus keta*. *O. keta* exhibits "run up" to its

birthplace for spawning, especially during floods. Because the eggs deposited by this

anadromous fish are nutritious and actively consumed by freshwater animals, the

location and timing of *O. keta* spawning events affect the spatiotemporal accumulation

of mobile consumers. In this study, we examined changes in temporal population

density in spawning *O. keta* and a mobile consumer (juvenile *O. masou masou*) in a

lowland, spring-fed tributary in northern Japan during a 48.5-mm autumn rainfall event.

In both species, population density increased, and then decreased, after the rainfall event.

In *O. keta*, these changes were closely associated with rainfall intensity, whereas in *O.*

masou masou the peak was delayed until 3 days after the rainfall event. A comparison

of the gut contents of *O. masou masou* sampled from a spring-fed tributary and an

adjacent non-spring-fed tributary indicated greater consumption of *O. keta* eggs in the

spring-fed tributary. These results suggested that preferential migration of *O. keta* into

spring-fed tributaries for spawning induces subsequent accumulation of juvenile *O.*

masou masou, in turn increasing *O. keta* egg consumption. These findings improve our

understanding of community dynamics during floods in a heterogeneous river network

environment.

Key words: Animal migration, Field ecology, Resource subsidy, Salmonids, Upwelling

groundwater

Introduction

Spring-fed streams, which are maintained by continuous groundwater upwelling, constitute unique ecosystems with stable temperature and flow regimes (Mattson et al. 1995; Sear et al. 1999; Lusardi et al. 2016). The narrow range of water temperature in spring habitats creates refugia for stenothermal glacial relict species (Reiss et al. 2016; Sun et al. 2020). Stable flow regimes in spring-fed streams are associated with less sediment runoff during floods than non-spring-fed streams, and the formation of fine substrates inhabited by abundant detritivorous macroinvertebrates (Sakai et al. 2020). Thus, the unique habitats of spring-fed streams enhance beta diversity in river networks (Reiss et al. 2016; Sakai et al. 2020).

The chum salmon *Oncorhynchus keta*, an important North Pacific fishery species, is an anadromous fish. After 2–4 years of growth in the ocean, *O. keta* exhibits "run up" to its birthplace for spawning from summer to winter (Masuda et al. 1984), especially during floods caused by rainfall events (Banks 1969). *O. keta* has preferentially selected upwelling water habitats with a stable warm temperature for spawning (Kobayashi 1968; Milligan et al. 1984; Geist et al. 2011); these upwelling springs form egg accumulation zones after rainfall events during the spawning season.

O. keta eggs are a significant and nutritious food resource for freshwater fishes. For example, juvenile salmonids rely heavily on eggs deposited by anadromous spawning salmon, including *O. keta* (Armstrong et al. 2013; Koshino et al. 2013; Bailey et al. 2019). This marine-derived resource subsidy is available both after floods in spring habitats, and spatiotemporal heterogeneity in its availability influences the migration of mobile consumers in rivers containing spawning salmon species (e.g., Armstrong et al. 2013).

Understanding migratory community dynamics in river networks is important

to preserve lotic system integrity (Ward et al. 1999; Elosegi et al. 2010); however, direct

observation of the migration of lotic animals, particularly fishes, during floods is

generally difficult and unsafe. In contrast, spring-fed streams are suitable for

investigating community dynamics during flood events due to their high stability.

Autumn rainfall can trigger upstream migration of *O. keta* for spawning, which

influences the foraging habitats of species that feed on their eggs. This interactional

process may represent an additional ecosystem function of spring-fed streams, i.e.,

structuring biota in river networks. However, substantial gaps in understanding remain

regarding the functional role of spring-fed streams in the dynamics of stream fishes

during floods.

Because springs generally provide purified water, spring-fed streams are

frequently reserved for use as a human water supply (LaMoreaux and Tanner 2001).

Elucidating the ecosystem functions of spring-fed streams would further highlight their

value. In the present study, we investigated temporal variation in stream fish dynamics

in a spring-fed tributary after a rainfall event. Although access to the surrounding

non-spring-fed streams was not feasible during the rainfall event, the spring-fed

tributary was relatively calm and could be safely accessed for surveys. We examined the

responses of spawning *O. keta* and juvenile *O. masou masou* to the flood event, and

associated interspecific interactions related to the consumption of *O. keta* eggs. The

findings of this study contribute to our understanding of community dynamics during

floods in heterogeneous river network environments.

Materials and methods

Study site

This study was conducted in the Shubuto River basin in Kuromatsunai,

Hokkaido, Japan (42.64° N, 140.34° E), which encompasses an area of 367 km^2 and

includes both montane and lowland regions. The region is underlain by sandstone and

mudstone, including Cenozoic fossil shells and tuff. The mean annual precipitation

between 2009 and 2018 was 1,615.8 mm, and the mean air temperature was 7.5°C,

measured at the Kuromatsunai AMeDAS automated weather station located 4 km

northwest of the study site. The dominant tree species in riparian zones within the study

area are *Salix* species and *Quercus crispula*, and the dominant understory plants are

Sasa kurilensis and *Reynoutria sachalinensis*. No dams or weirs prevent the migration

of fishes in the mainstem of the Shubuto River (Miyazaki and Terui 2016). The Shubuto

River system has a perennial spring-fed tributary with significant groundwater discharge

(ca. 0.1 m^3/s) in the lowland region (Kamiyama River, Fig. 1). This stream has no

artificial structures that inhibit longitudinal connections, and stream fishes thus migrate

between the spring-fed tributary and mainstem.

At this spring-fed tributary, we examined temporal changes in the population

density of two *Oncorhynchus* fishes (*O. keta* and *O. masou masou*) during a rainfall

event. On October 6–7, 2018, a temperate low-pressure system (formerly the Kong-rey

typhoon) released a total of 48.5 mm of rain at the study site. According to our visual

surveys performed immediately after the rainfall event, water levels increased

substantially in most non-spring-fed streams of the Shubuto River system; however, the

study tributary showed no dramatic increase in water level. We collected fish both

before (October 6) and after (October 7, 9, and 10) the rainfall event; hereafter, we refer

to these collection dates as "before," "after d1," "after d3," and "after d4", respectively.

We also collected the gut contents of *O. masou masou* in an adjacent perennial

non-spring-fed tributary (Utasai River, Fig. 1), for comparison between spring-fed and

non-spring-fed tributaries after the rainfall event. Stream width and water depth were

similar between the tributaries (Table 1); however, the non-spring-fed tributary had a

gravel-dominated streambed with a faster current. *O. keta* rarely migrates to the

non-spring-fed tributary during spawning seasons, but migrates yearly to the spring-fed

tributary; a preference for spring-fed habitats as spawning substrates in this species has

been reported previously (e.g., Kobayashi 1968; Milligan et al. 1986), leading to the

construction of a facility for capturing mature *O. keta* in this tributary. The facility

irreversibly traps mature *O. keta* in an artificial pool, and trapped individuals are

transported to another tributary of the Shubuto River system for egg collection and fry

rearing. Thus, eggs found in the spring-fed tributary are presumed to have spawned

within the tributary, and not drifted from the facility. On October 10 (after d4), when

both tributaries had ordinary flow conditions, we sampled *O. masou masou* gut

contents.

Hydrological monitoring

Water level and temperature loggers (HOBO CO-U20L-04; Onset, Bourne,

MA, USA) were fixed on streambeds of the tributaries using metal tubes and wire, and

hydraulic pressure was recorded once per hour. Before the rainfall event started, we

preliminarily recorded water levels at the monitoring points on October 6–7, 2018 using

a folding scale, and devised linear regression models of the relationship between water

level (cm) and hydraulic pressure (kPa). Based on these models, hydraulic pressure data

recorded in the loggers were transformed into water level data. For both study reaches,

the water level at 1:00 A.M. on October 6, 2018 was set at 0 cm, and hydrographs of the

ensuing water level changes were created. Hourly rainfall data were obtained from the

Kuromatsunai AMeDAS automated weather station.

Fish collection

A 40-m-long study reach was established in the spring-fed tributary to

investigate temporal changes in *O. keta* and *O. masou masou* population density during

a rainfall event. The Shubuto River *O. masou masou* population includes both

anadromous and resident individuals; most females and some males become

anadromous after approximately 2 years of growth in the river (Miyazaki 2017). Thus,

small individuals collected in the study area were mainly juveniles and resident males.

Because the rainfall event occurred during the reproductive period, we classified small

O. masou masou individuals (standard body length = 60–160 mm) as mature or juvenile

based on the presence or absence of nuptial coloration. *O. keta* is anadromous, and

generally runs up to the Shubuto River from mid-September, where it spawns until

mid-November after 2–8 years of growth in the North Pacific (Miyazaki 2017).

Therefore, individuals found in the river during the autumn reproduction season are

mature.

We conducted electrofishing using a backpack electrofisher (LR-20;

Smith-Root Inc., Vancouver, WA, USA). The upper and lower limits of the tributary

study reach were gently partitioned using fishnets to prevent fishes from entering or

exiting the reach during collection. Three-pass electrofishing was conducted from lower

to upper positions, to collect as many *Oncorhynchus* fish as possible. After collection,

the wetted width was measured at five points (10-m intervals) within the study reach, to

better represent the surveyed area during each collection (before, after d1, d3, and d4)

allow for accurate calculation of fish population density.

The population density of *O. masou masou* was estimated using the maximum

likelihood method proposed by Zippin (1956), because the number of collected

individuals decreased substantially between the first and third removals. The number of

collected *O. keta* individuals varied and exhibited no clear pattern. However, we

confirmed that each three-pass removal collected all individuals within the reach; this

was possible due to their large body size (standard length = 425–801 mm). Therefore,

we used the total number of collected individuals to directly estimate population density

for this species. The maximum likelihood method was applied using R software (v.

3.6.3; R Core Team 2020) with the *FSA* package (Ogle et al. 2020).

Gut content collection

In the early morning and evening of after d4, we collected *O. masou masou* for

gut content analysis using a stomach pump; a total of 31 and 11 individuals were

collected in the spring-fed and non-spring-fed tributaries, respectively. Each gut content

sample was immediately preserved in 70% ethanol. The body size distribution of the

collected *O. masou masou* was similar between tributaries. Fish were released into their

original habitats after gut content collection. To avoid interference with our population

density estimates in the spring-fed tributary, we performed population density sampling

in this tributary in the early morning of after d4, prior to gut content collection.

Gut contents were sorted and identified using a stereomicroscope (SZ61;

Olympus, Tokyo, Japan). Identified gut contents were dried in an oven (FC-610;

Advantec, Tokyo, Japan) at 60°C for 24 h and then weighted precisely to within 0.01

mg. To test for differences in the gut contents of *O. masou masou* between tributaries,

we analyzed the dry weight data using one-way analysis of variance (ANOVA) with

permutation tests, which allowed us to deal with unbalanced sample sizes. ANOVA was

performed using the R 3.6.3 software with the *anova.1way* function (Legendre 2007).

Results

196   Following the onset of the rainfall event, the water level in the spring-fed

tributary declined slightly, but was clearly lower overall than that in the non-spring-fed

tributary throughout the monitoring period (Fig. 2). Water levels in the non-spring-fed

tributary reflected changes in rainfall intensity, and peaked at 29 cm above the initial

water level (Fig. 2). A total of 175 individuals of *O. masou masou* and 35 individuals of

O. keta were collected in the spring-fed tributary. In both species, population density

increased after the rainfall event, but subsequently decreased (Fig. 3); that of *O. masou*

masou peaked at after d3, whereas for *O. keta* it decreased after d1. All collected *O.*

masou masou were juveniles, and all *O. keta* individuals were mature and reproductive.

205   At after d4, *O. masou masou* in the spring-fed tributary consumed significantly

206   more *O. keta* eggs, whereas those in the non-spring-fed tributary consumed more

207   hemipteran, hymenopteran, and trichopteran insects (Table 2). Due to the large

208   contribution of *O. keta* eggs, the total gut contents were higher in the spring-fed

209   tributary than in the non-spring-fed tributary (Table 2).

211   **Discussion**

212   The results showed that increase in population density of *O. keta* immediately

after the rainfall event, and the subsequent peak of population density of juvenile *O.*

masou masou was observed three days after the rainfall event in the spring-fed tributary.

The individuals of *O. masou masou* in the spring-fed tributary consumed more eggs of

O. keta compared to those in the adjacent non-spring-fed tributary. These results suggest

that preferential migration of *O. keta* into spring-fed tributaries for spawning induces

subsequent accumulation of juvenile *O. masou masou*, in turn increasing *O. keta* egg

consumption.

Typically, lowland spring-fed streams formed by groundwater discharge are

not deep (Sear et al. 1999), and their flow regimes are generally stable (Lusardi et al.

2016). In this study, water levels in the spring-fed tributary did not exhibit the same

marked increase observed in the non-spring-fed tributary during the rainfall event. This

stability creates a valuable opportunity for direct field observation of the dynamics of

aquatic animal communities during floods.

Hydrological monitoring indicated that the water level of the spring-fed

tributary decreased following the onset of rainfall. This change may be attributable to

the onset of water transmission into the *O. keta* egg collection facility, because local

fishermen know that this species actively migrates to upstream spawning habitats during

floods (Banks 1969). Our results also indicated that the population density of *O. keta*

peaked immediately after the rainfall intensity peak. The dependence of *O. keta* on

spring-fed habitats for spawning (Kobayashi 1968; Milligan et al. 1984; Geist et al.

2011) may explain the accumulation of this species in the spring-fed tributary during the

flood.

Although the population density of *O. keta* was sensitive to temporal changes

in rainfall intensity, the peak population density of *O. masou masou* was delayed until 3

days after the flood. We presume that this temporal increase in *O. masou masou*

population density was influenced by two factors: evacuation to flow refugia and

acquisition of *O. keta* eggs as food. Tributaries have been considered as flow refugia for

fishes (Koizumi et al. 2013); thus, some juvenile *O. masou masou* may migrate to

spring-fed tributaries to avoid floods. However, the study reach is located 500 m

upstream of the confluence of the tributary and the mainstem; this distance may be too

long for such evacuations. Makiguchi et al. (2009) reported that a related species, *O.*

masou formosanus, endured a massive typhoon event, which induced a 3-m water level

increase, by remaining within their original habitats in the mainstem. Therefore, the

moderate flood observed in this study may not have been a major driver of fish

evacuation to the spring-fed tributary.

The eggs of anadromous salmon species are a significant and nutritious food

resource for freshwater fishes (Armstrong et al. 2013; Koshino et al. 2013; Bailey et al.

2019). This food resource is likely targeted by juvenile *O. masou masou* accumulating

in the spring-fed tributary. Our gut content analysis indicated that there were

significantly more salmon eggs in the spring-fed tributary than in the adjacent

non-spring-fed tributary, where *O. keta* rarely migrates. The time lag between the

population density peaks of *O. keta* and *O. masou masou* may be attributable to

differences in their swimming abilities and/or the detection of egg production cues by *O.*

masou masou, perhaps related to the timing of *O. keta* sperm release and downward

dispersion. Because the present study lacks information on the redd distribution of *O.*

keta in the study streams before and after the rainfall event, whether the consumed eggs

by *O. masou masou* were spawned after the rainfall event or not was unknown.

Grasping the redd distribution can help explain the plausible process proposed here

more clearly.

Both spring-fed streams and upwelling water habitats function as spawning

habitats for *O. keta* (Kobayashi 1968; Milligan et al. 1984; Geist et al. 2011);

accumulation of *O. keta* in such habitats can occur during floods (Banks 1969). In this

study, we directly observed upward migration of *O. keta* into the spring-fed tributary

due to its stable flow regime. Our results indicate that the ecosystem functions of

spring-fed streams, with respect to the formation of a unique habitat with stable

temperature, provide mature *O. keta* with a suitable habitat and, in turn, provide other

species with marine-derived food resources. Interspecific interactions with juvenile *O.*

masou masou also occur in such habitats in spring. Thus, it is crucial to evaluate

community dynamics in river networks in spatiotemporally heterogeneous environments

(Armstrong et al. 2013). A robust sampling design with replications for tributaries and

rainfall events, lacked in this study, can unravel the ecosystem function of

heterogeneous river networks more clearly. The findings of the present study highlight

the importance of enhancing spatiotemporal heterogeneity in spring-fed habitats to

maintain interspecific interactions, which have implications for freshwater biodiversity

and resource management in river networks.

Acknowledgments: A portion of this study was supported by JSPS KAKENHI Grant

Numbers 26292181 and 19K20491, and Kuromatsunai Biodiversity Conservation

Research Grant (2017). Dr. Izumi Washitani provided invaluable comments on earlier

drafts of the manuscript. We thank the field assistance by Dr. Kosei Takahashi, Mr.

Hitoshi Saito, Kengo Ebihara and Katsuya Iwabuchi. The all fish investigations were

conducted with the permission of Hokkaido Prefecture. The authors have no competing

interests.

References

Armstrong JB, Schindler DE, Ruff CP, Brooks GT, Bentley KE, Torgersen CE (2013)

Diel horizontal migration in streams: Juvenile fish exploit spatial heterogeneity in

thermal and trophic resources. Ecology 94:2066-2075

Bailey CJ, Andersson LC, Arbeider M, Bradford K, Moore JW (2019) Salmon egg

subsidies and interference competition among stream fishes. Environ Biol Fish

102:915-926

Banks JW (1969) A review of the literature on the upstream migration of adult

Salmonids. J Fish Biol 1:85-136

Elosegi A, Díez J, Mutz M (2010) Effects of hydromorphological integrity on

biodiversity and functioning of river ecosystems. Hydrobiologia 657:199-215

Geist DR, Hanrahan TP, Arntzen EV, Mc Michael GA, Murray CJ, Chien YJ (2002)

Physicochemical characteristics of the hyporheic zone affect red site selection by

chum salmon and fall chinook salmon in the Columbia River. N Am J Fish

Manage 22:1077-1085

Kobayashi T (1968) Some observation on the natural spawning ground of chum and

pink salmon in Hokkaido. Scientific Reports of the Hokkaido Salmon Hatchery

22:7-13 (in Japanese with English abstract)

Koizumi I, Kanazawa Y, Tanaka Y (2013) The fishermen were right: experimental

evidence for tributary refuge hypothesis during floods. Zool Sci 30:375-379

Koshino Y, Kudo H, Kaeriyama M (2013) Stable isotope evidence indicates the

incorporation into Japanese catchments of marine-derived nutrients transported by

spawning Pacific Salmon. Freshw Biol 58:1864-1877

LaMoreaux PE, Tanner JT (2001) Springs and Bottled Waters of the World.

Springer-Verlag, Berlin, Germany

Legendre P (2007) One-way anova with permutation test.

http://adn.biol.umontreal.ca/~numericalecology/Rcode/. Accessed 5 February 2021

Lusardi RA, Bogan MT, Moyle PB, Dahlgren RA (2016) Environment shapes

invertebrate assemblage structure differences between volcanic spring-fed and

runoff rivers in northern California. Freshw Sci 35:1010-1022

Makiguchi Y, Liao LY, Konno Y, Nii H, Nakano K, Gwo JC, Onozato H, Huang YS,

Ueda H (2009) Site fidelity of and habitat use by the Formosan landlocked salmon

(*Oncorhynchus masou formosanus*) during typhoon season in Chichiawan stream,

Taiwan as assessed by nano-tag radio telemetry. Zool Stud 48:460-467

Masuda H, Amaoka K, Araga C, Uyeno T, Yoshino T (1984) The Fishes of the

Japanese Archipelago. Tokai University Press, Tokyo, Japan.

Mattson RA, Epler JH, Hein MK (1995) Description of benthic communities in Karst,

spring-fed streams of northern central Florida. J Kansas Entomol Soc 68:18-41

Milligan PA, Rublee WO, Cornett DD, Johnston RAC (1986) The distribution and

abundance of chum salmon (*Oncorhynchus keta*) in the upper Yukon River basin

as determined by a radio-tagging and spaghetti tagging program: 1982-1983.

Canadian Technical Report of Fisheries and Aquatic Sciences 1351

Miyazaki Y (2017) Fishes of Shubuto River System, Second Edition. Biodiversity

Council of Shiribeshi Region, Hokkaido, Japan. (in Japanese)

Miyazaki Y, Terui A (2016) Difference in habitat use between the two related goby

species of *Gymnogobius opperiens* and *Gymnogobius urotaenia*: a case study in

the Shubuto River System, Hokkaido, Japan. Ichthyol Res 63:317-323

Ogle D, Wheeler P, Dinno A (2020) Package 'FSA'.

https://cran.r-project.org/web/packages/FSA/. Accessed 5 February 2021

R Core Team (2021) R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Accessed 5 February 2021

Reiss M, Martin P, Gerecke R, von Fumetti S (2016) Limno-ecological characteristics

and distribution patterns of spring habitats and invertebrates from the Lowlands to

the Alps. Environ Earth Sci 75:1033

Sakai M, Natuhara Y, Fukushima K, Naito R, Miyashita H, Kato M, Gomi T (2013)

Responses of macroinvertebrate communities to 4 years of deer exclusion in first-

and second-order streams. Freshw Sci 32:563-575

Sakai M, Iwabuchi K (2020) Unique habitat and macroinvertebrate assemblage

structures in spring-fed stream: a comparison among clastic lowland tributaries and

mainstreams in northern Japan. bioRxiv https://doi.org/10.1101/2020.05.17.101014

Sear DA, Armitage PD, Dawson FH (1999) Groundwater dominated rivers. Hydrol

Process 13:255-276

Sun Y, Takemon Y, Yamashiki Y (2020) Freshwater spring indicator taxa of benthic

invertebrates. Ecohydrol Hydrobiol 20:622-631

Walton SE, Nunn AD, Probst WN, Bolland JD, Acreman M, Cowx IG (2017) Do fish

go with the flow? The effects of periodic and episodic flow pulses on 0+ fish

biomass in a constrained lowland river. Ecohydrology 10:e1777

Ward JV, Tockner K, Schiemer F (1999) Biodiversity of floodplain river ecosystems:

ecotones and connectivity. Regul Rives: Res Manage 15:125-139

Zippin C (1956) An evaluation of the removal method of estimating animal populations.

Biometrics 12:163-189

Table 1 Mean ± 1 standard deviation of environmental variables in the spring-fed and

non-spring-fed tributaries.

	Spring-fed tributary			Non-spring-fed tributary		
Variable	Mean	±	SD	Mean	±	SD
Stream width (m)	4.07	±	0.25	5.13	±	0.91
Water depth (cm)	15.02	±	3.60	16.64	±	4.99
Current velocity (cm/s)	12.43	±	7.09	23.72	±	10.35
Fine sediment deposition (%)*	80.63	±	9.81	22.50	±	15.92

364   *cover of fine sediment (< 2 mm) on streambed estimated using the method in Sakai et

365   al. (2013).

368   **Table 2** Mean ± 1 standard deviation of dry weight (mg) of gut contents in each order

369   and total in the spring-fed and non-spring-fed tributaries.

	Spring-fed tributary			Non-spring-fed tributary			F	P
	Mean	±	SD	Mean	±	SD	F	P
Amphipoda	0.52	±	1.04	0.00	±	0.00	2.63	0.14
Acari	0.01	±	0.03	0.00	±	0.00	0.98	0.35
Ephemeroptera	0.86	±	2.72	0.68	±	0.68	0.05	0.82
Plecoptera	0.01	±	0.04	0.02	±	0.06	0.49	1.00
Hemiptera	0.00	±	0.00	1.78	±	3.75	7.34	**0.01**
Hymenoptera	0.00	±	0.00	0.56	±	1.04	9.27	**0.02**
Coleoptera	0.00	±	0.00	0.03	±	0.11	0.09	0.25
Trichoptera	0.27	±	1.07	1.35	±	1.03	8.44	**0.01**
Diptera	1.22	±	2.04	0.75	±	0.70	0.54	0.45
Salmoniformes	9.30	±	12.78	0.00	±	0.00	5.73	**0.02**
Total	12.18	±	13.72	5.17	±	4.56	2.72	0.10

Bold characters indicate statistically significant difference between the tributaries,

according to one-way analyses of variances with permutation tests ($P < 0.05$).

Figure captions

Fig. 1 Map and photographs of the study tributaries.

Fig. 2 Hydrographs and water temperature data for spring-fed and non-spring-fed

tributaries during October 6–10, 2018. Circles indicate dates of fish collection for

population density estimates in the spring-fed tributary; double circle indicates the date

of additional fish collection in both tributaries for gut content analysis.

Fig. 3 Temporal changes in the population density of *Oncorhynchus masou masou* and

O. keta in the spring-fed tributary. Gray band indicates the 95% confidence interval for

O. masou masou.

Shubuto River

Kamiyama River
(Spring-fed tributary)

Utasai River
(Non-spring-fed tributary)

Horonai River

Hokkaido

Kuromatsunai

1 km

Fig. 2