Selection for the bacterial capsule in the absence of biotic and abiotic aggressions depends on growth conditions

Amandine Buffet[1], Eduardo P.C. Rocha[1] & Olaya Rendueles*[1]

[1] Microbial Evolutionary Genomics, Institut Pasteur, CNRS, UMR3525, Paris, 75015, France.

* olaya.rendueles-garcia@pasteur.fr

Keywords: capsule serotype/evolution/biotic stress/diversifying selection/

ABSTRACT

Extracellular capsules are the outermost layer of some cells and are prevalent across Bacteria. They protect the cell against both abiotic and biotic stresses such as bacteriophages and the host immune system. Yet, it is unclear if capsules contribute to fitness in the absence of external aggressions, in spite of the cost of production. Here, we enquire if there are conditions favouring the presence of the capsule in *Klebsiella,* where it is a major virulence factor. We allowed independent populations of 19 different strains, representing six different capsule serotypes, to evolve for twenty generations in different growth media. We observed that capsules were maintained in nutrient-poor media but were selected against in nutrient-rich media. Competitions with capsule mutants in nine different strains showed that capsules provide *ca* 20% of fitness advantage in nutrient poor-conditions. This is not caused by differences in maximal growth rate, nor death rates. We showed that most capsulated populations reached higher yields. We further characterized how growth conditions affected the role of the capsule during pathogenensis and whether this is conserved across *Klebsiella* strains. Our data reveals that the environment and the serotype strongly influences major virulence traits such as hypermucoviscosity and biofilm formation. Our data shows that the capsule can be selected for in situations lacking abiotic, but also biotic stresses. This raises the possibility that the forces selecting for the presence of a capsule are different from the ones selecting for rapid serotype variation.

IMPORTANCE

Bacterial capsules are a wide-spread virulence factor that limits efficacy of antimicrobial therapy. Whereas most studies focus on the role of the capsule in pathogenesis, very few have addressed the conditions under which the capsule is primarily selected for. Here, we show that

5   small changes in growth media have a strong impact in the maintenance of the capsule and the fitness advantage they confer. Our results suggest that the forces selecting for the capsule may be different from those selecting for its composition diversification, namely the immune system and phage predation. Our results further support that the role of the capsule in virulence may be a by-product of adaptation, hinting that there may be other functionalities

10   yet to be discovered for it.

INTRODUCTION

15   Most environments are complex and can be temporally or spatially heterogeneous, both in their abiotic (pH, nutrients, chemicals…) or biotic compositions (niche invasions, extinction events). This can represent an evolutionary challenge for microbes (1, 2). In order to survive, bacterial species can either move to less stressful territories, adapt to the new circumstances or resist until conditions are suitable (3). Notably, bacteria can produce a thick extracellular

20   layer – the capsule – that has been reported to enhance survival and tolerate external aggressions (4, 5). Extracellular capsules constitute the first barrier between the cell and its environment and are encoded in half of the bacterial genomes, in some archaea and yeast (6). Capsules fulfill many roles. They have been associated with resistance to abiotic stresses such as UV light and desiccation (4, 5). However, they are mostly studied for their role enhancing

25   survival under biotic stresses, including resistance to grazing protozoa (7), antibiotics (8, 9), antimicrobial peptides (10, 11), and host immune defences, such as human serum and phagocytes (12-14).

The majority of capsules belong to Group I or Wzx/Wzy-dependent capsules (6). These

30   capsules are high molecular weight polysaccharides made up of different repeat units of oligosaccharides. The different possible oligosaccharide combinations and residue modifications lead to different serotypes. Serotype diversity, even within-species, is very

large. This genetic diversity has been particularly studied in facultative pathogens such as *Streptococcus pneumoniae* (15), *Campylobacter jejuni* (16), and *Acinetobacter baumanii*

35   (17). In *K. pneumoniae,* there are at least 77 serologically-defined capsule serotypes (18-20), and more than 130 serotypes identified through comparative genomics (21, 22). This serotype diversity results from the faster evolutionary rates of capsular loci compared to those of the rest of the genome. The latter has been shown to occur by homologous recombination (23, 24) and horizontal gene transfer (25-27). As a result, the same capsular serotype can be found in

40   multiple lineages of *K. pneumoniae* and a monophyletic clade can include distinct serotypes (24, 28). Selection for rapid variation of capsule serotypes is thought to result from biotic stresses, notably because phages and the human adaptive immune system target the capsule in a serotype specific manner, which favours the emergence of novel serotypes (29). This process can be accelerated by vaccines which target specific capsule serotypes (16, 30, 31).

Many studies have assessed the role of the capsule during pathogenesis, yet little is known about the primary selective forces that underlie the emergence and maintenance of the capsule when bacteria are not under biotic stress. Since capsules are costly to produce, if they are selected to protect bacterial cells from phages, immune systems, or specific stresses, they

50   should be readily lost in the absence of these challenges. Consequently, serial passaging of *Klebsiella* cultures in nutrient-rich laboratory media (LB), as well as aging of cultures, readily leads to the emergence of non-capsulated clones (32, 33), indicating that the capsule can be easily lost.

55   To further investigate the conditions where the capsule is advantageous, we study bacteria from the genus Klebsiella. This genus is composed of ubiquitous bacteria able to colonize a large range of environments (34), due to a diverse metabolism (35) and the presence of a Wzx/Wzy capsule (21). To date, most, if not all isolated *Klebsiella*, have a capsule operon. *Klebsiella spp*., and more particularly *K. pneumoniae*, are opportunistic pathogens and major

60   multi-drug resistant (MDR) bacteria that cause both community-acquired and hospital-acquired infections in humans (36-38). The capsule is regarded as the major virulence factor of the genus (39). To test the maintenance of capsule production in several growth conditions, including host-related environments, we used nineteen *Klebsiella* strains representative of the species genetic diversity (35). These strains have different capsule serotypes (K-locus) and O-

65   antigens (O-locus), and include a closely-related *K. variicola* (strain #24), and two *K. quasipneumoniae* subsp. *similpneumoniae* (strains #44 & #214) (Table S1). We then

generated a panel of nine capsule mutants to address the precise fitness effects of the capsule in a range of conditions. Finally, we explored the importance of the serotypes and the genetic background of the strain in driving capsule-related phenotypes. The integration of these

70   results revealed that the capsule can be selected in the absence of biotic stresses.

RESULTS

The capsule is lost in nutrient rich environments

75   Early reports (32, 33) and our laboratory observations reveal that serial passaging of *Klebsiella* cultures in rich laboratory media (LB) generates spontaneous capsule mutants (40). This strongly suggests that capsule has a negative impact on fitness in laboratory environments. To understand the fitness cost of capsule in the absence of biotic pressures, we serially passaged 19 different *Klebsiella* strains belonging to six different serotypes (at least

80   three strains per serotype), amongst which the virulence-associated KL1, KL2 and the emergent KL107 associated to multi-drug resistance. We then measured the natural emergence of non-capsulated mutants after *ca*. 20 generations. We tested the emergence of non-capsulated mutants in five different growth media (see Methods). Artificial sputum medium (ASM) (41) and artificial urine medium (AUM) (42) mimic host-related nutritional

85   conditions. The remaining are typical laboratory media with decreasing levels of nutrients: rich Luria-Bertani medium (LB), minimal medium supplemented with 0.2% of glucose (M02), and minimal medium supplemented with 0.05% of glucose (M005). Beforehand, we verified that the strains were able to produce a capsule in our growth conditions by direct microscopic observation and by capsule quantification using the uronic acid method

90   (Methods, Figure S1). To initiate the experiment, we grew three independent clones of each strain for 16 hours in LB and then diluted (1:100) into each growth media. All 19 strains were then serially transferred into fresh media every 24 hours for three days. Although each growth media has different carrying capacities, cultures were always diluted to 1:100 and after 24h of growth, each media was saturated, ensuring that in all growth media, the different populations

95   underwent the similar number of generations (~ 20).

After three days, in 111 of the 290 independently evolved populations (38%) non-capsulated mutants emerged (Figure 1). Most ASM (74%) and LB (89%) lineages showed capsule mutants, and these accounted for 85 of the 111 populations. In contrast, non-capsulated

100 mutants were very rarely observed in the other three growth media (M02, M005, and AUM. These data suggest that it was the amount of nutrients present in the growth media that drove the maintenance of capsule rather than the opposition between host-related environments versus lab environments. Indeed, it was in the nutrient rich media (ASM and LB), where the capsule was readily lost, whereas it was consistently maintained in the three nutrient poor

105 media (M02, M005 and AUM). To further confirm this, we allowed the experiment to run for 30 days in LB and M02 for a KL2 strain (#26). The three replicate populations that evolved in M02 remained capsulated whereas populations evolving in LB rapidly lost their capsule and remained so until the end (Figure S2). Some strains also seem more prone to lose the capsule. An extreme example concerns strain #62 for which we found capsule mutants in all fifteen

110 independently-evolved populations across the five different growth media. We tested if these differences could be due to the amount of capsule produced by each strain, but the frequency of non-capsulated mutants does not correlate with the amount of capsule produced by each strain (P > 0.05, Spearman's rank correlation). We tested the relevance of both serotype and growth media on the emergence of capsule mutants using a generalized linear model ($R^2 =$

115 0.39). This revealed that the emergence of capsule mutants was both dependent on the growth media (P < 0.001) and on the serotype (P<0.02). This suggests the capsule is deleterious under rich media and selected for in poor media, a factor that depends at a lesser extent on the capsule serotype.

120 ## The capsule provides a fitness advantage in nutrient poor environments.

To precisely assess the costs and benefits of the capsule across growth media, we sought to perform competition experiments between capsulated (Cap+) and their respective isogenic non-capsulated (Cap-) mutants. We thus undertook the construction of in-frame and markerless deletions of *wza*, the outer membrane exporter of the capsule, a well-annotated

125 and essential core gene of the capsule operon. We used the allelic recombination strategy and aimed at deleting *wza* in 15 different strains from different serotypes, for which we had easily obtained *rcsB* in-frame deletion mutants. (RcsB is a capsule regulator and mutants have a reduced capsule production (43, 44). We obtained *wza* deletion mutants for five strains, from several serotypes (see Table S1). From these double-recombination events (see Methods), we

130 isolated both a mutant and a wild type. This ensures that the phenotype observed was directly due to the *wza* deletion and not to other off-target SNPs that could have accumulated in the genome during the mutant generation, and thus present in both the mutant and the wild type.

All experiments were performed in parallel with the parental strain, *Δwza* mutant and the associated wild type from the same double-recombination event. After several attempts, we

135 obtained *wza* deletion mutants in four other strains, but Illumina sequencing revealed off-target compensatory mutations only present in the mutant (and not in the associated wild type) (see Methods). Thus, because we cannot be certain that the phenotypes are exclusively due to the deletion of *wza*, we present general statistics in the text and supplemental figures, but only present results for the five clean mutants in the main figures (see Methods).

140 Microscopic observation and uronic acid quantification confirmed that all nine *Δwza* mutants did not produce a capsule (Figure S3). For all other seven strains, no *Δwza* mutants were obtained despite several attempts.

To assess the competitive indexes, or fitness, of the mutant strains, we performed direct

145 competition between wild type strains and their respective the *Δwza* mutant strains. They were initially mixed in a 1:1 ratio and allowed to compete in all five growth media for a 24-hour period. We estimated that the natural emergence of capsule mutants in rich media is rare during the first 24 hours and is thus negligible in the interpretation of our results (see Methods). Competition experiments showed a significant effect of the environment in the

150 fitness of the Cap + strains (Figure 2, Kruskal-Wallis, $P = 0.0046$). In agreement with our previous results that showed the maintenance of capsule in nutrient poor media, Cap+ strains had a strong fitness advantage in M02, M05 and AUM ($P < 0.05$, $N=8$, one-sided t-test, difference of 1). In this analysis, we excluded strain #55 due to a growth deficiency of the *Δwza* mutant (see Methods). The average fitness advantage provided by the capsule in

155 nutrient poor media is *ca.* 20%. Opposite to all other strains, both deletion mutants of KL24 strains showed increased fitness in AUM, suggesting a particular benefit of the capsule in these two strains in AUM (Figure S5). On the other hand, in nutrient rich media, we observe a significant decrease of Cap+ genotypes ($P= 0.07$ and $P = 0.05$ in LB and ASM, respectively, one-sided t-test, difference of 1). Decreased fitness in these growth media was significant in 5

160 out of 8 strains in LB and 4 out of 8 in ASM strains. More particularly, *K. variicola* is significantly less fit in nutrient rich media (-25%). Similarly, the KL1 strain (#56) is 18% less fit. Two other strains, KL2 and KL124, also have significant disadvantages in LB and ASM, respectively ($P<0.05$, one-sided t-test, difference of 1) (Figure 2 and Figure S5). Taken together, our results confirm that the capsule is advantageous in nutrient-poor media, but not

165 in rich media.

The capsule increases the population yield in nutrient poor growth media

We investigated how the capsule could increase fitness in nutrient poor environments by assessing the effect of mutants in a number of conditions. We first tested whether the capsule provided an advantage during starvation, by increasing cell survival, as shown in *Streptococcus pneumoniae* (45). To check this, we followed the survival of capsule mutants and wild type under starvation conditions (see Methods). We allowed cells to starve for three months and counted surviving CFUs every month. As expected, mortality rates of *Klebsiella* are very low compared to other Enterobacteria (46). Yet, we observed no significant differences between wild type and the capsule mutants after 30 or 60 days (Figure 3A and Figure S5A). To further check a potential effect of the *Klebsiella* capsule on survival, we also challenged the strains to desiccation. No effect of the capsule on cellular survival was observed during desiccation ($P > 0.05$, t test for significant difference between mutants and wild type, Figure S5B).

Given that no significant effect of the capsule on cellular survival was observed, we tested whether differences in growth rate could account for the fitness effects observed. We thus allowed all strains and their mutants to grow in microtiter plates in all growth media and calculated the minimum generation time (Figure 3B and Figure S6A). As expected, we observed significant differences in growth rate across strains and environments (GLM, P< 0.001 for both, $R^2 = 0.60$). However, no significant differences between the mutants and capsulated strains was observed (P = 0.41). Thus, increased fitness in nutrient poor media s is not due to faster growth. Close inspection of the growth curves revealed that strains with capsule in nutrient poor media reached higher absorbances (ODs), a proxy for population yield (Figure 3C and Figure S6B). Control experiments showed a strong correlation between absorbance and cell numbers (see Methods). Furthermore, direct plating of cells after 16 hours of growth in microtiter plates also revealed larger population sizes in capsulated strains than in mutants. Direct comparison of yield between mutants and the respective wild type showed significant difference for several strains in nutrient poor environments. Taken together, there is a statistically significant difference in yield between the mutants and the wild type in nutrient poor media ($P < 0.01$ in M02, $P < 0.001$ in M005 and $P = 0.1$ in AUM, paired t-test, $N = 8$), suggesting that the fitness benefits observed in competition are due to an increased yield or reproductive lifespan.

200   To further understand how capsulated bacteria achieve higher yields in poor media, we first verified that in our experimental conditions, non-capsulated strains did not exhibit higher death rates (Two-way ANOVA, P = 0.47, Figure S6C). We then hypothesized that capsulated bacteria could use their capsule as a nutrient source. To do so, we constructed new capsule

205   mutants (*ΔwcaJ* and *ΔwbaP,* initial glycosyl transferases*),* to interrupt capsule production at the first biosynthesis step and avoid capsule build up that could interfere with our results. We allowed the mutant strains to grow for 48 hours in minimal media with no source of carbon, in the presence of purified capsule added exogenously and of uronic acid used as a standard to quantify capsule production (Figure S7). We did not observe significant differences between

210   the treatments where the capsule was added exogenously and the respective controls (see Methods, see Figure S7). These results suggest that neither increased cellular death of mutants nor the ability to consume the capsule explain the observed higher yield.

Capsule mutants undergo different fates in host-mimicking media

215   Our previous results show that small changes in growth conditions can strongly affect selection for the capsule. We thus sought to further understand how these growth conditions affect the role of the capsule during pathogenesis and whether this is conserved across Klebsiella strains. We first addressed the ability of non-capsulated mutants to resist to human serum. Three Cap- mutants were significantly more sensitive to human serum than their Cap+

220   counterpart, whereas for the other two strains there was no significant difference (Figure 4A). A KL2 strain (#26) was resistant to human serum irrespective of its capsule (Figure 4A). One KL24 strain (#51) (Figure S8A) and *K. variicola* (strain #24) Cap- was as sensitive as the Cap+. The capsules of strain #211 and of strain #37, from the same serotype as strains #24 and #37, respectively, provide a significant degree of protection. These results strongly

225   suggest that the effect of the capsule on resistance to serum is strain-dependent (Multifactorial ANOVA, P < 0.001).

We then explored the ability of capsule mutants to survive in the presence of primary (sodium cholate –CHO-) and secondary (sodium deoxycholate –DCH-) bile salts. At physiological

230   concentrations (0.05%), we observed no significant sensitivity of *Klebsiella* to bile salts (Figure S8B). When treated with 10X physiological concentrations, some capsule mutants showed reduced viability, whereas others were not (Figure S8B).

Finally, the ability of niche colonization, specially within a host, is directly linked to the ability of the strains to adhere surfaces and proliferate, that is, to form a biofilm. Cells within biofilms are more resilient, tolerant to antibiotics and less exposed to the host immune system (47). We thus assessed the ability of the strains to form biofilm and how the capsule could impact this trait. It has been previously shown that in some strains of *Klebsiella*, capsule production masks the necessary fimbriae required to attach to the host mucosa (48, 49). On average, we do not observe a difference in the amount of biofilm that the capsulated and not capsulated mutants form (P > 0.05, $N = 9$, paired t-test, in all different growth media). However, we observe a strong negative correlation between the amount of biofilm the parental strain forms and the difference between the capsule mutant and its associated wild type (Figure 4B). Deletion of the capsule in strains with limited ability to form biofilm results in increased ability to form biofilm. In other words, the capsule tends to contribute more positively to biofilm formation in strains forming more biofilm. At the strain level, we observe that in some cases the non-capsulated mutants adhere significantly more across all environments (strain #56) (Figure S9), whereas other strains have a stronger ability to form biofilm when the capsule is present (strain #47) (Figure S9). This might either result from the absence of adhesion factors in the strain or it could also suggest that depending on the capsule composition, it could act as an adhesion factor. Overall, our data suggest that it is not the presence or absence of the capsule that affects the ability to form biofilm, but rather the amount of capsule expressed by a given strain.

Taken together, these results show that some properties like resistance to bile salts or human serum, are not inherent to the presence of the capsule, but are rather strain-dependent, or dependent on the serotype-genome interaction.

Complex interactions between the environment, genetic background and serotype

To further understand the interactions between the serotype, genetic background and environment, we used the original dataset of 19 strains, in which there are at least three strains belonging to each serotype. We analyzed different capsule-related variables in all five growth media. In addition to the amount of capsule production (Figure S2), the natural emergence of non-capsulated mutants (Figure 1), maximum growth rate, the maximum yield and biofilm

formation, we also measured the hypermucoviscosity of strains, a trait linked to
270 hypervirulence and driven, at least partly by the amount of capsule (50) (Figures S10-S12).
Firstly, we used Principal Components Analysis (PCA) to analyze how the different factors
(media, serotype or strain) explain the observed variance in the data. The first and second
principal components explained more than 60% of the variation observed (Figure 5A). We
observed no clear grouping of serotypes according to the two first axes. This suggests that the
275 serotype is not a major determinant of the capsule-related phenotypes (Figure S13). The
ellipses regrouping the points by type of media show a small overlap between both rich media
(ASM and LB) and these are clearly separated from the two poorest media (M005 and AUM).
This suggests that most of the variance is explained by the media, and more presumably by
the amount of nutrients available (Figure 5A). We then used multifactorial ANOVA to
280 analyze quantitatively the effect of the environment and the serotype on each trait. As
expected, we observed a significant effect of the environment in all measured traits, except in
biofilm formation ($R^2 = 0.57$, P = 0.06, Table S2). As suggested by the PCA, the influence of
the serotype in the capsule-associated traits is less important. Interestingly, the terms of
interaction between the environment and the serotype are not statistically significant,
285 suggesting that they are independent (Table S2).

Secondly, we sought to determine how the different capsule-related variables correlated with
each other. The PCA analysis already suggested the existence of such correlations (Figure
5B). As expected, we observe a strong correlation between hypermucoviscosity (HMV) and
290 the amount of capsule, in all individual growth media except in M005. Consistently with
previous reports (51, 52), we observe that the population yield is inversely proportional to the
growth rate (Figure 5B). This is significant in all growth media, except M005 (for which P =
0.07). Further, the amount of capsule is positively correlated with the maximum yield (Figure
5B), supporting our observation that the capsule contributes to increased yield of the
295 population. In contrast, there is a negative correlation between the amount of capsule and the
generation time. These results are driven by growth in nutrient poor environments (rho < -0.2
and rho > 0.2 in nutrient poor and nutrient rich environments) (Figure 5B). More precisely, in
poor media, bacteria with more capsule are able to grow faster that those producing less
capsule. This was not evident from the mutant analysis, as these showed no significant
300 differences in generation time between the capsulated and non-capsulated strains. Overall, our
results show that the capsule may impact several important cellular traits, like mucoviscosity
and growth, but this is strongly dependent on the environment.

DISCUSSION

The capsule is an extracellular structure commonly associated to the ability to resist biotic or abiotic aggressions, like the immune system, phages, or desiccation (10, 39, 53). Here, we explore further the conditions in which the capsule can be selected for, despite the cost of its production. Our results show that simple abiotic factors, such as nutrient limitation, can change dramatically whether the capsule is maintained in a population. More precisely, the capsule can be selected for in the absence of biotic stresses or strong abiotic aggressions. During selection for fast growth in nutrient-poor environments, capsulated cells have strong fitness advantages (*ca.* 20%). These challenges the current view that the main role of the capsule is to withstand aggressions. Our previous work already suggested that its role in virulence may be a by-product of adaptation, raising the question of its primary function in cells (6). Here we further show that, even in the absence of strong aggressions, the capsule provides a notorious advantage and may be selected because of the advantage it provides during growth.

We hypothesized that the cost of the capsule could be reflected in a reduced growth rate between of wild type strains compared to their respective capsule mutants. We, however, do not observe such differences, as calculated by the minimum generation time (Figure 3B and 6B). One possible explanation could be the lack of sensitivity of absorbance tests or to the large variation across biological replicates (over 10 %). However, our results are in line with studies performed in other species, for instance, in *Acinetobacter baumanii,* where the growth of wild type and capsule mutants in laboratory media was similar (54). Experiments performed in *S. pneumoniae* showed differences in growth rate between wild type and non-capsulated mutants, but these were dependent on capsule serotype. The loss of capsule 7B increased significantly the growth rate in rich media whereas the opposite was true for capsule 6B (55). The lack of observable differences may be due to the low cost of the production of capsule, or to the fine-tuning of its complex regulatory network (56). Indeed, the capsule could be regarded as an essential constituent of the physiology of *Klebsiella spp.* as, so far, few unencapsulated strains have been isolated, and the genome could have minimized the cost of its production. Despite the absence of differences between capsulated and unencapsulated strains, significant differences are observed across strains and environments, most notably, we

observe a negative correlation between the amount of capsule and generation time in nutrient poor environments (Figure 5B). Thus, bacteria producing more capsule are able to grow faster in nutrient poor environments, again highlighting the central role of the capsule in such environments and suggesting that that there may be other unknown functions to the capsule.

The increased fitness in nutrient-poor environments can be explained by the differences in population yield, or the increase of the reproductive lifespan of the population, as determined both by CFU and absorbance. In nutrient-poor environments, capsulated cells achieve higher yields and are fitter. Reduced yield in capsulated mutants was already observed in strains of serotype 6B of *S. pneumoniae* (55). How capsulated bacteria achieve a higher yield is still not clear, but could be due to several reasons. We first hypothesized that the capsule could increase survival in our cultures. The advantage provided by the capsule could be associated with a growth advantage, or reduced death, in stationary phase (GASP phenotype), a period in which birth and death rates are balanced. However, our results do not support this (Figure S6C). The phase referred to as "death phase" during growing cultures typically sets in after two or three days of batch culture in other Enterobacteria such as *E. coli* (57), and we observe higher yield after only 16 hours. Further, our long-term starvation experiments revealed no differences in survival between capsulated and non-capsulated strains. Previous reports had already shown that in the long term, *Klebsiella spp* are very resilient and have very slow death rates (58, 59), including in lake water at temperatures close to 0 (60) or during long-term starvation in PBS as a member of an oral community (46). Second, capsulated bacteria could use the capsules' polysaccharides to further multiply after consumption of all the environmental resources, as previously reported in *S. pneumoniae* (45). Addition of exogenous capsule to non-capsulated mutants did not result in a significant increase of CFU in a population, suggesting that it does not allow for higher population yields. Third, our analysis shows that biofilm formation, capsule amount and hypermucovicosity are positively correlated with yield (Figure 5B). This suggests that capsules could promote aggregation and thus cells could be present at higher local densities, due to limited dispersion and clumping during growth, which would in term, boost their individual fitness and increase total population yield. For example, dead capsulated cells would be dispersed less and could be more readily consumed by neighbouring growing cells.

Further analysis of our panel of strains reveals that capsule-associated features related to virulence are very variable across all strains and serotypes. First, we observe an overall strong

positive correlation between the capsule and hypermucoviscosity (HMV), a proxy of hypervirulence. This supports the hypothesis that HMV is a result of capsule overproduction. Yet, a recent study has shown that capsule overproduction is not required for HMV (61), suggesting that the role of the capsule in HMV is convoluted or dependent on the strain (56).

375   Indeed, we observe an effect of serotype in HMV (P = 0.002). Second, our results are in agreement with a previous study (62) in which cells from different serotypes were first exposed to depolymerases, to digest the capsule, and then tested for resistance to human serum and phagocytosis. Consistently, some non-capsulated strains were more sensitive to human serum, but others seem to be as resistant as the control cultures to which no

380   depolymerase was previously added. Resistance to human serum can also be due to the presence different O-antigens (63, 64). In our panel, only two strains #211 and #26 share the same O-antigen (O1v1). These strains have different resistance profiles, suggesting that O-antigen is not solely responsible for serum resistance. Third, we do not observe any protection of the capsule from bile salts at physiological levels, as *Klebsiella* strains here tested seem to

385   be naturally resistant to bile salts. When bile salt concentrations are increased by a factor of ten, the capsule significantly increases survival in some cells but not all. Overall, these results highlight that the capsules do not have the same function in all *Klebsiella* strains, and are thus, non-universal. Further, it highlights that the absence of capsule may not necessarily affect the potential of these strains for pathogenesis. A recent study showed that compared to

390   capsulated carbapenem-resistant *K. pneumoniae* ST258 isolates, the unencapsulated isolates formed more biofilm and could invade the epithelial bladder tissue more effectively. This resulted in the formation of intracellular infective reservoirs that lead to recurrent urinary infections (65).

395   To conclude, we show that the growth media affects the relative fitness of capsulated bacteria, which may justify selection for capsulated bacteria in nutritionally poor environments. One cannot ignore the possibility that such a selection pressure may drive capsule diversification, e.g. different capsules could be better adapted to certain poor environments because of their different propensity to aggregate or because of their organization at the cell surface (66, 67).

400   As previously suggested (67), the non-universality of some of the capsule functions across and within species suggests that there may be some functions that are very different to those that are commonly evoked during pathogenesis. It could also be that different serotypes are associated to specific functions or fitness advantages. This might lead to selection of serotype switching when bacteria colonize novel environments. Yet, it seems more likely that the rapid

405   turnover of capsular serotypes is due to biotic pressures, such as phage predation, cell-to-cell interactions, the host adaptive immune system, or protozoa grazing (29, 68). Taken together, our work suggests that the primary evolutionary forces selecting for the existence of a capsule may be very different to those selecting for its subsequent diversification, and highlights that much work is still needed to fully understand the primary role of capsules, both at the

410   molecular and at the evolutionary level.

MATERIALS AND METHODS

415   **Bacterial strains and growth conditions.** Bacterial strains and plasmids used in this study are listed in Table S1. Primers used for the construction of mutants are listed in Table S3. Bacteria were grown at 37° in Luria-Bertani (LB) agar plates or in 4 mL of liquid broth under vigorous shaking (250 rpm) unless indicated otherwise. Kanamycin (50 µg/ml) or Streptomycin (100 µg/mL for *E. coli* and 200 µg/ml for *Klebsiella*) were used in the selection

420   of the plasmids strain selection and cultures. Antibiotics were purchased from Sigma.

   Growth media description. AUM and ASM stand for artificial urine medium and artificial sputum medium, respectively, and were done as described previously (41, 42). Briefly, AUM is mainly composed of urea and peptone with trace amounts of lactic acid, uric acid, creatinine and peptone. ASM is composed of mucin, DNA, egg yolk and aminoacids. LB is

425   composed of tryptone and yeast extract. M02 and M005 indicate minimal M63B1 supplemented with 0.2% and 0.05% glucose (sole carbon source), respectively.

   Determination of serotype. Complete or draft genomes were serotyped using Kaptive with default settings (69).

   Natural loss of capsule. An overnight culture in LB was diluted at 1:100 in the different

430   environments (ASM, LB, M02, M005, AUM) in a final volume of 4 mL. Cultures were allowed to grow for 24 hours at 37°C and diluted again to 1:100 in fresh media. This was repeated for 3 times. Then, each culture in each environment was serially diluted and CFUs were counted (3 plates per sample, > 100 colonies per plate). Non-capsulated mutants are easily visualized by the naked eye as mutants produce smaller, rough and translucent

435   colonies. Results are expressed as a ratio of non-capsulated mutant colonies observed over the total plate count. The limit of detection of non-capsulated mutants in this assay was 0.22% ± 0.04 or 1:450 per biological replicate per environment. To ensure that the naturally-emerged non-capsulated colonies observed on plate are *bona fide* genetic mutants and not phase-variants, we randomly chose 10 populations, using the *sample* function in R. Three non-

440   capsulated clones from each population were randomly selected and restreaked twice. All clones derived from the original non-capsulated colony remained non-capsulated, demonstrating that the loss of the capsule is genetic.

Generation of *Δwza* capsule mutants. Isogenic capsule mutants were constructed by an in-frame deletion of *wza* by allelic exchange. Upstream and downstream sequences of the *wza* gene (> 500pb) were amplified using Phusion Master Mix (Thermo Scientific) then joined by overlap PCR. The resulting PCR product was purified using the QIAquick Gel Extraction Kit and then cloned with the Zero Blunt® TOPO® PCR Cloning Kit (Invitrogen) into competent *E. coli* DH5α strain. KmR colonies were isolated and checked by PCR. Cloned Zero Blunt® TOPO® plasmid was extracted using the QIAprep Spin Miniprep Kit, and digested for 2 hours at 37°C with ApaI and SpeI restriction enzymes and ligated with T4 DNA ligase (Promega) overnight at 16° to a double-digested pKNG101 plasmid. The ligation was transformed into competent *E. coli* DH5α pir strain, and again into *E. coli* MFD λ-pir strain (70), used as a donor strain for conjugation into *Klebsiella spp*. Conjugations were performed for 24 hours at 37°C. Single cross-over mutants (transconjugants) were selected on Streptomycin plates (200 µg/ml). To select for plasmid excision, transconjugants were grown for 16 hours at 25°C and double cross-over mutants were selected on LB without salt and supplemented with 5% sucrose at room temperature. To confirm the loss of the plasmid, colonies were tested for their sensitivity to streptomycin. From each double-recombination, a mutant and a wild type were isolated. All experiments were performed in parallel with the parental strain, *Δwza* mutant and the associated wild type, to ensure that the phenotype observed was directly due to the *Δwza* deletion and not to other potential SNPs that could have accumulated in the genome during the mutant generation, and thus present in both the mutant and the wild type. Deletion mutants were first verified by Sanger, and then checked for off-target mutations using Illumina. Three strains presented off-target mutations. Strain #63 (KL107) *Δwza* has a 15bp deletion upstream of *rfaH*, another capsule regulator. This inhibits the initiation of capsule synthesis altogether. Strain #51 (KL24) has a SNP in *pcaR*. There are no reports on a direct influence of this gene on capsule production. Strain #55 (KL2) has a SNP in *sacX* gene which results in delayed growth. Despite the fact that these strains are *bona fide* capsule mutants, as shown by microscopic observation and capsule quantification, the results are presented as supplemental figures.

Microscopy. Bacteria capsules were visualized using Indian Ink coloration on an exponential phase culture. Images were acquired on a Nikon Eclipse E400 camera using the Digital Net DN100 camera control unit and a 100X objective.

Capsule extraction and quantification. The bacterial capsule was extracted as described in (71) and quantified by using the uronic acid method (72). Briefly, 500 μL of an overnight culture was mixed with 100 μL of 1% Zwittergent 3-14 detergent in 100 mM citric acid (pH 2.0) and heated at 56°C for 20 minutes. After it was centrifuged for 5 min at 14,000 rpm and 300 μL of the supernatant was transferred to a new tube. Absolute ethanol was added to a final concentration of 80% and the tubes were placed on ice for 20 minutes. After a second wash with ethanol at 70%, the pellet was dried and dissolved in 250 μL of distilled water. The pellet was then incubated for 2 hours at 56°C. Polysaccharides were then quantified by measuring the amount of uronic acid. A 1,200 μL volume of 0.0125 M tetraborate in concentrated H_2SO_4 was added to 200 μL of the sample to be tested. The mixture was vigorously vortexed and heated in a boiling-water bath for 5 min. The mixture was allowed to cool, and 20 μL of 0.15% 3-hydroxydiphenol in 0.5% NaOH was added. The tubes were shaken, and 100 μL were transferred to a microtiter plate for absorbance measurements (520 nm). The uronic acid concentration in each sample was determined from a standard curve of glucuronic acid.

Competition assay. Mutants and their respective wild types were grown overnight in LB. They were then mixed in a 1:1 proportion. A sample was taken and used for serial dilution and CFU counting as a control of T_0. The co-culture was then diluted 1/100 in 4 mL of the different environments (i.e. LB, AUM, ASM, M02 and M005). After 24 hours of competition (T_{24}), each mixture was serially diluted and plated. Capsulated and non-capsulated colonies are clearly differentiated visually and counted separately. The competitive index of capsulated strains was calculated by dividing the ratio of CFU at T_{24} over T_0.

Growth curves. Overnight cultures were diluted at 1:100 in the aforementioned environments. 200 μl of each subculture was transferred in a 96-well microplate. Cell densities of cultures were estimated with a TECAN Genios™ plate reader. Absorbance (OD_{600}). Light absorbance values of each well reflect the average of 6 reads per well. Absorbance values from within-block technical replicates were averaged and these averages were used as statistically independent data points. Minimum generation times were estimated across replicates for the 1 hour interval (ΔT) spanning the fastest growth during exponential growth phase. This was calculated as:

$$\frac{1}{Td} = \frac{log_{10}(\text{maximum}_{OD600} / \text{minimum}_{OD600})}{log_{10}(2) * \Delta T}$$

Relative yields were calculated measuring ratios of the maximum absorbance of the assay compared to the reference (parental strain). The experiments were performed at least three independent times, with three technical replicates per experiment. The average of the three technical replicates per experiment was used for statistical purposes. Control experiments

515 were performed to assess the correlation between CFU and OD_{600} during growth for both capsulated (Spearman's rho = 1.0, P = 0.002) and non-capsulated mutants (Spearman's rho = 0.94, P = 0.016). To confirm the differences in yield observed in minimal medium between capsulated and non-capsulated strains, we performed a control experiment (N = 3) in which we grew the strains #56 and #24 in microtiterplates as described before. After 16h, we plated

520 an aliquot to count CFUs. Significantly less CFUs were observed in the non-capsulated mutants compared to wild type (P<0.05 for both strains).

Serum resistance. The ability of *Klebsiella* strains to resist killing by serum was performed as described previously (20, 43, 73). Briefly, bacteria were grown in LB to OD 600 nm of 1,

525 pelleted by centrifugation and resuspended in sterile PBS 1X. 200 µl of bacterial suspension was added to 400 µl of pre-warmed at 37°C human sera (Sigma-Aldrich S7023), and the mixture was incubated for 2h at 37°C. Control reactions were performed with serum inactivated by heat at 56°C for 30 min. Reactions were stopped by placing on ice, and viable bacterial counts were determined before and after incubation by serial dilution and CFU

530 plating.

Bile salts resistance. Bacteria were grown overnight in LB. As previously described (74), cultures were then serially diluted and plated in LB supplemented with either 0, 0.05% (physiological conditions) and 0.5% of CHO (primary bile salt, sodium cholate) or DCH

535 (secondary bile salt, sodium deoxycholate). Colonies were allowed to grow and surviving CFU were counted after 24 hours. The average of two technical replicates per experiment was used for statistical purposes.

Starvation. Bacteria were grown in LB to OD=0.2, 1mL of the culture was pelleted by

540 centrifugation and, resuspended in 400 µl of PBS 1X, as in (45). Cultures were allowed to

starve at room temperature. Aliquots were taken at 0, 30 and 60 days, and used for serial dilutions and CFU plating.

Biofilm Production. The ability of strains to form biofilm was performed as described in (75). Overnight cultures were diluted at 1:100 in the different growth media mentioned earlier. 100 µL of each diluted culture was transferred in triplicate into 96-well microplate wells and allowed to grow for 24 hours without shaking at 37°C. Cells were gently removed from the wells and washed once in distilled water. To stain biofilms, 125 µL of 1% crystal violet was added to each well for 20 minutes. The crystal violet was gently decanted and each well was washed three times with distilled water. After the plate was totally dry, the biofilm was solubilized for 10 min in 150 µL of mix with 80% ethanol and 20% acetone. The absorbance of the sample was read at OD_{570}. At least 3 biological independent replicates were performed for each sample.

Hypermucoviscosity. The optical density OD_{600} of overnight cultures were first determined by spectrophotometry as described in (43). The cultures were then sedimented by slow centrifugation (at 2,500 X g) for 5 min at room temperature. The optical density OD_{600} of the top 650 µL of supernatant was determined. Results were expressed as a ratio of the supernatant OD_{600} to that of the input culture.

Dessication test. Dessication assay was carried as previously described in (76). Briefly, 25 uL of exponentially-growing *Klebsiella* strains were deposited at the bottom of microtiterplates, and allowed the culture to dry under a laminar flow hood. Strains were then left either at 37° for one, two or four days or at room temperature for two, four, seven and 14 days. PBS was added to resuspend the cultures and diluted serially from CFU counts.

Quantification of dead cells. Cells were allowed to grow as described above with a minor modification; over day cultures were used to inoculate the 96-well microtiter plates. After 16 hours of growth in M02, dead cells were quantified using the LIVE/DEAD™ BacLight™ Bacterial Viability Kit from Invitrogen, following provider instructions. Images were acquired on a Nikon Eclipse E400 microscope using the Digital Net DN100 camera control unit and a 100X objective. Cells were counted using ImageJ software (https://imagej.net/). On average, 6 pictures were taken per sample (min = 3, max = 10) and *ca.* 550 cells per sample were observed (min = 161, max =1604). The experiment was performed four times.

Growth on exogenously added capsule. Capsules were extracted from either wild type strains or their respective Δ*wcaJ* or Δ*wbaP* (for strain 24) mutants and quantified as described above. To initiate growth, overnight cultures of Δ*wcaJ* mutants were diluted at 1:100 in M63B1 media. Media was supplemented with either 0.8 ng/uL of extracted capsule from wild type or an equivalent volume extracted from the Δ*wcaJ* or Δ*wbaP* (for strain 24) mutants. As a negative control, cultures were allowed to grow in M63B1 with no nutrients. In parallel, M63B1 was supplemented with glucuronic acid (the standard used for estimating capsule concentration) at 0.8 ng/uL final. Growth was allowed for 48 hours at 37° and 250 rpm. Bacterial counts were performed at T_0, T_{24} and T_{48} by serial dilution and plating. Results are expressed as the increase in CFU over 24h.

REFERENCES

1. Meyers LA, Bull JJ. 2002. Fighting change with change: adaptive variation in an uncertain world. Trends Ecol Evol 17:551-557.

2. Levins R. 1968. Evolution in changing environments; some theoretical explorations. Princeton University Press, Princeton, NJ.

3. Rittershaus ES, Baek SH, Sassetti CM. 2013. The normalcy of dormancy: common themes in microbial quiescence. Cell Host Microbe 13:643-51.

4. Ophir T, Gutnick DL. 1994. A role for exopolysaccharides in the protection of microorganisms from desiccation. Appl Environ Microbiol 60:740-5.

5. Tipton KA, Chin CY, Farokhyfar M, Weiss DS, Rather PN. 2018. Role of Capsule in Resistance to Disinfectants, Host Antimicrobials, and Desiccation in *Acinetobacter baumannii*. Antimicrob Agents Chemother 62: e01188-18.

6. Rendueles O, Garcia-Garcera M, Neron B, Touchon M, Rocha EPC. 2017. Abundance and co-occurrence of extracellular capsules increase environmental breadth: Implications for the emergence of pathogens. PLoS Pathog 13:e1006525.

7. Jung SY, Matin A, Kim KS, Khan NA. 2007. The capsule plays an important role in *Escherichia coli* K1 interactions with *Acanthamoeba*. Int J Parasitol 37:417-23.

8. Geisinger E, Huo W, Hernandez-Bird J, Isberg RR. 2019. *Acinetobacter baumannii*: Envelope Determinants That Control Drug Resistance, Virulence, and Surface Variability. Annu Rev Microbiol 73:481-506.

9. Fernebro J, Andersson I, Sublett J, Morfeldt E, Novak R, Tuomanen E, Normark S, Normark BH. 2004. Capsular expression in *Streptococcus pneumoniae* negatively affects

605   spontaneous and antibiotic-induced lysis and contributes to antibiotic tolerance. J Infect Dis 189:328-38.

10. Campos MA, Vargas MA, Regueiro V, Llompart CM, Alberti S, Bengoechea JA. 2004. Capsule polysaccharide mediates bacterial resistance to antimicrobial peptides. Infect Immun 72:7107-14.

610   11. Band VI, Weiss DS. 2015. Mechanisms of Antimicrobial Peptide Resistance in Gram-Negative Bacteria. Antibiotics (Basel) 4:18-41.

12. Podschun R, Ullmann U. 1992. Klebsiella capsular type K7 in relation to toxicity, susceptibility to phagocytosis and resistance to serum. J Med Microbiol 36:250-4.

13. Williams P, Lambert PA, Brown MR, Jones RJ. 1983. The role of the O and K antigens

615   in determining the resistance of *Klebsiella aerogenes* to serum killing and phagocytosis. J Gen Microbiol 129:2181-91.

14. Tomas JM, Benedi VJ, Ciurana B, Jofre J. 1986. Role of capsule and O antigen in resistance of *Klebsiella pneumoniae* to serum bactericidal activity. Infect Immun 54:85-9.

15. Bentley SD, Aanensen DM, Mavroidi A, Saunders D, Rabbinowitsch E, Collins M,

620   Donohoe K, Harris D, Murphy L, Quail MA, Samuel G, Skovsted IC, Kaltoft MS, Barrell B, Reeves PR, Parkhill J, Spratt BG. 2006. Genetic analysis of the capsular biosynthetic locus from all 90 pneumococcal serotypes. PLoS Genet 2:e31.

16. Guerry P, Poly F, Riddle M, Maue AC, Chen YH, Monteiro MA. 2012. Campylobacter polysaccharide capsules: virulence and vaccines. Front Cell Infect Microbiol 2:7.

625   17. Giguere D. 2015. Surface polysaccharides from *Acinetobacter baumannii*: Structures and syntheses. Carbohydr Res 418:29-43.

18. Mori M, Ohta M, Agata N, Kido N, Arakawa Y, Ito H, Komatsu T, Kato N. 1989. Identification of species and capsular types of Klebsiella clinical isolates, with special reference to *Klebsiella planticola*. Microbiol Immunol 33:887-95.

630   19. Orskov I. 1955. Serological investigations in the Klebsiella group. I. New capsule types. Acta Pathol Microbiol Scand 36:449-53.

20. Podschun R, Ullmann U. 1998. Klebsiella spp. as nosocomial pathogens: Epidemiology, taxonomy, typing methods, and pathogenicity factors. Clinical Microbiol Rev 11:589-603.

21. Follador R, Heinz E, Wyres KL, Ellington MJ, Kowarik M, Holt KE, Thomson NR.

635   2016. The diversity of *Klebsiella pneumoniae* surface polysaccharides. Microb Genom 2:e000073.

22. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, Holt KE. 2016. Identification of Klebsiella capsule synthesis loci from whole genome data. Microb Genom 2:e000102.

640 23. Mostowy RJ, Croucher NJ, De Maio N, Chewapreecha C, Salter SJ, Turner P, Aanensen DM, Bentley SD, Didelot X, Fraser C. 2017. Pneumococcal Capsule Synthesis Locus cps as Evolutionary Hotspot with Potential to Generate Novel Serotypes by Recombination. Mol Biol Evol 34:2537-2554.

24. Wyres KL, Gorrie C, Edwards DJ, Wertheim HFL, Hsu LY, Van Kinh N, Zadoks R,
645 Baker S, Holt KE. 2015. Extensive Capsule Locus Variation and Large-Scale Genomic Recombination within the *Klebsiella pneumoniae* Clonal Group 258. Genome Biol Evol 7:1267-1279.

25. Croucher NJ, Kagedan L, Thompson CM, Parkhill J, Bentley SD, Finkelstein JA, Lipsitch M, Hanage WP. 2015. Selective and genetic constraints on pneumococcal serotype
650 switching. PLoS Genet 11:e1005095.

26. McBride SM, Fischetti VA, LeBlanc DJ, Moellering RC, Gilmore MS. 2007. Genetic Diversity among *Enterococcus faecalis*. Plos One 2:e582.

27. Thrane SW, Taylor VL, Freschi L, Kukavica-Ibrulj I, Boyle B, Laroche J, Pirnay JP, Levesque RC, Lam JS, Jelsbak L. 2015. The Widespread Multidrug-Resistant Serotype O12
655 *Pseudomonas aeruginosa* Clone Emerged through Concomitant Horizontal Transfer of Serotype Antigen and Antibiotic Resistance Gene Clusters. MBio 6:e01396-15.

28. Wyres KL, Wick RR, Judd LM, Froumine R, Tokolyi A, Gorrie CL, Lam MMC, Duchene S, Jenney A, Holt KE. 2019. Distinct evolutionary dynamics of horizontal gene transfer in drug resistant and virulent clones of *Klebsiella pneumoniae*. PLoS Genet
660 15:e1008114.

29. Mostowy RJ, Holt KE. 2018. Diversity-Generating Machines: Genetics of Bacterial Sugar-Coating. Trends Microbiol 26:1008-1021.

30. Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, Bentley SD, Hanage WP, Lipsitch M. 2013. Population genomics of post-vaccine changes in
665 pneumococcal epidemiology. Nat Genet 45:656-63.

31. Lamb KE, Flasche S, Diggle M, Inverarity D, Greenhalgh D, Jefferies JM, Smith A, Edwards GFS, Denham B, McMenamin J, McDonald E, Mitchell TJ, Clarke SC, Robertson C. 2014. Trends in serotypes and sequence types among cases of invasive pneumococcal disease in Scotland, 1999-2010. Vaccine 32:4356-4363.

670   32. Randall WA. 1939. Colony and Antigenic Variation in *Klebsiella pneumoniae* Types A, B and C. J Bacteriol 38:461-77.

33. Julianelle LA. 1928. Bacterial Variation in Cultures of Friedlander's Bacillus. J Exp Med 47:889-902.

34. Bagley ST. 1985. Habitat association of Klebsiella species. Infect Control 6:52-8.

675   35. Blin C, Passet V, Touchon M, Rocha EPC, Brisse S. 2017. Metabolic diversity of the emerging pathogenic lineages of *Klebsiella pneumoniae*. Environ Microbiol 19:1881-1898.

36. Caneiras C, Lito L, Melo-Cristino J, Duarte A. 2019. Community- and Hospital-Acquired *Klebsiella pneumoniae* Urinary Tract Infections in Portugal: Virulence and Antibiotic Resistance. Microorganisms 7:138.

680   37. Lee CR, Lee JH, Park KS, Jeon JH, Kim YB, Cha CJ, Jeong BC, Lee SH. 2017. Antimicrobial Resistance of Hypervirulent *Klebsiella pneumoniae*: Epidemiology, Hypervirulence-Associated Determinants, and Resistance Mechanisms. Front Cell Infect Microbiol 7:483.

38. Bengoechea JA, Pessoa JS. 2019. *Klebsiella pneumoniae* infection biology: living to
685   counteract host defences. FEMS Microbiol Rev 43:123-144.

39. Paczosa MK, Mecsas J. 2016. *Klebsiella pneumoniae*: Going on the Offense with a Strong Defense. Microbiol Mol Biol Rev 80:629-61.

40. de Sousa JAM, Buffet A, Haudiquet M, Rocha EPC, Rendueles O. 2019. Modular prophage interactions driven by capsule serotype select for capsule loss under phage
690   predation. bioRxiv doi:10.1101/2019.12.17.878363:2019.12.17.878363.

41. Fung C, Naughton S, Turnbull L, Tingpej P, Rose B, Arthur J, Hu H, Harmer C, Harbour C, Hassett DJ, Whitchurch CB, Manos J. 2010. Gene expression of *Pseudomonas aeruginosa* in a mucin-containing synthetic growth medium mimicking cystic fibrosis lung sputum. J Med Microbiol 59:1089-100.

695   42. Brooks T, Keevil CW. 1997. A simple artificial urine for the growth of urinary pathogens. Lett Appl Microbiol 24:203-6.

43. Dorman MJ, Feltwell T, Goulding DA, Parkhill J, Short FL. 2018. The Capsule Regulatory Network of *Klebsiella pneumoniae* Defined by density-TraDISort. MBio 9:e01863-18.

700   44. Wacharotayankun R, Arakawa Y, Ohta M, Hasegawa T, Mori M, Horii T, Kato N. 1992. Involvement of *rcsB* in Klebsiella K2 capsule synthesis in *Escherichia coli* K-12. J Bacteriol 174:1063-7.

45. Hamaguchi S, Zafar MA, Cammer M, Weiser JN. 2018. Capsule Prolongs Survival of *Streptococcus pneumoniae* during Starvation. Infect Immun 86: e00802-17.

705 46. Baker JL, Hendrickson EL, Tang XY, Lux R, He XS, Edlund A, McLean JS, Shi WY. 2019. Klebsiella and Providencia emerge as lone survivors following long-term starvation of oral microbiota. Proc Natl Acad Sci USA 116:8499-8504.

47. Hall-Stoodley L, Costerton JW, Stoodley P. 2004. Bacterial biofilms: from the natural environment to infectious diseases. Nat Rev Microbiol 2:95-108.

710 48. Schembri MA, Blom J, Krogfelt KA, Klemm P. 2005. Capsule and fimbria interaction in *Klebsiella pneumoniae*. Infect Immun 73:4626-33.

49. Schembri MA, Dalsgaard D, Klemm P. 2004. Capsule shields the function of short bacterial adhesins. J Bacteriol 186:1249-57.

50. Lin TL, Yang FL, Yang AS, Peng HP, Li TL, Tsai MD, Wu SH, Wang JT. 2012. Amino
715 acid substitutions of MagA in *Klebsiella pneumoniae* affect the biosynthesis of the capsular polysaccharide. PLoS One 7:e46783.

51. Pfeiffer T, Schuster S, Bonhoeffer S. 2001. Cooperation and competition in the evolution of ATP-producing pathways. Science 292:504-7.

52. Lipson DA. 2015. The complex relationship between microbial growth rate and yield and
720 its implications for ecosystem processes. Front Microbiol 6:615.

53. Bengoechea JA, Sa Pessoa J. 2019. Klebsiella pneumoniae infection biology: living to counteract host defences. FEMS Microbiol Rev 43:123-144.

54. Russo TA, Luke NR, Beanan JM, Olson R, Sauberan SL, MacDonald U, Schultz LW, Umland TC, Campagnari AA. 2010. The K1 capsular polysaccharide of *Acinetobacter*
725 *baumannii* strain 307-0294 is a major virulence factor. Infect Immun 78:3993-4000.

55. Hathaway LJ, Brugger SD, Morand B, Bangert M, Rotzetter JU, Hauser C, Graber WA, Gore S, Kadioglu A, Muhlemann K. 2012. Capsule type of *Streptococcus pneumoniae* determines growth phenotype. PLoS Pathog 8:e1002574.

56. Walker KA, Miller VL. 2020. The intersection of capsule gene expression,
730 hypermucoviscosity and hypervirulence in *Klebsiella pneumoniae*. Curr Opin Microbiol 54:95-102.

57. Finkel SE. 2006. Long-term survival during stationary phase: evolution and the GASP phenotype. Nat Rev Microbiol 4:113-20.

58. Bossolan N, Godinho MJL, Volpon AGT. 2005. Growth and starvation of a strain of
735 *Klebsiella pneumoniae i*solated from a Brazilian oil formation. World J Microbiol Biotechnol 21:1471-1475.

59. Lappin-Scott HM, Cusack F, MacLeod A, Costerton JW. 1988. Starvation and nutrient resuscitation of *Klebsiella pneumoniae* isolated from oil well waters. J Appl Bacteriol 64:541-9.

60. Niemela SI, Vaatanen P. 1982. Survival in lake water of *Klebsiella pneumoniae* discharged by a paper mill. Appl Environ Microbiol 44:264-9.

61. Walker KA, Miner TA, Palacios M, Trzilova D, Frederick DR, Broberg CA, Sepulveda VE, Quinn JD, Miller VL. 2019. A *Klebsiella pneumoniae* Regulatory Mutant Has Reduced Capsule Expression but Retains Hypermucoviscosity. mBio 10:e00089-19.

62. Majkowska-Skrobek G, Latka A, Berisio R, Squeglia F, Maciejewska B, Briers Y, Drulis-Kawa Z. 2018. Phage-Borne Depolymerases Decrease *Klebsiella pneumoniae* Resistance to Innate Defense Mechanisms. Front Microbiol 9:2517.

63. Doorduijn DJ, Rooijakkers SH, van Schaik W, Bardoel BW. 2016. Complement resistance mechanisms of *Klebsiella pneumoniae*. Immunobiology 221:1102-9.

64. Merino S, Camprubi S, Alberti S, Benedi VJ, Tomas JM. 1992. Mechanisms of *Klebsiella pneumoniae* resistance to complement-mediated killing. Infect Immun 60:2529-35.

65. Ernst CM, Braxton JR, Rodriguez-Osorio CA, Zagieboylo AP, Li L, Pironti A, Manson AL, Nair AV, Benson M, Cummins K, Clatworthy AE, Earl AM, Cosimi LA, Hung DT. 2020. Adaptive evolution of virulence and persistence in carbapenem-resistant *Klebsiella pneumoniae*. Nat Med doi:10.1038/s41591-020-0825-4.

66. Phanphak S, Georgiades P, Li R, King J, Roberts IS, Waigh TA. 2019. Super-Resolution Fluorescence Microscopy Study of the Production of K1 Capsules by *Escherichia coli*: Evidence for the Differential Distribution of the Capsule at the Poles and the Equator of the Cell. Langmuir 35:5635-5646.

67. Rendueles O. 2020. Deciphering the role of the capsule of *Klebsiella pneumoniae* during pathogenesis: A cautionary tale. Mol Microbiol *in press*.

68. Cobey S, Lipsitch M. 2012. Niche and neutral effects of acquired immunity permit coexistence of pneumococcal serotypes. Science 335:1376-80.

69. Wick RR, Heinz E, Holt KE, Wyres KL. 2018. Kaptive Web: User-Friendly Capsule and Lipopolysaccharide Serotype Prediction for Klebsiella Genomes. J Clin Microbiol 56.

70. Ferrieres L, Hemery G, Nham T, Guerout AM, Mazel D, Beloin C, Ghigo JM. 2010. Silent mischief: bacteriophage Mu insertions contaminate products of *Escherichia coli* random mutagenesis performed using suicidal transposon delivery plasmids mobilized by broad-host-range RP4 conjugative machinery. J Bacteriol 192:6418-27.

71. Domenico P, Schwartz S, Cunha BA. 1989. Reduction of capsular polysaccharide production in *Klebsiella pneumoniae* by sodium salicylate. Infect Immun 57:3778-82.

72. Blumenkrantz N, Asboe-Hansen G. 1973. New method for quantitative determination of uronic acids. Anal Biochem 54:484-9.

775 73. Podschun R, Sievers D, Fischer A, Ullmann U. 1993. Serotypes, hemagglutinins, siderophore synthesis, and serum resistance of Klebsiella isolates causing human urinary tract infections. J Infect Dis 168:1415-21.

74. Tan YH, Chen Y, Chu W, Sham L, Gan YH. 2020. Cell envelope defects of different capsule-null mutants in K1 hypervirulent *Klebsiella pneumoniae* can affect bacterial
780 pathogenesis. Mol Microbiol *in press*.

75. Rendueles O, Travier L, Latour-Lambert P, Fontaine T, Magnus J, Denamur E, Ghigo JM. 2011. Screening of *Escherichia coli* species biodiversity reveals new biofilm-associated antiadhesion polysaccharides. MBio 2:e00043-11.

76. Chin CY, Tipton KA, Farokhyfar M, Burd EM, Weiss DS, Rather PN. 2018. A high-
785 frequency phenotypic switch links bacterial virulence and environmental survival in *Acinetobacter baumannii*. Nat Microbiol 3:563-569.

ACKNOWLEDGEMENTS

We thank Matthieu Haudiquet for critical reading of the manuscript. We thank Sylvain Brisse
790 for providing us with the necessary Klebsiella strains, and Christiane Forestier and Damien Balestrino for providing the pKNG101 plasmid used to generate the mutants. We would also like to thank Jean-Marc Ghigo and Christophe Beloin for the gift of *E. coli* S17 MFD λ-pir strain and *E. coli* DH5α-pir.

795 ## FUNDING

This work was supported by an ANR JCJC (Agence national de recherche) grant [ANR 18 CE12 0001 01 ENCAPSULATION] awarded to O.R.. The laboratory is funded by a Laboratoire d'Excellence 'Integrative Biology of Emerging Infectious Diseases' grant [ANR-10-LABX-62-IBEID], the INCEPTION programme, and the FRM [EQU201903007835]. The
800 funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

FIGURE LEGENDS

805 **Figure 1. Frequency of capsulated colonies across strains and growth media after three days of evolution.** Strains were diluted 1:100 and inoculated in fresh media every 24 hours during three days. Strain #212 was unable to grow in M005. Points represent the average loss of capsule across at least three independently evolving populations per strain. Error bars represent standard deviation across replicate populations.

Figure 2. Competition between capsule mutants and their associated wild type in different growth media. Each point represents the average of at least three independent replicates. Individual error bars for each strain are not displayed for clarity purposes. Colours

815 represent different serotypes (as in Figure 1), whereas the shape represent different species. Results for the strains with compensatory mutations are presented in the supplemental material (Figure S5). * P < 0.05 ; ** P< 0.01, Tukey post hoc and correction for multiple tests.

Figure 3. **Survival, growth rate and yield of strains in each environment. A**. Survival of parental, Δwza mutant and the associated wild type, relative to day 0. Data is $\log_{10}$ - transformed. Individual error bars for each strain are not displayed for clarity purposes. Colours represent different serotypes (as in Figure 1), whereas the shape represent different

825 species (ANOVA, P > 0.05). **B.** Minimum generation time of each strain in each environment, expressed in minutes. Error bars represent standard deviations of the mean, of at least three independent experiments. **C.** Maximum yield of the mutant and the wild type strains relative to the parental strain in each environment. Statistics were calculated using a paired t-test. * P < 0.05.

Figure 4. Fate of capsule mutants in host-related context. A. Resistance to human serum. Data is presented as relative survival to control (heat-inactivated human serum). * P< 0.05; ** P< 0.01; *** P<0.001 paired t-test. **B.** Correlation between the total amount of biofilm formed by the wild type and the difference in biofilm formation between the wild type and the

835 deletion mutant across. P-values correspond to Spearman's rank correlation. Results are

qualitatively similar when only taking into account the five strains with no off-target mutations. Absolute values of biofilm formation are shown in Figure S8.

Figure 5. The environment, not the serotype, explains most of the variance observed
840   **across capsule-associated traits. A**.PCA analyses of capsule-related phenotypes. Phenotypes are coloured by environment. Each individual point represents a strain. PCA was performed using *prcomp* function (options scale and center = TRUE) in R and ggbiplot package for visualization purposes. **B.** Correlation matrix of all traits. Correlation matrix of all traits. Numbers indicate Spearman's correlation. Stars identify significant correlations (P<0.05).

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Capsule quantification of each strain in each growth media by the uronic acid method.

Figure S2. Ratio of capsulated colonies throughout 30 days of evolution in two media. Strains were diluted 1:100 and inoculated in fresh media every 24 hours during 30 days. CFUs were counted every day. Each line represents an independent population (N = 3).

855   **Figure S3. Capsule quantification of each mutant by the uronic acid method.** This experiment was performed on overnight cultures grown in LB. Each dot corresponds to an independent experiment. The background noise of this assay is established around 1.5 ng/uL. Statistical tests: * P < 0.05; ** P < 0.01; *** P<0.001.

860   **Figure S4. Fitness advantage/disadvantage of capsulated strains in nutrient poor/rich media.** Competition between wild type and their isogenic capsule mutants in different growth media. Each point represents the average of at least three independent replicates. * P < 0.05 Tukey post hoc, p-values adjusted for multiple testing.

865   **Figure S5**. **Survival of strains under starvation and desiccation. A**. Survival of parental, mutant and wild type strains, relative to day 0. Data is $\log_{10}$ -transformed. Error bars are not displayed for clarity purposes. Colours represent different serotypes (as in Figure 1), whereas the shape represent different species. **B.** Cellular survival, in CFU, after 1, 2 or four days of desiccation at 37°C. Error bars for independent points are not shown for clarity purposes.

Figure S6. **Growth rate and yield of strains in each media. A.** Minimum generation time of each strain in each media, expressed in minutes. Error bars represent standard deviations of the mean, of at least three independent experiments. **B.** Maximum yield of the mutant and the wild type strains relative to the parental strain in each growth media. Statistics were

875   calculated using a paired t-test. * P < 0.05. **C.** Percentage of dead cells after 16 hours growth in a microtiter plate well. Each symbol represents an independent experiment. No significant differences are found across genotypes.

880 **Figure S7**. **Log$_{10}$-Increase of CFU after 48 hours of growth.** Growth of $\varDelta$wcaJ (26 and 56) and $\varDelta$wbaP (24) mutants was followed for 24 and 48 hours. Cells were grown in M63B1 without nutrient source (Control), with 0.8 ng/uL of purified capsule (Cap+), an equivalent volume of extraction of a non-capsulated strain (Cap-), and 0.8 ng/uL of glucuronic acid (the standard used to quantify the capsule). Individual points represent four independent

885 experiments.

Figure S8. Fate of capsule mutants in host-related context. A. Resistance to human serum. Data is presented as relative survival to control (heat-inactivated human serum). **B.** Resistance to bile salts at physiological concentrations (0.05%), and at 10X (0.5%). Results are presented

890 as Log$_{10}$ (relative to control, LB plates). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Figure S9. Biofilm formation of parental, mutant and wild type strains across growth media. P values reflect paired t-tests. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

895 **Figure S10. Growth rate and maximum yield of each population in each environment.**

Figure S11. Biofilm formation of all strains in all environments.

Figure S12.Mucoviscosity index of strains in all environments.

Figure S13. PCA analyses of capsule-related phenotypes. Phenotypes are coloured by serotype. Each individual point represents a strain. PCA was performed using *prcomp* function in R and ggbiplot package for visualization purposes.