Anaerobic microbial degradation of protein and lipid macromolecules in subarctic marine sediment

Claus Pelikan[1,2#], Kenneth Wasmund[1,2#*], Clemens Glombitza[3,4], Bela Hausmann[1,5,6], Craig W. Herbold[1], Mathias Flieder[1], and Alexander Loy[1,2,5*]

[1]*Division of Microbial Ecology, Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria.*

[2]*Austrian Polar Research Institute, Vienna, Austria.*

[3]*Center for Geomicrobiology, Department of Biology, Aarhus University, Aarhus, Denmark.*

[4]*ETH Zürich, Department of Environmental Systems Science, Zürich, Switzerland.*

[5]*Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna, Vienna, Austria*

[6]*Department of Laboratory Medicine, Medical University of Vienna, Vienna, Austria*

[#]Contributed equally.

*Correspondence: Kenneth Wasmund (kenneth.wasmund@univie.ac.at) and Alexander Loy (alexander.loy@univie.ac.at)

Running Head: Protein- and lipid-degraders in subarctic marine sediments.

Abstract

Microorganisms in marine sediments play major roles in marine biogeochemical cycles by mineralizing substantial quantities of organic matter from decaying cells. Proteins and lipids are abundant components of necromass, yet microorganisms that degrade them remain understudied. Here, we revealed identities, trophic interactions and genomic features of microorganisms that degraded ^{13}C-labelled proteins and lipids in cold anoxic microcosms with sulfidic subarctic marine sediment. Supplemented proteins and lipids were rapidly fermented to various volatile fatty acids within five days. DNA-stable isotope probing (SIP) suggested *Psychrilyobacter atlanticus* was an important primary degrader of proteins, and *Psychromonas* members were important primary degraders of both proteins and lipids. Closely related *Psychromonas* populations, as represented by distinct 16S rRNA gene variants, differentially utilized either proteins or lipids. DNA-SIP also showed ^{13}C-labeling of various *Deltaproteobacteria* within ten days, indicating trophic transfer of carbon to putative sulfate-reducers. Metagenome-assembled genomes revealed the primary hydrolyzers encoded secreted peptidases or lipases, and enzymes for catabolism of protein or lipid degradation products. *Psychromonas* were prevalent in diverse marine sediments, suggesting they are important players in organic carbon processing *in situ*. Together, this study provides an improved understanding of the metabolic processes and functional partitioning of necromass macromolecules among microorganisms in the seafloor.

Introduction

The majority of marine sediments that underlie the Earth's oceans are dominated by microorganisms mediating heterotrophic processes, which are primarily sustained by "pelagic-benthic" coupling [1]. This is driven by a constant supply of organic matter from planktonic organisms that thrive in the overlying water column and settle with particulate aggregates to the seafloor after their death [2]. The amount and composition of organic matter that reaches the seafloor is strongly dependent on water depth, whereby continental shelves and shallow sediments generally receive greater inputs compared to deep sea sediments [3]. In cold polar regions with lower microbial activities in the water column [4], a large fraction of the phytodetritus from spring bloom-like events reaches the underlying sediments [5]. Additionally, organic matter can originate from autochthonous microbial production [6] and from land-derived sources, which can be substantial in areas close to land and rivers, as well as polar regions subjected to melting [2, 7].

Organic matter in marine sediments is composed of an immense diversity of macromolecules, of which a substantial fraction can be enzymatically degraded and utilised as nutrient and energy sources by microorganisms [2]. Proteins and lipids typically constitute 10% and 5–10% of the organic matter found in marine sediments, respectively [8, 9]. Proteins, peptides and amino acids are important sources of nitrogen [10], particularly in sediment habitats with limited inorganic nitrogen sources [11]. Lipids generally consist of an alkyl chain that is ester- or ether-bound to a polar head group such as phospho-glycerol, or a glycerol that is glycosidically bound to a sugar moiety. They are major components of phytoplankton biomass [12, 13], of which the less labile fraction survives degradation in the water column [14] and is degraded by microorganisms in the underlying sediments [15]. Long-chain fatty acids released from lipid hydrolysis are energy-rich compounds [16], and can be expected to be favourable substrates for microorganisms.

Most macromolecules are too large to be directly imported into cells and must therefore

be degraded, at least partially, outside the cells [17]. Hydrolysis of macromolecules via the activity

of extracellular enzymes is the rate limiting step during organic matter mineralization in

heterotrophic marine sediments [17, 18]. Microorganisms utilize a compositionally and

functionally large diversity of extracellular hydrolases to facilitate the extracellular breakdown of

macromolecules [19]. For example, peptidases in sediments of Arctic fjords in Svalbard had

greater substrate ranges for different peptides and much higher activities (tens to hundreds of

thousands nmol l^{-1} hr^{-1}) than peptidases in the water column [20]. This showed that sediments act

as important biogeochemical hotspots for processing organic macromolecules. Interestingly,

peptidase activity was negatively correlated to amounts of phyto-detritus inputs, whereas lipase

activity was positively correlated [11]. This indicated that peptidases are excreted in times of

nutrient limitation, whereas lipases are excreted when organic matter availability is high [11].

The microbial degradation of organic macromolecules in anoxic marine sediments is a

complex inter-species process involving 'primary degraders' that break down larger

macromolecules into oligomers and monomers for fermentation to alcohols, lactate and/or short-

chain volatile fatty acids (VFAs), which are then mineralized to CH_4, CO_2 and/or H_2 [21]. 'Terminal

oxidizers' of fermentation products prevalent in marine sediments, such as sulfate-reducing

microorganisms (SRM), have been relatively well studied [22–27]. On the other hand, the key

microbial players responsible for the primary hydrolysis of different types of organic matter and

macromolecules are poorly understood. Most studies are based on predictions from genomic

analyses [28–31], whereas experimental evidence linking identities to functions are lacking.

In the present study, we investigated the identities, genomic features, and ecological

interactions of microbial taxa that may play important roles in protein and lipid macromolecule

degradation in subarctic marine sediments. We performed laboratory experiments whereby

sulfidic arctic marine sediments were incubated in microcosms under cold (4 °C), anoxic

conditions and supplemented with either [13]C-labelled proteins or [13]C-labelled lipids. The cold

conditions represented large depth expanses of the oceans and seafloor that are permanently

cold [32]. These microcosms were also incubated under conditions where sulfate reduction was

specifically inhibited, in order to elucidate the roles of SRM. Catabolism and assimilation of ^{13}C-

labelled substrates by the sediment microorganisms was investigated by DNA-based stable

isotope probing (DNA-SIP) and amplicon sequencing of 16S rRNA genes. The degradation of

organic matter was monitored by measuring concentrations of VFAs and sulfate. Genome-

resolved metagenomic analyses were used to predict secreted hydrolytic enzymes and

reconstruct organic matter degradation pathways encoded by the taxa identified by DNA-SIP.

This revealed (i) the identities and foraging strategies of several key protein- and lipid-consuming

bacteria in marine sediments, (ii) that niche partitioning among *Psychromonas* subspecies was

based on differential utilization of proteins and lipids, and (iii) and that SRM of the family

Desulfobacteraceae mainly utilized the macromolecule degradation intermediates, i.e., volatile

fatty acids.

Materials and methods

Sediment incubations

Sediment slurries were produced with sediment (0–30 cm below seafloor) from Greenland

(Nuuk fjord 'station 3', water depth 498 m, August 2013, 64°26′45″N, 52°47′39″W) mixed 1:1 (v/v)

with anoxic artificial seawater [33] containing approximately 28 mM sulfate. For each microcosm,

40 ml of this slurry was distributed into 250 ml serum vials under anoxic conditions in an anoxic

glove-box (nitrogen atmosphere containing approximately 2% hydrogen and 10% CO_2).

Microcosms were sealed with thick rubber stoppers and crimped, and flushed with N_2 after

removing from the anoxic glove-box. All subsequent subsampling was also performed within the

anoxic glove-box, with microcosms placed on ice-packs to minimize warming of incubations.

Triplicate microcosms were either supplemented with a single dose of: i) 300 µg C g^{-1} ^{13}C-labelled

Algal lipid mixture-99 atom% ^{13}C (ISOTEC, Sigma-Aldrich); ii) 300 µg C g^{-1} ^{13}C-labelled Algal

crude protein extract-98 atom% ^{13}C (ISOTEC, Sigma-Aldrich), iii) 300 µg C g^{-1} ^{13}C-labelled Algal

lipid mixture (as above) and the sulfate reduction inhibitor molybdate (28 mM; [34]); v) 300 µg C

g^{-1} ^{13}C-labelled Algal protein mixture (as above) and molybdate (28 mM); or v) were left without

supplementation (no-substrate controls). The microcosms were incubated at 4 °C and were

sampled after 2, 5, 10, 17, 25 and 48 days of incubation. Sediment slurry samples were taken for

analysis of volatile fatty acids and sulfate (1 ml), as well as for DNA-based analyses (0.5 ml).

Samples were stored on pre-cooled ice packs in the anoxic glove box and were transferred

immediately to the –80 °C freezer after sampling. Sampling, processing and molecular biological

analyses of sediment samples from Svalbard, Norway, are presented as Supplementary

information.

Determination of sulfate and volatile fatty acid concentrations

Sulfate concentrations in interstitial water samples of microcosms were determined by

capillary electrophoresis (P/ACETM MDQ molecular characterization system, Beckman Coulter)

with the CEofixTM anions 5 kit (Analis). Standards were produced by dissolving known

concentrations of Na_2SO_4 in artificial seawater. The concentrations of VFAs were determined as

follows: samples were defrosted, vortexed and centrifuged at 10 000 rpm for 10 minutes. 100 µl

of the supernatant was diluted 1:10 (v/v) in Milli-Q water. Syringe filters (Acrodisc IC grade filter,

d=13 mm, Suprapor® membrane with 0.2 µm pore size) were first rinsed with 10 ml Milli-Q water

followed by 0.5 ml of diluted sample to minimize further dilution with the rinsing water. Another

0.5 ml of sample were then filtered and measured by 2-dimensional ion chromatography-mass

spectrometry (IC-IC-MS; Dionex ICS-3000 coupled to an MSQ Plus™, both Thermo Scientific),

equipped with an Ion Pack™ AS 24 as the first column to separate bulk VFAs from chloride, and

an Ion Pack™ AS 11 HC as the second column to separate individual VFAs [35].

DNA extraction

For DNA-SIP gradients and PCR-based amplicon sequencing, DNA was extracted using

a combination of bead beating, cetyltrimethylammonium bromide-containing buffer and phenol-

chloroform extractions. Sediment slurry (0.5 ml) was added to Lysing Matrix E tubes (MP

Biomedicals) and were suspended in 0.675 ml cetyltrimethylammonium bromide containing

extraction buffer, as described previously [36]. The tubes were placed on a rotary shaker (200

rpm) and incubated at 37 °C for 30 minutes. The samples were supplemented with 75 µL of

sodium dodecyl sulfate (20% w/v) and were then incubated for 1 hour at 65 °C (tubes were

inverted every 20 minutes). Samples were subjected to two rounds of bead beating with a speed

setting 6 for 30 s using a FastPrep®-24 bead beater (MP Biomedicals). In between the beat

beating steps, samples were cooled on ice. Debris was pelleted by centrifugation at 6000 g for 10

minutes at 25 °C and supernatant was collected. An equal volume of phenol:chloroform:isoamyl

(25:24:1) alcohol (ROTH) was added to the supernatant and tubes were repeatedly inverted,

followed by centrifugation at 16 000 g for 10 min at 25 °C. The aqueous phase (upper layer) was

transferred into a clean 1.5 ml tube, supplemented with 0.6 volume of 2-propanol and incubated

for 1 hour at 4 °C to precipitate DNA. The precipitate was then pelleted by centrifugation at 16

000 g for 30 min at 4 °C. DNA pellets were washed with 250 μL of 70% (v/v) ethanol, air-dried

and resuspended in 50 μL Tris buffer [10 mM Tris-HCl (pH 8.0)]. For metagenome sequencing,

DNA was extracted from day 0 and from individual sediment microcosm samples at days 5, 17

and 25 (for both protein and lipid amended treatments) using the Ultra Pure Power Soil Kit (MoBio)

and following manufacturers' protocol.

DNA-SIP gradients

CsCl gradients with 5 μg DNA were prepared in a temperature controlled room at 23 °C

according to a previously published protocol [37]. The gradient mixtures were added to

ultracentrifuge tubes (Beckman Coulter) and were centrifuged in an Optima L-100XP

ultracentrifuge (Beckman Coulter) using the VTi 90 rotor for >48 hours at 146286 rcf (44100 rpm)

at 20 °C. After centrifugation, gradients were fractionated into 250 μl fractions by puncturing the

bottom of the tube with a sterile needle and adding water to the gradient tube at the top using a

sterile needle and a syringe pump (World Precision Instruments). The density of collected

fractions was determined by using a digital refractometer (AR 200, Reichert Analytical Instrument)

at 23 °C. The DNA-SIP fractions were considered heavy at densities >1.726 g ml^{-1} and light at

densities <1.720 g ml^{-1} (Supplementary Figure S1). Afterwards, DNA was precipitated with 500

μL sterile PEG 6000 (30% polyethylene glycol 6000 and 1.6 M NaCl) and 1 μL of glycogen (5 μg

ml^{-1}) and subsequently purified as described previously [37]. Bacterial DNA in SIP fractions was

quantified by quantitative PCR (qPCR) using the primers 341F 5′-CCT ACG GGA GGC AGC AG-

3′ and 534R 5′-ATT ACG CGG CGT GCT GGC A-3′. The 20 μL qPCR mix contained 1× IQ^{TM}

SYBR Green Supermix (BIO-RAD), 0.25 μM of each primer and 1 μL of 1:10 diluted DNA from

individual gradient fractions. The program used for thermal cycling on the iCycler thermal cycler

(Bio-Rad) consisted of: 3 min at 95 °C, followed by 39 cycles of 95 °C for 15 s, 60 °C for 30 s and

72 °C for 39 s, and was followed by a melting curve from 60 °C to 95 °C by increments of 0.5 °C

every 5 s.

Amplicon sequencing and analysis

Barcoded 16S rRNA gene amplicons for Illumina MiSeq sequencing were produced using

a previously established two-step PCR approach [38, 39]. Raw reads were processed as

described previously [38, 39]. The identity threshold used for OTU clustering was 97% identity.

Sub-OTU diversity at single-nucleotide resolution in the amplicon data was investigated using

cluster-free filtering [40]. Statistical significance of differences in OTU relative abundances

between treatments was determined using DESeq2 (version 1.10.1) [41] in the R software

environment (R core team, 2019). Additionally, an OTU was considered to be ^{13}C-enriched when

it was significantly more abundant in heavy SIP fractions compared to light fractions obtained

from the same incubation [42], which was also determined using DESeq2. The sequence

abundance of each OTU in the 'heavy' part (>1.726 g ml^{-1}) of the DNA-SIP gradient (numerator)

were compared to sequence abundance in the 'light' part (<1.720 g ml^{-1}) of the DNA-SIP gradient

(denominator). Specific samples used for comparisons are indicated in Supplementary Figure S1.

Only OTUs with more than 5 reads in 5 out of all 32 samples from gradient fractions were

considered during DESeq2 analyses. Results were extracted with the command:

217   `results(cooksCutoff = FALSE, altHypothesis = "greater")`

and were considered statistically significant if: (i) the false-discovery-rate (FDR)-adjusted p-value

was below 0.1, and (ii) the respective OTU was not significantly enriched in the heavy fractions

of the DNA-SIP gradient from the no-substrate incubations. OTU counts were 'rlog' transformed

with DESeq2 and heatmaps of enriched OTUs were created with R software package pheatmap

(version 1.0.8).

The presence of 16S rRNA gene sequences related (>97% identity) to specific organisms

of interest in publicly available amplicon-derived datasets were determined using the IMNGS

server [43], with 100 bp overlap as minimum.

Metagenome sequencing and differential coverage binning

Metagenome libraries were produced with the Nextera XT DNA Library Preparation Kit

(Illumina) and sequencing was performed at the Vienna Biocenter Core Facilities Next Generation

Sequencing facility (Vienna, Austria) on an Illumina Hiseq 2500 using HiSeq V4 chemistry with

the 125 bp paired-end mode. Reads were end trimmed at the first base with a q-score below 10

and reads with less than 50 bp were removed. For assembly, sequence coverage was normalized

across samples using bbnorm with an average read depth of 100 and a minimum read depth of

3 (BBmap version 33.57 http://sourceforge.net/projects/bbmap/). Normalized read files were

assembled with IDBA-UD [44] and SPAdes [45] using default parameters. Reads of each sample

(non-normalized) were mapped to each assembly using BWA [46], and coverage information was

obtained using SAMtools [47]. Differential coverage binning was performed with MetaBAT [48],

MaxBin [49], and CONCOCT [50]. Following the binning programs default parameters, only

contigs with a minimum length of 2500 and 1000 bp were used for binning with MetaBAT and

MaxBin/CONCOCT, respectively. The resulting metagenome-assembled genomes (MAGs)

generated from different assemblies and binning programs were aggregated with DASTool [51],

and de-replicated with dRep using default parameters [52]. MAGs that could be linked to ^{13}C-

incorporating OTUs during protein and lipid hydrolysis by 16S rRNA gene sequence identity were

further analysed. Selected MAGs were evaluated for completion and contamination using

CheckM [53], and classified using the Genome Taxonomy Database (GTDB) release 86 [54] and

GTDB-Tk version 0.1.3 [55]. A genome-based phylogenetic analysis of MAGs, the genome of the

type strain *Psychrilyobacter atlanticus* HAW-EB21 and closely related reference genomes, was

constructed from the concatenated marker protein alignment produced by CheckM [53]. The

phylogenetic tree was built from this concatenated marker alignment with IQ-TREE using

automatic substitution model selection (LG+F+I+G4) [56] and ultrafast bootstrap approximation

with 1000 replicates [57]. The tree was visualized with iTOL [58]. Average nucleotide identities

(ANI) and average amino acid identities (AAI) of selected MAGs and closely related reference

genomes were determined using all-vs-all FastANI 1.1 with default parameters [59] and

CompareM 0.0.23 (aai_wf) with default parameters (https://github.com/dparks1134/CompareM),

respectively.

Genome annotation

MAGs were annotated with the MicroScope annotation platform [60] and using the RAST

server [61]. The annotations of proteins were confirmed by DIAMOND searches [62] against the

NCBI-nr database (e-value 10^{-5}), hidden Markov model-based searches using InterProScan [63]

with the databases Pfam-A [64] and TIGRFAM [65], and online BLASTP searches against the

UniProt database [66]. Possible hydrolytic enzymes were examined for signal peptide sequences

that facilitate secretion from the cytoplasm and transmembrane helices using the Phobius online

server [67] and using PSORTb version 3.0 [68]. Furthermore, peptidases, lipases/esterases and

glycoside hydrolases were additionally compared to the databases MEROPS [69], ESTHER [70]

and CAZY [71], respectively, using DIAMOND searches (e-value $>10^{-5}$, identity ≥30%). Further,

searches were made against the NCBI-nr database using BLASTP [72] for certain proteins of

interest. The selection of potentially catabolic peptidases encoded by genomes and MAGs was

based on two recent studies about extracellular peptidases in marine sediments [29, 73]. In

addition, non-peptidase homologs and peptidases with regulatory functions, e.g., peptidases

involved in membrane protein remodelling, as indicated by MEROPS database descriptions, were

not considered (Supplementary Table S1). Potentially catabolic lipases/esterases were selected

based on the ESTHER database descriptions (Supplementary Table S1).

Phylogenetic analyses of 16S rRNA gene sequences

16S rRNA gene sequences were aligned and classified with the SINA aligner [74], using

the SILVA database release 128 [75]. Full-length sequences of close relatives of 16S rRNA OTUs

were extracted from the SILVA database and combined with the aligned 16S rRNA genes

sequences that were obtained from MAGs. A reference tree was then constructed using FastTree

[76]. Subsequently, 16S rRNA OTUS and partial 16S rRNA genes from MAGs were added to the

reference tree using the EPA algorithm [77] in RAxML [78]. Trees were visualized in iTOL [58].

Statistical analyses

Statistics for comparisons of VFAs and sulfate in treatment series versus controls were

determined with Student's T tests using the function t.test() in the R software environment (R core

team, 2014).

Data availability

All sequence datasets and metagenome-assembled genomes from the DNA-SIP

experiments are available under the NCBI-Genbank Bioproject PRJNA609450. DNA-SIP

amplicon sequencing datasets are available in the NCBI-Genbank Sequence Read Archive under

BioSample accession SAMN14253696 and SRA accessions SRR11221408-SRR11221561.

Metagenomic sequence reads are available in the NCBI-Genbank Sequence Read Archive under

BioSample accession numbers SAMN14421543–SAMN14421548. Metagenome-assembled

genomes are available in the NCBI-Genbank under BioSample accession numbers

SAMN14421524-SAMN14421532. Annotated MAGs and genomes from the MicroScope

annotation platform for key protein- and lipid-degrading organisms, and *Psychrilyobacter*

atlanticus DSM 19335, are publicly available in the MaGe-Microscope server

(https://mage.genoscope.cns.fr/). Amplicon sequencing datasets from Svalbard sediments are

available under NCBI-Genbank Bioproject PRJNA623111 and the BioSample accession numbers

SAMN14538997-SAMN14539076.

Results

Sulfate removal and volatile fatty acid turnover during protein and lipid degradation

Sulfate was largely turned-over after 48 days in all incubation treatments (down to 0.6–

2.4 mM), except in incubations where sulfate reduction was inhibited by molybdate, where it

remained between 18.1–24.2 mM (Figure 1A). Sulfate turnover was fastest between days 5 and

25, and was stimulated by the additions of proteins or lipids. Of all measured VFAs, acetate was

the most prominent, reaching concentrations of over 900 μM in protein amended microcosms on

day 17. Formation of the next most abundant VFA, formate, peaked at around 50 μM in

lipid/molybdate-amended microcosms. Supplementation of proteins to the incubations resulted in

significantly higher concentrations of acetate from days 2–5 compared to no-substrate controls

(Supplementary Table S2). While the stimulation of acetate production from lipid additions was

noticeably more than in no-substrate controls, the differences were not statistically significant

(Supplementary Table S2). Significantly more propionate, butyrate and isobutyrate were

produced in protein- and lipid-amended microcosms, as compared to the no-substrate controls

(mainly between 2–5 days) (Figure 1B, Supplementary Table S2). Molybdate-inhibited

incubations supplemented with proteins or lipids showed increased accumulation of formate,

butyrate, and isobutyrate, but decreased production of acetate and propionate.

^{13}C-labeling of OTUs and sub-OTUs during protein and lipid degradation

To evaluate incorporation of ^{13}C from the added ^{13}C-labelled proteins or lipids into the DNA

of microorganisms, bacterial 16S rRNA genes were quantified by qPCR across individual SIP

fractions (Supplementary Figure S1). Higher relative copy numbers were detected in the 'heavy'

fractions of the gradients (i.e., densities >1.726 g ml^{-1}) from both ^{13}C-protein and ^{13}C-lipid

incubations relative to corresponding no-substrate controls at days 5 and 10. This indicated ^{13}C-

uptake and incorporation into DNA at these early time points. At days 17 and 25, only very small

or no differences in relative 16S rRNA gene copy numbers between gradients from the ^{13}C-

substrate incubations and the no-substrate control incubations were observed.

Microbial community analysis of SIP fractions by 16S rRNA gene amplicon sequencing

identified five OTUs (Clostridia-JTB215 OTU 1, *Psychromonas* OTU 4, *Psychrilyobacter* OTU 5,

Fusibacter OTU 38 and *Photobacterium* OTU 54) that were significantly enriched in heavy SIP

fractions from the ^{13}C-protein incubations at day 5 (Figure 2A; Supplementary Table S3). At day

10, another 19 OTUs were significantly enriched in heavy SIP fractions from incubations with ^{13}C-

protein amendments (Figure 2A). These were affiliated with the classes *Deltaproteobacteria*

(OTUs 2, 19, 36, 44, 67, 183, 232, 892, 4719), *Gammaproteobacteria* (OTUs 80, 128, 205, 285)

and the phyla *Bacteroidetes* (OTUs 124 and 184), *Firmicutes* (OTUs 202 and 4050) and

Marinimicrobia (OTU 123), or were unclassified (OTU 312) (Supplementary Figure S2).

In SIP gradients from ^{13}C-lipid amended incubations at day 5, significant ^{13}C-enrichment

was revealed for only two gammaproteobacterial OTUs, i.e., *Psychromonas* OTU 4 and *Vibrio*

OTU 80 (Figure 2A). An analysis of the microdiversity within *Psychromonas* OTU 4 identified four

Psychromonas sub-OTUs (sub-OTUs 4, 192, 9 and 42) that responded differently in incubations

with protein or lipid amendments. The sub-OTUs 4 and 192 were significantly enriched in heavy

fractions of ^{13}C-lipid incubations (Figure 3A) and were more closely related to each other than to

sub-OTUs 9 and 43 (Figure 3B). Furthermore, the relative abundance of sub-OTUs 4 and 192

increased in the microcosms from ^{13}C-lipid incubations compared to no-substrate control

microcosms (Figure 3C). In contrast, the sub-OTUs 9 and 43 were significantly enriched in heavy

SIP fractions from ^{13}C-protein incubations, and sub-OTU 9 increased in relative abundance in the

microcosms from ^{13}C-protein incubations compared to no-substrate control microcosms (Figure

3C). No sub-OTU microdiversity was detected in any other OTU that was significantly enriched in

^{13}C. At day 10, we identified three additional OTUs enriched in the heavy SIP fractions from ^{13}C-

lipid incubations that were affiliated with the phylum *Firmicutes* (OTUs 1 and 4141) and the class

Deltaproteobacteria (OTU 13310) (Figure 2A).

We also compared the relative abundances of 16S rRNA OTUs among the protein- or

lipid-amended microcosms versus no-substrate control microcosms over time. Here, we restricted

the analysis to OTUs (*n*=26) that were enriched in ^{13}C from the protein- and lipid-amended

treatments (described above). Between days 5 and 10, ten OTUs (1, 2, 4, 5, 19, 54, 80, 285, 4141

and 13310) significantly increased in relative abundances in substrate-amended incubations as

compared to the no-substrate controls (Supplementary Figure S3A and Supplementary Table

S2). Most notably, several OTUs showed relatively large differences in relative abundances

between amended and no-substrate controls, and consequently had high relative abundances

among the overall communities. For instance, *Psychrilyobacter* OTU 5 reached relative

abundances of around 9–10% after 2–5 days in protein-amended microcosms, while staying around 3–4% in no-substrate controls. *Psychromonas* OTU 4 reached relative abundances of around 15–16% between 5–10 days after lipid amendments, while also reaching around 5–7% in the protein-amended microcosms from days 2–10. In contrast, *Psychromonas* OTU 4 stayed below 2.5% in the no-substrate controls over-time. The relatively fast (<10 days) and clear increases in relative abundances of these OTUs due to substrate additions therefore provided evidence for their direct involvement in substrate utilization for growth. On the contrary, the Clostridia-JTB215 OTU 1 also showed large increases in relative abundances from both protein- and lipid-amendments versus no-substrate controls, however, the response was relatively delayed, i.e., such increases developed mainly after 10 days.

Relative abundances of OTUs 1, 2, 67, 4050 and 4141 were significantly lower in microcosms with the sulfate-reduction-inhibitor molybdate than in microcosms without molybdate. This suggests that these OTUs are SRM or dependent on SRM activity.

Genomic evidence for extracellular hydrolysis of proteins and lipids

Samples from day 0, and from days 5, 17 and 25 from the lipid- and protein-amended microcosms were selected for metagenome sequencing. The aim was to recover genomes of organisms that were ^{13}C-labelled from protein or lipid treatments, and to specifically analyse their catabolic potentials for these macromolecules. Overall, nine metagenome-assembled genomes (MAGs) were recovered with >80% completeness and <5% contamination. Two had 16S rRNA sequences identical to bacterial OTUs that incorporated ^{13}C from protein or lipid hydrolysis, i.e., *Psychromonas* MAG GLG-1 and *Clostridia* MAG GPF-1 (Supplementary Figure S2). All MAGs that were affiliated with the genus *Psychrilyobacter* were very incomplete (<20%) (data not shown). Therefore, we analysed the publicly available genome of the type strain *Psychrilyobacter atlanticus* HAW-EB21. This strain was isolated from marine sediments [79], and had a 16S rRNA sequence that was identical to *Psychrilyobacter* OTU 5 that was labelled from protein amendments (Supplementary Figure S2). A *Desulfoluna* MAG GLD-1 was also analysed in detail because it encoded potential to degrade VFAs and/or lipids, although it could not be confidently

linked to any 16S rRNA OTU sequences and it is therefore presented in the Supplementary

results.

Psychromonas MAG GLG-1 was most closely related to *Psychromonas aquimarina* ATCC

BAA-1526 (ANI 80.7% of 41.8% aligned) (Supplementary Table S4) and encoded a 16S rRNA

sequence that was identical to *Psychromonas* sub-OTU 4 (Supplementary Figure S2). *Clostridia*

MAG GPF-1 was only distantly related to any publicly available genome (Supplementary Figure

S4), with an AAI of only 53.4% (of 37% aligned) (ANI was too low to calculate) to the most-related

genome of *Caloranaerobacter azorensis* DSM 13643 (Supplementary Table S4). A short 16S

rRNA fragment (92 bp) was retrieved from *Clostridia* MAG GPF-1 and was linked to the 16S rRNA

OTU 1 by phylogenetic placement into a reference tree (Supplementary Figure S2). The

genomes/MAGs of putative protein and/or lipid-degraders were subsequently analysed for

encoded capabilities to degrade proteins (*P. atlanticus* corresponding to 16S rRNA OTU 5), lipids

(*Psychromonas* MAG GLG-1 corresponding to sub-OTU 4) or both macromolecules (*Clostridia*

MAG GPF-1 corresponding to 16S rRNA OTU 1).

The genome of *Psychrilyobacter atlanticus* HAW-EB21 encodes a variety of peptidases

(n=24), which were likely involved in the breakdown of peptides for further catabolism of amino

acids (Supplementary Table S5). Two of these peptidases (i.e., M3 and M24) encoded signal

peptides for translocation across the cytoplasmic membrane (Figure 4). Overall, the genome of

P. atlanticus encodes an array of predicted peptide (n=19) and amino acid transporters (n=38),

especially when compared to the MAGs of the putative lipid degraders *Psychromonas* MAG GLG-

1 (peptide transporter, n=6; amino acid transporter, n=14) (Supplementary Table S5).

Furthermore, the genome of *P. atlanticus* encoded for proton-dependent peptide transportation

(Supplementary Table S5).

The lipid-metabolizing capability of *Psychromonas* MAG GLG-1 was supported by the

presence of genes for an extracellular phospholipase D/alkaline phosphatase and a

carboxylesterase with a signal peptide sequence, which corresponded to the ESTHER family

Carboxylesterase type-B (Supplementary Table S5). The long chain fatty acids that resulted from

extracellular degradation of acyl chain esters were likely taken-up by cells via two predicted long-

chain fatty acid transporters (Supplementary Table S5). Predicted glycerol transporters were not

encoded in the genome of *Psychromonas* MAG GLG-1. Because our study revealed

Psychromonas spp. to be potentially important lipid and protein degraders, we also investigated

available genomes and/or MAGs (*n*=16) from publicly available databases in order to examine

the prevalence of the encoded capacity to utilize extracellular proteins or lipids via extracellular

hydrolases (Supplementary Table S6). This identified predicted secreted phospholipases in most

genomes (*n*=10), while predicted secreted proteases/peptidases were restricted to 4 genomes

(Supplementary Figure S6).

Clostridia MAG GPF-1 encoded the largest number of peptidases (*n*=29) of all analysed

MAGs/genomes (Supplementary Table S5). Three of these peptidases (M20, M24 and S8

peptidase families) have predicted secretion signal peptides and might be excreted to the

extracellular space via the SEC pathway (Figure 4). Similar to *P. atlanticus*, *Clostridia* MAG GPF-

1 encoded a broad array of predicted peptide (*n*=14) and amino acid (*n*=24) transporters. In

accordance with the DNA-SIP results, *Clostridia* MAG GPF-1 also has the potential for catabolic

breakdown of lipids. The MAG encoded one potentially secreted catabolic esterase/lipase that

correspond to the ESTHER family Bacterial_EstLip_FamX, and two predicted membrane-bound

esterases/lipases that correspond to the ESTHER families Bacterial_EstLip_Fam and

Carboxylesterase, type B (Supplementary Table S5). The presence of genes for

esterases/lipases in the genome of *Clostridia* MAG GPF-1 indicated the potential of this MAG to

hydrolyse ester bonds between glycerol and fatty acids. Because genes encoding for known fatty

acid transport and catabolism were absent from the genome of *Clostridia* MAG GPF-1, this

bacterium might instead utilize glycerol. This was supported by an encoded glycerol-3-phosphate

transporter and a putative glycerol facilitator protein, which were also encoded directly adjacent

to a glycerol kinase, which catalyses the first step of the cytoplasmic catabolism of this substrate

(Supplementary Table S5).

***Intracellular catabolism of protein and lipid degradation intermediates**

The genome of *P. atlanticus* encoded 22 cytoplasmic peptidases with possible catabolic

roles (Supplementary Table S5). In comparison, the genomes of the putative lipid degrader, i.e.,

Psychromonas MAG GLG-1 encoded only 7 intracellular peptidases. In this study, we focused on

amino acid degradation processes that form glutamate and/or keto-acids for central carbon

metabolism and that could therefore explain incorporation of ^{13}C-carbon from the ^{13}C-labelled

proteins into DNA, as well as the development of VFAs in our incubations (Figure 4). The genome

of *P. atlanticus* encoded several aminotransferases and other amino acid degrading enzymes

(i.e., alanine dehydrogenases, cystathionine gamma-lyase, serine/threonine ammonia-lyase,

tryptophanase and methionine gamma-lyase) (Supplementary Table S5). These can convert

amino acids to the keto-acids pyruvate and oxobutanoate, which could then be fermented to

formate, and either acetate or propionate, respectively (Figure 4).

Psychromonas MAG GLG-1 encoded all enzymes required for beta-oxidation of fatty

acids, including a putative multienzyme complex that contains the two enzymes enoyl-CoA

hydratase/isomerase and 3-hydroxyacyl-CoA dehydrogenase (Supplementary Table S5).

Through beta-oxidation, even- and odd-chain fatty acids can be converted to propionyl- and

acetyl-CoA, which can be further fermented to acetate and propionate, respectively (Figure 4). In

comparison to *P. atlanticus* and *Clostridia* MAG GPF-1, which encoded strict fermentative

metabolisms, *Psychromonas* MAG GLG-1 also encoded a complete oxidative TCA cycle, which

enables a respiratory conversion of acetyl-CoA to CO_2 (Figure 4). When oxygen is available,

Psychromonas MAG GLG-1 might use one of the two encoded high affinity oxidases for

respiration, i.e., cbb3-type cytochrome *c* oxidase and cytochrome *bd*–type oxidase

(Supplementary Table S5). In oxygen-depleted sediments like in our microcosm incubations,

Psychromonas GLG-1 might ferment, or may respire using nitrate (dissimilatory nitrate reduction

to ammonium, DNRA) or fumarate as terminal electron acceptors (Supplementary Table S5).

Genes necessary for the intracellular degradation of glycerol were absent from *Psychromonas*

MAG GLG-1 (Supplementary Table S5). Furthermore, despite the presence of genes for several

extracellular glycosyl hydrolases in the genome of *Psychromonas* MAG GLG-1 and a complete

glycolysis pathway (Supplementary Table S5), enzymes for feeding galactose or glycerol

(subcomponents of Spirulina-derived galactosyl diglycerides), into glycolysis, were not identified.

The *Clostridia* MAG GPF-1 encoded the largest set of intracellular peptidases (n=26) of

all investigated genomes (Supplementary Table S5). The fermentation of amino acids by

Clostridia GPF-1 to propionate, acetate or formate, i.e., via aminotransferases and other amino

acid degrading enzymes (i.e., L-threonine 3-dehydrogenase, cystathionine β-synthase,

cystathionine gamma-lyase, serine/threonine ammonia-lyase and methionine gamma-lyase) was

mostly similar to the predicted routes used by *P. atlanticus*, although genes for lysine fermentation

were not present and the glutamate degradation pathways were incomplete (Supplementary

Table S5). In addition, the *Clostridia* MAG GPF-1 was the only MAG that encoded enzymes that

initiate valine degradation, i.e., two branched-chain amino acid transferases and the gene 2-keto-

isovalerate dehydrogenase that is part of the branched-chain α-keto acid dehydrogenase complex

(Supplementary Table S5). Besides the potential for amino acid degradation, *Clostridia* MAG

GPF-1 also encoded all genes necessary for the conversion of glycerol and glycerol-3-phosphate

to pyruvate (Supplementary Table S5), which may help explain with the ^{13}C enrichment of 16S

rRNA OTU 1 in ^{13}C-lipid amended incubations (Figure 2).

Environmental distributions of key taxa

To explore fine-scale depth distributions of the main primary-hydrolysing *Psychromonas*

and *Psychrilyobacter* species in sulfidic arctic marine sediments, we examined 16S rRNA-gene

and -cDNA derived sequence datasets from sediments of Smeerenbergfjord, Svalbard

(Supplementary Figure S6). This showed that *Psychromonas* sequences reached 1.7% and 1.9%

relative abundances in 16S rRNA-gene and -cDNA libraries in 0–1 centimetres below seafloor

(cmbsf) samples from Station GK, respectively. In a nearby core from Station J, they also reached

1.2% relative abundances in both 16S rRNA-gene and -cDNA libraries in 0–1 cmbsf samples. At

both sites, relative abundances of *Psychromonas* 16S rRNA-genes and -transcripts were very

low under 3 cmbsf (Supplementary Figure S6). Sequences from the *Psychrilyobacter* reached

only 0.3% in 0–1 cmbsf from Station J samples and comprised maximally only 0.05% of the 16S

rRNA transcripts in the same sample (results not shown). *Psychrilyobacter* sequences were under

0.03% in all other samples deeper than 1 cmbsf. Further, to gauge their prevalence in other

marine sediments, we examined sequences that were similar (>97% identity) to the

Psychromonas and *Psychrilyobacter* OTUs recovered in this study to sequences in publicly

available 16S rRNA gene datasets. This showed that sequences related to these organisms were

widely distributed in sediment sites around the worlds oceans (Supplementary Figure S7). For

example, *Psychromonas* sequences reached a maximum of 4% relative abundance in seagrass-

associated sediments (Nantucket Sound, USA).

Discussion

Proteins and lipids comprise a significant fraction (up to 10% each) of organic matter in

marine sediments [8, 9], and therefore represent major nutrient and energy sources for sediment

microbiomes. Our experiment showed that supplemented proteins and lipids were actively utilized

by microorganisms within 5 days of incubation. Both protein and lipid amendments induced rapid

accumulation of several VFAs, demonstrating primary fermentation of both macromolecules.

DNA-SIP further showed that ^{13}C-carbon from protein and lipid catabolism was incorporated into

the DNA of specific taxa within 5 days. This was also paralleled by sharp increases in the relative

abundances by many of the same taxa in the corresponding microcosms. We therefore reasoned

that taxa that were ^{13}C-labelled and stimulated to higher relative abundances (>1%) within the

first 5 days represented the main primary hydrolysers and fermenters of proteins or lipids. Thus,

the following discussion especially focuses on these primary hydrolysing taxa.

Overall, more diverse OTUs incorporated ^{13}C-carbon from proteins compared to lipids,

especially after 10 days of incubation, possibly due to a combination of: i) higher proportions and

bioavailability of proteins in sediments [8, 9], ii) greater nutritional value of proteins (e.g., nitrogen),

and/or iii) a greater biodegradability of proteins as compared to lipids [80]. Sediment

microorganisms are also known to efficiently salvage organic nitrogen sources such as amino

acids in times of low nutrient supply [11]. Therefore, some microorganisms may have salvaged

free peptides or amino acids released from the primary hydrolysers and became ^{13}C-labelled.

The *Psychromonas* 16S rRNA gene OTU 4 was notable in this study because our DNA-

SIP results showed that it was a prominent degrader of both lipids and proteins. Interestingly, we

could distinguish four sub-OTUs that each had distinct preferences for either lipids or proteins.

Thus, this demonstrated that closely related populations of *Psychromonas* in marine sediments

have very different preferences for macromolecules that require completely different catabolic

machinery. From the metagenomic analyses, the *Psychromonas* MAG GLG-1, which represented

the lipid-degrading sub-OTU 4 population, indeed had gene content that supported its capacity to

digest lipids and catabolize fatty acids. For instance, it encoded a secreted esterase/lipase to

digest lipids from the extracellular environment, as well as long-chain fatty acid transporters and

a beta-oxidation pathway to import and catabolize the fatty acids (Figure 4). Although we did not

recover any *Psychromonas* MAGs representative of the protein-degrading populations,

comparative genomic analyses of publicly available *Psychromonas* genomes and MAGs showed

predicted secreted lipases and/or peptidases are common among this genus. Diverse members

of this genus therefore have unique potentials to utilize lipids or proteins.

Our microbial community profiling showed that *Psychromonas* were relatively abundant

and active (based on 16S rRNA expression) in shallow sediments (<3 cmbsf) from arctic Svalbard.

Psychromonas were previously shown to be highly correlated with fluxes of fresh phyto-detritus

to surface sediments of the arctic Laptev Sea [81]. Together with our experimental and genomic

analyses, this suggests *Psychromonas* play an active role in the turn-over of 'fresh' and labile

organic matter delivered to surface marine sediments from the water column. This is inline with

previous experiments that showed *Psychromonas* were important players in the initial degradation

of whole Spirulina (cyanobacterial) necromass in arctic marine sediment [23]. Our study therefore

suggests that *Psychromonas* populations thrive in surface sediments by utilizing abundant

macromolecules that represent fresh and labile organic matter, e.g., lipids and/or proteins.

We further showed that *Psychrilyobacter* OTU 5 was one of the most prominent protein

and/or amino acid degraders in our experiment. Members of this genus were also shown to be

important degraders of whole Spirulina necromass in marine sediment experiments [23, 33]. Here,

we therefore deciphered their role in bulk organic matter degradation by showing that they are

efficient degraders of detrital protein/peptides. Correspondingly, genomic analyses showed that

the *Psychrilyobacter atlanticus* type strain HAW-EB21 [79], which had 100% 16S rRNA sequence

identity to OTU 5, encoded various routes for the degradation of peptides and for the subsequent

fermentation of amino acids (Figure 4) [79]. Their overall contributions to protein degradation in

marine sediments may, however, be less than the *Psychromonas* because they were not

abundant members of most sedimentary microbiomes analysed.

The Clostridia MAG GPF-1 (16S rRNA-OTU 1), had less than 54% AAI to any available

genome, and therefore our data represents the first insights into the potential metabolism of this

uncultured group. Overall, our analyses suggested these organisms are versatile nutrient utilizers.

The Clostridia OTU 1 population was ^{13}C-labelled in SIP analyses from both ^{13}C-protein and -lipid

incubations, and its relative abundances were also boosted by the additions of both substrates,

although to a much lesser extent than the *Psychromonas* or *Psychrilyobacter*. Our genomic

analyses indicated the potential to utilize peptides, amino acids and glycerol (Figure 4), thereby

highlighting its versatility. Further, OTU 1 also increased in no-substrate controls over time,

suggesting it either utilized natural organic matter in the microcosms, or fermentation products

produced from the degradation of the natural organic matter by other members of the community.

In addition to the relatively abundant taxa described above that were ^{13}C-enriched and

that could be linked to genomes or MAGs, we also identified ^{13}C-enrichment in several low

abundance taxa (<1% relative abundances), i.e., members of the genera *Photobacterium* (OTU

54), *Vibrio* (OTU 80) and *Fusibacter* (OTU 38). These taxa may have therefore played a lesser

role in the primary hydrolysis of the macromolecules and are therefore discussed briefly in the

Supplementary Discussion.

After the main primary degradation processes and increased formation of fermentation

products (i.e., VFAs) in the first 5 days, ongoing depletion of sulfate and inhibition of sulfate

depletion by molybdate indicated SRM were active. Various taxa including *Deltaproteobacteria*

related to known SRM became ^{13}C-labelled at day 10, which suggested transfer of carbon from

the added macromolecules to SRM. This was likely via oxidation of the VFAs by SRM, because

deltaproteobacterial SRM are known to be important oxidizers of fermentation products in marine

sediments [82–84]. The detection of more ^{13}C-labelled putative SRM of the *Deltaproteobacteria*

from protein incubations than from lipid incubations may have reflected the more diverse taxa

involved in primary degradation of the proteins, e.g., through more diverse metabolic interactions.

We also hypothesize the additional nitrogen supply from the proteins may have facilitated the

growth of more diverse *Deltaproteobacteria*, since none of the VFAs appeared to be limiting from

either protein or lipid treatments.

In conclusion, this study provided new insights into the identities and functions of various

protein- and lipid-degrading microorganisms in cold marine sediments and indicates

Psychromonas spp*.* are prominent players in the utilization of fresh detrital protein and/or lipid

macromolecules. The activity of these primary hydrolysers also facilitates relatively rapid transfer

of carbon and energy among trophic levels within the sediment microbial community.

Acknowledgements

The authors would like to thank the crew of the R/V Sanna 2013 summer sampling campaign,

which was funded by the Arctic Research Centre, Aarhus University, and Kasper Kjeldsen, Hans

Røy and Marit-Solveig Seidenkranz for providing the sediment samples. We thank captain (Stig

Henningsen) and first mate of MS Farm for their assistance during sampling of Svalbard

sediments in 2017, as well as the Kings Bay Marine Laboratory and the AWIPEV Arctic Research

Base in Ny-Ålesund for assistance and laboratory space. Furthermore the authors want to thank

Carmen Czepe from the Vienna Biocenter Core Facilities Next Generation Sequencing facility for

sequencing of metagenomic libraries on the Illumina Hiseq2500 instrument. This work was

financially supported by the Austrian Science Fund (P29426-B29 to KW; P25111-B22 to AL).

The authors declare no conflict of interest.

Figure legends

Figure 1. Concentration profiles of sulfate and fermentation products in subarctic marine

sediment microcosms. Concentrations of sulfate (A) and volatile fatty acids (B) were determined

in anoxic incubations with ^{13}C-protein or ^{13}C-lipids, and with or without the sulfate reduction

inhibitor molybdate. Control microcosms were incubated without organic substrates and without

molybdate.

Figure 2. 16S rRNA gene OTUs enriched in heavy ^{13}C-DNA-SIP gradient fractions. Heatmaps

showing the relative abundance of 16S rRNA gene OTUs in DNA-SIP gradient fractions from

anoxic sediment incubations amended with ^{13}C-proteins or ^{13}C-lipids and from unamended control

incubations. Significant (FDR-adjusted p-value was <0.1) ^{13}C-enrichment was determined by

differential abundance analysis between high density (^{13}C-enriched, >1.726 g ml^{-1}, pink blocks)

and low density (^{13}C-free, <1.720 g ml^{-1}, turquoise blocks) DNA, and is indicated by filled circles.

16S rRNA genes that were matched to 16S rRNA gene sequences of genomes/MAGs are

indicated in bold.

Figure 3. Differential ^{13}C-labeling and response of *Psychromonas* sub-OTUs in protein- and

lipid-amended sediment microcosms. (A) Heatmap showing the relative abundance of four

sub-OTUs of *Psychromonas* 16S rRNA OTU 4 in DNA-SIP gradients. Significant ^{13}C-enrichment

was determined by differential abundance analysis between high density (^{13}C-enriched) and low

density (^{12}C-enriched) DNA-SIP fractions. Sub-OTU 4, which was identical to the 16S rRNA gene

sequence of *Psychromonas* MAG GLG-1, is indicated in bold. (B) Phylogeny of *Psychromonas*

sub-OTUs. The sequences were aligned with MAFFT (Katoh et al., 2002) and the tree was

calculated with FastTree (Price et al., 2010). (C) Changes in relative abundance of *Psychromonas*

sub-OTUs in protein-amended, lipid-amended, and unamended sediment microcosms with or

without the sulfate reduction inhibitor molybdate.

Figure 4. Genome-inferred metabolic models of protein- and lipid-degrading bacteria in

arctic marine sediment. Schematic depiction of predicted metabolic properties of

Psychrilyobacter atlanticus, *Clostridia* MAG GPF-1, and *Psychromonas* MAG GLG-1, which

correspond to the 16S rRNA OTUs 5, 1, and 4, respectively. The models indicate macromolecule

degradation processes and enzymes that may explain the observed biogeochemical patterns and

the ^{13}C-enrichment of corresponding 16S rRNA OTUs in the DNA-SIP incubations. Enzymes that

are indicated in pale colour were not found in the respective genome/MAG. SS, Secretion system;

TCA, tricarboxylic acid cycle; TAT, Twin-arginine translocation pathway; SEC, Sec secretion

system; LIT, process has physiological evidence but the involved enzymes are not identified yet.

References

1.   Hop H, Pearson T, Hegseth EN, Kovacs KM, Wiencke C, Kwasniewski S, et al. The marine

ecosystem of Kongsfjorden, Svalbard. *Polar Res* 2002; **21**: 167–208.

2.   Arndt S, Jørgensen BB, LaRowe DE, Middelburg JJ, Pancost RD, Regnier P. Quantifying

the degradation of organic matter in marine sediments: A review and synthesis. *Earth-Sci*

Rev 2013; **123**: 53–86.

3.   Dunne JP, Sarmiento JL, Gnanadesikan A. A synthesis of global particle export from the

surface ocean and cycling through the ocean interior and on the seafloor. *Global*

Biogeochem Cycles 2007; **21**: 1–16.

4.   Christian JR, Karl DM. Bacterial ectoenzymes in marine waters: Activity ratios and

temperature responses in three oceanographic provinces. *Limnol Oceanogr* 1995; **40**:

1042–1049.

5. Fabiano M, Pusceddu A. Total and hydrolizable particulate organic matter (carbohydrates,

proteins and lipids) at a coastal station in Terra Nova Bay (Ross Sea, Antarctica). *Polar Biol*

1998; **19**: 125–132.

6. Bradley JA, Amend JP, LaRowe DE. Necromass as a Limited Source of Energy for

Microorganisms in Marine Sediments. *J Geophys Res Biogeosci* 2018; **123**: 577–590.

7. Wehrmann LM, Formolo MJ, Owens JD, Raiswell R, Ferdelman TG, Riedinger N, et al. Iron

and manganese speciation and cycling in glacially influenced high-latitude fjord sediments

(West Spitsbergen, Svalbard): Evidence for a benthic recycling-transport mechanism.

Geochim Cosmochim Acta 2014; **141**: 628–655.

8. Burdige DJ. Preservation of organic matter in marine sediments: controls, mechanisms,

and an imbalance in sediment organic carbon budgets? *Chem Rev* 2007; **107**: 467–485.

9. Hedges JI, Oades JM. Comparative organic geochemistries of soils and marine sediments.

Org Geochem 1997; **27**: 319–361.

10. McCarthy M, Pratum T, Hedges J, Benner R. Chemical composition of dissolved organic

nitrogen in the ocean. *Nature* 1997; **390**: 150–154.

11. Vetter YA, Deming JW. Extracellular enzyme activity in the Arctic Northeast Water polynya.

Mar Ecol Prog Ser 1994; **114**: 23–34.

12. Parsons TR, Stephens K, Strickland JDH. On the Chemical Composition of Eleven Species

of Marine Phytoplankters. *J Fish Res Board Can* 1961; **18**: 1001–1016.

13. Hudson BJ, Karis IG. The lipids of the alga Spirulina. *J Sci Food Agric* 1974; **25**: 759–763.

14. Wakeham SG, Lee C, Farrington JW, Gagosian RB. Biogeochemistry of particulate organic

matter in the oceans: results from sediment trap experiments. *Deep Sea Res A* 1984; **31**:

509–528.

15. Harvey HR, Rodger Harvey H, Fallon RD, Patton JS. The effect of organic matter and

oxygen on the degradation of bacterial membrane lipids in marine sediments. *Geochim*

Cosmochim Acta 1986; **50**: 795–804.

16. Sousa DZ, Smidt H, Alves MM, Stams AJM. Ecophysiology of syntrophic communities that

degrade saturated and unsaturated long-chain fatty acids. *FEMS Microbiol Ecol* 2009; **68**:

257–272.

17. Meyer-Reil L-A. Ecological Aspects of Enzymatic Activity in Marine Sediments.

Brock/Springer Series in Contemporary Bioscience. 1991. pp 84–95.

18. Beulig F, Røy H, Glombitza C, Jørgensen BB. Control on rate and pathway of anaerobic

organic carbon degradation in the seabed. *Proc Natl Acad Sci U S A* 2018; **115**: 367–372.

19. Arnosti C. Microbial extracellular enzymes and the marine carbon cycle. *Ann Rev Mar Sci*

2011; **3**: 401–425.

20. Arnosti C. Contrasting patterns of peptidase activities in seawater and sediments: An

example from Arctic fjords of Svalbard. *Mar Chem* 2015; **168**: 151–156.

21. Muyzer G, Stams AJM. The ecology and biotechnology of sulphate-reducing bacteria. *Nat*

Rev Microbiol 2008; **6**: 441–454.

22. Webster G, Watt LC, Rinna J, Fry JC, Evershed RP, Parkes RJ, et al. A comparison of

stable-isotope probing of DNA and phospholipid fatty acids to study prokaryotic functional

diversity in sulfate-reducing marine sediment enrichment slurries. *Environ Microbiol* 2006;

8: 1575–1589.

23. Müller AL, Pelikan C, de Rezende JR, Wasmund K, Putz M, Glombitza C, et al. Bacterial

interactions during sequential degradation of cyanobacterial necromass in a sulfidic arctic

marine sediment. *Environ Microbiol* 2018; **20**: 2927–2940.

24. Knoblauch C, Sahm K, Jørgensen BB. Psychrophilic sulfate-reducing bacteria isolated from

permanently cold arctic marine sediments: description of *Desulfofrigus oceanense* gen.

nov., sp. nov., *Desulfofrigus fragile* sp. nov., *Desulfofaba gelida* gen. nov., sp. nov.,

Desulfotalea psychrophila gen. nov., sp. nov. and *Desulfotalea arctica* sp. nov. *Int J Syst*

Bacteriol 1999; **49**: 1631–1643.

25. Sahm K, Knoblauch C, Amann R. Phylogenetic affiliation and quantification of psychrophilic

sulfate-reducing isolates in marine Arctic sediments. *Appl Environ Microbiol* 1999; **65**:

3976–3981.

26. Na H, Lever MA, Kjeldsen KU, Schulz F, Jørgensen BB. Uncultured Desulfobacteraceae

and Crenarchaeotal group C3 incorporate 13C-acetate in coastal marine sediment. *Environ*

Microbiol Rep 2015; **7**: 614–622.

27. Wasmund K, Mußmann M, Loy A. The life sulfuric: microbial ecology of sulfur cycling in
marine sediments. *Environ Microbiol Rep* 2017; **9**: 323–344.

28. Lloyd KG, Schreiber L, Petersen DG, Kjeldsen KU, Lever MA, Steen AD, et al. Predominant
archaea in marine sediments degrade detrital proteins. *Nature* 2013; **496**: 215–218.

29. Zinke LA, Glombitza C, Bird JT, Røy H, Jørgensen BB, Lloyd KG, et al. Microbial organic
matter degradation potential in Baltic Sea sediments influenced by depositional conditions
and in situ geochemistry. *Appl Environ Microbiol* 2018; **85**: e02164–18.

30. Orsi WD, Richards TA, Francis WR. Predicted microbial secretomes and their target
substrates in marine sediment. *Nat Microbiol* 2018; **3**: 32–37.

31. Baker BJ, Lazar CS, Teske AP, Dick GJ. Genomic resolution of linkages in carbon,
nitrogen, and sulfur cycling among widespread estuary sediment bacteria. *Microbiome*
2015; **3**: 14.

32. Boyer T, Levitus S, Garcia H, Locarnini RA, Stephens C, Antonov J. Objective analyses of
annual, seasonal, and monthly temperature and salinity for the World Ocean on a 0.25 grid.
International Journal of Climatology: A Journal of the Royal Meteorological Society 2005;
25: 931–945.

33. Graue J, Engelen B, Cypionka H. Degradation of cyanobacterial biomass in anoxic tidal-flat
sediments: a microcosm study of metabolic processes and community changes. *ISME J*
2012; **6**: 660–669.

34. Newport PJ, Nedwell DB. The mechanisms of inhibition of *Desulfovibrio* and
Desulfotomaculum species by selenate and molybdate. *J Appl Bacteriol* 1988; **65**: 419–
423.

35. Glombitza C, Pedersen J, Røy H, Jørgensen BB. Direct analysis of volatile fatty acids in
marine sediment porewater by two-dimensional ion chromatography-mass spectrometry.
Limnol Oceanogr Methods 2014; **12**: 455–468.

36. Dumont MG, Radajewski SM, Miguez CB, McDonald IR, Murrell JC. Identification of a
complete methane monooxygenase operon from soil by combining stable isotope probing
and metagenomic analysis. *Environ Microbiol* 2006; **8**: 1240–1250.

37. Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW, et al. DNA

767 stable-isotope probing. *Nat Protoc* 2007; **2**: 860–866.

768 38. Pelikan C, Herbold CW, Hausmann B, Müller AL, Pester M, Loy A. Diversity analysis of

769 sulfite- and sulfate-reducing microorganisms by multiplex *dsrA* and *dsrB* amplicon

770 sequencing using new primers and mock community-optimized bioinformatics. *Environ*

771 *Microbiol* 2016; **18**: 2994–3009.

772 39. Herbold CW, Pelikan C, Kuzyk O, Hausmann B, Angel R, Berry D, et al. Corrigendum: A

773 flexible and economical barcoding approach for highly multiplexed amplicon sequencing of

774 diverse target genes. *Front Microbiol* 2016; **7**: 870.

775 40. Tikhonov M, Leach RW, Wingreen NS. Interpreting 16S metagenomic data without

776 clustering to achieve sub-OTU resolution. *ISME J* 2015; **9**: 68–80.

777 41. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

778 seq data with DESeq2. *Genome Biol* 2014; **15**: 550.

779 42. Orsi WD, Smith JM, Liu S, Liu Z, Sakamoto CM, Wilken S, et al. Diverse, uncultivated

780 bacteria and archaea underlying the cycling of dissolved protein in the ocean. *ISME J* 2016;

781 **10**: 2158–2173.

782 43. Lagkouvardos I, Joseph D, Kapfhammer M, Giritli S, Horn M, Haller D, et al. IMNGS: A

783 comprehensive open resource of processed 16S rRNA microbial profiles for ecology and

784 diversity studies. *Sci Rep* 2016; **6**: 33721.

785 44. Peng Y, Leung HCM, Yiu SM, Chin FYL. IDBA-UD: a de novo assembler for single-cell and

786 metagenomic sequencing data with highly uneven depth. *Bioinformatics* 2012; **28**: 1420–

787 1428.

788 45. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new

789 genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*

790 2012; **19**: 455–477.

791 46. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

792 *Bioinformatics* 2009; **25**: 1754–1760.

793 47. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

794 Alignment/Map format and SAMtools. *Bioinformatics* 2009; **25**: 2078–2079.

795 48. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. *PeerJ* 2015; **3**: e1165.

49. Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover

genomes from multiple metagenomic datasets. *Bioinformatics* 2016; **32**: 605–607.

50. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning

metagenomic contigs by coverage and composition. *Nat Methods* 2014; **11**: 1144–1146.

51. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of

genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat*

Microbiol 2018; **3**: 836–843.

52. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic

comparisons that enables improved genome recovery from metagenomes through de-

replication. *ISME J* 2017; **11**: 2864–2868.

53. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res 2015; **25**: 1043–1055.

54. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A

standardized bacterial taxonomy based on genome phylogeny substantially revises the tree

of life. *Nat Biotechnol* 2018; **36**: 996–1004.

55. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* 2019.

56. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015;

32: 268–274.

57. Minh BQ, Nguyen MAT, von Haeseler A. Ultrafast approximation for phylogenetic

bootstrap. *Mol Biol Evol* 2013; **30**: 1188–1195.

58. Letunic I, Bork P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display

and annotation. *Bioinformatics* 2007; **23**: 127–128.

59. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018;

9: 5114.

60. Vallenet D, Calteau A, Cruveiller S, Gachet M, Lajus A, Josso A, et al. MicroScope in 2017:

an expanding and evolving integrated resource for community expertise of microbial

genomes. *Nucleic Acids Res* 2017; **45**: D517–D528.

61. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server:

rapid annotations using subsystems technology. *BMC Genomics* 2008; **9**: 75.

62. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat*

Methods 2015; **12**: 59–60.

63. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-

scale protein function classification. *Bioinformatics* 2014; **30**: 1236–1240.

64. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the

protein families database. *Nucleic Acids Res* 2013; **42**: D222–D230.

65. Haft DH, Selengut JD, White O. The TIGRFAMs database of protein families. *Nucleic Acids*

Res 2003; **31**: 371–373.

66. UniProt Consortium T. UniProt: the universal protein knowledgebase. *Nucleic Acids Res*

2018; **46**: 2699.

67. Kall L, Krogh A, Sonnhammer ELL. Advantages of combined transmembrane topology and

signal peptide prediction--the Phobius web server. *Nucleic Acids Res* 2007; **35**: W429–

W432.

68. Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, et al. PSORTb 3.0: improved protein

subcellular localization prediction with refined localization subcategories and predictive

capabilities for all prokaryotes. *Bioinformatics* 2010; **26**: 1608–1615.

69. Rawlings ND. MEROPS: the peptidase database. *Nucleic Acids Res* 2000; **28**: 323–325.

70. Lenfant N, Hotelier T, Velluet E, Bourne Y, Marchot P, Chatonnet A. ESTHER, the

database of the α/β-hydrolase fold superfamily of proteins: tools to explore diversity of

functions. *Nucleic Acids Res* 2013; **41**: D423–9.

71. Lombard V, Ramulu HG, Drula E, Coutinho PM, Henrissat B. The carbohydrate-active

enzymes database (CAZy) in 2013. *Nucleic Acids Res* 2014; **42**: D490–D495

72. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST

and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*

1997; **25**: 3389–3402.

73. Steen AD, Kevorkian RT, Bird JT, Dombrowski N, Baker BJ, Hagen SM, et al. Kinetics and

Identities of Extracellular Peptidases in Subsurface Sediments of the White Oak River

Estuary, North Carolina. *Appl Environ Microbiol* 2019; **85**.

74. Pruesse E, Peplies J, Glöckner FO. SINA: accurate high-throughput multiple sequence

alignment of ribosomal RNA genes. *Bioinformatics* 2012; **28**: 1823–1829.

75. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: improved data processing and web-based tools. *Nucleic Acids*

Res 2013; **41**: D590–6.

76. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for

Large Alignments. *PLoS One* 2010; **5**: e9490.

77. Berger SA, Krompass D, Stamatakis A. Performance, accuracy, and Web server for

evolutionary placement of short sequence reads under maximum likelihood. *Syst Biol* 2011;

60: 291–302.

78. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.

79. Zhao J-S, -S. Zhao J, Manno D, Hawari J. Psychrilyobacter atlanticus gen. nov., sp. nov., a

marine member of the phylum Fusobacteria that produces H2 and degrades nitramine

explosives under low temperature conditions. *Int J Syst Evol Microbiol* 2009; **59**: 491–497.

80. Wakeham SG, Canuel EA. Degradation and Preservation of Organic Matter in Marine

Sediments. *The Handbook of Environmental Chemistry*. 2006. pp 295–321.

81. Bienhold C, Boetius A, Ramette A. The energy–diversity relationship of complex bacterial

communities in Arctic deep-sea sediments. *ISME J* 2011; **6**: 724–732.

82. Glombitza C, Jaussi M, Røy H, Seidenkrantz M-S, Lomstein BA, Jørgensen BB. Formate,

acetate, and propionate as substrates for sulfate reduction in sub-arctic sediments of

Southwest Greenland. *Front Microbiol* 2015; **6**: 846.

83. Finke N, Vandieken V, Jørgensen BB. Acetate, lactate, propionate, and isobutyrate as

electron donors for iron and sulfate reduction in Arctic marine sediments, Svalbard. *FEMS*

Microbiology Ecology . 2007. , **59**: 10–22

84. Glombitza C, Egger M, Røy H, Jørgensen BB. Controls on volatile fatty acid concentrations

in marine sediments (Baltic Sea). *Geochim Cosmochim Acta* 2019; **258**: 226–241.

Figure 1. Concentration profiles of sulfate and fermentation products in subarctic marine sediment microcosms. Concentrations of sulfate (A) and volatile fatty acids (B) were determined in anoxic incubations with ^{13}C-protein or ^{13}C-lipids, and with or without the sulfate reduction inhibitor molybdate. Control microcosms were incubated without organic substrates and without molybdate. Error bars indicate standard deviation.

Figure 2. 16S rRNA gene OTUs enriched in heavy ^{13}C-DNA-SIP gradient fractions. Heatmaps showing the relative abundance of 16S rRNA gene OTUs in DNA-SIP gradient fractions from anoxic sediment incubations amended with ^{13}C-proteins or ^{13}C-lipids and from unamended control incubations. Significant (FDR-adjusted p-value was below 0.1) ^{13}C-enrichment was determined by differential abundance analysis between high density (^{13}C-enriched, >1.726 g ml^{-1}, pink blocks) and low density (^{13}C-free, <1.720 g ml^{-1}, turquoise blocks) DNA, and is indicated by filled circles. 16S rRNA genes that were matched to 16S rRNA gene sequences of genomes/MAGs are indicated in bold.

Figure 3. Differential [13]C-labeling and response of *Psychromonas* sub-OTUs in protein- and lipid-amended sediment microcosms. (A) Heatmap showing the relative abundance of four sub-OTUs of *Psychromonas* 16S rRNA OTU 4 in DNA-SIP gradients. Significant [13]C-enrichment was determined by differential abundance analysis between high density ([13]C-enriched) and low density ([12]C-enriched) DNA-SIP fractions. sOTU 4, which was identical to the 16S rRNA gene sequence of *Psychromonas* MAG GLG-1, is indicated in bold. (B) Phylogeny of *Psychromonas* sub-OTUs. The sequences were aligned with MAFFT (Katoh et al., 2002) and the tree was calculated with FastTree (Price et al., 2010). (C) Changes in relative abundance of *Psychromonas* sub-OTUs in protein-amended, lipid-amended, and unamended sediment microcosms with or without the sulfate reduction inhibitor molybdate.

Figure 4. Genome-inferred metabolic models of protein- and lipid-degrading bacteria in arctic marine sediment. Schematic depiction of predicted metabolic properties of *Psychrilyobacter atlanticus*, *Clostridia* MAG GPF-1, and *Psychromonas* MAG GLG-1, which correspond to the 16S rRNA OTUs 5, 1, and 4, respectively. The models indicate macromolecule degradation processes and enzymes that may explain the observed biochemical patterns and the [13]C-enrichment of corresponding 16S rRNA OTUs in the DNA-SIP incubations. Enzymes that are indicated in pale colour were not found in the respective genome/MAG. SS, Secretion system; TCA, tricarboxylic acid cycle; TAT, Twin-arginine translocation pathway; SEC, Sec secretion system; LIT, process has physiological evidence but the involved enzymes are not identified yet.