Disparate temperature-dependent virus – host dynamics for SARS-CoV-2 and SARS-CoV in the human respiratory epithelium

Philip V'kovski[1,2*], Mitra Gultom[1,2,3,4*], Jenna Kelly[1,2*], Silvio Steiner[1,2,3*], Julie Russeil[5], Bastien Mangeat[6], Elisa Cora[6], Joern Pezoldt[5], Melle Holwerda[1,2,3,4], Annika Kratzel[1,2,3], Laura Laloli[3,4], Manon Wider[4], Jasmine Portmann[1,2], Thao Tran[1,2,3], Nadine Ebert[1,2], Hanspeter Stalder[1,2], Rune Hartmann[7], Vincent Gardeux[5,8], Daniel Alpern[5,8], Bart Deplancke[5,8], Volker Thiel[1,2], Ronald Dijkman[1,2,4§]

[1] Institute of Virology and Immunology (IVI), Bern, Switzerland

[2] Department of Infectious Diseases and Pathobiology, Vetsuisse Faculty, University of Bern, Bern, Switzerland

[3] Graduate School for Biomedical Science, University of Bern, Bern, Switzerland

[4] Institute for Infectious Diseases, University of Bern, Bern, Switzerland

[5] Institute of Bioengineering, School of Life Sciences, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

[6] Gene Expression Core Facility (GECF), School of Life Sciences, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

[7] Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark

[8] Swiss Institute of Bioinformatics, Lausanne, Switzerland

*equal contribution

§corresponding author: Ronald Dijkman, email: ronald.dijkman@ifik.unibe.ch, phone: +41 31 664 07 83

Abstract: 158 words

Main text: 3519 words

Abstract

Since its emergence in December 2019, SARS-CoV-2 has spread globally and become a major public health burden. Despite its close phylogenetic relationship to SARS-CoV, SARS-CoV-2 exhibits increased human-to-human transmission dynamics, likely due to efficient early replication in the upper respiratory epithelium of infected individuals. Since different temperatures encountered in the human respiratory tract have been shown to affect the replication kinetics of several viruses, as well as host immune response dynamics, we investigated the impact of temperatures during SARS-CoV-2 and SARS-CoV infection in the human airway epithelial cell culture model. SARS-CoV-2, in contrast to SARS-CoV, replicated more efficiently at temperatures encountered in the upper respiratory tract, and displayed higher sensitivity to type I and type III IFNs. Time-resolved transcriptome analysis highlighted a temperature-dependent and virus-specific induction of the IFN-mediated antiviral response. These data reflect clinical features of SARS-CoV-2 and SARS-CoV, as well as their associated transmission efficiencies, and provide crucial insight on pivotal virus - host interaction dynamics.

Introduction

In December 2019, a new zoonotic coronavirus emerged in Wuhan, Hubei Province, China, which is referred to as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), and is the etiological agent of Coronavirus Disease 2019 (COVID-19) [1–3]. The novel coronavirus has a close phylogenetic relationship with SARS-CoV, which emerged in China in 2002/2003 and led to over 8000 confirmed cases worldwide, including 800 deaths [4]. SARS-CoV-2 differs from SARS-CoV by only 380 amino acids over its entire 30 kb genome and retains a high level of conservation in the receptor binding motifs that interact with the human receptor angiotensin converting enzyme 2 (ACE2) [5]. Despite these similarities, there are currently over 40 million confirmed cases of SARS-CoV-2 worldwide, including more than 1 million deaths [6] Moreover, although the cell surface receptor ACE2 and the serine protease TMPRSS2 have been demonstrated to serve as entry determinants for both SARS-CoV and SARS-CoV-2 [2,7–9], an accumulating body of evidence showing dissimilar human-to-human transmission dynamics and clinical courses between SARS-CoV-2 and SARS-CoV strongly suggests the presence of disparate virus-host dynamics during viral infection in the human respiratory epithelium [10–15].

The human conductive respiratory tract is lined by a pseudostratified, ciliated, columnar epithelium that contains mucin-producing goblet cells and represents a crucial barrier to constrain invading pathogens. The anatomical distance between the upper and lower respiratory conductive tract, and their different ambient temperatures (32-33°C and 37°C, respectively [16,17]), have previously been shown to influence the replication kinetics of diverse respiratory viruses, such as rhinoviruses, influenza viruses and coronaviruses [18–22]. Moreover, the anatomical disparity in ambient temperature also affects virus – host immune response dynamics, and thus potential human-to-human transmission dynamics [23]. Interestingly, SARS-CoV-2 has been detected earlier after infection than SARS-CoV in upper respiratory tissues of infected patients [10,13,14,24,25], suggesting that transmission kinetics and host innate immune response dynamics might differ between SARS-CoV and SARS-CoV-2 infections.

Since viral load may reflect the dynamic interaction between viral replication and inhibition by cellular defence mechanisms, we employed the human airway epithelial cell (hAEC) culture model to investigate the influence of different incubation temperatures on the viral replication kinetics and host immune response dynamics of both SARS-CoV and SARS-CoV-2 infections. Our study revealed that SARS-CoV-2 replication improved in hAEC cultures incubated at 33°C rather than 37°C and resulted in higher titers than SARS-CoV, while both viruses replicated equally efficiently at 37°C. Pretreatment of hAEC cultures with exogenous type I and III interferon (IFN) at different temperatures showed that SARS-CoV-2 and SARS-CoV are equally sensitive to both type I and III IFN, thereby exemplifying the relevance of early IFN signalling and innate immune responses to restrict viral infection. Importantly, a detailed temporal transcriptome analysis of infected hAEC cultures corroborated initial findings and uncovered characteristic innate immune response gene signatures relating to the viral replication efficiency of SARS-CoV and SARS-CoV-2 at different ambient temperatures. Altogether, these results provide the first in-depth fundamental insight on the virus-host innate immune response dynamics of SARS-CoV-2 and the closely phylogenetically related SARS-CoV in the respiratory epithelium, and likely reflect the clinical characteristics and transmission efficiencies of both viruses.

Results

Replication kinetics of SARS-CoV-2 and SARS-CoV at 33°C and 37°C

To assess the influence of the temperature variations that occur along the human respiratory tract, and to model virus-host interaction dynamics in distinct anatomical regions, we maintained well-differentiated hAEC cultures at either 33°C or 37°C throughout the experiment. hAEC cultures represent a well-characterized *in vitro* model that morphologically and functionally recapitulate the epithelial lining of the human respiratory tract *in vivo*. hAEC cultures from seven different human donors were inoculated with either SARS-CoV-2/München-1.1/2020/929 or SARS-CoV Frankfurt 1 isolates using a multiplicity of infection (MOI) of 0.1. The polarity of viral progeny release was monitored by collecting apical washes and basolateral medium in 24-hour intervals for a period of 96 hours. Since SARS-CoV-2 can be detected early after infection in the upper respiratory specimens of infected patients [14,24,25], we incubated virus-infected cultures at both 33°C and 37°C to mimic the ambient temperatures of the human upper and lower respiratory tract, respectively. At 37°C SARS-CoV and SARS-CoV-2 replicated to similar titers over the course of the infection (**Fig. 1a, b**). Interestingly, when assessing viral replication efficiency at 33°C rather than at 37°C, it was apparent that SARS-CoV-2 infection resulted in 10-fold higher titers released in the apical compartment between 72 and 96 hours post infection (hpi). In contrast, SARS-CoV replication at 33°C remained similar to replication at 37°C and showed no significant differences over the entire course of infection (**Fig. 1a, b**). Since the directionality of viral progeny release is crucial for subsequent virus spread and overall disease outcome, we also assessed whether SARS-CoV-2 was released to the apical surface, basolateral surface, or bilaterally. Similar to what we and others have observed previously for all other human coronaviruses, SARS-CoV-2 was predominantly released to the luminal surface (**Supp. Fig. 1a, b**) [26,27].

To assess whether the observed differential temperature-dependent replication efficiencies are a result of the number of cells infected by SARS-CoV or SARS-CoV-2, hAEC cultures were fixed at 96 hpi and processed for immunofluorescence analysis using antibodies directed against the SARS-CoV Nucleocapsid protein. Additionally, to discern potential preferential virus tropism to a distinct cell type, characteristic markers of the hAEC culture's architecture, such as the intercellular tight junctions (ZO-1) and the presence of cilia (β-tubulin), were also included. Microscopy investigations and automated image quantification revealed that in accordance with the more efficient replication kinetics of SARS-CoV-2 at 33°C, the fraction of SARS-CoV-2 infected cells increased significantly at 33°C compared to 37°C and to SARS-CoV. On the other hand, SARS-CoV did not show any significant change in the fraction of infected cells at 33°C compared to 37°C. Both viruses display a comparable fraction of infected cells at 37°C. (**Fig. 1c, d**). Notably, at 96 hpi the majority of SARS-CoV and SARS-CoV-2 antigen positive cells were not co-stained by the β-tubulin marker and were therefore qualified as non-ciliated cells (**Fig. 1c**). We previously observed that SARS-CoV infects both non-ciliated and ciliated cell populations [26], however, given that other reports show that SARS-CoV primarily targets ciliated cells [28], we analysed the localization of the entry receptor, ACE2, and β-tubulin markers by microscopy. Immunofluorescence analysis revealed that ACE2 is expressed in both ciliated and non-ciliated cell populations in uninfected hAEC cultures (**Supp. Fig. 2a**). In line with this, analysis of mRNA expression in non-infected hAEC cultures using single-cell RNA-sequencing (scRNA-seq)

confirmed that both *ACE2* and *TMPRSS2* mRNA are found in both secretory and ciliated cell
populations (**Supp. Fig. 2b-d**) [29]. Combined these results demonstrate that despite their shared
requirement on ACE2 and TMPRSS2 for entry into host cells, SARS-CoV and SARS-CoV-2 display
strong temperature-dependent variation in replication kinetics in hAEC cultures, suggestive of host
determinants intervening during post-entry stages of the viral life cycle. Importantly, the significantly
enhanced replication of SARS-CoV-2 at 33°C likely supports the increased replication in the upper
respiratory tract and transmissibility of SARS-CoV-2 compared to SARS-CoV.
133
Sensitivity of SARS-CoV-2 and SARS-CoV to IFN
The amount of viral progeny secreted from infected cells may reflect the dynamic interplay between
viral replication and its inhibition by cellular defence mechanisms, such as by different types of
interferon stimulated genes (ISGs). To examine whether the induction of ISGs differentially affects
SARS-CoV and SARS-CoV-2 replication, hAEC cultures were pretreated with 50 IU or 5 IU of
exogenous type I interferon (IFN-αA/D) and 50 or 5 ng of type III interferon (IFN-λ3), for 18 hours prior
to infection, at either 33°C or 37°C. Hereafter, the hAEC culture medium was replaced with IFN-free
medium and cells were infected with SARS-CoV and SARS-CoV-2 at an MOI of 0.1, at 33°C or 37°C,
for 72 hours. The titration of apically released virus revealed that the replication of both SARS-CoV-2
and SARS-CoV is severely restricted upon pretreatment with 50 IU of type I or 50 ng of III IFN at either
37°C or 33°C (**Fig. 2a**). Although, similar to previously reported observations [26], SARS-CoV seemed
less sensitive to type I IFN than to type III IFN pretreatment at 37°C (**Fig. 2a, 24 hpi**), these
differences were not statistically significant, which was further confirmed by pretreatment with 5 IU of
type I IFN or 5 ng of type III IFN (**Fig. 2b**). Consistently, SARS-CoV displayed a similar sensitivity to
type I and III IFN pretreatments at 33°C (**Fig. 2a, b**). In line with these results, SARS-CoV-2 was
equally sensitive to both doses of type I or III IFN pretreatments at either 37°C or 33°C (**Fig. 2a, b**).
However, overall SARS-CoV-2 replication kinetics were more severely impaired by IFN pretreatment
compared to SARS-CoV. In contrast, restriction of SARS-CoV-2 upon pretreatment with 10 IU of type
I and III IFNs was moderately relieved at 72 hpi (**Fig. 2b**).
The reduction of viral progeny titers in both IFN pretreatment conditions were corroborated by
immunofluorescence analysis at 72 hpi. Viral antigens were no longer detected upon immunostaining
of the type I or III IFN-pretreated hAEC cultures with anti-SARS-CoV Nucleocapsid protein antibodies
(**Fig. 2c**). Altogether, these results suggest that the viral replication kinetics of both SARS-CoV and
SARS-CoV-2 in the upper and lower airways are heavily dependent on innate immune responses
elicited by type I and III IFN, and that a potent IFN response can efficiently restrict viral replication of
SARS-CoV-2 in primary well-differentiated hAEC cultures.
160
Host response to SARS-CoV and SARS-CoV-2 infection in hAEC cultures at 33°C and 37°C.
The notable impact of incubation temperature on SARS-CoV and SARS-CoV-2 replication kinetics
(**Fig. 1**) and reduction in viral loads following both type I and III IFN pre-treatment (**Fig. 2**), prompted
the assessment of host transcriptional response dynamics to SARS-CoV and SARS-CoV-2 infections
at 33°C and 37°C. Cellular RNA was extracted from hAEC cultures infected with either SARS-CoV or
SARS-CoV-2 (MOI 0.1) and from uninfected hAEC cultures at 24, 48, 72, and 96 hpi. The latter was

then used for transcriptomics analysis following the Bulk RNA Barcoding and sequencing (BRB-
seq) protocol [30]. Data from seven different biological donors were collected and used to
perform pairwise comparisons between either SARS-CoV or SARS-CoV-2 virus-infected hAEC
cultures and unexposed hAEC cultures using three different approaches. First, pairwise comparisons
were performed by segregating sample by both temperature and time prior to analysis. This approach
identified a total of 185 differentially expressed genes (DEGs) for SARS-CoV and 351 DEGs for
SARS-CoV-2 ($Log_2FC \geq 1.5$, FDR ≤ 0.1), represented by a total of 401 unique genes (**Supp. Fig. 3a-**
d, Supp. Table 1).

To increase statistical power and identify DEGs that may have very low levels of expression in
uninfected hAEC cultures, two further approaches to differential expression analysis were also
performed. The first, which grouped samples by temperature, and then compared SARS-CoV or
SARS-CoV-2 virus-infected hAEC cultures to uninfected hAEC cultures, identified a total of 126 DEGs
for SARS-CoV at 33°C, 2 DEGs for SARS-CoV at 37°C, 161 DEGs for SARS-CoV-2 at 33°C, and 82
DEGs for SARS-CoV-2 at 37°C (**Fig. 3a, Supp. Table 2**). Hierarchical clustering of these SARS-CoV
and SARS-CoV-2 DEGs uncovered that irrespective of temperature and time, the overlap between the
host response induced by each virus was relatively low and the majority of upregulated DEGs were
present at 72 and 96 hpi (**Supp. Fig. 3a-d**). Interestingly, many DEGs identified at 72 and 96 hpi for
SARS-CoV-2 showed increased expression levels as early as 48 hpi at 37°C, but not at 33°C (**Fig.**
3b). Comparison of DEGs identified at each temperature for SARS-CoV-2 revealed a core group of 45
common upregulated ISG-related genes, including *IFNL1*, *CXCL10*, *CXCL11*, *IRF7*, *STAT1*, *IFI35*,
OASL, *CMP2K*, *HELZ2*, *MX1*, *DDX60*, *IFI44L*, *ISG15*, and *IFIT3* (**Supp. Table 3**). Finally, pairwise
comparisons were also performed for ungrouped samples between SARS-CoV or SARS-CoV-2 virus-
infected hAEC cultures and uninfected hAEC cultures (**Supp. Table 4**). In addition to the individual
comparisons for SARS-CoV-2, we also performed direct pairwise comparisons between SARS-CoV
and SARS-CoV-2. The latter revealed that the most contrasting differences observed between the two
viruses occurred at 96 hpi for 33°C, whereas they occurred at 72 hpi for 37°C (**Supp. Fig. 3e, f**).
DEGs identified in the SARS-CoV-2 versus SARS-CoV analysis using the aforementioned three
approaches are summarized in **Supp. Tables 5-7**.

To investigate the hierarchical clustering results from the SARS-CoV or SARS-CoV-2 versus
uninfected hAEC cultures in more detail, we annotated the top 25 upregulated DEGs for SARS-CoV-2
among the 16 distinct conditions (**Fig. 3b**). This revealed a clear temporal and temperature-dependent
host response profile for SARS-CoV-2 virus-infected hAEC cultures, a finding that was less apparent
for SARS-CoV. Furthermore, we found that *CXCL10*, *CXCL11*, and *TNFSF13B*, chemokines
responsible for immune cell recruitment to the site of infection from the bloodstream, were among the
core group of 45 common upregulated DEGs identified at both temperatures, albeit upregulation was
stronger at 37°C compared to 33°C. Conversely, no upregulation was observed during SARS-CoV-2
infection in hAEC cultures for canonical pro-inflammatory genes such as *TNF*, *IL-11*, *IL-18*, *IL-6*, and
IL1B (**Supp. Fig. 4a-b**). The pattern recognition receptor (PRR) *RIG-I/DDX58* and interferon-inducible
2'-5'-oligoadenylate synthetase-like protein (*OASL*), were also identified as some of the most
prominent DEGs that showed an earlier and higher expression level at 37°C (**Fig. 3b**).

To establish whether any biological pathways were significantly enriched over time at the
different ambient temperatures, pathway enrichment analysis was performed on all unique DEGs
detected in the comparison analyses for SARS-CoV or SARS-CoV-2. Similar to the hierarchical
clustering results, this analysis displayed a distinct temperature-dependent profile for SARS-CoV-2 at
72 and 96 hpi, specifically in diverse IFN and antiviral signalling pathways (**Fig. 3c**). Notably, none of
these antiviral pathways were significantly enriched during SARS-CoV infection. The 43 unique genes
associated with these enriched pathways, including *RIG-I/DDX58* and *OASL*, all displayed a clear
temporal and temperature-dependent expression pattern that was inversely associated with the
temperature-dependent viral kinetics difference for SARS-CoV-2 (**Fig. 3e, Fig. 1a, b, and Supp. Fig.**
5). Several individual genes were found to be associated with multiple enriched pathways, including
canonical ISG-related genes such as *OASL*, *STAT1*, *ISG15*, and *IFIT1*. The intersection among the
top 5 significantly enriched pathways for SARS-CoV-2 at 96 hpi for both temperatures, and 72 hpi for
37°C, is shown in **Fig. 3d**. A high degree of interconnectivity was observed among DEGs associated
with these 5 pathways, further highlighting the importance of the host antiviral response during SARS-
CoV-2 infection (**Fig. 3e**).

We previously observed, in the context of influenza A/H1N1 virus, that only a small fraction of
infected cells produces IFN during infection [31]. Since the majority of upregulated DEGs during SARS-
CoV-2 infection were genes induced downstream of the IFN pathway, we next sought to assess the
expression of individual type I and III IFN genes over time. Interestingly, although only *IFNL1* and
IFNB1 were classified as significantly upregulated DEGs in our analysis, the expression levels of
IFNL2 and *IFNL3* also followed a similar temperature-dependent pattern as the ISGs highlighted in
Fig. 3f. In contrast, and in agreement with previous results (**Fig. 3a, e**), SARS-CoV infection did not
induce type I or III IFNs at either 33°C or 37°C (**Fig. 3f**). Notably, we confirmed the expression of both
type I and III IFN receptors (*IFNAR1*, *IFNAR2*, *IFNLR1*, and *IL10RB*) in multiple cell types of non-
infected hAEC cultures using scRNA-seq (**Supp. Fig. 2e-h**). Together these results suggest that
SARS-CoV infection triggers only very mild IFN induction in hAEC cultures, whereas SARS-CoV-2
infection leads to stronger induction of multiple IFNs that is dominated by type III IFNs and dependent
on temperature [31].

Taken together, these results clearly demonstrate that SARS-CoV-2 and SARS-CoV induce
disparate, virus-specific, and temperature-dependent host responses that inversely correlate with the
previously observed dissimilar viral replication efficiencies at temperatures corresponding to the upper
and lower respiratory tract. The majority of DEGs are related to the antiviral and pro-inflammatory
response, which is much more pronounced in SARS-CoV-2 rather than SARS-CoV virus-infection, and
the delayed induction of these DEGs at 33°C coincides with the increased replication of SARS-CoV-2
at temperatures corresponding to the upper respiratory epithelium.

Discussion

In the current study, we demonstrate that the ambient temperatures reminiscent of the conditions in
the upper and lower respiratory tract have a profound influence on both viral replication and virus-host
dynamics, particularly innate immune responses, during SARS-CoV and SARS-CoV-2 infection in
human airway epithelial cells. Using an authentic *in vitro* model for the human respiratory epithelium

we demonstrated that SARS-CoV-2, in contrast to SARS-CoV, replicated up to 10-fold more efficiently
at temperatures encountered in the upper respiratory tract. Concordantly, significantly increased
amounts of Nucleocapsid-antigen positive cells were detected in these conditions. In addition, and
despite intrinsic donor-to-donor variations, SARS-CoV-2 and SARS-CoV were highly sensitive to
pretreatment with exogenous type I and type III IFNs. Importantly, temporal transcriptome analysis
showed a temperature-dependent induction of the IFN-mediated antiviral and pro-inflammatory
responses that inversely correlated with the observed replication kinetic efficiencies of both SARS-
CoV and SARS-CoV-2 at temperatures found in the upper and lower respiratory tract.

One of the most profound phenotypical characteristics of fulminant SARS-CoV-2 is the early
replication in the upper respiratory tract of infected individuals, which might facilitate the high
transmissibility of SARS-CoV-2 [14,24,25,32]. In contrast, SARS-CoV was shown to primarily replicate in
the lower respiratory tract and efficient transmissibility occurred at later stages of the clinical course
[10,12,13]. Additionally, SARS-CoV was demonstrated to inhibit the induction of interferon responses in
infected cells [33,34]. The data presented here reflect these aspects of SARS-CoV and SARS-CoV-2
infections and contribute to the understanding of the disparate human-to-human transmission
dynamics for both zoonotic coronaviruses. They provide a framework to address the parameters of the
molecular basis of exacerbations induced by SARS-CoV-2 infection in predisposed individuals.

SARS-CoV-2 replication was strongly potentiated at 33°C and significant differences in the
number of infected cells were observed between SARS-CoV and SARS-CoV-2 infected cultures.
Given that the viral S protein receptor binding motifs interacting with the human receptor ACE2 are
highly conserved between the two viruses and that both SARS-CoV and SARS-CoV-2 displayed a
similar cell tropism, the 380 amino acid dissimilarities, distributed across the entire genome and
distinguishing SARS-CoV-2 from SARS-CoV, might account for their differential replication efficiencies
[2,7–9]. Another factor that may have influenced our results is the 29-nucleotide truncation in the *ORF8*
gene of SARS-CoV Frankfurt-1, which was maintained in the SARS-CoV lineage that initiated the
international spread of SARS-CoV. Indeed, Muth and colleagues demonstrated that an intact *ORF8*
invoked 10-fold higher replication kinetics at 37°C in various cell culture models, including hAEC
cultures [35]. Therefore, besides comparing the replication of different SARS-CoV *ORF8* variants at the
temperature corresponding to the upper respiratory tract, it would be equally compelling to assess the
phenotypic influence of similar truncations in the *ORF8* gene of SARS-CoV-2, especially since several
SARS-CoV-2 isolates bearing a 382-nucleotide deletion truncating the *ORF8* gene have been
detected [36]. Such SARS-CoV-2 *ORF8* variants can be readily engineered using the reverse genetic
systems that were recently established for SARS-CoV-2 [32,37,38].

In our study, we report different temperature-dependent viral replication efficiencies for SARS-
CoV and SARS-CoV-2, inversely associated with the amplitude of the innate immune response, albeit
with a more pronounced phenotype for SARS-CoV-2. Our results are consistent with other reports
using undifferentiated primary cell-based systems where the induction of interferon responses was
detected between 24 and 48 hours post infection with SARS-CoV-2 [39]. In our study we analysed
SARS-CoV-2-infected differentiated AEC cultures derived from multiple donors until 96 hpi at two
different ambient temperatures. These data demonstrate a substantial induction of interferon and
inflammatory responses in the airway epithelium following SARS-CoV-2 infection, in line with recent

single-cell sequencing experiments performed on patient-derived samples [40]. Foxman and colleagues
elegantly described, in an analogous model to the human AEC cultures and by using common cold
viruses, that the PRR-mediated IFN response is influenced by temperature [23]. This may also apply in
the context of SARS-CoV-2 infections, however, due to the multifaceted intricate nature of virus-host
interactions it is likely that the efficient replication of SARS-CoV-2 and the concurrent expression of a
plethora of known coronavirus antagonists of the antiviral response also play a crucial role herein [41–46].
Nonetheless, we demonstrate that SARS-CoV-2 and SARS-CoV are sensitive to both type I and III
IFN. These data are supported by several studies investigating the outcome of IFN pre-treatment in
cultured cell lines [39,47,48] as well as the well-documented dominant antiviral role of type III IFN during
virus infection in the respiratory epithelium [31,49,50]. IFN lambda therefore represents an attractive
candidate for the development of intervention strategies against SARS-CoV-2 respiratory infections.

The findings reported here provide crucial insight into the profound impact of ambient
temperatures on pivotal virus – host interactions in the airway epithelium, which in turn influences the
host immune response to SARS-CoV-2 infection [51]. This work will likely lead to additional functional *in*
vivo studies delineating the efficacy of antiviral responses triggered by SARS-CoV and SARS-CoV-2
infections, as well as deciphering the influence of viral antagonists and physical conditions. Therefore,
the disparate viral replication efficiencies and host responses at 33°C and 37°C provide important
insights into the molecular basis of SARS-CoV-2 infection and should be exploited broadly to support
clinical interventions in COVID-19 patients.

Material and methods
Cells and human airway epithelial cell (hAEC) cultures
Vero-E6 cells (kindly provided by Doreen Muth, Marcel Müller, and Christian Drosten, Charité, Berlin,
Germany) were propagated in Dulbecco's Modified Eagle Medium-GlutaMAX supplemented with 1
mM sodium pyruvate, 10% (v/v) heat-inactivated fetal bovine serum (FBS), 100 µg/ml streptomycin,
100 IU/ml penicillin, 1% (w/v) non-essential amino acids and 15 mM HEPES (Gibco). Cells were
maintained at 37°C in a humidified incubator with 5% CO_2.

Primary human tracheobronchial epithelial cells were isolated from patients (>18 years old)
undergoing bronchoscopy or pulmonary resection at the Cantonal Hospital in St. Gallen, Switzerland,
or Inselspital in Bern, Switzerland, in accordance with ethical approval (EKSG 11/044, EKSG 11/103,
KEK-BE 302/2015, and KEK-BE 1571/2019). Isolation and culturing of primary material was performed
as previously described [52]. Briefly, cryopreserved cells were thawed and expanded for one week in
BEGM medium. After initial expansion phase, cells were transferred into in collagen type IV-coated
porous inserts (6.5 mm radius insert, Costar) in 24-well plates. Cells were expanded for another 2-3
days in BEGM in a liquid-liquid state. Once the cells reached confluency, the basolateral medium was
exchanged for ALI medium and the apical medium was removed to allow for the establishment of the
air-liquid-interface (ALI). Basolateral ALI medium was exchanged three times per week and apical side
was washed with Hanks balanced salt solution (HBSS, Gibco) once a week, until the development of a
fully differentiated epithelium (3-4 weeks), which was monitored by optical microscopy. Several
modifications to the original protocol were used. The concentrations of hydrocortisone for both BEGM
and ALI were increased to 0.48 µg/ml and BEGM was further supplemented with the inhibitors 1

μmol/L A83-01 (Tocris, USA), 3 μmol/L isoproterenol (Abcam, Cambridge, United Kingdom) and 5

μmol/L Y27832 (Tocris, USA) [53]. Basolateral ALI medium was exchanged three times per week and

apical side was washed with hanks balanced salt solution (HBSS, Gibco) once a week. hAEC cultures

were maintained at 37°C in a humidified incubator with 5% CO_2.

Viruses

SARS-CoV strain Frankfurt-1 (GenBank FJ429166) [35,54] and SARS-CoV-2 (SARS-CoV-2/München-

1.1/2020/929) [55] were kindly provided by Daniela Niemeyer, Marcel Müller, and Christian Drosten, and

propagated and titrated on Vero E6 cells.

Infection of hAEC cultures

Well-differentiated hAEC cultures were infected with 30,000 plaque-forming units (PFU) of either

SARS-CoV or SARS-CoV-2. Viruses were diluted in Hanks balanced salt solution (HBSS, Gibco),

inoculated on the apical side and incubated for 1 h at 33°C or 37°C. Afterwards, virus inoculum was

removed, and the apical surface washed three times with HBSS, whereby the third wash was collected

as the 1 hpi timepoint. The cells were incubated at the indicated temperatures in a humidified

incubator with 5% CO_2. Released virus progeny were monitored every 24 h by incubating 100 μl of

HBSS on the apical surface 10 min prior to the time point. The apical washes were collected, diluted

1:1 with virus transport medium (VTM), and stored at -80°C for later analysis. Basolateral medium was

collected at each time point and stored at -80°C for later analysis. Fresh ALI medium was then added

to the basolateral compartment. To analyze virus replication following interferon (IFN) exposure, hAEC

cultures were pretreated with recombinant universal type I IFN (100 or 10 IU/ml; Sigma) or

recombinant IFN-λ3 (100 or 10 ng/ml [56]) for 18 h from the basolateral side, prior to infection and

incubated at either 33°C or 37°C. As controls, untreated hAEC cultures were used. Shortly before

infection with SARS-CoV and SARS-CoV-2, the basolateral medium containing type I or type III IFN

was removed and replaced with medium without exogenous IFN.

Immunofluorescence analysis of infected hAECs

Well-differentiated hAEC cultures were fixed with 4% (v/v) neutral buffered formalin and processed as

previously described [52]. Cells were permeabilized in PBS supplemented with 50 mM NH_4Cl, 0.1%

(w/v) Saponin and 2% (w/v) Bovine Serum Albumin (CB). To detect SARS-CoV and SARS-CoV-2,

hAEC cultures were immunostained with a rabbit polyclonal antibody against SARS-CoV

Nucleocapsid protein (Rockland, 200-401-A50), which also cross-react with SARS-CoV-2. Cell

distribution of ACE2 were detected with a rabbit polyclonal antibody against ACE2 (ab15348, Abcam).

Alexa Fluor® 488-labeled donkey anti-rabbit IgG (H + L) (Jackson Immunoresearch) was used as

secondary antibody. Alexa Fluor® 647-labeled rabbit anti-β-tubulin (9F3, Cell Signaling Technology)

and Alexa Fluor® 594-labeled mouse anti-ZO1 (1A12, Thermo Fisher Scientific) were used to

visualize cilia and tight junctions, respectively. Antibodies were diluted in CB. All samples were

counterstained using 4',6-diamidino-2-phenylindole (DAPI, Thermo Fisher Scientific) to visualize the

nuclei. Samples were imaged on a DeltaVision Elite High-Resolution imaging system (GE Healthcare

Life Sciences) equipped with 60x oil immersion objective (1.4 NA), by acquiring 200-300 nm z-stacks

over the entire thickness of the sample. Images were deconvolved using the integrated softWoRx
software. For the quantification of infected cells, images were alternatively acquired using an EVOS FL
Auto 2 Imaging System equipped with a 20x air objective. All images were processed using FIJI
software packages [57]. Brightness and contrast were adjusted identically to their corresponding
controls. Figures were assembled using the FigureJ plugin [58]. Quantification of infected cells from four
donors was performed by morphological segmentation of individual cells using the ZO-1 staining and
the MorphoLibJ plugin in FIJI [59]. Each region of interest was used to measure the mean intensity in the
channel corresponding to the nucleocapsid staining. Cells with mean intensities > mean + 3 standard
deviations compared to the distribution of mock-infected cells were considered positive. On average
over 10^4 cells were analysed per donor and per condition.
381

Titration of apical and basolateral compartments

Viruses released in the apical or basolateral compartments were titrated by plaque assay on Vero-E6
cells. Shortly, $1.7*10^5$ cells/ml were seeded in 24-well plates one day prior to the titration and
inoculated with 10–fold serial dilutions of virus solutions. Inoculums were removed 1.5 hpi and
replaced with overlay medium consisting of DMEM supplemented with 1.2% Avicel (RC-581, FMC
biopolymer), 5% heat-inactivated FBS, 50 mg/ml streptomycin and 50 IU/ml penicillin. Cells were
incubated at 37 °C 5% CO_2 for 48 hours, fixed with 4% (v/v) neutral buffered formalin and stained with
crystal violet.
390

Bulk RNA Barcoding and sequencing (BRB-seq) and data analysis

Total cellular RNA from mock and virus-infected hAEC cultures was extracted with the NucleoMag
RNA kit (Macherey-Nagel) according manufacturers guidelines on a Kingfisher Flex Purification
system (Thermofisher). Total RNA concentration was quantified with QuantiFluor® RNA System
(Promega) according manufactures guidelines on a Cytation 5 multimode reader (Biotek). A total of
100 ng of total cellular RNA was used for the generation of Bulk RNA Barcoding and sequencing
(BRB-seq) libraries, and the subsequent sequencing on an Illumina HiSeq 4000 platform was done as
previously described at a depth of approximately 12 million raw reads per sample [30]. The sequencing
reads were demultiplexed using the BRB-seqTools suite and were aligned against a concatenation of
the human gene annotation of the human genome (hg38), SARS coronavirus Frankfurt 1 (AY291315)
and SARS-CoV-2/Wuhan-Hu1/2020 (NC_045512) viral genomes using STAR and HTSeq for
producing the count matrices. All downstream analyses were performed using R (version 3.6.1).
ComBat-seq was used with default settings to adjust for batch effects in the raw data and generate an
adjusted count matrix used for downstream analyses [60]. Library normalization and expression
differences between uninfected and virus-infected samples were then quantified using the DESeq2
package in R (version 1.28) with a fold change (FC) cut-off of ≥ 1.5 and a False Discovery Rate (FDR)
of ≤ 0.1 [61]. Due to the multi-factor design of these experiments, differential expression (DE) analysis
was performed using several approaches: 1) Samples were subset by temperature and time prior to
DE analysis (e.g. subset of samples at 33□C and 24 hpi). Infected samples were then compared to
uninfected samples using the design ~ Batch + Condition to identify DEGs; 2) Samples were subset by
temperature only prior to DE analysis (e.g. subset of samples for all time points at 33□C) and infected

samples were compared to uninfected samples using the design ~ Batch + Condition; 3) Samples
were kept together (not subset) and infected samples were compared to uninfected samples using the
design ~ Batch + Condition. Substantial overlap was observed for DEGs identified using these 3
approaches and results for each approach are reported in Supplemental tables 1, 2, 4, 5, 6, 7. Venn
diagrams were generated using the VennDiagram package in R for DEGs identified in approach 2
(**Fig. 3a**) and approach 1 (Supplementary Figure 3) [62].

Pathway enrichment analysis was performed using the clusterProfiler and ReactomePA
packages in R [63,64]. Significantly enriched pathways with a gene count > 1 and p-value of ≤ 0.05 were
visualized using the enrichplot package. The complete list of significantly enriched pathways identified,
as well as the genes associated with these pathways, are shown in Supplemental table 8. Additional
data analysis and visualization was performed using a variety of packages in R, including
ComplexHeatmap and ggplot2.[65]

Bioinformatic analysis of ACE2 and TMPRSS2 expression
For the analysis of *ACE2*, *TMPRSS2*, *IFNAR1*, *IFNAR2*, *IFNRL1*, and *IL10RB* mRNA expression we
reanalysed the previously obtained single-cell raw sequencing data [31]. The resulting unique molecule
identifier (UMI) count matrix of each individual sample was pre-processed, filtered individually and
merged in Seurat (v3.1). Data scaling, normalization and regressing out unwanted sources of variation
(number of UMI's, mitochondrial content, cell cycle phase) was performed using integrated
SCtransform option, followed by dimensional reduction using UMAP (Uniform Manifold Approximation
and Projection) embedding. For cell type annotation, the resulting integrated dataset was used for
unsupervised graph-based clustering to annotate the different cell types using both cluster-specific
marker genes and well-known canonical marker genes to match identified clusters with specific cell
types found in the respiratory epithelium, as previously described[31].

Statistical testing

Distribution testing was performed using Shapiro-Wilk normality test (>0.05), followed by computing the P value of the mean log10 PFU/ml at each timepoint between SARS-CoV and SARS-CoV-2 using a two-sided paired sample t test. Analyses were performed using R (version 3.6.1).

Data

Transcriptome data has been deposited in the Arrayexpress open-access public repository from the European Bioinformatics Institute (EMBL-EBI) under E-MTAB-9781, and scripts used for analysis and figure generation will be become available at Github upon publication.

Acknowledgements

We gratefully thank the École Polytechnique Fédérale de Lausanne (EPFL) and the University of Bern for providing special authorization to conduct our research during the SARS-CoV-2 outbreak. We are grateful to Sabina Berezowska and Irene Ramos-Centeno (Institute of Pathology, University of Bern) for providing the tissues via the Tissue Bank Bern. This work was supported by the European Commission (Marie Sklodowska-Curie Innovative Training Network "HONOURS"; grant agreement No 721367), the Swiss National Science Foundation (SNSF) grants 179260, 160780, 173085, 31CA30_196644, the National Center of Competence in Research (NCCR) on RNA and Disease funded by the SNSF, and the German Federal Ministry of Education and Research, project RAPID.

Author contributions

Conceptualization, R.D.; Investigation, P.V., M.G, S.S., J.K, J.R., B.M., E.C., J.P., M.H., A.K., L.L., M.W., J.P., T.T., N.E., H.S. and R.D., Resources, R.H., Writing – Original Draft, P.V. and R.D.; Writing – Review & Editing, P.V., J.K., V.T. and R.D.; Visualization, P.V., M.G., S.S., J.K. and R.D.; Supervision, V.G., D.A., B.D., V.T. and R.D.; Project Administration, R.D.; Funding Acquisition, V.T. and R.D.

Declaration of Interests

The authors declare no competing interests.

Figure legends

Figure 1. SARS-CoV and SARS-CoV-2 replication kinetics in hAEC cultures.

Well-differentiated hAEC cultures were infected with SARS-CoV (**a**) and SARS-CoV-2 (**b**) using 30,000 PFU or remain uninfected (mock), and were incubated at 37°C or 33°C. Inoculated virus was removed at 1 hpi and the apical side was washed. Cultures were further incubated at the indicated temperature. At the indicated time post infection, apical virus release was assessed by plaque titration (**a-b**). Data represent the mean ± 95% CI of hAEC cultures from seven different human donors. Individual points represent the average of two technical replicates. Values at 0 hpi indicate the titer of the inoculum used to infect the hAEC cultures, and values at 1 hpi indicate the remaining titer after the third wash. The p-values were computed by using two-sided paired sample t tests. At 96 hpi, hAEC

cultures were fixed and processed for immunofluorescence analysis using antibodies against SARS-
CoV Nucleocapsid protein (green), β-tubulin (cilia, red), ZO-1 (tight junctions, white) and DAPI (blue)
(**c**). Representative z-projections of one donor are shown. Scale bar, 20 microns. Infected cells were
quantified after segmentation of individual cells based on the ZO-1 staining and measuring the mean
intensity of the nucleocapsid protein staining (**d**). Data represent the mean ± 95% CI of multiple
images acquired from hAEC cultures derived from four different human donors. On average, more
than 10^4 cells per donor and per condition were analysed.

Figure 2. SARS-CoV and SARS-CoV replication upon IFN-I and IFN-III pretreatment
hAEC cultures were treated from the basolateral side with recombinant universal type I interferon (100
IU/ml or 10 IU/ml) or recombinant IFN-λ3 (100 ng/ml or 10 ng/ml) for 18 h. Before infection, medium
was removed and replaced with IFN-free medium and hAEC cultures were infected with SARS-CoV
and SARS-CoV-2 using 30,000 PFU, and were incubated at 37°C or 33°C. Inoculated virus was
removed at 1 hpi, and the apical side was washed. Cultures were further incubated at the indicated
temperature. At the indicated time, apical virus release was assessed by plaque titration (**a,b**). Data
represent the mean ± 95% CI of hAEC cultures from three different human donors. Individual points
represent the average of two technical replicates. Values at 0 hpi indicate the titer of the inoculum
used to infect the hAEC cultures, and values at 1 hpi indicate the remaining titer after the third wash.
The p-values were computed by using two-sided paired sample t tests. At 96 hpi, hAEC cultures
pretreated with 100 IU/ml type I IFN or 100 ng/ml type III IFN were fixed and processed for
immunofluorescence analysis using antibodies against SARS-CoV Nucleocapsid protein (green), β-
tubulin (cilia, red), ZO-1 (tight junctions, white) and DAPI (blue) (**c**). Representative z-projections of
one donor are shown. Scale bar, 20 microns.

Figure 3. Temperature-dependent host transcriptional response in SARS-CoV and SARS-CoV-2
virus-infected hAEC cultures.
Venn diagram showing the overlap among differentially expressed genes (DEG) identified in SARS-
CoV (purple) and SARS-CoV-2 (orange) virus-infected hAEC cultures at either 33°C or 37°C (**a**).
Hierarchical cluster analysis of DEGs identified in SARS-CoV and SARS-CoV-2 virus-infected hAEC
cultures at either 33°C or 37°C compared to uninfected hAEC cultures (Mock). Expression levels for
individual DEGs are shown in rows as the $\log_2$ mean normalized counts for seven human donors
stratified by condition, temperature, and hours post-infection (columns; representative colours shown
in legends). The top 25 DEGs for SARS-CoV-2 (ranked by adjusted p-values) are shown on the right
(y-axis) (**b**). Dotplot illustrating pathway enrichment analysis performed on the 16 distinct DEG profiles.
Significantly enriched pathways for SARS-CoV and SARS-CoV-2 are shown for both 33°C and 37°C
incubation temperatures at 72- and 96-hours post-infection (hpi). Dots were adjusted in size and
colour to illustrate the gene ratio and adjusted p-value (< 0.05) for a given pathway, respectively (**c**).
An UpSet plot visualizing the intersection of the number of individual genes (y-axis) found to be
associated with the significantly enriched pathways (x-axis). Intersections with 2 or more overlapping
genes are highlighted in orange (**d**). Gene-concept network plot illustrating the individual relationships
between DEGs and the top 5 significantly enriched pathways (orange) for SARS-CoV-2. Enriched

pathway hubs (orange) were adjusted in size to reflect the number of genes associated with each
respective pathway, whereas individual genes are coloured based on their expression level (fold
change relative to expression level in uninfected hAEC cultures) (**e**). Bar graph illustrating the $\log_2$
mean normalized expression levels over time for *IFNL1*, *IFNL2*, *IFNL3*, and *IFNB1*, at the respective
temperatures for Mock (uninfected) hAEC cultures (top two panels) and for SARS-CoV (middle two
panels) and SARS-CoV-2 (bottom two panels) virus-infected hAEC cultures. Bars were adjusted in
colour to illustrate the respective standard deviation (SD) among donors (**f**).
524

References

1.   Li, Q. *et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected
Pneumonia. *N. Engl. J. Med.* (2020). doi:10.1056/nejmoa2001316

2.   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin.
Nature **579**, 270–273 (2020).

3.   Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The
species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and
naming it SARS-CoV-2. *Nat. Microbiol.* 1–9 (2020). doi:10.1038/s41564-020-0695-z

4.   WHO | Summary of probable SARS cases with onset of illness from 1 November 2002 to 31
July 2003. Available at: https://www.who.int/csr/sars/country/table2004_04_21/en/. (Accessed:
23rd March 2020)

5.   Wu, A. *et al.* Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV)
Originating in China. *Cell Host Microbe* **27**, 325–328 (2020).

6.   WHO - Weekly epidemiological update - 3 November 2020. Available at:
https://www.who.int/publications/m/item/weekly-epidemiological-update---3-november-2020.
(Accessed: 3rd November 2020)

7.   Li, W. *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS
coronavirus. *Nature* **426**, 450–454 (2003).

8.   Glowacka, I. *et al.* Evidence that TMPRSS2 Activates the Severe Acute Respiratory Syndrome
Coronavirus Spike Protein for Membrane Fusion and Reduces Viral Control by the Humoral
Immune Response. *J. Virol.* **85**, 4122–4134 (2011).

9.   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked
by a Clinically Proven Protease Inhibitor. *Cell* (2020). doi:10.1016/j.cell.2020.02.052

10. Cheng, P. K. C. *et al.* Viral shedding patterns of coronavirus in patients with probable severe
acute respiratory syndrome. *Lancet* **363**, 1699–1700 (2004).

11. Peiris, J. S. M. *et al.* Coronavirus as a possible cause of severe acute respiratory syndrome.
Lancet **361**, 1319–1325 (2003).

12. Peiris, J. S. M. *et al.* Clinical progression and viral load in a community outbreak of
coronavirus-associated SARS pneumonia: A prospective study. *Lancet* **361**, 1767–1772
(2003).

13. Hung, I. F. N. *et al.* Viral loads in clinical specimens and SARS manifestations. *Emerg. Infect.*
Dis. **10**, 1550–7 (2004).

14. Wölfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* 1–10
(2020). doi:10.1038/s41586-020-2196-x

15. Zhou, F. *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in
Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054–1062 (2020).

16. McFadden, E. R. *et al.* Thermal mapping of the airways in humans. *J. Appl. Physiol.* **58**, 564–
570 (1985).

17. Lindemann, J., Leiacker, R., Rettinger, G. & Keck, T. Nasal mucosal temperature during
respiration. *Clin. Otolaryngol. Allied Sci.* **27**, 135–139 (2002).

18. Kendall, E. J. C., Bynoe, M. L. & Tyrrell, D. A. J. Virus isolations from common colds occurring
in a residential school. *Br. Med. J.* **2**, 82–86 (1962).

19. Tyrrell, D. A. J. & Bynoe, M. L. Cultivation of a Novel Type of Common-cold Virus in Organ
Cultures. *Br. Med. J.* **1**, 1467–1470 (1965).

20. Hoorn, B. & Tyrrell, D. A. J. A new virus cultivated only in organ cultures of human ciliated
epithelium. *Arch. Gesamte Virusforsch.* **18**, 210–225 (1966).

21. Corman, V. M. *et al.* Link of a ubiquitous human coronavirus to dromedary camels. *Proc. Natl.*
Acad. Sci. U. S. A. **113**, 9864–9 (2016).

22. Holwerda, M. *et al.* Determining the Replication Kinetics and Cellular Tropism of Influenza D
Virus on Primary Well-Differentiated Human Airway Epithelial Cells. *Viruses* **11**, (2019).

23. Foxman, E. F. *et al.* Temperature-dependent innate defense against the common cold virus
limits viral replication at warm temperature in mouse airway cells. *Proc. Natl. Acad. Sci. U. S.*
A. **112**, 827–832 (2015).

24. Zou, L. *et al.* SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N.*
Engl. J. Med. **382**, 1177–1179 (2020).

25. He, X. *et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat. Med.*
(2020). doi:10.1038/s41591-020-0869-5

26. Kindler, E. *et al.* Efficient replication of the novel human betacoronavirus EMC on primary
human epithelium highlights its zoonotic potential. *MBio* **4**, (2013).

27. Dijkman, R. *et al.* Isolation and Characterization of Current Human Coronavirus Strains in
Primary Human Epithelial Cell Cultures Reveal Differences in Target Cell Tropism. *J. Virol.* **87**,
6081–6090 (2013).

28. Jia, H. P. *et al.* ACE2 Receptor Expression and Severe Acute Respiratory Syndrome
Coronavirus Infection Depend on Differentiation of Human Airway Epithelia. *J. Virol.* **79**,
14614–14621 (2005).

29. Bertram, S. *et al.* TMPRSS2 Activates the Human Coronavirus 229E for Cathepsin-
Independent Host Cell Entry and Is Expressed in Viral Target Cells in the Respiratory
Epithelium. *J. Virol.* **87**, 6150–6160 (2013).

30. Alpern, D. *et al.* BRB-seq: Ultra-affordable high-throughput transcriptomics enabled by bulk
RNA barcoding and sequencing. *Genome Biol.* **20**, 71 (2019).

31. Kelly, J. N. *et al.* Comprehensive single cell analysis of pandemic influenza A virus infection in
the human airways uncovers cell-type specific host transcriptional signatures relevant for
disease progression and pathogenesis. *bioRxiv* 2020.04.03.014282 (2020).
doi:10.1101/2020.04.03.014282

32. Hou, Y. J. *et al.* SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the
Respiratory Tract. *Cell* **182**, 429-446.e14 (2020).

33. Channappanavar, R. *et al.* Dysregulated Type I Interferon and Inflammatory Monocyte-
Macrophage Responses Cause Lethal Pneumonia in SARS-CoV-Infected Mice. *Cell Host*
Microbe **19**, 181–193 (2016).

34. Gralinski, L. E. & Baric, R. S. Molecular pathology of emerging coronavirus infections. *J.*
Pathol. **235**, 185–195 (2015).

35. Muth, D. *et al.* Attenuation of replication by a 29 nucleotide deletion in SARS-coronavirus
acquired during the early stages of human-to-human transmission. *Sci. Rep.* **8**, (2018).

36. Su, Y. C. F. *et al.* Discovery and genomic characterization of a 382-nucleotide deletion in
ORF7B and orf8 during the early evolution of SARS-CoV-2. *MBio* **11**, 1–9 (2020).

37.   Thao, T. T. N. *et al.* Rapid reconstruction of SARS-CoV-2 using a synthetic genomics platform.
611   *Nature* 1–8 (2020). doi:10.1038/s41586-020-2294-9

612   38.   Xie, X. *et al.* An Infectious cDNA Clone of SARS-CoV-2. *Cell Host Microbe* **27**, 841-848.e3
613   (2020).

614   39.   Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development of
615   COVID-19. *Cell* **181**, 1036-1045.e9 (2020).

616   40.   Chua, R. L. *et al.* COVID-19 severity correlates with airway epithelium–immune cell interactions
617   identified by single-cell analysis. *Nat. Biotechnol.* (2020). doi:10.1038/s41587-020-0602-4

618   41.   Frieman, M. *et al.* Severe Acute Respiratory Syndrome Coronavirus ORF6 Antagonizes STAT1
619   Function by Sequestering Nuclear Import Factors on the Rough Endoplasmic Reticulum/Golgi
620   Membrane. *J. Virol.* **81**, 9812–9824 (2007).

621   42.   Züst, R. *et al.* Coronavirus non-structural protein 1 is a major pathogenicity factor: Implications
622   for the rational design of coronavirus vaccines. *PLoS Pathog.* **3**, 1062–1072 (2007).

623   43.   Kuri, T. *et al.* The ADP-ribose-1″-monophosphatase domains of severe acute respiratory
624   syndrome coronavirus and human coronavirus 229E mediate resistance to antiviral interferon
625   responses. *J. Gen. Virol.* **92**, 1899–1905 (2011).

626   44.   Züst, R. *et al.* Ribose 2'-O-methylation provides a molecular signature for the distinction of self
627   and non-self mRNA dependent on the RNA sensor Mda5. *Nat. Immunol.* **12**, 137–143 (2011).

628   45.   Kindler, E. *et al.* Early endonuclease-mediated evasion of RNA sensing ensures efficient
629   coronavirus replication. *PLoS Pathog.* **13**, (2017).

630   46.   Niemeyer, D. *et al.* The papain-like protease determines a virulence trait that varies among
631   members of the SARS-coronavirus species. *PLoS Pathog.* **14**, (2018).

632   47.   Lokugamage, K. G., Hage, A., Schindewolf, C., Rajsbaum, R. & Menachery, V. D. SARS-CoV-
633   2 is sensitive to type I interferon pretreatment. *bioRxiv* 2020.03.07.982264 (2020).
634   doi:10.1101/2020.03.07.982264

635   48.   Felgenhauer, U. *et al.* Inhibition of SARS–CoV-2 by type I and type III interferons. *J. Biol.*
636   *Chem.* **295**, 13958–13964 (2020).

637   49.   Davidson, S. *et al.* IFN λ is a potent anti-influenza therapeutic without the inflammatory side
638   effects of IFN α treatment . *EMBO Mol. Med.* **8**, 1099–1112 (2016).

639   50.   Galani, I. E. *et al.* Interferon-λ Mediates Non-redundant Front-Line Antiviral Protection against
640   Influenza Virus Infection without Compromising Host Fitness. *Immunity* **46**, 875-890.e6 (2017).

641   51.   Vabret, N. *et al.* Immunology of COVID-19: Current State of the Science. *Immunity* **52**, 910–
642   941 (2020).

643   52.   Jonsdottir, H. R. & Dijkman, R. Characterization of human coronaviruses on well-differentiated
644   human airway epithelial cell cultures. in *Coronaviruses: Methods and Protocols* **1282**, 73–87
645   (Springer New York, 2015).

646   53.   Gultom, M., Laloli, L. & Dijkman, R. Well-Differentiated Primary Mammalian Airway Epithelial
647   Cell Cultures. in *Methods in Molecular Biology* **2203**, 119–134 (Humana Press Inc., 2020).

648   54.   Pfefferle, S. *et al.* Reverse genetic characterization of the natural genomic deletion in SARS-
649   Coronavirus strain Frankfurt-1 open reading frame 7b reveals an attenuating function of the 7b
650   protein in-vitro and in-vivo. *Virol. J.* **6**, 131 (2009).

651   55.   Tran Thi Nhu Thao, Fabien Labroussaa, Nadine Ebert, Philip V'kovski, Hanspeter Stalder,
652   Jasmine Portmann, Jenna Kelly, Silvio Steiner, Melle Holwerda, Annika Kratze, Mitra Gultom,
653   Laura Laloli, Linda Hüsser, Manon Wider, Stephanie Pfaender, Dagny Hirt, Va, V. T. *et al.*
654   Rapid reconstruction of SARS-CoV-2 using a synthetic genomics platform. 2020.02.21.959817
655   (2020).

656   56.   Lauber, C. *et al.* Transcriptome analysis reveals a classical interferon signature induced by
657   IFNλ4 in human primary cells. *Genes Immun.* **16**, 414–421 (2015).

658   57.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**,
659   676–682 (2012).

660   58.   Mutterer, J. & Zinck, E. Quick-and-clean article figures with FigureJ. *J. Microsc.* **252**, 89–91
661   (2013).

662   59.   Legland, D., Arganda-Carreras, I. & Andrey, P. MorphoLibJ: integrated library and plugins for
663   mathematical morphology with ImageJ. *Bioinformatics* btw413 (2016).
664   doi:10.1093/bioinformatics/btw413

665   60.   Zhang, Y., Parmigiani, G. & Johnson, W. E. ComBat-seq: batch effect adjustment for RNA-seq
666   count data. *NAR Genomics Bioinforma.* **2**, (2020).

667   61.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
668   RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

669   62.   Chen, H. & Boutros, P. C. VennDiagram: a package for the generation of highly-customizable
670   Venn and Euler diagrams in R. *BMC Bioinformatics* **12**, 35 (2011).

671   63.   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. ClusterProfiler: An R package for comparing
672   biological themes among gene clusters. *Omi. A J. Integr. Biol.* **16**, 284–287 (2012).

673   64.   Yu, G. & He, Q. Y. ReactomePA: An R/Bioconductor package for reactome pathway analysis
674   and visualization. *Mol. Biosyst.* **12**, 477–479 (2016).

675   65.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
676   multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

Figure 1

Figure 2

Figure 3