1

Early-life stress facilitates the development of Alzheimer's disease pathology via angiopathy

Running title: Early-life stress-associated Alzheimer's disease

Tomoko Tanaka[1], Shinobu Hirai [1], Masato Hosokawa [2], Takashi Saito [3,4], Hiroshi Sakuma [5],

Takaomi Saido [3], Masato Hasegawa [2], Haruo Okado [1, *]

[1] Laboratory of Neural Development, Department of Psychiatry & Behavioral Science, Tokyo

Metropolitan Institute of Medical Science, Tokyo, Japan.

[2] Dementia Research Project, Department of Brain & Neuroscience, Tokyo Metropolitan

Institute of Medical Science, Tokyo, Japan.

[3] Laboratory for Proteolytic Neuroscience, RIKEN Center for Brain Science, Saitama, Japan.

[4] Department of Neurocognitive Science, Institute of Brain Science, Nagoya City University

Graduate School of Medical Science, Aichi, Japan.

[5] Child brain Project, Department of Brain & Neuroscience, Tokyo Metropolitan Institute of

Medical Science, Tokyo, Japan.

1   * Correspondence should be addressed to Haruo Okado, Neural Development Project,

2   Department of Brain Development and Neural Regeneration, Tokyo Metropolitan Institute of

3   Medical Science, Tokyo 156-8506, Japan. E-mail: okado-hr@igakuken.or.jp (H.O.)

5   Keyword: Alzheimer's disease, angiopathy, immune system, early life stress, amyloid-beta,

6   cognitive function

Abstract

Background: Alzheimer's disease (AD), a progressive neurodegenerative disorder, is a serious

social problem. Recently, several early-life factors have been associated with an increased risk

of a clinical diagnosis of AD.

Methods: We investigated the involvement of early-life stress in AD pathogenesis using

heterozygous the amyloid precursor protein (APP) mutant mice ($App^{NL-G-F/wt}$) and wild-type

($App^{wt/wt}$) mice. Maternal separation was used as an animal paradigm for early-life stress. Object

location and fear conditioning tests were performed to measure cognitive functions, in addition

to biochemical tests. Immunohistochemical analyses were performed after the behavioral tests.

Results: We found that maternal-separated $App^{wt/wt}$ mice showed narrowing of vessels and

decreased pericyte coverage of capillaries in prefrontal cortex, while maternal-separated App^{NL-}

$^{G-F/wt}$ mice additionally showed impairment of cognitive function, and earlier formation of Aβ

plaques and disruption of the blood–brain barrier. Severe activation of microglia was detected in

the maternal-separated $App^{NL-G-F/wt}$ mice and maternal-separated $App^{wt/wt}$ mice. At the early stage,

morphological changes and inflammatory responses were observed in the microglia of the

1   maternal-separated $App^{NL-G-F/wt}$ mice and maternal-separated $App^{wt/wt}$ mice, as well as

2   morphological changes in the microglia of the non-maternal-separated $App^{NL-G-F/wt}$ mice.

3   Conclusions: Microglia activation induced by maternal separation in combination with the APP

4   mutation may impairs the vascular system, leading to AD progression. These findings therefore

5   suggest that maternal separation causes early induction of AD pathology via angiopathy.

1 Introduction

The development of abnormal physiology in association with increased age is observed in the

progression of many disorders. Because daily habits and environmental stresses such as alcohol

intake and diet have the major influence on life-span, rather than genetic factors (Christensen et

al., 2006), acceleration of the aging process by such environmental factors is thought to increase

both the risk of developing a disease and mortality from a disease. Alzheimer's disease (AD) is

one such disease that is thought to be influenced by environmental factors; it is characterized by

progressive impairment of cognitive functions such as memory. It is considered that the

acquired predisposition to AD increases with age, and that trauma during adulthood increases

the likelihood of developing AD. Moreover, recent epidemiological research has suggested that

stressful environmental factors during the developmental period, including mistreatment,

neglect, and the loss of a parent, can also increase the likelihood of developing AD (Norton et

al., 2011; Radford et al., 2017; Seifan et al., 2015). However, as the research behind these

findings is generally the result of retrospective cohort studies, it is unclear whether early-life

stress affects the onset and progression of AD.

Recently, vascular impairment, a postmortem pathological finding in the brain, was reported

to occur before the onset of AD. Therefore, vascular impairment is considered as an initial

symptom of AD (Iadecola, 2017; Montagne et al., 2016; Wen et al., 2004). The neurovascular

unit (NVU) consists of neurons, endothelial cells, pericytes, microglia, and astrocytes. As the

NVU is important for the extrusion of amyloid beta, an impairment in extraction can cause

sporadic AD and cerebral amyloid angiopathy (Mawuenyega et al., 2010). In particular, the

capillary vessels in the brain play an important role in the extrusion of amyloid beta.

Dysfunction of the blood–brain barrier (BBB) and a decrease in amyloid beta clearance through

the cerebral vascular system is observed in AD (Zlokovic, 2011). Degeneration of cells making

up the BBB, such as pericytes, causes dysfunction of the BBB (Sweeney et al., 2016). However,

it remains unclear whether early-life stress results in dysfunction of the vascular system.

In this study, we investigated whether maternal separation, which can be used as an early-life

stress in rodents, facilitates the development of AD, with the purpose of understanding the

causal association between the development of AD and early-life stress. After finding

indications that maternal separation affects the development of AD, we next investigated the

changes induced by maternal separation, paying particular attention to capillary vessel

morphology in heterozygous the amyloid precursor protein (APP) mutant mice ($App^{NL-G-F/wt}$) and

1   wild-type ($App^{wt/wt}$) mice. As a result, we established that maternal separation facilitates the

2   development of AD. Furthermore, we found that impairment of capillaries, probably caused by

3   abnormalities in microglia, leads to dysfunction of the BBB, thereby facilitating the

4   development of clinical AD.

Materials and methods

All experimental procedures were approved by the Animal Experimentation Ethics Committee

of the Tokyo Metropolitan Institute of Medical Science (49040).

Animals

The original lines of App mutant mice ($App^{NL-G-F/NL-G-F}$) were established as a C57BL/6J

congenic line (a genetic background strain) by repeated back-crosses, as described previously,

and were obtained from RIKEN Center for Brain Science (Wako, Japan)(Saito et al., 2014). All

experiments were performed with $App^{NL-G-F/wt}$ and $App^{wt/wt}$ that were born from $App^{NL-G-F/wt}$ mice

x $App^{NL-G-F/wt}$ mice or $App^{NL-G-F/wt}$ mice x $App^{wt/wt}$ mice. $App^{NL-G-F/wt}$ mice with slow onset of

pathology were used because it was important to investigate the process of decline in function.

All mice were maintained under a 12:12 h light/dark cycle (lights on at 8:00 AM). All efforts

were made to minimize the number of animals used and their suffering. All experiments, except

for maternal separation, were performed in adolescence (postnatal day 45 (P 45)) or adulthood

(P90–P120).

Maternal separation

Maternal separation is usually used as a form of early-life stress in mice. Pups were separated

from their mother for 3 hours (AM 9:00-12:00) daily from P2 to P15. The separated pups were

taken to a different room and placed on a heat pat to maintain their body temperature. The

control mice had no experience of maternal separation. The body weights were not significantly

different between the non-maternal-separated and maternal-separated mice at several stages of

development.

Behavioral assessment

Male mice were allowed to habituate in the behavioral room for at least 1 hour before

commencement of the behavioral test. Mice showing outlier behavior > 2 standard deviations

from the mean or < 2 standard deviations from the mean were not analyzed by the behavioral

test.

The object location test consisted of three phases. All phases were performed under a light

intensity of 10 lux. On day 1, the animals were placed in an empty square box (50 cm x 50 cm)

for 10 minutes. On day 2, the animals were placed in the same box for 10 minutes with two

same identical unfamiliar objects. On day 3, the animals were placed in the same box, with one

of the two items being displaced to a novel location in the area. The discrimination index was

calculated as the ratio of the time spent exploring the novel place to that spent exploring the

familiar place: discrimination index = (novel place exploration time – familiar place exploration

time)/(novel place exploration time + familiar place exploration time). On each day, the

exploration time in each place was calculated automatically using a DVTrack video tracking

system (Muromachi, Tokyo, Japan).

The fear conditioning test consisted of three phases. On day 1, the animals were placed in a

triangular box for 5 minutes under a light intensity of 30 lux. On day 2, the animals were placed

into a square box with a stainless-steel grid floor and were allowed to explore the box freely for

2 minutes. Subsequently, a tone, which served as the conditioned stimulus (CS), was presented

for 30 sec. During the last 1 sec of CS presentation, a 0.75 mA electric shock was applied,

which served as the un-conditioning stimulus (US). Two more CS-US parings were presented

with a 1 minutes inter-stimulus interval under a light intensity of 100 lux. On day 3, the animals

were placed into the same square box as used on day2 for 5 minutes under a light intensity of

100 lux. On day 4, the animals were placed in the same triangular box as used on day 1 for 10

minutes and were allowed to explore the box freely for 2 minutes. Subsequently, CS was

presented for 30 sec. Seven more CS were presented with a 1 minutes inter-stimulus interval

under a light intensity of 100 lux. In each test, the percentage of freezing time and distance

travelled (cm) were calculated automatically using ImageFZ software (O'Hara & Co., Tokyo,

Japan). Freezing was taken to indicate that the mice remembered the box (context) or tone (cue)

in which they were exposed to the electrical shock.

Immunohistochemistry

At P45 or P120, male and female mice were anaesthetized with isoflurane and then

sequentially perfused through the heart with PBS followed by 4% paraformaldehyde (PFA) in

PBS. Their brains were post-fixed, cryoprotected in 20% sucrose at 4℃, and embedded in OCT

compound and frozen. Coronal sections of mouse brain were then cut using a cryostat (20 μm).

Free-floating sections containing PFC were activated by HistVT one (Nakalai., Kyoto, Japan) at

70℃ for 20 minutes, incubated in 0.4% Block-ase (DS Pharma Biomedical., Osaka, Japan) in

PBS at room temperature for 20 minutes, and then incubated with primary antibody

(Supplemental figure 1). All primary antibodies were diluted 1:500 in PBS containing 0.3%

Triton X-100. The sections were then incubated with secondary antibody (Jackson., Maine,

United State) (diluted 1:500 in PBS containing 0.3% Triton X-100) for 2 hours at room

temperature. All sections types were incubated with DAPI (Nacalai Tesque., Kyoto, Japan) for

nuclear staining. Fluorescence micrographs of the sections subjected to immunostaining

procedures were captured and digitized with a FluoView® FV3000 confocal laser scanning

microscope (Olympus., Tokyo, Japan). With the aim of quantifying the morphological change of

microglia, skeleton analysis and particle analysis were performed by ImageJ(Fernández-Arjona

et al., 2017; Xu et al., 2018).

Luminex analysis assay

Following the sacrifice of male and female mice, mouse PFC fractions were dissociated with

a tissue homogenizer in 100 μl of 0.5 × cell lysis buffer 2 diluted in PBS (R&D Systems.,

Minnesota, United State). Tissues were lysed at room temperature for 30 minutes, and debris

was then removed by centrifugation. Magnetic luminex assays were performed using a Mouse

Premixed Multi-Analysis Kit (R&D Systems., Minnesota, United State). Assay procedures were

carried out following manufacturer's instructions. Microplates were run on a Bio-Plex 200

system (Bio-Rad., California, United State).

Corticosterone assay

Male and female mice were subjected to a novel cage stress (Rattray et al., 2009), in which

each mouse was placed in isolation in a new cage with a smooth floor for 30 minutes at P45.

Blood samples were taken before stress and immediately after stress, and centrifuged at 3,300

rpm to separate serum. Corticosterone concentrations were measured using an enzyme-linked

immunosorbent assay (ELISA) (AssayPro., Missouri, United State).

Statistical data analysis

Group data are presented as the mean ± SEM. The statistical significance of between-group

differences was assessed by the Tukey–Kramer test, Dunnett's test, or the paired t-test, using

JMP software (SUS Institute., North Carolina, United State). The interaction between area and

distance was assessed using the polynomial regression model. The statistical significance of

differences between groups and over time was assessed by two-way repeated ANOVA followed

by a *post-hoc* Holm's test, using R software.

Results

Experimental design

Figure 1A shows the experimental schedule. To investigate whether stress in the early

developmental stage is involved in the onset and/or progression of AD, we performed maternal

separation for 3 hours daily during the period from P2 to P15 in $App^{wt/wt}$ and $App^{NL-G-F/wt}$ mice.

The $App^{NL-G-F/wt}$ mice show an AD-like phenotype in the late-aged stage (Saito et al., 2014). We

then performed behavioral tests and tissue sampling around P45 (adolescence) and P120 (adult).

Maternal separation induced impairment of cognitive function and facilitated formation of

amyloid beta plaque in the adult $App^{NL-G-F/wt}$ mice

Figure 1B illustrates the effect of maternal separation on object location test which evaluate

spatial memory (Denninger et al., 2018). Object location memory in the $App^{wt/wt}$ mice was not

altered by maternal separation, as indicated by the fact that the discrimination index was not

changed by maternal separation. By contrast, object location memory in the $App^{NL-G-F/wt}$ mice

was impaired by maternal separation, as indicated by a discrimination index close to zero. The

non-maternal-separated $App^{NL-G-F/wt}$ mice showed intact object location memory. These results

indicate that the combination of APP mutation and maternal separation influence object location

memory. Figure 1C indicates the effect of maternal separation on contextual and cued fear

memory. Contextual fear memory in the $App^{wt/wt}$ mice was not altered by maternal separation,

while that in the $App^{NL-G-F/wt}$ mice was impaired by maternal separation, as indicated by the

lower freezing time in the maternal-separated $App^{NL-G-F/wt}$ mice in comparison with the non-

maternal-separated $App^{NL-G-F/wt}$ and non-maternal separated $App^{wt/wt}$ mice. By contrast, tone-cued

fear memory was not impaired in any of the four groups, indicating that contextual memory,

which is related to prefrontal cortex–hippocampus signaling and is known to be necessary for

the encoding of spatial cues, was impaired (Kitamura et al., 2017). Figure 1D indicates the

effect of maternal separation on the formation of amyloid beta plaque. The formation of

amyloid beta plaque in the $App^{NL-G-F/wt}$ mice was facilitated in mice that underwent maternal

separation than in non-maternal-separated $App^{NL-G-F/wt}$ mice.

Maternal separation induces vascular system impairment in adult mice

Phenotypes showing decreased pericyte coverage, a narrowing of capillary diameter, and

impairment of the BBB are frequently observed in the postmortem brain of AD patients

(Halliday et al., 2016). Figure 2 illustrates the effect of maternal separation on the vascular

system. The maternal-separated $App^{wt/wt}$ and maternal-separated $App^{NL-G-F/wt}$ mice showed lower

pericyte coverage of capillaries in the prefrontal cortex (PFC) than the non-maternal-separated

controls in adult. Furthermore, maternal separation induced a narrowing of capillary diameter in

both the $App^{wt/wt}$ mice and $App^{NL-G-F/wt}$ mice (Figure 2A). The more vessel-associated microglia

were detected in the maternal-separated $App^{NL-G-F/wt}$ mice (Figure 2B). The result suggests

strengthening of the interaction between microglia and vessels in the maternal-separated App^{NL-}

$^{G-F/wt}$ mice. The vessel-associated microglia migrate towards cerebral vessel and cause increased

BBB permeability; Microglia, which can acquire a reactive phenotype that phagocytoses BBB

components, initiate leakage of systemic substances into the parenchyma (Haruwaka et al.,

2019). In our study, extra-vascular accumulation of fibrinogen deposits in the PFC was detected

only in the maternal-separated $App^{NL-G-F/wt}$ mice (Figure 2C). We speculate that this

accumulation was caused by leakage of fibrinogen due to impairment of the vascular system by

more cessel-associated phagocytotic microglia only in the maternal-separated $App^{NL-G-F/wt}$ mice.

Figure 2D shows the relationship between the sizes of the amyloid beta plaques and the

distances from the amyloid beta plaques to vessels in the maternal-separated and non-maternal-

separated $App^{NL\text{-}G\text{-}F/wt}$ mice that had mild amyloid beta plaque symptoms. The correlations

between the sizes of amyloid beta plaques and the distances from the amyloid beta plaques to

vessels were significant, which suggests the possibility that amyloid beta plaques were

generated around the vessels and caused disappearance of the vessels after disruption. We also

detected ameboid microglia surrounding amyloid beta plaques stained with tomato-lectin

(Acarin et al., 1994).

Either maternal separation or APP mutation caused activation of and morphological change in

microglia in adult mice

As the vascular system is affected by microglia, we examined the activity and morphology of

microglia. Figure 3 shows the effects of maternal separation on microglia. Maternal separation

caused activation of the microglia in the PFC of both $App^{wt/wt}$ mice and $App^{NL\text{-}G\text{-}F/wt}$ mice (Figure

3A). Although the numbers of microglia were not significantly different between any of the

groups, except for the microglia surrounded by amyloid beta plaque, the proportions of soma

and dendrites stained by Iba1 were lower in both $App^{wt/wt}$ mice and $App^{NL\text{-}G\text{-}F/wt}$ mice subjected to

maternal separation (Figure 3A) than in the corresponding controls. Maternal separation may

have induced morphological changes in both genotypes. More detailed analysis of the

morphological changes revealed differences between non-maternal-separated $App^{wt/wt}$ mice and

other groups (Figure 3B). Lower numbers of dendrite terminals and a shortening of the total

branch length were detected in the maternal-separated $App^{NL-G-F/wt}$ mice than in the non-

maternal-separated $App^{wt/wt}$ mice. On the other hand, the number of dendrite terminals was

lower in the non-maternal-separated $App^{NL-G-F/wt}$ mice than in the non-maternal-separated

$App^{wt/wt}$ mice (Figure 3B). The non-maternal-separated $App^{NL-G-F/wt}$ mice may have milder

microglial symptoms than the maternal-separated animals. Furthermore, the circularity and

solidity of microglia increased in only the maternal-separated $App^{NL-G-F/wt}$ mice. We speculated

that maternal separation in combination with the APP mutation induced ameboid-like severe

morphological change.

The maternal-separated $App^{wt/wt}$ and $App^{NL-G-F/wt}$ mice showed intact object location memory,

absence of amyloid beta plaque, and normal vascular morphology and pericyte in adolescence

To detect initial changes in the maternal-separated and/or mutated mice, we examined the

phenotypes at an earlier stage (P90), that is, in adolescence. Figure 4A shows the effect of

maternal separation on object location memory in adolescence. In all groups, the discrimination

index for object location memory indicated object location memory to be intact. Figure 4B

shows that amyloid beta plaque was absent in all groups. Figure 4C shows no impairment of

vascular morphology or pericytes in any of the groups. We detect no changes in cognitive

function, plaque formation and vascular system in the maternal-separated and/or mutated mice

at adolescence, which allow us to rule out that the maternal-separated and/or mutated mice are

model mice for developmental disorders.

Maternal separation and APP mutation cause functional and morphological change to

microglia in adolescence

Figure 5 indicates the effect of maternal separation on microglia in adolescence. The numbers

of activated microglia were not significantly different between each group (Figure 5A). A more

detailed analysis of morphological differences revealed lower numbers of dendrite terminals in

the maternal-separated $App^{NL-G-F/wt}$ mice than in the non-maternal-separated $App^{wt/wt}$ mice. In the

$App^{wt/wt}$ mice, the maternally separated mice also showed lower numbers of dendrite terminals.

Furthermore, lower numbers of dendrite terminals were detected in the non-maternal-separated

$App^{NL\text{-}G\text{-}F/wt}$ mice than in the non-maternal-separated $App^{wt/wt}$ mice (Figure 5B). On the other

hand, the numbers of vessel-associated microglia did not differ between the groups (Figure 5C).

To evaluate the pro-inflammatory properties of microglia, we measured the level of TNF-R1

protein in the PFC. Maternal-separated $App^{wt/wt}$ and $App^{NL\text{-}G\text{-}F/wt}$ mice showed higher levels of

TNF-R1 protein than the non-maternal-separated mice (Figure 5D). We detected changes

associated with both maternal separation and heterozygotic mutation in adolescence, indicating

the possibility that a change in microglia caused by both maternal separation and APP mutation

induces an AD-like phenotype. To assess the effect of maternal separation on the hypothalamus-

pituitary-adrenal axis in adolescence, we measured serum corticosterone levels induced by a

novel cage stress in maternal-separated and non-separated mice. Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$

and $App^{wt/wt}$ mice subjected to a novel cage stress in adolescence mice showed more

corticosterone release than non-maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ and $App^{wt/wt}$ mice (Figure 5E).

Discussion

In this study, we investigated whether maternal separation facilitates the development of AD,

with the aim of understanding the causal association between early-life stress and the

development and progression of AD. We also investigated for changes induced by maternal

separation, particularly in capillary vessels. We established that maternal separation facilitates

the impairment of spatial cognitive function and the formation of amyloid beta plaque in $App^{NL\text{-}}$

$^{G\text{-}F/wt}$ mice, with disruption of micro-capillaries, and we verified that early-life stress constitutes

a risk factor for AD. Furthermore, we found that morphological and functional changes to

microglia are early symptoms in our experimental model, and suggest the possibility that

impairment of the cerebral vascular system through microglia caused by combination of

maternal separation and APP mutation induces dysfunction in the BBB, thereby facilitating the

clinical condition of AD.

Research using the Childhood Trauma Questionnaire to assess childhood stress has suggested

that early-life stress may be associated with increased risk of AD and dementia (Radford et al.,

2017). However, because most previous reports, including the above research, were based on

retrospective cohort studies, it remains unclear whether early-life stress affects the onset and

progression of AD. Although it has been reported that maternal separation facilitates the

formation of amyloid beta plaque using an AD mouse model different from the one we used

(Hui et al., 2017), the pathway by which maternal separation induces increased amyloid beta

plaque remains unclear. We found that maternal separation facilitates the impairment of spatial

cognitive function and the formation of amyloid beta plaque through impairment of the cerebral

vascular system in $App^{NL\text{-}G\text{-}F/wt}$ mice. We propose that these maternally separated $App^{NL\text{-}G\text{-}F/wt}$

mice can be used as a stress-induced AD mouse model. As it is known that progression of AD in

humans is exacerbated by stress, such as a change in abode and hospital admission in old age,

our AD mouse model may contribute to understanding the mechanism of stress-induced AD and

how it can be prevented.

We found that maternal separation mildly impaired the cerebral capillary vascular system.

However, dysfunction of the BBB, which is a more serious symptom, was detected in the

maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice and was related to both maternal separation and genetic

factors. Several mechanisms are involved in clear amyloid beta clearance (Chen et al., 2017).

One mechanism involves transcytosis through cells of the BBB. Another mechanism involves

phagocytosis by microglia. The reduction in amyloid beta clearance due to impairment of the

vascular system and impaired microglial phagocytoses of amyloid beta may induce

inflammation and increase fibrin generation due to coagulation factor XII (FXII) activation by

amyloid beta. The interaction of amyloid beta with fibrin (or fibrinogen) can lead to increased

fibrin deposition in cerebral blood vessels, and these accumulated fibrin deposits may induce

microinfarcts, which result in hemorrhage, inflammation, and BBB disruption, all of which are

commonly observed in AD (Ahn et al., 2017). Our result that disappearance of vessel around

larger amyloid plaque (Fig. 2D) may be caused by the interaction of amyloid beta with fibrin.

We report that maternal separation induces impairments to the cerebral vascular system, with

impairments to pericytes and narrowing of vessels in adulthood. Both of these phenotypes are

detected in the postmortem brains of human patients with AD (Halliday et al., 2016). Therefore,

these findings indicate a relationship between early-life stress and AD pathology. It has been

demonstrated that both antenatal and postnatal stress delay the development of the BBB during

weaning (Gomez-Gonzalez and Escobar, 2009). We detected no impairment in vascular

morphology or in pericytes in adolescence. Our findings that early-life stress induces

impairments in adult mice is in accord with the Development Origins of Health and Disease

(DOHaD) hypothesis, which purports that exposure to certain environmental conditions during

critical periods of development and growth may have significant adverse effects on short- and

long-term health. As the ultrastructural breakdown of capillaries and a decrease in blood flow

are detected in AD patients and AD mouse models (de la Torre, 2004), it is likely that

impairment of the cerebral vascular system develops before the onset of AD pathogeny in

$App^{NL-G-F/wt}$ mice.

We detected morphological changes in microglia and changes in their pro-inflammatory

properties (which are mediated by the activation of TNF-R1) at an early stage, while we did not

detect amyloid plaque formation, changes to the vascular system, or changes in behavior.

Microglia have been reported to show morphological changes in response to the clinical

condition of AD (Yamada and Jinno, 2013). An increase in corticosterone in response to stress

has been reported to be related to a change in microglia (Rincon-Cortes and Sullivan, 2014). In

our study, maternal-separated $App^{NL-G-F/wt}$ and maternal-separated $App^{wt/wt}$ mice showed

increased corticosterone release in response to a novel cage stress at adolescence, which

suggests that maternal separation affects the stress response. In fact, the production of

corticosterone in response to stress facilitates the release of monocytes into the circulation from

bone marrow, which in turn facilitates monocyte adherence to vessels (Niraula et al., 2018). The

release of monocytes upregulates IL-1β and stimulates endothelial cells (McKim et al., 2018).

Haruwaka et al. hypothesized that endothelial cells may respond to systemic inflammation and

release molecules to trigger microglia migration and phenotype changes. They also reported that

chemokine (C-C motif) ligand 5 (CCL5), which is secreted by endothelial cells, attracts

microglia to vessels. Recently, plasma from aged mice was shown to affect the activity of

microglia, referred to as neuroinflammation, and increase VCAM1 expression on endothelial

cell, a major cell type of the BBB, in young mice (Yousef et al., 2019). The authors suggested

that circulating cytokines and chemokines with detrimental effects on the brain give rise to these

changes in microglia and endothelial cells. Considering that increased age is crucial for the

development of AD, increases in inflammatory signals in the plasma may induce AD. On the

other hand, the non-maternal-separated $App^{NL-G-F/wt}$ mice also showed morphological changes in

adolescence. Our data support the hypothesis that impairment of the vascular system by amyloid

beta oligomers induces activation of microglia (Nortley et al., 2019). As microglia and the

vascular system are deeply linked, the possibility that an undetectable alteration of the vascular

system led to a change in the microglia, which then led to the development of AD via an

interaction between microglia and the vascular system, cannot be ruled out. The changes in

microglia induced by maternal separation and those induced by APP mutation may occur via

different mechanisms. Microglia activation induced by maternal separation in combination with

$App^{NL-G-F/wt}$ mice may impairs the vascular system, leading to AD progression.

In the present study, we established that maternal separation facilitates the development of

AD. Furthermore, we identified changes in microglia induced by maternal separation and/or

genetic factors as an early phenotype, which probably preceded the impairment of the vascular

system in the stress-induced AD mouse model. These results indicate that microglia could have

potential use as biomarkers and as targets for new therapies against AD.

Author contributions

T.T. and H.O. designed the research. T.T. performed and analyzed all experiments. T.S. and T.S.

generated and provided App mutant mice ($App^{NL\text{-}G\text{-}F/NL\text{-}G\text{-}F}$). S.H., M.H., H.S., and M.H. helped

with this study and made important suggestions. T.T. generated all the figures and tables and

wrote the manuscript. H.O. edited the manuscript and supervised the study.

Acknowledgments

The authors thank Dr. Makoto Hashimoto for advice. We would like to thank Bioedit Ltd

(https://www.bioedit.com) for English language editing. This work was partially supported by

JSPS KAKENHI Grant Number 18H02537 for Haruo Okado, and by JSPS KAKENHI Grant

Number 17K16408 and Research Grant for Public Health Science for Tomoko Tanaka.

Competing interests

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

References

Acarin, L., Vela, J.M., González, B., Castellano, B., 1994. Demonstration of

poly-N-acetyl lactosamine residues in ameboid and ramified microglial cells

in rat brain by tomato lectin binding. J. Histochem. Cytochem. 42, 1033-

1041.

Ahn, H.J., Chen, Z.L., Zamolodchikov, D., Norris, E.H., Strickland, S., 2017.

Interactions of ß-amyloid peptide with fibrinogen and coagulation factor XII

may contribute to Alzheimer's disease. Curr. Opin. Hematol. 24, 427-431.

Chen, G.F., Xu, T.H., Yan, Y., Zhou, Y.R., Jiang, Y., Melcher, K., Xu, H.E.,

2017. Amyloid beta: structure, biology and structure-based therapeutic

development. Acta Pharmacol. Sin. 38, 1205-1235.

Christensen, K., Johnson, T.E., Vaupel, J.W., 2006. The quest for genetic

determinants of human longevity: challenges and insights. Nature reviews.

Genetics 7, 436-448.

de la Torre, J.C., 2004. Is Alzheimer's disease a neurodegenerative or a

vascular disorder? Data, dogma, and dialectics. Lancet Neurol. 3, 184-190.

Denninger, J.K., Smith, B.M., Kirby, E.D., 2018. Novel Object Recognition

and Object Location Behavioral Testing in Mice on a Budget. J Vis Exp.

Fernández-Arjona, M.D.M., Grondona, J.M., Granados-Durán, P.,

Fernández-Llebrez, P., López-Ávalos, M.D., 2017. Microglia Morphological

Categorization in a Rat Model of Neuroinflammation by Hierarchical

Cluster and Principal Components Analysis. Front. Cell. Neurosci. 11, 235.

Gomez-Gonzalez, B., Escobar, A., 2009. Altered functional development of

the blood-brain barrier after early life stress in the rat. Brain Res. Bull. 79,

376-387.

Halliday, M.R., Rege, S.V., Ma, Q., Zhao, Z., Miller, C.A., Winkler, E.A.,

Zlokovic, B.V., 2016. Accelerated pericyte degeneration and blood-brain

barrier breakdown in apolipoprotein E4 carriers with Alzheimer's disease. J.

Cereb. Blood Flow Metab. 36, 216-227.

Haruwaka, K., Ikegami, A., Tachibana, Y., Ohno, N., Konishi, H., Hashimoto,

A., Matsumoto, M., Kato, D., Ono, R., Kiyama, H., Moorhouse, A.J.,

Nabekura, J., Wake, H., 2019. Dual microglia effects on blood brain barrier

permeability induced by systemic inflammation. Nat Commun 10, 5816.

Hui, J., Feng, G., Zheng, C., Jin, H., Jia, N., 2017. Maternal separation exacerbates Alzheimer's disease-like behavioral and pathological changes in adult APPswe/PS1dE9 mice. Behav. Brain Res. 318, 18-23.

Iadecola, C., 2017. The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease. Neuron 96, 17-42.

Kitamura, T., Ogawa, S.K., Roy, D.S., Okuyama, T., Morrissey, M.D., Smith, L.M., Redondo, R.L., Tonegawa, S., 2017. Engrams and circuits crucial for systems consolidation of a memory. Science 356, 73-78.

Mawuenyega, K.G., Sigurdson, W., Ovod, V., Munsell, L., Kasten, T., Morris, J.C., Yarasheski, K.E., Bateman, R.J., 2010. Decreased clearance of CNS beta-amyloid in Alzheimer's disease. Science 330, 1774.

McKim, D.B., Weber, M.D., Niraula, A., Sawicki, C.M., Liu, X., Jarrett, B.L., Ramirez-Chan, K., Wang, Y., Roeth, R.M., Sucaldito, A.D., Sobol, C.G., Quan, N., Sheridan, J.F., Godbout, J.P., 2018. Microglial recruitment of IL-1beta-producing monocytes to brain endothelium causes stress-induced anxiety. Mol. Psychiatry 23, 1421-1431.

Montagne, A., Nation, D.A., Pa, J., Sweeney, M.D., Toga, A.W., Zlokovic, B.V., 2016. Brain imaging of neurovascular dysfunction in Alzheimer's disease. Acta Neuropathol. 131, 687-707.

Niraula, A., Wang, Y., Godbout, J.P., Sheridan, J.F., 2018. Corticosterone Production during Repeated Social Defeat Causes Monocyte Mobilization from the Bone Marrow, Glucocorticoid Resistance, and Neurovascular Adhesion Molecule Expression. J. Neurosci. 38, 2328-2340.

Nortley, R., Korte, N., Izquierdo, P., Hirunpattarasilp, C., Mishra, A., Jaunmuktane, Z., Kyrargyri, V., Pfeiffer, T., Khennouf, L., Madry, C., Gong, H., Richard-Loendt, A., Huang, W., Saito, T., Saido, T.C., Brandner, S., Sethi, H., Attwell, D., 2019. Amyloid beta oligomers constrict human capillaries in Alzheimer's disease via signaling to pericytes. Science 365.

Norton, M.C., Smith, K.R., Ostbye, T., Tschanz, J.T., Schwartz, S., Corcoran, C., Breitner, J.C., Steffens, D.C., Skoog, I., Rabins, P.V., Welsh-Bohmer, K.A., 2011. Early parental death and remarriage of widowed parents as risk factors for Alzheimer disease: the Cache County study. Am. J. Geriatr.

Psychiatry 19, 814-824.

Radford, K., Delbaere, K., Draper, B., Mack, H.A., Daylight, G., Cumming, R., Chalkley, S., Minogue, C., Broe, G.A., 2017. Childhood Stress and Adversity is Associated with Late-Life Dementia in Aboriginal Australians. Am. J. Geriatr. Psychiatry 25, 1097-1106.

Rattray, I., Scullion, G.A., Soulby, A., Kendall, D.A., Pardon, M.C., 2009. The occurrence of a deficit in contextual fear extinction in adult amyloid-over-expressing TASTPM mice is independent of the strength of conditioning but can be prevented by mild novel cage stress. Behav. Brain Res. 200, 83-90.

Rincon-Cortes, M., Sullivan, R.M., 2014. Early life trauma and attachment: immediate and enduring effects on neurobehavioral and stress axis development. Front. Endocrinol. (Lausanne) 5, 33.

Saito, T., Matsuba, Y., Mihira, N., Takano, J., Nilsson, P., Itohara, S., Iwata, N., Saido, T.C., 2014. Single App knock-in mouse models of Alzheimer's disease. Nat. Neurosci. 17, 661-663.

Seifan, A., Schelke, M., Obeng-Aduasare, Y., Isaacson, R., 2015. Early Life Epidemiology of Alzheimer's Disease - A Critical Review. Neuroepidemiology 45, 237-254.

Sweeney, M.D., Ayyadurai, S., Zlokovic, B.V., 2016. Pericytes of the neurovascular unit: key functions and signaling pathways. Nat. Neurosci. 19, 771-783.

Wen, Y., Yang, S., Liu, R., Simpkins, J.W., 2004. Transient cerebral ischemia induces site-specific hyperphosphorylation of tau protein. Brain Res. 1022, 30-38.

Xu, H., Rajsombath, M.M., Weikop, P., Selkoe, D.J., 2018. Enriched environment enhances ß-adrenergic signaling to prevent microglia inflammation by amyloid-ß. EMBO Mol. Med. 10.

Yamada, J., Jinno, S., 2013. Novel objective classification of reactive microglia following hypoglossal axotomy using hierarchical cluster analysis. J. Comp. Neurol. 521, 1184-1201.

Yousef, H., Czupalla, C.J., Lee, D., Chen, M.B., Burke, A.N., Zera, K.A., Zandstra, J., Berber, E., Lehallier, B., Mathur, V., Nair, R.V., Bonanno, L.N., Yang, A.C., Peterson, T., Hadeiba, H., Merkel, T., Korbelin, J., Schwaninger,

1   M., Buckwalter, M.S., Quake, S.R., Butcher, E.C., Wyss-Coray, T., 2019.

2   Aged blood impairs hippocampal neural precursor activity and activates

3   microglia via brain endothelial cell VCAM1. Nat. Med. 25, 988-1000.

4   Zlokovic, B.V., 2011. Neurovascular pathways to neurodegeneration in

5   Alzheimer's disease and other disorders. Nat. Rev. Neurosci. 12, 723-738.

Figure Legends

Figure. 1 Maternal separation induces Alzheimer's disease-like behavior and amyloid beta

plaque formation.

(A) Time schedule for maternal separation and other investigations. (B) Spatial memory was

measured using an object location test. Behavioral layout of the object recognition test for the

four experimental groups. Maternal-separated $App^{NL-G-F/wt}$ mice showed impairment of spatial

memory. *p < 0.05 Maternal-separated $App^{NL-G-F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$

mice; n = 10/ Non- maternal-separated $App^{wt/wt}$ mice, n = 13/ Non-maternal-separated App^{NL-G-}

$^{F/wt}$ mice, n = 8/ Maternal-separated $App^{wt/wt}$ mice, n = 11/ Maternal-separated $App^{NL-G-F/wt}$ mice

(Dunnett's test). (C) Contextual memory and cued memory were measured using a fear-

conditioning test. Behavioral layout of the fear conditioning test for the four experimental

groups. Maternal-separated $App^{NL-G-F/wt}$ mice showed impairment of contextual memory.

Contextual test: Main effect of group, $F_{(3,68)}$ = 8.4507, p = 0.0001; main effect of time, $F_{(4,272)}$ =

5.3835, p = 0.0003; interaction of group × time, $F_{(12,272)}$ =1.0939, p = 0.3651; ** p < 0.01

Maternal-separated $App^{NL-G-F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice, *** p < 0.001

Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non-maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice; n = 19/ Non-

maternal-separated $App^{wt/wt}$ mice , n = 24/ Non-maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice , n = 10/

Maternal-separated $App^{wt/wt}$ mice , n = 22/ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice (two-way

repeated ANOVA followed by Holm's test). (D) The formation of amyloid beta plaque detected

by 82E1-biotin (cyan). The formation of amyloid beta plaque was facilitated by maternal

separation in maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice. *** $p < 0.001$ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$

mice vs other groups n = 4/each group (Tukey-Kramer test). Scale bar = 100 μ m.

Figure. 2 Maternal separation induced impairment to the vascular system.

(A) The health of the vascular system was evaluated by tomato-lectin labelling of endothelial

cells (green) and CD13 labelling of pericytes (magenta). Maternal separation induced lower

pericyte coverage in $App^{wt/wt}$ mice and $App^{NL-G-F/wt}$ mice. * p < 0.05, Maternal-separated $App^{wt/wt}$

vs non-maternal-separated $App^{wt/wt}$ mice, ** p < 0.01 Maternal-separated $App^{NL-G-F/wt}$ vs non-

maternal-separated $App^{wt/wt}$ mice, ** p < 0.01, Maternal-separated $App^{wt/wt}$ vs non-maternal-

separated $App^{NL-G-F/wt}$ mice, *** p < 0.001 Maternal-separated $App^{NL-G-F/wt}$ vs non-maternal-

separated $App^{NL-G-F/wt}$ mice. Furthermore, maternal separation also induced a narrowing of

vascular diameter in both phenotypes. * p < 0.05 Maternal-separated $App^{wt/wt}$ mice vs non-

maternal-separated $App^{wt/wt}$ mice, *** p < 0.001 Maternal-separated $App^{NL-G-F/wt}$ mice vs non-

maternal-separated $App^{wt/wt}$ mice, ** p < 0.01 Maternal-separated $App^{NL-G-F/wt}$ mice vs non-

maternal-separated $App^{NL-G-F/wt}$ mice; n = 3/each group (Tukey-Kramer test). Scale bar = 20 μ

m. (B) Microglia (White arrow in Left image) were detected by microglial labelling with Iba1

(magenta) and DAPI (cyan). Vessel-associated microglia (Grey square in Right image) were

detected by microglial labelling with Iba1 (magenta) and endothelial cell labelling with tomato-

lectin (green). In maternal-separated $App^{NL-G-F/wt}$ mice, vessel associated microglia were

increased. * p < 0.05 Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non- maternal-separated $App^{wt/wt}$

mice; n = 3/ Non-maternal-separated $App^{wt/wt}$ mice, n = 3/ Non-maternal-separated $App^{NL\text{-}G\text{-}F/wt}$

mice, n =3/ Maternal-separated $App^{wt/wt}$ mice, n =4/ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice

(Dunnett's test). (C) The integrity of the BBB was evaluated by tomato-lectin labelling of

endothelial cells (green) and fibrinogen (magenta). Maternal separation caused accumulation of

fibrinogen outside of vessels in maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice. * p < 0.05 Maternal-

separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice; n = 3/each group (Dunnett

test). Scale bar = 20 μ m. (D) The distance from the amyloid beta plaques to vessels and the

area of amyloid beta plaque were determined using 82E1-biotin labelling of amyloid beta

(magenta) and tomato-lectin labelling of endothelial cells (green). The area of amyloid beta

plaque showed a significant association with the distance from the amyloid beta plaque to

vessels in non-maternal-separated and maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice. Small amyloid beta

plaques are indicated with cyan squares, and amyloid beta plaques with a size midway between

small and big amyloid beta plaques are indicated by grey squares. $r^2 = 0.987186$ p < 0.001; n =

12 (polynomial regression). Scale bar = 20 μ m.

Figure. 3 Maternal separation and genetic factors activated and changed the morphology of

microglia.

(A)   Microglial morphology and activation were detected by labelling microglia with Iba1

(magenta) and labelling activation with CD68 (cyan). Maternal separation increased in activated

microglia in $App^{wt/wt}$ and $App^{NL-G-F/wt}$ mice. * p < 0.05 Maternal-separated $App^{wt/wt}$ mice vs non-

maternal-separated $App^{wt/wt}$ mice, * p < 0.05 Maternal-separated $App^{NL-G-F/wt}$ mice or non-

maternal-separated $App^{wt/wt}$ mice. * p < 0.05 Maternal-separated $App^{NL-G-F/wt}$ mice or non-

maternal-separated $App^{NL-G-F/wt}$ mice (Tukey-Kramer test). Furthermore, occupancy of soma and

dendrites in maternal-separated $App^{wt/wt}$ and maternal-separated $App^{NL-G-F/wt}$ mice was lower. n =

3/ Non-maternal-separated $App^{wt/wt}$ mice, n = 3/ Non-maternal-separated $App^{NL-G-F/wt}$ mice, n

=3/ Maternal-separated $App^{wt/wt}$ mice, n =4/ Maternal-separated $App^{NL-G-F/wt}$ mice (Tukey-

Kramer test). Scale bar = 10 μ m. (B) Detailed morphological analysis was performed using

the ImageJ plugin AnalyzeSkeleton. Images show examples of the outlines of microglia.

Maternal separation and APP mutation decreased dendritic terminals in $App^{wt/wt}$ and $App^{NL-G-F/wt}$

mice. * p < 0.05 Non-maternal-separated $App^{NL-G-F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$

mice, ** p < 0.01 Maternal-separated $App^{wt/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice,

***$p < 0.001$ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice

(Dunnett test). Furthermore, the total branch length in maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice and

maternal-separated $App^{wt/wt}$ mice were shorter. * $p < 0.05$ Maternal-separated $App^{wt/wt}$ mice vs

non-maternal-separated $App^{wt/wt}$ mice, ** $p < 0.01$ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non-

maternal-separated $App^{wt/wt}$ mice. The circularity of microglia increased in only the maternal-

separated $App^{NL\text{-}G\text{-}F/wt}$ mice. ** $p < 0.01$ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non-maternal-

separated $App^{wt/wt}$ mice. The solidity of microglia increased in only the maternal-separated

$App^{NL\text{-}G\text{-}F/wt}$ mice. ** $p < 0.01$ Maternal-separated $App^{NL\text{-}G\text{-}F/wt}$ mice vs non-maternal-separated

$App^{wt/wt}$ mice; n =4/each group (Dunnett test).

Figure. 4. Alzheimer's disease-like phenotypes did not begin during adolescence in maternal-

separated App$^{NL-G-F/wt}$ mice, and impairment of the vascular system did not begin during

adolescence in all groups.

(A) Spatial memory was measured using an object location test. Maternal-separated $App^{NL-G-F/wt}$

mice showed normal cognitive function. n = 9/ Non-maternal-separated $App^{wt/wt}$ mice, n = 11/

Non-maternal-separated $App^{NL-G-F/wt}$ mice, n = 12/ Maternal-separated $App^{wt/wt}$ mice, n = 11/

Maternal-separated $App^{NL-G-F/wt}$ mice (B) The formation of amyloid beta plaque detected by

82E1-biotin labelling of amyloid beta. No amyloid beta plaque was detected in maternal-

separated $App^{NL-G-F/wt}$ mice. n = 3/each group (C) The health of the vascular system was

evaluated by tomato-lectin labelling of endothelial cells (green) and CD13 labelling of pericytes

(magenta). All groups showed normal pericyte coverage and a healthy vascular diameter. n =

3/each group. Scale bar = 20 μ m.

Figure 5. Maternal separation and genetic factors induced morphological changes in microglia

and maternal separation in combination with genetic factors led to abnormality in the stress

response during adolescence.

(A) Microglial morphology and activation were detected by labelling with Iba1 (magenta) and

CD68 (cyan), respectively. Increased activation of microglia was not observed in maternal-

separated $App^{wt/wt}$ and maternal-separated $App^{NL-G-F/wt}$ mice. n = 3/each group. Scale bar = 10 μ

m. (B) Detailed morphological analysis was performed using the ImageJ. Maternal separation

and APP mutation decreased dendritic terminals in $App^{wt/wt}$ and $App^{NL-G-F/wt}$ mice. *** p < 0.001

Non-maternal-separated $App^{NL-G-F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice, ** p < 0.01

Maternal-separated $App^{wt/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice, ** p < 0.01

Maternal-separated $App^{NL-G-F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice (Dunnett test). (C)

Microglia (white arrow in left image) were detected by microglial labelling with Iba1 (magenta)

and DAPI (cyan). Vessel-associated microglia (Grey square in right image) were detected by

microglial labelling with Iba1 (magenta) and endothelial cell labelling with tomato-lectin

(green). The numbers of vessel-associated microglia were similar across the four experimental

groups. n = 3/each group. Scale bar = 20 μ m. (D) The protein level of TNF-R1 was higher in

maternal-separated $App^{wt/wt}$ and maternal-separated $App^{NL-G-F/wt}$ mice. p = 0.058 Maternal-

separated $App^{wt/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice, p = 0.054 Maternal-separated

$App^{NL-G-F/wt}$ mice vs non-maternal-separated $App^{wt/wt}$ mice n = 3/each group (Dunnett test). (E)

The response to novel cage stress was measured to assess an effect of maternal separation on the

HPA axis. Not only $App^{NL-G-F/wt}$ mice but also $App^{wt/wt}$ mice showed that increased stress caused

by maternal separation induced corticosterone release. Maternal-separated $App^{NL-G-F/wt}$ mice: F

$_{(1,2)}$ = 75.414, p = 0.0130; Maternal-separated $App^{wt/wt}$ mice: $F_{(1,2)}$ = 22.035, p = 0.0425. * p <

0.05 vs corticosterone before novel cage stress; n = 3/each group (Paired t-test).

(A)

Maternal separation
or
Control

P0 P2   P15   P45   P120

Behavioral test
Tissue and blood
sampling

Behavioral test
Tissue sampling

Control

Maternal
separation

(B)

Day 1   Day 2   Day 3

Familiar place

Novel place

Discrimination index

0.6

-0.6

3-4 months

$App^{wt/wt}$
□ Control
■ Maternal separation
$App^{NL-G-F/wt}$
□ Control
■ Maternal separation

(C)

Day 3 Contextual test

Freezing time (%)

1   5

Day 4 Tone-cued test

Freezing time (%)

1   10   Time (s)

3-4 months

$App^{wt/wt}$
□ Control
■ Maternal separation
$App^{NL-G-F/wt}$
□ Control
■ Maternal separation

(D)

$App^{wt/wt}$ $App^{NL-G-F/wt}$

Control

Maternal
separation

82E-1 (Aβ)

Aβ plaque in
prefrontal cortex (%)

2.5

***
***
***

4 months

$App^{wt/wt}$
□ Control
■ Maternal separation
$App^{NL-G-F/wt}$
□ Control
■ Maternal separation

(A)

$App^{wt/wt}$ $App^{NL-G-F/wt}$

Control

Maternal separation

CD68 CD68 (activation marker)/Iba1 (microglia)

4 months
$App^{wt/wt}$
□ Control
■ Maternal separation
$App^{NL-G-F/wt}$
□ Control
■ Maternal separation

% of reactive microglia

Number of microglia

% of Soma and dendrite area

p = 0.0512
p = 0.0109

(B)

$App^{wt/wt}$ $App^{NL-G-F/wt}$

Control

Maternal separation

Dendrite terminal /cell

Circularity

Total branch length (μm)

Solidity

3-4 months
$App^{wt/wt}$
□ Control
■ Maternal separation
$App^{NL-G-F/wt}$
□ Control
■ Maternal separation

(A)

Discrimination index

1.5 months
□ Control
■ Maternal separation
App $^{NL-G-F/wt}$
□ Control
■ Maternal separation

(B)

Aβ plaque in prefrontal cortex (%)

1.5 months
□ Control
■ Maternal separation
App $^{NL-G-F/wt}$
□ Control
■ Maternal separation

(C)

App $^{wt/wt}$ *App* $^{NL-G-F/wt}$

Control

Maternal separation

% of pericyte coverage

Vascular diameter (μm)

1.5 months
□ Control
■ Maternal separation
App $^{NL-G-F/wt}$
□ Control
■ Maternal separation

tomato-lection (Vessel)/CD13 (Pericyte)