1

Evolutionary Genetic Species Detected in Prokaryotes by Applying the K/θ

Ratio to DNA Sequences

Using K/θ to delimit bacterial species

C. William Birky, Jr.

Department of Ecology and Evolutionary Biology, The University of Arizona, Tucson,

Arizona, United States of America

Abstract

In previous papers in this series, I described the "evolutionary genetic species

concept" which is based on population genetic theory and should be applicable to any

organism. I also described a species criterion, the K/θ ratio, that delimits independently

evolving evolutionary species based on single-gene sequences. I then illustrated its

application to sexual and asexual eukaryotes. Here, I show how the evolutionary genetic

species concept and the K/θ ratio can be applied to bacteria, using the sequences from

some genes of the core genome that are rarely, if ever, transferred horizontally between

different species. This completes the demonstration that K/θ is a general method for

species delimitation, applicable to all organisms. Also, it adds to the evidence that

bacteria have species in the most general sense, even though they have the ability to

exchange genes across species boundaries. Finally, I show that a published critique of the

use of $K/\theta \geq 4$ as a criterion for independently evolving species rests on two errors in the

application of population/evolutionary genetic theory.

34 Funding: The author's personal funds were the sole support of the research, and the

35 author was solely responsible for study design, data collection and analysis, decision to

36 publish, and preparation of the manuscript.

37 Competing Interests: The author has declared that no competing interests exist.

38 * E-mail: birky@u.arizona.edu

Introduction

A prominent feature of the biological world is discontinuous variation, with

clusters of similar organisms differing significantly from organisms in other clusters.

These clusters, called species and given different binomial names, were originally

recognized by their different phenotypes. With the development of population genetic

theory, species have been recognized as populations with distinct gene pools underlying

their phenotypic differences. When gene sequences are used to reconstruct evolutionary

history in the form of phylogenetic trees, different species appear as different clades of

similar sequences separated by deep gaps in the tree.

This background is widely, but not universally, accepted. Doubters often focus on

the gradual divergence of sister populations from each other, wondering at what point the

y should be called different species. Even among those who agree that species are real,

distinct populations, there is disagreement on how to define and recognize them, given

the gradual divergence of recently separated populations. Many authors simply point to

"deep gaps" in a phylogenetic tree, usually of gene sequences, as evidence for different

species, but this raises the question: "How deep is deep enough?"

This question can be answered by thinking of species as independently evolving

populations. This is arguably the most general model of species (species concept),

applicable in principle to all organisms and consistent with population and evolutionary

genetic theory [1-8]. Different species are populations that evolve independently of each

other, in the sense that they are independent arenas for mutation, natural selection, and

random genetic drift. In other words, a new mutant allele arising in one population cannot

spread to the other, or if it does move from one species to the other by hybridization, it is

very unlikely to be fixed in that population. Moreover, if a shared allele is fixed or lost in

one population by drift and/or selection, this does not affect the probability that it will be

fixed or lost in the other. I call this the Evolutionary Genetic Species Concept (EGSC) to

distinguish it from the Evolutionary Species Concept (EvSC) of Simpson (1). The EvSC

as it is usually worded requires that we know, or assume, that the populations in question

will continue to evolve independently in the indefinite future. The requirement for future

independence should not be part of a species concept because is unknowable; for

instance, a trait difference required for adaptation of species A to a different niche from

species B could be lost in the future. (However, it should be noted that in 1953 Simpson

did allow that speciation might occur in allopatry without ecological differences in "some

probably rare cases" [1, p. 171].

In practice, a pair of sister species are recognized as soon as they have diverged to

the point where samples of individuals can be shown to come from two independently

evolving populations. This can be done using a number of different criteria for

independent evolution. Prominent examples that have been applied to prokaryotes as well

as eukaryotes are reproductive isolation and ecological isolation, both of which obviously

result in independent evolution of the isolated populations. However, both of these

species criteria have conceptual and practical problems. Neither one is generally

applicable; reproductive isolation is relevant only for organisms with significant amounts

of sexual reproduction, and neither criterion includes the possibility that species may be

evolving independently simply because they are physically isolated from each other.

Moreover, neither is readily applicable to large-scale biodiversity surveys such as the

Barcode of Life based on DNA sequences of one or a few genes isolated from many

individuals.

Although DNA sequences are widely used to delimit species in prokaryotes, there

is a great deal of controversy about how this should be done, and even about whether

bacteria really have species. Some of this controversy mirrors similar debates about

species in eukaryotes. However, much of it arises from the phenomenon of horizontal

gene transfer (HGT), in which segments of genes, whole genes, and even large segments

of chromosomes are transferred between distantly related bacteria, in addition to the more

common forms of bacterial gene exchange via conjugation, transformation, and

transduction. This problem can be solved by distinguishing between core genes and

auxiliary genes [9,10] The core genes are responsible for basic functions such as DNA

and protein synthesis and metabolic pathways which are common to all members of

closely related populations adapted to similar niches. Auxiliary genes include genes that

confer pathogenicity to specific hosts, resistance to certain antibiotics, or adaptation to a

novel ecological niche. Auxiliary genes are often exchanged between distantly related

bacteria, but core genes are not.

In this paper, as in previous publications [3-8], I argue that the EGSC can be applied

to all organisms, sexual and asexual, prokaryote and eukaryote. I also show how the K/θ

ratio, first used to discover and delimit EG species in eukaryotes based on DNA sequence

data from a single gene, can be applied to bacteria if a core protein-coding gene is used.

In particular, it could be applied to large-scale diversity surveys of bacteria using such a

gene. Such surveys are routine for eukaryotes, and can now be done with bacteria

regardless of whether they can be cultured in the lab, thanks to the development of

methods that allow amplification of genes from single bacterial cells isolated by dilution

or flow cytometry from environmental samples (e.g. [11,12]]. The successful

identification of evolutionary genetic species in bacteria using the K/θ ratio (this paper)

as well as methods such as the Generalized Mixed Yule Coalescent (GMYC; [4,13]) adds

to the growing literature that demonstrates that EG species are a general phenomenon

found in varying degrees to all organisms, sexual and asexual, prokaryotes and

eukaryotes.

The K/θ test is designed to distinguish between two kinds of clades: (i) those

separated by shallow gaps formed within a single species by random genetic drift or

transient physical isolation, and (ii) those separated by deep gaps produced by adaptation

to different ecological niches and/or by long-term physical isolation, i.e. different species.

Simulations [14] and population genetic theory [15] show that 95% of the gaps between

clades within a species are expected to have a depth of < 4N$_e$ generations, while gaps

between species clades can be of any depth. It is thus possible to recognize a pair of

clades as different species, with probability ≥ 0.95, if they are separated by gaps of ≥ 4N$_e$

generations. At that point the sequence difference between the two clades is $K \geq 8N_e\mu$

and the ratio K/θ ≥ 8Neu/2Neu ≥ 4 (3-8). (Barraclough's [16] claim to have found errors

in my use of K/θ is erroneous, as I show below in Discussion.)

This result is based on expected probabilities and does not account for uncertainty

introduced by finite sample sizes. The relevant sampling theory has been developed by

Noah Rosenberg [17, 18], who showed that it holds for any pair of sister clades, i.e.

reciprocally monophyletic samples n1 and n2, provided the sample sizes are at least n1 =

3 and n2 = 2. The power of these very small samples may seem surprising, but it arises

from their first being shown to be reciprocally monophyletic.

To illustrate how the K/θ ratio is used to detect evolutionary genetic species in

prokaryotes, I use the set of sequences of six different housekeeping/core genes

previously used by Wertz et al. [19, 20] to delimit species in enteric bacteria. This was

one of the first multilocus data sets used to discover bacterial species. It provides an

opportunity to apply the K/θ method to single genes and compare the results of using

different genes to each other and to the named taxonomic species defined by classical

phenotypic traits. Note that these data are used simply to demonstrate the application of

the K/θ method and are not intended to be definitive statements about these specific

bacterial species.

Materials and Methods

John Wertz kindly provided partial sequences of six housekeeping core genes

(*gapA*, *groEL*, *gyrA*, *ompA*, *pgi*, and *16S* rRNA) from seven named species of enteric

bacteria (*Citrobacter freundii* (abbreviated CF), *Enterobacter cloacae* (EB), *Escherichia*

coli (EC), *Hafnia alvei* (HA), *Klebsiella oxytoca* (KO), *Klebsiella pneumonia* (KP),

Serratia plymuthica (SP), and the outgroup *Vibrio cholera* (VC).

The aligned sequences for each gene were visualized in MacClade [21] where all

the sequences of the gene were trimmed to the same length and all internal sites with gaps

in any one sequence were removed. Consequently, all pairwise sequence differences for a

given gene are based on the same regions of the gene. Final sequence lengths were: *gapA*,

540 bp; *groEL*, 348 bp; *gyrA*, 634 bp; *ompA*, 433 bp; *pgi*, 242 bp; *16S*, 289 bp. One

internal gap of 27 bp in the *ompA* gene of the outgroup *V. cholerae* was retained because

to remove it from all sequences would cause an unnecessary loss of sequence information

in the ingroup.

To discover species using the trimmed sequences, I followed the procedure

outlined by Birky [8] for each gene separately. Briefly,

1. First, I used PAUP* [22] to find a bootstrapped neighbor-joining tree in which all

clades were supported by ≥70% of 1000 bootstrap replicas. A Neighbor-joining tree was

used to rearrange the sequences in the PAUP* file to reflect the order of sequences in the

tree; this greatly simplifies subsequent calculations in Excel, but note that the tree itself is

not used in subsequent calculations. Both corrected and uncorrected sequence

differences were then exported from PAUP* to an Excel spreadsheet where all

subsequent calculations were performed. Starting at the tips of the tree, each pair of sister

clades is tested separately in steps 2-7, to determine the probability that the clades are

samples from independently evolving species. This process is continued with deeper and

deeper clades until species are found. For clades that are members of a non-bifurcating

tree structure such as a polytomy (A, B, C) or ladder ((A,B)C), I compare A and B first,

then compare C to whichever one of those is closest to C.

2. For the two candidate clades in a pair, the mean pairwise difference d between

sequences is multiplied by the sample size correction n/(n-1) where n is the number of

sequences in the clade, to get the nucleotide diversity π. Note that nucleotide diversity

is defined as a function of the uncorrected pairwise distance. When $d = 0$, I used a non-

zero estimate of π by assuming that one sequence differs from the others by 1/L where L

is the sequence length; then $\pi = 2/Ln(n-1)$.

3. For each clade, I then estimate $\theta = 2Ne\mu$ by $\pi/(1 - 4\pi/3)$.

4. Calculate K = mean pairwise sequence difference between the two clades, corrected

for multiple hits.

5. Calculate K/θ for the pair of clades. When the values of θ for the two clades differ, I

use the larger value to get a conservative estimate of K/θ. If K/$\theta \geq 4$, I conclude that the

individuals in those clades came from different species, with probability ≥ 0.95.

6. The preceding steps are based on expected probabilities and do not account for

uncertainty introduced by finite sample sizes. This can be done as follows, using results

obtained by Noah Rosenberg [17]. Using the K/θ ratio and the numbers $n1$ and $n2$ of

individuals in the two clades, we want to find the probability that the individuals were

sampled from populations that have been evolving independently long enough to become

reciprocally monophyletic. This is best done using a table available on request from Noah

Rosenberg or me; alternatively, the values can be estimated from Figure 6 in Rosenberg

[17].

It is worth repeating that the phylogenetic tree used in step 1 is only used for two

purposes: (1) to facilitate the identification of sister clades, and (2) to inform the

organization of the Excel spreadsheet of pairwise sequence differences with the most

similar sequences together.

205 Results

The Wertz et al. [20] data consist of 5 protein- coding genes plus part of the 16S

RNA gene. I will first illustrate the application of the K/θ ratio by showing the detailed

analysis for *pgi*, the protein-coding gene that resolves the most species. The high

resolution suggests that this gene segregated before most of the other genes in most of the

speciation events, by chance. I will then summarize the results obtained with each of the

other genes. Following Wertz et al. [20], I consider the single isolate of *S. marcescens* as

belonging to the same taxonomic species as *S. plymuthica* because for all the genes in this

study it is identical or nearly so to the *S. plymuthica* isolates.

214 Finding species with *pgi*

The Neighbor-joining (NJ hereafter) tree in Figure 1 shows that the uncorrected

pgi sequences form a number of clades that supported that are seen in $\geq 63\%$ of 1000

bootstraps; most have 100% support. The taxonomic species *K. oxytoca*, KO; *E. cloacae*,

EB; *H. alvea*, HA; and *S. plymuthica* SP, SM all form clades with 100% bootstrap

support, but some of these in turn contain two or more strongly supported clades. I

applied the K/θ test to each well-supported pair of sister clades, starting at the bottom of

the tree in Figure 1. These clades are also well-supported in the NJ tree using corrected

sequence differences. The Excel table for *H. alvea* and *S. plymuthica* is shown in Table 1

as an example of the data.

.   — 0.01 substitutions/site

Fig 1. NJ tree of *pgi* sequences. CF, EC, KP, KO, EB, SM, and HA are taxonomic

species. Closed circles indicate clades that are different EG species delimited by the

K/method.

1.   The Excel table for *pgi* from isolates of *H. alvea* and *S. plymuthica* is shown in

229   Table 1. The full tables for all genes are available on request from the author. To

230   illustrate the procedure, I begin with clade HA2,4, which has weak support (69%

231   of bootstraps; Its sister group is the singlet HA3. Using the formula for a

maximum estimate of π and θ when d = 0, I calculate that θ = 0.00836 and from

the table K = 0.0021 so K/θ = 0.25 and these are not different species. This result

was obvious from inspection and normally would not warrant a calculation. Next

I ask if clade HA1-4 is a different species from clade (SM1,SP1,SP2). The

pairwise distances for SM1, SP1, and SP2 are zero while those for HA1-4 are not

all zero; I prefer to use the latter because they do not require any correction. For

this clade d = 0.00625, π = 0.00833, and θ = 0.00843. K is 0.03032 so K/θ =

36.0 and these are samples from different clades with probability > 0.988.

Because K/θ >> 4, these specimens are also samples from different independently

species with P >> 0.95. Note that these results are obvious by inspection of the

branch lengths in an NJ tree like that in Figure 1 in which branch lengths are

proportional to the number of substitutions. The branches connecting the (SM,SP)

clade to the HA clade are clearly more than four times the length of the branches

connecting any two specimens within the HA clade.

251   Table 1. Matrix of pairwise sequence differences for *pgi* from *Serratia plymuthica* and

252   *Hafnia alvea*.

	SM1	SP1	SP2	HA1	HA2	HA3	HA4
SM1	-						

SP1	0.0000	-				
SP2	0.0000	0.0000	-			
HA1	**0.3022**	**0.3022**	**0.3022**	-		
HA2	**0.3010**	**0.3010**	**0.3010**	0.0083	-	
HA3	**0.3087**	**0.3087**	**0.3087**	0.0125	0.0042	-
HA4	**0.3010**	**0.3010**	**0.3010**	0.0083	0.0000	0.0042	-

Uncorrected sequence differences d and corrected differences K (bold type).

2. The taxonomic species *C. freundii* (CF) forms two clades, CF1,5 and CF3,4, each with

100% bootstrap support. Using Excel, I calculate the mean corrected sequence difference

between these clades, K = 0.07618. CF1,5 has the larger value of θ = 0.0437, giving K/θ

= 1.74. In Rosenberg's table, when K/θ = 1.7 and the sample sizes are 2 and 2, the

probability that these two clades are samples from reciprocally monophyletic populations

is only 0.57. Moreover, because K/θ is much less than 4, the probability that the samples

came from different species is low. This is the more conservative test in terms of

splitting species because it uses the larger of the two values of θ. On the other hand, the

clade CF3,4 has the smaller θ = 0.00167, giving K/θ = 4.5. From Rosenberg's table,

$P = 0.97$ that these are samples from different clades; furthermore, because K/$\theta > 4$, the

probability is very high that they are from different independently evolving species.

However, this is based on the less conservative estimate of θ. Because only one of the

two possible estimates of θ support the hypothesis of different species, I conclude that the

pgi gene shows weak support for two cryptic species within *C. freundii*. As we will see

below, some, but not all, other genes strongly support splitting. This suggests that there is

enough recombination in *C. freundii* to cause different genes to complete segregation at

different times, by chance.

3. The taxonomic species *E. coli* (EC) is also consists of a clade (EC1-5,ECMG) with

100% bootstrap support and a singleton EC6. Only the clade can be used to estimate θ,

but since the 6 sequences in the clade are identical, π and thus θ cannot be calculated

directly. For these cases, one can conservatively estimate these values by assuming that

one of the 6 sequences actually differs from the others at a single site. Then using the

calculation given in Materials and Methods, one can calculate $\pi = 0.001377$, $\theta = 0.00166$,

and K = 0.07000 so K/θ = 42.3 and there is strong evidence that the clades EC1-5,MG

and EC6 are samples from different clades and from different cryptic species. This

procedure is likely to overestimate π and θ and hence underestimate K/θ, so in this sense

it is conservative with respect to splitting species.

4. Next we consider the clade consisting of *pgi* from KP isolates. Starting at the tips, I

compared (KP3,4) with KP5. Using the same procedure as above to get a non-zero value

for π, K/θ = 8.38, providing evidence that these are samples from two different clades

and species. Comparing (KP3,4) to the other singleton, KP2, I find K/θ = 11.6 and these

are also from different species. Thus KP is divided into three species by these gene

sequences.

5. Finally, the sequence data from this gene confirm that the isolates identified as *K.*

oxytoca (KO) and E. cloacae (EB) are evolutionary genetic species.

To summarize, the sequences of *pgi* confirm that KO, HA, EB, and SM,P are

species; split EC into 2 species; and split KP into 3 species.

Finding species with *groEL*

The *groEL* gene (Figure 2) confirms that KO, EB, SM, and HA are species. It

splits CF into two species, CF1,5 and CF3,4. EC is confirmed as a separate species from

its sister clade CF2-4 if one uses θ from CF2-4 , but not if one uses θ from EC. Although

groEL sequences split KP into KP1,4 and KP2,6 if one uses θ from KP2,6 they do not if

one uses the larger θ from KP1,4.

Fig 2. NJ tree of *groEL* sequences. CF, EC, KO, EB, KP, HA, and SM,SP are taxonomic

species. Closed circles indicate EG species.

Finding species with *gyrA*

301   The gene *gyrA* (Figure 3) confirms that KO, EB, EC, HA, and SM are

302   Evolutionary Genetic Species. The sequences of this gene split the taxonomic species CF

303   into two EG species, CF1,5 and CF2-4, and also split KP into KP1,4 and KP2,6.

Fig 3. NJ tree of *gyrA* sequences. Closed circles indicate EG species.

Finding species with *gapA*

The *gapA* gene (Figure 4) confirms that CF, KO, EC, SM,SP, EB, and HA are

different species. KP consists of two clades, KP1,3,4 and KP2,6, and a singlet KP5 which

are samples from three different species.

Fig 4. NJ tree of *gapA* sequences. Closed circles indicate EG species.

Finding species with *ompA*

The *ompA* gene (Figure 5) provides evidence that the CF, EC, EB, HA, and

SP,SM sequences each represent a single species. Unlike some of the other genes, it does

not split any of these species. In contrast, the KO and KP sequences are polyphyletic.

Although the KO sequences are very similar with a diversity commensurate with that of

other species in these data, the KP sequences are very divergent from each other.

Moreover, KP5 is basal to KP1-4 as well as to KO; thus KP is polyphyletic, suggestive of

incomplete lineage sorting.

Fig 5. NJ tree of *ompA* sequences. Closed circles indicate EG species (these will be

changed to red to match the other figures).

Finding species with *16S*

The *16S* sequences of the taxonomic species *E. coli* and *H. alvia* contained

regions that were not present in the other species and for which I could find no

explanation, so these two taxonomic species were omitted from the analysis of the *16S*

sequences. Also, all pairwise differences among the ingroup sequences were ≤ 0.063,

including those between species, so no corrections for multiple hits were applied. The

K/θ test verified that the CF1,5, KO, CF2-4, EB, and KP sequence sets, and the SP1

singlet, each represent evolutionary genetic species with probabilities >0.99. However,

the CFl and CF5 sequences did not; the clade was supported by only 56% of bootstrap

replicas, so no K/θ test was done. The NJ tree is shown in Fig 6.

Fig 6. NJ tree of *16S* sequences. Closed circles indicate EG species.

Discussion

A huge literature has accumulated on the nature of species and how they should be

defined and discovered (reviewed in [23,24]. Even the very existence of species has been

debated, largely because the process of speciation is usually a gradual one in which two

populations of one species diverge from each other until they reach the point where they

can be recognized as different species. Nowhere has the debate been more intensive than

in the prokaryotes. Although the Bacteria and Archaea came later to the party than the

Eukaryota, they brought their own entertainment with them, in the form of novel means

of sharing genes with close relatives (transformation, transduction, and conjugation).

Prokaryotes can even receive new kinds of genes from distant relatives (Horizontal Gene

Transfer, hereafter HGT).

My focus here, and in several preceding papers on discovering species in asexual

and sexual eukaryotes [3-8] is on two problems: (i) identifying a model of species, i.e. a

species concept, that is general enough to cover all organisms, whether prokaryotes or

eukaryotes, and all degrees and kinds of sexuality; and (ii) based on the model, testing a

general method of assigning specimens to the same or different species (a species

criterion), using gene sequences.

16SrRNA genes are found in all organisms and evolve slowly enough to be useful

in constructing phylogenetic trees across the tree of life. They have also been used to

define Operational Taxonomic Units in bacteria, especially in environmental sequencing

projects. Unfortunately, the same evolutionary conservatism that makes rRNA genes

useful for deep taxonomy also often means that they have too little variation within and

between closely-related species to be useful for species delimitation [25, 26]. Using

16SrRNA sequences from nearly the same set of isolates as in the present study, Riley *et*

al. [19, 20] found that all of the taxonomic species formed clades with >50% bootstrap

support in a maximum likelihood tree, except CF1,5 which was polyphyletic. My results

with the 16S sequences differ; CF2-4 form a species distinct from CF1 and CF5, which

are singlets.

Eckert and colleagues [26] used nearly complete 16S sequences to look for a

barcode gap in Cyanobacteria, using the Automatic Barcode Gap Discovery (ABGD)

program to find gaps in pooled sequences from some but not all taxonomic groups.

However, rather than testing pairs of sister clades, Eckert et al. pooled between 34 and

2448 sequences from each of 16 genera. This will obscure the barcode gap if different

pairs of species have gaps in different places, i.e. if some species have values of θ that

overlap the values of K in other groups. It is also possible that some of the groups

examined by Eckert et al. have undergone a recent adaptive radiation such that K values

are very small, as the authors noted.

The results with the protein-coding genes and isolates EB, EC, HA, KO, and

SM/P are straightforward: each of the genes shows that these five taxonomic species,

discovered by traditional methods of bacterial taxonomy using morphology and various

phenotypic tests, are also EG species. The situation with CF is more complex. All five

genes show two clades, (CF1,5) and (CF2-4) with 89-100% bootstrap support except

gapA where the support for the (CF2-4) clade is only 64%. Moreover the K/θ ratio for

this pair of clades is greater than 4, signifying different species, for *gyrA*, *pgi*, and *groEL*

but not for *gapA* or *ompA*.

Deep gaps in trees separating species: how deep is deep

enough?

In the literature of new species descriptions, and in papers proposing the splitting

of species, it is common to refer to deep gaps between clades in phylogenetic trees as

evidence that the clades should be assigned to different species. The decision as to

whether the gap between two clades A and B is deep enough to assign A and B to

different species, as opposed to indicating a relatively shallow gap within a single

species, is generally made using the systematist's intuition. Occasionally the systematist

compares the gap between A and B to the gap between named species C and D,

concluding that A and B are different species if the gap between them appears to be as

deep or deeper than the gap between C and D. However, this assumes that species C and

D are well defined. The K/θ method avoids making these questionable assumptions by

providing a decision criterion that is soundly based on population genetic theory.

Not All Well-Supported Clades Are Species

400   These analyses illustrate the danger of assuming that all clades are species. For

401   example, in Fig 1 the clade EB1,3,5 has 90% bootstrap support and yet belongs to the

402   same species as EB4. If one studied *E. cloacae* EB only, one could easily draw a tree

403   with the Y axis stretched out, so that the branch connecting the clade to EB4 looks very

404   long and might appear to show two species connected by a long branch.

Applicability of the K/θ method

It is clear that the Evolutionary Genetic Species Concept is applicable to bacteria.

It is also clear that within the framework of the EGSC, species can be discovered using

DNA sequences analyzed with the K/θ method. It is important to note that the K/θ test

was deliberately designed to work with a single gene or sequence. This is appropriate

because sequencing multiple genes from every specimen is too expensive and time-

consuming for routine application in species discovery, and will continue to be too

expensive in the foreseeable future. It will be vastly more difficult, and often impossible,

in the case of environmental sequences, which are of necessity not tied to individual

organisms. If the sequences of two or more genes are used, it is important to remember

that in a sexual organism different genes may become reciprocally monophyletic at

different times. I recommend that the K/θ test be applied to each gene separately; if any

gene places the specimens in two different clades and $K/\theta \geq 4$, this would be strong

evidence that the clades represent different species. I do not recommend concatenating

gene sequences without first testing whether the signal from a gene that has not yet

segregated can obscure the signal from a gene that has become reciprocally

monophyletic, i.e. could cause the specimens to be incorrectly assigned to a single

species.

Multiple comparisons
423

I used a probability of 0.05 for hypothesis testing, understanding that when this is

applied to many different cases, I expect to make a wrong decision in approximately 5%

of the cases. This could be accounted for by using a correction such as the Bonferroni. On

the other hand, many of the the results are significant with a much higher probability. I

choose not to use a correction and simply accept the risk of a small number of wrong

conclusions, less than one in twenty for this work as a whole.

When small sample sizes can provide strong conclusions
430

Rosenberg [17] showed that small sample sizes can provide strong evidence that

the entire populations from which they were taken are reciprocally monophyletic. This is

because the probabilities are contingent upon the samples themselves being reciprocally

monophyletic. Of course this assumes that the samples are representative of the

populations, and do not, for example, come from a local sub-population that happens to

be very different from the rest of the population. Thus one must always bear in mind the

possibility that the specimens being studied are not representative. On the other hand, the

possibility of erroneously splitting a species into two species based on a non-

representative sample is likely to be small when K is very large. This is because, if a

large K gives a K/θ value less than 4, the θ value must also be much larger than the

generally observed values of < 0.05. There are parallels in traditional taxonomy, where

new species are often described on the basis of one or a few specimens, especially when

the new specimen is very different from the most similar described species. In this case

we usually assume that additional new specimens will not have intermediate phenotypes

that will cause us to combine the new specimen with a described species. If we had a

single specimen of an African lion and two specimens of an ocelot, traditional taxonomy

would probably recognize them as different species based on the absence of intermediate

phenotypes, even without additional evidence. If gene sequences from these specimens

had K/θ ≥ 4, this would indicate an absence of intermediate sequences and we would

have similar objective evidence that they were different species.

451   Use of trees and choice of tree-making methods

452   When implementing the K/θ method, the use of phylogenetic trees is limited to (i)

453   finding sister pairs of clades for comparison; (ii) guiding the arrangement of sequences in

454   the Excel spreadsheet to facilitate the calculation of K/θ; and (iii) spotting obvious cases

455   of K/θ based on branch lengths. I use Neighbor-joining for tree construction because it is

456   fast and easy to do with PAUP*, but presumably a different distance-based method could

457   be used.

458   Genes or phenotypes in bacterial species descriptions?

459   The bacterial rules of nomenclature do not specifically require phenotypic data.

However, the report of an *ad hoc* committee on species definition in bacteria [27] calls

for the use of qualitative phenotypic traits as well as 16S rDNA sequences to delimit

species. A quick survey of some recent papers in the *International Journal of Systematic*

and Evolutionary Microbiology shows that while rDNA sequences are now commonly

used in descriptions of new bacterial species, they are used essentially like a collection of

traits or to form phylogenetic trees. When these trees, or pairwise sequence differences,

or both are used to define trees, there is usually no attempt to apply a theoretically

justifiable method to distinguish sequence differences between species from differences

within species. Instead, this is based on the judgment or intuition of the investigators.

469 One gene or many?

There is a tendency in the recent literature to extoll the virtues of using many

genes to discover species. But using more than one gene has several drawbacks. First, it

costs more and will not be useful if one gene is sufficient to show that two samples come

from different species. Second, if one gene is a core gene, adding another gene which is

not a core gene and is involved in LGT can obscure the signal of speciation.

Ideally, to discover closely related species we would like to use the gene or genes

that segregated earliest in those particular populations; in this way we would catch

speciation in its early stages and discover closely related species. This is a real advantage

to using multiple gene sequences, which would increase the chance that at least one of

them would have segregated early. How important this depends on the amount of sexual

reproduction in the bacteria in question. At one extreme, if there is little or no

recombination, or extreme inbreeding, all genes will segregate at the same time and any

core gene should be as good as any other. At the other extreme, if there is extensive

outcrossing and recombination, different genes may segregate at very different times after

two populations begin evolving independently. In fact, a given gene may segregate early

in one such population and late in the sister population. This is a likely explanation for

the observation that CF1 and CF5 form a separate species from CF2, 3, and 4 for all

protein-coding genes except *ompA*. Stackebrandt et al. [27] speak favorably of using

protein-coding genes, claiming that there is a consensus that at least five such genes are

necessary, but there is no such consensus in the sense of a number that is agreed on or at

least accepted by nearly everyone in the field. More importantly, Stackebrandt et al. do

not consider the possibility that these genes may be at different stages of segregation,

which is the simplest explanation of the different trees supported by different genes

Wertz et al. data.

494   There is also a potential problem with concatenated gene sequences: a significant

495   K/θ ratio in one gene may be obscured by a small, insignificant K/θ ratio in another gene.

496   In other words, the signal of species divergence in one gene that segregated early after

497   speciation may be swamped by the signal in another gene that has not yet completed

498   segregation with 95% confidence.

499   I suggest that the best method for large-scale diversity studies would be to use a

500   single protein-coding gene to discover species and then, for groups of special economic,

501   medical, ecological, or evolutionary importance or interest, test additional genes to see

502   they separate more species pairs because they segregated earlier in those pairs. Then it

503   would often be useful to look for phenotypic or molecular markers that could be used to

504   identify those species without requiring sequencing.

Stability of EG species

506   A nice feature of species discovery using $K/\theta \geq 4$ is that additional specimens are

unlikely to fill in the gap between species and require them to be lumped. That is because

the large difference between K and θ reduces the probability that other specimens that are

intermediate between the two species and fill in the gap between them will ever be

discovered. Put another way, it is unlikely that we will find specimens and lineages that

join the two clades we identify as species. If this did happen, the single species produced

by joining the two clades would have exceptionally large nucleotide diversity. It seems

likely that, once two populations are assigned to different species using $K/\theta \geq 4$, those

species are unlikely to be lumped in the future. It is perhaps more likely that data from

additional genes or from new specimens will cause a species to be split in the future, and

this has been seen repeatedly in the literature. But this is true for species delimited by any

method, using any kind of data; in science, new information often forces us to change our

conclusions.

519   Other methods used to discover bacterial species with single

520   loci

Fixed cutoffs of 95%, 99%, or other levels of sequence similarity in rRNA genes

are widely used to delimit species or potential species in the bacterial literature. These

cutoffs share with $K/\theta \geq 4$ the advantage of allowing one to assign sequences from

uncultivated or uncultivatable bacteria, or environmental sequences, to species. They are

based mainly on the observed levels of sequence similarity among already described

species, and as such they are only good to the extent that the described species are real.

Moreover, cutoffs are generally assumed to apply to uniformly over all species, but this

assumes that all species have the same within-species variation, i.e. the same values of

nucleotide diversity; this is not true.

Most of the methods proposed for using DNA sequence data to delimit species are

global methods, i.e. in a phylogenetic tree of gene sequences from many specimens, they

look for the transition point from species divergence to divergence between individuals

within species. One of these, GMYC [13] uses coalescent theory to detect the transition

point. Barraclough et al. [28] used GMYC to discover species from environmental 16S

rDNA sequences of bacteria but the method has not been applied to sequences from

already-described taxonomic species of prokaryotes. Poisson Tree Picking (PTP) [29]

attempts to identify significant changes in the numbers of substitutions along branches,

which are expected to be significantly higher between species than within species. To

my knowledge, PTP has not been applied to prokaryotes.

Ecosim [30] detects species produced by selective sweeps during adaptation to

different habitats. It assumes a Poisson model of substitutions, in contrast to K/θ and

GMYC that allow more complex and presumably more realistic models to be used. The

theory behind Ecosim includes random drift and selective sweeps but not less drastic

forms of selection. AdaptML [31] is another local method. It identifies clades associated

with ecological parameters that might be relevant to ecological species. It detects species

only if they are adapted to different niches that have already been identified. These

methods will not detect species that evolve independently because they are physically

isolated as in the case of *Sulfolobus islandicus* in hot springs [32] unless they are also

adapted to different niches. In other words, they may fail to detect some evolutionarily

independent species.

These global methods are all subject to error if different species have different

levels of within-species variation, i.e. different values of nucleotide diversity. This is in

contrast to $K/\theta \geq 4$ which is a local method focusing on sister clades and thus on putative

sister species. Such pairwise comparisons have the advantage that sister species are more

likely to have similar values of θ. In fact, the problem of delimiting species in general

really is a local problem. We have no problem assigning specimens of the grizzly bear

and the mule deer to different species, but it is more difficult to decide whether the

Alaskan brown bear and polar bear are really different species, e.g. [33].

Microscopic organisms do have species

Rossberg et al. [34] claimed that organisms smaller than about 1mm can not have species,

which the present paper clearly refutes for bacteria. Rossberg *et al.* focused on

meiofauna, but their computer simulations apply equally to bacteria as well as to asexual

and sexual eukaryotes. Remarkably, they failed to cite any of the nearly two centuries of

taxonomic literature describing species in microscopic invertebrates, protists, algae,

fungi, etc. A likely source of their erroneous conclusion is their use of carrying capacities

instead of the effective population size, which is appropriate for the critical parameter

$N_e\mu$ and is very much smaller than the carrying capacity. In fact, their Figure 3 shows the

signal of species when they used values of the carrying capacity that are more in line with

known effective population sizes.

Limitations of the K/θ method

In order to study the diversity of cyanobacteria from hot springs in Yellowstone

National Park, Ward et al. [35] did environmental sequencing of clone libraries from the

16SrRNA/ITS region from PCR-amplified DNA. Their Figure 4 shows a neighbor-

joining tree of ITS sequences of bacteria identified as *Synechococcus*. They found 10

putative ecotypes of *Synechococcus*, which are thus different species adapted to different

niches. However, it is evident by inspection of the tree that only the three clades called A,

A', and B satisfy the $K/\theta \geq 4$ criterion. This could happen if most of the ecotypes have

diverged too recently to pass the the $K/\theta \geq 4$ criterion for independent evolution. In this

case ecotype analysis can divide species more finely because it delimits species at an

earlier stage in the speciation process.

581   Using K/θ in Formal Descriptions of New Species

I have shown that the K/θ ratio can be used to delimit evolutionary species in

bacteria, illustrating the procedure with an exemplar data set. However, this does not

mean that the species so delimited conform to current practice in bacterial systematics, or

that the species so detected can be formally described in conformance with the current

rules of taxonomic nomenclature for bacteria. Of course the same is true of species

delimited with the GMYC method that also detects EG species, or with the method of

Cohan [36], which detects a subset of EG species that are separated by ecology.

Unfortunately, the rules of bacterial systematics codified in the International Code of

Nomenclature of Bacteria are not explicitly based in population genetic theory, even

though they pay lip service to the BSC which defines species as reproductively isolated

populations in outcrossing sexual organisms.

593   Response to the critique of Barraclough [37]

Barraclough [37] claimed to have found two errors in my calculation that a K/θ

value of 4 or greater is strong evidence that samples belonging to two different

evolutionary species. First, citing Edwards and Beerli [38], he claimed that my "K from

the genealogy overestimates the true divergence time, however, due to sampling of the

polymorphism from the ancestral common ancestor species" and that K is equal to 2Tμ

plus the θ of the ancestor. However, Edwards and Beerli said that the addition of θ is

necessary when K denotes the time elapsed since two populations diverged, but it is not

appropriate when estimating the time T since allelic divergence, as I am doing here and in

previous papers using K/ θ. Second, Barraclough refers to "a mitochondrial marker with

θ = Nμ ." (In correspondence, he said that in this section he used N to refer to the

effective population size rather than the census size.) If this were true, then the upper

95% confidence limit of K/ θ would be 8. But by definition $\theta = 2N_e\mu$ [39]. This is

because the sequence difference between two gene copies in two specimens A and B

from the same species sharing a most recent common ancestor O is equal to the sequence

difference $N_e\mu$ between A and O plus the sequence difference between B and O, which is

also $N_e\mu$. Consequently, after $4N_e$ generations, $K/\theta = 8N_e\mu/2N_e\mu = 4$, and not 8.

Could we use K/ θ as a measure of the progress of speciation?

If we adopt the evolutionary species concept, then speciation is usually a gradual

process. (Exceptions can include rare dispersal to isolated sites such as islands, or

speciation by hybridization or polyploidization.) To the extent that speciation is gradual,

the K/ θ ratio is expected to gradually increase. Consequently, if we have two populations

that we suspect are undergoing speciation, K/ θ might be used as a measure of the

progress of speciation. It would presumably be non-linear, as suggested by the

nonlinearity of progress toward reciprocal monophyly of the two populations [17].

Species are universal

619 In this paper and previous papers in this series [3-8] I have shown that Evolutionary

620 Genetic species are found in bacteria and in both asexual and sexual eukaryotes. The

621 arguments in this paper should also apply to Archaea. I strongly suspect that viruses also

622 form such species, and that extraterrestrial organisms, if they exist, will likewise form

623 populations that evolve independently from each other.

Acknowledgements

626 I thank John Wertz for providing the aligned sequences.

References

1. Simpson GG The species concept. Evolution 1951; 5:285-298.

2. Wiens JJ. What is speciation and how should we study it? Am Nat 2004;163(6):914-
23.

3. Birky, CWJr, Wolf C, Maughan H, Herbertson L, Henry E. Speciation and selection
without sex. Hydrobiologia 2005; 546:29-45.

4. Barraclough TG, Birky CWJ, Burt A. Diversification in sexual and asexual
organisms. Evolution. 2003;57(9):2166-72.

5. Birky, CWJr, Adams J, Gemmel M, Perry J. Using population genetic theory and
DNA sequences for species detection and identification in asexual organisms PLoS
One 2010; 5:e10609.

6. Birky, CWJr, Ricci C, Melone G, Fontaneto D. Integrating DNA and traditional
taxonomy to describe diversity in poorly studied microscopic animals: new species of

the genus Abrochtha Bryce, 1910 (Rotifera: Bdelloidea: Philodinavidae). Zool J Lin

Soc 2011; 161:723-734.

7. Schön I, Pinto RL, Halse S, Smith AJ, Martens K, Birky CWJr. Cryptic species in

putative ancient asexual darwinulids (Crustacea, Ostracoda). PLoS One 2012;

7:e39844.

8. Birky CWJr. Species detection and identification in sexual organisms using

population genetic theory and DNA sequences. PLoS One 2013; 8:e52544.

9. Lan R, Reeves PR. Intraspecies variation in bacterial genomes: the need for a species

genome concept. Trends in Microbiology. 2000;8(9):396-401.

10. Lan R, Reeves PR. When does a clone deserve a name? A perspective on bacterial

species based on population genetics. Trends in Microbiology. 2001;9(9): 419-24.

11. Otteson EA, Hong JW, Quake SR, Leadbetter JR. Microfluidic digital PCR enables

multigene analysis of individual environmental bacteria. Science. 2006;314:1464-7.

12. Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Andeerson IJ, Cheng J-F, et al.

Insights into the phylogeny and coding potential of microbial dark matter. Nature.

2013;499:431-7.

13. Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, et al.

Sequence-based species delimiation for the DNA taxonomy of undescribed insects.

Syst Biol. 2006;55:1-15.

14. Avise JC. Molecular Markers, Natural History and Evolution. New York: Chapman

& Hall, Inc.; 1994.

15. Hudson RR, Coyne JA. Mathematical consequences of the genealogical species

concept. Evolution. 2002;56:1557-65.

16. Barraclough TG. The Evolutionary Biology of Species. Godfray CJ, Dunne JA,

Sheldon BC, editors. Oxford, United Kingdom: Oxford University Press; 2019.

17. Rosenberg NA. The shapes of neutral gene genealogies in two species: Probabilities

of monophyly, paraphyly, and polyphyly in a coalescent model. Evolution.

2003;57(7):1465-77.

18. Rosenberg NA. Statistical tests for taxonomic distinctiveness from observations of

monophyly. Evolution. 2007;61(2):245-476.

19. Riley MA, Lizotte-Waniewski M. Population Genomics and the Bacterial Species

Concept. In: Gogarten MB, Gogarten JP, Olendzenski L, editors. Horizontal Gene

Transfer: Genomes in Flux. Methods in Molecular Biology. New York City: Humana

Press; 2009. p. 367-77.

20. Wertz JE, Goldstone C, Gordon DM, Riley MA. A molecular phylogeny of enteric

bacteria and implications for a bacterial species concept. J. Evol. Biol. 2003; 16:1236-

1248.

21. Maddison DR, Maddison WP. 2000. MacClade 4. Sinauer Associates, Sunderland,

Massachusetts.

22. Swofford DL. 2002. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other

Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts

23. Zachos FA. Species Concepts in Biology: Springer International Publishing;2016.

220 p. 32.

24. Wilkins JS. Species: University of California Press; 2011. 320 pp.

25. Tang CQ, Leasi f, Obertegger U, Kieneke A, Barraclough TG, Fontaneto D. The

widely used small subunit 18S rDNA molecule greatly underestimates true diversity

in biodiversity surveys of the meiofauna. Proceedings of the National Academy of

Sciences USA. 2012;109(40):16208–12.

26. Eckert EM, Fontaneto D, Coci M, Callieri C. Does a barcoding gap exist in

prokaryotes? Evidences from species delimitation in Cyanobacteria. Life. 2015;5:50-

64. doi: 10.3390/life5010050.

27. Stackebrandt E, Frederiksen W, Garrity GM, Grimont PAD, Kampfer P, Maiden

MCJ, et al. Report of the ad hoc committee for the re-evaluation of the species

definition in bacteriology. Int. J. Syst. Evol. Micr. 2001;52:1043-7.

28. Barraclough TG, Hughes M, Ashford-Hodges N, Fujisawa T. Inferring evolutionarily

significant units of bacterial diversity from broad environmental surveys of single-

locus data. Biol. Lett. 2009; 5:425-8.

29. Zhang J, Kapli P, Pavlidis P, Stamatakis A. A general species delimitation methode

with applications to phylogenetic plasements. Bioinformatics. 2013;29(22): 2869-76.

30. Koeppel A, Perry EB, Sikorski J, Krizanc D, Waarner A, Ward DM, et al. Identifying

the fundamental units of bacterial diversity: A paradigm shift to incorporate ecology

into bacterial systematics. Proceedings of the National Academy of Sciences USA.

2008;105(7):2504-9.

31. Hunt DE, David LA, Gevers D, Preheim SP, Alm EJ, Polz MF. Resource partitioning

and sympatric differentiation among closely related bacterioplankton. Science.

2008;320:1081-5.

32. Reno ML, Held NL, Fields CJ, Burke PV, Whitaker RJ. Biogeography of the

Sulfolobus islandicus pan-genome. Proc Nat Acad Sci USA. 2009;l106:8605-10.

33. Cahill JG, Green RE, Fulton TL, Stiller M, Jay F. Genomic evidence for island

population conversion resolves conflicting theories of polar bear evolution. PLoS

Genetics. 2013;9:e31003345.

34. Rossberg AG, Rogers T, McKane AJ. Are there species smaller than 1 mm? Proc.

Roy. Soc. B. 2013;280:20131248.

35. Ward DM, Bateson MM, Ferris MJ, Kuhl M, Wieland A, Koeppel A, et al.

Cyanobacterial ecotypes in the microbial mat community of Mushroom Spring

(Yellowstone National Park, Wyoming) as species-like units linking microbial

community composition, structure and function. Phil. Trans. R. Soc. B.

2006;361:1997-2008.

36. Cohan FM. Towards a conceptual and operational union of bacterial systematics,

ecology, and evolution. Phil. Trans. R. Soc. B. 2006; 361:1985-86.

37. Barraclough TG. The Evolutionary Biology of Species. Godfray CJ, Dunne JA,

Sheldon BC, editors. Oxford, United Kingdom: Oxford University Press; 2019.

38. Edwards SV, Beerli P. Perspective: Gene Divergence, Population Divergence, and the

Variance in Coalescence Time in Phylogeographic Studies. Evolution.

2000;54;6:1839-54.

39. Nei M, Li W-H. Mathematical model for studying genetic variation in terms of

restriction endonucleases. Proc. Nat. Acad. Sci. USA1979;76;10:5269-5273.