Evolutionary Genetic Species Detected in Prokaryotes by Applying the Κ/θ

Ratio to DNA Sequences

Running title: Using Κ/θ to delimit bacterial species

C. William Birky, Jr.*

Department of Ecology and Evolutionary Biology, The University of Arizona, Tucson,

Arizona 85721, United States of America

*Corresponding author

Email: birky@arizona.edu

Heather Maughan

Ronin Institute, Montclair, New Jersey 07043, United States of America
14
15   Orchid ID: 0000-0002-9421-8689
16
17

18   Keywords: bacterial species; evolutionary genetic species concept; Κ/θ species criterion

Abstract

Previous papers in this series described the "evolutionary genetic species concept" which is based on population genetic theory and should be applicable to any organism. Also described was a species criterion, the K/θ ratio, that delimits independently evolving evolutionary species based on single-gene sequences, and its application to sexual and asexual eukaryotes was illustrated. Here, we show how the evolutionary genetic species concept and the K/θ ratio can be applied to bacteria, using the sequences from some genes of the core genome that are rarely, if ever, transferred horizontally between different species. This completes the demonstration that K/θ is a general method for species delimitation, applicable to all organisms. Also, it adds to the evidence that bacteria have species in the most general sense, even though they have the ability to exchange genes across species boundaries. Finally, we show that a published critique of the use of $K/\theta \geq 4$ as a criterion for independently evolving species rests on two errors in the application of population/evolutionary genetic theory.

Introduction

A prominent feature of the biological world is discontinuous variation, with clusters of similar organisms differing significantly from organisms in other clusters. These clusters, called species and given different binomial names, were originally recognized by their different phenotypes. With the development of population genetic theory, species have been recognized as populations with distinct gene pools underlying their phenotypic differences. When gene sequences are used to reconstruct evolutionary history in the form of phylogenetic trees, different species appear as different clades of similar sequences separated by deep gaps in the tree.

This background is widely, but not universally, accepted. Doubters often focus on the gradual divergence of sister populations from each other, wondering at what point they should be called different species. Even among those who agree that species are real, distinct populations, there is disagreement on how to define and recognize them, given the gradual divergence of recently separated populations. Many authors simply point to "deep gaps" in a phylogenetic tree, usually of gene sequences, as evidence for different species, but this raises the question: "How deep is deep enough?"

This question can be answered by thinking of species as independently evolving populations. This is arguably the most general model of species (species concept), applicable in principle to all organisms and consistent with population and evolutionary genetic theory [1-8]. Different species are populations that evolve independently of each other, in the sense that they are independent arenas for mutation, natural selection, and

random genetic drift. In other words, a new mutant allele arising in one population cannot

spread to the other, or if it does move from one species to the other by hybridization, it is

very unlikely to be fixed in that population. Moreover, if a shared allele is fixed or lost in

one population by drift and/or selection, this does not affect the probability that it will be

fixed or lost in the other. This is referred to as the Evolutionary Genetic Species Concept

(EGSC) to distinguish it from the Evolutionary Species Concept (EvSC) of Simpson [7].

The EvSC as it is usually worded requires that we know, or assume, that the populations

in question will continue to evolve independently in the indefinite future. The

requirement for future independence should not be part of a species concept because it is

unknowable; for instance, a trait difference required for adaptation of species A to a

different niche from species B could be lost in the future. (However, it should be noted

that in 1953 Simpson did allow that speciation might occur in allopatry without

ecological differences in "some probably rare cases" [[7], p. 171]).

In practice, a pair of sister species are recognized as soon as they have diverged to

the point where samples of individuals can be shown to come from two independently

evolving populations. This can be done using a number of different criteria for

independent evolution. Prominent examples that have been applied to prokaryotes as well

as eukaryotes are reproductive isolation and ecological isolation, both of which obviously

result in independent evolution of the isolated populations. However, both of these

species criteria have conceptual and practical problems. Neither one is generally

applicable; reproductive isolation is relevant only for organisms with significant amounts

of sexual reproduction, and neither criterion includes the possibility that species may be

evolving independently simply because they are physically isolated from each other.

Moreover, neither is readily applicable to large-scale biodiversity surveys or species

discovery that relies on only sequences, as in the Barcode of Life [9] or Multi-Locus

Sequence Typing projects [10].

90   Although DNA sequences are widely used to delimit species in prokaryotes, there

is a great deal of controversy about how this should be done, and even about whether

bacteria really have species. Some of this controversy mirrors similar debates about

species in eukaryotes. However, much of it arises from the phenomenon of horizontal

gene transfer (HGT), in which segments of genes, whole genes, and even large segments

of chromosomes are transferred between distantly related bacteria via common

mechanisms of bacterial gene exchange: conjugation, transformation, and transduction.

This problem can be solved by distinguishing between core genes and auxiliary genes, as

core genes are less likely to be transferred or retained if transferred [11, 12]. Furthermore,

HGT doesn't appear to affect the bacterial species identification method based on average

nucleotide identity (ANI), where the nucleotide identity of each pair of orthologous genes

from two genomes is calculated and then averaged over all genes shared between the two

genomes [13]. The use of numerous genes is expected to diminish the effects of HGT on

sequence-based approaches to delimiting species. Although ANI-based methods have

proven to be useful, they are not based on theory but derive from empirical observations.

106   In this paper, as in previous publications [1-6], we argue that the EGSC can be

applied to all organisms, sexual and asexual, prokaryote and eukaryote. We also show

how the K/θ ratio, first used to discover and delimit EG species in eukaryotes based on

DNA sequence data from a single gene, can be applied to bacteria if a core protein-

coding gene or 16S rRNA gene is used. In particular, it could be applied to large-scale

diversity surveys of bacteria using such a gene. Such surveys are routine for eukaryotes,

and can now be done with bacteria regardless of whether they can be cultured in the lab,

thanks to the development of methods that allow amplification of genes from single

bacterial cells isolated by dilution or flow cytometry from environmental samples (e.g.

[14, 15]]. The successful identification of evolutionary genetic species in bacteria using

the K/θ ratio (this paper) as well as methods such as the Generalized Mixed Yule

Coalescent (GMYC; [1, 16]) adds to the growing literature that demonstrates that EG

species are a general phenomenon found in varying degrees in all organisms, sexual and

asexual, prokaryotes and eukaryotes.

The K/θ test is designed to distinguish between two kinds of clades: (i) those

separated by shallow gaps formed within a single species by random genetic drift or

transient physical isolation, and (ii) those separated by deep gaps produced by adaptation

to different ecological niches and/or by long-term physical isolation, i.e. different species.

Simulations [17] and population genetic theory [18] show that 95% of the gaps between

clades within a species are expected to have a depth of $< 4N_e$ generations (where N_e is the

effective population size), while gaps between species clades can be of any depth. It is

thus possible to recognize a pair of clades as different species, with probability ≥ 0.95, if

they are separated by gaps of $\geq 4N_e$ generations. At that point the sequence difference

between the two clades is $K \geq 8N_e\mu$ (where μ is the mutation rate per base pair) and the

ratio K/θ ≥ 8N$_e$μ/2N$_e$μ ≥ 4 (3-8). (Barraclough's [19] claim to have found errors in this

theory is erroneous, as we show below in Discussion.)

This result is based on expected probabilities and does not account for uncertainty

introduced by finite sample sizes. The relevant sampling theory has been developed by

Noah Rosenberg [20, 21], who showed that it holds for any pair of sister clades, i.e.

reciprocally monophyletic samples n1 and n2, provided the sample sizes are at least n1 =

3 and n2 = 2. The power of these very small samples may seem surprising, but it arises

from their first being shown to be reciprocally monophyletic.

To illustrate how the K/θ ratio is used to detect evolutionary genetic species in

prokaryotes, we use the set of sequences of six different housekeeping/core genes

previously used by Wertz et al. [22, 23] to delimit species in enteric bacteria. This was

one of the first multilocus data sets used to discover bacterial species. It provides an

opportunity to apply the K/θ method to single genes and compare the results of using

different genes to each other and to the named taxonomic species defined by classical

phenotypic traits. Note that these data are used simply to demonstrate the application of

the K/θ method and are not intended to be definitive statements about these specific

bacterial species.

Materials and Methods

151 John Wertz kindly provided partial sequences of six housekeeping core genes

152 (*gapA*, *groEL*, *gyrA*, *ompA*, *pgi*, and 16S rRNA) from seven named species of enteric

153 bacteria (*Citrobacter freundii* (abbreviated CF), *Enterobacter cloacae* (EB), *Escherichia*

154 *coli* (EC), *Hafnia alvei* (HA), *Klebsiella oxytoca* (KO), *Klebsiella pneumonia* (KP),

155 *Serratia plymuthica* (SP), and the outgroup *Vibrio cholera* (VC)).

157   The aligned sequences for each gene were visualized in MacClade [24] where all

158 the sequences of the gene were trimmed to the same length and all internal sites with gaps

159 in any one sequence were removed. Consequently, all pairwise sequence differences for a

160 given gene are based on the same regions of the gene. Final sequence lengths were: *gapA*,

161 540 bp; *groEL*, 348 bp; *gyrA*, 634 bp; *ompA*, 433 bp; *pgi*, 242 bp; *16S*, 289 bp. One

162 internal gap of 27 bp in the *ompA* gene of the outgroup *V. cholerae* was retained because

163 to remove it from all sequences would cause an unnecessary loss of sequence information

164 in the ingroup.

166   To discover species using the trimmed sequences, we followed the procedure

167 outlined by Birky [2] for each gene separately. Briefly,

168 1. First, we used PAUP* [25] to find a bootstrapped Neighbor-joining tree in which all

169 clades were supported by $\geq$70% of 1000 bootstrap replicas. A Neighbor-joining tree was

170 used to rearrange the sequences in the PAUP* file to reflect the order of sequences in the

171 tree; this greatly simplifies subsequent calculations in Excel, but note that the tree itself is

172 not used in subsequent calculations. Both corrected and uncorrected sequence

173 differences were then exported from PAUP* to an Excel spreadsheet where all

174 subsequent calculations were performed. Starting at the tips of the tree, each pair of sister

175 clades is tested separately in steps 2-7, to determine the probability that the clades are

176 samples from independently evolving species. This process is continued with deeper and

deeper clades until species are found. For clades that are members of a non-bifurcating

tree structure such as a polytomy (A, B, C) or ladder ((A,B)C), we compare A and B first,

then compare C to whichever one of those is closest to C.

2. For the two candidate clades in a pair, the mean pairwise difference *d* between

sequences is multiplied by the sample size correction n/(n-1) where *n* is the number of

sequences in the clade, to get the nucleotide diversity π. Note that nucleotide diversity

is defined as a function of the uncorrected pairwise distance. When $d = 0$, we used a non-

zero estimate of π by assuming that one sequence differs from the others by 1/L where L

is the sequence length; then $\pi = 2/Ln(n-1)$.

3. For each clade, we then estimate $\theta = 2N_e\mu$ by $\pi/(1 - 4\pi/3)$.

4. Calculate K = mean pairwise sequence difference between the two clades, corrected

for multiple substitutions.

5. Calculate K/θ for the pair of clades. When the values of θ for the two clades differ, we

use the larger value to get a conservative estimate of K/θ. If K/$\theta \geq 4$, we conclude that the

individuals in those clades came from different species, with probability ≥ 0.95.

6. The preceding steps are based on expected probabilities and do not account for

uncertainty introduced by finite sample sizes. This uncertainty is considered as follows,

using results obtained by Noah Rosenberg [20]. Using the K/θ ratio and the numbers $n1$

and $n2$ of individuals in the two clades, we want to find the probability that the

individuals were sampled from populations that have been evolving independently long

enough to become reciprocally monophyletic. This is best done using a table available on

request from Noah Rosenberg or the first author; alternatively, the values can be

estimated from Figure 6 in Rosenberg [20].

It is worth repeating that the phylogenetic tree used in step 1 is only used for two

purposes: (1) to facilitate the identification of sister clades, and (2) to inform the

organization of the Excel spreadsheet of pairwise sequence differences with the most

similar sequences together. The calculations could be done without the tree, which would

require identifying pairs of most similar sequences by eye in the spreadsheet. Thus, the

K/θ method is not affected by discrepancies between distance-based phylogenetic

inference methods.

Results

The Wertz et al. [23] data consist of 5 protein-coding genes plus part of the 16S

rRNA gene. We will first illustrate the application of the K/θ ratio by showing the

detailed analysis for *pgi*, the protein-coding gene that resolves the most species. The high

resolution suggests that this gene segregated before most of the other genes in most of the

speciation events, by chance. We will then summarize the results obtained with each of

the other genes. Following Wertz et al. [23], we consider the single isolate of *S.*

marcescens as belonging to the same taxonomic species as *S. plymuthica* because for all

the genes in this study it is identical or nearly so to the *S. plymuthica* isolates.

Finding species with *pgi*

The Neighbor-joining (NJ hereafter) tree in Figure 1 shows that the uncorrected

pgi sequences form a number of clades that are seen in $\geq$ 63% of 1000 bootstraps; most

have 100% support. The taxonomic species *K. oxytoca*, KO; *E. cloacae*, EB; *H. alvea*,

HA; and *S. plymuthica* SP, SM all form clades with 100% bootstrap support, but some of

these in turn contain two or more strongly supported clades. We applied the K/θ test to

each well-supported pair of sister clades, starting at the bottom of the *pgi* tree in Figure 1.

These clades are also well-supported in the NJ tree using corrected sequence differences.

The Excel table for *H. alvea* and *S. plymuthica* is shown in Table 1 as an example of the

data. The full tables for all genes are available on request from the first author.

1. To illustrate the procedure, we begin with clade HA2,4, which has weak support

(69% of bootstraps); its sister group is the singlet HA3. Using the formula for a

maximum estimate of π and θ when d = 0, we calculate that θ = 0.00836 and

from the table K = 0.0021 so K/θ = 0.25 and these are not different species. This

result was obvious from inspection and normally would not warrant a calculation.

Next we ask if clade HA1-4 is a different species from clade (SM1,SP1,SP2). The

pairwise distances for SM1, SP1, and SP2 are zero while those for HA1-4 are not

all zero; we prefer to use the latter because they do not require any correction. For

this clade d = 0.00625, π = 0.00833, and θ = 0.00843. K is 0.03032 so K/θ =

36.0 and these are samples from different clades with probability > 0.988.

Because K/θ >> 4, these specimens are also samples from different independently

species with P >> 0.95. Note that these results are obvious by inspection of the

branch lengths in an NJ tree like that shown for *pgi* in Figure 1 in which branch

lengths are proportional to the number of substitutions. The branches connecting

the (SM,SP) clade to the HA clade are clearly more than four times the length of

the branches connecting any two specimens within the HA clade.

2. The taxonomic species *C. freundii* (CF) forms two clades, CF1,5 and CF3,4, each

with 100% bootstrap support. Using Excel, we calculate the mean corrected

sequence difference between these clades, K = 0.07618. CF1,5 has the larger

value of θ = 0.0437, giving K/θ = 1.74. In Rosenberg's table, when K/θ = 1.7

and the sample sizes are 2 and 2, the probability that these two clades are samples

from reciprocally monophyletic populations is only 0.57. Moreover, because K/θ

is much less than 4, the probability that the samples came from different species is

low. This is the more conservative test in terms of splitting species because it

uses the larger of the two values of θ. On the other hand, the clade CF3,4 has the

smaller θ = 0.00167, giving K/θ = 4.5. From Rosenberg's table, $P = 0.97$ that

these are samples from different clades; furthermore, because K/$\theta > 4$, the

probability is very high that they are from different independently evolving

species. However, this is based on the less conservative estimate of θ. Because

only one of the two possible estimates of θ support the hypothesis of different

species, we conclude that the *pgi* gene shows weak support for two cryptic species

within *C. freundii*. As we will see below, some, but not all, other genes strongly

support splitting. This suggests that there is enough recombination in *C. freundii*

to cause different genes to complete segregation at different times, by chance.

3. The taxonomic species *E. coli* (EC) also consists of a clade (EC1-5,ECMG) with

100% bootstrap support and a singleton EC6. Only the clade can be used to

estimate θ, but since the 6 sequences in the clade are identical, π and thus θ

cannot be calculated directly. For these cases, one can conservatively estimate

these values by assuming that one of the 6 sequences actually differs from the

others at a single site. Then using the calculation given in Materials and Methods,

one can calculate π = 0.001377, θ = 0.00166, and K = 0.07000 so K/θ = 42.3 and

there is strong evidence that the clades EC1-5,MG and EC6 are samples from

different clades and from different cryptic species. This procedure is likely to

overestimate π and θ and hence underestimate K/θ, so in this sense it is

conservative with respect to splitting species.

4. Next we consider the clade consisting of *pgi* from KP isolates. Starting at the tips,

we compared (KP3,4) with KP5. Using the same procedure as above to get a non-

zero value for π, $K/\theta = 8.38$, providing evidence that these are samples from two

different clades and species. Comparing (KP3,4) to the other singleton, KP2, we

find $K/\theta = 11.6$ and these are also from different species. Thus KP is divided into

three species by these gene sequences.

5. Finally, the sequence data from this gene confirm that the isolates identified as *K.*

oxytoca (KO) and *E. cloacae* (EB) are evolutionary genetic species.

To summarize, the sequences of *pgi* confirm that KO, HA, EB, and SM,P are species;

split EC into 2 species; and split KP into 3 species.

Finding species with *groEL*

284   The *groEL* gene (Figure 1) confirms that KO, EB, SM, and HA are species. It

285   splits CF into two species, CF1,5 and CF3,4. EC is confirmed as a separate species from

286   its sister clade CF2-4 if one uses θ from CF2-4, but not if one uses θ from EC. *groEL*

287   sequences split KP into KP1,4 and KP2,6 if one uses θ from KP2,6; however, they do not

288   make this split if one uses the larger θ from KP1,4.

Finding species with *gyrA*

The gene *gyrA* (Figure 1) confirms that KO, EB, EC, HA, and SM are Evolutionary Genetic Species. The sequences of this gene split the taxonomic species CF into two EG species, CF1,5 and CF2-4, and also split KP into KP1,4 and KP2,6.

Finding species with *gapA*

The *gapA* gene (Figure 1) confirms that CF, KO, EC, SM,SP, EB, and HA are different species. KP consists of two clades, KP1,3,4 and KP2,6, and a singlet KP5 which are samples from three different species.

Finding species with *ompA*

The *ompA* gene (Figure 1) provides evidence that the CF, EC, EB, HA, and SP,SM sequences each represent a single species. Unlike some of the other genes, it does not split any of these species. In contrast, the KO and KP sequences are polyphyletic. Although the KO sequences are very similar with a diversity commensurate with that of other species in these data, the KP sequences are very divergent from each other. Moreover, KP5 is basal to KP1-4 as well as to KO; thus KP is polyphyletic, suggestive of incomplete lineage sorting.

Finding species with 16S

The 16S sequences of the taxonomic species *E. coli* and *H. alvia* contained regions that were not present in the other species and for which we could find no explanation, so these two taxonomic species were omitted from the analysis of the 16S sequences. Also, all pairwise differences among the ingroup sequences were ≤ 0.063, including those between species, so no corrections for multiple hits were applied. The

311 K/θ test verified that the CF1,5, KO, CF2-4, EB, and KP sequence sets, and the SP1

312 singlet, each represent evolutionary genetic species with probabilities >0.99. However,

313 the CFl and CF5 sequences did not; the clade was supported by only 56% of bootstrap

314 replicas, so no K/θ test was done. The NJ tree is shown in Figure 1.

Comparison with ANI

To compare these results with ANI values we used d, the mean pairwise difference for all

bacterial sequences for each gene. ANI is equal to 1-d, multiplied by 100 to make it a

percentage. The resulting values for the genes included in our analysis ranged from 95.9

to 100% ANI, all greater than the 95% cutoff for within-species differences [26]. Note

that these are values for single genes, not whole genomes, but if each gene is

representative of its genome, this means that species delimited with the K/θ $\geq$ 4 criterion

will also be identified as species by the accepted ANI criterion.

Discussion

Our focus here, and in several preceding papers on discovering species in asexual

and sexual eukaryotes [1-6], is on two problems: (i) identifying a model of species, i.e. a

species concept, that is general enough to cover all organisms, whether prokaryotes or

eukaryotes, and all degrees and kinds of sexuality; and (ii) based on the model, testing a

general method of assigning specimens to the same or different species (a species

criterion), using gene sequences. We showed that the Evolutionary Genetic species

concept and the K/theta ratio can be used to delimit bacterial species using DNA

sequences, and that this procedure often splits species defined by traditional systematics.

The results with the protein-coding genes and isolates EB, EC, HA, KO, and

SM/P are straightforward: each of the genes shows that these five taxonomic species,

discovered by traditional methods of bacterial taxonomy using morphology and various

phenotypic tests, are also EG species. The situation with CF is more complex. All five

genes show two clades, (CF1,5) and (CF2-4) with 89-100% bootstrap support except

gapA where the support for the (CF2-4) clade is only 64%. Moreover the K/θ ratio for

this pair of clades is greater than 4, signifying different species, for *gyrA*, *pgi*, and *groEL*

but not for *gapA* or *ompA*.

Deep gaps in trees separating species: how deep is deep enough?

In the literature of new species descriptions, and in papers proposing the splitting

of species, it is common to refer to deep gaps between clades in phylogenetic trees as

evidence that the clades should be assigned to different species. The decision as to

whether the gap between two clades A and B is deep enough to assign A and B to

different species, as opposed to indicating a relatively shallow gap within a single

species, is generally made using the systematist's intuition. Occasionally the systematist

compares the gap between A and B to the gap between named species C and D,

concluding that A and B are different species if the gap between them appears to be as

deep or deeper than the gap between C and D. However, this assumes that species C and

D are well defined. The K/θ method avoids making these assumptions by providing a

decision criterion that is soundly based on population genetic theory.

Barcode gaps

If one plots a frequency distribution of pairwise sequence differences between the

DNA sequences of a set of specimens, one often finds a bimodal distribution; the space

between the modes is called a barcode gap. This is the basis of the Barcode of Life

project. Eckert and colleagues [27] used nearly complete 16SrRNA sequences to look for

a barcode gap in Cyanobacteria, using the Automatic Barcode Gap Discovery (ABGD)

program; they found gaps in some but not all taxonomic groups. However, rather than

testing pairs of sister clades, Eckert et al. pooled between 34 and 2448 sequences from

each of 16 genera. This will obscure the barcode gap if different pairs of species have

gaps in different places, i.e. if some species have values of θ that overlap the values of

K. It is also possible that some of the groups examined by Eckert et al. have undergone a

recent adaptive radiation such that K values are very small, as the authors noted. In

contrast, Jain et al. [28] found a pronounced gap in plots of ANI from 91,761 whole

genome sequences from GenBank; the gap stretched from 83-95% ANI and contained

only 0.2%. of the pairwise comparisons.

Not all well-supported clades are species

372   These analyses illustrate the danger of assuming that all clades are species. For

373   example, in the *pgi* tree shown in Figure 1 the clade EB1,3,5 has 90% bootstrap support

374   and yet belongs to the same species as EB4. If one studied *E. cloacae* EB only, one could

375   easily draw a tree with the Y axis stretched out, so that the branch connecting the clade to

376   EB4 looks very long and might appear to show two species connected by a long branch.

Applicability of the K/θ method

It is clear that the Evolutionary Genetic Species Concept is applicable to bacteria.

It is also clear that within the framework of the EGSC, species can be discovered using

DNA sequences analyzed with the K/θ method. It is important to note that the K/θ test

was deliberately designed to work with a single gene or sequence. If the sequences of two

or more genes are used, it is important to remember that in a sexual organism (including

bacteria with high levels of recombination) different genes may become reciprocally

monophyletic at different times. We recommend that the K/θ test be applied to each gene

separately; if any gene places the specimens in two different clades and $K/\theta \geq 4$, this

would be strong evidence that the clades represent different species. We do not

recommend concatenating gene sequences without first testing whether the signal from a

gene that has not yet segregated can obscure the signal from a gene that has become

reciprocally monophyletic, i.e. could cause the specimens to be incorrectly assigned to a

single species.

The K/θ method complements the ANI method of species delineation. ANI-based

species cutoffs rely on ANI having a bimodal distribution with one cluster of interspecies

ANI values, a second cluster of intraspecies ANI values, and a gap in between (e.g., Fig 3

in ref [28]). This empirical observation is predicted by the population genetic theory that

underpins the K/θ method. Moreover, the K/θ method may be used to validate ANI

cutoffs or as a second method for cases where ANI values overlap the cutoff (e.g., 95%)

and thus do not support clear species separation.

398   Multiple comparisons

We used a probability of 0.05 for hypothesis testing, understanding that when this

is applied to many different cases, we expect to make a wrong decision in approximately

5% of the cases. This could be accounted for by using a correction such as the

Bonferroni. On the other hand, many of the results are significant with a much higher

probability. We choose not to use a correction and simply accept the risk of a small

number of wrong conclusions, less than one in twenty for the cases in this paper.

405 **When small sample sizes can provide strong conclusions**

Rosenberg [20] showed that small sample sizes can provide strong evidence that

the entire populations from which they were taken are reciprocally monophyletic. This is

because the probabilities are contingent upon the samples themselves being reciprocally

monophyletic. Of course this assumes that the samples are representative of the

populations, and do not, for example, come from a local sub-population that happens to

be very different from the rest of the population. Thus one must always bear in mind the

possibility that the specimens being studied are not representative. On the other hand, the

possibility of erroneously splitting a species into two species based on a non-

representative sample is likely to be small when K is very large. This is because, if a

large K gives a K/θ value less than 4, the θ value must also be much larger than the

generally observed values of < 0.05. There are parallels in traditional taxonomy, where

new species are often described on the basis of one or a few specimens, especially when

the new specimen is very different from the most similar described species. In this case

we usually assume that additional new specimens will not have intermediate phenotypes

that will cause us to combine the new specimen with a described species. If we had a

single specimen of an African lion and two specimens of an ocelot, traditional taxonomy

would probably recognize them as different species based on the absence of intermediate

phenotypes, even without additional evidence. If gene sequences from these specimens

had K/θ ≥ 4, this would indicate an absence of intermediate sequences and we would

have similar objective evidence that they were different species.

Use of trees and choice of tree-making methods

When implementing the K/θ method, the use of phylogenetic trees is limited to (i)

finding sister pairs of clades for comparison; (ii) guiding the arrangement of sequences in

the Excel spreadsheet to facilitate the calculation of K/θ; and (iii) spotting obvious cases

of K/θ based on branch lengths. We use Neighbor-joining for tree construction because it

is fast and easy to do with PAUP*, but presumably a different distance-based method

could be used.

Genes or phenotypes in bacterial species descriptions?

The bacterial rules of nomenclature do not specifically require phenotypic data.

However, the report of an *ad hoc* committee on species definition in bacteria [29] calls

for the use of qualitative phenotypic traits as well as 16S rDNA sequences to delimit

species. A quick survey of some recent papers in the *International Journal of Systematic*

and Evolutionary Microbiology shows that while rDNA sequences are now commonly

used in descriptions of new bacterial species, they are used essentially like a collection of

traits or to form phylogenetic trees. When these trees, or pairwise sequence differences,

or both are used to define trees, there is usually no attempt to apply a theoretically

justifiable method to distinguish sequence differences between species from differences

within species. Instead, this is based on the judgment or intuition of the investigators.

One gene or many?

There is a tendency in the recent literature to extoll the virtues of using many

genes to discover species. But using more than one gene has several drawbacks. First, it

costs more and will not be useful if one gene is sufficient to show that two samples come

from different species. Second, if one gene is a core gene, adding another gene which is

not a core gene and is involved in HGT can obscure the signal of speciation.

Ideally, to discover closely related species we would like to use the gene or genes

that segregated earliest in those particular populations; in this way we would catch

speciation in its early stages and discover closely related species. This is a real advantage

to using multiple gene sequences, which would increase the chance that at least one of

them would have segregated early. How important this is depends on the amount of

sexual reproduction in the bacteria in question. At one extreme, if there is little or no

recombination, or extreme inbreeding, all genes will segregate at the same time and any

core gene should be as good as any other. At the other extreme, if there is extensive

outcrossing and recombination, different genes may segregate at very different times after

two populations begin evolving independently. In fact, a given gene may segregate early

in one such population and late in the sister population. This is a likely explanation for

the observation that CF1 and CF5 form a separate species from CF2, 3, and 4 for all

protein-coding genes except *ompA*. Stackebrandt et al. [29] speak favorably of using

protein-coding genes, claiming that there is a consensus that at least five such genes are

necessary, but there is no such consensus in the sense of a number that is agreed on or at

least accepted by nearly everyone in the field. More importantly, Stackebrandt et al. [29]

do not consider the possibility that these genes may be at different stages of segregation,

which is the simplest explanation of the different trees supported by different genes in the

Wertz et al. [23] data.

There is also a potential problem with concatenated gene sequences: a significant

K/θ ratio in one gene may be obscured by a small, insignificant K/θ ratio in another gene.

In other words, the signal of species divergence in one gene that segregated early after

speciation may be swamped by the signal in another gene that has not yet completed

segregation with 95% confidence.

We suggest that the best method for large-scale diversity studies would be to use a

single protein-coding gene to discover species and then, for groups of special economic,

medical, ecological, or evolutionary importance or interest, test additional genes to see if

they separate more species pairs because they segregated earlier in those pairs. Then it

would often be useful to look for phenotypic or molecular markers that could be used to

identify those species without requiring sequencing.

480 Stability of EG species

A nice feature of species discovery using $K/\theta \geq 4$ is that additional specimens are

unlikely to fill in the gap between species and require them to be lumped. That is because

the large difference between K and θ reduces the probability that other specimens that are

intermediate between the two species and fill in the gap between them will ever be

discovered. Put another way, it is unlikely that we will find specimens and lineages that

join the two clades we identify as species. If this did happen, the single species produced

by joining the two clades would have exceptionally large nucleotide diversity.

Consequently, once two populations are assigned to different species using $K/\theta \geq 4$, those

species are unlikely to be lumped in the future. It is perhaps more likely that data from

additional genes or from new specimens will cause a species to be split in the future, and

this has been seen repeatedly in the literature. But this is true for species delimited by any

method, using any kind of data; in science, new information often forces us to change our

conclusions.

Other methods used to discover bacterial species with single

loci

496   Most of the methods proposed for using DNA sequence data to delimit species are

global methods, i.e. in a phylogenetic tree of gene sequences from many specimens, they

look for the transition point from species divergence to divergence between individuals

within species. One of these, GMYC [16] uses coalescent theory to detect the transition

point. Barraclough et al. [30] used GMYC to discover species from environmental 16S

rDNA sequences of bacteria but the method has not been applied to sequences from

already-described taxonomic species of prokaryotes. Poisson Tree Picking (PTP) [31]

attempts to identify significant changes in the numbers of substitutions along branches,

which are expected to be significantly higher between species than within species. To

our knowledge, PTP has not been applied to prokaryotes.

506   Ecosim [32] detects species produced by selective sweeps during adaptation to

different habitats. It assumes a Poisson model of substitutions, in contrast to K/θ and

GMYC that allow more complex and presumably more realistic models to be used. The

theory behind Ecosim includes random drift and selective sweeps but not less drastic

forms of selection. AdaptML [33] is another local method. It identifies clades associated

with ecological parameters that might be relevant to ecological species. It detects species

only if they are adapted to different niches that have already been identified. These

methods will not detect species that evolve independently because they are physically

isolated as in the case of *Sulfolobus islandicus* in hot springs [34] unless they are also

adapted to different niches. In other words, they may fail to detect some evolutionarily

independent species.

517   These global methods are all subject to error if different species have different

levels of within-species variation, i.e. different values of nucleotide diversity. This is in

contrast to $K/\theta \geq 4$ which is a local method focusing on sister clades and thus on putative

sister species. Such pairwise comparisons have the advantage that sister species are more

likely to have similar values of θ. In fact, the problem of delimiting species in general

really is a local problem. We have no problem assigning specimens of the grizzly bear

and the mule deer to different species, but it is more difficult to decide whether the

Alaskan brown bear and polar bear are really different species, e.g. [35].

Limitations of the K/θ method

526   In order to study the diversity of cyanobacteria from hot springs in Yellowstone

National Park, Ward et al. [36] did environmental sequencing of clone libraries from the

16SrRNA/ITS region from PCR-amplified DNA. Their Figure 4 shows a neighbor-

joining tree of ITS sequences of bacteria identified as *Synechococcus*. They found 10

putative ecotypes of *Synechococcus*, which are thus different species adapted to different

niches. However, it is evident by inspection of the tree that only the three clades called A,

A', and B satisfy the $K/\theta \geq 4$ criterion. This could happen if most of the ecotypes have

diverged too recently to pass the the $K/\theta \geq 4$ criterion for independent evolution. In this

case ecotype analysis can divide species more finely because it delimits species at an

earlier stage in the speciation process.

Using K/θ in Formal Descriptions of New Species

We have shown that the K/θ ratio can be used to delimit evolutionary species in

bacteria, illustrating the procedure with an exemplar data set. However, this does not

mean that the species so delimited conform to current practice in bacterial systematics, or

that the species so detected can be formally described in conformance with the current

rules of taxonomic nomenclature for bacteria. Of course the same is true of species

delimited with the GMYC method that also detects EG species, or with the method of

Cohan [37], which detects a subset of EG species that are separated by ecology.

Unfortunately, the rules of bacterial systematics codified in the International Code of

Nomenclature of Bacteria are not explicitly based in population genetic theory, even

though they pay lip service to the BSC which defines species as reproductively isolated

populations in outcrossing sexual organisms.

Response to the critique of Barraclough [19]

Barraclough [19] claimed to have found two errors in Birky's [1-5] calculation

that a K/θ value of 4 or greater is strong evidence of samples belonging to two different

evolutionary species. First, citing Edwards and Beerli [38], he claimed that "K from the

genealogy overestimates the true divergence time…due to sampling of the polymorphism

from the ancestral common ancestor species" and that K is equal to 2Tμ plus the θ of the

ancestor. However, Edwards and Beerli said that the addition of θ is necessary when K

denotes the time elapsed since two populations diverged, but it is not appropriate when

estimating the time T since allelic divergence, as we have done here and Birky did in

previous papers using K/ θ. Second, Barraclough refers to "a mitochondrial marker with

$\theta = N\mu$." In correspondence, he said that in this section he used N to refer to the effective

population size rather than the census size, so he is claiming that $\theta = N_e\mu$. If this were

true, then the upper 95% confidence limit of K/θ would be 8. But by definition $\theta =$

$2N_e\mu$ [39]. This is because the sequence difference between two gene copies in two

specimens A and B from the same species sharing a most recent common ancestor O is

equal to the sequence difference $N_e\mu$ between A and O plus the sequence difference

between B and O, which is also $N_e\mu$. Consequently, after $4N_e$ generations, K/$\theta =$

$8N_e\mu/2N_e\mu = 4$, and not 8.

Could we use K/θ as a measure of the progress of speciation?

If we adopt the evolutionary species concept, then speciation is usually a gradual

process. (Exceptions can include rare dispersal to isolated sites such as islands, or

speciation by hybridization or polyploidization.) To the extent that speciation is gradual,

the K/θ ratio is expected to gradually increase. Consequently, if we have two populations

that we suspect are undergoing speciation, K/θ might be used as a measure of the

progress of speciation. It would presumably be non-linear, as suggested by the

nonlinearity of progress toward reciprocal monophyly of the two populations [20].

Species are universal

575   In this paper and previous papers in this series [1-6] we have shown that Evolutionary

576   Genetic species are found in bacteria and in both asexual and sexual eukaryotes. The

577   arguments in this paper should also apply to Archaea. We strongly suspect that viruses

578   also form such species, and that extraterrestrial organisms, if they exist, will likewise

579   form populations that evolve independently from each other.

Funding information

The personal funds of CWB were the sole support of the research.

Author contributions

CWB was responsible for study design, data collection and analysis, decision to publish,

and preparation of the manuscript. HM assisted with manuscript writing and figure

preparation.

Conflicts of interest

The authors have declared that no competing interests exist.

Acknowledgements

We thank John Wertz for providing the aligned sequences.

References

1.   **Barraclough TG, Birky CW, Jr., Burt A**. Diversification in sexual and asexual organisms. *Evolution* 2003;57(9):2166-2172.

2.   **Birky CW, Jr.** Species detection and identification in sexual organisms using population genetic theory and DNA sequences. *PLoS One* 2013;8(1):e52544.

3.   **Birky CW, Jr., Adams J, Gemmel M, Perry J**. Using population genetic theory and DNA seuqences for species detection and identification in asexual organisms. . *PLoS ONE* 2010;5:e10609.

4.   **Birky CW, Jr., Ricci C, Melone G, Fontaneto D**. Integrating DNA and
traditional taxonomy to describe diversity in poorly studied microscopic animals: new
species of the genus Abrochtha Bryce, 1910 (Rotifera: Bdelloidea: Philodinavidae).
Zoological Journal of the Linnean Society 2011;161:723-734.

5.   **Birky CW, Jr., Wolf C, Maughan H, Herbertson LK, Henry E**. Speciation and
selection without sex. *Hydrobiologica* 2005;546:1-17.

6.   **Schon I, Pinto RL, Halse S, Smith AJ, Martens K et al.** Cryptic species in
putative ancient asexual darwinulids (Crustacea, Ostracoda). *PLoS One*
2012;7(7):e39844.

7.   **Simpson GG**. The species concept. *Evolution* 1951;5:285-298.

8.   **Wiens JJ**. What is speciation and how should we study it? *American Naturalist*
2004;163(6):914-923.

9.   International Barcode of Life. https://ibol.org/. [accessed.

10.   **Maiden MC**. Multilocus sequence typing of bacteria. *Annu Rev Microbiol*
2006;60:561-588.

11.   **Lan R, Reeves PR**. Intraspecies variation in bacterial genomes: the need for a
species genome concept. *Trends Microbiol* 2000;8(9):396-401.

12.   **Lan R, Reeves PR**. When does a clone deserve a name? A perspective on
bacterial species based on population genetics. *Trends Microbiol* 2001;9(9):419-424.

13.   **Konstantinidis KT, Tiedje JM**. Genomic insights that advance the species
definition for prokaryotes. *Proc Natl Acad Sci U S A* 2005;102(7):2567-2572.

14.   **Ottesen EA, Hong JW, Quake SR, Leadbetter JR**. Microfluidic digital PCR
enables multigene analysis of individual environmental bacteria. *Science*
2006;314(5804):1464-1467.

15.   **Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ et al.** Insights
into the phylogeny and coding potential of microbial dark matter. *Nature*
2013;499(7459):431-437.

16.   **Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP et al.**
Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Syst*
Biol 2006;55(4):595-609.

17.   **Avise JC**. *Molecular Markers, Natural History and Evolution*. New York:
Chapman & Hall, Inc.; 1994.

18.   **Hudson RR, Coyne JA**. Mathematical consequences of the genealogical species
concept. *Evolution* 2002;56(8):1557-1565.

19.   **Barraclough TG**. *The Evolutionary Biology of Species*. Oxford, United
Kingdom: Oxford University Press; 2019.

20.   **Rosenberg NA**. The shapes of neutral gene genealogies in two species:
probabilities of monophyly, paraphyly, and polyphyly in a coalescent model. *Evolution*
2003;57(7):1465-1477.

21.   **Rosenberg NA**. Statistical tests for taxonomic distinctiveness from observations
of monophyly. *Evolution* 2007;61(2):317-323.

22.   **Riley MA, Lizotte-Waniewski M**. Population genomics and the bacterial species
concept. *Methods Mol Biol* 2009;532:367-377.

23.   **Wertz JE, Goldstone C, Gordon DM, Riley MA**. A molecular phylogeny of
enteric bacteria and implications for a bacterial species concept. *J Evol Biol*
2003;16(6):1236-1248.

24.   **Maddison DR, Maddison WP**. MacClade 4. Sinauer Associates, Sunderland,
Massachusetts; 2000.

25.   **Swofford DL**. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other
Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts; 2002.

26.   **Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P et al.**
DNA-DNA hybridization values and their relationship to whole-genome sequence
similarities. *Int J Syst Evol Microbiol* 2007;57(Pt 1):81-91.

27.   **Eckert EM, Fontaneto D, Coci M, Callieri C**. Does a barcoding gap exist in
prokaryotes? Evidences from species delimitation in cyanobacteria. *Life (Basel)*
2014;5(1):50-64.

28.   **Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S**. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.
Nat Commun 2018;9(1):5114.

29.   **Stackebrandt E, Frederiksen W, Garrity GM, Grimont PAD, Kampfer P et
al.** Report of the ad hoc committee for the re-evaluation of the species definition in
bacteriology. *Int J Syst Evol Microbiol* 2002;52(Pt 3):1043-1047.

30.   **Barraclough TG, Hughes M, Ashford-Hodges N, Fujisawa T**. Inferring
evolutionarily significant units of bacterial diversity from broad environmental surveys of
single-locus data. *Biol Lett* 2009;5(3):425-428.

31.   **Zhang J, Kapli P, Pavlidis P, Stamatakis A**. A general species delimitation
method with applications to phylogenetic placements. *Bioinformatics* 2013;29(22):2869-
2876.

32.   **Koeppel A, Perry EB, Sikorski J, Krizanc D, Warner A et al.** Identifying the
fundamental units of bacterial diversity: a paradigm shift to incorporate ecology into
bacterial systematics. *Proc Natl Acad Sci U S A* 2008;105(7):2504-2509.

33.   **Hunt DE, David LA, Gevers D, Preheim SP, Alm EJ et al.** Resource
partitioning and sympatric differentiation among closely related bacterioplankton.
Science 2008;320(5879):1081-1085.

34.   **Reno ML, Held NL, Fields CJ, Burke PV, Whitaker RJ**. Biogeography of the
Sulfolobus islandicus pan-genome. *Proc Natl Acad Sci U S A* 2009;106(21):8605-8610.

35.   **Cahill JA, Green RE, Fulton TL, Stiller M, Jay F et al.** Genomic evidence for
island population conversion resolves conflicting theories of polar bear evolution. *PLoS
Genet* 2013;9(3):e1003345.

36.   **Ward DM, Bateson MM, Ferris MJ, Kuhl M, Wieland A et al.** Cyanobacterial
ecotypes in the microbial mat community of Mushroom Spring (Yellowstone National
Park, Wyoming) as species-like units linking microbial community composition,
structure and function. *Philos Trans R Soc Lond B Biol Sci* 2006;361(1475):1997-2008.

37.   **Cohan FM**. Towards a conceptual and operational union of bacterial systematics,
ecology, and evolution. *Philos Trans R Soc Lond B Biol Sci* 2006;361(1475):1985-1996.

38.   **Edwards SV, Beerli P**. Perspective: gene divergence, population divergence, and
the variance in coalescence time in phylogeographic studies. *Evolution* 2000;54(6):1839-
1854.

39.   **Nei M, Li WH**. Mathematical model for studying genetic variation in terms of
restriction endonucleases. *Proc Natl Acad Sci U S A* 1979;76(10):5269-5273.
693

Figures and Tables

698　**Table 1.** Matrix of pairwise sequence differences for *pgi* from *Serratia plymuthica* and

699　*Hafnia alvea*.

	SM1	SP1	SP2	HA1	HA2	HA3	HA4
SM1	-						
SP1	0.0000	-					
SP2	0.0000	0.0000	-				
HA1	**0.3022**	**0.3022**	**0.3022**	-			
HA2	**0.3010**	**0.3010**	**0.3010**	0.0083	-		
HA3	**0.3087**	**0.3087**	**0.3087**	0.0125	0.0042	-	
HA4	**0.3010**	**0.3010**	**0.3010**	0.0083	0.0000	0.0042	-

700　Uncorrected sequence differences *d* and corrected differences K (bold type).

Figure 1. NJ trees of *pgi, groEL, gyrA, gapA, ompA* or 16S rRNA sequences. CF, EC,

KP, KO, EB, SM, and HA are taxonomic species. Closed circles indicate clades that are

705   different EG species delimited by the K/θ method, with circle colors representing

706   different taxonomic species. All scale bars indicate 0.02 substitutions per site except for

707   those of *groEL* and *ompA*, which represent 0.04 and 0.2 substitutions per site,

708   respectively.