Conserved Molecular Function and Regulatory Subfunctionalization of the LORELEI Gene Family in Brassicaceae

Jennifer A. Noble[1], Ming-Che James Liu[2], Thomas A. DeFalco[3], Martin Stegmann[4,8], Kara McNamara[2], Brooke Sullivan[2], Khanhlinh K. Dinh[2], Nicholas Khuu[2], Sarah Hancock[1], Shin-Han Shiu[5], Cyril Zipfel[3,4], Mark A. Beilstein[1]*, Alice Y. Cheung[2,6,7], and Ravishankar Palanivelu[1]*

[1]School of Plant Sciences, University of Arizona, Tucson, AZ 85721, USA

[2]Department of Biochemistry and Molecular Biology, University of Massachusetts, Amherst, MA 01003, USA

[3]Institute of Plant and Microbial Biology, Zurich-Basel Plant Science Center, University of Zurich, Zurich, Switzerland

[4]The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, NR4 7UH, United Kingdom

[5]Department of Plant Biology, Michigan State University, East Lansing, MI 48824, USA

[6]Molecular and Cell Biology Program, University of Massachusetts, Amherst, MA 01003, USA

[7]Plant Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA

[8]Department of Phytopathology, Technical University of Munich, School of Life Sciences Weihenstephan, Freising, Germany

*Corresponding author: Email: mbeilstein@email.arizona.edu, rpalaniv@email.arizona.edu

Abstract

A signaling complex comprising members of the LORELEI (LRE)-LIKE GPI-anchored protein (LLG) and *Catharanthus roseus* RECEPTOR-LIKE KINASE 1-LIKE (*Cr*RLK1L) families perceive RAPID ALKALINIZATION FACTOR (RALF) peptides and regulate growth, development, reproduction, and immunity in *Arabidopsis thaliana*. Duplications in each component, which potentially could generate thousands of combinations of this signaling complex, are also evident in other angiosperms. Widespread duplication in angiosperms raises the question what evolutionary mechanisms underlie the expansion and retention of these gene families, as duplicated genes are typically rendered non-functional. As genetic and genomic resources make it a tractable model system, here we investigated this question using *LLG* gene family evolution and function in Brassicaceae. We first established that the *LLG* homologs in the Brassicaceae resulted from duplication events that pre-date the divergence of species in this family. Complementation of vegetative phenotypes in *llg1* by LRE, LLG2, and LLG3 showed that the molecular functions of LLG homologs in *A. thaliana* are conserved. We next tested the possibility that differences in gene expression (regulatory subfunctionalization), rather than functional divergence, played a role in retention of these duplicated genes. For this, we examined the function and expression of *LRE* and *LLG1* in *A. thaliana* and their single copy ortholog in *Cleome violacea* (*Clevi LRE/LLG1*), a representative species outside the Brassicaceae, but from the same order (Brassicales). We showed that expression of *LLG1* and *LRE* did not overlap in *A. thaliana* and that *Clevi-LRE/LLG1* expression in *C. violacea* encompassed all the expression domains of *A. thaliana LRE* + *LLG1*. Still, complementation experiments showed that *LLG1* rescued reproductive phenotypes in *lre* and that *Clevi LRE/LLG1* rescued both vegetative and reproductive phenotypes in *llg1* and *lre.* Additionally, we found that expression of *LLG2* and *LLG3* in *A. thaliana* have also diverged from the expression of their corresponding single copy ortholog (*Clevi LLG2/*LLG3) in *C. violacea*. Our findings demonstrated how regulatory subfunctionalization, rather than functional divergence, underlies the retention of the *LLG* gene family in Brassicaceae. Our findings on the regulatory divergence and

55 functional conservation provide an experimental framework to characterize the
56 combinatorial assembly and function of this critical plant cell signaling complex.

Introduction

How cells communicate with each other to ensure coordinated growth and development
remains a fundamental question in eukaryotes. The signaling complex comprising
members of FERONIA (FER), LORELEI (LRE), and RAPID ALKALINIZATION FACTOR
(RALF) families is rapidly emerging as one of the best characterized cell–cell signaling
models in Arabidopsis [1]. FER is a receptor kinase (RK) of the *Catharanthus roseus*
RECEPTOR-LIKE KINASE 1-LIKE (*Cr*RLK1L) family, LRE is a membrane-associated
glycosylphosphatidylinositol (GPI)-anchored protein, and RALF is a small cysteine-rich
peptide (CRP). In *Arabidopsis thaliana* reproduction, LRE localizes in the synergid cells
of the ovule and interacts with FER and the FER–LRE complex functions at the interface
of the synergid cell and pollen tube to mediate pollen tube reception and release of sperm
cells to effect double fertilization [2-10]. Consequently, ~80% of ovules remain unfertilized
and only ~20% of seeds are produced in *fer* or *lre* mutant pistils [2-10]. FER-based
signaling pathway in ovules is also important to prevent late-arriving pollen tubes from
entering an ovule that has already engaged with a pollen tube and thus prevent
polyspermy [11]. In *A. thaliana* pollen and pollen tubes, the integrity of these cells during
reproduction is dependent on a signaling complex involving the *Cr*RLK1Ls ANXUR1
(ANX1), ANX2, BUDDHA'S PAPER SEAL1 (BUPS1), and BUPS2, along with the peptide
hormones RALF4/19 and LRE homologs LORELEI-LIKE GPI-ANCHORED PROTEIN2
(LLG2) and LLG3 [12-15].
In addition to functioning in reproduction, variants of the *Cr*RLK1L–LLG–RALF
signaling complex mediate diverse processes in vegetative tissues, stress responses,
and plant immunity [6, 12, 16-18]. The FER–LLG1–RALF1 complex regulates a Rho-
GTPase complex to produce reactive oxygen species (ROS) and is critical for root hair
growth and hypocotyl elongation [6, 10, 18]. The FER–LLG1–RALF23 complex regulates
pattern recognition receptor complex formation to modulate the perception of the

pathogen-associated molecular patterns (PAMPs) flagellin and elongation factor thermo

unstable (EF-TU) via the immunogenic epitopes flg22 and elf18, respectively. In this

context, RALF23 binds to FER–LLG1 to suppress the scaffolding function of FER, thereby

inhibiting ROS production and immunity [16, 17].

In *A. thaliana*, the GPI-anchored protein (GPI-AP), receptor kinase, and small CRP

components of the trimeric *Cr*RLK1L–LLG–RALF signaling complex are encoded by 17,

4, and 37 genes, respectively [19]. These could potentially form 2,516 unique

combinations that function in myriad cell types and at different developmental time points.

Additionally, this core trimeric complex may interact with other signaling components to

form a multimeric signaling complex and mediate a variety of cellular processes [15].

Additional components that could interact with this trimeric complex include LEUCINE-

RICH REPEAT EXTENSIN (LRX) proteins, which can directly bind RALF peptides [20].

LRXs are required for maintaining pollen tube integrity [13] and form a complex with FER

that is important for vacuolar expansion [21]. However, a comprehensive evolutionary

analysis complemented with functional and expression studies has been done for any of

the members of the *Cr*RLK1L–LLG–RALF signaling complex. Consequently, how the

expansion of these gene families contributes to the functional diversification of this critical

signaling complex and what factors affect the maintenance of duplicate members of these

gene families remain unknown.

We addressed this question by producing a phylogeny for the *LLG* family members

and characterizing their patterns of functional and regulatory evolution. We identified

orthologs of the four-member *A. thaliana LLG* gene family and showed that they are

conserved throughout the Brassicaceae. We used complementation assays and showed

that the molecular functions of GPI-AP proteins encoded by the *LLG* gene family are

conserved. Examination of the function and expression of *LRE* and *LLG1* family members

in *A. thaliana* and their single copy orthologs in *Cleome violacea* showed that regulatory

divergence (i.e. differences in gene expression), rather than functional divergence, likely

contributed to the retention of *LRE* and *LLG1* in *A. thaliana* and possibly played a key role

in the diversification of this signaling complex.

Results

LRE, *LLG1*, *LLG2*, and *LLG3* are maintained in the Brassicaceae

To study the evolution of the *LLG* gene family, we obtained full-length coding sequences (CDS) for orthologs of *LRE*, *LLG1*, *LLG2*, and *LLG3* in eleven species from the Brassicaceae. We also identified orthologs from three species outside the Brassicaceae, but from the same order (Brassicales): *Carica papaya* (Caricaceae), *Tarenaya hassleriana* (Cleomaceae), and *Cleome violacea* (Cleomaceae). Only a single ortholog of the *LLG* family was found in *Amborella trichopoda*, a basal angiosperm, which was used as the outgroup in this analysis (S1 Table). We generated an alignment of all of the full-length CDS, then inferred phylogeny using maximum likelihood methods. The resulting tree was rooted with the single-copy gene from *A. trichopoda*.

Among species of Brassicaceae, *LRE* and its orthologs formed a monophyletic group sister to the clade containing all *LLG1* orthologs (Fig 1). We identified single-copy *LRE/LLG1* orthologs from *C. papaya*, *T. hassleriana*, and *C. violacea* that were sister to the *LRE* + *LLG1* clade, consistent with the possibility that *LRE* and *LLG1* in the Brassicaceae are products of the alpha whole-genome duplication (WGD) [22] (Fig 1).

Our approach identified *LLG2* and *LLG3* orthologs in all the species of Brassicaceae we analyzed with two exceptions: no *LLG2* ortholog was identified in *Brassica rapa* and no LLG3 ortholog was identified in *Aethionema arabicum* (S1 Table). Absence of an ortholog could be due to gene loss, incomplete genome sequencing coverage, or because our approach failed to identify the putative orthologs (see Methods). Our phylogenetic analysis also found that the single-copy *LLG2/LLG3* orthologs in *T. hassleriana*, and *C. violacea* are sister to *LLG3* orthologs in Brassicaceae. This suggested that the duplication occurred prior to the split of *LLG2* and *LLG3* and that these species have an *LLG3* ortholog but likely lack an *LLG2* ortholog (Fig 1). However, this conclusion is based on branches of the phylogenetic tree that are only supported by low bootstrap values. Additional data from species in Brassicaceae and Cleomaceae are required to determine whether the duplication that led to the *LLG2* and *LLG3* clades in Brassicaceae occurred early in the history of Brassicaceae, or whether the duplication predated the split between Brassicaceae and Cleomaceae. Still, our phylogenetic

146　analyses pointed to maintenance of four copies of *LLG* in the genomes of species in the

147　Brassicaceae following at least two duplication events (Fig 1).

149　The molecular functions of LLG1 are likely conserved in LRE, LLG2, and LLG3

Following a whole-genome duplication, the duplicated genes are initially redundant;

accumulating mutations quickly render most gene duplicates non-functional, with a half-

life of a few million years [23]. For instance, only about 15% of duplicated genes were

retained in *A. thaliana* [24, 25]. However, our phylogenetic analysis showed that several

duplicates have been retained in the *LLG* family; these genes may have developed non-

redundant functions. Therefore, we examined whether divergence in molecular function

caused by differences in the transcribed genic regions could explain the maintenance of

the *LLG* gene family in the Brassicaceae.

To this end, we expressed *A. thaliana LRE*, *LLG2*, or *LLG3* from the *LLG1*

promoter in the *llg1-2* mutant and tested if they could complement the vegetative

development defects of *llg1-2* mutants [6, 16, 17, 26]. As a positive control in these

experiments, we used *llg1-2* mutant carrying a transgenic construct with *LLG1* expressed

from its own promoter. Expression of *LLG3* or *LRE* restored root hair phenotypes

comparable to the expression of *LLG1* (Fig 2A, S1B Fig) and complemented hypocotyl

length and epidermal pavement cell defects in *llg1-2* seedlings (Figs 2B,3A-3E, S1C Fig).

Additionally, seedlings expressing either of these two transgenes showed restored

RALF1 sensitivity (Fig 3G, S2 Fig). Notably, LRE complemented insensitivity to RALF1 in

llg1-2, restoring RALF1-induced root growth inhibition (Fig 3G). Expression of *LLG2* also

complemented vegetative defects in *llg1-2* mutants, as rosette size was restored to wild

type levels (Fig 4A).

In addition to vegetative phenotypes, *fer* and *llg1* mutants are defective in immune

responses, as they show reduced responsiveness to several PAMPs, including the

bacterial elicitors elf18 and flg22 [16, 17, 26]. We found that responses to flg22 and elf18

in *llg1-2* plants expressing *LLG2* from the *LLG1* promoter were restored to levels

comparable to those expressing *LLG1* from *LLG1* promoter, indicating that LLG2 can

substitute for LLG1 and perform its molecular functions (Fig 4B, S3 Fig).
Complementation of *llg1-2* with *LLG2* similarly restored responsiveness to exogenous
RALF23 peptide to levels detected in *llg1-2* plants expressing *LLG1* (Fig 4C). Taken
together, these results demonstrated that the molecular functions of LLG1 are mostly
indistinguishable from those of LRE, LLG2, or LLG3. Hence, the retention of *LLG* paralogs
in Brassicaceae was unlikely to have been due to divergence in molecular function
caused by differences in their transcribed genic regions.

LRE and LLG1 have distinct expression patterns in *A. thaliana*

The *LLG* gene family members are differentially expressed in *A. thaliana*, with *LLG1*, *LRE*,
and *LLG2/LLG3* primarily expressed in vegetative, female, and male reproductive tissues,
respectively [9, 17]. Consistent with this, we found that putative transcription factor
binding sites in the promoters of *LLG* gene family members are distinct and showed
considerable variation (S4 Fig). Based on these findings, we considered the possibility
that divergence in expression of the *LLG* gene family (regulatory divergence), rather than
divergence in molecular function, underlies the maintenance of the *LLG* gene family in
the Brassicaceae. To test this possibility, we performed detailed, cell-specific expression
of *LRE* and *LLG1*, as they are closely related paralogs (Fig 1) that are also differentially
expressed [9]. Additionally, we chose to use these two genes to investigate this possibility
because *lre* and *llg1* single mutants have well-defined, non-overlapping phenotypes [6,
9, 17], allowing reciprocal complementation experiments to be performed (see below) to
test the functional divergence aspect of the hypothesis. By contrast, *llg2* and *llg3* single
mutants do not show phenotypes; phenotypes were detected only in a *llg2 llg3* double
mutant, thus making these genes not useful for reciprocal functional tests [6, 9, 17].
Previously, RT-PCR experiments indicated that expression of *LRE* is more tightly
restricted than *LLG1*, as *LRE* is primarily expressed in reproductive tissues and *LLG1* is
expressed throughout plant development [9]. Still, *LRE* and *LLG1* expression overlapped
in at least three tissues: 8-day-old seedlings, unfertilized pistils, and pollinated pistils [9,
27]. Thus, it is unknown if the domains of *LRE* and *LLG1* expression are indeed distinct
within these multicellular tissues. We therefore examined cell-specific expression of *LRE*

and *LLG1* using transcriptional fusions of the *LRE* or *LLG1* promoters to β-glucuronidase

(GUS). We characterized GUS expression in three *pLRE::GUS* lines (this study), and in

a previously reported *pLLG1::GUS* line [6]. GUS expression was examined in the

following tissues where either *LRE* or *LLG1* is expressed [9, 27]: 8-day-old seedlings, 21-

day-old seedlings, unpollinated pistils 24 hours after emasculation (HAE), and pollinated

pistils at 13.5 hours after pollination (HAP) and 18 HAP (Fig 5).

In 8-day-old seedlings, *pLLG1::GUS* was expressed in true leaves, the hypocotyl,

and to a lesser extent in roots (Fig 5A and 5B). However, we were unable to detect GUS

staining at this timepoint in any of the three *pLRE::GUS* lines (Fig 5A and 5C). The

pLRE::GUS expression results were not consistent with the previous RT-PCR results

obtained using 8-day-old seedlings [9, 27]. Perhaps the *LRE* promoter sequence used in

this construct did not include all the *cis*-regulatory elements required for expression in 8-

day-old seedlings. Alternatively, *pLRE::GUS* may be expressed in these tissues, but at

levels below the detection limit of our assay. In 21-day-old seedlings, *pLRE::GUS*

expression matched the RT-PCR results [9]. *pLLG1::GUS* was expressed in newly

emerged true leaves (Fig 5E) and *pLRE::GUS* was not detected in any cell or tissues of

21-day-old seedlings (Fig 5F).

In unpollinated pistils, *pLRE::GUS* was strongly expressed in the synergid cells

(Fig 5G and 5H). After pollination, *pLRE::GUS* expression was weaker in the zygote at

13.5 HAP (Fig 5J) and in proliferating endosperm at 18 HAP (Fig 5I). The cell-specific

expression of *pLRE::GUS* at these stages was consistent with the results obtained using

pLRE::GFP [27]. *pLLG1::GUS* was expressed only in the septum and nectaries in

unpollinated and pollinated pistils (Fig 5G, 5I, 5K, and 5M). These results demonstrated

that *LRE* and *LLG1* are not expressed in the same cells in seedlings and pistils. These

findings are consistent with the regulatory divergence hypothesis and suggested that

expression differences between *A. thaliana LRE* and *LLG1* likely contributed to their

retention post duplication.

LLG1 complements the reproductive functions of LRE in *A. thaliana*

Reciprocal complementation experiments are one way to investigate if functional divergence likely contributed to their retention post duplication. We showed that LRE can substitute for LLG1 molecular functions in vegetative development (Fig 2). To investigate if LLG1 can perform the reproductive functions of LRE, we expressed LLG1 from the *LRE* promoter, and transformed it into *lre-7* plants. To allow us to visualize the protein, we also fused LLG1 to citrine Yellow Fluorescent Protein (c*YFP*). In three homozygous single-locus insertion *pLRE::LLG1-cYFP* lines, LLG1-cYFP localized to the filiform apparatus and puncta in the synergid cells (S5 Fig), similar to what we previously reported for LRE using *pLRE::LRE-cYFP* (S5 Fig; [7]. All three homozygous single-locus insertion *pLRE::LLG1-cYFP* lines restored seed set defects in *lre-7* to levels detected in wild type or *pLRE::LRE-cYFP* (Fig 6B). Additionally, when the transgenic plant was used as the female parent in a cross to wild type, there was a significantly increased transmission of *pLRE::LLG1-cYFP* transgene in the progeny of the cross, showing that LLG1-cYFP complemented the defects in the *lre-7* female gametophyte (Table 1); no such increase in transmission was observed in the progeny from a reciprocal cross, when pollen from the transgenic plant was crossed to wild type (Table 1). Based on these results, we concluded that LLG1 expressed under the *LRE* promoter complements the reproductive defects in *lre-7* mutants, and that the molecular functions of LRE and LLG1 are mostly conserved.

The single copy LRE/LLG1 ortholog in *C. violacea* can substitute for both LLG1 and LRE in *A. thaliana*

Based on reciprocal complementation experiments in *A. thaliana*, we concluded that LRE and LLG1 can perform each other's molecular functions. Additionally, phylogenetic analysis (Fig 1) indicated that these paralogs were a product of the alpha WGD that occurred at the base of Brassicaceae [22]. Taken together, these results raise the possibility that the molecular functions performed by the single copy LRE/LLG1 ortholog in *C. violacea*, a member of the Cleomaceae, sister family to Brassicaceae (Fig 1), are conserved in LRE and LLG1. To test this prediction, we fused *Clevi-LRE/LLG1* to cYFP, and expressed it from the *A. thaliana LLG1* or *LRE* promoters (S1A and S5 Figs), then

transformed these constructs into *llg1-2* and *lre-7* plants, respectively. We found that Clevi-LRE/LLG1 expressed from the *LLG1* promoter complemented root hair defects (Fig 2a and S1B Fig), epidermal cell defects (Fig 3F), and hypocotyl lengths in dark-grown *llg1-2* seedlings (Fig 2B, S1C Fig) to levels seen in *llg-1-2* lines carrying LLG1. Clevi-LRE/LLG1 also complemented *llg1-2* insensitivity to RALF1-induced root growth inhibition comparable to levels seen in *llg1-2* lines carrying LLG1 (Fig 3G and S2 Fig), which also suggests that the *Cr*RLK1L–LLG–RALF signaling complex is conserved outside of the Brassicaceae.

In all three *pLRE::Clevi-LRE/LLG1-cYFP* single-locus insertion transgenic lines, we detected partial complementation of reproductive defects in *lre-7*, as the seed set in these lines were significantly higher than that in *lre-7*, yet significantly lower when compared to wild-type (6B Fig)*.* Additionally, when the transgenic plants were used as the female parent in a cross to wild type, there was a significantly increased transmission of *pLRE::Clevi-LRE/LLG1-cYFP* transgene in the progeny of the cross, showing that Clevi LRE/LLG1-cYFP complemented the defects in the *lre-7* female gametophyte (Table 2); no such increase in transmission was observed in the progeny from a reciprocal cross, when pollen from the transgenic plant was crossed to wild type (Table 2). Based on these results, we concluded that the single-copy *LRE/LLG1* in *C. violacea* partially complemented *lre-7* reproductive phenotypes.

Partial complementation was perhaps due to relatively lower protein levels of Clevi-LRE/LLG1 compared to LRE as revealed by the cYFP fusion proteins levels in the filiform apparatus of the synergid cells in these lines (S5G Fig). Increased sequence divergence in the single-copy *LRE/LLG1* ortholog in *C. violacea* compared to *LRE* rather than *LLG1* is another possibility for partial complementation, as the branch length leading to the *LRE* clade was longer than the branch length leading to *LLG1*, indicating more substitutions post duplication in LRE compared to LLG1 (Fig 1). Since LRE and LLG1 were able to complement each other's functions (Figs 2–4), a third possibility is that the changes that that affect the ability to complement occurred along the branch leading to *C. violacea*. Nevertheless, our results indicate that the *C. violacea* single-copy LRE/LLG1 ortholog is capable of complementing the functions of both *A. thaliana* LRE and LLG1.

The single copy *LRE/LLG1* ortholog in *C. violacea* shows broad expression in both vegetative and reproductive tissues

Conserved molecular functions in *LRE* orthologs coupled with non-overlapping expression of *pLRE::GUS* and *pLLG1::GUS* in *A. thaliana* indicate an important role for regulatory divergence in the maintenance of the *LLG* gene family paralogs in Brassicaceae. Such divergence is a hallmark of regulatory sub-functionalization [28, 29]. In this case, the ancestral single copy gene would have been expressed in vegetative and reproductive tissues and post duplication, this expression pattern would have been partitioned between the descendant paralogs.

To test this prediction, we characterized the expression of the single copy *LRE/LLG1* ortholog in *C. violacea* (*Clevi-LRE/LLG1*; Fig 7). We performed RT-PCR on cDNA isolated from the following three stages of *C. violacea* development in which *LLG1* is expressed, but *LRE* is not expressed, in *A. thaliana*: 1) rosette leaves from 30-day-old plants (equivalent in size to 21-day-old *A. thaliana* rosette leaves; Fig 5), 2) anthers and pollen [9, 17], and 3) mature emasculated pistils without ovules that still included the septum and nectaries (Fig 5). In these RT-PCR experiments, we also included ovules isolated from mature emasculated pistils in which *LRE*, but not *LLG1*, is expressed in *A. thaliana* (Fig 5).

Clevi-LRE/LLG1 was expressed in all developmental stages that we tested and *Clevi-LRE/LLG1* expression encompassed all the expression domains of *A. thaliana LRE + LLG1* (Fig 7). The RT-PCR products amplified were sequenced to confirm that they were indeed full-length transcripts of *Clevi-LRE/LLG1*. Expression analysis in *A. thaliana* and *C. violacea* showed that the ancestral expression pattern of *Clevi-LRE/LLG1* is divided between the paralogs, leading to non-overlapping expression in *A. thaliana*. This provided additional support to the hypothesis that regulatory sub-functionalization underlies *LLG* paralog retention in Brassicaceae.

Discussion

Regulatory divergence likely led to the retention of *LRE* and *LLG1*

Multiple mechanisms have been proposed to explain the retention of duplicated genes, as progressive degeneration of one member of a paralogous set of genes is the default outcome (non-functionalization; [30]). Duplicated genes may acquire new functions (neo-functionalization) and hence may be retained [31]. A third mechanism of retention is sub-functionalization via sequence variations in the protein-coding or regulatory regions of the genes, by which the ancestral functions could be shared between the duplicated copies, resulting in non-overlapping expression of each paralog [29]. The advent of high-throughput sequencing and large-scale transcriptomic studies have allowed evaluation of expression divergence in duplicated genes in *A. thaliana* [28, 32-34], *Glycine max* [35], and *Zea mays* [36]. Although these studies generated critical evidence in support of regulatory sub-functionalization and/or regulatory neo-functionalization, they also relied on comparisons of reconstructed expression of the ancestral expression states rather than direct expression analysis of a species with a single copy ortholog. Similarly, functional complementation studies, particularly those across species, are critical for establishing the mechanisms contributing to duplicate retention; however, very few genes have been subjected to such studies.

Here, we demonstrated that a single copy ortholog from *C. violacea*, which diverged prior to the alpha WGD that gave rise to *LRE* and *LLG1* in the Brassicaceae, is able to fully substitute for LLG1 in vegetative tissues and can partially perform the functions of LRE in synergid cells. Expression analyses of *LRE*, *LLG1*, and *Clevi-LRE/LLG1* support our hypothesis that after gene duplication, the *LRE* and *LLG1* clades experienced regulatory sub-functionalization yielding non-overlapping expression patterns [33]. The combined expression domains of *LRE* and *LLG1* together with the *Clevi-LRE/LLG1* expression domains suggest that a single-copy *LRE/LLG1* played multiple roles in the common ancestor of the Brassicaceae and Cleomaceae. Based on

these results, we propose that the complementary expression of *LRE* and *LLG1* led to
the retention of these paralogs in Brassicaceae, as loss of either copy would cause loss
of expression in certain tissues and lower fitness [29].

Expression divergence is also a key player in the evolution of *LLG2* and *LLG3* in
Brassicaceae, as we found that the single-copy *LLG2/LLG3* ortholog in *C. violacea* is
expressed in all the tissues examined (Fig 7) even though *A. thaliana LLG2/LLG3* are
primarily expressed in pollen and pollen tubes [9, 17]. Consistent with these results, highly
diverged transcription factor binding sites were found in the promoter regions of *LLG2*
and *LLG3* in *A. thaliana* compared to the single copy ortholog *LLG2/LLG3* in *C. violacea*
(S4 Fig).

Besides expression divergence, in this study we showed that in *A. thaliana,*
members of the *LLG* family share molecular functions, as every member of this family
can substitute for LLG1 in vegetative tissues and immune responses and LLG1 can
function in ovules in place of LRE. We also showed that *Clevi* LRE/LLG1 can substitute
for LLG1 and LRE. Taking the results of expression and molecular function analyses
together, we propose that conserved molecular functions and expression divergence are
the keys to the expansion and retention of the *LLG* gene family in Brassicaceae.

The *LLG* Gene family may be co-evolving with the *Cr*RLK1L and RALF families

Division of ancestral expression is one mechanism that allows duplicated genes to be
retained [29]. Given our results, it is possible that the genes encoding the other members
of the trimeric complex (*Cr*RLK1Ls and RALFs) show similar divisions of expression
domains. Additional phylogenetic analyses combined with expression analyses will be
required to understand the evolution of expression domains in *CrRLK1L* and *RALF*
families in the Brassicaceae. *RALFs* and *CrRLK1Ls* are members of large gene families,
which poses a challenge in efforts to study all the possible combinations of the co-receptor
complex [5, 37]. However, phylogenetic analysis coupled with functional and expression
studies, similar to that performed in this study, may offer a viable approach to address
this challenge.

FER functions with both LRE and LLG1, and correspondingly, *FER* is expressed
in both *LRE* and *LLG1* expression domains [6, 9]. Given that specific *Cr*RLK1Ls, RALFs,
and LLGs play distinct biological roles, it may be that *Cr*RLK1Ls and RALFs have co-
evolved with members of the LLG family to perform their functions in different tissues. *In*
vitro binding of RALF23 with LLG1, LLG2, and LLG3 but not LRE provide support for this
possibility [17]. Methods such as evolutionary rate covariation have been used to link co-
evolution with functional associations [38]. Such methods, in combination with the
phylogenetics, expression analyses, and molecular genetic assays used in this study will
prove invaluable in further characterizing members of this critical signaling complex.

Materials and Methods

Identifying CDS for the orthologs of the *LLG* gene family

CDS of putative orthologs of the *LLG* gene family and single-copy *LRE/LLG1* and
LLG2/LLG3 orthologs were obtained through the Comparative Genomics (CoGe)
Platform using CoGeBLAST (tBLASTx) with *A. thaliana LRE* or *LLG1* nucleotide CDS as
the search query using standard parameters [39-41]. For each result, if annotations for
CDS were available, then they were downloaded directly using "FeatView" on CoGe. In
cases of incomplete CDS or when no CDS annotations were available, a 5–7-kb region
surrounding the sequence of interest was downloaded and then aligned to the original
BLAST query from *A. thaliana LRE* and *LLG1* to identify the entire putative CDS using an
exon-by-exon approach with Geneious Alignment (R11.1.2) (https://www.geneious.com).

To identify putative orthologs from our tBLASTx results, we performed reciprocal
BLASTs to *A. thaliana*, followed by alignments and phylogenetic trees using *A. thaliana*
LRE, *LLG1*, *LLG2*, and *LLG3* sequences (see below). Putative orthologs were determined
based on the most closely related *A. thaliana* paralog. To find *LRE/LLG1* and *LLG2/LLG3*
single-copy orthologous genes in *Tarenaya hassleriana*, *Cleome violacea*, and *Carica*
papaya, we used tBLASTx of *A. thaliana LRE* and *LLG1* nucleotide CDS as the search
queries. We built an alignment of these sequences, *LRE*, and its paralogs in *A. thaliana*,
and used the alignment to build a phylogenetic tree. Only two loci were identified for each
genome: one corresponding to *LRE/LLG1*, the other corresponded to *LLG2/LLG3*;

therefore, we named the single-copy genes after their corresponding phylogenetic group.
We found in polyploid species that there were typically additional copies of paralogs. We
named these numerically, without any particular preference. We identified a single-copy
gene in *Amborella trichopoda*, as tBLASTx with any *LLG* gene family member in *A.*
thaliana only identified the single-copy ortholog in *A. trichopoda*. A list of the genomes
and corresponding gene IDs for putative orthologs are presented in Table S1.

CDS alignments and phylogenies

Alignments were built using the standard parameters for MUSCLE 3.8.425 in Geneious
R11.1.2, followed by manual curation using Geneious [42, 43]. From CDS alignments, we
built phylogenetic trees using the RAxML 8.2.11 plugin in Geneious with the following
parameters: GTR GAMMA nucleotide model; rapid bootstrapping and search for best-
scoring ML tree algorithm, with 100 bootstrap replicates; and starting with a completely
random tree [44]. Phylogenetic trees were rooted with *Amborella trichopoda*, which
served as the outgroup. The alignment and trees resulting from these analyses were
deposited to TreeBASE (Accession URL during review: https://treebase.org/treebase-
web/search/study/summary.html?id=25583&x-access-
code=2ede7b91dbd6e347d5c22132b139cc70&agreement=ok and Final URL after
publication: http://purl.org/phylo/treebase/phylows/study/TB2:S25583).

Plant materials and growth conditions

A. thaliana and *C. violacea* seeds were liquid sterilized in the following manner: 10–300
seeds were placed into a 1.5-mL microcentrifuge tube with 1 mL of 70% EtOH and
vortexed for 3 seconds at maximum speed at least 3 times over the course of 3–5 minutes
to ensure all seeds were sufficiently exposed to the sterilizing solution. The 70% EtOH
solution was discarded and replaced with 1 mL of sterilization solution (50% bleach, 0.2%
Tween-20 [Sigma-Aldrich, Catalog # P9416-100ML]), then vortexed for 3 seconds at
maximum speed, at least 3 times over 3–5-minutes. The sterilization solution was

discarded, and seeds were washed three times with 1 mL of ice-cold autoclaved dH$_2$O
each time. Using a 1-mL pipette, seeds were plated on 1/2X Murashgi and Skoog (MS)
plates (Carolina Biological Supply Co., Catalog # 195703), with 1% sucrose for seedling
growth assays and with corresponding antibiotics for transmission assays.

Seeds on plates were stratified for 3 days in the dark and at 4 °C, then plates were
moved to a Percival growth chamber maintained at 21 °C with continuous light (75–100
µmol·m^{-2}·s^{-1}). Ten-to-fourteen-day-old seedlings were transplanted to soil and grown in
the following conditions: 16 h light (100–120 µmol·m^{-2}·s^{-1}) at 21 °C and 8 h dark at 18 °C
as described [45].

Columbia (Col-0) is the ecotype of all *A. thaliana* seeds used in this study.
pLRE::GUS was transformed into the Col-0 background. *pLRE::LRE-cYFP*, *pLRE::LLG1-*
cYFP, and *pLRE::Clevi-LRE/LLG1-cYFP* are all in the homozygous *lre-7* mutant
background and were selected on plates containing hygromycin B (20 µg/mL;
PhytoTechnology Laboratories, Catalog # H397) and glufosinate ammonium (10 µg/mL;
Oakwood Chemical, Catalog # 044851). The *pLLG1::SP-mRFP-LLG1* and *pLLG1::SP-*
mRFP-LLG2 lines were transformed into *llg1-2* plants by floral dip, and transformants
were selected on 1x MS-agar with 1% sucrose, supplemented with 25 µg/mL kanamycin
or 10 µg/mL glufosinate ammonium, respectively. Seedlings for seedling growth inhibition
assays were grown in 12 h light (120 120 µmol·m^{-2}·s^{-1}) at 19–21 °C. Plants for ROS burst
assays were grown in 10 h light (150 µmol·m^{-2}·s^{-1}) at 20 °C.

Cloning transgenic constructs

The *pLLG1::LRE-HA-GPI* and *pLLG1::LLG3* transgenic constructs were prepared using
the primers and DNA templates indicated in Supplemental table S4, and contained the
LLG1 promoter region (~2 kb upstream of ATG) as presented in the previously published
constructs [6]. For SP-mRFP-LLG1 expression *in planta*, the native *LLG1* promoter was
amplified from seedling genomic DNA using primers
"ccaagcttgcatgccGTCGTTGTCCCAGATTCGTCG" and
"gatctagagtcgaccGGTTCTTTGTTGGTTACAGGAGAAGTCAC" for subsequent cloning
into the Gateway destination vector pGWB1 [46] using Infusion cloning (Takara Bio) and

the SdaI/SbfI restriction enzyme (Thermo Scientific). The *SP-mRFP-LLG1* construct was

synthesized with attached attB1/attB2 sites (Twist Bioscience) and subsequently

recombined into pDONR-Zeo using BP II Clonase (Invitrogen). The resulting pDONR-Zeo

SP-mRFP-LLG1 construct was recombined with *pGWB1-pLLG1*. The *pLLG1::SP-mRFP-*

LLG2 construct was synthesized (Thermo Scientific) with attB1/attB2 sites added and

cloned into *pDONR-Zeo* via BP II Clonase (Invitrogen) and was subsequently recombined

with *pGWB601* [47] for *in planta* expression.

The *pLRE::LLG1-cYFP, pLRE::GUS, pLRE::Clevi-LRE/LLG1-cYFP,* and

pLLG1::Clevi-LRE/LLG1-cYFP transgenic lines were created by replacing the genomic

sequence of *LRE* in the *pLRE::LRE-cYFP* construct with the gene of interest [7]. The

LLG1 promoter region for *pLLG1::Clevi-LRE/LLG1-cYFP* PCR is the same as previously

published [6]. PCR was performed with PrimeSTAR GXL DNA Polymerase (TaKaRa Bio

Inc.; Catalog # R050A) using primers and DNA templates listed in Table S4. The amplified

PCR products were cloned into *pLRE::LRE-cYFP* plasmid linearized with *SpeI-HF* (NEB,

Catalog # R3133S) and *AscI* (NEB, Catalog # R0558S) by using the In-Fusion HD Cloning

Plus system (Clontech, Catalog # 639645). The recombinant plasmids were transformed

into Stellar Competent Cells (Clontech, Catalog # 636763), and positive colonies were

selected on LB plates containing spectinomycin (100 µg/mL, Sigma-Aldrich, Catalog #

85555).

Constructs generated were sequence verified (Eton Bioscience, Inc.) before being

transformed into *Agrobacterium tumefaciens* (GV3101 pMP90 strain). The positive colony

selected for transforming into *A. thaliana* was also verified by colony PCR for the presence

of the transgene.

Plant transformation

491 Transformation solution containing *Agrobacterium tumefaciens* (GV3101 pMP90 strain)

492 harboring the desired transgene was either sprayed onto *A. thaliana* inflorescences or

493 applied by the floral dip method [48]. Hygromycin-resistant T_1 transformants were

494 selected as described [49].

Isolation of single-locus insertion lines

497 For each construct, single-locus insertion lines were isolated as described [7].

Scoring cYFP in mature unpollinated pistils

cYFP expression in mature unpollinated pistils were scored as described [7]. Samples
were mounted in 5% glycerol with a coverslip, and YFP expression in synergid cells was
scored by epifluorescence on a Zeiss Axiophot microscope with a GFP filter (excitation
HQ 470/40 and emission HQ 525/50). Images were acquired with Picture Frame
(Optronics).
Confocal images of the filiform apparatus in synergid cells were taken using a
Leica SP5 confocal laser scanning microscope. For cYFP imaging, samples were excited
with a 488-nm laser line, and emission spectra between 510 and 550 nm were collected.
YFP images were processed with Leica Application Suite X and ImageJ software
(http://imagej.nih.gov/ij/).

Complementation of root hair and hypocotyl length phenotypes in *llg1* seedlings

Root hair analysis was performed as previously described [18]. Root hairs located
between 1.5 and 3.5 mm from the primary root tip of four-day-old seedlings were observed
with a stereoscope. The number of normal and defective (stunted and collapsed) root
hairs was scored. The hypocotyl length assay was performed as described [6]. Three-
day-old dark-grown seedlings were imaged using Epson Perfection V370 Photo Scanner
at 600 dpi resolution and the length of hypocotyl was measured with Image J.

Propidium iodide staining of pavement cells

Six-day-old seedlings were stained with 50 μg/mL propidium iodide (PI) for 20 mins and then excess stain was removed by washing with ddH$_2$O. Images were acquired using confocal microscopy on a NIKON A1 Spectral System and analyzed by the NIS-Elements AR Analysis Software (V 5.02) with 40x objective. Images were acquired using optimal laser power and gain with excitation wavelength 561 nm and emission wavelength 595 nm.

RALF1-induced root growth inhibition assays

The root sensitivity assays with RALF1 treatment were performed as described [6, 50] : three-day-old light-grown seedlings were treated with RALF1 for 2 days at concentrations indicated in Figs 3, S2. Primary root length was measured using Image J at the beginning and end of treatments to obtain growth during treatment.

RALF23-induced seedling growth inhibition assays

Surface-sterilized T2 *pLLG1::SP-mRFP-LLG1* and *pLLG1::SP-mRFP-LLG2* seeds were selected on 1x MS-agar with 1% sucrose supplemented with kanamycin or glufosinate ammonium, respectively, alongside Col-0 and *llg1-2* on 1x MS-agar with 1% sucrose without selection. Five-day-old seedlings were then transferred to liquid 1x MS with 1% sucrose in sterile 48-well plates, with or without 1 μM RALF23 peptide (Scilight) [17]. Seedling fresh weight was measured after 7 days of growth in liquid medium.

ROS burst measurements

ROS burst measurements were performed as described previously [17]. Briefly, twelve 4-mm leaf discs from 4.5-week-old plants were harvested and equilibrated overnight in 96-well plates in sterile, deionized water. The next day, water was replaced with PAMP

solution (100 nM flg22 or elf18 peptide plus 10 µg/mL horse radish peroxidase and 100

µM luminol). ROS was immediately measured using a charge-coupled device camera

(Photek). Total ROS production was calculated as the sum of Random Light Units value

over 40 minutes of PAMP treatment.

Seed set scoring

Unfertilized, viable (enlarged after fertilization), and aborted ovules in siliques (10 days

after pollination) were scored as described [9]. Three to five self-pollinated siliques

located between 5th and 15th siliques from the bottom of the main branch of an *A. thaliana*

plant were scored.

GUS staining of *pLRE::GUS* and *pLLG1::GUS* transgenic plants

Tissues were stained for GUS activity as described [51]. Stained samples were mounted

in 50% glycerol and observed for epifluorescence on a Zeiss Axiovert 100 microscope

and images were taken using Metamorph (Version 7; Molecular Devices) using a red-

green-blue filter with autoexposure settings. For each tissue, at least five samples were

observed for each genotype.

Transcription factor binding site analysis in the promoters of *LLG* gene family

members in *A. thaliana* and *C. Violacea*

DAP-seq data were obtained from the Plant Cistrome Database

(http://neomorph.salk.edu/dev/pages/shhuang/dap_web/pages/index.php) in the form of

MEME motif file format. These 838 motifs were scanned against the putative promoter

sequences (1 kb upstream of the transcriptional start, not overlapping with neighboring

genes) of the *LLG* gene family members in *A. thaliana* (*LRE*, *LLG1*, *LLG2*, and *LLG3)*

and the single-copy orthologs *LRE/LLG1* and *LLG2/LLG3* in *Cleome violacea* with MAST

[52] using default parameters. The scanned results were used to determine the presence

of motif sites. A motif site was considered present in a putative promoter sequence if its

mapping p-value was <1e-3. The motif site information was combined for all motifs to

generate a motif presence (1)/absence (0) matrix for hierarchical clustering using the

heatmap.2 function in R.

RNA isolation and RT-PCR of *Cleome violacea*

The following tissues were collected from *C. violacea* for RNA isolation: rosette leaves

from 30-day-old plants, mature pistils 24 HAE with ovules removed, and mature ovules

24 HAE removed from the pistils. For each tissue type, three biological replicates were

collected. For rosette tissues, 3 rosette leaves were collected for each biological replicate.

Unlike *A. thaliana*, where every pistil has the potential to mature in every flower, we

observed that not all flowers in *C. violacea* develop a mature pistil. We determined mature

pistils alternate every 2–4 flowers; therefore, to collect pistils and ovules we first

monitored the maturation pattern and then emasculated 2–3 pistils which may mature.

Twenty-four hours later, mature pistils were dissected, ovules collected separately from

the other pistil tissues (septum, transmitting tract, stigma, style, carpel walls, nectaries).

100 ovules were collected for each biological replicate. For pistil without ovule tissues, 4

pistils were harvested for each biological replicate. All tissues were flash-frozen in liquid

nitrogen and stored at -80 °C until RNA extraction.

RNA was isolated using RNeasy Plant Mini Kit (Qiagen, Catalog # 74904)

according to the manufacturer's instructions and treated with RNase-free DNase I (Life

Technologies, Catalog # AM2222) to remove residual genomic DNA. Samples were

cleaned up using RNeasy MinElute Cleanup Kit (QIAGEN, Catalog #74204) and tested

for RNA integrity by Agilent Bioanalyzer 2100 (Agilent Technologies). cDNA was reverse

transcribed from 550 ng of total RNA using SuperScript IV First-Strand Synthesis System

(ThermoFisher Scientific, Catalog # 18091050).

PCR was performed with 20 ng of cDNA for each reaction, using the following PCR

cycle conditions: 1. 98 °C for 2 minutes; 2. 95 °C for 30 seconds, 3. 56 °C for 20 seconds,

4. 72 °C for 1 minute 10 seconds, 5. Repeat cycles 2–5 for 34 cycles, 6. 72 °C for 10

minutes, 7. Hold at 4 °C. PCR products were analyzed by gel electrophoresis on a 1%

agarose gel with ethidium bromide in 1X TAE buffer. Gel images were processed using
ImageJ. *Clevi-LRE/LLG1* and the control gene *Clevi-ACTIN2* were amplified using
primers listed in Table S4. Two technical replicates of each biological replicate were
performed.

Image processing

ImageJ was used to assemble image panels, insert scale bars, and prepare figures.

Genomes and accession numbers

A. thaliana LRE (*At4g26466*), *LLG1* (*At5g56170*), *LLG2* (*At2g20700*), and *LLG3*
(*At4g28280*) were obtained from The Arabidopsis Information Resource (Version:
TAIR10). *LRE, LLG1, LLG2, LLG3, LRE/LLG1,* and *LLG2/LLG3* orthologs were identified
using the Comparative Genomics platform (CoGe) [40, 41]. Genome references and
accession numbers can be found in Table S1.

Acknowledgements

We acknowledge Dr. Patrick Edger from Michigan State University for providing us
access to the *Cleome violacea* genome on CoGe and Dr. Jocelyn Hall from the University
of Alberta for providing *Cleome violacea* seeds. We thank Ramin Yadegari (University of
Arizona) for the Zeiss Axiophot microscope. We thank the past and present members of
the Palanivelu Lab for discussions. J.A.N. was supported by the following: IGERT
Comparative Genomics Program at the University of Arizona (Award ID: 0654435); NSF
Graduate Research Fellowship: Grant DGE-1143953; the Boynton Graduate Fellowship;
the American Society of Plant Biologists; and the University of Arizona Graduate College
Office of Diversity and Inclusion. Additional support for this work was provided by an NSF
grant to R.P. (IOS-1146090) and University of Arizona Undergraduate Biology Research
Program fellowship to S.H. (private donors). This work was also supported by the

University of Zürich (C.Z.), the European Research Council under the Grant Agreement

773153 (grant IMMUNO-PEPTALK to C.Z.), as well as fellowships from the European

Molecular Biology Organization, the Natural Sciences and Engineering Research Council

of Canada and the Deutsche Forschungsgemeinschaft (fellowships EMBO-LTF 100-2017

and NSERC PDF-532561-2019 to T.A.D.; DFG STE 2448/1 to M.S.T.). Work in A.Y.C.

lab was supported by Natural Science Foundation (IOS-1645854 and MCB-1715764) to

A.Y.C. and Hen-Ming Wu, and the National Institute of Food and Agriculture, U.S.

Department of Agriculture, the Center for Agriculture, Food and the Environment under

project number MAS00525. The contents are solely the responsibility of the authors and

do not necessarily represent the official views of the USDA or NIFA. We also thank

H.M.W. for contribution of LLG1 and LLG3 constructs. K.M. was supported by the Torrey

Summer Research Scholarship from UMass Plant Biology Program and the Linda Slakey

Summer Research Scholarship from UMass BMB department. This study was also

supported by funding from NSF pollen RCN grant for sponsoring activities and meetings

that forged collaborations between A.Y.C. and R.P. labs (MCB0955910).

Author contributions

JAN, MJL, TAD, MS, KM, BS, KD, NK, and SH planned and designed the research in

consultation with and guidance of their respective principal investigators, performed

experiments, collected and/or analyzed data. SS performed the transcription factor

binding site analysis in the promoters of *LLG* family members. JAN and MAB performed

all the evolutionary analysis reported in this study. JAN, RP, MJL, MS, TAD, SS, CZ,

MAB, and AYC wrote the manuscript with input from and revisions by all authors.

References

1.   Johnson MA, Harper JF, Palanivelu R. A Fruitful Journey: Pollen Tube Navigation from Germination to Fertilization. Annual Review of Plant Biology. 2019. doi: 10.1146/annurev-arplant-050718-100133.

2.   Capron A, Gourgues M, Neiva LS, Faure JE, Berger F, Pagnussat G, et al. Maternal control of male-gamete delivery in *Arabidopsis* involves a putative GPI-anchored protein encoded by the *LORELEI* gene. Plant Cell. 2008;20(11):3038-49. PubMed PMID: 19028964.

3.   Escobar-Restrepo J-M, Huck N, Kessler S, Gagliardini V, Gheyselinck J, Yang W-C, et al. The FERONIA Receptor-like Kinase Mediates Male-Female Interactions During Pollen Tube Reception. Science. 2007;317(5838):656-60. doi: 10.1126/science.1143562.

4.   Huck N, Moore JM, Federer M, Grossniklaus U. The Arabidopsis mutant feronia disrupts the female gametophytic control of pollen tube reception. Development. 2003;130(10):2149-59. doi: 10.1242/dev.00458.

5.   Kessler SA, Lindner H, Jones DS, Grossniklaus U. Functional analysis of related CrRLK1L receptor-like kinases in pollen tube reception. EMBO reports. 2015;16(1):107-15. doi: 10.15252/embr.201438801.

6.   Li C, Yeh F-L, Cheung AY, Duan Q, Kita D, Liu M-C, et al. Glycosylphosphatidylinositol-anchored proteins as chaperones and co-receptors for FERONIA receptor kinase signaling in Arabidopsis. eLife. 2015;4:e06587. doi: 10.7554/eLife.06587.

7.   Liu X, Castro C, Wang Y, Noble J, Ponvert N, Bundy M, et al. The Role of LORELEI in Pollen Tube Reception at the Interface of the Synergid Cell and Pollen

Tube Requires the Modified Eight-Cysteine Motif and the Receptor-Like Kinase

FERONIA. The Plant Cell. 2016;28(5):1035-52. doi: 10.1105/tpc.15.00703.

8.   Rotman N, Rozier F, Boavida L, Dumas C, Berger F, Faure JE. Female control of

male gamete delivery during fertilization in *Arabidopsis thaliana*. Curr Biol.

2003;13(5):432-6. PubMed PMID: 12620194.

9.   Tsukamoto T, Qin Y, Huang Y, Dunatunga D, Palanivelu R. A role for LORELEI,

a putative glycosylphosphatidylinositol-anchored protein, in Arabidopsis thaliana double

fertilization and early seed development. Plant J. 2010;62(4):571-88. PubMed PMID:

20163554.

10.   Duan Q, Kita D, Johnson EA, Aggarwal M, Gates L, Wu HM, et al. Reactive

oxygen species mediate pollen tube rupture to release sperm for fertilization in

Arabidopsis. Nature communications. 2014;5:3129. doi: 10.1038/ncomms4129.

PubMed PMID: 24451849.

11.   Duan Q, Liu M-CJ, Kita D, Jordan SS, Yeh F-LJ, Yvon R, et al. FERONIA

controls pectin- and nitric oxide-mediated male–female interaction. Nature. 2020. doi:

10.1038/s41586-020-2106-2.

12.   Ge Z, Bergonci T, Zhao Y, Zou Y, Du S, Liu M-C, et al. Arabidopsis pollen tube

integrity and sperm release are regulated by RALF-mediated signaling. Science. 2017.

doi: 10.1126/science.aao3642.

13.   Mecchia MA, Santos-Fernandez G, Duss NN, Somoza SC, Boisson-Dernier A,

Gagliardini V, et al. RALF4/19 peptides interact with LRX proteins to control pollen tube

growth in Arabidopsis. Science. 2017. doi: 10.1126/science.aao5467.

14.   Feng H, Liu C, Fu R, Zhang M, Li H, Shen L, et al. LORELEI-LIKE GPI-

ANCHORED PROTEINS 2/3 regulate pollen tube growth as chaperones and

coreceptors for ANXUR/BUPS receptor kinases in Arabidopsis. Molecular plant. 2019.

doi: https://doi.org/10.1016/j.molp.2019.09.004.

15.   Ge Z, Dresselhaus T, Qu L-J. How CrRLK1L Receptor Complexes Perceive

RALF Signals. Trends in plant science. 2019;24(11):978-81. doi:

10.1016/j.tplants.2019.09.002.

16.   Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton

N, et al. The receptor kinase FER is a RALF-regulated scaffold controlling plant immune

signaling. Science. 2017;355(6322):287-9. doi: 10.1126/science.aal2541.

17.   Xiao Y, Stegmann M, Han Z, DeFalco TA, Parys K, Xu L, et al. Mechanisms of

RALF peptide perception by a heterotypic receptor complex. Nature.

2019;572(7768):270-4. doi: 10.1038/s41586-019-1409-7.

18.   Duan Q, Kita D, Li C, Cheung AY, Wu H-M. FERONIA receptor-like kinase

regulates RHO GTPase signaling of root hair development. Proceedings of the National

Academy of Sciences. 2010;107(41):17821-6. doi: 10.1073/pnas.1005366107.

19.   Campbell L, Turner SR. A Comprehensive Analysis of RALF Proteins in Green

Plants Suggests There Are Two Distinct Functional Groups. Frontiers in plant science.

2017;8:37. Epub 2017/02/09. doi: 10.3389/fpls.2017.00037. PubMed PMID: 28174582;

PubMed Central PMCID: PMCPMC5258720.

20.   Moussu S, Broyart C, Santos-Fernandez G, Augustin S, Wehrle S, Grossniklaus

U, et al. Structural basis for recognition of RALF peptides by LRX proteins during pollen

tube growth. Preprint at https://wwwbiorxivorg/content/101101/695874v1 2019. doi:

10.1101/695874.

21.   Dünser K, Gupta S, Herger A, Feraru MI, Ringli C, Kleine-Vehn J. Extracellular

matrix sensing by FERONIA and Leucine-Rich Repeat Extensins controls vacuolar

expansion during cellular elongation in Arabidopsis thaliana. The EMBO Journal.

2019;38(7):e100353. doi: 10.15252/embj.2018100353.

22.   Bowers JE, Chapman BA, Rong J, Paterson AH. Unravelling angiosperm

genome evolution by phylogenetic analysis of chromosomal duplication events. Nature.

2003;422(6930):433-8. Epub 2003/03/28. doi: 10.1038/nature01521. PubMed PMID:

12660784.

23.   Lynch M, Conery JS. The evolutionary demography of duplicate genes. Journal

of Structural and Functional Genomics. 2003;3(1):35-44. doi:

10.1023/A:1022696612931.

24.   Blanc G, Hokamp K, Wolfe KH. A Recent Polyploidy Superimposed on Older

Large-Scale Duplications in the Arabidopsis Genome. Genome Research.

2003;13(2):137-44.

25.   Maere S, De Bodt S, Raes J, Casneuf T, Van Montagu M, Kuiper M, et al.

Modeling gene and genome duplications in eukaryotes. Proceedings of the National

Academy of Sciences of the United States of America. 2005;102(15):5454. doi:

10.1073/pnas.0501102102.

26.   Shen Q, Bourdais G, Pan H, Robatzek S, Tang D. Arabidopsis

glycosylphosphatidylinositol-anchored protein LLG1 associates with and modulates

FLS2 to regulate innate immunity. Proceedings of the National Academy of Sciences.

2017. doi: 10.1073/pnas.1614468114.

27.   Wang Y, Tsukamoto T, Noble JA, Liu X, Mosher RA, Palanivelu R. Arabidopsis

LORELEI, a Maternally Expressed Imprinted Gene, Promotes Early Seed Development.

Plant physiology. 2017;175(2):758-73. doi: 10.1104/pp.17.00427.

28.   Duarte JM, Cui L, Wall PK, Zhang Q, Zhang X, Leebens-Mack J, et al.

Expression pattern shifts following duplication indicative of subfunctionalization and

neofunctionalization in regulatory genes of Arabidopsis. Mol Biol Evol. 2006;23(2):469-

78. Epub 2005/11/11. doi: 10.1093/molbev/msj051. PubMed PMID: 16280546.

29.   Force A, Lynch M, Pickett FB, Amores A, Yan Y-l, Postlethwait J. Preservation of

Duplicate Genes by Complementary, Degenerative Mutations. Genetics.

1999;151(4):1531-45.

30.   Panchy N, Lehti-Shiu M, Shiu S-H. Evolution of Gene Duplication in Plants. Plant

physiology. 2016;171(4):2294. doi: 10.1104/pp.16.00523.

31.   Ohno S. Evolution by Gene Duplication: Springer-Verlag; 1970.

32.   De Smet R, Sabaghian E, Li Z, Saeys Y, Van de Peer Y. Coordinated Functional

Divergence of Genes after Genome Duplication in Arabidopsis thaliana. The Plant Cell.

2017;29(11):2786. doi: 10.1105/tpc.17.00531.

33.   Liu S-L, Baute GJ, Adams KL. Organ and cell type-specific complementary

expression patterns and regulatory neofunctionalization between duplicated genes in

Arabidopsis thaliana. Genome biology and evolution. 2011;3:1419-36. Epub

2011/11/04. doi: 10.1093/gbe/evr114. PubMed PMID: 22058183.

34.   Zou C, Lehti-Shiu MD, Thomashow M, Shiu S-H. Evolution of Stress-Regulated

Gene Expression in Duplicate Genes of Arabidopsis thaliana. PLOS Genetics.

2009;5(7):e1000581. doi: 10.1371/journal.pgen.1000581.

35.   Roulin A, Auer PL, Libault M, Schlueter J, Farmer A, May G, et al. The fate of

duplicated genes in a polyploid plant genome. The Plant Journal. 2013;73(1):143-53.

doi: 10.1111/tpj.12026.

36.   Hughes TE, Langdale JA, Kelly S. The impact of widespread regulatory

neofunctionalization on homeolog gene evolution following whole-genome duplication in

maize. Genome Research. 2014;24(8):1348-55. doi: 10.1101/gr.172684.114.

37.   Sharma A, Hussain A, Mun BG, Imran QM, Falak N, Lee SU, et al.

Comprehensive analysis of plant rapid alkalization factor (RALF) genes. Plant

physiology and biochemistry : PPB / Societe francaise de physiologie vegetale.

2016;106:82-90. Epub 2016/05/08. doi: 10.1016/j.plaphy.2016.03.037. PubMed PMID:

27155375.

38.   Findlay GD, Sitnik JL, Wang W, Aquadro CF, Clark NL, Wolfner MF. Evolutionary

Rate Covariation Identifies New Members of a Protein Network Required for Drosophila

melanogaster Female Post-Mating Responses. PLoS Genet. 2014;10(1):e1004108. doi:

10.1371/journal.pgen.1004108.

39.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment

search tool. J Mol Biol. 1990;215(3):403-10. Epub 1990/10/05. doi: 10.1016/s0022-

2836(05)80360-2. PubMed PMID: 2231712.

40.   Lyons E, Freeling M. How to usefully compare homologous plant genes and

chromosomes as DNA sequences. The Plant journal : for cell and molecular biology.

2008;53(4):661-73. Epub 2008/02/14. doi: 10.1111/j.1365-313X.2007.03326.x. PubMed

PMID: 18269575.

41.   Lyons E, Pedersen B, Kane J, Alam M, Ming R, Tang H, et al. Finding and

Comparing Syntenic Regions among Arabidopsis and the Outgroups Papaya, Poplar,

and Grape: CoGe with Rosids. Plant physiology. 2008;148(4):1772. doi:

10.1104/pp.108.124867.

42.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 2004;32(5):1792-7. Epub 2004/03/23. doi:

10.1093/nar/gkh340. PubMed PMID: 15034147; PubMed Central PMCID:

PMCPMC390337.

43.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time

and space complexity. BMC Bioinformatics. 2004;5(1):113. doi: 10.1186/1471-2105-5-

113.

44.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 2014;30(9):1312-3. doi:

10.1093/bioinformatics/btu033.

45.   Kessler SA, Shimosato-Asano H, Keinath NF, Wuest SE, Ingram G, Panstruga

R, et al. Conserved molecular components for pollen tube reception and fungal

invasion. Science. 2010;330(6006):968-71. PubMed PMID: 21071669.

46.   Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, et al.

Development of series of gateway binary vectors, pGWBs, for realizing efficient

construction of fusion genes for plant transformation. Journal of Bioscience and

Bioengineering. 2007;104(1):34-41. doi: https://doi.org/10.1263/jbb.104.34.

47.   Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, et al. Improved

Gateway Binary Vectors: High-Performance Vectors for Creation of Fusion Constructs

in Transgenic Analysis of Plants. Bioscience, Biotechnology, and Biochemistry.

2007;71(8):2095-100. doi: 10.1271/bbb.70216.

48.   Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. The Plant journal : for cell and molecular biology.

1998;16(6):735-43. Epub 1999/03/09. doi: 10.1046/j.1365-313x.1998.00343.x. PubMed

PMID: 10069079.

49.   Harrison SJ, Mott EK, Parsley K, Aspinall S, Gray JC, Cottage A. A rapid and

robust method of identifying transformed Arabidopsis thaliana seedlings following floral

dip transformation. Plant Methods. 2006;2(1):19. doi: 10.1186/1746-4811-2-19.

50. Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR. A Peptide Hormone
and Its Receptor Protein Kinase Regulate Plant Cell Expansion. Science.
2014;343(6169):408-11. doi: 10.1126/science.1244454.

51. Johnson MA, Kost B. Pollen tube development. Methods Mol Biol. 2010;655:155-
76. PubMed PMID: 20734260.

52. Bailey TL, Gribskov M. Combining evidence using p-values: application to
sequence homology searches. Bioinformatics. 1998;14(1):48-54. Epub 1998/04/01. doi:
10.1093/bioinformatics/14.1.48. PubMed PMID: 9520501.

Figure Legends

Fig. 1. The *LLG* gene family is maintained in the Brassicaceae family.

A maximum likelihood phylogenetic tree of full-length CDS of single copy orthologs of *LRE* and *LLG1* (*LRE/LLG1*) and *LLG2* and *LLG3* (*LLG2/LLG3)* in *Carica papaya* (pink and grey boxes) and two species in the Cleomaceae (*Cleome violacea* and *Tarenaya hassleriana*) (purple and red boxes, respectively) and 11 species in the Brassicaceae. The *LRE* and its orthologs formed a distinct clade (blue box) from *LLG1* and its orthologs (orange box). *LLG2* and its orthologs (green box) formed a separate clade than *LLG3* and its orthologs (yellow box). The phylogenetic tree was generated with 100 bootstrap replicates and was rooted using a single copy ortholog identified in *Amborella trichopoda*. Only bootstrap values ≥ 50 are represented along each branch.

Fig 2. LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complemented vegetative defects in *llg1*.

(A) Complementation of root hair defects in *llg1-2* seedlings. Root hairs were scored as normal or defective in at least 10 seedlings in each indicated single-insertion line carrying LRE-HA, LLG3, or Clevi-LRE/LLG1-cYFP fusion protein. (B) Complementation of short hypocotyl lengths in dark grown *llg1-2* mutant seedlings. Hypocotyl length was assayed in T2 seedlings from selfed seeds of single insertion lines carrying LRE-HA, LLG3, or Clevi-LRE/LLG1-cYFP fusion protein. Quantification of hypocotyl lengths was done in at least three trials of 20-25 seedlings for each line. Error bars represent ±SD.

Fig 3. LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complemented epidermal defects and RALF1 insensitivity in *llg1-2* seedlings.

(A-F) Epidermal pavement cells of 6-day-old *llg1-2* seedlings expressing LRE-HA, LLG3, or Clevi-LRE/LLG1-cYFP showed restored normal pavement cell morphology like that seen in wild-type (Col-0) and *llg1-2* seedlings expressing LLG1. In each genotype, 5-10 seedlings were stained with Propidium Iodide (PI) and visualized with confocal

microscopy. (G) Percentage of root growth after RALF1 treatment of three-day-old

seedlings as described in S2 Fig. Root length was measured two days after RALF1

treatment, and three trials were performed. Error bars represent ±SD.

Fig 4. mRFP-LLG1 and mRFP-LLG2 complemented defects in *llg1-2*.

(A) Expression of *pLLG1::SP-mRFP-LLG*1 or *pLLG1::SP-mRFP-LLG2* restores rosette

size in 4.5-week-old *llg1-2* plants. (B) ROS production in response to flg22 (left) or elf18

(right) is restored in SP-mRFP-LLG1 or SP-mRFP-LLG2 lines driven by the *LLG1* native

promoter. Total Relative Light Unit (RLU) over 40 minutes of exposure to 100 nM flg22 or

elf18 treatment is displayed. Letters indicate significantly different values (n=12 leaf discs,

two-way ANOVA with Tukey test, flg22 $p<0.0001$; elf18 $p=0.0009$). Error bars show ±SD.

(C) RALF23 sensitivity is restored in seedlings expressing SP-mRFP-LLG1 or SP-mRFP-

LLG2. Letters indicate significantly different values (n=16 seedlings, two-way ANOVA

with Tukey test, $p<0.0001$). Error bars show ±SD.

Fig 5. *pLRE::GUS* and *pLLG1::GUS* showed non-overlapping expression in

vegetative and reproductive tissues.

(A-F) *pLLG1::GUS* was expressed, while *pLRE::GUS* was not, in vegetative tissues. In 8-

day-old seedlings, *pLLG1::GUS* was expressed in true leaves, hypocotyls, and roots (A-

C). In 21-day-old seedlings, *pLLG1::GUS* was expressed in the epicotyl, the hypocotyl,

and weakly expressed in roots (D-F). (G) At 24 HAE, *pLRE::GUS* and *pLLG1::GUS* were

both expressed in pistils but in different cell-types. *pLRE::GUS* was expressed in synergid

cells, while *pLLG1::GUS* was expressed in septum. Close up of area marked in red

rectangles are shown below. (H-I) *pLRE::GUS* was expressed in synergid cells at 24

hours after emasculation (HAE) (H) but *pLLG1::GUS* is not expressed in the ovule (I). (J-

M) *pLRE::GUS* and *pLLG1::GUS* showed non-overlapping expression after pollination.

Mature unpollinated pistils were pollinated with Col-0 pollen and collected at 13.5 HAP

(J-K) or 18 HAP (I-M) and stained for GUS activity. (J-K) At 13.5 HAP, *pLRE::GUS* was

expressed in the micropylar end of the female gametophyte (J), while *pLLG1::GUS*

continues to be expressed in the septum (K). (L-M) At 18 HAP, *pLRE::GUS* and
pLLG1::GUS were both expressed in pollinated pistils but in different cell-types.
pLRE::GUS was expressed in the zygote and developing endosperm nuclei, while
pLLG1::GUS was expressed in septum.

Fig 6. LLG1-cYFP and Clevi-LRE/LLG1-cYFP complemented reproductive defects
894
in *lre*.

(A) Images of opened siliques of indicated genotypes in *A. thaliana.* A representative
unfertilized ovule (*) and viable (V) or aborted (A) seed is marked in the *lre* silique. (B)
LLG1-cYFP complemented lre mutant seed set defects in self-pollinated pistils of
indicated three independent transformants (ANOVA, p = 0.18). (C) Clevi-LRE/LLG1-cYFP
partially complemented lre mutant seed set defects in self-pollinated pistils of indicated
three independent transformants (pairwise two-tailed t-tests, p > 0.05). (B,C) Number in
the middle of each column refers to the number of ovules/seeds scored. Groups sharing
same lowercase letters are similar to each other in statistical tests.

Fig 7. *Clevi-LRE/LLG1* and *Clevi-LLG2/LLG3* are expressed in vegetative and
reproductive tissues of *Cleome violacea*.

RT-PCR of full-length *Clevi-LRE/LLG1* or *Clevi-LLG2/LLG3* in cDNAs isolated from 30-
day-old rosette leaves, anther and pollen, emasculated pistils without ovules, and ovules
from emasculated pistils of *Cleome violacea*. A homolog of *A. thaliana ACTIN2* (*Clevi-*
ACTIN2) was used as a control in these experiments. gDNA, genomic DNA isolated from
of Cleome leaves was used as a positive control in PCR portion of the RT-PCR
experiment. RT-PCR was repeated with two additional biological replicates with similar
results and the amplified bands were sequenced to confirm the identity of amplified
cDNAs.

Supplemental Information

Fig S1. LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complemented vegetative defects in *llg1-2* seedlings.

(A) Diagrams of the *pLLG1::LRE-HA, pLLG1::LLG3,* and *pLLG1::Clevi-LRE/LLG1-cYFP* constructs. (B) Root hair length defects were restored when LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP were expressed in *llg1-2* seedlings to wild-type (Col-0) levels. Images were taken from T2 seedlings of single insertion lines. Quantification of data shown here are reported in Fig 2A. (C) LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complementation of short hypocotyl length phenotype in *llg1-2* seedlings. Representative images of wild-type (Col-0) or T2 seedlings of single-locus insertion lines of *pLLG1::LRE-HA, pLLG1::LLG3,* or *pLLG1::Clevi-LRE/LLG1-cYFP* are shown. Quantification of data shown here are reported in Fig 2B.

Fig S2. LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complemented root hair lengths in *llg1-2* seedlings prior to RALF treatment and increased RALF1 sensitivity after treatment.

Root lengths before and after RALF1 treatment were measured in single representative lines from *LRE-HA, LLG3,* and *Clevi-LRE/LLG1-cYFP*. Root lengths in three-day-old seedlings were measured, then treated with 0 µM RALF1 (untreated), 0.5 µM RALF1, or 1 µM RALF1. Roots were measured two days after RALF1 treatments with three trial replicates. Error bars represent ±SD.

Fig S3 ROS production kinetics in response to flg22 (top) or elf18 (bottom) is restored in *pLLG1::SP-mRFP-LLG1* or *pLLG1::SP-mRFP-LLG2* lines.

ROS burst levels in response to 100 nM flg22 (top) or elf18 (bottom) over time is indicated in Relative Light Unit (RLU). In each trace of indicated genotypes, data shown is average of n=12 leaf discs, ±SE

Fig S4 Distribution of putative transcription factor (TF) binding sites in the putative promoter regions of LRE homologs in *Arabidopsis thaliana* and *Cleome violacea*.

(A) TFs with putative binding sites as determined using DNA affinity purification-sequencing (DAP-seq) are shown as present (yellow boxes, *p* < 1e-4) or absent (blue boxes). The TF family and gene name information were based on the Plant Cistrome Database. *: amplified DAP-seq where secondary DNA modifications were removed. (B) Table showing the frequency of putative TF binding site occurrence in the promoters of *LRE*, *LLG1*, and the single copy ortholog *LRE/LLG1* in *Cleome violacea.* (C) Table showing the frequency of putative TF binding site occurrence in the promoters of *LLG2*, *LLG3*, and the single copy ortholog *LLG2/LLG3* in *Cleome violacea.*

Fig S5 LLG1-cYFP and Clevi-LRE/LLG1-cYFP were expressed in synergid cells and localized to the FA.

(A-B) Diagrams of the *pLRE::LLG1-cYFP* and *pLRE::Clevi-LRE/LLG1-cYFP* constructs. (C) A diagram of a mature ovule with a 7-celled female gametophyte. Synergid cells are located in the micropylar end of the ovule, adjacent to the egg cell. The finger-like projections of the FA are shown in yellow. A red arrow points to the FA. (D) In mature unpollinated pistils, LRE-cYFP is expressed in the synergid cells, with localization in the puncta in the synergid cell cytoplasm and in the FA. The ovule is outlined in light gray dashed line, while the female gametophyte is outlined in dark gray dashed line. The red rectangle marks the synergid cells. (E) Close-up image of the LRE-cYFP in the synergid cells marked by the red rectangle in Fig C, outlined in dark gray dashed line. YFP localized in the FA and puncta in the synergid cells. (F) Close-up image of the LLG1-cYFP in the synergid cells with YFP localization in the puncta and the FA of the synergid cells. (g) Close-up image of the Clevi-LRE/LLG1-cYFP in the synergid cells. YFP is weakly expressed in the FA, but was not present elsewhere in the synergid cells, including the puncta

Table S1. List of genes and genomes.

973 **Table S2.** List of primers used in this study.

Table 1. Enhanced transmission of the *pLRE::LLG1-cYFP* transgene through the *lre-7* female gametophyte.

Parents		Observed No. of progeny		Transmission Efficiency (TE) Analysis		
Female parent+	Male parent+	HygR*	HygS*	TE (R/S)	χ2†	P-value
WT	*LRE-cYFP-23*	129	116	1.11	0.34	0.56#
LRE-cYFP-23	*WT*	156	31	5.03	47.13	6.64E-12
WT	*LLG1-cYFP-A2*	213	171	1.25	2.30	0.13#
LLG1-cYFP-A2	*WT*	163	24	6.79	60.05	9.25E-15
WT	*LLG1-cYFP-10*	60	71	0.85	0.46	0.50#
LLG1-cYFP-10	*WT*	179	30	5.97	60.95	5.87E-15
WT	*LLG1-cYFP-11*	147	79	1.86	10.47	0.001#
LLG1-cYFP-11	*WT*	138	16	8.63	57.32	3.71E-14

+ Line Numbers refer to three independent transformants in the *lre-7/lre-7* background containing single insertion of the *pLRE::LLG1-cYFP* transgene. Genotype of each transgenic line is heterozygous for the transgene (*pLRE::LLG1-cYFP/+*) and homozygous for the *lre-7* mutation (*lre-7/lre-7*).

* Hygromycin resistant (HygR) and susceptible (HygS) progeny. Hygromycin resistance gene is linked to the construct carrying the *pLRE::LLG1-cYFP* transgene.

TE, Transmission efficiency was calculated as the ratio of hygromycin resistance (R) to susceptibility (S) in the progeny of the indicated cross

† χ2 is calculated based on an expected segregation ratio of hygromycin resistant to susceptibility of 1:1

No significant deviation from 1:1 segregation through the male gametophyte indicates that pollen parent contains a single insertion of the *pLRE::LLG1-cYFP* transgene. Additional details on our protocol to isolate single insertion lines can be found in the methods.

Table 2. Enhanced transmission of the *pLRE::Clevi-LRE/LLG1- cYFP* transgene through the *lre-7* female gametophyte.

Parents		Observed No. of progeny		Transmission Efficiency (TE) Analysis		
Female parent+	Male parent+	HygR*	HygS*	TE (R/S)	χ2†	P-value
WT	*LRE-cYFP-23*	212	182	1.16	1.14	0.28#
LRE-cYFP-23	*WT*	335	43	7.79	132.56	1.13E-30
WT	*Clevi-LRE/LLG1-cYFP-10*	127	130	0.98	0.02	0.89#
Clevi-LRE/LLG1-cYFP-10	*WT*	87	52	1.67	4.49	0.03
WT	*Clevi-LRE/LLG1-cYFP-11*	150	113	1.33	2.62	0.10#
Clevi-LRE/LLG1-cYFP-11	*WT*	109	52	2.10	10.44	0.001
WT	*Clevi-LRE/LLG1-cYFP-12*	90	88	1.02	0.01	0.92#
Clevi-LRE/LLG1-cYFP-12	*WT*	84	28	3	14.93	0.0001

+ Line Numbers refer to three independent transformants in the *lre-7/lre-7* background containing single insertion of the *pLRE::Clevi-LRE/LLG1-cYFP* transgene. Genotype of each transgenic line is heterozygous for the transgene (*pLRE::Clevi-LRE/LLG1-cYFP/+*) and homozygous for the *lre-7* mutation (*lre-7/lre-7*).

* Hygromycin resistant (HygR) and susceptible (HygS) progeny. Hygromycin resistance gene is linked to the construct carrying the *pLRE::Clevi-LRE/LLG1-cYFP* transgene.

TE, Transmission efficiency was calculated as the ratio of hygromycin resistance (R) to susceptibility (S) in the progeny of the indicated cross

† χ2 is calculated based on an expected segregation ratio of hygromycin resistant to susceptibility of 1:1

Fig 1 The *LLG* gene family is maintained in the Brassicaceae family.

A maximum likelihood phylogenetic tree of full-length CDS of single copy orthologs of *LRE* and *LLG1* (*LRE/LLG1*) and *LLG2* and *LLG3* (*LLG2/LLG3)* in *Carica papaya* (pink and grey boxes) and two species in the Cleomaceae (*Cleome violacea* and *Tarenaya hassleriana*) (purple and red boxes, respectively) and 11 species in the Brassicaceae. The *LRE* and its orthologs formed a distinct clade (blue box) from *LLG1* and its orthologs (orange box). *LLG2* and its orthologs (green box) formed a separate clade than *LLG3* and its orthologs (yellow box). The phylogenetic tree was generated with 100 bootstrap replicates and was rooted using a single copy ortholog identified in *Amborella trichopoda*. Only bootstrap values ≥ 50 are represented along each branch.

Fig 2 LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complemented vegetative defects in *llg1*.

(a) Complementation of root hair defects in *llg1-2* seedlings. Root hairs were scored as normal or defective in at least 10 seedlings in each indicated single-insertion line carrying LRE-HA, LLG3, or Clevi-LRE/LLG1-cYFP fusion protein.

(b) Complementation of short hypocotyl lengths in dark grown *llg1-2* mutant seedlings. Hypocotyl length was assayed in T2 seedlings from selfed seeds of single insertion lines carrying LRE-HA, LLG3, or Clevi-LRE/LLG1-cYFP fusion protein. Quantification of hypocotyl lengths was done in at least three trials of 20-25 seedlings for each line. Error bars represent ±SD.

Fig 3 LRE-HA, LLG3, and Clevi-LRE/LLG1-cYFP complemented epidermal defects and RALF1 insensitivity in *llg1-2* seedlings.

(a-f) Epidermal pavement cells of 6-day-old *llg1-2* seedlings expressing LRE-HA, LLG3, or Clevi-LRE/LLG1-cYFP showed restored normal pavement cell morphology like that seen in wild-type (Col-0) and *llg1-2* seedlings expressing LLG1. In each genotype, 5-10 seedlings were stained with Propidium Iodide (PI) and visualized with confocal microscopy.

(g) Percentage of root growth after RALF1 treatment of three-day-old seedlings as described in Supplemental Figure 2. Root length was measured two days after RALF1 treatment, and three trials were performed. Error bars represent ±SD.

Fig 4 mRFP-LLG1 and mRFP-LLG2 complemented defects in *llg1-2*.

(a) Expression of *pLLG1::SP-mRFP-LLG*1 or *pLLG1::SP-mRFP-LLG2* restores rosette size in 4.5-week-old *llg1-2* plants.

(b) ROS production in response to flg22 (left) or elf18 (right) is restored in SP-mRFP-LLG1 or SP-mRFP-LLG2 lines driven by the *LLG1* native promoter. Total Relative Light Unit (RLU) over 40 minutes of exposure to 100 nM flg22 or elf18 treatment is displayed. Letters indicate significantly different values (n=12 leaf discs, two-way ANOVA with Tukey test, flg22 p<0.0001; elf18 p=0.0009). Error bars show ±SD.

(c) RALF23 sensitivity is restored in seedlings expressing SP-mRFP-LLG1 or SP-mRFP-LLG2. Letters indicate significantly different values (n=16 seedlings, two-way ANOVA with Tukey test, p<0.0001). Error bars show ±SD.

Fig 5 *pLRE::GUS* and *pLLG1::GUS* showed non-overlapping expression in vegetative and reproductive tissues.

(a-f) *pLLG1::GUS* was expressed, while *pLRE::GUS* was not, in vegetative tissues. In 8-day-old seedlings, *pLLG1::GUS* was expressed in true leaves, hypocotyls, and roots (a-c). In 21-day-old seedlings, *pLLG1::GUS* was expressed in the epicotyl, the hypocotyl, and weakly expressed in roots (d-f).

(g) At 24 HAE, *pLRE::GUS* and *pLLG1::GUS* were both expressed in pistils but in different cell-types. *pLRE::GUS* was expressed in synergid cells, while *pLLG1::GUS* was expressed in septum. Close up of area marked in red rectangles are shown below.

(h-i) *pLRE::GUS* was expressed in synergid cells at 24 hours after emasculation (HAE) (H) but *pLLG1::GUS* is not expressed in the ovule (i).

(j-m) *pLRE::GUS* and *pLLG1::GUS* showed non-overlapping expression after pollination. Mature unpollinated pistils were pollinated with Col-0 pollen and collected at 13.5 HAP (j-k) or 18 HAP (l-Mm and stained for GUS activity.

(j-k) At 13.5 HAP, *pLRE::GUS* was expressed in the micropylar end of the female gametophyte (j), while *pLLG1::GUS* continues to be expressed in the septum (k).

(l-m) At 18 HAP, *pLRE::GUS* and *pLLG1::GUS* were both expressed in pollinated pistils but in different cell-types. *pLRE::GUS* was expressed in the zygote and developing endosperm nuclei, while *pLLG1::GUS* was expressed in septum.

Fig 6 LLG1-cYFP and Clevi-LRE/LLG1-cYFP complemented reproductive defects in *lre*.

(a) Images of opened siliques of indicated genotypes in *A. thaliana.* A representative unfertilized ovule (*) and viable (V) or aborted (A) seed is marked in the *lre* silique.

(b) LLG1-cYFP complemented lre mutant seed set defects in self-pollinated pistils of indicated three independent transformants (ANOVA, p = 0.18).

(c) Clevi-LRE/LLG1-cYFP partially complemented lre mutant seed set defects in self-pollinated pistils of indicated three independent transformants (pairwise two-tailed t-tests, p > 0.05).

(b,c) Number in the middle of each column refers to the number of ovules/seeds scored. Groups sharing same lowercase letters are similar to each other in statistical tests.

Fig 7 *Clevi-LRE/LLG1* and *Clevi-LLG2/LLG3* are expressed in vegetative and reproductive tissues of *Cleome violacea*.

RT-PCR of full-length *Clevi-LRE/LLG1* or *Clevi-LLG2/LLG3* in cDNAs isolated from 30-day-old rosette leaves, anther and pollen, emasculated pistils without ovules, and ovules from emasculated pistils of *Cleome violacea*. A homolog of *A. thaliana ACTIN2* (*Clevi-ACTIN2*) was used as a control in these experiments. gDNA, genomic DNA isolated from of Cleome leaves was used as a positive control in PCR portion of the RT-PCR experiment. RT-PCR was repeated with two additional biological replicates with similar results and the amplified bands were sequenced to confirm the identity of amplified cDNAs.