Titration of *in-cellula* affinities of protein-protein interactions

David Cluet, Blandine Vergier, Nicolas-Pierre Levy, Lucie Dehau, Alexandre Thurman, Ikram

Amri and Martin Spichty[#]

Laboratoire de Biologie et de Modélisation de la Cellule, Ecole Normale Supérieure de Lyon,

CNRS, Université Lyon 1, Université de Lyon; 46 allée d'Italie; 69364 Lyon cedex 07; France.

#) corresponding author. New address: Laboratoire d'Innovation Moléculaire et Applications,

Université de Strasbourg – Centre National de la Recherche Scientifique – Université de Haute-

Alsace, 3 bis rue Alfred Werner, 68057 Mulhouse Cedex, France; Phone: +33 389 33 68 62;

Email: martin.spichty@uha.fr.

KEYWORDS: protein-protein interactions, quantitative yeast-two hybrid, flow cytometry,

single-cell analysis.

A genetic assay permits simultaneous quantification of two interacting proteins and their bound fraction at the single-cell level using flow cytometry. *In-cellula* **affinities of protein-protein interactions can be extracted from the acquired data through a titration-like analysis. The applicability of this approach is demonstrated on a diverse set of interactions with proteins from different families and organisms and with** *in-vitro* **dissociation constants ranging from picomolar to micromolar.**

The quest for methods that permit rapid and reliable determination of the affinity of protein-protein interactions (PPI) is unbroken. In contrast to biochemical *in-vitro* methods such as Isothermal Titration Calorimetry (ITC) and Surface Plasmon Resonance (SPR) that require purified proteins, quantitative genetic assays rely on the expression of the proteins of interest in cells. Many of these assays[1–6] are inspired by the yeast two-hybrid (Y2H) technique[7–9] which is based on the *in-cellula* expression of two proteins, usually named Bait and Prey, fused to an DNA-binding domain (BD) and an activation domain (AD), respectively. Upon physical interaction of the BD-Bait and AD-Prey proteins, a functional transcription factor is reconstituted that drives the expression of a reporter gene. The stronger the interaction, the higher should be the expression level of the reporter.[1] However, the expression level of the AD-Bait and BD-Prey play an important role, too.[10]

We recently introduced a quantitative yeast-two hybrid system (qY2H) that permits for the first time simultaneous quantification of BD-Bait, AD-Prey and the reporter at the single-cell level without the need of any antibodies or purified proteins.[10] Instead, we take advantage of fluorescent fusion proteins that can be detected by standard flow cytometers. Here we show how this qY2H method can be exploited to perform *in-cellula* affinity titrations by applying the following two important improvements:

1) Cellular contents of fluorescent proteins are determined in units of Molecules of Equivalent Soluble Fluorochrome (MESF), so that measured quantities become independent of the applied apparatus setup. It facilitates the future transferability of the qY2H measurements to other flow cytometers and allows researchers to consistently compare their results. Our reference fluorochrome is the yeast Enhanced Green Fluorescent protein (yEGFP) for which commercial calibration beads exist. The fluorescence intensity of TagRFP is converted into units of MESF of EGFP using independent calibration experiments with a fluorescent tandem protein BD-TagRFP-EGFP (see "Methods").

2) We analyze the data by a titration-like procedure which allows the straightforward extraction of *in-cellula* dissociation constants for Bait:Prey interactions. In a proof of concept, we apply this *in-cellula* titration approach to a diverse set of PPIs with dissociation constants ranging from 117 pM to 17 µM (Table 1). As in *in-vitro* SPR experiments, each PPI can be measured by Y2H in two different orientations (by exchanging Bait and Prey). Here we study only the orientation that produced the higher reporter level.[10] This orientation is considered as the molecular configuration with the higher accessibility of the PPI binding interface.[1]

In our qYH2 experiments, diploid yeast cells with constitutive expression of BD-Bait and induced expression of AD-Prey are cultured for two hours. Then, their fluorescence intensity is measured by flow cytometer in the three channels corresponding to TagRFP (BD-Bait), EGFP (AD-Prey), and TagBFP (reporter). Due to phenotypic variations, BD-Bait and AD-Prey are expressed at different levels among these cells which can be exploited to "prepare samples" for a titration. By gating, we can split the global heterogeneous ensemble of cells into several homogenous subensembles (bins). Each bin contains only cells within two specific, narrow intervals of red and green fluorescence intensity centered at values R and G, respectively. Assuming a linear relationship between fluorescence intensity and molecule

numbers, R and G can be considered as measures for the mean cellular content of BD-Bait and

AD-Prey in the corresponding bin.

With the mean value of the blue fluorescence intensity, we can calculate for each bin

the normalized reporter level φ. It is obtained by forming the ratio of the expression level for

the interaction of interest, $E_{\text{interaction}}$ (Fig. 1A) and the level for a covalent BD-AD fusion,

$E_{\text{colvalent}}$ (Fig. 1B). This normalization renders φ dimensionless and independent of the

acquisition apparatus (assuming again a linear relationship between molecule number and

fluorescence intensity). Most importantly, we consider that φ reflects the time-averaged

fraction of BD-Bait bound by AD-Prey during the reaction (as explained in the caption of Fig.

1). Thus, titration curves can be obtained when φ is plotted as a function of G while keeping R

fixed (Fig. 1C). The curves can be fitted with the following Langmuir-type equation:

85
$$\varphi(G) \cong \beta \frac{G/\alpha}{K'_{\text{d}} + R + G/\alpha}$$

Eq. 1

where K'_{d} is the *in-cellula* dissociation constant (in units of MESF of EGFP) and α and β are

dimensionless parameters that empirically account for the fact that φ is a time-integrated

property. The parameter α reduces the final cellular content of AD-Prey (measured at the end

of the reaction, G) to the time-averaged content (over the entire reaction course, $<G>$). Since

the induction kinetics under the GAL1-promotor in yeast[11] displays a quadratic-like time

dependence (for short induction times), a reasonable choice for α is 3 $[<G> = _0\int^1 G\, t^2\, dt = G/3]$.

The prefactor β, on the other hand, integrates differences in the expression kinetics of the

reporter for $E_{\text{interaction}}$ (induced expression) and $E_{\text{colvalent}}$ (constitutive expression). It can be

determined experimentally by monitoring φ for $G \rightarrow \infty$ using a high-affinity couple (such as

BD-Barstar29F/AD-BarnaseH102A).

We recommend that the titrations are carried out with the lowest possible value of

$R=R_{min}$ (as defined by the detection limit of TagRFP by flow cytometery, see "Methods"). It

limits overexpression and associated protein burden effects.[12] Furthermore, the auto-activation

potential of the BD-Bait fusion is kept at a minimum, too.[13] Most importantly, it mimics the

condition of *in-vitro* affinity titration experiments[14] where the concentration of the titrated

species (here BD-Bait) is kept fixed and as low as possible to avoid saturation effects. For the

titrations with $R=R_{min}$ the parameters $\alpha=3$ and $\beta=1.35$ were used to extract the K'_d-values.

Despite substantial differences between our *in-cellula* system and *in-vitro* setups (as

previously discussed[10] in detail), the *in-cellula* affinities strongly correlate with those from *in-*

vitro measurements ($R^2=0.91$, Fig. 1D). The slope of the regression line is 0.84. Other *in-cellula*

assays usually find lower correlation coefficients (< 0.9) and significantly lower values for the

slope of the regression line (0.2-0.6).[4,5,15–17] This is even more remarkable if one considers that

the tested set of PPIs in this work is significantly more diverse. It may indicate a higher

sensitivity for the qY2H titration approach; more testing will be necessary to confirm this

surmise.

The presented protocol is robust as witnessed by the small error bars in the titration

curves (Fig. 1c). All steps of the protocol have been optimized in liquid phase that can be easily

automated for the use of microplates and integrated within robotic pipelines. It sets the stage

for high-throughput affinity screenings of PPIs using cross-mating approaches[18,19] with libraries

of yeast clones. As an outlook, affinity-based networks[20] can be created by attributing weights

to the PPI edges according to their effective affinities. It contrasts standard Y2H screens that

yield networks with only binary information (YES or NO). The topology of edge-weighted

spring-embedded networks[21] may help identifying key pathways within the network, and how

they change as a function of environmental conditions (stress, metabolism, *etc*). Thus, we

anticipate that high-throughput qY2H affinity data would boost the modelling of interactomes

and thereby advance significantly systems biology.

Methods

The qY2H experiments, acquisitions by flow cytometry and analyses were carried out as described in our previous study[10] with the following particularities. About 10^7 cells were cultured per experiment and interaction (including the covalent BD-AD fusion and the control sample BD-Empty / AD-Empty). To ensure that these cells have been indeed transfected with all three vectors, we selected for the analysis large (=growing) cells with a forward scatter range 75 000 < FSC-H < 125 000; "H" indicates signal height. Furthermore, only cells with a red fluorescence intensity of 800 ± 100 TagRFP-H were analysed. This bin is located just above the 95% threshold of the non-fluorescent cells,[10] and therefore defines R_{min}.

The mean Tag BFP-H value was then calculated for bins of varying G values from -500 to 25500 yEGFP-H (bin size 1000). For each bin we calculated:

$$\varphi(G) = \frac{\langle F_{BFP,interaction}\rangle_G - \langle F_{BFP,CTRL}\rangle_G}{\langle F_{BFP,covalent}\rangle_G - \langle F_{BFP,CTRL}\rangle_G}$$

Eq. 2

where $<F_{BFP,X}>_G$ is the mean blue fluorescence intensity. The subscripted X refers to the physical interaction, covalent fusion or control couple. The control couple BD-Empty / AD-Empty[10] permits to remove the background of the reporter system.

Finally, G values were converted into MESF of EGFP using calibration beads (Ozyme, reference 632594) following the manufactor's protocol. For the conversion of R_{min}, we performed independent calibration measurements with diploid yeast cells expressing the fluorescent tandem fusion protein BD-TagRFP-yEGFP (under the same condition as the qY2H experiments). Cells with a red fluorescence intensity of $R_{min} = 800 ± 100$ TagRFP-H displayed a mean green fluorescence intensity of 370 000 MESF of EGFP.

Experiments and analyses were performed at least three times for each interaction and averaged titration curves were least-square fitted with Eq. 1.

148 **Acknowledgments**

149   This work was supported by the Fonds Recherche of the Ecole Normale Supéreure de Lyon.

150
151

Graphical elements

154
Figure 1: The qY2H system and its application to affinity titrations
(A) In our qY2H system, red-fluorescent BD-Bait interacts with green-fluorescent AD-Prey to reconstitute a
transcription factor that drives the expression of a blue-fluorescent reporter. Our hypothesis is that the expression
level of the reporter, $E_{interacting}$, reflects the number of BD-molecules bound to the promotor corrected by the
fraction of BD-Bait bound to AD-Prey. This fraction is influenced by the affinity between the Bait and the Prey,
but also by the expression levels of BD-Bait and AD-Prey. (B) When the activation domain is covalently linked
to the DNA-binding domain, the expression level $E_{covalent}$ depends only on the number of BD-Bait molecules bound
the promotor. Thus, when forming the quantity φ by dividing $E_{interacting}$ with $E_{covalent}$, we obtain a measure for the
fraction of BD-molecules bound by an AD-Prey molecule. To determine $E_{covalent}$, we constructed a BD-AD fusion
protein. Unfortunately, the activation domain B42 (as used in **a**) turned out to be toxic for our yeast strains when
used in the BD-AD construct. Instead, we used the activation domain B112. Difference in the activation potential
between B42 and B112 are integrated in the parameter β of Eq. 1. (C) The quantity φ can be monitored as a
function of different levels of EGFP Molecules of Equivalent Soluble Fluorochrome (MESF) corresponding to
different cellular levels of AD-Prey. In these titrations, the level of BD-Bait is kept fixed at the lowest possible
value (see "Methods"). For the interaction TEM/BLIP1 (cyan line) the titration can be preformed only up to one
third of the titrant quantity due to expression problems of AD-BLIP1.[10] Green triangles at the top vertical axis
indicate the position of used calibration beads. (D) When the titration curves are fitted with Eq. 1, we can extract
the dissociation constant in units of MESF (K_d'). The estimated K_d'-values show a remarkable correlation with
the dissociation constants measured from alternative *in-vitro* experiments (with matching symbols of Table 1).

Table 1: Investigated protein-protein interactions and their *in-vitro* affinities (K_d).

Bait proteins							Prey proteins				
Organism	Family	Name	Mutant	MW	K_d (nM)	Symbol	Organism	Family	Name	Mutant	MW
B. amylo-liquefaciens	RNAse inhibitor	Barstar	WT	10	320 [a]	■	*B. amylo-liquefaciens*	RNAse	Barnase	H102A	12
			Y29A	10	420 [a]	▲					
			Y29F	10	117 [a]	●					
			W38F	10	4 000 [a]	□					
			D35A	10	25 000 [h]	◉					
			D39A	10	420 000 [b]	▲					
H. sapiens	GTPase	HRas	G12V & C186A	21	122 000 [c]	■	*H. sapiens*	Kinase	CRaf RBD	WT	9
					11 000 [d]	▲				A85K	9
H. sapiens	Kinase regulatory subunit	CksHs1	WT	10	77 000 [e]	■	*H. sapiens*	Kinase	CDK2	WT	34
E.coli	β-Lactamase	TEM	WT	31	15 000 [f]	◉	*S. clavuligerus*	β-Lactamase inhibitor	BLIP1	WT	21
HIV1	Virulence factor	Nef	LAI	23	11 400 000 [g]	■	*H. sapiens*	Kinase	SRC SH3	WT	7
H. sapiens	Adapter	Grb2 SH3	WT	7	17 000 000 [i]	◆	*M. musculus*	Nucleotide exchange factor	Vav1 SH3	WT	8

[a] ITC, 50mM Tris/HCl, pH 8 at 25°C.[22]
[b] Mean values from two studies[22,23] with ITC, 24mM Hepes, pH 8, 1 mM DTT at 25°C.
[c] Mean values from four studies of Ras G12V (without the membrane anchor): SPR, 50 mM Tris/HCl, pH 7.4, 100mM NaCl, 5 mM MgCl$_2$;[24] SPR, 50 mM Tris/HCl, pH 7.4, 100mM NaCl, 5 mM MgCl$_2$;[25] SPR, 10 mM Hepes, pH 7.4, 150 mM NaCl, 2 mM MgCl$_2$, and 0.01% Nonidet P-40 25°C;[26] ITC, 50 mM Hepes, pH 7.4, 125 mM NaCl, 5 mM MgCl$_2$,25°C.[27]
[d] ITC, 50 mM Hepes, pH 7.4, 125 mM NaCl, 5 mM MgCl$_2$,25°C.[27] The dissociation constant of the CRaf RBD A85K mutant was measured with HRas WT loaded with a GTP-analogue. The mutant HRas G12V is known to decrease the dissociation constant for the interaction with CRaf RBD WT by a factor of 11.[28] The given value applies the same correction factor.
[e] SPR, 10 mM Hepes, 3.4 mM EDTA, 150 mM NaCl, 0.001% surfactant P20, pH 7.4.[29]
[f] SPR, 10 mM Hepes, 3.4 mM EDTA, 150 mM NaCl, 0.05% surfactant P20, pH 7.4.[30]
[g] ITC, 20 mM phosphate buffer, pH 7.5, 150 mM NaCl, 2 mM EGTA, and 5 mM DTT, 25°C.[31]
[h] SPR, 10 mM Hepes-Na, pH 7.4, 0.15 M NaCl, 3 mM EDTA, and 0.005% (v/v) Tween 20, 25 °C.[32]
[i] SPR, 25°C.[33]
[h] Free-energy calculations.[10]

References

1. Estojak, J., Brent, R. & Golemis, E. A. Correlation of two-hybrid affinity data with in vitro measurements. *Mol. Cell. Biol.* **15**, 5820–5829 (1995).

2. Endoh, H., Walhout, A. J. & Vidal, M. A green fluorescent protein-based reverse two-hybrid system: application to the characterization of large numbers of potential protein-protein interactions. *Methods Enzymol.* **328**, 74–88 (2000).

3. Möckli, N. & Auerbach, D. Quantitative β-galactosidase assay suitable for high-throughput applications in the yeast two-hybrid system. *BioTechniques* **36**, 872–876 (2004).

4. Jeong, K. J., Seo, M. J., Iverson, B. L. & Georgiou, G. APEx 2-hybrid, a quantitative protein-protein interaction assay for antibody discovery and engineering. *Proc. Natl. Acad. Sci.* **104**, 8247–8252 (2007).

5. Dutta, S., Koide, A. & Koide, S. High-throughput Analysis of the Protein Sequence–Stability Landscape using a Quantitative Yeast Surface Two-hybrid System and Fragment Reconstitution. *J. Mol. Biol.* **382**, 721–733 (2008).

6. Zolghadr, K. *et al.* A fluorescent two-hybrid assay for direct visualization of protein interactions in living cells. *Mol. Cell. Proteomics MCP* **7**, 2279–2287 (2008).

7. Fields, S. & Song, O. A novel genetic system to detect protein protein interactions. *Nature* **340**, 245–246 (1989).

8. Gyuris, J., Golemis, E., Chertkov, H. & Brent, R. Cdi1, a human G1 and S phase protein phosphatase that associates with Cdk2. *Cell* **75**, 791–803 (1993).

9. Stynen, B., Tournu, H., Tavernier, J. & Van Dijck, P. Diversity in Genetic In Vivo Methods for Protein-Protein Interaction Studies: from the Yeast Two-Hybrid System to the Mammalian Split-Luciferase System. *Microbiol. Mol. Biol. Rev.* **76**, 331–382 (2012).

10. Cluet, D. *et al.* A quantitative tri-fluorescent yeast two-hybrid system: from flow

cytometry to in-cellula affinities. *Mol. Cell. Proteomics* (2020)

doi:10.1074/mcp.TIR119.001692.

11. Li, J. *et al.* Green fluorescent protein in Saccharomyces cerevisiae: Real-time studies of

the GAL1 promoter. *Biotechnol. Bioeng.* **70**, 187–196 (2000).

12. Bolognesi, B. & Lehner, B. Reaching the limit. *eLife* **7**, e39804 (2018).

13. Rajagopala, S. V. & Uetz, P. Analysis of Protein–Protein Interactions Using High-

Throughput Yeast Two-Hybrid Screens. in *Network Biology: Methods and Applications*

(eds. Cagney, G. & Emili, A.) 1–29 (Humana Press, 2011).

14. Pollard, T. D. A Guide to Simple and Informative Binding Assays. *Mol. Biol. Cell* **21**,

4061–4067 (2010).

15. Hu, X., Kang, S., Chen, X., Shoemaker, C. B. & Jin, M. M. Yeast Surface Two-hybrid for

Quantitative *in Vivo* Detection of Protein-Protein Interactions via the Secretory Pathway.

J. Biol. Chem. **284**, 16369–16376 (2009).

16. Younger, D., Berger, S., Baker, D. & Klavins, E. High-throughput characterization of

protein–protein interactions by reprogramming yeast mating. *Proc. Natl. Acad. Sci.* **114**,

12166–12171 (2017).

17. Colas, P., Cohen, B., Ferrigno, P. K., Silver, P. A. & Brent, R. Targeted modification and

transportation of cellular proteins. *Proc. Natl. Acad. Sci.* **97**, 13720–13725 (2000).

18. Kolonin, M. G., Zhong, J. & Finley, R. L. Interaction mating methods in two-hybrid

systems. *Methods Enzymol.* **328**, 26–46 (2000).

19. Chen, J., Carter, M. B., Edwards, B. S., Cai, H. & Sklar, L. A. High throughput flow

cytometry based yeast two-hybrid array approach for large-scale analysis of protein-

protein interactions. *Cytometry A* **81A**, 90–98 (2012).

20. Yugandhar, K. & Gromiha, M. M. Analysis of protein-protein interaction networks based
on binding affinity. *Curr. Protein Pept. Sci.* **17**, 72–81 (2016).

21. Kamada, T. & Kawai, S. An algorithm for drawing general undirected graphs. *Inf.*
Process. Lett. **31**, 7–15 (1989).

22. Schreiber, G. & Fersht, A. R. Energetics of protein-protein interactions: Analysis ofthe
Barnase-Barstar interface by single mutations and double mutant cycles. *J. Mol. Biol.*
248, 478–486 (1995).

23. Frisch, C., Schreiber, G., Johnson, C. M. & Fersht, A. R. Thermodynamics of the
interaction of barnase and barstar: changes in free energy versus changes in enthalpy on
mutation 1 1Edited by J. Karn. *J. Mol. Biol.* **267**, 696–706 (1997).

24. Herrmann, C., Horn, G., Spaargaren, M. & Wittinghofer, A. Differential Interaction of the
Ras Family GTP-binding Proteins H-Ras, Rap1A, and R-Ras with the Putative Effector
Molecules Raf Kinase and Ral-Guanine Nucleotide Exchange Factor. *J. Biol. Chem.* **271**,
6794–6800 (1996).

25. Block, C., Janknecht, R., Herrmann, C., Nassar, N. & Wittinghofer, A. Quantitative
structure-activity analysis correlating Ras/Raf interaction in vitro to Raf activation in
vivo. *Nat. Struct. Mol. Biol.* **3**, 244–251 (1996).

26. Fischer, A. *et al.* B- and C-RAF display essential differences in their binding to Ras: the
isotype-specific N terminus of B-RAF facilitates Ras binding. *J. Biol. Chem.* **282**, 26503–
26516 (2007).

27. Kiel, C. *et al.* Improved Binding of Raf to Ras·GDP Is Correlated with Biological
Activity. *J. Biol. Chem.* **284**, 31893–31902 (2009).

28. Kiel, C. Untersuchung von Ras/Effektor-Komplexen mit gezielt veränderten
elektrostatischen Eigenschaftens. (Dissertation, Ruhr-Universität Bochum, Fachbereich
Biochemie, 2003).

29. Bourne, Y. *et al.* Crystal Structure and Mutational Analysis of the Human CDK2 Kinase

Complex with Cell Cycle–Regulatory Protein CksHs1. *Cell* **84**, 863–874 (1996).

30. Albeck, S. & Schreiber, G. Biophysical Characterization of the Interaction of the β-

Lactamase TEM-1 with Its Protein Inhibitor BLIP [†]. *Biochemistry (Mosc.)* **38**, 11–21

(1999).

31. Arold, S. *et al.* RT Loop Flexibility Enhances the Specificity of Src Family SH3 Domains

for HIV-1 Nef [†,‡]. *Biochemistry (Mosc.)* **37**, 14683–14691 (1998).

32. Sato, Y. *et al.* Characterization of the Interaction between Recombinant Human Peroxin

Pex3p and Pex19p: *IDENTIFICATION OF TRP-104 IN Pex3p AS A CRITICAL*

RESIDUE FOR THE INTERACTION. *J. Biol. Chem.* **283**, 6136–6144 (2008).

33. Nishida, M. Novel recognition mode between Vav and Grb2 SH3 domains. *EMBO J.* **20**,

2995–3007 (2001).