***Toxoplasma* FER1 is a versatile and dynamic mediator of differential**

microneme trafficking and microneme exocytosis

Daniel N.A. Tagoe[1§†], Adeline Ribeiro E Silva[1§], Allison A. Drozda[1†], Isabelle Coppens[2],

Bradley I. Coleman[1†], Marc-Jan Gubbels[1*]

[1]Department of Biology, Boston College, Chestnut Hill, MA 02467, USA

[2]Department of Molecular Microbiology and Immunology, Johns Hopkins University Bloomberg

School of Public Health, Baltimore, MD 21205, USA

[*]Correspondence to Marc-Jan Gubbels, gubbelsj@bc.edu

[§]Daniel N.A. Tagoe and Adeline Ribeiro E Silva contributed equally to this work

[†]Present addresses: Daniel N.A. Tagoe, CANbridge Pharmaceuticals Inc., Burlington, MA;

Allison A. Drozda, Mosaic Biosciences, Boulder, CO; Bradley. I Coleman, Flagship Pioneering,

Cambridge, MA

Running title: *Toxoplasma* FER1 acts on micronemes
21
22
23
24

Abstract

Toxoplasma gondii is a polarized cell concentrating several secretory organelles at the apical pole. The secretory micronemes come in two sub-populations differentiated by dependence on the Rab5A/C in their biogenesis. Calcium-dependent exocytosis of micronemes occurs at the very apical tip and is critical for parasite egress from its host cell, adhesion and invasion of the next cell. Ferlins represent a protein family with roles in exocytosis containing multiple Ca^{2+}-sensing C2 domains. We determined that *T. gondii*'s ferlin 1 (FER1) localized dynamically to the parasite's secretory pathway. FER1 function was dissected by dominant negative overexpression strategies. We demonstrated that FER1 traffics microneme organelles along the following trajectories:1. From the *trans*-Golgi-endosomes network to the subpellicular cortex; 2. Along the cortex to the apical end; 3. To the apical tip for fusion with the plasma membrane; 4. Retrograde transport allowing microneme recycling from mother to daughter; 5. Differential microneme sub-population traffic. Finally, FER1 overexpression triggers a microneme exocytosis burst, supporting the notion that the radially organized micronemes at the apical tip comprise a readily-releasable microneme pool. In summary, FER1 is pivotal for dynamic microneme trafficking, acts differently on the two microneme subpopulations, and acts on the plasma membrane fusion step during microneme exocytosis.

Keywords: exocytosis, ferlin, microneme, trafficking, *Toxoplasma*

Introduction

In humans, the apicomplexan parasite *Toxoplasma gondii* causes birth defects, vision loss, myocarditis and encephalitis. Lytic replication cycles unfolding through repetitive rounds of host cell invasion, intracellular replication and host cell egress are central to the pathogenesis of toxoplasmosis[1]. The micronemes are pivotal for successful host cell invasion as they contain adhesion molecules facilitating gliding motility and host cell attachment[2,3]. In addition, the *Toxoplasma gondii* micronemes encode a pore-forming protein, PLP1, that permeabilizes both the parasitophorous vacuole and host plasma membrane, which is required for host cell egress[4,5]. The parasite's signal transduction pathways controlling the correct timing of micronemes secretion comprises cGMP, Ca^{2+} and phosphatidic acid (PA)[6], accompanied by a crucial cAMP-mediated switch between the intracellular and extracellular states[7].

The micronemes are localized along the apical cortex in association with the subpellicular microtubules emanating from the apical end[8,9]. Upon activation of exocytosis, micronemes move into the conoid, a tubulin basket at the apical tip of the parasites, to fuse with the plasma membrane. Following secretion, microneme protein complexes are embedded in the plasma membrane with the extracellular domains serving as adhesion domains and the cytoplasmic tail engaging with actin filaments that are transported in an apical to basal direction by a myosin motor[10]. A set of radial micronemes organized and anchored just below the conoid is believed to be a readily-releasable pool of micronemes[11-13]. Furthermore, microneme exocytosis is dosed to support prolonged periods of gliding motility enabling migration cross the biological barriers in between host cells[14] as only ~20% of all microneme proteins can be triggered to secrete at any given time[15,16]. Dosing is most likely achieved by gradually trafficking micronemes toward the conoid[15]. Finally, microneme exocytosis is balanced with active endocytosis[17], whereas during cell division micronemes of the mother are re-directed into the forming daughters[18].

Microneme biogenesis required protein trafficking through the secretory pathway comprising sequential passage through the endoplasmic reticulum (ER), Golgi apparatus, trans-Golgi network (TGN) and an endosome-like compartment (ELC)[19-21]. Microneme proteins undergo proteolytic processing to remove their pro-peptide inside the plant like vacuole (PLV or VAC), an acidic compartment[22-25]. Trafficking proteins from the ELC to micronemes and other destinations is facilitated by an unconventional endosomal system[20,21]. Protein sorting to both microneme and rhoptry secretory organelles requires sortilin (SORTLR) in the *trans*-Golgi[26],

trafficking to the rhoptries and dense granules REMIND[27], whereas the HOPS/CORVET

complex and Rab7 are involved in PLV/VAC routing[28]. Moreover, adaptor complex AP1 is

involved in microneme and rhoptry protein trafficking but has a more general function across

other vesicular trafficking events[29]. Although some rhoptry specific targeting signals have been

identified[30,31], specific sorting signals for microneme proteins are still elusive. Specific Rab

GTPases have been associated with some aspects of microneme protein trafficking, and

differentiate two sub-populations of micronemes with a different protein content: one of which is

Rab5A/C-dependent, and one that is Rab5A/C-independent[9]. Consequently, specific microneme

proteins end up in different, non-overlapping microneme sub-populations. To date, no

differential function or behavior for these distinct microneme populations have been described,

leaving the rationale for this distinction unresolved.

In all well-studied Ca^{2+} triggered exocytosis systems, proteins containing double C2

(DOC2) domains execute the Ca^{2+}-mediated vesicle fusion[32]. The Apicomplexa encode an

unconventional, long DOC2 domain protein essential for microneme exocytosis[33]. In addition,

two ferlin proteins, FER1 and FER2, are conserved across the Apicomplexa[34,35]. FER1 in

Plasmodium berghei, named ferlin-like protein (PbFLP), is reported to be essential for male

gametocyte egress[36], whereas *T. gondii* FER2 is required for rhoptry secretion[34]. The ferlins

make up a unique branch of the DOC2 domain protein family because of their size (200-240

kDa) and contain five to seven C2 domains, organized in C2 pairs to form 2-3 DOC2 domains.

The extended C2 repertoire in ferlins has broadened their functional spectrum beyond membrane

fusion to vesicle trafficking and membrane repair[37]. Mammalian ferlins are differentiated by

their sub-cellular localization at either the plasma membrane or on intracellular compartments,

which relates to their function in late endosomal transit versus trans-Golgi recycling[38].

Mammalian otoferlin, essential for neurotransmitter release from the inner hair cells (IHC) in the

auditory system, has been most widely studied. Otoferlin functions as both a scaffolding protein

in the secretory pathway as well as in the actual membrane fusion during exocytosis[39-42]. Here

we examined *T. gondii* FER1 and reveal roles in microneme protein trafficking, microneme

organelle dynamics and exocytosis. Furthermore, FER1 acts differentially on the two different

microneme populations.

Results

1. *Toxoplasma* FER1 localizes to the secretory pathway

TgFER1 (TGGT1_309420) is a protein of 1425 amino acids harboring six C2 domains organized

in three pairs and a tail-anchored transmembrane domain (**Fig. 1a**). A FerI domain found in most

ferlins is present between the C2A and C2B domains, however no function has been associated

with this domain[43]. In the genome wide fitness screen for the *Toxoplasma* lytic cycle, FER1 has a

severe fitness score of -4.77, indicating that this protein is essential[44]. The HyperLOPIT sub-

cellular localization atlas did not contain an assignment for TgFER1[45]. Hence, to localize FER1

we generated a polyclonal antiserum against the central C2DE domain, which is the most

evolutionary diverse sequence of the protein compared to the other ferlins (**Fig. 1a**). By western

blot, the α-FER1 serum reacts with the full-length FER1 protein with a predicted molecular

weight of 159 kDa (**Fig. 1b**). Additional bands at approximately 120 and 30 kDa were detected,

which most likely represent fragments of the full length FER1 protein, although cross reactivity

with other proteins cannot be excluded. By IFA, since we anticipated a function in microneme

exocytosis, we co-stained FER1 in both intracellular (**Fig. 1c**) and extracellular (**Fig. 1d**)

parasites with MIC2 antiserum to highlight the micronemes. FER1 and MIC2 were physically

distinct in intracellular parasites with FER1 localized apical of the nucleus. In extracellular

parasites, FER1 localized throughout the cytoplasm, suggesting a re-localization in response to

the environment. This coincides with the activation of microneme exocytosis and gliding

motility[46]. The absence of FER1 co-localization with the micronemes suggests FER1 behaves

like the mammalian ferlins in *trans*-Golgi recycling[38].

To further define the sub-cellular compartment to which FER1 localizes in intracellular

parasites we employed a series of *T. gondii* secretory pathway markers (**Fig. 1e-g**). We evaluated

FER1 overlap with these markers using the Pearson correlation coefficient (PCC; ranging from 1

for perfect positive to -1 for perfect negative co-localization). Most of the FER1 signal overlaps

with a marker for the *trans*-Golgi apparatus (GalNAc[29]) and the TGN-endosomal compartment

(dynamin related protein B: DrpB[47]). Surprisingly, no colocalization is observed with Sortilin,

which is another marker for the *trans*-Golgi[26,47-51]. IFAs showing the relative distribution of the

different markers in wild-type parasites are provided in **Supplementary Figure 1**. Although

depletion of Sortilin does affect both microneme and rhoptry biogenesis[26], it has a much stronger

connection to sorting proteins to the rhoptries[19,29]. Thus, FER1 and sortilin could occupy

different domains in the *trans*-Golgi, with the former catering to microneme and the latter to the

rhoptry proteins. Taken together, in intracellular parasites FER1 localizes to the *trans*-Golgi-

endosome network, whereas upon egress the signal disperses into the cytoplasm, supporting a

role in the endosome-related trafficking pathway of microneme proteins.

2.   Generation and validation of conditionally lethal FER1 mutant lines

To dissect the function of FER1 we attempted conditional depletion approaches by either placing

the gene under a tetracycline regulatable promoter or fusing the auxin-inducible degron at either

the N- or C-terminus. None of these efforts were successful, and neither were efforts to insert

tags within the open reading frame, indicating that FER1 is extremely sensitive to perturbation.

As an alternative, we resorted to a dominant negative (DN) approach. We generated an α-tubulin

promoter driven N-terminal fusion of the destabilization domain (DD) linked to a Myc epitope

that can be conditionally stabilized with Shield-1[52]. We established constructs without the 21 C-

terminal amino acids encoding the TM domain (DD-Myc-FER1ΔTM) as well as a control

construct encoding the full-length FER1 protein (DD-Myc-FER1[FL]) (**Fig. 2a**). We were able to

generate stable parasite lines with both constructs in absence of Shield-1. Correct expression of

the proteins was confirmed by western blot (**Fig. 2b**). We showed by plaque assay that

overexpression of both the ΔTM and FL alleles caused parasite lethality (**Fig. 2c**). Collectively,

these data align with a critical role of FER1 in the lytic cycle.

3.   FER1ΔTM overexpression causes a microneme exocytosis defect

To determine the cause of the lethal defect upon FER1ΔTM overexpression, we evaluated key

events in the lytic cycle. Host cell invasion was strongly reduced, consistent with an anticipated

FER1 role in microneme exocytosis (**Fig. 3a**). We observed a mild reduction in the parasite

multiplication rate, with a slight but significant (p=0.031) accumulation of 4-parasites/vacuole

(**Fig. 3b**). However, no specific delay was observed in a particular cell cycle stage (**Fig. 3c**),

suggesting that the overall rate through cell division is somewhat reduced. We triggered

microneme-exocytosis mediated egress with Ca^{2+}-ionophore A23187 and assessed the integrity

of the parasitophorous vacuole membrane (PVM) using GRA3 as marker together with IMC3

marking the parasite's cytoskeleton (**Fig. 3d**). We observed a 90% egress reduction in the

overexpression condition, which was confirmed using pharmacological triggers in different

points in the signaling pathway (**Suppl Fig. S2a,b**). Together, this convincingly supports a

microneme defect (**Fig. 3e**). Trail assays were performed to further functionally assess

microneme exocytosis and we determined that gliding motility is abrogated (**Fig. 3f**). Finally, we

determined differential exocytosis of the Rab5A/C-dependent (MIC3, 5, 8) and -independent

(MIC2, M2AP,etc.) microneme sub-populations[9]. The latter sub-population was tested by release

of processed MIC2 in the medium under various triggers, whereas the former was visualized by

MIC3, 5 and 8 protein exposure on the parasite's surface. Overall, we see a sharp reduction in

exocytosis to nearly undetectable levels of both microneme populations in the FER1ΔTM

overexpressing mutant (**Fig. 3g,h**; **Suppl Fig. S2c**). However, we detected propranolol induced

exocytosis at a 10-fold lower level than the non-induced control (**Fig. 3g**). Additionally, a small

amount of MIC3 on the surface of induced parasites was observed, indicating a minimal

exocytosis capacity (**Fig. 3h**). Since propranolol triggers the Ca^{2+}-independent PA pathway, this

section of the pathway appears to be still functional. Taken together, this suggests that FER1 acts

in the Ca^{2+}-dependent events in the microneme protein secretory pathway.

4.   FER1ΔTM overexpression causes a microneme trafficking defect

By design, we anticipated that FER1ΔTM would mislocalize. IFA in the DD-Myc-FER1ΔTM

line revealed a reproducible signal accumulation in a defined mid-apical region upon Shield-1

induction (**Fig. 4a**). In parallel, we generated a YFP-tagged version of the FER1ΔTM allele to

directly track accumulation. We observe no YFP signal without Shield-1 whereas under Shield-1

YFP accumulates in the mid-apical region as seen with the Myc construct, displaying a PCC of

0.968 co-localization with MIC2 (**Fig. 4b**; **Suppl Fig. S3a**). Co-staining with Rab5A/C-

dependent (MIC5) class showed perfect overlap with the accumulated YFP signal indicating that

all microneme proteins are misdirected (**Fig. 4c**; **Suppl Fig. S3b**). The α-FER1 signal also co-

localizes with the YFP signal, suggesting that endogenous FER1 is also diverted from its *trans*-

Golgi-endosome localization (**Fig. 4d**). Collectively, these data indicate that both the Rab5A/C-

dependent and -independent microneme sub-populations are controlled by FER1.

207   Since all known micronemes trafficking mutants present defects in rhoptry protein

trafficking[23], we interrogated the rhoptries by IFA. ROP proteins remained localized to rhoptries,

displaying their normal morphology and distribution (**Fig. 4e**). This distinction makes the DN-

FER1ΔTM unique by disrupting microneme trafficking while leaving the rhoptries intact.

The mid-apical localization is reminiscent of the position of various compartments of the

Toxoplasma secretory pathway. To differentiate whether the accumulation is due to a protein

trafficking problem, or is of another nature, we again used a series of compartment-specific

markers (**Fig. 4f, g**). No overlap was detected between the PLV/VAC (VP1) and FER1 (YFP)

signals (**Fig. 4f**, **Suppl Fig. S3c**). Furthermore, no co-localization was observed with the Golgi

or endosomal compartments. All markers (GRASP55, DrpB, Rab5A, Rab7) displayed their

normal morphology (**Fig. 4g**). Instead, DN-FER1ΔTM accumulated in an uncharacterized

compartment, beyond known trafficking steps toward the micronemes and, after splitting from

rhoptry protein trafficking[23].

To reveal the sites of FER1ΔTM and micronemes accumulation, we performed a two-

step ultrastructure expansion microscopy (pan-ExM). We stained the micronemes with M2AP

antiserum and as a guide co-stained general protein density with the NHS-ester. In parent

parasites, micronemes are observed in both channels as protein-dense small, elongated foci

aligned along the parasite's cortex and a set radially arranged at the apical pole (**Fig. 4h**). In

mutant parasites, both channels reveal mature micronemes mis-localized in the mid-apical

region. Transmission electron microscopy (TEM: **Fig. 4i**) confirmed that micronemes with

normal ultrastructure are absent from the cortex but aggregate in the parasite's apical cytoplasm.

Immune-electron microscopy (IEM) using MIC2 antiserum firmed up this assessment. Gold

particles positioned MIC2 in the aggregated microneme structures (**Suppl Fig. S2d**). These could

either be the result of retrograde trafficked mature micronemes released from the cortex, or, an

incomplete organelle biogenesis and/or recycling of micronemes from mother to daughters

during cell division.

In summary, overexpression of DN-FER1ΔTM leads to mis-localization of fully formed

micronemes without disrupting the rhoptries, indicating that FER1 functions in the endosome-

related trafficking and/or biogenesis of micronemes beyond the TGN[19].

5. DD-YFP-FER1ΔTM overexpression retracts micronemes from the cortex

To differentiate between the retrograde trafficking from the biogenesis scenarios, we asked

whether the aggregated microneme are proteolytically processed by removal of the pro-peptide

in the ELC/PLV compartments[22,25]. For that, specific antisera against the pro-peptide of M2AP

and the mature M2AP protein were used[53]. No difference in relative abundance of the pro-M2AP

versus the total amount of M2AP protein was observed (**Fig. 5a**). Furthermore, by IFA we

observed normal MIC5 pro-peptide containing proteins in the ELC compartment[53], and no co-

localization with the FER1 accumulation (**Fig. 5b**). These observations indicate that the mis-

localized micronemes contain fully mature proteins.

Since pro-MIC5 manifests predominantly in budding parasites, the co-staining with

FER1 revealed another intriguing phenomenon: no DD-YFP-FER1ΔTM accumulates in newly

forming parasites. We validated this by triple staining with YFP, α-pro-MIC5 and α-IMC3 (**Fig.**

5c). Moreover, representative images demonstrate that YFP migrates basally toward the residual

body in parasites undergoing division (**Fig. 5d**). We never observed any new micronemes in the

apical periphery of daughter buds either (**Fig. 5e**). Collectively, we observed that mature

micronemes progress normally in forming daughters, yet new micronemes do not assemble in the

daughters. This suggest that microneme proteins end up in the YFP-FER1ΔTM pile up formed in

the mother and cannot be recycled to the daughter cells.

This raised the next intriguing question: is microneme aggregation independent on cell

division? This model implies that micronemes can detach from the cortex and be directed toward

the FER1ΔTM microneme aggregate. To this end we performed time courses to determine

induction and reversion kinetics of the Shield-1 induced phenotype in both intracellular and

extracellular parasites. Microneme mis-localization kinetics were over 95% complete within two

hours (**Fig. 5f** top panel), which is much shorter than the 6.5 hr cell division cycle, indicating that

the process is cell cycle independent[54]. The same kinetics were also observed in extracellular

parasites which are arrested in a non-division G1a/0 state (**Fig. 5f** bottom panel)[55]. Finally, time

lapse microscopy allowed us to visualize the cell division independent re-localization of the

micronemes following Shield-1 induction. In DD-YFP-FER1ΔTM co-expressing a MIC3-

mCherryRFP marker we observed that microneme retraction and accumulation with YFP-

FER1ΔTM started after 1.5 hr of Shield-1 induction (**Supplementary Movie S1** and **Fig. 5g**).

Moreover, we confirmed that microneme retraction was independent of daughter budding as

tracked with a marker enriched in daughter cytoskeletons, IMC3-mCherryRFP (**Supplementary**

Movie S2 and **Suppl Fig. S3**).

Taken together, our results indicate that intact organelle microneme trafficking is cell

cycle independent and bidirectional: either in apical direction to the site of exocytosis or, from

the peripheral cortex to the site of aggregation that could be a relevant natural process when
micronemes are recycled from mother to daughters[18].

6. FER1^{FL} overexpression results in premature, untriggered microneme exocytosis
We dissected the phenotype associated to DD-Myc-FER1^{FL}, by assessing the localization of
FER1, which upon Shield-1 induction presents on membrane structures throughout the parasite.
(**Fig. 6a**). We asked whether these parasites display changes in their microneme exocytosis
ability using the same set of secretagogues applied previously (**Fig. 3g**). We observe strong
exocytosis under all conditions (**Fig. 6b**). Although slight variations were observed in the
sensitivity to different secretagogues (**Fig. 3g; Suppl Fig. S1c**), they were not significant.
Next, microneme localization and morphology was assessed by IFA, starting with the
Rab5A/C-dependent sub-population (MIC5, MIC8). Non-induced parasites display the typical
apical microneme distribution. Upon FER1^{FL} overexpression, a modest redistribution of this sub-
population to a prominent basal end appeared, while the cortical apical signal became more
diffuse (**Fig. 6c**). Staining for the Rab5A/C-independent population (MIC2, M2AP) revealed a
dramatically different change in localization pattern. By pan-ExM, the microneme signal was
strongly concentrated at the apical end in induced intracellular parasites (**Fig. 6d**; **Fig. 4h**) and
correspond to the radial micronemes[12]. Radial micronemes are anchored firmly in the apical
region and believed to be primed for exocytosis[11]. To obtain a higher resolution we resorted to
both IEM and TEM. Here we observe an accumulation of micronemes at the apical end,
confirming the apical MIC signals seen by light microscopy. The densely packed micronemes in
the apical region just below the conoid often displayed an elongated morphology (**Fig. 6e**),
whereas several micronemes were squeezed inside the conoid (**Suppl Fig. S4**). Further below,
we do observe micronemes not stained with MIC2 (**Fig. 6e**), which likely represent the
Rab5A/C-dependent sub-population that become scattered as observed by IFA (**Fig. 6c**). Thus, it
appears that overexpression of FER1^{FL} has a differential effect on the two microneme sub-
populations: the MIC2 population is driven to the apical end, whereas the MIC8 population
becomes scattered and accumulates basally. Furthermore, much fewer MIC2 micronemes were
observed. We tested relative MIC2 abundance by western blot (**Fig. 6f, Suppl Fig. S5**) in the
parental, FER1^{FL} and FER1ΔTM strains and found that FER1^{FL} parasites displayed a significant
decrease of MIC2, with negligible changes upon Shield-1 treatment. No changes were observed

in FER1ΔTM parasites. These data suggest a role for FER1 in directing microneme sub-

populations to different locations, which is a phenomenon not observed before.

To further dissect these phenotypes we tested specific functional capacities. We showed

that parasite egress could be triggered by Shield-1 addition in FER1[FL] overexpressing parasites

(**Fig. 6g**). This implies microneme exocytosis as PLP1 released through the micronemes

facilitates egress[4]. To determine if FER1 overexpression can trigger microneme exocytosis

independently of other signals, we tested if egress could happen in small vacuoles which lacked

an acidified PVM[5] or accumulated secreted diacylglycerol kinase 2 (DGK2) inside the PV[56],

which are both required for egress[6]. To this end we modified the standard red-green invasion

assay into an invasion+egress assay exploiting the same differential staining of intracellular vs

extracellular parasites[57]. Following 1 hr standard invasion, non-invaded parasites were washed

away, and Shield-1 was added for two hrs (**Fig. 6h** top). Nearly 50% of vacuoles with single

parasites have egressed (**Fig 6h.** bottom), which largely mimics the 60% egress rate seen after 2

hrs of induction from large vacuoles. Collectively, these data strongly support that Shield-1

induced microneme exocytosis is only due to FER1 overexpression and not due to activation of

the signaling pathways leading to egress. This implies that the most apically located radial

micronemes act as a readily-releasable vesicle pool.

7. Putative Ca^{2+}-binding residues are critical for FER1 function

To probe the mechanism of FER1 beyond the TM domain we analyzed the individual C2

domains for their functional potential. C2 domains fold into β-sheets connected by three loops

that can bind to proteins or insert in membranes, which can be modulated upon binding Ca^{2+}[32,58-

60]. Five key residues in the loops, of which the three central amino acids carry most weight[61], can

predict an association with phospholipids or Ca^{2+}. Across FER1's C2 domains, we detected Ca^{2+}-

binding potential only in the C2D that carries three Asp residues and a supportive Glu in the

conserved loop positions to accommodate two Ca^{2+} ions (**Fig. 7a,b**). To perturb the Ca^{2+} binding

capacity we mutated all three C2D conserved Asp residues to Ala and conditionally

overexpressed this 3DA sequence in the FER1[FL] and FER1ΔTM DD-YFP constructs (**Fig. 7c,d**).

We observed dramatic difference in MIC2 and MIC8 localization across the various mutant

alleles, to which end we quantified where in the parasite the signals are distributed (**Fig. 7E-H**).

Comparing the MIC2 signal in the overexpressed wild type (wt-3D) with the overexpressed 3DA

allele in the FER1^{FL} construct we observed a modest less pronounced apical tip accumulation

(**Fig. 7E**). In the ΔTM context, however, the effect of the 3DA mutation is a sharp reduction in

overall MIC2 signal that is still somewhat accumulated in the mid-apical region (**Fig. 7G**). For

MIC8 in the FER1^{FL} construct, we observe a mild increase in mid apical signal and a depletion

of the basal signal in the 3DA allele relative to the wt-3D allele, suggesting MIC8 trafficking

into the apical direction (**Fig. 7F**). In the ΔTM construct, we observed a much more variable

phenotype for MIC8 in the 3DA allele, which still generally looks like the mid apical

accumulation (**Fig. 7H**).

Overall, in FER1^{FL} overexpression the 3DA allele reduces the strong apical shift of MIC2

to the radial micronemes, suggesting that replenishment of the radial micronemes is mediated by

the Ca^{2+}-binding capacity of FER1. On the other hand, the MIC8 micronemes trend in the other

direction away from the very apical end. In the FER1ΔTM overexpressing parasites, MIC2

expression appears to be dependent on Ca^{2+}-binding capacity. MIC8 dispersal becomes even

more pleomorphic in the 3DA allele and FER1ΔTM context than the wt-3D allele. Thus, the

Ca^{2+}-binding capacity in the C2D domain mediates apical migration of MIC2 micronemes for

putative replenishment and, might regulate MIC2 recycling during division. The MIC8

micronemes, if anything, move in the opposite direction and become dispersed upon disrupting

Ca^{2+}-binding capacity in the FER1ΔTM context. Collectively, the function of the C2D and TM

domains in FER1 confer differential roles on the dynamics of the two microneme populations,

both in microneme position and abundance resulting in a very versatile modulating role for FER1

in microneme biology.

Discussion

The use of a series of FER1 overexpressing DN alleles revealed its multi-faceted role throughout

the lytic cycle (**Fig. 8**).

The kinetics of microneme re-direction upon DN-FER1ΔTM overexpression established

that matured micronemes are almost instantly released from the cortex, supposedly the

subpellicular microtubules, and traffic to a cytoplasmic location. Although similar phenomena

were reported upon the disruption of the vacuolar-proton ATPase (v-ATPases) in *Toxoplasma*,

they contained immature microneme proteins[25] unlike what we observed. Also, actin dependent

trafficking of micronemes from mother to daughters has been reported[18], which further supports

the feasibility and a biological function for the retrograde microneme organelle transport we

observe. Moreover, the effect we see is specific for the micronemes, unlike most other mutants in

the endosomal legs of microneme trafficking, which invariable also affect rhoptry protein

trafficking[26,29]. Overall, FER1 provides the first mechanistic insight in the microneme trafficking

beyond the upstream leg shared with the rhoptries.

Overexpression of FER1^{FL} supports a direct role in membrane fusion in the actual

exocytosis. The first piece of evidence is that Shield-1 induction leads to fast parasite egress

(**Fig. 6g,h**). In parallel to other secretory systems[62,63], we believe that only the primed or so-

called readily-releasable microneme pool is released in this mutant, but that additional secretion

requires renewed priming (e.g. Ca^{2+}-dependent events such as potential phosphorylation of

FER1[64]). The radial micronemes, accumulated right below the conoid, make a good candidate for

this readily-releasable microneme pool[12,13]. Radial micronemes are tightly anchored as they were

the only set of micronemes remaining upon VPS9 knock-down[11]. However, we cannot exclude

that only one or two micronemes are pre-docked and engaged to complete fusion with the plasma

membrane.

MIC2 microneme population squeezes into the apical end of the parasite, indicating

anterograde transport along the cortex was triggered as well. A similar microneme apical

migration was observed upon knock out of AP1, although that was not unique to the micronemes

and acted much more widely across many aspects of vesicular trafficking[29]. Since the AP1

mutant did not lead to microneme exocytosis, we conclude that the secretion burst, and apical

movement are independent events facilitated by FER1 overexpression. We interpret the apical

movement as part of the process of secretory vesicle replenishment. By comparison, in mammals

a role for otoferlin in replenishment of synaptic vesicles is supported, which mimics this

particular function of FER1 in *T*. gondii[13,42,65].

The mis-localizing micronemes upon overexpression of the ΔTM allele, make us assert

that the TM domain is associated with microneme organelle trafficking. Further support is

provided by the vesicular localization in the gametocyte cytoplasm reported for the FER1

ortholog PbFLP in *P*. berghei[36]. C2 domains appear to directly engage the micronemes. Analysis

of FER1 C2 domains pointed at C2D with potential for Ca^{2+} binding and no indicators of

phospholipid binding. Among mammals, the C2D domain of otoferlin has been demonstrated to

bind Ca^{2+}[39] and interacts with MyosinVI[66] allowing vesicle transport[40]. This engagement

connects to the actin-myosin dependent microneme recycling during cell division[18], although

there is currently no indication this process is Ca^{2+}-dependent. In general, low Ca^{2+} promotes

intramolecular protein-protein interactions among otoferlin C2 domains, whereas high Ca^{2+}

triggers interaction with phospholipids[40] and SNARE proteins *in vitro*[67,68]. For *Toxoplasma*

FER1, we suggest that the C2 domains function in protein-protein interactions to expose or hide

functional domains upon signaling conditions. Indeed, we observe distinct effects on the two

microneme sub-populations, indicating that these micronemes have different proteins exposed on

their cytoplasmic facing membrane. *Plasmodium falciparum* also harbors two different

microneme populations, with different contents that, unlikely *T. gondii*, display distinct

exocytosis kinetics[69]. How *P. falciparum* regulates this or whether protein trafficking is

differentiated by Rab5A is currently unknown. In future work we will pursue the nature of this

microneme specific biology.

In conclusion, FER1 is a key player in several different aspects of microneme biology.

FER1's Golgi-endosomal localization suggests it is involved in microneme biogenesis, likely at

the vesicular trafficking level. In addition, it can direct microneme organelle traffic differentially

based on microneme population as well as differentially during microneme secretion versus cell

division. Finally, FER1 has a direct role in membrane fusion facilitating exocytosis. Thus, FER1

is a versatile player that may be a tool to understand the intricate details of microneme biology.

Material and Methods

Parasites and host cells

Transgenic derivatives of the RH strain were maintained in human foreskin fibroblasts (HFF) or

hTERT immortalized HFF cells as previously described[70]. Parasite transfections and selections

use 1 µM pyrimethamine, 20 µM chloramphenicol, 5 µg/ml FUDR, or a combination of 25

mg/ml mycophenolic acid and 50 mg/ml xanthine (MPA/X). All parasite lines were cloned by

limiting dilution.

Generation of constructs and parasite lines

All primer sequences are provided in **Supplementary Table S1**; all plasmids used are provided

in **Supplementary Table S2**. Expression plasmids fusing ddFKBP destabilization domain (DD)

with FER1 were generated from tub-DD-Myc-YFP/sagCAT plasmid[52] by replacing YFP with

the PCR amplified FER1 CDS (primer pair 1573/1574) by *Avr*II and *Xma*I restriction enzymes to

generate tub-DD-Myc-FER1[FL]/sagCAT and in tub DD-YFP-TgNek1-2(MCS)/sagCAT[71] to

generate tub-DD-YFP-FER1[FL]/sagCAT. FER1ΔTM constructs were generated by amplifying a

3' section without TM domain (primer pair 1651/1652; deletion of the C-terminal 21 aa) and

cloning the product in the FER1-FL plasmids using *Nhe*I and *Xma*I. The FER1 Ca^{2+}-binding

mutants in the C2D domain were generated by Q5 site directed mutagenesis kit (NEB) using

primers 4833/4834 to change positions A1622 and A1634 to C resulting in Asp codon 542, 545

changes to Ala, and primers 5892/5893 to change position A1814 to C resulting in Asp codon

605 to Ala.

Plasmid tub-YFP-FER1(TM) encoding only the 31 most C-terminal aa of FER1

including the TM domain was cloned by PCR amplification using primer pair 4786/4788 and

cloned by Gibson assembly into *Bgl*II/*Avr*II digested tub-YFPYFP/sagCAT plasmid[34].

Plasmid tub-IMC3mCherry/DHFR was cloned by swapping IMC3mCherry from tub-

IMC3mCherry/sagCAT[72] with *Pme*I/*Avr*II into tub-YFPYFP(MCS)/DHFR[34].

To generate pmic3-MIC3-Cherry/DHFR 1.3 kb of promoter region together with 1.2 kb

of the ORF encoding genomic locus was PCR amplified from genomic DNA (primer pair

4864/4865) and Gibson assembled into *Pme*I and *Avr*II digested tub-mCherry$_2$/DHFR[73].

Parasites expressing endogenously tagged SORTLR_Ollas/HXGPRT were generated by

inserting a PCR fragment (amplified with primer pair 6014/6015) targeting the 3'-SORTLR

locus using a SmOllas/HXGPRT template. Insertion at the C-terminus was facilitated by

CRISPR/Cas9 using a pU6 plasmid[74] containing primer 6013 as guide RNA.The

SmOllas/HXGPRT template was produced by amplifying the SmOLLAS from the

pCAG_mRuby2_smFP OLLAS plasmid (Addgene cat no. 59761[75]) using primer pair 5688/5689,

while the HXGPRT resistance cassette was amplified from the mAID-3xMyc-3'hxgprt-5'dhfr-

HXGPRT-3'dhfr plasmid[76] (primer pair 5690/4769). PCR fragments were cloned into the

previously *Not*I and *Kpn*I digested ptub-YFP-YFP/sagCAT plasmid[72,77], via Gibson assembly

(NEBuilder® HiFi 478 DNA Assembly Master Mix; NEB) (**Suppl Fig. S7**).

Antiserum generation

TgFER1 amino acids 669-877 including the diverse C2 domain DE were PCR amplified using primers Ava-LIC-Fer1-F/R and fused to a 6xHis tag in plasmid pAVA0421[78], expressed in *Escherichia coli* BL21, purified by Ni-NTA chromatography (Invitrogen), and used to immunize a guinea pig (Covance, Inc). Serum was affinity purified as described previously[79] against recombinant His6-TgFER1.

Live-cell microscopy

For monitoring egress (P30-YFP and GCaMP3 expression), parasites were grown in hTERT confluent 15 mm glass bottom cell culture dish (MatTek Corporation, cat #801002) for 30 hrs and then induced with 2 µM Shield-1 for 90 min at 37ºC. Dishes were live cell imaged on a Zeiss Axiovert 200M inverted microscope for 15 min at 2 images per minute. To monitor DD-YFP-FER1ΔTM accumulation a Leica TCS SP5 scanning confocal microscope with incubation chamber in the Boston College Imaging Core in consultation with Bret Judson. Upon addition of 1 µM Shield-1 images were captured every 5 mins for 3 hrs. All images were acquired, analyzed and adjusted using Leica, Volocity (Quorum Technologies) and/or ImageJ/FIJI software[80,81].

(Immuno-) fluorescence microscopy

Indirect immunofluorescence assays were performed on intracellular parasites grown for 18 hrs in 6-well plate containing coverslips confluent with HFF cells fixed with 100% methanol (unless stated otherwise) using primary antisera as listed in **Supplementary Table S3**. Alexa 488 (A488) or A594 conjugated goat α-mouse, α-rabbit, α-rat, or α-guinea pig were used as secondary antibodies (1:500, Invitrogen). DNA was stained with 4',6-diamidino-2-phenylindole (DAPI).

For intracellular IFAs, parasites were allowed to invade and replicate for 24 h after which 1 µM Shield-1 was applied for 18 hr (DD-Myc/YFP-FERΔTM) or 2 µM Shield-1 for 3 hr (DD-Myc-FER[FL]). Extracellular parasites grown ± Shield-1 for 18 hr were harvested by mechanical lysis and captured on Poly-L-lysine coated coverslips. A Zeiss Axiovert 200 M wide-field fluorescence microscope was used to collect images, which were deconvolved and adjusted for phase contrast using Volocity software). Co localization assays were imaged on a Zeiss AxioImager Z2 microscope with a sCMOS camera (Flash 4, Hamamatsu) using an Apotome2.

Apotome images were processed in ZEN Blue using automatic processing while Pearson

Correlation Coefficient (PCC) were measure using ImageJ.

Pan expansion microscopy

Pan expansion microscopy assays were performed basically as described[82,83]. Intracellular

parasites were grown for 18 hrs in 6-well plate containing HFF cells confluent coverslips. and

fixed with 4% PFA for 20 min at RT. Samples were then incubated in the 2x fixing/anchoring

solution (2% acrylamide; 1.4% formaldehyde in 1×PBS) for 5 h at 37 °C in a dessicator box.

Excess fixing/anchoring solution was washed with 1xPBS before sealing the coverslips in the

gelation chamber (as described in[82]). 90 µl of first gelation solution (19% sodium acrylate; 10%

acrylamide; 0.1% DHEBA; 0.25% TEMED; 0.25% APS in 1xPBS) was added directly on the

cells, before covering with a new coverslip that becomes the lid inducing the formation of a

homogeneous height gel. The gelation chamber was incubated 15 min at RT, then placed in a

humid chamber for 1.5 hrs at 37°C. After gelation, the coverslips were removed from the

gelation chamber and placed in a 6-well plate filled with 2 ml of denaturation buffer (200 mM

SDS; 200 mM NaCl; 50 mM Tris; pH 6.8) and incubated for 15 min at RT while shaking until

the gel detached from the coverslip. Next, gels were transferred in a 2 ml Eppendorf tube with 1

ml fresh denaturation buffer and incubated for 1 hr at 37°C. After denaturation, the gels were

placed in a Petri dish, washed once with 1xPBS before starting the first expansion with ddH_2O.

Water was replaced three times in 30 min intervals before full expansion overnight. Gels were

cut in approximately 1 cm² pieces and placed in a 6-well plate filled with 1.5 ml neutral hydrogel

solution (10% acrylamide; 0.05% DHEBA; 0.05% TEMED; 0.05% APS in ddH_2O) and

incubated 2x20 min at RT while shaking. Excess monomer solution was removed by pressing

carefully with Kimwipes on both sides of the gel. Gels were placed on a microscope slide and

covered with a coverslip before incubation in a humid chamber for 1 hr at 37°C.

For the second expansion, gels were placed in a 6-well plate and washed once with

1xPBS before two incubation with 1 ml second gelation solution (19% sodium acrylate; 10%

acrylamide; 0.1% bis-acrylamide; 0.05% TEMED; 0.05% APS in 1xPBS) for 15 min rocking on

ice. Gels were placed on slides and excess solution removed as described above, then covered

with coverslips and places in humid chamber for 1 hr at 37°C. Finally, the first and neutral gels

were dissolved by 2 ml of 0.2 M NaOH for 1 hr at RT while shaking. Dissolution was followed

by several washes with 1xPBS until pH becomes neutral. For antibody staining, small pieces of

gel were cut and incubated with α-M2AP (1:250) followed by AlexaFluor 594 (1:200)

antibodies, overnight at RT followed by 3 hrs at 37°C the next day. Antibodies were diluted in

2% BSA and washed 3x20 min with PBS-Tween (0.1%). Finally, NHS-ester 488 staining was

performed at 20 µg/ml in 1xPBS for 1 hr. Washing was performed as described above, before a

final expansion overnight in ddH$_2$0. Gels were imaged in 35 mm Mattek on a Zeiss LSM 880

using Airyscan and a Plan-APOCHROMAT (x40; water). Z-stacks were collected for all gels

using 0.2 µm step size through the entire depth of parasites. Post processing of Airyscan images

was completed in ZEN Blue and ImageJ.

Electron microscopy

For ultrastructural observations of *T. gondii*-infected cells by thin-section transmission electron

microscopy (TEM), infected cells were fixed in 2.5% glutaraldehyde in 0.1 mM sodium

cacodylate (EMS) and processed as described[84]. Ultrathin sections of infected cells were stained

before examination with a Philips CM120 EM (Eindhoven, the Netherlands) under 80 kV. For

immunoelectron microscopy (IEM) samples were prepared as described before[34]. Sections were

immunolabeled with MIC2 MAb 6D10 in 1% fish skin gelatin and then with goat anti-IgG

antibodies, followed by 10-nm protein A-gold particles before examination with a Philips

CM120 electron microscope under 80 kV.

Host cell invasion

Extracellular parasites from 80% naturally lysed flask were induced with 2 µM Shield-1 for 90

min at 37°C before they were allowed to invade HFF host cells for 10 min at 37°C[7]. The

red/green invasion assay was performed as described[85] using Alexa594- and Alexa488-

conjugated SAG1 MAb T41E5. Three images were taken per biological replicate (3 biological

replicates total) on an EVOS FL (Life Technologies). The number of invaded versus uninvaded

parasites were enumerated manually for at least 300 parasites per counted sample.

Secretagogue induced egress

The egress assay was performed essentially as described previously[34]. Parasites were grown in

HFF monolayers for 30 h after which the phenotype was induced with 1 µM Shield-1 for 18 hrs.

Egress was triggered with 1-3 µM A23187, 500 µM propranolol, 50-150 µM BIPPO (kindly

shared by Dr. Jeff Dvorin) 0.01-0.1% saponin or DMSO for 5 min at 37°C, followed by IFA

with IMC3 and GRA3 antisera. Egressed, permeabilized and intact vacuoles were counted[56]. 100

vacuoles were counted for each experiment and three biological replicates were performed.

Shield-1 induced egress

Parasites were inoculated on HFF coverslips and allowed to grow for 30 hrs and then induced

with either 1 µM Shield-1 for 2 hrs or 1 µM calcium ionophore A23187 for 5 min as a control,

prior to fixation with 100% methanol and IFA staining using MAb 9E10 cMyc and rabbit αMIC8

(antisera details in **Supplementary Table S1**). DNA was stained with 4',6-diamidino-2-

phenylindole (DAPI). The number of intact vacuoles per 20 fields was enumerated. Three

independent biological replicates per experiment were performed.

Combined invasion-egress Assay

Parasites were grown for 36 hours, mechanically lysed in standard ED1 parasite medium and

allowed to invade coverslips coated in an HFF monolayer for 1 hr. All unattached parasites were

then washed off with a PBS and coverslips were incubated in 1 µM Shield-1 or vehicle control

for an additional 2 hrs. Coverslips were then fixed with PFA and IFA was performed as

described for a typical red-green invasion assay. The number of parasites was counted in 10

fields and four independent biological replicates per experiment were performed.

Microneme secretion by Western blotting

Microneme secretion by western blot was performed as published[16]. Freshly lysed parasites were

resuspended in DMEM/FBS and transferred to a 96-well polystyrene round-bottom plate

(CELLTREAT Scientific Products). Secretion was induced by 1-3 µM A23187, 500 µM

propranolol, 1% ethanol or DMSO for 5 min at 37°C. Constitutive microneme secretion was

assessed by incubation without secretagogue at 37°C for 60 min. Supernatants were probed by

western blot using 6D10 MIC2 MAb and TG17.43 GRA1 MAb and HRP conjugated secondary

antiserum. Signals were quantified using a densitometer. Three independent biological replicates

per experiment were performed.

Microneme abundance by Western blotting

Intracellular parasites were stimulated overnight with 0, 1 or 2 µM Shield-1 (AOBIOUS) at

37°C. The day after, intracellular parasites were washed with DMEM/FBS and released from

host cell using a 26G needle. Parasites were resuspended in RIPA buffer and probed by western

blot using 6D10 MIC2 Mab and 12G10 Tubulin Mab and HRP conjugated secondary antiserum.

Signals were quantified for three independent biological replicates.

Microneme secretion by IFA

IFA on parasites exposed to a host cell monolayer was performed as reported[34]. Parasites were

resuspended in Endo buffer and spun onto HFF cells in a 6-well plate at 28*g, 5 min, RT and

allowed to settle for 20 min at 37°C. Secretion was induced by replacing the buffer with 3% FBS

in DMEM and 10 mM HEPES (pH 7.2) and incubation at 37°C for 5 min. PBS-washed

coverslips were fixed with 4% formaldehyde and 0.02% glutaraldehyde, and subjected to IFA

with anti-MIC3, -MIC5, -MIC8 or MIC10 in the presence of 0.02% saponin.

Gliding motility trail assay

Trail assays were performed as previously described[28]. Parasites were induced with 1 µM Shield-

1 for 18 hr, mechanically released, resuspended in ED1 with 1 µM Shield-1 and incubated on

poly-L-lysine coated coverslips for 15 min at 37°C. Parasites were fixed with 4% formaldehyde

and 0.02% glutaraldehyde and stained with DG52 SAG1 MAb (64) to visualize trails.

Statistics.

Student's paired *t*-test, one-way and two-way analysis of variance (ANOVA) using post hoc

Tukey correction were performed.

Acknowledgements

We thank Bret Judson and Dr. Patrick Autissier of the Boston College Imaging and Flow

Cytometry Cores, respectively, for infrastructure and support, Dr. Sander Groffen for assistance

with molecular modeling, Klemens Engelberg for establishing the *Toxoplasma* pan-ExM

protocol, Emily Stoneburner, Natalie Sandlin, and Elizabeth Gray for technical support, Drs.

Gustavo Arrizabalaga, Peter Bradley, Vern Carruthers, Iain Cheeseman, Jean-François

Dubremetz, Wassim Daher, Jeff Dvorin, Maryse Lebrun, Sebastian Lourido, Sabrina Marion,

Silvia Moreno, Naomi Morrissette, Jeroen Saeij, David Sibley, and Gary Ward for sharing

reagents. We also thank Kim Zichichi at the EM Facility at Yale University for processing and

immunolabeling our samples.

618   This study was supported by National Science Foundation (NSF) Major Research

619   Instrumentation grant 1626072, National Institutes of Health grants AI108251 (B.I.C.),

620   AI060767 (I.C.), AI122042 (M.-J.G), AI099658 (M.-J.G.), and AI122923 (M.-J.G.), and

621   American Cancer Society grant RSG-12-175-01-MPC (M.-J.G.).

Author contributions

DNAT performed all functional experiments on the DD-[Myc/YFP]-FER1ΔTM parasites,

imaging of the DD-Myc-FER1[FL] line, and time lapse microscopy, ARES performed αFER1

localizations, pan-ExM, MIC2 quantification by western blot, generated DDD-AAA mutants and

quantified their phenotypes, AAD performed all functional studies on the DD-Myc-FER1[FL]

parasites, IC performed all electron microscopy studies, BIC cloned FER1 cDNA, established

the overexpression plasmids and assessed the initial phenotype, DNAT, ARES, AAD and MJG

designed the experiments and interpreted the data, ARES and MJG wrote the manuscript and all

authors reviewed and edited the manuscript.

Competing Interests

634   The authors declare no competing interests.

References

639   1   Blader, I. J., Coleman, B. I., Chen, C. T. & Gubbels, M. J. Lytic Cycle of Toxoplasma
640   gondii: 15 Years Later. *Annu Rev Microbiol* **69**, 463-485, doi:10.1146/annurev-micro-
641   091014-104100 (2015).
642   2   Carruthers, V. B. & Tomley, F. M. Microneme proteins in apicomplexans. *Subcell
643   Biochem* **47**, 33-45 (2008).
644   3   Sheiner, L. *et al.* Toxoplasma gondii transmembrane microneme proteins and their
645   modular design. *Mol Microbiol*, doi:10.1111/j.1365-2958.2010.07255.x (2010).

4   Kafsack, B. F. *et al.* Rapid membrane disruption by a perforin-like protein facilitates
parasite exit from host cells. *Science* **323**, 530-533, doi:1165740 [pii]
10.1126/science.1165740 (2009).
5   Roiko, M. S., Svezhova, N. & Carruthers, V. B. Acidification Activates Toxoplasma
gondii Motility and Egress by Enhancing Protein Secretion and Cytolytic Activity. *PLoS*
Pathog **10**, e1004488, doi:10.1371/journal.ppat.1004488 (2014).
6   Bullen, H. E., Bisio, H. & Soldati-Favre, D. The triumvirate of signaling molecules
controlling Toxoplasma microneme exocytosis: Cyclic GMP, calcium, and phosphatidic
acid. *PLoS Pathog* **15**, e1007670, doi:10.1371/journal.ppat.1007670 (2019).
7   Uboldi, A. D. *et al.* Protein kinase A negatively regulates Ca2+ signalling in Toxoplasma
gondii. *PLoS Biol* **16**, e2005642, doi:10.1371/journal.pbio.2005642 (2018).
8   Leung, J. M. *et al.* Stability and function of a putative microtubule-organizing center in
the human parasite Toxoplasma gondii. *Mol Biol Cell* **28**, 1361-1378,
doi:10.1091/mbc.E17-01-0045 (2017).
9   Kremer, K. *et al.* An overexpression screen of Toxoplasma gondii Rab-GTPases reveals
distinct transport routes to the micronemes. *PLoS Pathog* **9**, e1003213,
doi:10.1371/journal.ppat.1003213 (2013).
10   Jacot, D. *et al.* An Apicomplexan Actin-Binding Protein Serves as a Connector and Lipid
Sensor to Coordinate Motility and Invasion. *Cell Host Microbe* **20**, 731-743,
doi:10.1016/j.chom.2016.10.020 (2016).
11   Sakura, T. *et al.* A Critical Role for Toxoplasma gondii Vacuolar Protein Sorting VPS9
in Secretory Organelle Biogenesis and Host Infection. *Sci Rep* **6**, 38842,
doi:10.1038/srep38842 (2016).
12   Paredes-Santos, T. C., de Souza, W. & Attias, M. Dynamics and 3D organization of
secretory organelles of Toxoplasma gondii. *Journal of structural biology* **177**, 420-430,
doi:10.1016/j.jsb.2011.11.028 (2012).
13   Saha, S., Coleman, B. I., Dubey, R., Blader, I. J. & Gubbels, M. J. Two
Phosphoglucomutase Paralogs Facilitate Ionophore-Triggered Secretion of the
Toxoplasma Micronemes. *mSphere* **2**, doi:10.1128/mSphere.00521-17 (2017).
14   Wetzel, D. M., Chen, L. A., Ruiz, F. A., Moreno, S. N. & Sibley, L. D. Calcium-
mediated protein secretion potentiates motility in Toxoplasma gondii. *J Cell Sci* **117**,
5739-5748 (2004).
15   Lentini, G., Dubois, D. J., Maco, B., Soldati-Favre, D. & Frenal, K. The roles of Centrin
2 and Dynein Light Chain 8a in apical secretory organelles discharge of Toxoplasma
gondii. *Traffic*, doi:10.1111/tra.12673 (2019).
16   Carruthers, V. B. & Sibley, L. D. Mobilization of intracellular calcium stimulates
microneme discharge in Toxoplasma gondii. *Mol Microbiol* **31**, 421-428 (1999).
17   Gras, S. *et al.* An endocytic-secretory cycle participates in Toxoplasma gondii in motility.
PLoS Biol **17**, e3000060, doi:10.1371/journal.pbio.3000060 (2019).
18   Periz, J. *et al.* A highly dynamic F-actin network regulates transport and recycling of
micronemes in Toxoplasma gondii vacuoles. *Nat Commun* **10**, 4183,
doi:10.1038/s41467-019-12136-2 (2019).
19   Venugopal, K. & Marion, S. Secretory organelle trafficking in Toxoplasma gondii: A
long story for a short travel. *Int J Med Microbiol* **308**, 751-760,
doi:10.1016/j.ijmm.2018.07.007 (2018).

20   Sangare, L. O. *et al.* Unconventional endosome-like compartment and retromer complex
in Toxoplasma gondii govern parasite integrity and host infection. *Nat Commun* **7**,
11191, doi:10.1038/ncomms11191 (2016).
21   Tomavo, S., Slomianny, C., Meissner, M. & Carruthers, V. B. Protein trafficking through
the endosomal system prepares intracellular parasites for a home invasion. *PLoS Pathog*
9, e1003629, doi:10.1371/journal.ppat.1003629 (2013).
22   Thornton, L. B. *et al.* An ortholog of Plasmodium falciparum chloroquine resistance
transporter (PfCRT) plays a key role in maintaining the integrity of the endolysosomal
system in Toxoplasma gondii to facilitate host invasion. *PLoS Pathog* **15**, e1007775,
doi:10.1371/journal.ppat.1007775 (2019).
23   Dubois, D. J. & Soldati-Favre, D. Biogenesis and secretion of micronemes in
Toxoplasma gondii. *Cell Microbiol* **21**, e13018, doi:10.1111/cmi.13018 (2019).
24   McGovern, O. L., Rivera-Cuevas, Y., Kannan, G., Narwold, A. J. & Carruthers, V. B.
Intersection of endocytic and exocytic systems in Toxoplasma gondii. *Traffic* **19**, 336-
353, doi:10.1111/tra.12556 (2018).
25   Stasic, A. J. *et al.* The Toxoplasma Vacuolar H(+)-ATPase Regulates Intracellular pH
and Impacts the Maturation of Essential Secretory Proteins. *Cell Rep* **27**, 2132-2146
e2137, doi:10.1016/j.celrep.2019.04.038 (2019).
26   Sloves, P. J. *et al.* Toxoplasma sortilin-like receptor regulates protein transport and is
essential for apical secretory organelle biogenesis and host infection. *Cell Host Microbe*
11, 515-527, doi:10.1016/j.chom.2012.03.006 (2012).
27   Houngue, R. *et al.* Toxoplasma membrane inositol phospholipid binding protein
TgREMIND is essential for secretory organelle function and host infection. *Cell Rep* **43**,
113601, doi:10.1016/j.celrep.2023.113601 (2023).
28   Morlon-Guyot, J., Pastore, S., Berry, L., Lebrun, M. & Daher, W. Toxoplasma gondii
Vps11, a subunit of HOPS and CORVET tethering complexes, is essential for the
biogenesis of secretory organelles. *Cellular Microbiology*, doi:10.1111/cmi.12426
(2015).
29   Venugopal, K. *et al.* Dual role of the Toxoplasma gondii clathrin adaptor AP1 in the
sorting of rhoptry and microneme proteins and in parasite division. *PLoS Pathog* **13**,
e1006331, doi:10.1371/journal.ppat.1006331 (2017).
30   Bradley, P. J. & Boothroyd, J. C. The pro region of Toxoplasma ROP1 is a rhoptry-
targeting signal. *Int J Parasitol* **31**, 1177-1186 (2001).
31   Bradley, P. J., Li, N. & Boothroyd, J. C. A GFP-based motif-trap reveals a novel
mechanism of targeting for the Toxoplasma ROP4 protein. *Mol Biochem Parasitol* **137**,
111-120 (2004).
32   Martens, S. & McMahon, H. T. Mechanisms of membrane fusion: disparate players and
common principles. *Nat Rev Mol Cell Biol* **9**, 543-556, doi:nrm2417 [pii]
10.1038/nrm2417 (2008).
33   Farrell, A. *et al.* A DOC2 protein identified by mutational profiling is essential for
apicomplexan parasite exocytosis. *Science* **335**, 218-221, doi:335/6065/218 [pii]
10.1126/science.1210829 (2012).
34   Coleman, B. I. *et al.* A member of the ferlin calcium sensor family is essential for
Toxoplasma gondii rhoptry secretion. *MBio* **9**, e01510-01518 (2018).
35   Tagoe, D. N. A. *et al.* Ferlins and TgDOC2 in Toxoplasma Microneme, Rhoptry and
Dense Granule Secretion. *Life (Basel)* **11**, doi:10.3390/life11030217 (2021).

36 Obrova, K., Cyrklaff, M., Frank, R., Mair, G. R. & Mueller, A. K. Transmission of the
malaria parasite requires ferlin for gamete egress from the red blood cell. *Cell Microbiol*,
e12999, doi:10.1111/cmi.12999 (2018).
37 Lek, A., Evesson, F. J., Sutton, R. B., North, K. N. & Cooper, S. T. Ferlins: regulators of
vesicle fusion for auditory neurotransmission, receptor trafficking and membrane repair.
Traffic **13**, 185-194, doi:10.1111/j.1600-0854.2011.01267.x (2012).
38 Redpath, G. M., Sophocleous, R. A., Turnbull, L., Whitchurch, C. B. & Cooper, S. T.
Ferlins Show Tissue-Specific Expression and Segregate as Plasma Membrane/Late
Endosomal or Trans-Golgi/Recycling Ferlins. *Traffic* **17**, 245-266, doi:10.1111/tra.12370
(2016).
39 Roux, I. *et al.* Otoferlin, defective in a human deafness form, is essential for exocytosis at
the auditory ribbon synapse. *Cell* **127**, 277-289, doi:10.1016/j.cell.2006.08.040 (2006).
40 Pangrsic, T., Reisinger, E. & Moser, T. Otoferlin: a multi-C2 domain protein essential for
hearing. *Trends Neurosci* **35**, 671-680, doi:10.1016/j.tins.2012.08.002 (2012).
41 Hams, N., Padmanarayana, M., Qiu, W. & Johnson, C. P. Otoferlin is a multivalent
calcium-sensitive scaffold linking SNAREs and calcium channels. *Proc Natl Acad Sci U
S A* **114**, 8023-8028, doi:10.1073/pnas.1703240114 (2017).
42 Michalski, N. *et al.* Otoferlin acts as a Ca(2+) sensor for vesicle fusion and vesicle pool
replenishment at auditory hair cell ribbon synapses. *eLife* **6**, doi:10.7554/eLife.31013
(2017).
43 Lek, A., Lek, M., North, K. N. & Cooper, S. T. Phylogenetic analysis of ferlin genes
reveals ancient eukaryotic origins. *BMC evolutionary biology* **10**, 231, doi:10.1186/1471-
2148-10-231 (2010).
44 Sidik, S. M. *et al.* A Genome-wide CRISPR Screen in Toxoplasma Identifies Essential
Apicomplexan Genes. *Cell* **166**, 1423-1435 e1412, doi:10.1016/j.cell.2016.08.019
(2016).
45 Barylyuk, K. *et al.* A Comprehensive Subcellular Atlas of the Toxoplasma Proteome via
hyperLOPIT Provides Spatial Context for Protein Functions. *Cell Host Microbe* **28**, 752-
766, doi:10.1016/j.chom.2020.09.011 (2020).
46 Bisio, H. & Soldati-Favre, D. Signaling Cascades Governing Entry into and Exit from
Host Cells by Toxoplasma gondii. *Annu Rev Microbiol* **73**, 579-599,
doi:10.1146/annurev-micro-020518-120235 (2019).
47 Breinich, M. S. *et al.* A Dynamin Is Required for the Biogenesis of Secretory Organelles
in Toxoplasma gondii. *Curr Biol* **19**, 277-286, doi:S0960-9822(09)00611-3 [pii]
10.1016/j.cub.2009.01.039 (2009).
48 Drozdowicz, Y. M. *et al.* Isolation and characterization of TgVP1, a type I vacuolar H+-
translocating pyrophosphatase from Toxoplasma gondii. The dynamics of its subcellular
localization and the cellular effects of a diphosphonate inhibitor. *J Biol Chem* **278**, 1075-
1085 (2003).
49 Miranda, K. *et al.* Characterization of a novel organelle in Toxoplasma gondii with
similar composition and function to the plant vacuole. *Mol Microbiol* **76**, 1358-1375,
doi:MMI7165 [pii] 10.1111/j.1365-2958.2010.07165.x (2010).
50 Luo, S., Ruiz, F. A. & Moreno, S. N. The acidocalcisome Ca2+-ATPase (TgA1) of
Toxoplasma gondii is required for polyphosphate storage, intracellular calcium
homeostasis and virulence. *Mol Microbiol* **55**, 1034-1045, doi:MMI4464 [pii]
10.1111/j.1365-2958.2004.04464.x (2005).

783 51 Francia, M. E. *et al.* A Toxoplasma gondii protein with homology to intracellular type
784   Na(+)/H(+) exchangers is important for osmoregulation and invasion. *Exp Cell Res*,
785   doi:S0014-4827(11)00125-X [pii] 10.1016/j.yexcr.2011.03.020 (2011).

786 52 Herm-Gotz, A. *et al.* Rapid control of protein level in the apicomplexan Toxoplasma
787   gondii. *Nat Methods* **4**, 1003-1005 (2007).

788 53 Brydges, S. D., Harper, J. M., Parussini, F., Coppens, I. & Carruthers, V. B. A transient
789   forward-targeting element for microneme-regulated secretion in Toxoplasma gondii.
790   *Biology of the Cell* **100**, 253-264, doi:10.1042/Bc20070076 (2008).

791 54 Nishi, M., Hu, K., Murray, J. M. & Roos, D. S. Organellar dynamics during the cell cycle
792   of Toxoplasma gondii. *J Cell Sci* **121**, 1559-1568 (2008).

793 55 Zarringhalam, K., Ye, S., Lou, J., Rezvani, Y. & Gubbels, M. J. Cell cycle-regulated
794   ApiAP2s and parasite development: the Toxoplasma paradigm. *Curr Opin Microbiol* **76**,
795   102383, doi:10.1016/j.mib.2023.102383 (2023).

796 56 Bisio, H., Lunghi, M., Brochet, M. & Soldati-Favre, D. Phosphatidic acid governs natural
797   egress in Toxoplasma gondii via a guanylate cyclase receptor platform. *Nat Microbiol* **4**,
798   420-428, doi:10.1038/s41564-018-0339-8 (2019).

799 57 Kafsack, B. F., Beckers, C. & Carruthers, V. B. Synchronous invasion of host cells by
800   Toxoplasma gondii. *Mol Biochem Parasitol* **136**, 309-311 (2004).

801 58 Martens, S. Role of C2 domain proteins during synaptic vesicle exocytosis. *Biochem Soc*
802   *Trans* **38**, 213-216, doi:BST0380213 [pii] 10.1042/BST0380213 (2010).

803 59 Verdaguer, N., Corbalan-Garcia, S., Ochoa, W. F., Fita, I. & Gomez-Fernandez, J. C.
804   Ca(2+) bridges the C2 membrane-binding domain of protein kinase Calpha directly to
805   phosphatidylserine. *EMBO J* **18**, 6329-6338, doi:10.1093/emboj/18.22.6329 (1999).

806 60 Chapman, E. R. & Davis, A. F. Direct interaction of a Ca2+-binding loop of
807   synaptotagmin with lipid bilayers. *J Biol Chem* **273**, 13995-14001,
808   doi:10.1074/jbc.273.22.13995 (1998).

809 61 Davis, D. B., Doherty, K. R., Delmonte, A. J. & McNally, E. M. Calcium-sensitive
810   phospholipid binding properties of normal and mutant ferlin C2 domains. *J Biol Chem*
811   **277**, 22883-22888, doi:10.1074/jbc.M201858200 (2002).

812 62 Kaeser, P. S. & Regehr, W. G. Molecular mechanisms for synchronous, asynchronous,
813   and spontaneous neurotransmitter release. *Annual review of physiology* **76**, 333-363,
814   doi:10.1146/annurev-physiol-021113-170338 (2014).

815 63 Kaeser, P. S. & Regehr, W. G. The readily releasable pool of synaptic vesicles. *Curr*
816   *Opin Neurobiol* **43**, 63-70, doi:10.1016/j.conb.2016.12.012 (2017).

817 64 Meese, S. *et al.* Activity-Dependent Phosphorylation by CaMKIIdelta Alters the Ca(2+)
818   Affinity of the Multi-C2-Domain Protein Otoferlin. *Front Synaptic Neurosci* **9**, 13,
819   doi:10.3389/fnsyn.2017.00013 (2017).

820 65 Pangrsic, T. *et al.* Hearing requires otoferlin-dependent efficient replenishment of
821   synaptic vesicles in hair cells. *Nat Neurosci* **13**, 869-876, doi:10.1038/nn.2578 (2010).

822 66 Heidrych, P. *et al.* Otoferlin interacts with myosin VI: implications for maintenance of
823   the basolateral synaptic structure of the inner hair cell. *Human molecular genetics* **18**,
824   2779-2790, doi:10.1093/hmg/ddp213 (2009).

825 67 Ramakrishnan, N. A., Drescher, M. J., Morley, B. J., Kelley, P. M. & Drescher, D. G.
826   Calcium regulates molecular interactions of otoferlin with soluble NSF attachment
827   protein receptor (SNARE) proteins required for hair cell exocytosis. *J Biol Chem* **289**,
828   8750-8766, doi:10.1074/jbc.M113.480533 (2014).

68   Johnson, C. P. & Chapman, E. R. Otoferlin is a calcium sensor that directly regulates
SNARE-mediated membrane fusion. *J Cell Biol* **191**, 187-197,
doi:10.1083/jcb.201002089 (2010).
69   Healer, J., Crawford, S., Ralph, S., McFadden, G. & Cowman, A. F. Independent
translocation of two micronemal proteins in developing Plasmodium falciparum
merozoites. *Infect Immun* **70**, 5751-5758, doi:10.1128/IAI.70.10.5751-5758.2002 (2002).
70   Roos, D. S., Donald, R. G., Morrissette, N. S. & Moulton, A. L. Molecular tools for
genetic dissection of the protozoan parasite Toxoplasma gondii. *Methods Cell Biol* **45**,
27-63 (1994).
71   Chen, C. T. & Gubbels, M. J. The Toxoplasma gondii centrosome is the platform for
internal daughter budding as revealed by a Nek1 kinase mutant. *J Cell Sci* **126**, 3344-
3355, doi:10.1242/jcs.123364 (2013).
72   Anderson-White, B. R. *et al.* A family of intermediate filament-like proteins is
sequentially assembled into the cytoskeleton of Toxoplasma gondii. *Cell Microbiol* **13**,
18-31 (2011).
73   Schaeffer, M. *et al.* Dynamic imaging of T cell-parasite interactions in the brains of mice
chronically infected with Toxoplasma gondii. *J Immunol* **182**, 6379-6393,
doi:182/10/6379 [pii] 10.4049/jimmunol.0804307 (2009).
74   Sidik, S. M., Huet, D. & Lourido, S. CRISPR-Cas9-based genome-wide screening of
Toxoplasma gondii. *Nature protocols* **13**, 307-323, doi:10.1038/nprot.2017.131 (2018).
75   Viswanathan, S. *et al.* High-performance probes for light and electron microscopy. *Nat*
Methods **12**, 568-576, doi:10.1038/nmeth.3365 (2015).
76   Lou, J. *et al.* Single cell expression and chromatin accessibility of the Toxoplasma gondii
lytic cycle identifies AP2XII-8 as an essential controller of a ribosome regulon. *bioRxiv*
2023.10.06.561197, doi:https://doi.org/10.1101/2023.10.06.561197 (2023).
77   Gubbels, M. J., Li, C. & Striepen, B. High-throughput growth assay for Toxoplasma
gondii using yellow fluorescent protein. *Antimicrob Agents Chemother* **47**, 309-316
(2003).
78   Alexandrov, A. *et al.* A facile method for high-throughput co-expression of protein pairs.
Mol Cell Proteomics **3**, 934-938 (2004).
79   Gubbels, M. J., Vaishnava, S., Boot, N., Dubremetz, J. F. & Striepen, B. A MORN-repeat
protein is a dynamic component of the Toxoplasma gondii cell division apparatus. *J Cell*
Sci **119**, 2236-2245, doi:10.1242/jcs.02949 (2006).
80   Abramoff, M. D., Magalhaes, P. J. & Ram, S. J. Image Processing with ImageJ.
Biophotonics International **11**, 36-42 (2004).
81   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
82   Louvel, V. *et al.* iU-ExM: nanoscopy of organelles and tissues with iterative
ultrastructure expansion microscopy. *Nat Commun* **14**, 7893, doi:10.1038/s41467-023-
43582-8 (2023).
83   M'Saad, O. & Bewersdorf, J. Light microscopy of proteins in their ultrastructural context.
Nat Commun **11**, 3850, doi:10.1038/s41467-020-17523-8 (2020).
84   Coppens, I. & Joiner, K. A. Host but not parasite cholesterol controls Toxoplasma cell
entry by modulating organelle discharge. *Mol Biol Cell* **14**, 3804-3820,
doi:10.1091/mbc.E02-12-0830 (2003).

85   Carey, K. L., Westwood, N. J., Mitchison, T. J. & Ward, G. E. A small-molecule
approach to studying invasive mechanisms of Toxoplasma gondii. *Proc Natl Acad Sci U*
S A **101**, 7433-7438 (2004).
86   Jimenez, J. L. & Bashir, R. In silico functional and structural characterisation of ferlin
proteins by mapping disease-causing mutations and evolutionary information onto three-
dimensional models of their C2 domains. *Journal of the neurological sciences* **260**, 114-
123, doi:10.1016/j.jns.2007.04.016 (2007).
881
882
883

Figure 1

Figure 1. FER1 localizes to the secretory pathway. (**a**) Schematic representation of TgFER1. Yellow letters mark the C2 domains; the aa 669-877 region used to generate a specific antiserum is marked in green; FERI is a domain conserved in most ferlins with unknown function; TM is the transmembrane domain. (**b**) Cropped western blot analysis with the affinity purified guinea pig polyclonal antiserum generated against the FER1 region marked in panel A. Total lysate of wild type (RHΔKu80) parasites was loaded. (**c**) IFA of intracellular parasites and (**d**) extracellular parasites, co-stained for FER1 and MIC2 using respectively affinity purified FER1 antiserum and α-MIC2 serum. Samples were fixed with 4% PFA (**e**) Analysis of affinity purified FER1 antiserum localization in intracellularly replicating parasites by IFA co-stained with α-SERCA serum marking the endoplasmic reticulum (ER), GRASP55-RFP expressing parasites marking the *cis*-Golgi, NAGTI-YFP expressing parasites for the middle-Golgi, GalNac-YFP, and SORTLR-smOLLAS expressing parasites for the trans-Golgi, α-DrpB serum marking the *trans*-Golgi endosomes, α-Pro-M2AP serum marking the endosome like compartment (ELC) and, α-VP1 serum marking the plant like vacuole (PLV/VAC) and acidocalcisomes. PCC: Pearson correlation coefficients (yellow) for each image. (**f**) Pearson correlation coefficients reflect the level of FER1 colocalization with the compartment specific marker, n=10. (**g**) Schematic representation of FER1 localization through the secretory pathway. All IFA panels represent the wild type RHΔKu80 strain. See **Supplementary Figure S1** for marker cross-calibration.

Figure 2

 Figure 2. Generation and validation of conditional FER1 overexpression parasite lines. (**a**)

 Schematic representation of the overexpression constructs driven by the strong constitutive α-

 tubulin promoter (*ptub*). DD: destabilization domain; Myc: cMyc epitope tag; TM:

 transmembrane domain; FL: full length. (**b**) Western blot analysis of the overexpression parasite

 lines. Polyclonal guinea pig FER1 antiserum as in **Fig 1**. Monoclonal antibody12G10

 recognizing α-tubulin was used as loading control. Parasites were induced with 1 µM Shield-1

 for 24 hrs. (**c**) Plaque assays of infected HFF monolayers grown for 7 days ±1 µM Shield-1.

Figure 3

Figure 3. Phenotypic characterization of parasites overexpressing DD-Myc-FER1ΔTM. (**a**) Red-green invasion assay reveal an invasion defect. n=3+std. (**b, c**) Cell division and cell cycle progression analysis. Cell cycle stages quantified in the right of panel **c** are shown on the left. "C" and "B" represent the number of centrosomes and daughter buds, respectively. Parasites

925 were allowed to invade for 2 hrs upon which 1 µM Shield-1 was added for 18 hrs. p=0.031 (*t*-

926 test) accumulation of parasite in the 4-cells/vacuole stage. n=3+std. (**d,e**) Induced egress assays.

927 Parasites grown for 30 hrs and induced for 18 hrs with 1 µM Shield-1 were triggered for egress

928 with calcium ionophore A23187 (or vehicle control), fixed and stained with α-IMC3 (parasite

929 cortex) and α-GRA3 (PVM) sera and scored for status of vacuole permeabilization and/or egress

930 (**d**; representative image; arrows mark holes in the PVM of permeabilized vacuoles). n=3±std.

931 Parasites were not under Shield-1 pressure during the 5 min pharmacological incubation. (**f**) Trail

932 assay using α-SAG1 serum to assess gliding motility. Parasites were induced for 18 hrs ± 1 µM

933 Shield-1, mechanically released from host cells and kept under 1 µM Shield-1 throughout the 30

934 min gliding experiment at 37ºC. (**g**) Assessment of microneme secretion by western blot

935 detection of MIC2 released in the supernatant under various triggers. Parasites were induced 18

936 hrs with 1 µM Shield-1 and harvested by physical release from the host cell. 10%: 10% of total

937 lysate; const.: 1 hr constitutive secretion at 37ºC (no secretagogue); 1% ethanol; 2 µM A23187;

938 500 µM propranolol. DMSO is the vehicle control for A23187. Induced secretion for 5 min at

939 37ºC. Bottom of panel represents quantified secretion normalized to the GRA1 signal and to the

940 10% loading control for each condition. n=3+std. Parent line controls in **Supplementary Figure**

941 **S2**. Parasites were not under Shield-1 pressure during secretion assay. (**h**) Secretion of the

942 Rab5A/C-dependent microneme population was assessed by IFA on non-permeabilized parasites

943 induced for 18 hrs ± 1 µM Shield-1, mechanically released, and exposed to fresh host cells for 5

944 min at 37ºC. Parasites were not under Shield-1 pressure during assay.

Figure 4

Figure 4. Overexpression of dominant negative FER1ΔTM constructs result in microneme mis-

localization. **(a, b)** Overexpression of DD-Myc-FER1ΔTM (**a**) or DD-YFP-FER1ΔTM (**b**) leads

to accumulation of microneme protein, visualized with α-MIC2, in a central, apical location.

YFP and MIC2 co-localization was assesses by Pearson correlation coefficient. (**c**) Microneme

proteins of the Rab5A/C-dependent trafficking pathway, visualized with α-MIC5, also

accumulate in the FER1 compartment. (**d**) α-FER1 serum confirms exclusive accumulation in

the microneme protein compartment. (**e**) ROP proteins do not accumulate and rhoptry

morphology is normal. (**f**) Co-localization of VP1 and the YFP accumulation is not detected. (**g**)

Markers for *cis*-Golgi, *trans*-Golgi network (TGN), as well as early (EE) and late (LE) endosome

markers localize normally and do no co-localize with the YFP accumulation. In all IFA

experiments parasites were treated with 1 µM Shield-1 for 18 hrs. (**h**) Pan-expansion microscopy

of RHΔku80 parent line and DD-YFP-FER1ΔTM overexpressing parasites induced for 18 hrs

with 1 µM Shield-1. Compiled stacks of 20 slices are shown. Green arrowhead marks the radial

set of microneme proteins in the RHΔku80 parasites. (**i**) TEM of DD-Myc-FER1ΔTM

overexpressing parasites induced for 16 hrs with 1 µM Shield-1. Dotted circle marks cytoplasmic

microneme accumulation in the apical cytosol, which is enlarged in the inset.

Figure 5

 Figure 5. Microneme trafficking dynamics upon DD-YFP-FER1ΔTM overexpression. (**a**)

 Normal maturation by pro-peptide processing in the ELC/PLV compartments was demonstrated

 by western blot using serum against the M2AP pro-peptide (α-proM2AP) and the mature protein

(α-M2AP). α-tubulin serves as loading control. (**b**) IFA showing the normal localization of

proMIC5 in the ELC/PLV compartment in newly forming daughter buds using α-proMIC5. (**c**)

Induced mutants co-stained with YFP, proMIC5, and IMC3 antiserum to track the timing and

localization of proMIC5. Arrowheads mark the proMIC5 compartment within daughter buds.

DD-YFP-FER1ΔTM is exclusively present in the mother parasites**. **(**d**) Induced mutants co-

stained with IMC1 antiserum to track division stages as indicated. Arrowheads mark the mother

parasite's YFP accumulation migrating into a basal direction during progression of daughter

budding. Note the absence of YFP in the daughter buds. (**e**) Induced mutants co-stained with

YFP, MIC2, and IMC3 antisera to track the localization of mature MIC2 protein through cell

division. DD-YFP-FER1ΔTM consistently colocalize with MIC2 throughout the division stages.

(**f**) Time courses of the incidence of vacuoles or individual parasites with mis-localizing

micronemes in intracellular and extracellular DD-YFP-FER1ΔTM parasites visualized through

α-MIC2 IFA. At least 100 vacuoles (intracellular) or parasites (extracellular) were counted per

time point in reversion experiments. n=3±std (intracellular) or n=5±std (extracellular). (**g**) Select

panels from time lapse experiment with DD-YFP-FER1ΔTM parasites co-transfected with

MIC3-mCherry to track microneme localization dynamics. At t=0, 2 μM Shield-1 was added.

Yellow asterisk marks the first time point at which the piling up of YFP-FER1 could be

convincingly observed; while purple asterisk marks the first time where MIC3-mCherry can be

seen re-localizing from the apical cortex to the central apical localization co-localizing with

YFP-FER1. Panels from supplementary movie S1. In all experiments parasites were treated with

1 μM Shield-1 for 18 hrs.

Figure 6

Figure 6. DD-Myc-FER1^{FL} overexpression results in apical microneme migration and a burst of
signaling-independent microneme release. (**a**) Co-staining of α-Myc and α-FER1 sera by IFA.
Total projection of deconvolved image is shown. (**b**) MIC2 secretion in the supernatant was
assessed by western. Parasites were induced 6 hrs with 1 μM Shield-1 and harvested by physical
release from the host cell. 10%: 10% of total lysate; const.: 1 hr constitutive secretion at 37ºC;
1% ethanol; 2 μM A23187; 500 μM propranolol. DMSO is the vehicle control for A23187.

Induced secretion for 5 min at 37ºC. Bottom panel represents quantified secretion normalized to

the GRA1 signal and to the 10% loading control. n=3+std. Parent line controls in **Suppl Fig S2**.

(**c**) Immunofluorescence of DD-Myc-FER1^{FL} co-stained with α-MIC5/α-MIC8 and α-FER1.

The Rab5A/C-dependent sub-population becomes scattered upon phenotype induction, although

an extreme apical focal point remains. Arrowheads mark microneme signal enriched at the basal

end of the parasites. (**d**) Pan-expansion microscopy of DD-YFP-FER1^{FL} parasites induced for

18hrs with 1 µM Shield-1. Compiled stack of 10 slices is shown. Arrowheads indicate the radial

set of micronemes. (**e**) IEM of DD-Myc-FER1^{FL} overexpressing parasites stained with MIC2

antibody (10 nm gold particles) induced for 16 hrs with 1 µM Shield-1. (**f**) Quantification of

MIC2 protein expression levels assessed by western blot across parent line and mutants using α-

tubulin as standard. Quantification was performed using Image Lab software by BioRad.

n=3±std. Two-way ANOVA, ** p-value< 0.005. See **Suppl Fig S5** for all western blots. (**g**)

Time course of Shield-1 induced egress. Parasites grown for 30 hrs in fibroblasts and induced

with 1 µM Shield-1 for the indicated times, fixed with 100% methanol and stained with α-Myc,

α-MIC8 and DAPI. Intact vacuoles per field were scored as proxy for egress. n=3±std. (**h**)

Modified red-green invasion assay. Top schematic shows parasites were allowed to invade for 1

hr at 37ºC followed by a wash to remove non-invaded parasites and 2 hrs 1 µM Shield-1

induction followed by the red-green invasion assay. Lower panel shows the relative number of

parasites observed outside the host cell. n=4+std. ANOVA with post-hoc Tukey HSD.

Figure 7

Figure 7. FER1-mediated trafficking of micronemes relies on predicted Ca^{2+}-binding residues. (**a**) Sequence analysis of the five conserved key positions (#1-5) in the C2A-F domain loops interfacing with Ca^{2+}[86]. D or E residues (red) stabilize Ca^{2+}, N or Q (bleu) are expected to support phospholipid binding. Positions 2, 3, and 4 shaded in grey are more strongly conserved than positions 1 and 5. Yellow highlighted residues were mutated to A to abolish the predicted Ca^{2+}-binding capacity in the C2D domain resulting in the mutant FER1$^{(3DA)}$ allele. (**b**) Models of the C2D domain loops and putative Ca^{2+} binding capacity. Yellow highlighted residues as in panel A. (**c, d**) Transient overexpression DD-YFP-FER1ΔTM$^{(3DA)}$ DD-YFP-FER1$^{FL-(3DA)}$, respectively, co-stained for YFP, MIC2 and MIC8, as indicated. Each panel contains transfected (YFP positive) and non-transfected (YFP negative) parasite examples. Following electroporation, parasites were allowed to invade for 2 hrs before 1 µM Shield-1 induction for 18 hrs. (**e-h**) Quantification of MIC2 (**e, g**) and MIC8 (**f, h**) signal localization for wild-type and mutant allele overexpression as indicated. Fluorescence was quantified in the four regions as illustrated in the inserted IFA image in panel E. At least 20 parasites were imaged for each condition. Open symbols represent outlier data points which were excluded from statistical analysis. Error bars display average ±std. One way ANOVA: * $p < 0.05$, ** $p < 0.01$, *** $p <$ 0.001, **** $p < 0.0001$. Exact numbers are provided as Data source.

1042 **Figure 8**

1043

1044

1045

Figure 8. Working model of FER1 mediated microneme events. FER1 mediates the following

trafficking steps: 1. from the ELC to the subpellicular cortex (actin dependent?); 2. microneme

migration to the apical tip (along the subpellicular microtubules?); 3. microneme membrane

fusion with the plasma membrane; 4. recycling of micronemes into the budding daughters (actin

dependent); 5. differential trafficking dynamics of the two micronemes populations differentiated

by their involvement of Rab5A/C in their biogenesis. We propose that the radial micronemes

represent a readily releasable vesicle pool primed for secretion. ELC: endosome-like

compartment; IMC: inner membrane complex; MT: microtubules; PLV: plant like vacuole or

VAC.

Supplementary information

Supplementary Movie S1. Time lapse experiment of DD-YFP-FER1ΔTM parasites co-

transfected with MIC3-mCherry to track the micronemes. At t=0, 2 μM Shield-1 was added and

images collected every 5 minutes. Scale bars 80 μm.

Supplementary movie S2. Time lapse experiment of DD-YFP-FER1ΔTM parasites co-

transfected with IMC3-mCherry to track cell division status. At t=0, 2 μM Shield-1 was added

and images collected every 5 minutes. Scale bars 80 μm.

Supplementary Figure S1

Supplementary Figure S1. Calibration of Golgi and endosome markers. (**a**) Co-transfected
parasites with the markers as indicated. (**b**) Schematic representation of collected insights.

17 **Supplementary Figure S2**

Supplementary Figure S2. Egress assays and microneme secretion. (**a**) Schematic overview of

signaling toward microneme secretion and egress, highlighting pharmacological agents acting on

various events. (**b**) Induced egress assays. Parasites grown for 30 hrs in fibroblasts and induced

for 18 hrs with 1 µM Shield-1 were triggered for egress with the pharmacologicals as indicated,

fixed and stained with α-IMC3 (parasite cortex) and α-GRA3 (PVM) sera and scored for status

of vacuole permeabilization and/or egress. n=3±std. Parasites were not under Shield-1 pressure

during the 5 min pharmacological incubation. (**c**) Assessment of microneme secretion by western

blotting by detecting MIC2 release in the supernatant under various triggers. Parasites were

induced 18 hrs with 1 µM Shield-1 and harvested by physical release from the host cell. 10%:

10% of total lysate; const.: 1 hr constitutive secretion at 37ºC in absence of secretagogue; 1%

ethanol; 2 µM A23187; 500 µM propranolol. DMSO is the vehicle control for A23187. Induced

secretion for 5 min at 37ºC. Bottom of panel represents quantified secretion normalized to the

GRA1 signal and to the 10% loading control for each condition. n=3±std.

Supplementary Figure S3

Supplementary Figure S3. Additional localization data in the DD-YFP-FER1 Δ TM dominant

negative mutant. (**a**) Additional microneme proteins in the MIC2 group. (**b**) Additional protein in

41   the MIC3/5/8 group. (**c**) NHE3 is a specific marker of the PLV compartment[1]. Parasites were

42   treated with 1 µM Shield-1 for 18 hrs. (**d**) IEM with MIC2 antibody (10 nm gold particles) of

43   DD-YFP-FER1ΔTM overexpressing parasites induced for 7 hrs with 3 µM Shield-1. Dotted

44   circles mark atypical microneme accumulations in the cytosol.

Supplementary Figure S4

Supplementary Figure S4. Select panels from time lapse experiment of DD-YFP-FER1ΔTM

parasites co-transfected with IMC3-mCherry to track cell division. At t=0, 2 µM Shield-1 was

added. The first time point at which the piling up of YFP could be convincingly observed is

marked with a yellow asterisk. Panels from supplementary movie S2.

Supplementary Figure S5

Supplementary Figure S5. (**a**) IEM of DD-Myc-FER1^{FL} overexpressing parasites stained with

MIC2 antibody (10 nm gold particles) induced for 16 hrs with 1 µM Shield-1. MIC signal

localizes to apically accumulated micronemes which display a stretched or extended

morphology. (**b, c**) TEMs of extracellular DD-Myc-FER1^{FL} overexpressing parasites induced for

3 hrs with 2 µM Shield-1. Cross section in **b** shows the extended feature or the densely packed

microneme organelles, whereas **c** shows the radial micronemes just below the conoid, again with

a slightly stretched or extended appearance. Accolade brackets labeled 'C' mark the conoid.

Supplementary Figure S6

Supplementary Figure S6. Western blot used to quantify the amount of MIC2 in each parasite

strain and under the conditions as indicated. Blots were probed simultaneously with MIC2 and

α-tubulin (loading control) antisera.

Supplementary Figure S7

74
75
76
Supplementary Figure S7. Sortilin_SmOLLAS/RHΔku80 generation and PCR validation. (**a**)

Schematic representation of the one step strategy that simultaneously tags sortilin at the C

terminus with SmOLLAS tag and introduces the hxgprt selectable marker. Double homologous

recombination is facilitated by 30 bp homologous flanks (highlighted in bold) and a

CRISPR/Cas9 generated dsDNA break. (**b**) Diagnostic PCR reactions as indicated on parental

(RHΔKu80), and clonal strain (mutant), and a no DNA negative control (control). Primer pair

localizations are marked in panel a using the following primer pairs: PCR 1: primers 6068 and

6069; PCR 2: primers 6068 and 6070; (see **Supplementary Table S1** for primer sequences).

Expected lengths are respectively: 1225 and 2371 bp.

 Supplementary Table S1. Description of oligonucleotides used. Restriction enzyme sites
87 underlined.
88
89 *Cloning of tub-YFP-TM(FER1)*
90 4786   FER1-TM-F(Gib)   GCATGGACGAGCTGTACAAGCCTAGGAGCGCCATTATGAGTGGGCTG
91 4788   3'dhfr-Not-R(Gib)   CGAATTGGAGCTCCACCGCGGTGGCGGCCGC
92
93 *Cloning of tub-DD-YFP-FER1^{FL}/sagCAT*
94 1573   YFP-109420_F   GGTCCCCTAGGATGGCGGCGAAGGCGATC
95 1574   YFP-109420_R   GGTCCCCCGGGTTATTTGAGGAGAAACATAACAAAGATGACTAGCGC
96 1635   5xTy_F(avrII)   GGACCCCTAGGGAGGTCCATACTAACCAGGATCCACTTGACGGCGAGG
97 1636   DD_R(MfeI-EcoRV)   GGACCGATATCTTACAATTGTTCCAGTTTTAGAAGCTCCACACGG
98
99 *Internal primers used to for FER1 CDS Sanger sequencing*
100 2717   Fer1-RACE-F   CGCACCAACGAGAGGCGCTCGCAGCTA
101 2748   FER1_SeqF2   ATGCGTTCCCTGTTTCGAT
102 2749   FER1_SeqF3   GTACAGCCTCCGCTCTCTTG
103 2750   FER1_SeqF4   TGTCTGTGTCTCTCCCCAAA
104
105 *Cloning of tub-DD-YFP-FER1ΔTM/sagCAT*
106 1651   FLPnoTM(NheI)_F   GGTCCGCTAGCCCTGAAGAGGAAACCTACG
107 1652   FLPnoTM(XmaI)_R   GGTCCCCCGGGTTACGTCCATTTCAGCCCACTCATAATGGC
108
109 *Site Directed Mutagenesis of C2D*
110 4833   SDM-FER1-1-2-F   TCCGGCTCCCTACATCCAAGTGGAACTG
111 4834   SDM-FER1-1-2-R   AACACGGCGTAGCCGACAGCTGGGAG
112 5892   SDM-FER1-3-F   CCGTCGTCCTCGGCTCTGTTCAGT
113 5893   SDM-FER1-3-R   GCCGACGATAGGAGGGAGTTCGAT
114
115 *6His-FER1 for antiserum generation*
116 2787   Ava-LIC-Fer1-F   GGGTCCTGGTTCGATGTCGACGAAGGAAGGCAACATTC
117 2903   Ava-LIC-Fer1-R   CTTGTTCGTGCTGTTTATTAGAGTTGCTCAGGCGTGAAGC
118
119 *Cloning of pmic3-MIC3-Cherry-DHFR*
120 4864   gibpMIC3-MIC3-F   GGAGATGACGAGACGTTTAAACGAGGAAAGTGGGGGGGGGGGG
121 4865   gibpMIC3-MIC3-R   GTTATCCTCCTCGCCCTTGCTCACCCTAGGGAGACTCGCTGGGTACCTTCGTC
122
123 *Cloning of SORTLR_OLLAS/HX*
124 6013   pU6_SORTLR   TCTGGTAAATGGGGATGTCAAGTTGTAGGAGCTTCCGCCGAAGAGTTTTAGA-
125   GCTAGAAATAGCAAGTT
126 6014   SORTLR-Ollas-F   GATAACGTCGAACTTCTTAAACGTCTCCTAGGATGGAGCAAAAACTGATATC-
127   GGAG
6015   SORTLR-Ollas-R   TTCAGAGACGACACACGAGAAAGACAGGATATCATCCTGCAAGTGCATAGAA
5688   OLLAS-s (PmeI)   AAGGGAACAAAAGCTGGGTACCGTTTAAACGGGTCCGGATTTGCCAATGAAC
5689   OLLAS-as   CATATGCCCGGGTTACTATTTGCCCATCAGTCTGGGTC
5690   3'hx_s   ACTGATGGGCAAATAGTAACCCGGGCATATGTAGAAAAGT
4769   dhfr-as   CCACCGCGGTGGCGGCCGCCTCCTGCAAGTGCATAGAAGGA
6068   Screen_SORTLR_Fwd   GCGGAAGATCCGGAGATCAG
6069   Screen_SORTLR_Rvs   GTGTCTGCGAGTGTCCAGAG
6070   Screen_Ollas_Rvs   GCCCGAGGCTCATGGGGAAG
136
137
138
139

140 **Supplementary Table S2.** Description of plasmids used.

Plasmid	Source and reference
mAID-3xMyc-3'hxgprt-5'dhfr-HXGPRT-3'dhfr	in house[2]
pCAG_mRuby2_smFP OLLAS	Addgene[3]
ptub-DD-YFP-FER1ΔTM/sagCAT	this paper
ptub-DD-Myc-FER1ΔTM/sagCAT	this paper
ptub-DD-Myc-FER1ΔTM$^{(3DA)}$/sagCAT	this paper
ptub-DD-Myc-FER1^{FL}/sagCAT	this paper
ptub-DD-Myc-FER1$^{FL\,(3DA)}$/sagCAT	this paper
ptub-YFP-YFP(MCS)/sagCAT	in house[4]
ptub-YFP-TM^{FER1}/sagCAT	this paper
ptub-IMC3-mCherryRFP/sagCAT	in house[5]
pmic3-MIC3-mCherryRFP/hxgprtDHFR	this paper
ptub-GalNAc-YFP/sagCAT	David S. Roos[6]
ptub-GRASP55-RFP/sagCAT	David S. Roos[7]
ptub-NAGTI-YFP/sagCAT	David S. Roos[6,7]
Rab5A-HA OE Transient overexpression	Sabrina Marion[8]
Rab7-HA OE Transient overexpression	Sabrina Marion[8]

143 **Supplementary Table S3.** Description of antibody and antisera used.

Name (monoclonal)	Species	Dilution (IFA)	Dilution (western)	Source and reference
hCentrin2	rabbit	1:1000		Iain Cheeseman, unpublished
DrpB	rat	1:2000		Peter Bradley[9]
HA (3F10)	rat	1:3000		Roche
GFP	rabbit	1:500		Torrey Pines Biolabs
GRA1 (TG17.43)	mouse	1:500	1:15000	BioVision.com[10]
GRA3 (T62ch)	mouse	1:500		Maryse Lebrun, Jean-François Dubremetz[11]
FER1	guinea pig	1:1000	1:1000	This paper
IMC3	rat	1:2000	1:1000	In house[5]
cMyc (9E10)	mouse	1:50	1:50	Santa Cruz Biotech
MIC2 (6D10)	mouse	1:2000	1:8000	David Sibley[12]
MIC2	rabbit	1:1000	1:8000	David Sibley[12]
MIC3	rabbit	1:100	1:400	Jean-François Dubremetz[13]
MIC5	rabbit	1:500	1:10000	Vern Carruthers[14]
MIC8	rabbit	1:1000		Dominique Soldati[15]
MIC10	rabbit	1:500	1:15000	Vern Carruthers[16]
M2AP	rabbit	1:1000	1:1000	Vern Carruthers[17]
NHE3	guinea pig	1:1500		Gustavo Arrizabalaga[1]
OLLAS	rat	1:1000		Invitrogen
proM2AP	rabbit	1:1000	1:1000	Vern Carruthers[17]
proMIC5	rabbit	1:500		Vern Carruthers[18]
proROP4	rabbit	1:750		Gary Ward[19]
ROP2,3,4	mouse	1:250		Maryse Lebrun[20]
SAG1 (DG52)	mouse	1:500		Jeroen Saeij[5,21]
SAG1 (T41E5)	mouse & rabbit	1:500		Jean-François Dubremetz[22]
SERCA	mouse	1:5000		David Sibley[23]
α-tubulin (12G10)	mouse	1:2000	1:100	Developmental Studies Hybridoma Bank[24]
Tg-β-tubulin	rabbit	1:500		Naomi Morrissette[25]
VP1	rabbit	1:4000		Silvia Moreno[26]

146
147

148   **References**
149
150   1   Francia, M. E. *et al.* A Toxoplasma gondii protein with homology to intracellular type
151   Na(+)/H(+) exchangers is important for osmoregulation and invasion. *Exp Cell Res*,
152   doi:S0014-4827(11)00125-X [pii] 10.1016/j.yexcr.2011.03.020 (2011).
153   2   Jingjing Lou, Y. R., Argenis Arriojas, David Degras, Caroline D. Keroack, Manoj T.
154   Duraisingh, Kourosh Zarringhalam, Marc-Jan Gubbels. Single cell expression and
155   chromatin access of the Toxoplasma gondii lytic cycle identifies AP2XII-8 as an essential
156   pivotal controller of a ribosome regulon. *BioRXiv* (2023).
157   3   Viswanathan, S. *et al.* High-performance probes for light and electron microscopy. *Nat*
158   *Methods* **12**, 568-576, doi:10.1038/nmeth.3365 (2015).
159   4   Gubbels, M. J., Li, C. & Striepen, B. High-throughput growth assay for Toxoplasma
160   gondii using yellow fluorescent protein. *Antimicrob Agents Chemother* **47**, 309-316
161   (2003).
162   5   Anderson-White, B. R. *et al.* A family of intermediate filament-like proteins is
163   sequentially assembled into the cytoskeleton of Toxoplasma gondii. *Cell Microbiol* **13**,
164   18-31 (2011).
165   6   Wojczyk, B. S. *et al.* cDNA cloning and expression of UDP-N-acetyl-D-
166   galactosamine:polypeptide N-acetylgalactosaminyltransferase T1 from Toxoplasma
167   gondii. *Mol Biochem Parasitol* **131**, 93-107, doi:10.1016/s0166-6851(03)00196-8 (2003).
168   7   Hartmann, J. *et al.* Golgi and centrosome cycles in Toxoplasma gondii. *Mol Biochem*
169   *Parasitol* **145**, 125-127 (2006).
170   8   Venugopal, K. *et al.* Dual role of the Toxoplasma gondii clathrin adaptor AP1 in the
171   sorting of rhoptry and microneme proteins and in parasite division. *PLoS Pathog* **13**,
172   e1006331, doi:10.1371/journal.ppat.1006331 (2017).
173   9   Breinich, M. S. *et al.* A Dynamin Is Required for the Biogenesis of Secretory Organelles
174   in Toxoplasma gondii. *Curr Biol* **19**, 277-286, doi:S0960-9822(09)00611-3 [pii]
175   10.1016/j.cub.2009.01.039 (2009).
176   10   Cesbron-Delauw, M. F. *et al.* Molecular characterization of a 23-kilodalton major antigen
177   secreted by Toxoplasma gondii. *Proc Natl Acad Sci U S A* **86**, 7537-7541 (1989).
178   11   Dubremetz, J. F., Achbarou, A., Bermudes, D. & Joiner, K. A. Kinetics and pattern of
179   organelle exocytosis during Toxoplasma gondii/host-cell interaction. *Parasitol Res* **79**,
180   402-408 (1993).
181   12   Wan, K. L., Carruthers, V. B., Sibley, L. D. & Ajioka, J. W. Molecular characterisation
182   of an expressed sequence tag locus of Toxoplasma gondii encoding the micronemal
183   protein MIC2. *Mol Biochem Parasitol* **84**, 203-214 (1997).
184   13   Garcia-Reguet, N. *et al.* The microneme protein MIC3 of Toxoplasma gondii is a
185   secretory adhesin that binds to both the surface of the host cells and the surface of the
186   parasite. *Cell Microbiol* **2**, 353-364 (2000).
187   14   Brydges, S. D. *et al.* Targeted deletion of MIC5 enhances trimming proteolysis of
188   Toxoplasma invasion proteins. *Eukaryot Cell* **5**, 2174-2183, doi:EC.00163-06 [pii]
189   10.1128/EC.00163-06 (2006).
190   15   Meissner, M. *et al.* A family of transmembrane microneme proteins of Toxoplasma
191   gondii contain EGF-like domains and function as escorters. *J Cell Sci* **115**, 563-574
192   (2002).

16   Hoff, E. F. *et al.* Toxoplasma gondii: molecular cloning and characterization of a novel
18-kDa secretory antigen, TgMIC10. *Exp Parasitol* **97**, 77-88,
doi:10.1006/expr.2000.4585 (2001).
17   Harper, J. M. *et al.* A cleavable propeptide influences Toxoplasma infection by
facilitating the trafficking and secretion of the TgMIC2-M2AP invasion complex.
Molecular biology of the cell **17**, 4551-4563, doi:10.1091/mbc.E06-01-0064 (2006).
18   Herm-Gotz, A. *et al.* Rapid control of protein level in the apicomplexan Toxoplasma
gondii. *Nat Methods* **4**, 1003-1005 (2007).
19   Carey, K. L., Jongco, A. M., Kim, K. & Ward, G. E. The Toxoplasma gondii rhoptry
protein ROP4 is secreted into the parasitophorous vacuole and becomes phosphorylated
in infected cells. *Eukaryot Cell* **3**, 1320-1330 (2004).
20   Sadak, A., Taghy, Z., Fortier, B. & Dubremetz, J. F. Characterization of a Family of
Rhoptry Proteins of Toxoplasma-Gondii. *Molecular and biochemical parasitology* **29**,
203-211, doi:Doi 10.1016/0166-6851(88)90075-8 (1988).
21   Burg, J. L., Perelman, D., Kasper, L. H., Ware, P. L. & Boothroyd, J. C. Molecular
analysis of the gene encoding the major surface antigen of Toxoplasma gondii. *Journal of
immunology* **141**, 3584-3591 (1988).
22   Couvreur, G., Sadak, A., Fortier, B. & Dubremetz, J. F. Surface antigens of Toxoplasma
gondii. *Parasitology* **97 (Pt 1)**, 1-10 (1988).
23   Nagamune, K., Beatty, W. L. & Sibley, L. D. Artemisinin induces calcium-dependent
protein secretion in the protozoan parasite Toxoplasma gondii. *Eukaryot Cell* **6**, 2147-
2156, doi:EC.00262-07 [pii] 10.1128/EC.00262-07 (2007).
24   Jerka-Dziadosz, M. *et al.* Cellular polarity in ciliates: persistence of global polarity in a
disorganized mutant of Tetrahymena thermophila that disrupts cytoskeletal organization.
Dev Biol **169**, 644-661 (1995).
25   Morrissette, N. S. & Sibley, L. D. Disruption of microtubules uncouples budding and
nuclear division in Toxoplasma gondii. *J Cell Sci* **115**, 1017-1025 (2002).
26   Miranda, K. *et al.* Characterization of a novel organelle in Toxoplasma gondii with
similar composition and function to the plant vacuole. *Mol Microbiol* **76**, 1358-1375,
doi:MMI7165 [pii] 10.1111/j.1365-2958.2010.07165.x (2010).
223