Efficient Dilution-to-Extinction isolation of novel virus-host model systems for

fastidious heterotrophic bacteria.

Holger H. Buchholz[1], Michelle Michelsen[1], Luis M. Bolaños[1], Emily Browne[1], Michael J. Allen[1,2],

Ben Temperton[1*]

[1]University of Exeter, School of Biosciences, Exeter, UK;

[2]Plymouth Marine Laboratory, Plymouth, UK

*Corresponding Author

b.temperton@exeter.ac.uk

Running Title: Cultivation of viruses of fastidious bacteria

Keywords: bacteriophages, high-throughput culturing, marine microbiology, virus-host systems,

Western English Channel

Abstract

Microbes and their associated viruses are key drivers of biogeochemical processes in marine and soil biomes. While viruses of phototrophic cyanobacteria are well-represented in model systems, challenges of isolating marine microbial heterotrophs and their viruses have hampered experimental approaches to quantify the importance of viruses in nutrient recycling. A resurgence in cultivation efforts has improved the availability of fastidious bacteria for hypothesis testing, but this has not been matched by similar efforts to cultivate their associated bacteriophages. Here, we describe a high-throughput method for isolating important virus-host systems for fastidious heterotrophic bacteria that couples advances in culturing of hosts with sequential enrichment and isolation of associated phages. Applied to six monthly samples from the Western English Channel, we first isolated one new member of the globally dominant bacterial SAR11 clade and three new members of the methylotrophic bacterial clade OM43. We used these as bait to isolate 117 new phages including the first known siphophage infecting SAR11, and the first isolated phage for OM43. Genomic analyses of 13 novel viruses revealed representatives of three new viral genera, and infection assays showed that the viruses infecting SAR11 have ecotype-specific host-ranges. Similar to the abundant human-associated phage ɸCrAss001, infection dynamics within the majority of isolates suggested either prevalent lysogeny or chronic infection, despite a lack of associated genes; or host phenotypic bistability with lysis putatively maintained within a susceptible subpopulation. Broader representation of important virus-host systems in culture collections and genomic databases will improve both our understanding of virus-host interactions, and accuracy of computational approaches to evaluate ecological patterns from metagenomic data.

Introduction

It is estimated that viral predation kills ~15% of bacterial cells in marine surface water each day [1] and is a major contributor to nutrient recycling via the viral shunt, where marine viruses make cell-bound nutrients available to the neighbouring microbial community through viral lysis of host cells [2, 3]. Viruses are key players in the modulation of carbon fluxes across the oceans (150 Gt/yr), increasing particle aggregation and sinking to depth [2, 4], and accounting for 89% of the variance in carbon export from surface water to the deep ocean [5]. Viruses alter host metabolism through auxiliary metabolic genes (AMGs), increasing and altering the cellular carbon intake of infected cells [6]. Virus-host interactions also increase co-evolutionary rates of both predator and prey via Red Queen dynamics [7, 8]. While recent metagenomic advances have provided major insight into global viral diversity and abundance [9–12], mechanistic understanding of virus-host interactions in ecologically important taxa is reliant on experimental co-culturing of model systems. In cyanobacteria, such systems have shown that viruses increase duration of photosynthetic function [13] and can inhibit CO_2 fixation, providing direct evidence that viruses of abundant phototrophs play an important role in nutrient cycling and global carbon budgets [14]. Furthermore, isolation of new viruses provides complete or near-complete viral genomes, with concrete evidence of known hosts. Such systems are critical to the development, ground-truthing and application of computational methods to identify and classify viral genomes in metagenomic data (e.g. VirSorter [15]; VirFinder [16]; MARVEL [17]); quantify boundaries for viral populations [12, 18] and genera (VConTACT2 [19]); understand the importance of AMGs in altering nutrient flux in natural communities [20] and to predict host ranges of uncultured viruses in metagenomic data (e.g. WIsH [21, 22]).

Viruses of primary producers, such as cyanophages are both well represented with model systems and well-studied in the laboratory. In contrast, virus-host model systems for similarly important and abundant marine heterotrophic bacteria are rare. Isolated viruses infecting heterotrophs are heavily biased towards those with fast-growing, copiotrophic hosts that grow readily on solid agar, enabling the use of plaque assays for viral isolation. Such systems are not representative of the vast majority of heterotrophs in nutrient-limited soil and aquatic environments, which are dominated by slow-growing, oligotrophic taxa with few regulatory mechanisms and complex auxotrophies that limit growth on solid media [23–26]. Advances in Dilution-to-Extinction culturing of ecologically important hosts have enabled the cultivation of many fastidious bacterial taxa that are not amenable to growth on solid media from soil [27], marine [28, 29] and freshwater environments [30]. Without plaque assays to facilitate isolation and purification of viral isolates, cultivation of viruses infecting fastidious taxa in liquid media is challenging, and further exacerbated by the slow growth rates and complex nutrient requirements of their hosts. Paucity of such model systems introduces significant bias in our understanding of viral influence on global carbon biogeochemical cycles. Therefore, it is important that the efforts to isolate heterotrophic bacterial taxa for experimentation and synthetic ecology is matched by efforts to isolate their associated viruses.

Here, we adapted recent advances in Dilution-to-Extinction culturing of hosts [29], and protocols to isolate viruses from liquid media [31] to improve efficiency of cultivating novel virus-host systems for fastidious taxa: First, we selected the ecologically significant SAR11 and OM43 heterotrophic marine clades as models for viral isolation; Second, we used sequential enrichment of viruses from natural communities on target hosts to improve rates of viral isolation [32]; Third, we replaced the requirement for time-intensive epifluorescent microscopy with identification of putative viral infection by comparing infected and uninfected hosts by flow cytometry, followed by confirmation using Transmission Electron Microscopy (TEM). These clades are abundant and important to global

carbon biogeochemistry [23, 33–37], but little is known about their associated viruses. In the case of

viruses infecting SAR11, two challenges limit our ability to evaluate host-virus ecology in natural

communities: (1) Assembly of abundant and microdiverse genomes from viral metagenomes presents

a challenge to short-read assembly methods, resulting in underrepresentation in subsequent datasets

[12, 18]. This was demonstrated in the successful isolation of the first known pelagiphages by

culturing, including the globally dominant HTVC010P, which, prior to its isolation, was entirely

missed in marine viromes [38]; (2) poor representation of viral taxa in databases limits our capacity

to accurately train machine learning approaches for *in silico* host prediction [12, 39, 40]. This results

in either a lack of host information for abundant viral contigs, or worse, incorrect assignment of host

to viral contigs, confounding ecological interpretation of data. Some important taxa, such as the

OM43 clade, which plays an important role in oxidation of volatile carbon associated with

phytoplankton blooms [33, 35, 41], lack any isolated viruses with experimentally confirmed hosts. In

addition, both SAR11 and OM43 represent model organisms for genome streamlining as a result of

nutrient-limited selection [42]. The effect of genome minimalism on viral infection dynamics is

poorly understood, but critical to evaluating the impact of predator-prey dynamics on global marine

carbon budgets. In this study, novel SAR11 and OM43 representatives from the Western English

Channel were isolated and used as bait to isolate associated viruses. We increased the initial

concentration of viruses in natural seawater samples by tangential flow filtration, followed by

inoculation of cultures and one to three rounds of sequential enrichment on target hosts in 96-well

plates. This yielded 117 viral isolates from 218 inoculated cultures from seven monthly water samples

(September 2018 to July 2019). A subsample of putative viral isolates for both clades were sequenced,

providing 13 novel viral genomes, including the first known siphovirus to infect SAR11 and the first

known virus-host model for OM43.

Results and Discussion

113

114 **Isolation of a novel SAR11 strain and three new OM43 strains from the Western English**

115 **Channel to use as bait for phage isolation**

116 Dilution-to-Extinction culturing for host taxa using natural seawater-based medium was performed

117 from a water sample collected in September 2017 from the Western English Channel, and yielded the

118 first SAR11 strain (named H2P3α) and the first three OM43 strains (named C6P1, D12P1, H5P1)

119 from this region. The full-length 16S rRNA gene of *Pelagibacter sp.* H2P3α was 100% identical to

120 that of the warm water SAR11 ecotype *Pelagibacter bermudensis* HTCC7211 (subclade 1a.3) and

121 was considered to be a local variant [43] (Supplementary Fig. 1A). All three novel OM43 isolates

122 were most closely related to *Methylophilales sp.* HTCC2181, a streamlined member of the OM43

123 clade with a 1.3 Mbp genome, isolated from surface water of the North Eastern Pacific [44] (C6P1

124 96.17%, D12P1 96.62%, H5P1 97.79% nucleotide identity across the full 16S rRNA gene)

125 (Supplementary Fig. 1B). The average nucleotide identity of the 16S rRNA gene of isolates CP61,

126 D12P1 and H5P1 to each other was ~98.46% (Supplementary Table 1), suggesting they are

127 representatives of the same genus [45].

128

129 **An efficient, low-cost method of isolating new viruses yielded 117 new viral isolates for SAR11**

130 **and OM43 taxa**

131 Using the four new hosts from above and established SAR11 isolates *Pelagibacter ubique* HTCC1062

132 (subclade 1a.1) and HTCC7211 (subclade 1a.3), we developed an optimised viral isolation pipeline

133 (Fig. 1) and applied it to six monthly water samples from the Western English Channel, taken between

134 September 2018 to April 2019 (Supplementary Table 2). Each month, we concentrated a natural viral

135 community with tangential flow filtration and used it to inoculate one to two 96-well Teflon plates

136 containing host cultures at ~10^6 cells·mL^{-1}. Plates were monitored by flow cytometry and growth of

137 putatively infected cultures was compared to those of unamended controls over the course of ~2

weeks to account for the slow growth rates of SAR11 and OM43 [35, 46]. An additional sample was

taken in July 2019 to attempt viral isolation on OM43 strains D12P1 and H5P1. Out of a total of 218

cultures amended with concentrated viral populations, 117 viruses were isolated, purified and still

infective after at least three passages, with repeatable differences observed in cytograms of infected

cultures and control cultures (Supplementary Fig. 2-5). This represents an overall isolation efficiency

of 53% and an average yield of 18 viruses per environmental sample (Fig. 2). For 90% of inoculated

SAR11 cultures (94 out of 105) we observed evidence of viral infection, and the putative viral isolate

could be propagated and purified, fulfilling Koch's postulates for confirming a pathogenic agent. For

OM43, 23 out of 113 (20%) inoculated cultures yielded positive infections that could be similarly

propagated. All viral isolations required between one and three rounds of virus enrichment

(Supplementary Table 3) before changes in host growth curves between infected and uninfected

cultures could be observed. This suggests putatively rare viruses can be enriched within three rounds

to a level at which infection can be observed on a flow cytometer.

New viruses represent novel viral populations and support established ANI cut-offs for

ecologically discrete viral ecotypes

Due to the rate-limiting step of culturing sufficient biomass for extraction of viral DNA, we

subselected 16 viral isolates based on availability in November 2018 across four different hosts

(HTCC1062, HTCC7211, H2P3α and H5P1) for Illumina sequencing to >30-fold coverage (Table

1). Three out of 16 sequenced samples (~19%, two from host HTCC7211, one from OM43 host

H5P1) failed to assemble into single viral contigs, in line with previously reported failure rates of 18-

39% for phages of *Escherichia coli* and *Salmonella spp.* [47]. For 11 of the remaining 13 samples (12

from SAR11 hosts and one from OM43), each individual sequence assembly was identified as a

complete viral genome by VirSorter [15] and 95-100% complete using CheckV [48]. All assemblies

yielded a single viral contig (categories 1 or 2, >15kbp) per sample, indicating that our purification

process was effective in recovering pure viral isolates. Interestingly, all viral isolate genomes were classified by VirSorter as Category 2 ("likely virus") rather than Category 1 ("most confident"), despite being complete - indicating either a lack of viral hallmark genes or a lack of enrichment of viral genes on the contigs. This finding matches our own observations for other isolated viruses infecting SAR11 (data not shown), and suggests VirSorter classification of pelagiphages is conservative. Viral isolates were named in accordance with current ICTV guidelines for prokaryotic virus taxonomy [49], with names selected from Norse mythology and folklore, and contemporary culture (Table 1).

Viral populations are defined as discrete ecological and evolutionary units that form non-overlapping clouds of sequence space, and previous work in cyanophage populations have shown that viral populations can be delineated into populations using an average nucleotide identity (ANI) cutoff of 95% [50]. Pairwise ANI was calculated between the thirteen successfully sequenced viral genomes from this study and 85 other known or putative pelagiphages [39, 51, 52]. Pairwise ANI ranged between 77.5-100%, with a discrete distribution between 96.4-100.0% (Supplementary Fig. 6, Supplementary Table 4). This is in agreement with previous work in cyanophages [50, 53] and supports the broad use of proposed boundary cutoffs to define viral populations within viral metagenomic assemblies [10, 11]. At the proposed ANI cutoffs of 95% over 85% length [18], our 13 new viruses clustered into six viral populations, ranging from singletons to a viral population with four members (Table 1). Phages within the same populations were all isolated from the same environmental sample and on the same host, in agreement with their classification as discrete ecological and evolutionary units. All viruses isolated in this study formed their own exclusive viral populations, with no representatives from either known isolates [38, 52] or fosmid-derived [39] genomes from other studies, indicating that a high degree of viral population diversity remains to be discovered in the Western English Channel and beyond.

Current pelagiphage isolates can be organised into five distinct phylogenetic clades

To evaluate isolate diversity at higher taxonomic organisation, we picked one representative from each of our six viral populations and compared them to previous isolates and fosmid-derived phage sequences using three approaches: First, phylogenetic analyses were performed based on conserved genes in known pelagiphage isolates and closely-related taxa from viral metagenomic surveys [11, 12]; Second, raw hypergeometric probability of shared gene content was calculated (to capture broader relationships and account for genomic mosaicism) [54]; Third, genomes were organised into ICTV-recognized genera using vConTACT2. vConTACT2 initially derives viral clusters using a hypergeometric approach, with subsequent refinement with Euclidean-distance-based hierarchical clustering to split mismatched, 'lumped' viral clusters [19]. All three approaches were congruent - clustering on probability of shared gene content organised pelagiphage genomes into four main clusters and numerous singleton genomes (Fig. 3). This was broadly supported by phylogenetic (Supplementary Fig. 7) and vConTACT2 (Supplementary Fig. 8) classification approaches. Cluster A contained 23 members (nine from fosmid-derived contigs [39]; eleven previously isolated pelagiphages [52, 55]) and Pelagibacter phages *Ran* (EXVC014P), *Bylgja* (EXVC010P) and *Eistla* (EXVC025P) from this study. Cluster B contained two previously isolated pelagiphages, one fosmid-derived contig and Pelagibacter phage *Eyrgjafa* (EXVC018P) from this study. All viruses in Clusters A and B were assigned to a single viral genus by vConTACT2 that also contained 12 previously isolated pelagiphages [38, 52]. Cluster C only contained fosmid-derived contigs from the Mediterranean [39], with no isolated representatives, marking it an important target for future isolation attempts. Cluster D contained eight fosmid-derived contigs, Pelagibacter phage HTVC010P, and Pelagibacter phage *Greip* (EXVC021P) from this study. Pelagibacter phage *Kolga* (EXVC016S) from this study and the only known Pelagimyophage HTVC008M from a previous study [38] fell outside the four main clusters. VConTACT2 split Cluster D, leaving Pelagibacter phages *Greip* and

Kolga as members of two singleton clusters, suggesting they are the first cultured representatives of novel viral genera and distinct from the globally ubiquitous Pelagibacter phage HTVC010P.

Novel pelagiphages are ecotype-specific and persist in the community.

Community composition analysis using 16S rRNA genes showed that during the sampling period (September 2018 to July 2019) the SAR11 contribution relative to the total number of sequences ranged from a minimum of 28.0% in July 2019 to a maximum of 42.2% in September 2018 (Supplementary Fig.9). At the clade level, SAR11 composition was relatively stable over time, except for clade II roughly doubling its relative contribution from 15.1% to about 29.6% between February and April. The SAR11 community was dominated by clade I overall throughout the sampling period. Within clade I, the ratio between SAR11 subclade Ia.1 (cold-water ecotype) and Ia.3 (warm-water ecotype) showed that the warm-water ecotype dominated from September to November as well as July (Figure 2A). During the coldest months in the Western English Channel (February to April) the cold-water ecotype became as abundant as the warm-water ecotype with roughly a 1:1 ratio. Overall, temperature ranged from 9.3 °C - 14.1 °C from October to April, when isolations were attempted on all strains (Supplementary Table 2).   HTCC1062 and HTCC7211 have specific growth rates of ~0.22 and ~0.12 divisions per day at 10 °C, respectively [46]. Our new SAR11 isolate from the Western English Channel, *Pelagibacter sp.* H2P3α, showed similar growth rates to *Pelagibacter bermudensis* HTCC7211 (Supplementary Fig. 10), with specific growth rates of ~0.10, ~0.45 and ~0.84 divisions per day at 10, 18 and 25 °C, respectively. Therefore, measured *in situ* temperatures during our sampling period were sufficient to support slow growth of warm-water ecotypes even during winter months, potentially providing sufficient prey to support a population of warm-water ecotype specific phages. This result shows that slow growth of warm water ecotypes at *in situ* temperatures is possible, supporting the finding of the 16S community analysis that this ecotype persists throughout the year.

An alternative explanation is that isolated viruses have a broad host range that encompasses both

warm- and cold-water ecotypes of the SAR11 subclade Ia. We tested the host range (Table 2) of six

pelagiphages (one of each viral population), isolated from samples in October 2018 and November

2018 (*in situ* water temperature of 14.8 °C and 14.2 °C respectively) across the three SAR11 strains

(Table 1). Pelagibacter phages *Eistla, Eyrgjafa* and *Greip* all infected cold-water ecotype HTCC1062

exclusively, while Pelagibacter phages *Ran* and *Kolga* only infected warm-water ecotypes

HTCC7211 and H2P3α. Pelagibacter phage *Bylgia* was the only virus that could infect both warm

and cold-water ecotypes. Therefore, our new pelagiphages appear to be broadly ecotype-specific,

confirming previous findings [52]. Our results suggest overall that pelagiphages persist in the water

column throughout the year in sufficient densities to be isolated by our enrichment method, despite

ecotypic specificity and fluctuations in warm- and cold-water ecotype community contributions of

SAR11 subclade 1a. If concentration and enrichment of viruses during isolation is sufficient to

successfully isolate even low abundance phages then a comprehensive library of representative phage

isolates could be generated with relatively modest sampling effort across a few locations.

Pelagibacter phages Kolga and Aegir- the first siphovirus infecting SAR11

The 25 previously known viral isolates infecting SAR11 comprise 24 podoviruses and one myovirus

[38, 52, 56]. Previous cultivation efforts for viruses of SAR11 have not isolated any siphoviruses, nor

are any known from viral metagenomic studies. We isolated and sequenced the two Pelagibacter

phages *Kolga* and *Aegir* using host H2P3α as bait (Fig. 4A-C). Transmission electron microscopy

(TEM), which showed evidence of a long tail (Fig. 4B), suggested classification as the first reported

siphoviruses infecting members of the SAR11 clade.

Pelagibacter phages *Aegir* and *Kolga* were classified as members of the same population using a

boundary cutoff of 95% ANI over 85% contig length, however, *Aegir* had a length of 18,297 bp

compared to 48,659 bp in *Kolga*, therefore we considered *Aegir* to be a partial genome of the same

viral population. *Kolga* did not share a significant number of genes with known SAR11 podoviruses

(Fig. 3), and did not cluster with other pelagiphages using hypergeometric analysis based on shared

gene content. VConTACT2 also grouped *Aegir* and *Kolga* into one cluster without any other known

viruses, suggesting it represents a novel viral genus. A number of genes found in *Kolga* were shared

with other known siphoviruses (associated with different hosts) such as the Bordetella phage LK3.

Screening of contigs from the Global Ocean Virome dataset [11] identified six contigs from various

ecological zones which shared a viral cluster with *Kolga*, but which belonged to different viral

populations based on network analysis using vConTACT2. Phylogenetic analysis of all Pelagibacter

phage TerL genes indicates that the closest known relatives to *Kolga* and *Kolga*-like contigs are

members of the Myoviridae family, supporting a classification as a distinct and novel viral group

(Fig. 4C, Supplementary Fig. 7). In *Kolga* 67% of encoded genes could not be functionally annotated,

and out of all hypothetical genes identified on *Kolga*, only three hypothetical genes were shared with

SAR11 podoviruses. In contrast, on average ~90% of genes without known function identified within

our novel SAR11 podoviruses are shared between different SAR11 podoviruses. *Kolga* possesses a

tail tip J protein (Fig. 4A), often found in phages with long non-contractile tails such as E. coli phage

λ, where it plays a role in DNA injection during cell entry and tail assembly [57]. *Kolga* also encodes

a small S21 subunit of the 30S ribosomal gene structurally similar to the ones found in Pelagibacter

phage HTVC008M, Pelagimyophage-like contigs [58], and hosts HTCC7211 and H2P3α. Encoding

ribosomal genes is a feature found in numerous myoviruses and siphoviruses [59]. The S21 gene is

involved in translation initiation and needed for the binding of mRNA [60]. Virally-encoded S21

genes may provide a competitive advantage for the phage as it could replace cellular S21 and assist

in the translation of viral transcripts. *Kolga* may need its S21 gene for shifting the translational frame,

as it has been shown that for some members of the Caudovirales the production of tail components is

dependent on programmed translational frameshifting [61]. Given the constitutive nature of gene

expression in genomically streamlined bacteria [62], genes such as S21 may also provide the virus

with a mechanism to manipulate host metabolism in the absence of typical promoters and repressors.

First methylophages for marine OM43 isolated

Isolation of novel phages for the OM43 clade yielded 23 positive infections, with efficiencies ranging

from 0% (no viruses isolated on host C6P1) to 45% on H5P1 (Fig. 2). To the best of our knowledge

these are the first reported viruses infecting members of the OM43 clade. One explanation for the

lower efficiency of isolation of phages infecting OM43 is simply one of lower host abundance

concomitant with lower phage abundance in the viral community, reducing the likelihood of infective

viruses coming into contact with susceptible and permissive cells. OM43 are closely associated with

metabolism of extracellular substrates from phytoplankton blooms [63], but have low abundance

outside of phytoplankton spring blooms [33]. Our water samples were not associated with high *in-*

situ fluorescence (used as proxy measurement for phytoplankton), and missed the April 2019 spring

bloom by about two weeks (Supplementary Fig. 11). Based on 16S community analysis, the OM43

contribution to the bacterial community (0.7%) was lowest during sampling in April 2019 and highest

in October 2018 (1.9%). Relative to all other OM43, H5P1 was the most abundant OM43 in the

Western English Channel, contributing more than half of all OM43 in November (Fig. 2B). This

could explain the higher success rate of isolating phages on host H5P1 compared to C6P1 and D12P1.

Sequencing of the first OM43 phages isolate, Methylophilales phage *Venkman* (EXVC282S),

returned a single genome 38,624 bp long (31.9% GC content), which was linear but complete

(Supplementary Table 5). *Venkman* encodes genes (Fig. 5A) with similar synteny and function to the

siphovirus P19250A (38,562 bp) that infects freshwater *Methylophilales* LD28, which are often

considered a freshwater variant of OM43 [64, 65]. Unlike the siphovirus P19250A, TEM images

indicated that OM43 phage *Venkman* had a short tail (Fig. 5B) similar to podoviruses, though it is

possible that tail structures were lost during grid preparation. *Venkman* shared a viral population with

a 23kbp contig from the Western English Channel virome assembled from short-read data

(WEC_HYBRID_01170), suggesting this viral type does not suffer from the issues of high abundance

and microdiversity that challenge assembly of some pelagiphages. Phylogenetic analysis of

concatenated TerL and exonuclease genes indicated that *Venkman* is most closely related to P19250A

and other siphoviruses infecting different Proteobacteria (Fig. 5C). However, branch support values

were low, despite numerous attempts to refine the tree with different approaches (see Supplementary

Methods). We identified a number of common phage proteins such as a capsid protein, terminase,

nucleases and tail structural proteins, and the remaining 54% of genes were hypothetical.

VConTACT2 assigned OM43 phage *Venkman* and LD28 phage P19250A to the same genus-level

cluster, therefore the OM43 phage *Venkman* may be a marine variant of the freshwater LD28

siphovirus P19250A.

Global abundance of novel isolates highlights niche-specificity and low representation in

existing datasets

To evaluate relative global abundance of our new phage isolates, existing virome datasets from the

Global Ocean Virome survey [11] and the Western English Channel [12] were randomly subsampled

to 5 million reads and competitively recruited against genomes of viral population representatives

from this study as well as previously isolated pelagiphages (see Supplementary methods). Overall,

with the exception of Pelagibacter phage HTVC010P, pelagiphages were poorly represented in

samples from temperate-tropical mesopelagic (TT-MES) and Antarctic (ANT) ecological zones, as

defined in [11] (Fig. 6). Phages isolated in this study were neither globally ubiquitous, nor abundant

in the single Western English Channel virome (with the exception of phage *Ran*). Phages *Bylgja,*

Himinglaeva, Eistla and *Eyrgjafa* did not achieve the minimum cutoff of 40% genome coverage to

be classified as present [18] in any of the viromes tested. *Ran* was the only phage isolated in this study

with representation in at least two samples from temperate-tropical epipelagic (TT-EPI) zones,

concordant with its host specificity of warm-water SAR11 ecotypes (*Pelagibacter bermudensis* HTCC7211 and H2P3α from this study (Table 2). Similarly, the Western English Channel virome was taken in September 2016, when waters are usually highly stratified after summer heating [66] and warm water ecotypes of SAR11 dominate the microbial community (Fig. 2A). Other viral populations from this study also isolated on warm water SAR11 ecotypes (phages *Kolga, Bylgja* and *Himinglaeva*) were either absent or below limits of detection in this sample. This coupled with the low global abundance of these viruses suggests that pelagiphage communities comprise few highly abundant taxa and a long rare tail. Pelagibacter phage *Greip* was detected in seven samples, six of which were Arctic (ARC) samples from a discrete region of the Arctic characterized by low nutrient ratios [11]. In three of those samples (191_SRF, 193_SRF, 196_SRF), HTVC010P was not detected and in the other three (206_SRF, 208_SRF and 209_SRF) *Greip* was 1.5 to 6.6-fold more abundant than HTVC010P, identifying *Greip* as an abundant arctic pelagiphage ecotype. The fact that *Greip* was isolated on the cold-water ecotype of SAR11 (Pelagibacter ubique HTCC1062) and does not infect either warm-water ecotype HTCC7211 or H2P3 supports the hypothesis that host niche specificity shapes the phylogeography of associated viral taxa.

In the Western English Channel viral metagenomes, OM43 phage *Venkman* was the third most abundant (2,699 RPKM) after Pelagibacter phages HTVC027P (5,023 RPKM) and HTVC010P (5,026 RPKM) identifying it as an ecologically important virus in this coastal microbial community (Fig. 6). In contrast, neither *Venkman* nor LD28 phage P19250A were identified in the majority of samples from GOV2, except in one sample (67_SRF) in the South Atlantic Ocean (257 RPKM against *Venkman*). Sample 67_SRF was also classified as a coastal biome by the authors [11]. These results coincide with previous reports that OM43, and presumably their associated viruses as a consequence of host abundance, are important in some coastal regions, but largely absent in open-ocean systems [33, 41].

Unusual host-virus dynamics are prevalent in isolated phages

Pelagibacter phages *Bylgja*, *Eyrgjafa*, *Ran* and *Eistla* all encode endonucleases and exonucleases (Supplementary Fig. 12) and cluster by shared protein content with other pelagiphages (Cluster A and B), such as e.g. HTVC011P and HTVC025P, shown previously to integrate into host genomes (Fig. 3) [52]. We therefore predict that all our phages within Clusters A and B are temperate phages. *Eyrgjafa* also encodes a tRNA-Leu, that has 85% nucleotide identity over the first 34 bases of the tRNA-Leu of its host HTCC1062, suggesting a putative integration site into the host genome [67]. To date, 16 of the 29 viruses previously isolated on SAR11 strains have either been shown to be capable of lysogeny, and/or encode genes associated with a temperate infection cycle. In contrast, viruses such as the Pelagibacter phages *Greip,* and the abundant HTVC010P in Cluster D do not possess any genes associated with lysogeny, and would therefore be classified as exclusively lytic. Viruses in Clusters A and B (putatively temperate phage), were of much lower abundance in the environment compared HTVC010P which was among the most abundant pelagiphages in the Western English Channel (Fig. 6), suggesting a possible ecological difference between two groups of viruses with different infectivity strategies.

Interestingly, growth curves of hosts infected with Pelagibacter phage *Greip* and other isolated phages deviated from the expected decay in cell abundance associated with viral lysis and previously observed in isolated pelagiphages [38] (Fig.1 Bii). The first pelagiphages HTVC010P and HTVC008M were isolated from the warm waters of the Sargasso Sea, and HTVC011P and HTVC019P were isolated from the colder waters of the Oregon Coast. All four strains were propagated on the cold-water SAR11 ecotype *P. ubique* HTCC1062. In all cases, host density was reduced from ~8×10^6 cells·mL^{-1} at T_0 to <10^6 cells·mL^{-1} over a 60-72h period. Viruses isolated from warm waters took 17% longer than those from cold water to do so [38], suggesting that suboptimal hosts reduced the rate of infection as shown in cyanophages [68]. In contrast, infection

dynamics of our isolates often resulted in host density of infected cultures growing to a steady state,

but at a lower cell density than uninfected cells. (Supplementary Video 1), irrespective of cluster,

population assignment or evidence of genes associated with temperate lifestyles. Out of 117 viruses

isolated in this study, only 16 infections reduced host abundance below their inoculum density of 10^6

cells·mL^{-1}. In 53 infections, densities of infected cells increased to within an order of magnitude of

uninfected cells (Supplementary Fig. 13), but demonstrated clear evidence of viral infection in

cytograms, TEMs and subsequent recovery of viral genomes in selected samples.

Similar patterns of infection were recently reported in the extremely abundant bacteriophage

ɸCrAss001 found in the human gut, where infection of exponentially growing cells of *B. intestinalis*

919/174 did not result in complete culture lysis, but caused a delay in stationary phase onset time and

final density, despite lacking genes associated with lysogeny. As with our study, the authors observed

that this only occurred in liquid culture, and isolation of the virus required numerous rounds of

enrichment. They postulated that the virus may cause a successful infection in only a subset of host

cells, with the remainder exhibiting alternative interactions such as pseudolysogeny or dormancy

[32]. The prevalence of similar infection dynamics in the phages isolated in this study offer two

intriguing possibilities: (i) Many of the viruses isolated in this study are either not fully lytic, but fall

somewhere on the continuum of persistence [69]. This could be controlled by genes currently lacking

a known function, with lysogeny (and associated superinfection immunity) favoured at high cell

density, which would support the Piggyback-the-Winner hypothesis [70, 71]; (ii) the steady-state of

host and virus densities observed here are an indicator of host phenotypic bistability in these

streamlined heterotrophic taxa. Viral propagation occurring in only a subset of cells could explain the

requirement of multiple rounds of enrichment before sufficient viral load is reached to be able to

observe lytic infection on the host population. Either strategy, or a combination of both, would

provide an ecological advantage of long-term stable coexistence between viruses and hosts and offer

an explanation of the paradox of stable high abundances of both predator and prey across global

oceans [12, 38, 72, 73]. Infection in a subset of the population could also explain the low lytic activity

observed in pelagiphages *in situ*, despite high host densities [74]; and the small dynamic range and

decoupled abundances of SAR11 and virioplankton in the Sargasso Sea [75]. Limited lysis of

subpopulations of hosts such as SAR11 and OM43 that specialize in the uptake of labile carbon

enriched through viral predation [76, 77] could facilitate efficient intra-population carbon recycling

and explain the limited influence of SAR11 and associated viral abundances on carbon export to the

deep ocean [5]. We propose the moniker the 'Soylent Green Hypothesis' for this mechanism, after

the 1973 cult film in which the dead are recycled into food for the living. Further investigation

leveraging our new virus-host model will provide greater insight into viral influence on ocean carbon

biogeochemistry.

In conclusion, our method coupled Dilution-to-Extinction cultivation of hosts and associated viruses,

resulting in the isolation of three new strains of OM43; a Western English Channel variant of a warm

water ecotype of SAR11; the first known methylophages for OM43; the first siphovirus infecting

SAR11, as well as eleven other viruses infecting this important marine heterotrophic clade and >100

more isolates to be sequenced and explored. The described method represents an efficient and cost-

effective strategy to isolate novel virus-host systems for experimental evaluation of co-evolutionary

dynamics of important fastidious taxa from marine and other biomes. Coupling these methods to

existing advances in host cultivation requires minimal additional effort and will provide valuable

representative genomes to improve success rates of assigning putative hosts to metagenomically-

derived viral contigs. Broader representation of model systems beyond cyanophages and viruses of

copiotrophic, *r*-strategist hosts will reduce bias in developing methods to delineate viral population

boundaries [78, 79], increasing the accuracy with which we derive ecological meaning from viral

metagenomic data. We therefore hope that this method will enable viruses to be included in the

current resurgence of cultivation efforts to better understand the biology and ecology of phages, and

the influence of the world's smallest predators on global biogeochemistry.

Methods Summary

A complete description of the materials and methods is provided in the Supplementary Information.

Four bacterial strains (*Methylophilales sp*. C6P1, D12P1 and H5P1; *Pelagibacter sp*. H2P3α) were

isolated from Western English Channel station L4 seawater samples using Dilution-to-Extinction

methods [29]. All four bacteria and two additional SAR11 strains *Pelagibacter bermudensis*

HTCC7211 and *Pelagibacter ubique* HTCC1062 were used as bait to isolate phages from six monthly

Western English Channel L4 seawater samples (50°15.00N; 4°13.00W). Briefly, water samples were

concentrated for viruses using tangential flow filtration (100k Da Hydrosart membrane) and used as

viral inoculum (10% v/v) in exponentially growing cultures of host bacteria in artificial seawater

medium [80] in 96-well Teflon plates (Radleys, UK). Cells of the resulting lysate were filtered out

(0.1 μm PVDF syringe filters) and the filtrate was used as viral inoculum in another round of isolation.

This process was repeated until viral infection could be detected by flow cytometry - comparing

cytograms and maximum density of infected cultures against uninfected cultures. Phages were

purified   by   dilution-to-extinction   methods   (detailed   protocol   available   here:

dx.doi.org/10.17504/protocols.io.c36yrd). Phage genomes were sequenced using Illumina 2x150 PE

sequencing, assembled and manually annotated as described in [81]. Phylogenetic classification of

phages was performed on concatenated shared genes using a combination of Bayesian inference trees,

maximum likelihood trees and shared-gene likelihood analyses, depending on the availability of

appropriate taxa. ICTV-recognised genera based on shared gene content were assigned with

VConTACT2 [19]. The relative abundance of novel phages in the Western English Channel [12] and

Global Ocean viromes [11] was estimated (RPKM) by competitive read recruitment of five million

randomly subsampled reads against pelagiphage and methylophage genomes from this study and
others [38, 52, 64, 82].

Data availability All reads can be found in the SRA database under BioProject number
PRJNA625644 as BioSamples SAMN14604128 - SAMN14604140. Annotated phage genomes are
deposited as GenBank submissions under accession numbers MT375519 - MT375531. Sequences
used for phylogenetic analysis are deposited under https://github.com/ViralPirates/viral-dte.

Acknowledgements We would like to thank Christian Hacker and the Bioimaging Centre of the
University of Exeter for performing the TE microscopy and imaging. We would also like to thank the
crew of the *R/V Plymouth Quest* and our collaborators at PML for collecting water samples, and the
driver Magic for delivering water samples from Plymouth to Exeter. Genome sequencing was
provided by MicrobesNG (http://www.microbesng.uk) which is supported by the BBSRC (grant
number BB/L024209/1). This project used equipment funded by the Wellcome Trust Institutional
Strategic Support Fund (WT097835MF), Wellcome Trust Multi-User Equipment Award
(WT101650MA) and BBSRC LOLA award (BB/K003240/1). Bioinformatic analyses were
conducted using the high-performance computing, ISCA, provided by the University of Exeter.

Funding information The efforts of Holger Buchholz in this work were funded by the Natural
Environment Research Council (NERC) GW4+ Doctoral Training program. Michelle Michelsen and
Ben Temperton were funded by NERC (NE/R010935/1) and by the Simons Foundation BIOS-
SCOPE program.

Conflict of interests The authors declare no conflict of interest.

References

1. Eric Wommack K, Colwell RR. Virioplankton: Viruses in Aquatic Ecosystems. *MICROBIOLOGY AND*

MOLECULAR BIOLOGY REVIEWS 2000; **64**: 69–114.

2. Weitz JS, Stock CA, Wilhelm SW, Bourouiba L, Coleman ML, Buchan A, et al. A multitrophic model

to quantify the effects of marine viruses on microbial food webs and ecosystem processes. *ISME J*

2015; **9**: 1352–1364.

3. Weitz JS, Wilhelm SW. Ocean viruses and their effects on microbial communities and biogeochemical

cycles. *F1000 Biol Rep* 2012; **4**: 17.

4. Jover LF, Effler TC, Buchan A, Wilhelm SW, Weitz JS. The elemental composition of virus particles:

implications for marine biogeochemical cycles. *Nat Rev Microbiol* 2014; **12**: 519–528.

5. Guidi L, Chaffron S, Bittner L, Eveillard D, Larhlimi A, Roux S, et al. Plankton networks driving

carbon export in the oligotrophic ocean. *Nature* 2016; **532**: 465–470.

6. Warwick-Dugdale J, Buchholz HH, Allen MJ, Temperton B. Host-hijacking and planktonic piracy: how

phages command the microbial high seas. *Virol J* 2019; **16**: 15.

7. Weitz JS, Hartman H, Levin SA. Coevolutionary arms races between bacteria and bacteriophage. *Proc*

Natl Acad Sci U S A 2005; **102**: 9535–9540.

8. Ignacio-Espinoza JC, Ahlgren NA, Fuhrman JA. Long-term stability and Red Queen-like strain

dynamics in marine viruses. *Nat Microbiol* 2020; **5**: 265–271.

9. Brum JR, Hurwitz BL, Schofield O, Ducklow HW, Sullivan MB. Seasonal time bombs: dominant

temperate viruses affect Southern Ocean microbial dynamics. *ISME J* 2016; **10**: 437–449.

10. Roux S, Brum JR, Dutilh BE, Sunagawa S, Duhaime MB, Loy A, et al. Ecogenomics and potential

biogeochemical impacts of globally abundant ocean viruses. *Nature* 2016; **537**: 689–693.

11. Gregory AC, Zayed AA, Sunagawa S, Wincker P, Sullivan MB, Temperton B, et al. Marine DNA Viral

Macro- and Microdiversity from Pole to Pole. *Cell* 2019; **177**: 1109–1123.

12. Warwick-Dugdale J, Solonenko N, Moore K, Chittick L, Gregory AC, Allen MJ, et al. Long-read viral

metagenomics captures abundant and microdiverse viral populations and their niche-defining genomic

islands. *PeerJ* 2019; **7**: e6800.

13. Mann NH, Cook A, Millard A, Bailey S, Clokie M. Bacterial photosynthesis genes in a virus. *Nature*

2003; **424**: 741.

14. Puxty RJ, Millard AD, Evans DJ, Scanlan DJ. Viruses Inhibit CO2 Fixation in the Most Abundant

Phototrophs on Earth. *Curr Biol* 2016; **26**: 1585–1589.

15. Roux S, Enault F, Hurwitz BL, Sullivan MB. VirSorter: mining viral signal from microbial genomic

data. *PeerJ* 2015; **3**: e985.

16. Ren J, Ahlgren NA, Lu YY, Fuhrman JA, Sun F. VirFinder: a novel k-mer based tool for identifying

viral sequences from assembled metagenomic data. *Microbiome* 2017; **5**: 69.

17. Amgarten D, Braga LPP, da Silva AM, Setubal JC. MARVEL, a Tool for Prediction of Bacteriophage

Sequences in Metagenomic Bins. *Front Genet* 2018; **9**: 304.

18. Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an in silico evaluation

of metagenome-enabled estimates of viral community composition and diversity. *PeerJ* 2017; **5**: e3817.

19. Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Taxonomic assignment

of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nat Biotechnol* 2019.

20. Hurwitz BL, Brum JR, Sullivan MB. Depth-stratified functional and taxonomic niche specialization in

the 'core' and 'flexible' Pacific Ocean Virome. *ISME J* 2015; **9**: 472–484.

21. Galiez C, Siebert M, Enault F, Vincent J, Söding J. WIsH: who is the host? Predicting prokaryotic hosts

from metagenomic phage contigs. *Bioinformatics* 2017; **33**: 3113–3114.

22. Ahlgren NA, Ren J, Lu YY, Fuhrman JA, Sun F. Alignment-free d_2^* oligonucleotide frequency

dissimilarity measure improves prediction of hosts from metagenomically-derived viral sequences.

Nucleic Acids Res 2017; **45**: 39–53.

23. Sun J, Steindler L, Thrash JC, Halsey KH, Smith DP, Carter AE, et al. One carbon metabolism in

SAR11 pelagic marine bacteria. *PLoS One* 2011; **6**: e23973.

24. Becker JW, Hogle SL, Rosendo K, Chisholm SW. Co-culture and biogeography of Prochlorococcus and

SAR11. *ISME J* 2019.

25. Kraemer S, Ramachandran A, Colatriano D, Lovejoy C, Walsh DA. Diversity and biogeography of

SAR11 bacteria from the Arctic Ocean. *ISME J* 2020; **14**: 79–90.

26. Robert M. Morris*, Michael S. Rappe´ *, Stephanie A. Connon*, Kevin L. Vergin*, William A.

Siebold*, Craig A. Carlson† & Stephen J. Giovannoni*. SAR11 clade dominates ocean surface

bacterioplankton communities. *Nature* 2002; **420**.

27. Bartelme RP, Custer JM, Dupont CL, Espinoza JL, Torralba M, Khalili B, et al. Influence of Substrate

Concentration on the Culturability of Heterotrophic Soil Microbes Isolated by High-Throughput

Dilution-to-Extinction Cultivation. *mSphere* 2020; **5**.

28. Connon SA, Giovannoni SJ. High-throughput methods for culturing microorganisms in very-low-

nutrient media yield diverse new marine isolates. *Appl Environ Microbiol* 2002; **68**: 3878–3885.

29. Henson MW, Pitre DM, Weckhorst JL, Celeste Lanclos V, Webber AT, Cameron Thrash J. Artificial

Seawater Media Facilitate Cultivating Members of the Microbial Majority from the Gulf of Mexico.

American Society for Microbiology 2016; **1**: 1–9.

30. Henson MW, Lanclos VC, Faircloth BC, Thrash JC. Cultivation and genomics of the first freshwater

SAR11 (LD12) isolate. *ISME J* 2018; **12**: 1846–1860.

31. Nagasaki K, Bratbak G. Isolation of viruses infecting photosynthetic and nonphotosynthetic protists.

Manual of aquatic viral ecology ASLO 2010; 92–101.

32. Shkoporov AN, Khokhlova EV, Fitzgerald CB, Stockdale SR, Draper LA, Ross RP, et al. ΦCrAss001

represents the most abundant bacteriophage family in the human gut and infects Bacteroides intestinalis.

Nat Commun 2018; **9**: 4781.

33. Morris RM, Longnecker K, Giovannoni SJ. Pirellula and OM43 are among the dominant lineages

identified in an Oregon coast diatom bloom. *Environ Microbiol* 2006; **8**: 1361–1370.

34. Ramachandran A, Walsh DA. Investigation of XoxF methanol dehydrogenases reveals new

methylotrophic bacteria in pelagic marine and freshwater ecosystems. *FEMS Microbiol Ecol* 2015; **91**.

35. Halsey KH, Carter AE, Giovannoni SJ. Synergistic metabolism of a broad range of C1 compounds in

the marine methylotrophic bacterium HTCC2181. *Environ Microbiol* 2012; **14**: 630–640.

36. Reintjes G, Arnosti C, Fuchs B, Amann R. Selfish, sharing and scavenging bacteria in the Atlantic

Ocean: a biogeographical study of bacterial substrate utilisation. *ISME J* 2019; **13**: 1119–1132.

37. Carini P, White AE, Campbell EO, Giovannoni SJ. Methane production by phosphate-starved SAR11

chemoheterotrophic marine bacteria. *Nat Commun* 2014; **5**: 4346.

38. Zhao Y, Temperton B, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC, et al. Abundant SAR11

viruses in the ocean. *Nature* 2013; **494**: 357–360.

39. Mizuno CM, Rodriguez-Valera F, Kimes NE, Ghai R. Expanding the marine virosphere using

metagenomics. *PLoS Genet* 2013; **9**: e1003987.

40. Martinez-Hernandez F, Fornas O, Lluesma Gomez M, Bolduc B, de la Cruz Peña MJ, Martínez JM, et

al. Single-virus genomics reveals hidden cosmopolitan and abundant viruses. *Nat Commun* 2017; **8**:

15892.

41. Taubert M, Grob C, Howat AM, Burns OJ, Dixon JL, Chen Y, et al. XoxF encoding an alternative

methanol dehydrogenase is widespread in coastal marine environments. *Environ Microbiol* 2015; **17**:

3937–3948.

42. Giovannoni SJ, Cameron Thrash J, Temperton B. Implications of streamlining theory for microbial

ecology. *ISME J* 2014; **8**: 1553–1565.

43. Schwalbach MS, Tripp HJ, Steindler L, Smith DP, Giovannoni SJ. The presence of the glycolysis

operon in SAR11 genomes is positively correlated with ocean productivity. *Environ Microbiol* 2010;

12: 490–500.

44. Giovannoni SJ, Hayakawa DH, Tripp HJ, Stingl U, Givan SA, Cho J-C, et al. The small genome of an

abundant coastal ocean methylotroph. *Environ Microbiol* 2008; **10**: 1771–1782.

45. Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer K-H, et al. Uniting the classification

of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Microbiol*

2014; **12**: 635–645.

46. Polyphasic taxonomy of Marine bacteria. *studylib.net*. https://studylib.net/doc/5577926/polyphasic-

taxonomy-of-marine-bacteria. Accessed 11 May 2020.

588 47. Olsen NS, Hendriksen NB, Hansen LH, Kot W. A New High-Throughput Screening Method for

589    Phages: Enabling Crude Isolation and Fast Identification of Diverse Phages with Therapeutic Potential.

590    *PHAGE* 2020; **1**: 137–148.

591 48. Nayfach S, Camargo AP, Eloe-Fadrosh E, Roux S, Kyrpides N. CheckV: assessing the quality of

592    metagenome-assembled viral genomes. *bioRxiv* . 2020. , 2020.05.06.081778

593 49. Adriaenssens EM, Sullivan MB, Knezevic P, van Zyl LJ, Sarkar BL, Dutilh BE, et al. Taxonomy of

594    prokaryotic viruses: 2018-2019 update from the ICTV Bacterial and Archaeal Viruses Subcommittee.

595    *Arch Virol* 2020; **165**: 1253–1260.

596 50. Deng L, Ignacio-Espinoza JC, Gregory AC, Poulos BT, Weitz JS, Hugenholtz P, et al. Viral tagging

597    reveals discrete populations in Synechococcus viral genome sequence space. *Nature* 2014; **513**: 242–

598    245.

599 51. Våge S, Storesund JE, Thingstad TF. SAR11 viruses and defensive host strains. *Nature* . 2013. , **499**:

600    E3–4

601 52. Zhao Y, Qin F, Zhang R, Giovannoni SJ, Zhang Z, Sun J, et al. Pelagiphages in the Podoviridae family

602    integrate into host genomes. *Environ Microbiol* 2018.

603 53. Marston MF, Martiny JBH. Genomic diversification of marine cyanophages into stable ecotypes.

604    *Environ Microbiol* 2016; **18**: 4240–4253.

605 54. Lima-Mendez G, Van Helden J, Toussaint A, Leplae R. Reticulate representation of evolutionary and

606    functional relationships between phage genomes. *Mol Biol Evol* 2008; **25**: 762–777.

607 55. Giovannoni S, Temperton B, Zhao Y. Giovannoni et al. reply. *Nature* . 2013. , **499**: E4–5

608 56. Zhang Z, Qin F, Chen F, Chu X, Luo H, Zhang R, et al. Novel pelagiphages prevail in the ocean.

609    *bioRxiv* . 2019. , 2019.12.14.876532

610 57. Tam W, Pell LG, Bona D, Tsai A, Dai XX, Edwards AM, et al. Tail tip proteins related to

611    bacteriophage λ gpL coordinate an iron-sulfur cluster. *J Mol Biol* 2013; **425**: 2450–2462.

612 58. Zaragoza-Solas A, Rodriguez-Valera F, López-Pérez M. Metagenome Mining Reveals Hidden Genomic

613    Diversity of Pelagimyophages in Aquatic Environments. *mSystems* 2020; **5**.

614 59. Mizuno CM, Guyomar C, Roux S, Lavigne R, Rodriguez-Valera F, Sullivan MB, et al. Numerous

cultivated and uncultivated viruses encode ribosomal proteins. *Nat Commun* 2019; **10**: 752.

60. Van Duin J, Wijnands R. The function of ribosomal protein S21 in protein synthesis. *Eur J Biochem*

1981; **118**: 615–619.

61. Xu J, Hendrix RW, Duda RL. Conserved translational frameshift in dsDNA bacteriophage tail assembly

genes. *Mol Cell* 2004; **16**: 11–21.

62. Giovannoni SJ, Thrash JC, Temperton B. Implications of streamlining theory for microbial ecology.

ISME J 2014; **8**: 1553–1565.

63. Neufeld JD, Boden R, Moussard H, Schäfer H, Murrell JC. Substrate-specific clades of active marine

methylotrophs associated with a phytoplankton bloom in a temperate coastal environment. *Appl Environ*

Microbiol 2008; **74**: 7321–7328.

64. Moon K, Kang I, Kim S, Kim S-J, Cho J-C. Genome characteristics and environmental distribution of

the first phage that infects the LD28 clade, a freshwater methylotrophic bacterial group. *Environ*

Microbiol 2017; **19**: 4714–4727.

65. Salcher MM, Neuenschwander SM, Posch T, Pernthaler J. The ecology of pelagic freshwater

methylotrophs assessed by a high-resolution monitoring and isolation campaign. *ISME J* 2015; **9**: 2442–

2453.

66. Sargeant SL, Murrell JC, Nightingale PD, Dixon JL. Seasonal variability in microbial methanol

utilisation in coastal waters of the western English Channel. *Mar Ecol Prog Ser* 2016; **550**: 53–64.

67. Bailly-Bechet M, Vergassola M, Rocha E. Causes for the intriguing presence of tRNAs in phages.

Genome Res 2007; **17**: 1486–1495.

68. Enav H, Kirzner S, Lindell D, Mandel-Gutfreund Y, Béjà O. Adaptation to sub-optimal hosts is a driver

of viral diversification in the ocean. *Nat Commun* 2018; **9**: 4698.

69. Weitz JS, Li G, Gulbudak H, Cortez MH, Whitaker RJ. Viral invasion fitness across a continuum from

lysis to latency. *Virus Evol* 2019; **5**: vez006.

70. Knowles B, Silveira CB, Bailey BA, Barott K, Cantu VA, Cobian-Guemes AG, et al. Lytic to temperate

switching of viral communities. *Nature* 2016; **531**: 466–470.

71. Silveira CB, Rohwer FL. Piggyback-the-Winner in host-associated microbial communities. *NPJ*

Biofilms Microbiomes 2016; **2**: 16010.

72.   Kang I, Oh HM, Kang D, Cho JC. Genome of a SAR116 bacteriophage shows the prevalence of this

phage type in the oceans. *Proc Natl Acad Sci U S A* 2013; **110**: 12343–12348.

73.   Martinez-Hernandez F, Fornas Ò, Lluesma Gomez M, Garcia-Heredia I, Maestre-Carballa L, López-

Pérez M, et al. Single-cell genomics uncover Pelagibacter as the putative host of the extremely abundant

uncultured 37-F6 viral population in the ocean. *ISME J* 2018.

74.   Alonso-Sáez L, Morán XAG, Clokie MR. Low activity of lytic pelagiphages in coastal marine waters.

ISME J 2018; **12**: 2100–2102.

75.   Parsons RJ, Breitbart M, Lomas MW, Carlson CA. Ocean time-series reveals recurring seasonal

patterns of virioplankton dynamics in the northwestern Sargasso Sea. *ISME J* 2012; **6**: 273–284.

76.   Middelboe M, Lyck PG. Regeneration of dissolved organic matter by viral lysis in marine microbial

communities. *Aquat Microb Ecol* 2002; **27**: 187–194.

77.   Middelboe M, Jorgensen NOG. Viral lysis of bacteria: an important source of dissolved amino acids

and cell wall compounds. *J Mar Biol Assoc U K* 2006; **86**: 605–612.

78.   Brum JR, Sullivan MB. Rising to the challenge: accelerated pace of discovery transforms marine

virology. *Nat Rev Microbiol* 2015; **13**: 147–159.

79.   Gregory AC, Solonenko SA, Ignacio-Espinoza JC, LaButti K, Copeland A, Sudek S, et al. Genomic

differentiation among wild cyanophages despite widespread horizontal gene transfer. *BMC Genomics*

2016; **17**: 930.

80.   Carini P, Steindler L, Beszteri S, Giovannoni SJ. Nutrient requirements for growth of the extreme

oligotroph 'Candidatus Pelagibacter ubique' HTCC1062 on a defined medium. *ISME J* 2013; **7**: 592–

602.

81.   Salisbury A, Tsourkas PK. A Method for Improving the Accuracy and Efficiency of Bacteriophage

Genome Annotation. *Int J Mol Sci* 2019; **20**.

82.   Zhang Z, Qin F, Chen F, Chu X, Luo H, Zhang R, et al. Culturing novel and abundant pelagiphages in

the ocean. *Environ Microbiol* 2020.

Exeter Viral Collection Number	Virus	Host	Viral Population Representative	Hypergeometric cluster	Genome	Length (bp)	G + C %	Taxon	Accession number	Simplified phonetic spelling	Meaning and origin of the names
EXVC025P	Eistla	HTCC1062	Eistla	Cluster A	Circular	39638	32.7	Podoviridae	MT375521	aɪsːtlaː	"stormy one", Giantess in the poetic Edda
EXVC018P	Eyrgjafa	HTCC1062	Eyrgjafa	Cluster B	Circular	38005	32.6	Podoviridae	MT375523	eːɪrgjaːfaː	"scar donor", Giantess in the poetic Edda
EXVC020P	Gjalp	HTCC1062	Eyrgjafa	Cluster B	Circular	37857	32.5	Podoviridae	MT375524	gjaːlp	"Roaring one", Giantess in the poetic Edda
EXVC021P	Greip	HTCC1062	Greip	Cluster D	Circular	34916	31.5	Podoviridae	MT375525	graɪp	"Grasp", Giantess in the poetic Edda
EXVC014P	Ran	H2P3α	Ran	Cluster A	Circular	41529	34.1	Podoviridae	MT375530	ran	"plundering", Goddess of the Sea
EXVC013S	Aegir	H2P3α	Kolga	Singleton	Fragment	18297	31.1	Siphoviridae	MT375519	æːgɪr	"Sea", God of the Sea
EXVC012P	Jörmungand	H2P3α	Ran	Cluster A	Circular	41529	34.1	Podoviridae	MT375528	jɔrmunˌgand	"huge monster", Giant sea serpent
EXVC016S	Kólga	H2P3α	Kolga	Singleton	Circular	48659	30.5	Siphoviridae	MT375529	kɔlgaː	"Cool-wave", Daughter of Ran and Aegir
EXVC019P	Unn	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375531	ʊːn	"Wave", Daughter of Ran and Aegir
EXVC015P	Hroenn	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375527	ɦrøn	"Wave", Daughter of Ran and Aegir
EXVC010P	Bylgja	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375520	bɪːlgjaː	"Billow"-wave, Daughter of Ran and Aegir
EXVC011P	Himinglæva	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375526	hiːmɪŋɡlæfa	"transparent-on-top"-wave, Daughter of Ran and Aegir
EXVC282S	Venkman	H5P1	Venkman	Singleton	Linear	38624	34.4	Siphoviridae	MT375522	vɛŋkmæn	Bill Murray's Character in the Ghostbusters movie

Phage	H2P3	HTVC1062	HTVC7211
Eistla		+	
Eyrgjafa		+	
Greip		+	
Ran	+		+
Kolga	+		+
Bylgja	+	+	+

A

(i) Increasing Viral Concentration

Tangential Flow Filtration

Viral Sample

2 L Seawater

(ii) Enrichment

Infection

Filtration

(iii) Purification

B

(i) Flow Cytometry

V1-L 0.2% V1-R 99.8%

Day 18

No-virus control

V1-L 47.1% V1-R 52.9%

Infected SAR11 culture

Events Count

Side Scatter (SSC-A)

Green Fluorescence (FITC-A)

10^3 10^4 10^4 10^5

(ii) Host Cell Growth Curves

Control SAR11 culture (no viruses)

Infected SAR11 culture

Typical infection

$Log10(cells.ml^{-1})$

Day

(iii) TE Microscopy

100 nm 100 nm

100 nm 100 nm

A

	September	October	November	February	March	April	July		
								Ia.1 Ia.3 ratio	
								Isolation success rates	
SAR11 HTCC7211		4/7	3/4	8/8	8/8	5/8		28/35	80%
HTCC1062	8/8	4/4	4/4	8/8	8/8	5/8		37/40	92.5%
H2P3α		4/4	4/4	7/7	8/8	6/7		29/30	96.6%

B

								H5P1 other OM43 ratio	
OM43 C6P1		0/7		0/8	0/8	0/8		0/31	0%
D12P1		0/4		0/8	0/8	0/8	2/8	2/36	5%
H5P1		6/7	7/7	0/8	6/8	2/8	0/8	21/46	45.6%

Successful isolations / Attempted isolations

A

Head morphogenesis

Tail Tip J

Tail fibre

RNR

Capsid

Thymidylate synthase

Lysozyme

48,784 bp

Ribosomal S21

Terminase

B

50 nm

100 nm

100 nm

C

Bordetella phage LK3

Pelagibacter Podoviridae

GOV2_74774

0.36

0.76

Pelagibacter phage Kolga

GOV2_16560

0.66

GOV2_49791

0.98

GOV2_50383

0.27

0.88

GOV2_49937

0.79

GOV2_01467

0.9

0.82

GOV2_38049

Pelagibacter Myoviridae

ARC
ANT
TT-EPI
TT-MES

0.3

A

RNAP II mediator
Peptidase
Stabilisation
Peptidase
Portal protein
38,624 bp

Protease
Holin
Terminase
Endonuclease

DNA replication
Tail proteins
Capsid proteins

B

500 nm

500 nm

100 nm

100 nm

C

Pasteurella phage Pm86
Mannheimia phage vB_MhS_535AP2
Tetrasphaera phage TJE1
Pseudomonas phage PS-1DNA
Pseudomonas phage vB_Pae_CF23a
Pseudomonas phage AF
Faith5x5
Asapag
Apricot
Denise
0.74
0.76
0.96
0.99
Sinorhizobium phage phiLM21
0.95
Lutum
98.0
0.92
0.7
0.8
Yersinia phage vB_Yen_X1
Arthrobacter phage Isolde
0.85
0.86
0.92
0.92
0.75
0.8
0.6
0.69
0.8
Paracoccus phage Shpa
0.94
Acinetobacter phage phiAC-1
0.91
Xanthomonas phage CP2
Aeromonas phage D3
Pseudomonas phage JBD44
OM43 phage Venkman EXVC282S
WEC_HYBRID_01170
Burkholderia phage PE067
LD28 phage P19250A

Siphoviridae

Myoviridae
0.3

Podoviridae

Fig. 1 Workflow for high-throughput isolation **(A)** i. Increasing concentration of viruses in water

samples by tangential flow filtration (TFF); ii. Initial infection of host cultures to enrich the sample

for specific viruses; iii. Purification of viral isolates through three rounds of Dilution-to-Extinction;

(B) Initial screen of viral infections using **(**i). Flow cytometry, by comparing populations of no-virus

controls and infected cultures; (ii). Comparing growth curves of no-virus control culture (HTCC1062)

against infected SAR11 cultures; (iii). Confirming the presence of viruses in infected SAR11 cultures

using transmission electron microscopy: Top left: HTCC1062 no-virus control, bottom left: infected

HTCC1062, top right: aggregated cellular debris and viruses, bottom right: virus found in infected

HTCC1062 culture.

Fig. 2 Summary of success rates for each bacterial host as well as relative contribution to the clade

community during sampling based on 16S rRNA analysis: (**A**) Three SAR11 strains HTCC1062,

HTCC7211 and H2P3α and (**B**) three OM43 strains C6P1, D12P1 and H5P1. Circles and associated

numbers represent number of successful isolations (i.e. successful passage through three rounds of

Dilution-to-Extinction), over the total number of attempts made. Bars at the top of each panel

represent the relative abundance (SAR11 subclades 1a.1 and 1a.3, and OM43 strains for A and B,

respectively) of strains in the microbial community sampled at the time of collection (based on 16S

rRNA community profiling). Percentages on the right represent total successes for all attempts per

host strain.

Fig. 3 Hypergeometric probability of shared gene content between known pelagiphage genomes

identified four main viral phylogenetic clusters (outlined with red boxes).

Fig. 4 The first reported siphovirus infecting SAR11 (Pelagibacter phage *Kolga* EXVC016S) isolated

on novel host *Pelagibacter sp.* H2P3α. (**A**) Gene map of the 48,784 bp genome, which contains 80%

hypothetical genes without known function; (**B**) Transmission Electron Micrographs of *Kolga* (left

and bottom) and an H2P3α host cell infected with *Kolga* (top right).   (**C**) Unrooted maximum

likelihood tree of *terL* genes found in pelagiphages. Branches containing members of the Podoviridae

and Myoviridae family infecting SAR11 are collapsed for clarity (full tree is available in

Supplementary Fig. 7). Closely related contigs prefixed with 'GOV' represent viral contigs from the

Global Ocean Virome dataset and the ecological zone from which they were assembled is marked:

Temperate-tropical epipelagic (TT-EPI); temperate-tropical mesopelagic (TT-MES); Antarctic

(ANT), Arctic Ocean (ARC).

Fig. 5 The first reported cultured virus known to infect a member of the OM43 clade (Methylophilales phage *Venkman* EXVC282S) which was isolated on host strain H5P1 from this study; **(A)** Gene map displaying protein coding genes. Gene function is colour coded as follows: Structural genes (blue); DNA replication (red); lysis (green); transcription (turquoise); packaging (orange); hypothetical genes (grey). **(B)** TEM images of infected and chaining H5P1 cells (top left), uninfected H5P1 chaining cells (top right), *Venkman* viral particles (bottom left and right); **(C)** Maximum likelihood tree (500 bootstraps) of concatenated viral *terL* and exonuclease genes. Host families of the phages are indicated on the figure.

Fig.6 Relative abundance (reads recruited per kilobase of contig per million reads (RPKM)) of known pelagiphage isolates and methylophage genomes in viromes from the Western English Channel [12] and the Global Ocean Virome [11]. Samples are organised by decreasing latitude per ecological zone: ARC: Arctic; TT-EPI: temperate-tropical epipelagic; TT-MES: temperate-tropical mesopelagic; ANT: Antarctic; WEC: Western English Channel.

Table 1. Summary of phages isolated and sequenced in this study.

Table 2. Host infectivity of viral populations isolated and sequenced in this study. *Pelagibacter sp.* H2P3α and *Pelagibacter bermudensis* HTCC7211 are warm-water ecotypes of SAR11 subclade 1a; HTCC1062 is a cold-water ecotype. *Pelagibacter* phage *Bylgja* was the only virus capable of infecting both ecotypes.

Supplementary Information

Supporting Methods

Water sampling: A total of 20 L of seawater was collected in rosette-mounted Niskin bottles at a
depth of 5m from the Western Channel Observatory (WCO;
http://www.westernchannelobservatory. org.uk/) coastal station 'L4' (50°15.00N; 4°13.00W) on
the following dates: 2018-09-24, 2018-10-17, 2018-11-05, 2019-02-11, 2019-03-11, 2019-04-01,
2019-07-22 (for details see Supplementary Table 2). Seawater was transferred immediately to a
clean 2 L acid-washed polycarbonate (PC) Nalgene bottles (Thermo Fisher Scientific, Waltham,
USA) and placed in a cooler box (Igloo, Katy, USA) at ambient temperature. Upon return to shore,
water was transported to the University of Exeter for immediate processing (Two hours maximum
duration from collection to processing).

Isolation of SAR11 strain H2P3a and OM43 strains: 1 L seawater was collected as described
above in September 2017, filtered through a 142 mm Merck Millipore pore-size 0.2 µm PC filter
and autoclaved in 2 L Duran bottles (DWK Life Sciences GmbH, Mainz, Germany) with the lid
tightly shut to maintain carbon chemistry [1]. Upon cooling, nutrients were added to make natural
seawater medium (NSW): 1 mM NH_4Cl, 10 µM K_2HPO_4, 1 µM $FeCl_3$, 25 µM glycine, 25 µM
methionine, 100 µM pyruvate [2]. 1 mL of medium was placed into each 2 mL well in three sterile
96-well acid washed Teflon plates (Radleys, UK). Surface water from station L4 was filtered
through a 142 mm Merck Millipore 0.2 µm PC filter to remove larger plankton and bacteria and
enrich for smaller bacteria. A 200 µL aliquot was stained with SybrGreen and then quantified on
a C6 Accuri flow cytometer. The remaining retained fraction was diluted in NSW to a density of
~ 1 cell·µL^{-1}. 1 µL of diluted inoculum was added to each well, with eight wells left blank as
negative controls and eight wells with undiluted low-nucleic acid community fraction as positive
controls. Plates were covered and incubated in the dark at 15 °C for three months and checked
monthly for positive growth by flow cytometry as described above. Positive wells (>10^6 cells·mL^{-1}
) were transferred into 50 mL cultures in NSW medium and passaged three times once they
achieved >10^6 cells·mL^{-1}). To verify culture purity and to taxonomically classify new isolates, the
16S rRNA gene was amplified from 10 µL of exponentially growing culture, heated to 95°C for
10 minutes, followed by PCR (25 cycles, 30 s @ 98 °C; 45 s @ 59 °C; 45 s @ 72 °C; final extension
of 90 s @ 72 °C) using commercially available 27F and 1492R primers by Eurofins Genomics.
Amplicon DNA was purified for Sanger sequencing (Eurofins) using a QIAquick PCR purification
kit (Qiagen) following the manufacturer's standard protocol. 16S rRNA sequences were compared
to known bacterial sequences in the SILVA SSU database with SINA [3]. Sequences were aligned
using T-coffee [4], curated using Gblocks [5] and maximum likelihood trees for both alignments
were created using PhyML with standard settings and 500 bootstraps [6]. The trees were visualised
using FigTree (v1.4.4, available https://github.com/rambaut/figtree).

Host cultivation: SAR11 strains HTCC1062 and HTCC7211 were kindly provided by the
Giovannoni lab (Oregon State University, USA) and were kept in continuous culture in 50 mL
polycarbonate flasks at 15 °C on artificial seawater medium ASM1 [7] throughout the duration of
the study. Following isolation, cultures of OM43 and H2P3α were maintained on ASM1; medium
for OM43 was further amended with 100 mM MeOH and 32 µL of commercially available MEM
amino acid solution per 1.6 L of medium (Sigma, M5550), to improve cellular yields and reduce
occurrences of chaining cells (observed through EM, Fig. 5, and a phenotype of alanine starvation
in SAR11 [2]), respectively.
49
Viral concentration and inoculation: Monthly samples of 2 L of surface water were collected from
station L4 as described above and sequentially filtered through a 142 mm Whatman GF/D filter
(2.7 µm pore size) and a 142 mm 0.2 µm pore polycarbonate filter (Merck Millipore) using a
peristaltic pump, to remove larger sized bacterioplankton. The filtrate was concentrated to 50 mL
(40-fold concentration) using a 50R VivaFlow tangential flow filtration unit (Sartorius Lab
instruments, Goettingen, Germany). The concentrate was filtered through a 0.1 µm pore PVDF
membrane syringe-filter to remove any residual small cells and used as inoculum for viral
isolation. 1 mL of exponentially growing bacterial host in ASM1 medium (amended with nutrients
for OM43 strains as described for host cultivation above) was placed into each 2 mL well of a 96-
well, acid washed Teflon plate. Eight wells were used as blank medium controls (no cell controls).
100 µL of viral inoculum was added to each well in the 96-well plate, with eight wells amended
with 10% (v/v) medium instead of viral inoculum to serve as no-virus controls. The plates were
incubated for ~2 weeks until no-virus controls reached maximum cell density (~2 weeks for strains
HTCC1062, HTCCC7211 and H2P3α; about 1 week for OM43 strains H5P1, C6P1, D12P1).
Growth was monitored at the end of the incubation period by flow cytometry; incubation periods
were determined by the average growth times and data of the strains used for infections. Cytograms
of virus-amended wells and no-virus controls were compared to identify infections as described
below. A detailed protocol of the viral isolation process is available on www.protocols.io (DOI:
dx.doi.org/10.17504/protocols.io.c36yrd). Once the no-virus controls reached maximum cell density,
inoculated wells that did not show signs of infection were pooled (according to host type), filtered
through a 0.1 µm pore PVDF membrane syringe (Durapore) filter to remove the cells and then
used as fresh inoculum in a new 96-well plate of target host. This was repeated up to three times
to amplify host-specific viral particles to a density whereby signs of viral infectivity could be
observed on the flow cytometer.
74
Identification of positive infection via flow cytometry: We observed positive infections in cells via
flow cytometry by comparing cytograms of wells inoculated with viruses against no-virus controls.
Cytograms of wells containing infected cultures were identifiable by (1) a population shift of up
to a 10-fold increase in green fluorescence compared to uninfected cultures; (2) an increase in the
'noise' fraction (presumably from cellular debris following lysis as well as viral particles) over
time (Supplementary Video 1); and (3) a reduced maximum cell density (ranging from 10 to 1000-

fold depending on initial cell density and viral inoculum concentration) in infected cultures
compared to no-virus controls (Fig. 1B). The presence of viral-like particles in wells where a shift
in green fluorescence, increased noise, and reduced maximum cell density were observed, were
verified by TEM and reinfection of new cultures in fresh medium.

Viral purification: Following identification of a positive infection within a well, the contents of
the well were filtered through a 0.1 µm pore-size PVDF syringe filter (Durapore) and used as
inoculum in three rounds of viral purification. Briefly, a 96-well Teflon plate was inoculated with
bacterial hosts as described previously. A 10-fold serial dilution series (from 10^0-10^{-10}) of the viral
inoculum in ASM1 was added to a single row of the plate at 10 % v/v, with one well per row as
no-virus controls (total of eight no-virus controls/plate). Plates were incubated at 18 °C for ~2
weeks until the no-virus controls reached maximum cell density and then screened for signs of
viral infection using flow cytometry as described above. For each dilution series, the well amended
with the lowest number of viruses that showed positive signs of infection was identified, and used
as the inoculum in another round of viral purification [8]. Using this format, we were able to purify
eight to twelve viral isolates simultaneously per plate.

Viral isolation costs and handling time. For each sample (processed environmental water sample),
all steps of the initial viral isolation process (counting one to two plates for a total of three times)
required ~6 hours run time on a flow cytometer. For the subsequent three rounds of purification,
one plate was required for each host-sample combination (another ~6 hours of cytometer run time
in total for one host-sample combination). Initial isolation of viruses from environmental water
samples and three rounds of purification took ~10 weeks of incubation time in total. Between all
steps, our protocol required ~7 hours of handling time per sample. Following three rounds of viral
purification, generating sufficiently high viral titres to extract enough DNA for sequencing
(approximately two weeks incubation time, and roughly four hours handling time over two days)
was the rate-limiting step, and so required subselection of available viruses for sequencing. Future
advances in DNA extraction efficiency, reducing DNA input requirements for sequencing and/or
automation of viral DNA extraction will enable all isolated viruses to be sequenced. We estimate
the cost of isolating a single virus is ~£20 for cultivation, flow cytometry and DNA extraction
consumables, and ~£50-100 for genome sequencing at 30-fold coverage required for successful
assembly, giving a total cost of £70-120 per sequenced viral isolate, not including the costs for
person-hours. Thus, our protocol provides a high-throughput and scalable approach to viral
isolation.

Transmission Electron Microscopy of viral isolates. For ultrastructural analysis, virus particles
were transferred onto pioloform-coated electron microscopy (EM) copper grids (Agar Scientific,
Standsted, UK) by floating the grids on droplets of virus-containing suspension for 3 min.
Following a series of four washes on droplets of deionized water, the bound virus particles were
contrasted with 2 % (w/v) uranyl acetate in 2 % (w/v) methyl cellulose (mixed 1:9) on ice for 8

min and the grids then air-dried on a wire loop after carefully removing excess staining solution
with a filter. Dried grids were inspected with a JEOL JEM 1400 transmission electron microscope
operated at 120 kV and images taken with a digital camera (ES 1000W CCD, Gatan, Abingdon,
UK).
125
Host ranges of viral isolates. To test the infectivity of phage isolates against different hosts, 2 mL
96-well Teflon plates were prepared with 1 mL of exponentially growing bacterial host
(HTCC1062, HTCC7211, H2P3α) in ASM1 medium. As medium control, ASM1 without any
bacteria was added to eight wells. 100 µL of phage inoculum was added to eight wells per host
type. As no-virus control, for each host further eight wells were left without any viruses added to
them. The plate was incubated for ~2 weeks and analysed for signs of infection using flow
cytometry as described above. A phage was classified as unable to infect a certain host if none of
the eight amended replicates showed signs of infection.
134
DNA preparation and sequencing of viral isolates. 50 mL ASM1 (amended with 100 mM MeOH
and amino acid solution for OM43 as described above) in 250 mL acid-washed, polycarbonate
flasks were inoculated with host cultures to 10^6 cells·mL^{-1}. The cultures were infected with 10%
v/v viruses in ASM1 medium and incubated until after host cell lysis was detected using flow
cytometry. The cultures were transferred to 50 mL Falcon tubes and the cellular fraction was
removed by centrifugation (GSA rotor, Thermo Scientific 75007588) at 8,500 rpm/10,015 x g for
120 minutes. The supernatant was subsequently filtered through pore-size 0.1 µm PVDF syringe
filter membranes to remove any remaining smaller cellular debris. Phages in the filtrate were
precipitated using a modified version of an established PEG8000/NaCl DNA isolation method [9].
Briefly, the filtered phage lysate was transferred into 50 mL Falcon tubes with pre-weighted 5 g
PEG8000 and 3.3 g NaCl, shaken until both dissolved and incubated on ice overnight. The phage
particles were then pelleted by centrifugation at 8,500 rpm/10,015 x g for 90 min at 4 °C.
Supernatant was discarded and phage particles resuspended by rinsing Falcon tubes twice with 1
mL SM buffer (100 mM NaCl, 8 mM MgSO$_4$·7H$_2$O, 50 mM Tris-Cl). DNA was extracted using
the Wizard DNA Clean-Up system (Promega) following manufacturer's instructions, but instead
of TE we used pre-heated 60 °C nuclease free water to elute DNA. DNA was sequenced by
MicrobesNG (Birmingham, UK) with Illumina paired-end HiSeq 2500. Default settings were used
for all programs listed below, unless specified otherwise. All reads were trimmed, quality-
controlled and error-corrected using bbmap and tadpole [10]. Contigs were assembled using
SPAdes v3.13 [11] and evaluated with QUAST [12]. The trimmed reads were mapped back against
the contigs for scaffolding using bowtie2 [13], BamM (alignment 0.9, identity 0.95) (available at
https://github.com/minillinim/BamM) and samtools [14]. Viral contigs were confirmed with
VirSorter (categories 1 or 2, >15kbp) and only accepted if the representative contig showed mean
depth coverage an order of magnitude greater than the next highest recruiting contig (to filter out
any cellular DNA carryover). Gene calls returned by VirSorter were imported into DNA Master
for manual curation [15]. Additional gene calls were made using GenMark [16], GenMarkS [17],

GenMarkS2 [18], GenMark.hmm [19], GenMark.heuristic [20], Glimmer v.3.02 [21], and
Prodigal v.2.6.3 [22]. All gene calls were listed and compared using a scoring system which
evaluates gene length, gene overlap and coding potential of ORFs [15]. ORFs were then annotated
using BLASTp against the NCBI's non-redundant protein sequences [23], Phmmer [24] against
UniProts UniProtKB and uniprotrefprot [25], Swissprot [26] as well as InterProScan [27] and Pfam
[28]. Virally-encoded tRNAs were identified with tRNAScan-SE v2.0 using settings `tRNAscan-
SE -qQ --detail -o# -m# -f# -l# -c tRNAscan-SE.conf -s# -B` [29] to identify putative
tRNA genes. Matching tRNA genes (e.g. viral and host tRNA-Leu) were then aligned using
BLASTn [30]. All viral genomes were organised into viral clusters (ICTV-recognised genera)
based on shared gene content with VConTACT2 [31], and into populations using a boundary cutoff
of 95% ANI over 85% contig length [32].

Evaluation of gene-sharing networks with hypergeometric testing. For the hypergeometric testing
we gathered all annotated genomes from this study, other known pelagiphage isolates [33, 34] and
we arbitrarily picked one representative contig from each cluster of complete phage genomes that
were retrieved from a metagenomic Mediterranean deep chlorophyll maximum sample, that the
authors speculated to be infective towards 1A SAR11s and included them in the phylogenetic
analysis [35]. The contigs were imported into anvio v6.1 [36] where open reading frames were
identified internally using Prodigal v2.6.3. The contigs were then used to run a pangenome analysis
using the --use-ncbi-blast option. The binning function was then used to export the protein
clusters, from which a matrix of the sum of shared proteins between all phages was created. The
hypergeom function in the sciPy.stats python package was used to calculate the probability of
sharing a protein between phages, based on which we created an average correlation cluster map.

Viral phylogenetic analysis. First we identified additional contigs to include in the phylogenetic
analysis. The Global Ocean Virome (GOV2) [37] and a virome from the Western English Channel
[38] were screened for contigs that belonged to the same population (95% ANI over 80% length
using ClusterGenomes.pl - https://github.com/simroux/ClusterGenomes) or the same genus
(clustered by VConTACT2 [39] into the same viral cluster, using default parameters) as isolate
genomes. Additionally, all available genomes from Pelagibacter phage isolates were included, as
well as selected fosmid derived contigs from Mediterranean metagenomes and 26 sequences from
metagenomic mining that were identified as putative Pelagimyophages [33–35, 40, 41]. All
sequences were subjected to gene calling using Prodigal v2.6.3. (default parameters) and screened
for TerL genes by aligning all proteins against known TerL genes in annotated genomes (belonging
to clusters A, B and D, as well as siphovirus *Kolga*) from this study using Protein BLAST (default
parameter) [22, 30]. The TerL gene of Bordetella phage LK3 was added as an outgroup. The TerL
genes were aligned within the Phylogeny.fr online service [42] opting for MUSCLE alignment
[43] and Gblocks curation [5] with default settings. A maximum-likelihood tree was calculated
with   PhyML   [44,   45],   visualised   using   FigTree   (v.1.4.4,   available   at

http://tree.bio.ed.ac.uk/software/figtree/) and edited in Inkscape (www.inkscape.org) for aesthetics
(font adjustments and addition of GOV2 metadata).

We only identified a single additional contig from metagenomes using the approach described
above that was similar to OM43 phage *Venkman*. Only one other known phage for a related host
(LD28 phage P19250A [46]) was available. These three contigs as well as all 14,883 viral contigs
publicly available in January 2020 on http://millardlab.org/ were gene called using Prodigal v2.6.3.
[22]. All called genes were compared to vFams [47] using hmmsearch [48] and sorted by E-value
for best hits. The best matching gene sequences that were identified as similar to shared vFams
found in *Venkman* and P19250A were important to the Phylogeny.fr online service [42]. Sequences
of were aligned individually using T-coffee [4], curated using Gblocks [5] and concatenated
manually. Maximum likelihood trees for both alignments were created using PhyML with standard
settings and 500 bootstraps [6, 44]. The trees were visualised with FigTree (v.1.4.4) and edited in
Inkscape (www.inkscape.org) for aesthetics. All sequences used for phylogenetic analysis are
available at https://github.com/ViralPirates/viral-dte.

Calculation of Average Nucleotide Identity. All-vs-all average nucleotide identity (ANI) for our
new isolates and existing pelagiphages using nucmer v.4.1 with the --nooptimize flag, followed by
show-coords to convert the delta file into tabular format. ANI was calculated by taking the mean
of all matching regions between two isolates.

Viral metagenomic recruitment. For assessing relative abundances of phage contigs in global
datasets, a virome from the Western English Channel and all samples of the Global Ocean
Virome dataset (GOV2) were used for recruitment [37, 38]. Metagenomic reads were
downloaded from the European Nucleotide Archive ERR2625613 and subsampled to 5 million
reads using bbmap's `reformat.sh` command. A bowtie2 [13] index of dereplicated contigs was
created from 25 known pelagiphage genomes from isolates [33, 34, 41], LD28 phage P19250A
[46] and from this study six viral population representative genomes isolated on SAR11 hosts
and one Methylophilales phage *Venkman* isolated on OM43 host. Reads were mapped against the
genomes for each metagenome sample using bowtie2 using these commands: `bowtie2 --seed`
`42 --non-deterministic`. To calculate coverage and Reads Per Kilobase of contig per Million
reads (RPKM) we used coverm [reference], with the following commands: `coverm contig --`
`bam-files *.bam --min-read-percent-identity 0.9 --methods rpkm --min-covered-`
`fraction 0.4.` To minimize false positive detection rates [32], contigs that did not meet the
40% minimum genome coverage in a given sample were given an RPKM value of 0.

Evaluation of isolation efficiency as a function of matching host and virus geography. In total, 105
attempts were made for isolating viruses for SAR11 across the three host strains (HTCC7211: 35,
HTCC1062: 40; H2P3α: 30). For each strain, the number of successful attempts were recorded in
a vector, v_s, of length n where n is the number of attempts made to isolate a virus, such that $v_i=1$ if

a successful attempt was made and v_i=0 if an unsuccessful attempt was made. To test whether
observed rates of success for H2P3 were significantly elevated compared to those for HTCC1062
and HTCC7211, the observed rates were evaluated against the null hypothesis that host strain did
not make a difference to isolation success. Briefly, for each pair (H2P3 vs HTCC1062; H2P3α vs
HTCC7211), the two vectors v_{s1} and v_{s2} were concatenated, and then randomly subsampled 999
times with replacement to a size of the smallest value of n between the vectors. The number of
successes in each subsample were counted and the distribution of values across the 999 bootstraps
was recorded. P-values were calculated by evaluating how many times within the 999 bootstraps
a value as extreme as the observed value had been recorded.

16S rRNA high-throughput amplicon sequencing and analysis. Genomic DNA was extracted using
Qiagen DNeasy PowerWater Kit (REF 14900-50-NF) from biomass retained on 0.2 µm PC filters
following the manufacturer's protocol with minor modifications to increase the yield. Step 7 was
modified from a 5 minute to 10 minute vortex bead beat. Step 21 was changed to have a 2 minute
incubation with EB warmed to 55C. Amplification of the hypervariable V4 region of the 16S rRNA
gene was performed using the 515fB (5'-GTGYCAGCMGCCGCGGTAA-3') and 806rB (5'-
GGACTACNVGGGTWTCTAAT-3') primers. Libraries for each reaction were done attaching
dual indices and Illumina adapters with the NexTera XT Index Kit (Illumina Inc.) Purified libraries
were pooled and sequenced in a MiSeq platform. Primer sequences from paired-end fastq files
were trimmed using CutAdapt [49]. Trimmed sequences were quality filtered, dereplicated and
merged with DADA2 R package (version 1.8 [50]). Taxonomic assignment was performed using
the "assignTaxonomy" command within dada2 pipeline and silva_nr_v132 database [51] as
training set. SAR11 ASVs were extracted for further characterization using Phyloassigner v6.166
[52] and oligotyping [53]. SAR11 phylogenetic database SAR11_Phy_DB used in this study is
available at https://www.github.com/lbolanos32/NAAMES_2020. Relative contribution barplots
were done in R using ggplot2 [54] and edited in inkscape (www.inkscape.org) for aesthetics.
261

Supplementary References

1. Henson MW, Pitre DM, Weckhorst JL, Celeste Lanclos V, Webber AT, Cameron Thrash J. Artificial

Seawater Media Facilitate Cultivating Members of the Microbial Majority from the Gulf of Mexico.

American Society for Microbiology 2016; **1**: 1–9.

2. Carini P, Steindler L, Beszteri S, Giovannoni SJ. Nutrient requirements for growth of the extreme

oligotroph 'Candidatus Pelagibacter ubique' HTCC1062 on a defined medium. *ISME J* 2013; **7**:

592–602.

3.   Pruesse E, Peplies J, Glöckner FO. SINA: accurate high-throughput multiple sequence alignment of

ribosomal RNA genes. *Bioinformatics* 2012; **28**: 1823–1829.

4.   Notredame C, Higgins DG, Heringa J. T-Coffee: A novel method for fast and accurate multiple

sequence alignment. *J Mol Biol* 2000; **302**: 205–217.

5.   Talavera G, Castresana J. Improvement of phylogenies after removing divergent and ambiguously

aligned blocks from protein sequence alignments. *Syst Biol* 2007; **56**: 564–577.

6.   Guindon S, Lethiec F, Duroux P, Gascuel O. PHYML Online--a web server for fast maximum

likelihood-based phylogenetic inference. *Nucleic Acids Res* 2005; **33**: W557–9.

7.   Carini P, Campbell EO, Morré J, Sañudo-Wilhelmy SA, Thrash JC, Bennett SE, et al. Discovery of a

SAR11 growth requirement for thiamin's pyrimidine precursor and its distribution in the Sargasso

Sea. *ISME J* 2014; **8**: 1727–1738.

8.   Nagasaki K, Bratbak G. Isolation of viruses infecting photosynthetic and nonphotosynthetic protists.

Manual of aquatic viral ecology ASLO 2010; 92–101.

9.   Solonenko N. Isolation of DNA from phage lysate. *protocols.io* 2016.

10.   Bushnell B, Rood J, Singer E. BBMerge – Accurate paired shotgun read merging via overlap. *PLOS*

ONE . 2017. , **12**: e0185056

11.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new

genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012; **19**:

455–477.

12.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome

assemblies. *Bioinformatics* 2013; **29**: 1072–1075.

13.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012; **9**: 357–

359.

14.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map

format and SAMtools. *Bioinformatics* 2009; **25**: 2078–2079.

15. Salisbury A, Tsourkas PK. A Method for Improving the Accuracy and Efficiency of Bacteriophage

Genome Annotation. *Int J Mol Sci* 2019; **20**.

16. Borodovsky M, McIninch J. GenMark: Parallel gene recognition for both DNA strands. *Computers*

Chemistry 1993; **17**: 123–133.

17. Besemer J, Lomsadze A, Borodovsky M. GeneMarkS: a self-training method for prediction of gene

starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic*

Acids Res 2001; **29**: 2607–2618.

18. Lomsadze A, Gemayel K, Tang S, Borodovsky M. Modeling leaderless transcription and atypical

genes results in more accurate gene prediction in prokaryotes. *Genome Res* 2018; **28**: 1079–1089.

19. Zhu W, Lomsadze A, Borodovsky M. Ab initio gene identification in metagenomic sequences.

Nucleic Acids Res 2010; **38**: e132.

20. Besemer J, Borodovsky M. Heuristic approach to deriving models for gene finding. *Nucleic Acids*

Res 1999; **27**: 3911–3920.

21. Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and endosymbiont

DNA with Glimmer. *Bioinformatics* 2007; **23**: 673–679.

22. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene

recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**: 119.

23. Pruitt KD, Tatusova T, Maglott DR. NCBI reference sequences (RefSeq): a curated non-redundant

sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* 2007; **35**: D61–5.

24. Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. HMMER web server: 2018 update.

Nucleic Acids Res 2018; **46**: W200–W204.

25. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 2019; **47**:

D506–D515.

26. Bairoch A, Apweiler R. The SWISS-PROT protein sequence database and its supplement TrEMBL

in 2000. *Nucleic Acids Res* 2000; **28**: 45–48.

27.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale

321   protein function classification. *Bioinformatics* 2014; **30**: 1236–1240.

322   28.   Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the protein

323   families database. *Nucleic Acids Res* 2014; **42**: D222–30.

324   29.   Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of transfer

325   RNA genes. *Nucleic Acids Res* 2016; **44**: W54–7.

326   30.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol

327   Biol* 1990; **215**: 403–410.

328   31.   Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Taxonomic

329   assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nat

330   Biotechnol* 2019; **37**: 632–639.

331   32.   Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an in silico

332   evaluation of metagenome-enabled estimates of viral community composition and diversity. *PeerJ*

333   2017; **5**: e3817.

334   33.   Zhao Y, Temperton B, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC, et al. Abundant SAR11

335   viruses in the ocean. *Nature* 2013; **494**: 357–360.

336   34.   Zhao Y, Qin F, Zhang R, Giovannoni SJ, Zhang Z, Sun J, et al. Pelagiphages in the Podoviridae

337   family integrate into host genomes. *Environ Microbiol* 2018.

338   35.   Mizuno CM, Rodriguez-Valera F, Kimes NE, Ghai R. Expanding the marine virosphere using

339   metagenomics. *PLoS Genet* 2013; **9**: e1003987.

340   36.   Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced

341   analysis and visualization platform for 'omics data. *PeerJ* 2015; **3**: e1319.

342   37.   Gregory AC, Zayed AA, Sunagawa S, Wincker P, Sullivan MB, Temperton B, et al. Marine DNA

343   Viral Macro- and Microdiversity from Pole to Pole. *Cell* 2019; **177**: 1109–1123.

38.   Warwick-Dugdale J, Solonenko N, Moore K, Chittick L, Gregory AC, Allen MJ, et al. Long-read

metagenomics reveals cryptic and abundant marine viruses. *bioRxiv* . 2018. , 345041

39. Bolduc B, Jang HB, Doulcier G, You Z-Q, Roux S, Sullivan MB. vConTACT: an iVirus tool to

classify double-stranded DNA viruses that infect Archaea and Bacteria. *PeerJ* 2017; **5**: e3243.

40. Zaragoza-Solas A, Rodriguez-Valera F, López-Pérez M. Metagenome Mining Reveals Hidden

Genomic Diversity of Pelagimyophages in Aquatic Environments. *mSystems* 2020; **5**.

41. Zhang Z, Qin F, Chen F, Chu X, Luo H, Zhang R, et al. Culturing novel and abundant pelagiphages

in the ocean. *Environ Microbiol* 2020.

42. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, et al. Phylogeny.fr: robust

phylogenetic analysis for the non-specialist. *Nucleic Acids Res* 2008; **36**: W465–9.

43. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic*

Acids Res 2004; **32**: 1792–1797.

44. Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast, accurate, and

powerful alternative. *Syst Biol* 2006; **55**: 539–552.

45. Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by

maximum likelihood. *Syst Biol* 2003; **52**: 696–704.

46. Moon K, Kang I, Kim S, Kim S-J, Cho J-C. Genome characteristics and environmental distribution

of the first phage that infects the LD28 clade, a freshwater methylotrophic bacterial group. *Environ*

Microbiol 2017; **19**: 4714–4727.

47. Skewes-Cox P, Sharpton TJ, Pollard KS, DeRisi JL. Profile hidden Markov models for the detection

of viruses within metagenomic sequence data. *PLoS One* 2014; **9**: e105067.

48. Eddy S. HMMER: profile HMMs for protein sequence analysis. 1998.

49. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 2011; **17**: 10–12.

50. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-

resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**: 581–583.