Efficient Dilution-to-Extinction isolation of novel virus-host model systems for

fastidious heterotrophic bacteria.

Holger H. Buchholz[1], Michelle L Michelsen[1], Emily Browne[1], Michael J. Allen[1,2], Ben Temperton[1*]

[1]University of Exeter, School of Biosciences, Exeter, UK;

[2]Plymouth Marine Laboratory, Plymouth, UK

*Corresponding Author

b.temperton@exeter.ac.uk

Running Title: Cultivation of viruses of fastidious bacteria

Keywords: bacteriophages, high-throughput culturing, marine microbiology, virus-host systems, Western English Channel

Abstract

Microbes and their associated viruses are key drivers of biogeochemical processes in marine and soil biomes. While viruses of phototrophic cyanobacteria are well-represented in model systems, challenges of isolating marine microbial heterotrophs and their viruses have hampered experimental approaches to quantify the importance of viruses in nutrient recycling. A resurgence in cultivation efforts has improved the availability of fastidious bacteria for hypothesis testing, but this has not been matched by similar efforts to cultivate their associated bacteriophages. Here, we describe a high-throughput method for isolating important virus-host systems for fastidious heterotrophic bacteria that couples advances in culturing of hosts with sequential enrichment and isolation of associated phages. Applied to six monthly samples from the Western English Channel, we first isolated a new SAR11 and three new members of the methylotrophic bacterial clade OM43, and used these as bait to isolate 117 new phages including the first known siphophage infecting SAR11, and the first known phages for OM43. Genomic analyses of 13 novel viruses revealed representatives of three new viral genera, and infection assays showed that SAR11 viruses have ecotype-specific host-ranges. Similar to the abundant human-associated phage ɸCrAss001, infection dynamics within the majority of isolates suggested either prevalent lysogeny, irrespective of detectable associated genes, and/or host phenotypic bistability, with lysis putatively maintained within a susceptible subpopulation. Broader representation of virus-host systems in culture collections and genomic databases will improve both our understanding of virus-host interactions, and accuracy of computational approaches to evaluate ecological patterns from metagenomic data.

Introduction

It is estimated that viral predation kills ~15% of bacterial cells in marine surface water each day [1]

and is a major contributor to nutrient recycling via the viral shunt, where marine viruses make cell-

bound nutrients available to the neighbouring microbial community through viral lysis of host cells

[2, 3]. Viruses are key players in the modulation of carbon fluxes across the oceans (150 Gt/yr),

increasing particle aggregation and sinking to depth [2, 4], and accounting for 89% of the variance in

carbon export from surface water to the deep ocean [5]. Viruses alter host metabolism through

auxiliary metabolic genes (AMGs), increasing and altering the cellular carbon intake of infected cells

[6]. Virus-host interactions also increase co-evolutionary rates of both predator and prey via Red

Queen dynamics [7, 8]. While recent metagenomic advances have provided major insight into global

viral diversity and abundance [9–12], mechanistic understanding of virus-host interactions in

ecologically important taxa is reliant on experimental co-culturing of model systems. In

cyanobacteria, such systems have shown that viruses increase duration of photosynthetic function

[13] and can inhibit CO_2 fixation, providing direct evidence that viruses of abundant phototrophs

have a direct and major role in nutrient regulation and global carbon cycling [14]. Furthermore,

isolation of novel viral model systems provides complete or near-complete viral genomes with known

hosts, critical to the development, ground-truthing and application of computational methods to

identify viral genomes in metagenomic data (e.g. VirSorter [15]; VirFinder [16]; MARVEL [17]);

quantify boundaries for viral populations [12, 18] and genera (VConTACT2 [19]); understand the

importance of AMGs in altering nutrient flux in natural communities [20] and to predict host ranges

of uncultured viruses *in situ* (e.g. WIsH [21, 22]).

In comparison to viruses of primary producers, such as cyanophage, which are well represented with

model systems and well-studied in the laboratory, virus-host model systems for similarly important

and abundant marine heterotrophic bacteria are rare. Available models of viruses infecting

heterotrophs are heavily biased towards those with a fast-growing copiotrophic host that grow readily

on solid agar, enabling the use of plaque assays to isolate associated viruses. Such systems are not

representative of the vast majority of heterotrophs in nutrient-limited soil and aquatic environments,

which are dominated by slow-growing, oligotrophic taxa with few regulatory mechanisms and

complex auxotrophies that do not lend themselves to growth on solid or rich medium [23–26].

Advances in Dilution-to-Extinction culturing of ecologically important hosts have enabled the

cultivation of many fastidious bacterial taxa that are not amenable to growth on solid media from

soil, marine [27, 28] and freshwater environments [29]. Without plaque assays to identify infection

and purification of new viral taxa, cultivation of associated viruses is challenging, and exacerbated

by the slow growth rates and complex nutrient requirements of their hosts. Paucity of such model

systems introduces significant bias in our understanding of viral influence on global carbon

biogeochemical cycles. Therefore, it is important that the efforts to isolate heterotrophic bacterial taxa

for experimentation and synthetic ecology is matched by efforts to isolate their associated viruses.

Here, we adapted recent advances in Dilution-to-Extinction culturing of hosts [28] and protocols to

isolate viruses from liquid media [30] to improve efficiency of recovering novel virus-host systems

for fastidious taxa: First, we used sequential enrichment of viruses from natural communities on target

hosts to improve rates of viral isolation [31]; Second, we replaced the requirement for time-intensive

epifluorescent microscopy with comparative flow cytometry between infected and uninfected hosts

to screen for putative viral infection, followed by confirmation using Transmission Electron

Microscopy (TEM). We selected the ecologically important, genomically streamlined SAR11 and

OM43 heterotrophic marine clades as models for viral isolation. These clades are abundant and

important to global carbon biogeochemistry [23, 32–36], but little is known of their associated viral

diversity. In the case of viruses infecting SAR11, challenges in assembling abundant and

microdiverse genomes from viral metagenomes limit our ability to determine associated host-virus

ecology and representatives for *in silico* host prediction [12, 37, 38], particularly for novel species

that lack any representation in the databases. This was demonstrated in the successful isolation of the

first known pelagiphages by culturing, including the globally dominant HTVC010P, which, prior to

its isolation, was entirely missed in marine viromes [39]. To the best of our knowledge there are no known viruses for OM43, which plays an important role in remineralisation of volatile carbon associated with phytoplankton blooms [32, 34, 40]. As well as their importance in experimentally evaluating the impact of predator-prey dynamics on global marine carbon biogeochemistry, these systems are critical to understand the influence of genome minimalism on host-virus co-evolution. Novel SAR11 and OM43 representatives from the Western English Channel were isolated and used as bait to isolate associated viruses. Initial concentration of viruses in inocula by tangential flow filtration, followed by one to three rounds of sequential enrichment on target hosts in 96-well plates yielded 117 viral isolates from 218 inoculated cultures from seven monthly water samples from September 2018 to July 2019. 94 out of 105 (90%) inoculated SAR11 cultures yielded positive viral infections. 23 out of 113 (20%) inoculated OM43 cultures yielded positive infections. A subsample of positive infections for both clades were subsequently purified and sequenced, revealing the first known siphovirus infecting SAR11 and providing the first known host-virus model for OM43.

Results and Discussion

Isolation of a novel SAR11 strain and three new OM43 strains from the Western English Channel to use as bait for phage isolation

Dilution-to-Extinction culturing for host taxa using natural seawater-based medium on three 96-well plate samples in September 2017 yielded the first SAR11 strain to be isolated from the Western English Channel (named H2P3α) and the first three new OM43 strains (named C6P1, D12P1, H5P1) from the Western English Channel. The full-length 16S rRNA gene of H2P3α was 100% identical to that of the warm water SAR11 ecotype *Pelagibacter bermudensis* HTCC7211 (subclade 1A) and was considered to be a local variant [41] (Supplementary Fig. 1A). All three novel OM43 isolates (named CP61, D12P1, H5P1) were most closely related to *Methylophilales* sp. HTCC2181 (C6P1 96.17%,

D12P1 96.62%, H5P1 97.79% nucleotide identity across the full 16S rRNA gene), a streamlined

member of the OM43 clade with a 1.3 Mbp genome, isolated from surface water of the North Eastern

Pacific [42] (Supplementary Fig. 1B). The average identity of the 16S rRNA gene of isolates CP61,

D12P1 and H5P1 to each other was ~98.46% (Supplementary Table 1), suggesting they are

representatives of the same genus [43].

An efficient, low-cost method of isolating new viruses yielded 117 new viral isolates for SAR11

and OM43 taxa

Using the four new hosts and established SAR11 isolates *Pelagibacter ubique* HTCC1062 (cold-

water ecotype) and HTCC7211 (warm water ecotype), we developed an optimised viral isolation

pipeline (Fig. 1) and applied it to six monthly water samples from the Western English Channel, taken

between September 2018 to April 2019 (Supplementary Table 2). Briefly, each month, we inoculated

one to two 96-well Teflon plates containing our four new hosts, plus SAR11 isolates HTCC1062 and

HTCC7211 at ~10^6 cells per mL with a natural viral community concentrated by tangential flow

filtration. Plates were monitored by flow cytometry and growth of putatively infected cultures was

compared to those of uninfected controls over the course of ~2 weeks, due to the slow growth rates

of SAR11 and OM43 [34, 44]. Additional samples were taken July 2019 to attempt viral isolation on

OM43 strains D12P1 and H5P1. In total, out of 218 cultures inoculated with concentrated natural

viral communities for initial isolation, 117 viruses were purified and still infective after at least three

passages, evidenced by differences in cytograms between cultures inoculated with viruses and control

cultures (Supplementary Fig. 2-5), demonstrating an overall efficiency of 53% and an average yield

of 18 isolates per sample (Fig. 2). All viral isolations required between one and three rounds of host

enrichment (Supplementary Table 3) before changes in host growth curves between infected and

uninfected cultures could be observed by flow cytometry, suggesting a minimal viral density is

required to identify infected wells using this approach, but this threshold can be reached through

successive rounds of enrichment.

For each sample, all steps of the initial viral isolation process (counting one to two plates for a total

of three times) required ~6 hours run time on a flow cytometer. For the subsequent three rounds of

purification, one plate was required for each host-sample combination (another ~6 hours of cytometer

run time in total for one host-sample combination). Both initial isolation of viruses from

environmental water samples and three rounds of purification took ~10 weeks of incubation time in

total. Between all steps, our protocol required ~7 hours of handling time per sample. Following three

rounds of viral purification, generating sufficiently high viral titres to extract enough DNA for

sequencing (approximately two weeks incubation time, and roughly four hours handling time over

two days) was the rate-limiting step, and so required subselection of available viruses for sequencing.

Future advances in DNA extraction efficiency, reducing DNA input requirements for sequencing

and/or automation of viral DNA extraction will enable all isolated viruses to be sequenced. We

estimate the cost of isolating a single virus is ~£20 for cultivation, flow cytometry and DNA

extraction consumables, and ~£50-100 for genome sequencing at 30-fold coverage required for

successful assembly, giving a total cost of £70-120 per sequenced viral isolate, not including the costs

for person-hours. Thus, our protocol provides a high-throughput and scalable approach to viral

isolation.

Novel pelagiphages are ecotype-specific and persist in the community

We hypothesised that the density dependence of predator/prey dynamics would result in greater

isolation efficiency of viruses on the cold-water SAR11 ecotype, HTCC1062, during winter months,

when this host is more abundant. Contrary to our expectations, the isolation efficiencies (number of

successes / number of attempts) were: 80% for HTCC7211, 93% for HTCC1062 and 96.6% for

H2P3α, but these were not significantly different (see Supporting Methods: HTCC1062 vs. H2P3α

(P=0.5, 999 bootstraps); HTCC1062 vs. HTCC7211 ($P=0.18$, 999 bootstraps)). Temperature ranged

from 9.3 °C - 14.1 °C from October to April, when isolation was attempted on all strains

(Supplementary Table 2). Neither temperature, ecotype nor an ecotype*temperature interaction term

were significantly correlated with isolation efficiency based on a generalized linear model using a

quasi-binomial error distribution (see Supplementary Methods). One explanation for this result is that

host ecotype abundances are constant throughout the year and unaffected by temperature. HTCC1062

and HTCC7211 have specific growth rates of ~0.22 and ~0.12 divisions per day at 10 °C, respectively

[44], and our new SAR11 isolate from the Western English Channel, H2P3α, showed similar growth

rates to HTCC7211 (Supplementary Fig. 6), with specific growth rates of ~0.10, ~0.45 and ~0.84

divisions per day at 10, 18 and 25 °C, respectively. Measured lower specific growth rates of warm

water ecotypes at winter temperatures, and evidence of global ecotype niche partitioning [45] suggest

that uniform ecotype abundance across seasons in the Western English Channel is unlikely, but that

slow growth of warm water ecotypes at *in situ* temperatures is possible.

An alternative explanation is that isolated viruses have a broad host range that encompasses both

warm- and cold-water ecotypes. We tested the host range (Table 1) of six pelagiphage population

representatives out of the 12 pelagiphages isolated from samples in October 2018 and November

2018 (*in situ* water temperature of 14.8 °C and 14.2 °C respectively) across the three SAR11 strains

(Table 2). Viruses *Eistla, Eyrgjafa* and *Greip* all infected cold-water ecotype HTCC1062,

exclusively, while *Ran* and *Kolga* only infected warm-water ecotypes HTCC7211 and H2P3α.

Therefore, our new pelagiphages appear to be broadly ecotype-specific, confirming previous findings

[46], with important consequences for their global distribution. *Bylgia* was the only virus that could

infect both warm and cold-water ecotypes. Ecotypic specificity at first seems at odds with our ability

to isolate viruses on the warm water SAR11 ecotype at *in situ* temperature far below its optimal

growth temperature. A possible explanation is that pelagiphages are ecotype-specific, but persist in

the water column throughout the year in sufficient densities to be isolated by our enrichment method.

Measured *in situ* temperatures during our sampling period were sufficient to support slow growth of warm-water ecotypes even during winter months, potentially providing sufficient prey to support a population of warm-water ecotype specific phages. If concentration and enrichment of viruses during isolation is sufficient to successfully isolate even low abundance phages then a comprehensive library of representative phage isolates could be generated with relatively modest sampling effort across a few locations.

First methylophages for marine OM43 isolated.

Viral isolation of novel methylophages for the OM43 clade yielded 23 positive infections, with efficiencies ranging from 0% (no viruses isolated on host C6P1) to 45% on H5P1 (Fig. 2). To the best of our knowledge these are the first reported viruses infecting members of the OM43 clade. One explanation for the lower efficiency of isolation of OM43 viruses is simply one of lower host abundance concomitant with lower phage abundance in the viral community, reducing the likelihood of infective viruses coming into contact with susceptible and permissive cells. OM43 are closely associated with metabolism of extracellular substrates from phytoplankton blooms [47], but have low abundance outside of phytoplankton spring blooms [32]. Our water samples were not associated with high *in-situ* fluorescence (used as proxy measurement for phytoplankton), and missed the April 2019 Spring bloom by about two weeks (Supplementary Fig. 7). Therefore, OM43 and their viruses were likely of lower abundance at sampling times, decreasing the success rate of viral isolation from these samples.

New viruses represent novel viral populations and support established ANI cut-offs for ecologically discrete viral ecotypes

Following three rounds of purification through Dilution-to-Extinction, due to the rate-limiting step of extracting viral DNA, we subselected 16 viral isolates based on availability in November 2018 across four different hosts (HTCC1062, HTCC7211, H2P3α and H5P1) for sequencing at >30-fold coverage

(Table 2). Three out of 16 sequenced samples (two from host HTCC7211, one from OM43 host H5P1)

failed to assemble into single viral contigs. For 11 of the remaining 13 samples (12 from SAR11 hosts

and one from OM43) each individual sequence assembly was identified as a complete viral genome

by VirSorter [15] and 95-100% complete using CheckV [48]. OM43 phage *Venkman* possesses a

linear but complete genome (Supplementary Table 4). Sequence data yielded a single viral genome

(categories 1 or 2, >15kbp) per sample, indicating that our purification process was effective in

recovering pure viral isolates.

Viral populations are defined as discrete ecological and evolutionary units that form non-overlapping

clouds of sequence space, and average nucleotide identity (ANI) cutoffs of >95% have been proposed

from cyanophage studies to delineate viral populations [49]. To evaluate whether our new isolates

represented discrete ecological units with respect to other known pelagiphages, pairwise ANI was

calculated between the thirteen successfully sequenced viral genomes from this study and 85 other

known or putative pelagiphages [37, 46, 50]. Pairwise ANI ranged between 77.5-100%, with a

discrete distribution between 96.4-100.0% (Supplementary Fig. 8, Supplementary Table 5). This is

in agreement with previous work in cyanophages [49, 51] and thus supports their application to

define viral populations beyond cyanophages for viral genomes derived from metagenomic assembly

[10, 11]. At proposed ANI cutoffs of 95% over 85% length [18], the new viruses clustered into six

viral populations, ranging from singletons to a viral population with four members (Table 2). Phages

within the same populations were all isolated from the same source water sample and on the same

host, in agreement with their classification as discrete ecological and evolutionary units. Interestingly,

viral population membership was exclusive to viruses isolated in this study, with no associated

populations containing representatives from either known isolates [39, 46] or fosmid-derived [37]

genomes from other studies, including pelagiphage HTVC010P [39], which is ubiquitous and

dominant throughout global oceans, including the Western English Channel [12]. Such a distinct

separation between populations associated with viruses sampled in this study and all other

pelagiphages, either in culture or assembled from metagenomes, suggests a high degree of viral

population diversity remains to be discovered in the Western English Channel and beyond.

Current pelagiphage isolates can be organised into five distinct phylogenetic clades

To evaluate isolate diversity at higher taxonomic organisation, chosen representatives of each of the

six viral populations were compared to previous isolates and fosmid-derived phage sequences using

three approaches: First, where possible, phylogenetic analysis was performed based on conserved

genes; Second, to capture broader relationships and account for genomic mosaicism, raw

hypergeometric probability of shared gene content; Finally genomes were organised into ICTV-

recognized genera using vConTACT2 (Supplementary Fig. 9), which initially derives viral clusters

using a similar hypergeometric approach followed by refinement with Euclidean-distance-based

hierarchical clustering to split mismatched, 'lumped' viral clusters [19]. All three approaches were

congruent - clustering on probability of shared gene content organising pelagiphage genomes into

four main clusters and one singleton genome from this study (*Kolga*) (Fig. 3A-C), which was broadly

supported by phylogenetic and vConTACT2 classifications. Cluster A contained 23 members (nine

from fosmid-derived contigs [37]; eleven previously isolated pelagiphages [46, 52]), including *Ran*,

Bylgja and *Eistla* from this study. Cluster B contained two previously isolated pelagiphages, one

fosmid-derived contig and *Eyrgjafa* from this study. All viruses in Clusters A and B were assigned to

a single viral genus by vConTACT2 that also contained 12 previously isolated pelagiphages [39, 46].

Cluster C only contained fosmid-derived contigs from the Mediterranean [37], with no isolated

representatives, making it an important location for future isolation attempts of different hosts and

viruses using our method. Cluster D contained eight fosmid-derived contigs, pelagiphage

HTVC010P, and a new representative, *Greip*, from this study (Fig. 3B). VConTACT2 split Cluster

D, leaving *Greip* and *Kolga* as members of two singleton clusters, suggesting they are the first

cultured representatives of novel viral genera.

Kolga - the first siphovirus infecting SAR11

The 25 previously known viral isolates infecting SAR11 comprise 24 podoviruses and one myovirus [39, 46, 53]. Previous cultivation efforts for viruses of SAR11 has not isolated any siphoviruses, nor are any known from viral metagenomic studies. We isolated and sequenced the two viruses *Kolga* and *Aegir* using H2P3α as bait, which to the best of our knowledge are the first known siphoviruses infecting members of the SAR11 clade (Fig. 4A-C). The phages appeared to have a long tail on transmission electron microscopy (TEM) images and are thus morphologically indicative of belonging to the *Siphoviridae* (Fig. 4B). *Aegir* and *Kolga* were members of the same population using a boundary cutoff of 95% ANI over 85% contig length, however, *Aegir* had a length of 18,297 bp compared to 48,659 bp in *Kolga*, therefore we considered *Aegir* to be a partial genome of the same viral population. *Kolga* did not share a significant number of genes with known SAR11 podoviruses (Fig. 3B), and did not cluster with other pelagiphages using a hypergeometric analysis based on gene content. VConTACT2 also grouped *Aegir* and *Kolga* into one cluster without any other known viruses, suggesting it represents a novel viral genus. A number of genes found in *Kolga* were shared with other known siphoviruses, and phylogenetic analysis of concatenated genes shows *Kolga* is most closely related to LK3, a known siphovirus infecting the Betaproteobacterial genus *Bordetella* (Fig. 4C). This suggests that *Kolga* is more closely related to siphoviruses of different hosts compared to podoviruses infecting the same host. 67% of genes encoded by Kolga could not be functionally annotated, and out of all hypothetical genes identified on *Kolga*, only three hypothetical genes were shared with SAR11 podoviruses. In contrast, for on average ~90% of genes without known function identified within our novel SAR11 podoviruses are shared between different pelagiphages. *Kolga* was the 10th most abundant viral isolate in our Western English Channel virome (Fig. 3D). *Kolga* possesses a tail tip J protein (Fig. 4A), often found in phages with long non-contractile tails such as λ, where it plays a role in DNA injection during cell entry and tail assembly [54]. *Kolga* also encodes

a small S21 subunit of the 30S ribosomal gene structurally similar to the ones found in HTVC008M,

and hosts HTCC7211 and H2P3α, a feature found in numerous myoviruses and sipoviruses [55]. S21

is involved in translation initiation and needed for the binding of mRNA [56]. Virally-encoded S21

genes may provide a competitive advantage for the phage as it could replace cellular S21 and assist

in the translation of viral transcripts. *Kolga* may need its S21 gene for shifting the translational frame,

as it has been shown that for some members of the Caudovirales the production of tail components is

dependent on programmed translational frameshifting [57]. Given the constitutive nature of gene

expression in genomically streamlined bacteria [58], genes such as S21 may provide the virus with a

mechanism to manipulate host metabolism in the absence of typical promoters and repressors.

Kolga and Greip are the most abundant of the novel viruses in Western English Channel

To evaluate relative abundance of our new isolates in the Western English Channel, we used

competitive mapping of metagenomic reads from a Western English Channel virome taken in

September 2016 (*in situ* surface water temperature of ~16.1 °C) [12] against existing pelagiphage

genomes and those from this study using FastViromeExplorer [59]. This identified HTVC010P as

the most abundant virus within the known pelagiphages (536,531 Reads Per Million [RPM - see

Supplementary Methods]). Other isolates from this and previous studies recruited 1-3 orders of

magnitude fewer reads than HTVC010P (684 - 53,591 RPM), indicating that viral community

structures for pelagiphages in the Western English Channel consist of a small number of viral species

that are highly abundant, and long tail of lower abundance members (Fig. 3D). *Greip* and *Kolga* were

the two most abundant of our new viral isolates in the Western English Channel. Their abundance

could be driven in part by the fact that either the *in situ* surface water temperature when the virome

sample was taken matches the optimum growth temperature of host HTCC1062 (~0.44 divisions per

day at 16 °C) [44], in line with proposed Kill-the-Winner predator/prey dynamics [60], and/or that

genomic novelty promotes recruitment of reads closely related taxa that have no other representation

in the reference dataset [39]. Future work will evaluate the global abundance of these and newly

sequenced isolates across spatial (Global Ocean Viromes) and temporal gradients (an ongoing time-

series project in the Western English Channel) to evaluate their biogeography.

Venkman - the first known virus infecting OM43

Our isolation efforts in the Western English Channel yielded the first cultured bacteriophage infecting

the methylotroph OM43 (Fig. 5A-C). Named *Venkman*, in homage to the actor Bill Murray in the

film Ghostbusters, this virus has a genome 38,624 bp long (31.9% GC content), encoding genes (Fig.

5A) with similar synteny and function to the siphovirus P19250A (38,562 bp) that infects freshwater

Methylophilales LD28, which are often considered a freshwater variant of OM43 [61, 62]. Unlike

P19250A, TEM images indicated that *Venkman* had a short tail (Fig. 5B) similar to podoviruses,

though it is possible that tail structures had been lost during grid preparation. Phylogenetic analysis

of concatenated TerL and exonuclease genes further indicated that these genes in *Venkman* are most

closely related to other siphoviruses infecting different Proteobacteria (Fig. 5C). However, branch

support values were low, despite numerous attempts to refine the tree with different approaches (see

Supplementary Methods). We identified a number of common phage proteins such as a capsid

protein, terminase, nucleases and tail structural proteins, the remaining 54% of genes were

hypothetical. VConTACT2 assigned *Venkman* and P19250 to the same genus-level cluster, therefore

Venkman may be a marine variant of the freshwater phage P19250. In the Western English Channel

viral metagenomes, *Venkman*-like viruses were the second most abundant viruses (148,681 RPM)

after HTVC010P (536,531 RPM) identifying them as ecologically important viruses in this coastal

microbial community (Fig. 3B).

Unusual host-virus dynamics are prevalent in isolated phages

Viral isolates *Bylgja*, *Eyrgjafa*, *Ran* and *Eistla* (Cluster A and B) are presumed to be temperate phages, with all encoding endonucleases and exonucleases (Supplementary Fig. 10) and clustering by shared protein content with other pelagiphages, such as e.g. HTVC011P and HTVC025P, shown previously to integrate into host genomes (Fig. 3B) [46]. *Eyrgjafa* encodes a tRNA-Leu, that has 85% nucleotide identity over the first 34 bases of the tRNA-Leu of its host HTCC1062, suggesting a putative integration site into the host genome [63]. To date, 16 of the 29 viruses isolated on SAR11 strains have either been shown to be capable of lysogeny, and/or encode genes associated with a temperate infection cycle. In contrast, viruses such as *Greip,* and the abundant HTVC010P in Cluster D (Fig. 3B) do not possess any genes associated with lysogeny, and would therefore presumed to be exclusively lytic. Viruses in Clusters A and B, presumed temperate, were of much lower abundance in the environment compared to *Greip* or HTVC010P in Cluster D which were among the most abundant pelagiphages in the Western English Channel (Fig. 3D), suggesting a possible ecological difference between two groups of viruses with different infectivity strategies.

Interestingly, growth curves of hosts infected with *Greip* and other isolated phages deviated from the expected decay in cell abundance associated with viral lysis and previously observed in isolated pelagiphages [39]. The first pelagiphages HTVC010P and HTVC008M were isolated from the warm waters of the Sargasso Sea, and HTVC011P and HTVC019P were isolated from the colder waters of the Oregon Coast. All four strains were propagated on the cold-water SAR11 ecotype *P. ubique* HTCC1062. In all cases, host density was reduced from ~8×10^6 cells per mL at T_0 to <10^6 cells per mL over a 60-72h period, although viruses from warm waters took 17% longer than those from cold water to do so [39], suggesting that suboptimal hosts reduced the rate of infection as shown in cyanophages [64]. In contrast, infection dynamics of our isolates often resulted in host density of infected cultures growing to a steady state, but at a lower cell density than uninfected cells. (Supplementary Video 1), irrespective of cluster or population assignment. Out of 117 viruses isolated in this study, only 16 infections reduced host abundance below their inoculum density of 10^6 cells per

mL. In 53 infections, densities of infected cells increased to within an order of magnitude of uninfected cells (Supplementary Fig. 11), but demonstrated clear evidence of viral infection in cytograms, TEMs and subsequent recovery of viral genomes in selected samples.

Similar patterns of infection were recently reported in the extremely abundant bacteriophage ɸCrAss001 found in the human gut, where infection of exponentially growing cells of *B. intestinalis* 919/174 did not result in complete culture lysis, but caused a delay in stationary phase onset time and final density, despite lacking genes associated with lysogeny. As with our study, the authors observed that this only occurred in liquid culture, and isolation of the virus required numerous rounds of enrichment. They postulated that the virus may cause a successful infection in only a subset of host cells, with the remainder exhibiting alternative interactions such as pseudolysogeny or dormancy [31]. The prevalence of similar infection dynamics in the phages isolated in this study offer two intriguing possibilities: (i) Many of the viruses isolated in this study are either not fully lytic, but fall somewhere on the continuum of persistence [65], controlled by genes currently lacking a known function, with lysogeny (and associated superinfection immunity) favoured at high cell density, in support of the Piggyback-the-Winner hypothesis [66, 67]; (ii) the steady-state of host and virus densities observed here are an indicator of host phenotypic bistability in these streamlined heterotrophic taxa. Viral propagation occurring in only a subset of cells could explain the requirement of multiple rounds of enrichment before sufficient viral load is reached to be able to observe lytic infection on the host population. Either strategy, or a combination of both, would provide an ecological advantage of long-term stable coexistence between viruses and hosts and offer an explanation of the paradox of stable high abundances of both predator and prey across global oceans [12, 39, 68, 69]. Infection in a subset of the population could also explain the low lytic activity observed in pelagiphages *in situ*, despite high host densities [70]; and the small dynamic range and decoupled abundances of SAR11 and virioplankton in the Sargasso Sea [71]. Limited lysis of subpopulations of hosts such as SAR11 and OM43 that specialise in the uptake of labile carbon

enriched through viral predation [72, 73] would facilitate efficient intra-population carbon recycling

and explain the limited influence of SAR11 and associated viral abundances on carbon export to the

deep ocean [5]. We propose the moniker the 'Soylent Green Hypothesis' for this mechanism, after

the 1973 cult film in which the dead are recycled into food for the living. Further investigation

leveraging our new virus-host model will provide greater insight into viral influence on ocean carbon

biogeochemistry.

In conclusion, our method coupled Dilution-to-Extinction cultivation of hosts and associated viruses,

resulting in the isolation of three new strains of OM43; a Western English Channel variant of a warm

water ecotype of SAR11; the first known methylophages for OM43; the first siphovirus infecting

SAR11, as well as eleven other viruses infecting this important marine heterotrophic clade and >100

more isolates to be sequenced and explored. The described method represents an efficient and cost-

effective strategy to isolate novel virus-host systems for experimental evaluation of co-evolutionary

dynamics of important fastidious taxa from marine and other biomes. Coupling these methods to

existing advances in host cultivation requires minimal additional effort and will provide valuable

representative genomes to improve success rates of assigning putative hosts to metagenomically-

derived viral contigs. Broader representation of model systems beyond cyanophages and viruses of

copiotrophic, r-strategist hosts will reduce bias in developing methods to delineate viral population

boundaries [74, 75], increasing the accuracy with which we derive ecological meaning from viral

metagenomic data. We therefore hope that this method will enable viruses to be included in the

current resurgence of cultivation efforts to better understand the biology and ecology of phages, and

the influence of the world's smallest predators on global biogeochemistry.

Methods Summary

A complete description of the materials and methods is provided in the Supplementary Information. Four bacterial strains (*Methylophilales sp*. C6P1, D12P1 and H5P1; *Pelagibacter sp*. H2P3α) were isolated from Western English Channel station L4 seawater samples using Dilution-to-Extinction methods [28]. All four bacteria and two additional SAR11 strains *Pelagibacter bermudensis* HTCC7211 and *Pelagibacter ubique* HTCC1062 were used as bait to isolate phages from six monthly Western English Channel L4 seawater samples (50°15.00N; 4°13.00W). Briefly, water samples were concentrated for viruses using tangential flow filtration and used as viral inoculum (10% v/v) in exponentially growing cultures of host bacteria in artificial seawater medium [76] in 96-well Teflon plates (Radleys, UK). Cells of the resulting lysate were filtered out and the filtrate was used as viral inoculum in another round of isolation. This process was repeated until viral infection could be detected by flow cytometry - comparing cytograms and maximum density of infected cultures against uninfected cultures. Phages were purified by dilution-to-extinction methods (detailed protocol available here: dx.doi.org/10.17504/protocols.io.c36yrd). Phage genomes were sequenced using Illumina 2x150 PE sequencing, assembled and manually annotated as described in [77]. Phylogenetic classification of phages was performed on concatenated shared genes using a combination of Bayesian inference trees, maximum likelihood trees and shared-gene likelihood analyses, depending on the availability of appropriate taxa. ICTV-recognised genera based on shared gene content were assigned with VConTACT2 [19]. The relative abundance of novel phages in the Western English Channel was calculated by competitive read recruitment using FastViromeExplorer v1.3 [59] against an existing virome from the same location [12].

Data availability All reads can be found in the SRA database under BioProject number PRJNA625644 as BioSamples SAMN14604128-SAMN14604140. Annotated phage genomes are deposited as GenBank submissions under accession numbers MT375519- MT375531.

Acknowledgements We would like to thank Christian Hacker and the Bioimaging Centre of the University of Exeter for performing the TE microscopy and imaging. We would also like to thank the

crew of the *R/V Plymouth Quest* and our collaborators at PML for collecting water samples, and the

driver Magic for delivering water samples from Plymouth to Exeter. Genome sequencing was

provided by MicrobesNG (http://www.microbesng.uk) which is supported by the BBSRC (grant

number BB/L024209/1).

Funding information The efforts of Holger Buchholz in this work were funded by the Natural

Environment Research Council (NERC) GW4+ Doctoral Training program. Michelle Michelsen and

Ben Temperton were funded by NERC (NE/R010935/1) and by the Simons Foundation BIOS-

SCOPE program.

Conflict of interests The authors declare no conflict of interest.

References

1. Eric Wommack K, Colwell RR. Virioplankton: Viruses in Aquatic Ecosystems. *MICROBIOLOGY AND*

MOLECULAR BIOLOGY REVIEWS 2000; **64**: 69–114.

2. Weitz JS, Stock CA, Wilhelm SW, Bourouiba L, Coleman ML, Buchan A, et al. A multitrophic model

to quantify the effects of marine viruses on microbial food webs and ecosystem processes. *ISME J*

2015; **9**: 1352–1364.

3. Weitz JS, Wilhelm SW. Ocean viruses and their effects on microbial communities and biogeochemical

cycles. *F1000 Biol Rep* 2012; **4**: 17.

4. Jover LF, Effler TC, Buchan A, Wilhelm SW, Weitz JS. The elemental composition of virus particles:

implications for marine biogeochemical cycles. *Nat Rev Microbiol* 2014; **12**: 519–528.

5. Guidi L, Chaffron S, Bittner L, Eveillard D, Larhlimi A, Roux S, et al. Plankton networks driving

carbon export in the oligotrophic ocean. *Nature* 2016; **532**: 465–470.

6. Warwick-Dugdale J, Buchholz HH, Allen MJ, Temperton B. Host-hijacking and planktonic piracy: how

phages command the microbial high seas. *Virol J* 2019; **16**: 15.

7.   Weitz JS, Hartman H, Levin SA. Coevolutionary arms races between bacteria and bacteriophage. *Proc*

Natl Acad Sci U S A 2005; **102**: 9535–9540.

8.   Ignacio-Espinoza JC, Ahlgren NA, Fuhrman JA. Long-term stability and Red Queen-like strain

dynamics in marine viruses. *Nat Microbiol* 2020; **5**: 265–271.

9.   Brum JR, Hurwitz BL, Schofield O, Ducklow HW, Sullivan MB. Seasonal time bombs: dominant

temperate viruses affect Southern Ocean microbial dynamics. *ISME J* 2016; **10**: 437–449.

10.   Roux S, Brum JR, Dutilh BE, Sunagawa S, Duhaime MB, Loy A, et al. Ecogenomics and potential

biogeochemical impacts of globally abundant ocean viruses. *Nature* 2016; **537**: 689–693.

11.   Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A, et al. Marine DNA

Viral Macro- and Microdiversity from Pole to Pole. *Cell* 2019; **177**: 1109–1123.e14.

12.   Warwick-Dugdale J, Solonenko N, Moore K, Chittick L, Gregory AC, Allen MJ, et al. Long-read viral

metagenomics captures abundant and microdiverse viral populations and their niche-defining genomic

islands. *PeerJ* 2019; **7**: e6800.

13.   Mann NH, Cook A, Millard A, Bailey S, Clokie M. Bacterial photosynthesis genes in a virus. *Nature*

2003; **424**: 741.

14.   Puxty RJ, Millard AD, Evans DJ, Scanlan DJ. Viruses Inhibit CO2 Fixation in the Most Abundant

Phototrophs on Earth. *Curr Biol* 2016; **26**: 1585–1589.

15.   Roux S, Enault F, Hurwitz BL, Sullivan MB. VirSorter: mining viral signal from microbial genomic

data. *PeerJ* 2015; **3**: e985.

16.   Ren J, Ahlgren NA, Lu YY, Fuhrman JA, Sun F. VirFinder: a novel k-mer based tool for identifying

viral sequences from assembled metagenomic data. *Microbiome* 2017; **5**: 69.

17.   Amgarten D, Braga LPP, da Silva AM, Setubal JC. MARVEL, a Tool for Prediction of Bacteriophage

Sequences in Metagenomic Bins. *Front Genet* 2018; **9**: 304.

18.   Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an in silico evaluation

of metagenome-enabled estimates of viral community composition and diversity. *PeerJ* 2017; **5**: e3817.

19.   Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Taxonomic assignment

of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nat Biotechnol* 2019.

20.   Hurwitz BL, Brum JR, Sullivan MB. Depth-stratified functional and taxonomic niche specialization in

the 'core' and 'flexible' Pacific Ocean Virome. *ISME J* 2015; **9**: 472–484.

21. Galiez C, Siebert M, Enault F, Vincent J, Söding J. WIsH: who is the host? Predicting prokaryotic hosts

from metagenomic phage contigs. *Bioinformatics* 2017; **33**: 3113–3114.

22. Ahlgren NA, Ren J, Lu YY, Fuhrman JA, Sun F. Alignment-free d_2^* oligonucleotide frequency

dissimilarity measure improves prediction of hosts from metagenomically-derived viral sequences.

Nucleic Acids Res 2017; **45**: 39–53.

23. Sun J, Steindler L, Thrash JC, Halsey KH, Smith DP, Carter AE, et al. One carbon metabolism in

SAR11 pelagic marine bacteria. *PLoS One* 2011; **6**: e23973.

24. Becker JW, Hogle SL, Rosendo K, Chisholm SW. Co-culture and biogeography of Prochlorococcus and

SAR11. *ISME J* 2019.

25. Kraemer S, Ramachandran A, Colatriano D, Lovejoy C, Walsh DA. Diversity and biogeography of

SAR11 bacteria from the Arctic Ocean. *ISME J* 2020; **14**: 79–90.

26. Robert M. Morris*, Michael S. Rappe´ *, Stephanie A. Connon*, Kevin L. Vergin*, William A.

Siebold*, Craig A. Carlson† & Stephen J. Giovannoni*. SAR11 clade dominates ocean surface

bacterioplankton communities. *Nature* 2002; **420**.

27. Connon SA, Giovannoni SJ. High-throughput methods for culturing microorganisms in very-low-

nutrient media yield diverse new marine isolates. *Appl Environ Microbiol* 2002; **68**: 3878–3885.

28. Henson MW, Pitre DM, Weckhorst JL, Celeste Lanclos V, Webber AT, Cameron Thrash J. Artificial

Seawater Media Facilitate Cultivating Members of the Microbial Majority from the Gulf of Mexico.

American Society for Microbiology 2016; **1**: 1–9.

29. Henson MW, Lanclos VC, Faircloth BC, Thrash JC. Cultivation and genomics of the first freshwater

SAR11 (LD12) isolate. *ISME J* 2018; **12**: 1846–1860.

30. Nagasaki K, Bratbak G. Isolation of viruses infecting photosynthetic and nonphotosynthetic protists.

Manual of aquatic viral ecology ASLO 2010; 92–101.

31. Shkoporov AN, Khokhlova EV, Fitzgerald CB, Stockdale SR, Draper LA, Ross RP, et al. ΦCrAss001

represents the most abundant bacteriophage family in the human gut and infects Bacteroides intestinalis.

Nat Commun 2018; **9**: 4781.

32. Morris RM, Longnecker K, Giovannoni SJ. Pirellula and OM43 are among the dominant lineages

identified in an Oregon coast diatom bloom. *Environ Microbiol* 2006; **8**: 1361–1370.

33.　Ramachandran A, Walsh DA. Investigation of XoxF methanol dehydrogenases reveals new

methylotrophic bacteria in pelagic marine and freshwater ecosystems. *FEMS Microbiol Ecol* 2015; **91**.

34.　Halsey KH, Carter AE, Giovannoni SJ. Synergistic metabolism of a broad range of C1 compounds in

the marine methylotrophic bacterium HTCC2181. *Environ Microbiol* 2012; **14**: 630–640.

35.　Reintjes G, Arnosti C, Fuchs B, Amann R. Selfish, sharing and scavenging bacteria in the Atlantic

Ocean: a biogeographical study of bacterial substrate utilisation. *ISME J* 2019; **13**: 1119–1132.

36.　Carini P, White AE, Campbell EO, Giovannoni SJ. Methane production by phosphate-starved SAR11

chemoheterotrophic marine bacteria. *Nat Commun* 2014; **5**: 4346.

37.　Mizuno CM, Rodriguez-Valera F, Kimes NE, Ghai R. Expanding the marine virosphere using

metagenomics. *PLoS Genet* 2013; **9**: e1003987.

38.　Martinez-Hernandez F, Fornas O, Lluesma Gomez M, Bolduc B, de la Cruz Peña MJ, Martínez JM, et

al. Single-virus genomics reveals hidden cosmopolitan and abundant viruses. *Nat Commun* 2017; **8**:

15892.

39.　Zhao Y, Temperton B, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC, et al. Abundant SAR11

viruses in the ocean. *Nature* 2013; **494**: 357–360.

40.　Taubert M, Grob C, Howat AM, Burns OJ, Dixon JL, Chen Y, et al. XoxF encoding an alternative

methanol dehydrogenase is widespread in coastal marine environments. *Environ Microbiol* 2015; **17**:

3937–3948.

41.　Schwalbach MS, Tripp HJ, Steindler L, Smith DP, Giovannoni SJ. The presence of the glycolysis

operon in SAR11 genomes is positively correlated with ocean productivity. *Environ Microbiol* 2010;

12: 490–500.

42.　Giovannoni SJ, Hayakawa DH, Tripp HJ, Stingl U, Givan SA, Cho J-C, et al. The small genome of an

abundant coastal ocean methylotroph. *Environ Microbiol* 2008; **10**: 1771–1782.

43.　Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer K-H, et al. Uniting the classification

of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Microbiol*

2014; **12**: 635–645.

44.　Polyphasic taxonomy of Marine bacteria. *studylib.net*. https://studylib.net/doc/5577926/polyphasic-

taxonomy-of-marine-bacteria. Accessed 11 May 2020.

45. Delmont TO, Kiefl E, Kilinc O, Esen OC, Uysal I, Rappé MS, et al. Single-amino acid variants reveal

evolutionary processes that shape the biogeography of a global SAR11 subclade. *Elife* 2019; **8**.

46. Zhao Y, Qin F, Zhang R, Giovannoni SJ, Zhang Z, Sun J, et al. Pelagiphages in the Podoviridae family

integrate into host genomes. *Environ Microbiol* 2018.

47. Neufeld JD, Boden R, Moussard H, Schäfer H, Murrell JC. Substrate-specific clades of active marine

methylotrophs associated with a phytoplankton bloom in a temperate coastal environment. *Appl Environ*

Microbiol 2008; **74**: 7321–7328.

48. Nayfach S, Camargo AP, Eloe-Fadrosh E, Roux S, Kyrpides N. CheckV: assessing the quality of

metagenome-assembled viral genomes. *bioRxiv* . 2020. , 2020.05.06.081778

49. Deng L, Ignacio-Espinoza JC, Gregory AC, Poulos BT, Weitz JS, Hugenholtz P, et al. Viral tagging

reveals discrete populations in Synechococcus viral genome sequence space. *Nature* 2014; **513**: 242–

245.

50. Våge S, Storesund JE, Thingstad TF. SAR11 viruses and defensive host strains. *Nature* . 2013. , **499**:

E3–4

51. Marston MF, Martiny JBH. Genomic diversification of marine cyanophages into stable ecotypes.

Environ Microbiol 2016; **18**: 4240–4253.

52. Giovannoni S, Temperton B, Zhao Y. Giovannoni et al. reply. *Nature* . 2013. , **499**: E4–5

53. Zhang Z, Qin F, Chen F, Chu X, Luo H, Zhang R, et al. Novel pelagiphages prevail in the ocean.

bioRxiv . 2019. , 2019.12.14.876532

54. Tam W, Pell LG, Bona D, Tsai A, Dai XX, Edwards AM, et al. Tail tip proteins related to

bacteriophage λ gpL coordinate an iron-sulfur cluster. *J Mol Biol* 2013; **425**: 2450–2462.

55. Mizuno CM, Guyomar C, Roux S, Lavigne R, Rodriguez-Valera F, Sullivan MB, et al. Numerous

cultivated and uncultivated viruses encode ribosomal proteins. *Nat Commun* 2019; **10**: 752.

56. Van Duin J, Wijnands R. The function of ribosomal protein S21 in protein synthesis. *Eur J Biochem*

1981; **118**: 615–619.

57. Xu J, Hendrix RW, Duda RL. Conserved translational frameshift in dsDNA bacteriophage tail assembly

genes. *Mol Cell* 2004; **16**: 11–21.

58. Giovannoni SJ, Thrash JC, Temperton B. Implications of streamlining theory for microbial ecology.

ISME J 2014; **8**: 1553–1565.

59. Tithi SS, Aylward FO, Jensen RV, Zhang L. FastViromeExplorer: a pipeline for virus and phage

identification and abundance profiling in metagenomics data. *PeerJ* 2018; **6**: e4227.

60. Thingstad TF, Lignell R. Theoretical models for the control of bacterial growth rate, abundance,

diversity and carbon demand. *Aquat Microb Ecol* 1997; **13**: 19–27.

61. Moon K, Kang I, Kim S, Kim S-J, Cho J-C. Genome characteristics and environmental distribution of

the first phage that infects the LD28 clade, a freshwater methylotrophic bacterial group. *Environ*

Microbiol 2017; **19**: 4714–4727.

62. Salcher MM, Neuenschwander SM, Posch T, Pernthaler J. The ecology of pelagic freshwater

methylotrophs assessed by a high-resolution monitoring and isolation campaign. *ISME J* 2015; **9**: 2442–

2453.

63. Bailly-Bechet M, Vergassola M, Rocha E. Causes for the intriguing presence of tRNAs in phages.

Genome Res 2007; **17**: 1486–1495.

64. Enav H, Kirzner S, Lindell D, Mandel-Gutfreund Y, Béjà O. Adaptation to sub-optimal hosts is a driver

of viral diversification in the ocean. *Nat Commun* 2018; **9**: 4698.

65. Weitz JS, Li G, Gulbudak H, Cortez MH, Whitaker RJ. Viral invasion fitness across a continuum from

lysis to latency. *Virus Evol* 2019; **5**: vez006.

66. Knowles B, Silveira CB, Bailey BA, Barott K, Cantu VA, Cobian-Guemes AG, et al. Lytic to temperate

switching of viral communities. *Nature* 2016; **531**: 466–470.

67. Silveira CB, Rohwer FL. Piggyback-the-Winner in host-associated microbial communities. *NPJ*

Biofilms Microbiomes 2016; **2**: 16010.

68. Kang I, Oh HM, Kang D, Cho JC. Genome of a SAR116 bacteriophage shows the prevalence of this

phage type in the oceans. *Proc Natl Acad Sci U S A* 2013; **110**: 12343–12348.

69. Martinez-Hernandez F, Fornas Ò, Lluesma Gomez M, Garcia-Heredia I, Maestre-Carballa L, López-

Pérez M, et al. Single-cell genomics uncover Pelagibacter as the putative host of the extremely abundant

uncultured 37-F6 viral population in the ocean. *ISME J* 2018.

70. Alonso-Sáez L, Morán XAG, Clokie MR. Low activity of lytic pelagiphages in coastal marine waters.

ISME J 2018; **12**: 2100–2102.

71. Parsons RJ, Breitbart M, Lomas MW, Carlson CA. Ocean time-series reveals recurring seasonal

patterns of virioplankton dynamics in the northwestern Sargasso Sea. *ISME J* 2012; **6**: 273–284.

72. Middelboe M, Lyck PG. Regeneration of dissolved organic matter by viral lysis in marine microbial

communities. *Aquat Microb Ecol* 2002; **27**: 187–194.

73. Middelboe M, Jorgensen NOG. Viral lysis of bacteria: an important source of dissolved amino acids

and cell wall compounds. *J Mar Biol Assoc U K* 2006; **86**: 605–612.

74. Brum JR, Sullivan MB. Rising to the challenge: accelerated pace of discovery transforms marine

virology. *Nat Rev Microbiol* 2015; **13**: 147–159.

75. Gregory AC, Solonenko SA, Ignacio-Espinoza JC, LaButti K, Copeland A, Sudek S, et al. Genomic

differentiation among wild cyanophages despite widespread horizontal gene transfer. *BMC Genomics*

2016; **17**: 930.

76. Carini P, Steindler L, Beszteri S, Giovannoni SJ. Nutrient requirements for growth of the extreme

oligotroph 'Candidatus Pelagibacter ubique' HTCC1062 on a defined medium. *ISME J* 2013; **7**: 592–

602.

77. Salisbury A, Tsourkas PK. A Method for Improving the Accuracy and Efficiency of Bacteriophage

Genome Annotation. *Int J Mol Sci* 2019; **20**.

Fig. 1 Workflow for high-throughput isolation **(A)** i. Increasing concentration of viruses in water

samples by TFF; ii. Initial infection of host cultures to enrich the sample for specific viruses; iii.

Purification of viral isolates through 3 rounds of Dilution-to-Extinction; **(B)** Initial screen of viral

infections using i. Flow cytometry, by comparing populations of no-virus controls and infected

cultures; ii. Comparing growth curves of no-virus control culture (HTCC1062) against infected

SAR11 cultures; iii. Confirming the presence of viruses in infected SAR11 cultures using TE

microscopy: top left: HTCC1062 no-virus control, bottom left: infected HTCC1062, top right:

aggregated cellular debris and viruses, bottom right: virus found in infected HTCC1062 culture.

Successful isolations / Attempted isolations

Fig. 2 Summary of success rates for each bacterial host (three SAR11 and three OM43) and water sample combination used for infections; the numerator indicates successful isolation of a virus from that water sample (i.e. successful passage through three rounds of Dilution-to-Extinction), denominator shows the isolation attempts made for a host-water sample combination (i.e. how many cultures were treated with viral samples after multiple rounds of enrichment).

Fig. 3 Phylogeny and abundance of the first sequenced pelagiphages from the Western English

Channel **(A)** Hypergeometric testing on viral genomes by shared protein content identified four viral

clusters and numerous singletons; **(B)** Bayesian inference phylogenetic tree of pelagiphage major

capsid proteins from cluster A and B (most diverse pelagiphage cluster, includes most known

pelagiphages isolated previously [39, 46] as well as derived from metagenomes [37]), outgroup

Cluster D; **(C)** Bayesian inference phylogenetic tree of pelagiphage major capsid proteins from

Cluster D (containing the most abundant pelagiphages, including HTVC010P) with Cluster A as

outgroup; **(D)** Metagenomic recruitment (reads recruited per million reads (RPM)) of isolate and

metagenomically-derived viral genomes against a virome from the Western English Channel [12].

Fig. 4 The first reported siphovirus *Kolga* infecting SAR11 and isolated on novel host H2P3α ; **(A)**

gene map of the 48,784 bp genome, which contains 80% hypothetical genes without known function;

(B) TEMs of *Kolga* (top left and bottom) and a H2P3α host cell infected with *Kolga* (top right); **(C)**

Bayesian inference phylogenetic tree of concatenated genes (Pyrophosphohydrolase, thymidylate

synthase, two ribonucleotide reductase subunits), all branch support values of 1 have been omitted

for clarity

Fig. 5 Evidence for the first reported virus EXEVC282S *Venkman* infecting a member of the OM43

clade (H5P1); **(A)** Gene map displaying protein coding genes. Structural genes are shown in blue,

DNA replication genes in red, lysis related genes in green, transcription genes in turquoise, packaging

genes in orange, hypothetical genes are grey; **(B)** TEM images of: infected and chaining H5P1 cells

(top left), uninfected H5P1 chaining cells (top right), *Venkman* viral particles (bottom left and right);

(C) Maximum likelihood tree (500 bootstraps) of concatenated viral TerL and exonuclease genes,

host families of the phages are indicated on the figure.

Table 1. Host infectivity of viral populations isolated and sequenced in this study. H2P3 and

HTCC7211 are warm-water ecotypes of *Pelagibacter spp*. subclade 1a; HTCC1062 is a cold-water

ecotype of *Pelagibacter spp.* Phage *Bylgja* is the only virus known to infect both ecotypes.

Phage	H2P3	HTVC1062	HTVC7211
Eistla		+	
Eyrgjafa		+	
Greip		+	
Ran	+		+
Kolga	+		+
Bylgja	+	+	+

700   Table 2. Summary of phages isolated and sequenced in this study.

Virus	Host	Viral Population Representative	Hypergeometric cluster	Genome	Length (bp)	G + C %	Taxon	Accession number	Simplified phonetic spelling	Meaning and origin of the names
Eistla	HTCC1062	Eistla	Cluster A	Circular	39638	32.7	Podoviridae	MT375521	aɪsːtlaː	"stormy one", Giantess in the poetic Edda
Eyrgjafa	HTCC1062	Eyrgjafa	Cluster B	Circular	38005	32.6	Podoviridae	MT375523	eːɪrgjaːfaː	"scar donor", Giantess in the poetic Edda
Gjalp	HTCC1062	Eyrgjafa	Cluster B	Circular	37857	32.5	Podoviridae	MT375524	gjaːlp	"Roaring one", Giantess in the poetic Edda
Greip	HTCC1062	Greip	Cluster D	Circular	34916	31.5	Podoviridae	MT375525	graɪp	"Grasp", Giantess in the poetic Edda
Ran	H2P3α	Ran	Cluster A	Circular	41529	34.1	Podoviridae	MT375530	ran	"plundering", Goddess of the Sea
Aegir	H2P3α	Kolga	Singleton	Fragment	18297	31.1	Siphoviridae	MT375519	æːgir	"Sea", God of the Sea
Jörmungand	H2P3α	Ran	Cluster A	Circular	41529	34.1	Podoviridae	MT375528	jɔrmun,gand	"huge monster", Giant sea serpent
Kólga	H2P3α	Kolga	Singleton	Circular	48659	30.5	Siphoviridae	MT375529	kɒlga:	"Cool-wave", Daughter of Ran and Aegir
Unn	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375531	ʊːn	"Wave", Daughter of Ran and Aegir
Hroenn	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375527	hrøn	"Wave", Daughter of Ran and Aegir
Bylgja	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375520	bːlgja:	"Billow"-wave, Daughter of Ran and Aegir
Himinglæva	H2P3α	Bylgja	Cluster A	Circular	41069	33.5	Podoviridae	MT375526	hiːmɪŋglæfa	"transparent-on-top"-wave, Daughter of Ran and Aegir
Venkman	HSP1	Venkman	Singleton	Linear	38624	34.4	Siphoviridae	MT375522	vɛŋkmæn	Bill Murray's Character in the Ghostbusters movie

701