1 Title

2 **The hepatic compensatory response to elevated systemic sulfide**

3 **promotes diabetes.**

4 Authors

Roderick N. Carter[1], Matthew T.G. Gibbins[1,a], Martin E. Barrios-Llerena[1,b], Stephen E. Wilkie[1,2],

Peter L. Freddolino[3], Marouane Libiad[3,c], Victor Vitvitsky[3], Barry Emerson[1,d], Thierry Le

Bihan[4,e], Madara Brice[1], Huizhong Su[5,f], Scott G. Denham[1], Natalie Z.M. Homer[1], Clare Mc

Fadden[1,g], Anne Tailleux[6], Nourdine Faresse[7,h], Thierry Sulpice[7], Francois Briand[7], Tom

Gillingwater[8], Kyo Han Ahn[9], Subhankar Singha[9,i], Claire McMaster[10], Richard C. Hartley[10], Bart

Staels[6], Gillian A. Gray[1], Andrew J. Finch[5,j], Colin Selman[2], Ruma Banerjee[3], Nicholas M.

Morton[1*]

[1]University/British Heart Foundation Centre for Cardiovascular Science, University of Edinburgh, Queen's

Medical Research Institute, Edinburgh, EH16 4TJ, United Kingdom

[2]Glasgow Ageing Research Network (GARNER), Institute of Biodiversity, Animal Health and Comparative

Medicine, University of Glasgow, Glasgow, G12 8QQ, United Kingdom

[3]The Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, MI 48109, USA

[4]SynthSys - Systems and Synthethic Biology, Edinburgh, EH9 3JD, United Kingdom

[5]Cancer Research UK Edinburgh Centre, MRC Intitute of Genetics & Molecular Medicine, University of

Edinburgh, Western General Hospital, Edinburgh, EH4 2XR, United Kingdom

[6]Université de Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U101-EGID, F-59000, Lille, France

[7]Physiogenex S.A.S, Prologue Biotech, 516 rue Pierre et Marie Curie, 31670 Labège, France

[8]College of Medicine & Veterinary Medicine, University of Edinburgh, Old Medical School (Anatomy), Teviot

Place, Edinburgh, EH8 9AG, United Kingdom

[9]Department of Chemistry, POSTECH, 77 Cheongam-Ro, Nam-Gu, Pohang, Gyungbuk, 37673, South Korea

[10]School of Chemistry, Joseph Black Building, University of Glasgow, Glasgow, G12 8QQ, United Kingdom

Current affiliations where different from the place where work for the manuscript was carried out;

[a]Cambridge Biomedical Campus, Royal Papworth Hospital NHS Foundation Trust, Cardiology department,

Cambridge, CB2 0AY, United Kingdom

[b]The International Clinical Research Centre, St. Anne's University Hospital, Brno, 656 91, Czech Republic

[c]Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, 91198,

Gif-sur-Yvette cedex, France

[d]BD Research Centre Ireland, Co. Limerick, Castletroy, Ireland

[e]Rapid Novor Inc, 44 Gaukel St, Kitchener, ON N2G 4P3, Canada

[f]Wellcome Trust Centre for Mitochondrial Research, Newcastle University, NE2 4HH, United Kingdom

[g]Springer Nature Campus, London, N1 9FN, United Kingdom

[h]D.I.V.A-expertise, 1 place Pierre Potier, 31100 Toulouse, France

[i]Institute of Advanced Studies and Research, JIS University, Kolkata 700091, India

[j]Cancer Research UK Barts Centre, London, EC1M 6BQ, United Kingdom

[*]**Lead Contact details:** Nicholas M. Morton

Molecular Metabolism Group, University/British Heart Foundation Centre for Cardiovascular Science,

University of Edinburgh, Queen's Medical Research Institute, Edinburgh, EH16 4TJ, United Kingdom.

nik.morton@ed.ac.uk

Conflict of interests statement

The authors have declared that no conflict of interest exists.

Abstract

Impaired hepatic glucose and lipid metabolism are hallmarks of type−2 diabetes. Increased sulfide production from cysteine, or sulfide−donor compounds, may beneficially regulate hepatic metabolism. Disposal of sulfide through the sulfide oxidation pathway (SOP) is critical for maintaining sulfide within a safe physiological range. We show that mice lacking the liver−enriched mitochondrial SOP enzyme thiosulfate sulfur−transferase ($Tst^{-/-}$ mice) exhibit high circulating sulfide, increased gluconeogenesis, hypertriglyceridemia and fatty liver, despite whole−body insulin−sensitisation. Unexpectedly, hepatic sulfide levels were normal in $Tst^{-/-}$ mice, a result of homeostatic induction of mitochondrial sulfide disposal and glutathione excretion associated with net suppression of protein persulfidation and nuclear respiratory factor−2 target proteins. Proteomic and persulfidomic profiling converged on gluconeogenesis and hepatic lipid metabolism and revealed a selective deficit in medium−chain fatty acid oxidation in $Tst^{-/-}$ mice. We reveal a critical role for TST in hepatic metabolism that raises implications for sulfide-donor strategies in the context of liver function and metabolic disease.

Introduction

The prevalence of Type 2 diabetes (T2D) continues to soar in parallel with that of obesity[1]. Increased hepatic glucose production and aberrant hepatic lipid metabolism are cardinal features of T2D[2,3]. Dysregulation of hepatic nutrient metabolism in T2D is a promising area for therapeutic intervention because it precipitates the more severe liver pathologies that manifest along the spectrum of non-alcoholic fatty liver disease (NAFLD), steatosis, steatohepatitis and hepatocellular carcinoma[4].

Hydrogen sulfide (hereafter referred to as sulfide), an endogenously produced gaseous signalling molecule[5–8], has recently emerged as a modulator of nutrient metabolism[9–12]. Enzymatic sulfide production from sulfur amino acids is catalysed by cystathionine beta–synthase; CBS, and cystathionine gamma lyase; CTH[13,14] and by 3-mercaptopyruvate sulfur transferase; MPST[15–17]. Thioredoxin-mediated reduction of cysteine persulfides on proteins also regulates free sulfide and cysteine persulfide levels[18]. Endogenously produced and exogenously administered sulfide specifically influences hepatic glucose and lipid metabolism[19,20]. Thus, *in vitro,* treatment of murine hepatocytes with NaHS, or overexpression of rat *Cth* in HepG2 liver cells, increased glucose production through increased gluconeogenesis and reduced glycogen storage[21]. Conversely, glucose production was lower in hepatocytes from *Cth* gene knockout mice (*Cth*[−/−] mice) that exhibit low sulfide production[21]. Elevation of sulfide with NaHS administration *in vivo* reduced cholesterol and triglyceride accumulation in the liver of high fat diet (HFD)–fed mice[22]. In contrast, inter-cross of sulfide production–deficient *Cth*[−/−] mice with the hyperlipidemic *Apoe*[−/−] mouse strain (*Cth*[−/−]*Apoe*[−/−] mice) produced a phenotype of elevated plasma cholesterol following exposure to an atherogenic diet[23]. Consistent with their higher cholesterol, *Cth*[−/−]*Apoe*[−/−] mice developed fatty streak lesions earlier than *Apoe*[−/−] mice, and this effect was reversed

by NaHS administration[23]. Sulfide may also indirectly impact hepatic nutrient metabolism

through its effect on hepatic artery vasorelaxation and thus liver perfusion[24,25]. The

apparently beneficial effects of sulfide administration in multiple disease indications has led

to a major drive towards development of targeted H_2S–donor molecules as a therapeutic

approach[26,27]. However, an often-overlooked aspect of net sulfide exposure, key to the

efficacy of therapeutic H_2S–donors, is that it is regulated through its oxidative disposal. Thus,

endogenous sulfide exposure is actively limited to prevent mitochondrial respiratory

toxicity[28–30]. Sulfide is rapidly oxidized[31,32] through the mitochondrial sulfide oxidation

pathway (SOP), consisting of sulfide quinone oxidoreductase (SQOR), persulfide dioxygenase

(ETHE1/PDO) and thiosulfate sulfurtransferase (TST, also known as rhodanese)[31,33,34]. The

liver is highly abundant in SOP enzymes, and is a major organ of whole body sulfide disposal[32].

Mice lacking the *Ethe1* gene (*Ethe1*$^{-/-}$ mice) die of fatal sulfide toxicity[30], consistent with its

critical role in sulfide oxidation and the severe pathological consequences of unchecked

sulfide build-up in tissues. However, the importance of mitochondrial TST in the SOP *in vivo*

remains obscure. In contrast to *Ethe1*$^{-/-}$ mice, *Tst*$^{-/-}$ mice were grossly normal despite

exhibiting substantially elevated blood sulfide levels, as implied by qualitative measures[35].

This revealed an important yet distinct role for TST in the SOP for the first time *in vivo*.

Nevertheless, *Tst*$^{-/-}$ mice showed an apparently diabetogenic impairment of glucose

tolerance[35], consistent with the concept that increased sulfide promotes hepatic glucose

production[21]. As *Tst* deficiency represents a model of chronic but viable sulfide elevation,

determining the molecular mechanisms driving their aberrant metabolic profile can provide

important insights into the optimal range for therapeutic sulfide exposure, particularly in light

of the current interest in developing mitochondrially–targeted sulfide–donors[36,37]. To this end

we sought to define the impact of *Tst* deficiency on the underlying molecular pathways that

impact hepatic metabolism.

Results

Tst$^{-/-}$ mice exhibit increased hepatic gluconeogenesis and dyslipidaemia despite mild peripheral insulin sensitisation.

TST *mRNA* expression is highest in the liver (e.g. it is ~8-fold higher than in white adipose tissue http://biogps.org/#goto=genereport&id=22117). We therefore hypothesised that liver TST deficiency was the principal driver of the impaired glucose tolerance previously observed in *Tst$^{-/-}$* mice[35]. *Tst$^{-/-}$* mice exhibited higher glucose levels than C57BL/6J controls in response to a pyruvate challenge, consistent with higher hepatic glucose production (Figure 1A). We next tested phosphoenolpyruvate carboxykinase (PEPCK) activity, a key enzyme of *de-novo* hepatic glucose synthesis, and found it was higher in liver homogenates from *Tst$^{-/-}$* mice (Figure 1B). Next, we performed a 1 hour $^{13}C_3$-pyruvate metabolite–pulse incorporation experiment in isolated hepatocytes cultured in $^{12}C_3$-pyruvate–free medium. Hepatocytes from *Tst$^{-/-}$* mice displayed ^{13}C labelling consistent with increased metabolism of pyruvate to oxaloacetate – a critical early step in gluconeogenesis. Specifically, aspartate, which is derived from pyruvate via oxaloacetate was significantly increased in the *Tst$^{-/-}$* (Figure 1C). A trend towards higher $^{13}C_3$ malate, and lower $^{13}C_2$ acetyl–CoA was also observed (Figure S1A and S1B). $^{13}C_3$ Lactate was similar between genotypes, suggesting a similar activity of glycolytic disposal of pyruvate through lactate dehydrogenase (Figure S1A and S1B). Total pool sizes for all measured metabolites were similar between genotypes (Figure S1C). Consistent with increased endogenous glucose production in the *Tst$^{-/-}$* mice, fasting plasma glucose was higher in *Tst$^{-/-}$* mice relative to 6J mice during the pre-clamp 3-^{3}H glucose tracer infusion phase (60-90 minutes post tracer) of euglycemic, hyperinsulinemic (EH) clamp experiments (Figure 1D, Table S1A). Higher plasma glucose levels in *Tst$^{-/-}$* mice under these conditions

was not explained by lower glucose utilization in $Tst^{-/-}$ mice; glycogen synthesis and

glycolysis were comparable between genotypes across 60–90 minutes (Table S1A). Glucose

turnover – a derived parameter used to infer glucose production – was also comparable

between genotypes (Table S1A). However, derivation of glucose turnover requires that

glucose levels are stable during the period in which it is calculated. In our pre-clamp baseline

period, a highly significant effect of time (Figure 1D) indicated that this assumption was not

met, and thus true endogenous glucose production cannot be inferred from the glucose

turnover parameter in this instance. Combined with the pyruvate tolerance, PEPCK activity

and $^{13}C_3$-pyruvate pulse data, higher fasting glucose levels in the $Tst^{-/-}$ mice, given

comparable glucose utilization, is most likely due to higher endogenous glucose production

in the $Tst^{-/-}$ mice.

We next wished to explore whether the changes to glucose metabolism were driven by insulin

resistance. Liver glycogen, a marker of long–term carbohydrate storage typically impaired

with insulin resistance, was comparable between $Tst^{-/-}$ and C57BL/6J control mice (Figure

S2A). Despite unchanged steady–state markers of hepatic insulin sensitivity, impaired glucose

tolerance, previously described in the $Tst^{-/-}$ mice[35], suggested that whole body, and usually

hepatic, insulin–resistance was present. We investigated this using the euglycaemic clamp

where, unexpectedly, we observed whole–body insulin sensitisation under these short–term

steady–state conditions. During the clamp, when insulin was high, and blood glucose levels

were maintained constant, the glucose infusion rate was comparable between genotypes

(Table S1B). However, an increase in whole–body glucose uptake (integral glucose) by tissues

in the $Tst^{-/-}$ mice was apparent (Figure 1E, Table S1B), supporting increased peripheral

insulin–sensitivity, with a directionally consistent trend for increased glucose uptake into

several tissues. We confirmed this finding using standard insulin tolerance tests, where the

glucose decrement in response to insulin was greater in $Tst^{-/-}$ mice (Figure 1F, Figure S2B).

Together these data demonstrate a net increase in dynamic whole–body insulin sensitivity,

despite increased hepatic glucose output in $Tst^{-/-}$ mice. Finally, we assessed whole–body

glucose homeostasis with the EH–clamp method after chronic HFD feeding. Under these

conditions, $Tst^{-/-}$ mice maintained increased heptic glucose output (Figure 1D) but showed

convergence of the insulin–sensitivity profile to that of the C57BL/6J mice.

We also assessed whether Tst deficiency was associated with impaired lipid metabolism,

another hallmark of diabetes. Fast protein liquid chromatography analysis of triglyceride

levels and their lipoprotein distribution revealed significantly higher total plasma triglycerides

in $Tst^{-/-}$ mice (Figure 1G). The higher triglyceride was selectively associated with an

increased VLDL triglyceride fraction (Figure 1G), consistent with a dominant liver-driven

impairment in lipid metabolism[38]. Total and distinct lipoprotein fraction plasma cholesterol

levels were similar between genotypes (Figure S2C and S2D), suggestive of a triglyceride–

selective effect of Tst deficiency on hepatic lipid efflux. HFD significantly increased liver lipid

content of C57BL/6J mice but did not further increase the elevated lipid levels in the liver of

$Tst^{-/-}$ mice (Figure 1H-1I).

TST deficiency elicits compensatory hepatic sulfide disposal mechanisms that drive reduced

global protein persulfidation.

A role for TST in the disposal of sulfide was suggested by its participation in the SOP[31,34] and

supported *in vivo* by the qualitatively higher blood sulfide of $Tst^{-/-}$ mice[35]. Here, we

quantified circulating sulfide, showing approximately 10–fold elevation in the blood and

plasma of $Tst^{-/-}$ mice (Table 1A and 1B). Thiosulfate, an oxidised metabolite of sulfide [39,40]

and a TST substrate[41], was approximately 20–fold higher in plasma (Table 1B), and profoundly

higher (450–fold) in urine (Table 1C) of $Tst^{-/-}$ mice compared to C57Bl/6J mice. Reduced

glutathione (GSH) levels were ~2–fold higher in the plasma of $Tst^{-/-}$ mice (Table 1B). To

determine any direct hepatic contribution to the elevated systemic sulfide *in vivo*, whole

blood was sampled from the inferior vena cava (IVC) (Table 1D). IVC sulfide levels tended to

be higher in the $Tst^{-/-}$ mice, but the magnitude of the increase (~3-fold) did not parallel that

in trunk blood (~10-fold), suggesting liver was not a major source of the elevated circulating

sulfide. Surprisingly, liver homogenate sulfide, thiosulfate, cysteine and GSH levels were

similar between $Tst^{-/-}$ and C57BL/6J mice (Table 1E). Further, cultured hepatocytes from

$Tst^{-/-}$ and C57BL/6J mice exhibited similar intracellular sulfide levels, as estimated using P3,

a sulfide–selective fluorescent probe[42] (Table 1F). Mitochondrial sulfide levels in liver

reported by MitoA/MitoN[43] were similarly unchanged between genotypes (Table 1G). The

apparently unaltered hepatic steady-state sulfide levels, despite higher circulating sulfide,

suggested a profound homeostatic mechanism was invoked in the liver of $Tst^{-/-}$ mice. We

assessed respiratory sulfide disposal (antimycin–sensitive) and found this was markedly

increased in hepatocytes from $Tst^{-/-}$ mice, whereas antimycin–insensitive sulfide disposal

was relatively reduced compared to hepatocytes from C57BL/6J mice (Table S2). Isolated liver

mitochondria from $Tst^{-/-}$ hepatocytes also exhibited a higher sulfide disposal rate (Table S2).

In addition, cysteine and GSH was excreted at higher levels from $Tst^{-/-}$ hepatocytes under

basal conditions and after stimulation of sulfur amino acid metabolism by addition of

methionine (Figure 2A and 2B). Consistent with higher GSH turnover, hepatocytes from $Tst^{-/-}$

mice showed resistance to exogenous H_2O_2–mediated mitochondrial ROS production

(Figure S3). We next determined the global hepatic protein persulfidation profile, the major

post–translational modification mediated by sulfide[44–46]. Mass spectrometry analysis of

maleimide–labelled liver peptides revealed a greater abundance of peptides with a lower

persulfidation level (under-persulfidated) in the liver of $Tst^{-/-}$ mice (Figure 2C). We

confirmed this using semi-quantitative western–blot analysis on pulled down maleimide–

labelled proteins (Figure 2D). Gene Ontology (GO) analysis of proteins exhibiting altered

persulfidation levels revealed 28 terms different between $Tst^{-/-}$ and C57BL/6J mice by log–

fold change (Table 2). Major under-persulfidated terms (20) included "*FAD-binding, methyl*

transferase, peroxisome, acyl-CoA dehydrogenase activity and transaminase" while the less

abundant over-persulfidated terms (8) were dominated by terms related to "*Nicotinamide*

metabolism". Given evidence for altered gluconeogenesis above, we also performed a

pathway specifc analysis of the persulifidome. Persulfidation of proteins involved in

gluconeogenesis showed a mixture of under and over–persulfidated cysteines with a strong

enrichment of over-persulfidation , a pattern that was significantly different from the overall

persulfidation profile of the liver from $Tst^{-/-}$ mice (Figure S4A). We further found that the

gluconeogenesis pathway also exhibited a significantly higher magnitude of change

(independent of direction of change) in persulfidation than in the overall profile of the liver

of $Tst^{-/-}$ mice (Figure S4B).

The hepatic proteome of Tst$^{-/-}$ mice reveals a distinct molecular signature of altered sulfur

and mitochondrial nutrient metabolism.

To gain molecular insight into the mechanisms underlying the apparently diabetogenic liver

phenotype in ND-fed $Tst^{-/-}$ mice, we performed a comparative iTRAQ proteomic analysis of

whole liver with ND-fed C57BL/6J mice. Kyoto Encyclopedia of Genes and Genomes (KEGG)

analysis (at significance threshold P < 0.05) revealed 4 up–regulated pathways in liver of Tst^{-}

$^{/-}$ mice related to amino acid metabolism, including sulfur amino acids, and sulfur metabolism

(Table 3A). GO analysis (at threshold P < 0.05) revealed 95 significantly up–regulated terms in

liver of $Tst^{-/-}$ mice (Table S3A). Of the top 14 up–regulated terms, 7 referred to amino acid metabolism and 1 referred to the organellar term '*mitochondrion*'. KEGG pathway analysis revealed 27 down–regulated pathways in the liver of $Tst^{-/-}$ mice (Table 3B) including Phase 1 and 2 detoxification pathways (Cytochrome P450s, Glutathione and Glucuronidation) and '*Lysosome*', '*Protein processing in the Endoplasmic reticulum'* organellar terms. 213 GO terms were significantly down–regulated in $Tst^{-/-}$ mice (Table S4B). Among the most significant down–regulated terms were phase 2 detoxification pathways such as *'glutathione binding'* and *'glutathione transferase activity'* and *'endoplasmic reticulum'*. We validated a representative subset of protein changes (Figures S5A and S5D), noting broadly consistent direction of change in the key pathways. At the single protein level, the most robust change we observed was increased MPST protein in whole liver (Figure S5A and S5D) and mitochondrial sub-fractions (Figure S5B and S5D). This change was remarkable as mRNA levels for *Mpst* were lower in $Tst^{-/-}$ mice (Figure S5C), likely as a result of loss of proximal *Mpst* promoter function; *Mpst* is a paralog of *Tst*[47] juxtaposed approximately 1kb from the *Tst* gene. Protein levels for other sulfide producing and disposal enzymes were comparable between genotypes (Table S4). Given the evidence for glucose and lipid metabolic changes in the $Tst^{-/-}$ mice, we undertook a further focussed comparison of canonical proteins in these pathways (Table S5). Four GO terms were enriched in liver from $Tst^{-/-}$ mice, all of which were down–regulated; *'Lipid metabolic process'*, *'fatty acid beta-oxidation'*, *'Acyl-CoA dehydrogenase activity' and 'Acyl-CoA hydrolase activity'* (Table S5). Notably, supportive of largely unchanged hepatic insulin sensitivity, canonical insulin–regulated proteins[48, 49] were expressed comparably between genotypes (Tables S6 and S7).

*Hepatic protein expression in Tst$^{-/-}$ mice is consistent with lower NRF2 activation***.**

We performed a transcription factor binding site (TFBS) enrichment analysis in the promoters

of proteins that are up–regulated in the liver of $Tst^{-/-}$ mice to look for potential hub

transcriptional drivers of the proteome profile (Figure S6A). This revealed a statistically

significant under-representation of TFBS for the sulfide–responsive[50,51] NRF2 transcription

factor (Figure S6A). To explore whether there was evidence for reduced NRF2 activation in

the $Tst^{-/-}$ liver, we looked at the relative expression of 47 NRF2–regulated proteins in the

proteome of ND-fed $Tst^{-/-}$ and ND-fed C57Bl/6J mice. While 37 of these target proteins were

similar in expression between the genotypes, 10 NRF2 target proteins were lower in the liver

of $Tst^{-/-}$ mice (Figure S6B) and none of the 47 NRF2 target proteins were increased in $Tst^{-/-}$

liver.

Proteome of TST deficiency versus HFD response in C57BL/6J mice reveals distinct regulation

of lipid metabolism, sulfide metabolism and detoxification pathways.

We examined mechanistic commonalities between the diabetogenic hepatic phenotype of

$Tst^{-/-}$ mice and that induced by the diabetogenic HFD–feeding regimen in C57BL/6J mice.

Note, ND-fed $Tst^{-/-}$ mice were in a pre-existing diabetogenic state (Figure 1) that does not

further worsen with HFD (Figure 1H-I, Table S1), suggesting gross phenotypic convergence of

the two genotypes after HFD. We compared the identity and direction of change of the 188

proteins differentially expressed in ND-fed $Tst^{-/-}$ mice (versus ND-fed C57BL/6J mice; Figure

3A) to proteins that were differentially expressed in response to HFD in C57BL/6J mice (432

proteins; Figure 3A). There was a striking 67% overlap in individual proteins (126) in this

comparison (Figure 3A). When we analysed these two protein signatures for directionally

shared pathways, one up-regulated KEGG pathway *'Glycine, serine and threonine metabolism'*

(Table S8A) and 12 down-regulated KEGG pathways, including *'drug metabolism'* and

'endoplasmic reticulum' (Table S8B) were common to the liver of the ND-fed $Tst^{-/-}$ and HFD-

fed C57BL/6J mice. Consistent with a pre-existing HFD–like proteome, the dynamic response

to HFD in the liver of $Tst^{-/-}$ mice was muted, relative to that observed in C57BL6J mice (106

proteins, a 4–fold lower response; Figure 3B). Focussing on the sulfide pathway, MPST and

SUOX were increased by HFD in 6J and $Tst^{-/-}$ mice (Table S9). The HFD–induced increase in

MPST was less pronounced in the liver of $Tst^{-/-}$ mice, likely a reflection that it is already

elevated in ND–fed $Tst^{-/-}$ mice. We then considered contrasting, rather than congruent,

proteomic responses arising from TST deficiency versus HFD responses in C57BL/6J mice to

illuminate potential novel pathways underlying the otherwise functionally similar

diabetogenic hepatic $Tst^{-/-}$ phenotype. 5 KEGG pathways (Table S10A) and 4 GO terms (Table

S10B) were regulated in the opposite direction in this comparison. Strikingly, the GO terms

were all related to lipid metabolism, which were up–regulated in the HFD response but down–

regulated with TST deficiency (Table S10A-B). An organelle–focussed protein analysis showed

shared up-regulation of mitochondrial and endoplasmic reticulum pathways between TST

deficiency (Figure 3C upper row) and C57BL/6J HFD-responses (Figure 3C lower row), but a

striking discordance in peroxisomal protein pathways (up-regulated by HFD, down-regulated

TST deficiency) and nuclear proteins (down-regulated by HFD, up-regulated TST deficiency;

Figure 3C).

The $Tst^{-/-}$ liver proteome and persulfidome converge on transamination and lipid oxidation

pathways.

To assess whether conservation of changes at protein and post-translational modification

levels can illuminate key regulatory hubs driving the hepatic phenotype we ran a congruence

analysis of the proteome and persulfidome. We found that GO terms referring to *amino acid*,

lipid metabolism and *peroxisome* were significantly regulated at both the level of protein

abundance and persulfidation status (Table 4).

Tst$^{-/-}$ hepatocytes exhibit elevated mitochondrial respiration and a defect in medium–chain

fatty acid oxidation.

Enhanced respiratory sulfide disposal was found from *Tst$^{-/-}$* hepatocytes, and enrichment of

mitochondrial proteins was suggested from the liver proteome of the *Tst$^{-/-}$* mice (Table S4).

We therefore sought to determine if TST deficiency affected respiratory function and

substrate utilisation of the hepatocyte. Analysis of electron micrographs prepared from the

liver of ND-fed *Tst$^{-/-}$* mice and C57BL/6J controls showed morphologically normal

mitochondria (Figure 4A). Basal respiration, comprising ATP–linked and leak respiration, was

significantly higher in hepatocytes from *Tst$^{-/-}$* mice (Figure 4B–4D). Maximal hepatocyte

respiratory capacity and non–respiratory oxygen consumption was similar between

genotypes (Figures S7A and S7B). In line with phenotypic convergence following HFD,

hepatocyte respiration was comparable between genotypes from HFD-fed mice (Figures S7C-

S7H). A unique feature of the liver from *Tst$^{-/-}$* mice that contrasted with HFD response in

C57BL/6J mice was a decrease in proteins and persulfidation levels of proteins in lipid

oxidation pathways. We therefore investigated hepatocyte respiration of lipids. Using a low

pyruvate (100 μM) medium to reveal respiratory dependency on other substrates, we showed

that CPT1A–mediated mitochondrial oxidation of endogenous long chain fatty acids (LCFA;

etomoxir-inhibited) was similar between genotypes (Figure 4E). Next, we by–passed CPT1A–

mediated LCFA transfer, and revealed a marked deficit in respiration stimulated by the

medium chain fatty acid octanoate in hepatocytes from *Tst$^{-/-}$* mice (Figure 4F). A similar

experiment adding back pyruvate revealed comparable stimulation of respiration between

genotypes (Figure S6I). In amino acid free media, combined glutamine–, aspartate– and

alanine–stimulated hepatocyte respiration was comparable between genotypes (Figure S6J).

Discussion

Elevated TST expression in adipose tissue was recently identified as a genetic mechanism

driving metabolically protective leanness in mice[35]. Conversely, mice with genetic deletion of

Tst (*Tst*$^{-/-}$ mice) exhibited impaired glucose tolerance relative to C57BL/6N controls[35]. A key

observation from the *Tst*$^{-/-}$ mouse was qualitative evidence for elevated blood sulfide

levels[35], which implicated the mitochondrial sulfurtransferase as playing an important role in

the sulfide oxidation pathway (Figure S8) for the first time *in vivo*. However, the mechanistic

link between *Tst* deficiency and impaired glucose tolerance remained unclear. The *Tst*$^{-/-}$

mice had a subtle adipose tissue phenotype, with comparable fat accumulation upon high fat

feeding, suggesting alternate organs as a potential source of impaired glucose homeostasis.

Moreover, TST is most highly abundant in the liver, consistent with its role in detoxification

of cyanide and related molecules. Indeed, hepatic TST mRNA levels are ~8-fold higher than

adipose tissue, with the difference in enzyme activity likely substantially exceeding this, per

gram of tissue. Based upon the reported effects of sulfide to increase hepatic glucose

production[20,21] and modulate lipid metabolism[22,23], we hypothesised that TST deficiency

predominantly drives a diabetogenic state through increased hepatic sulfide exposure.

Consistent with this, we found evidence supportive of increased gluconeogenesis, steatosis,

elevated plasma VLDL triglycerides and, quantitatively, a 10-fold higher circulating blood

sulfide level in *Tst*$^{-/-}$ mice. Unexpectedly, and in spite of the markedly increased circulating

sulfide levels, steady-state sulfide level was normal in the liver of *Tst*$^{-/-}$ mice. Moreover, we

found evidence for multiple mechanisms for increased, not decreased, hepatic sulfide

disposal, reduced downstream sulfide signalling and novel molecular explanations for the

apparently diabetogenic phenotype. Our data are consistent with the liver being a major site

for sulfide disposal rather than production[32]. Moreover, our data suggest that the liver of

$Tst^{-/-}$ mice has over-compensated in its response to maximise hepatic sulfide removal, thus

engendering detrimental metabolic consequences indirectly. This response is achieved

through a combination of distinct compartmentalised cellular responses including increased

respiratory sulfide disposal and export of cysteine and GSH and is further evidenced by the

reduction of global hepatic protein persulfidation and depletion of sulfide-responsive[51] NRF2

target proteins. Additionally, up-regulation of translation and recruitment of MPST to the

mitochondria of $Tst^{-/-}$ mice is observed. This response, in face of reduced transcription of

Mpst due to *Mpst* promoter disruption by the *Tst* null allele, suggests a powerful post-

transcriptional cellular sulfide sensing mechanism. Moreover, if MPST is compensating for

TST–mediated sulfide disposal in this context, this suggests a subversion of normal MPST

function away from sulfide production[52,53] [54,55]. Intriguingly, TST expression levels were

elevated in the liver of $Mpst^{-/-}$ mice, providing further support for an exquisite reciprocal

compensatory mechanism between these two enzymes[56].

The unexpected finding of normal hepatic sulfide levels in the $Tst^{-/-}$ mice led us to discover

that the metabolic phenotype we observed was driven by the very mechanisms invoked to

maintain sulfide within a normal range rather than sulfide excess *per se*. Several observations

were consistent with this. For example, the major amino acid pathways increased in the liver

proteome of $Tst^{-/-}$ mice were transaminases involved in metabolism of GSH that support

increased export of sulfur equivalents as GSH (and cysteine). These same transaminases

support gluconeogenesis by redirecting Krebs cycle intermediates[57–59]. Re–programming of

amino acid metabolism for sulfide disposal, with knock–on effects to drive hepatic glucose

production are suggested, rather than any change to amino acid–linked mitochondrial

respiration in hepatocytes. This is supported by the shift in pyruvate metabolism towards

aspartate. In addition, glutathione-S-transferases (GST) that inhibit gluconeogenesis[60] were

lower in the liver of $Tst^{-/-}$ mice. Further, NRF2 activation, which represses gluconeogenesis[61]

appears lower in liver of $Tst^{-/-}$ mice. The involvement of NRF2 in the $Tst^{-/-}$ liver phenotype

is further supported by the phenotype of $Nrf2^{-/-}$ mice that also exhibited steatohepatitis, in

the absence of insulin resistance[62]. However, we note that NRF2 signalling can be complex

and dependent upon dietary context; $Nrf2^{-/-}$ mice showed improved glucose tolerance after

a high–fat diet[63] suggesting any contribution of a NRF2 signalling deficit in the liver of the $Tst^{-/-}$

mice changes upon high fat feeding. Beyond altered pyruvate flux, we also showed that the

hepatocytes of $Tst^{-/-}$ mice exhibited defective lipid metabolism. Specifically, medium–chain

fatty acid (MCFA) oxidation was impaired, associated with selective reduction of protein levels

and persulfidation levels of lipid catabolic enzymes. This represents a novel mechanism

linking altered sulfide metabolism to lipid oxidation, hepatic lipid accumulation and

dyslipidaemia. Consistent with impaired MCFA oxidation defects as a driver, at least in part,

of the phenotype of $Tst^{-/-}$ mice, steatosis is observed in medium–chain acyl-CoA

dehydrogenase $(Mcad)^{-/-}$ mice[64] and dyslipidemia is found in MCADD deficient humans[65].

The data we present adds to a growing understanding of the link between sulfide regulating

genes and nutrient metabolism that has hitherto focussed on the enzymes of sulfide

production. Specifically, we provide support for the importance of the sulfide oxidising

pathway as a regulator of cellular sulfide exposure. Unexpectedly, the data reveal novel

cellular mechanisms that are engaged to homeostatically regulate sulfide disposal and that

can impact upon cell energetics and nutrient metabolism.

Our findings may have implications for potentially unexpected side effects of sulfide donor

therapeutics. In normal mice, *in vivo* sulfide administration for 4 weeks post–HFD partially

reversed hepatic lipid accumulation invoked by chronic (16 weeks) HFD[22]. No evidence was

provided for whether sulfide disposal mechanisms were altered[22]. This efficacious sub-

chronic sulfide administration regimen contrasts with our genetic model of chronic sulfide

elevation as a driver of dysregulated metabolism and NAFLD. Clearly, the normal mice in the

Na_2S administration studies had a fully functional SOP, suggesting the presence of TST is

required to achieve the beneficial metabolic effects of Na_2S administration. This is also

consistent with the apparently low sulfide signalling status (evidenced by lower

persulfidation, NRF2 target protein abundance) in the liver of the $Tst^{-/-}$ mice. The benefits

of elevated sulfide cannot be realised, perhaps because the major mediator of those effects

is missing and the alternate mechanisms invoked do not fully compensate (e.g. MPST), or

actively drive aberrant nutrient metabolism. Comparable studies of glucose and lipid

metabolism after manipulation of other sulfide regulating genes are limited. However in a

contrasting model of reduced sulfide production ($Cth^{-/-}$ mice), plasma triglycerides were

lowered[23], consistent and opposite to what we observed with the $Tst^{-/-}$ mice. The hepatic

sulfide disposal status of the $Cth^{-/-}$ mouse model is unknown, but our findings predict a

suppression of the SOP to spare the limited endogenous sulfide produced. Intriguingly they

also predict a knock-on effect on nutrient homeostasis due to reduced metabolic demand of

the TST/SOP axis. A more direct model informing on the effects of impairment of the sulfide

disposal pathway is deficiency of the key mitochondrial SOP enzyme ETHE1. $Ethe1^{-/-}$ mice

suffer fatal sulfide toxicity[30] and therefore comparable metabolic studies are lacking.

However, one notable observation is that $Ethe1^{-/-}$ mice have an apparently 10-fold higher

liver sulfide exposure than control mice[30], in contrast to the normalised hepatic sulfide levels

of $Tst^{-/-}$ mice. Circulating sulfide levels were not reported for comparison, but the

presumably relatively lower systemic sulfide levels of $Tst^{-/-}$ mice appear to have permitted

an effective homeostatic sulfide disposal response in the liver to avoid toxicity, albeit with a

metabolic cost. Consequently, the liver of $Tst^{-/-}$ mice has a distinct functional and proteomic

profile to that of the $Ethe1^{-/-}$ mice. For example, in the liver of $Tst^{-/-}$ and $Ethe1^{-/-}$ mice[66],

proteins of the glutathione–S–transferase Mu type (GSTM) and peroxiredoxin (PRDX) families

were altered, but sometimes in the opposite direction or with alteration of distinct protein

sub-classes. A notable difference is also observed in amino acid metabolism. The liver of

$Ethe1^{-/-}$ mice exhibited increased expression of enzymes of branched chain amino acid

metabolism[66], distinct from the predominantly glutathione–related amino acid pathways that

are increased in liver of $Tst^{-/-}$ mice. Beyond sulfide, TST may also have distinct cellular roles

that affect metabolism such as mitoribosomal synthesis, ROS attenuation and modulation of

mitochondrial iron-sulfur clusters[67–71].

Given the pro-diabetogenic liver phenotype in $Tst^{-/-}$ mice, its was surprising that insulin

signalling in the liver appeared normal, and peripheral insulin sensitivity was, if anything,

increased. There are precedents for increased hepatic glucose production independent of

insulin resistance, as found in the $Nrf2^{-/-}$ mice[62] and as driven by the transcription factor

ChREBP[72,73]. There is also evidence to support insulin–sensitising effects of sulfide

administration *in vivo* in mice and rats[74–76], consistent with a sulfide–mediated insulin–

sensitisation of non-hepatic tissues in $Tst^{-/-}$ mice. Higher circulating GSH in $Tst^{-/-}$ mice may

also promote peripheral insulin–sensitisation[77,78]. Clearly, the net balance of glucose

production from the liver and its peripheral disposal remains abnormal in $Tst^{-/-}$ mice.

Indeed, the baseline metabolic phenotype of $Tst^{-/-}$ mice resembles in many ways that of a

normal mouse fed a HFD. Thus, we expected to gain insight into conserved pro-diabetogenic

mechanisms from the overlap between the liver proteome of $Tst^{-/-}$ mice and that of HFD-

fed C57BL/6J mice. However, upon closer inspection this was not entirely the case, with some

lipid metabolism and peroxisomal proteins changing in the opposite direction to those

observed in HFD–fed C57BL/6J mice. Unlike a HFD state, which is associated with dominant hepatic insulin resistance, the increased hepatic glucose production in ND-fed $Tst^{-/-}$ mice occurs despite of normal hepatic insulin sensitivty. The significant changes in persulfidation of transaminase and gluconeogenesis proteins in the liver of $Tst^{-/-}$ mice suggest a coordinated cross–talk across metabolic pathways underlies this atypical metabolic phenotype.

Sulfide donor therapeutics were proposed as a clinical strategy for improving cardiovascular health[27,79,80]. Elevated endogenous sulfide was also implicated in the beneficial metabolic effects of caloric restriction[11,81–85]. Our results suggest that chronic sulfide elevation may have unintended detrimental consequences, driving liver glucose production and fat accumulation to undesirable levels. This caveat may be fortunately limited to cases where SOP proteins are compromised through rare genetic effects – such as TST variants[86,87]. More broadly, a number of drugs or supplements are known to increase cyanide, which may dominantly inhibit TST activity and result in secondary sulfide overexposure. These include nitroprusside[88] and amygdalin[89,90]. Indeed the TST metabolite thiosulfate is commonly co-administered with nitroprusside to prevent cyanide toxicity[91]. Furthermore, dietary and environmental exposure to cyanogenic compounds[92], e.g. smoking[93] or cyanogenic diets[94] may interfere with normal TST function and could lead to increased sensitivity to sulfide therapeutics. In contrast, we have shown that administration of the TST substrate thiosulfate can ameliorate diabetes[35] further underlining the potential utility of targeting the SOP in metabolic disease. As with all therapeutic strategies, a careful cost-benefit analysis is required. A comparable case of relevance are the statins, one of the most potent and widely used drugs to prevent atherosclerosis, which also carry a higher risk for diabetes[95]. The full impact of TST manipulation on opposing metabolic pathways requires further study. Our current study

472   sheds light on the underlying hepatic mechanisms invoked for sulfide disposal that are

473   relevant to current sulfide–donor strategies and may inform on routes to reduce their

474   potential metabolic side-effects.

Methods

477   The methods for all data and work presented in this manuscript is provided as supplementary

478   information.

Author Contributions

480   N.M.M. and R.N.C conceived experiments. R.N.C., M.T.G.G, M. B.-L., A. M.-C., M.L., V.V., B.E.,

481   T. LeB., M.B., S.H., S.D., N.H., C. Mc F., A.T., N. F., T.G., performed experiments. R.N.C., M. B.-

482   L., P.F., V.V., T. LeB., N.H., T.S., F.B., T.G., R.C.H., B.S., G.G., A.J.F., C.S., R.B. and N.M.M.

483   analysed and interpreted data and commented on the manuscript. K.H.A., S.S. generated

484   reagents. C.McM. and R.C.H. generated reagents. R.N.C. and N.M.M. wrote the manuscript.

Acknowledgements

486   This work was funded by a Wellcome Trust New Investigator Award to NMM (100981/Z/13/Z)

487   and Diabetes UK grant (17/0005697). RCH was funded by a Medical Research Council

488   Discovery Award (MC-PC-15076). We would like to thank Professor Ken Olsen and Eric DeLeon

489   (Notre Dame) for advice establishing sulfide measurement by probe.

References

References

1.   OMS. GLOBAL REPORT ON DIABETES WHO Library Cataloguing-in-Publication Data.

ISBN. 2014;978:92-94. http://www.who.int/about/licensing/.

2.   Lewis GF, Carpentier A, Adeli K, Giacca A. Disordered fat storage and mobilization in

the pathogenesis of insulin resistance and type 2 diabetes. *Endocr Rev*.

2002;23(2):201-229. doi:10.1210/edrv.23.2.0461

3.   Consoli A, Nurjhan N, Capani F, Gerich J. Predominant role of gluconeogenesis in

increased hepatic glucose production in NIDDM. *Diabetes*. 1989;38(5):550-557.

doi:10.2337/diab.38.5.550

4.   Caron S, Verrijken A, Mertens I, et al. Transcriptional activation of apolipoprotein CIII

expression by glucose may contribute to diabetic dyslipidemia. *Arterioscler Thromb*

Vasc Biol. 2011;31(3):513-519. doi:10.1161/ATVBAHA.110.220723

5.   Abe K, Kimura H. The possible role of hydrogen sulfide as an endogenous

neuromodulator. *J Neurosci*. 1996;16(3):1066-1071.

http://www.ncbi.nlm.nih.gov/pubmed/8558235.

6.   Wang R. Physiological Implications of Hydrogen Sulfide: A Whiff Exploration That

Blossomed. *Physiol Rev*. 2012;92(2):791-896. doi:10.1152/physrev.00017.2011

7.   Mishanina T V., Libiad M, Banerjee R. Biogenesis of reactive sulfur species for

signaling by hydrogen sulfide oxidation pathways. *Nat Chem Biol*. 2015;11(7):457-

464. doi:10.1038/nchembio.1834

8.   Filipovic MR, Zivanovic J, Alvarez B, Banerjee R. Chemical Biology of H2S Signaling

through Persulfidation. *Chem Rev*. 2017. doi:10.1021/acs.chemrev.7b00205

9.   Desai KM, Chang T, Untereiner A, Wu L. Hydrogen sulfide and the metabolic

syndrome. *Expert Rev Clin Pharmacol*. 2011;4(1):63-73. doi:10.1586/ecp.10.133

10. Szabo C. Roles of Hydrogen Sulfide in the Pathogenesis of Diabetes Mellitus and Its

Complications. *Antioxid Redox Signal*. 2011;17(1):68-80. doi:10.1089/ars.2011.4451

11. Hine C, Harputlugil E, Zhang Y, et al. Endogenous hydrogen sulfide production is

essential for dietary restriction benefits. *Cell*. 2015. doi:10.1016/j.cell.2014.11.048

12. Carter RN, Morton NM. Cysteine and hydrogen sulphide in the regulation of

metabolism: Insights from genetics and pharmacology. *J Pathol*. 2016;238(2):321-332.

doi:10.1002/path.4659

13. Chen X, Jhee KH, Kruger WD. Production of the neuromodulator H2S by

cystathionine ??-synthase via the condensation of cysteine and homocysteine. *J Biol*

Chem. 2004. doi:10.1074/jbc.C400481200

14. Singh S, Padovani D, Leslie RA, Chiku T, Banerjee R. Relative contributions of

cystathionine ??-synthase and ??-cystathionase to H2S biogenesis via alternative

trans-sulfuration reactions. *J Biol Chem*. 2009. doi:10.1074/jbc.M109.010868

15. Shibuya N, Tanaka M, Yoshida M, et al. 3-Mercaptopyruvate Sulfurtransferase

Produces Hydrogen Sulfide and Bound Sulfane Sulfur in the Brain. *Antioxid Redox*

Signal. 2009. doi:10.1089/ars.2008.2253

16. Mikami Y, Shibuya N, Kimura Y, Nagahara N, Ogasawara Y, Kimura H. Thioredoxin and

dihydrolipoic acid are required for 3-mercaptopyruvate sulfurtransferase to produce

hydrogen sulfide. *Biochem J*. 2011. doi:10.1042/BJ20110841

17. Yadav PK, Yamada K, Chiku T, Koutmos M, Banerjee R. Structure and kinetic analysis

of H2S production by human mercaptopyruvate sulfurtransferase. *J Biol Chem*. 2013.

doi:10.1074/jbc.M113.466177

18. Wedmann R, Onderka C, Wei S, et al. Improved tag-switch method reveals that

thioredoxin acts as depersulfidase and controls the intracellular levels of protein

persulfidation. *Chem Sci*. 2016;7(5):3414-3426. doi:10.1039/c5sc04818d

19. Mani S, Cao W, Wu L, Wang R. Hydrogen sulfide and the liver. *Nitric Oxide - Biol*

Chem. 2014. doi:10.1016/j.niox.2014.02.006

20. Pichette J, Gagnon J. Implications of Hydrogen Sulfide in Glucose Regulation: How

H2S Can Alter Glucose Homeostasis through Metabolic Hormones. *Oxid Med Cell*

Longev. 2016;2016. doi:10.1155/2016/3285074

21. Zhang L, Yang G, Untereiner A, Ju Y, Wu L, Wang R. Hydrogen sulfide impairs glucose

utilization and increases gluconeogenesis in hepatocytes. *Endocrinology*. 2013.

doi:10.1210/en.2012-1658

22. Wu D, Zheng N, Qi K, et al. Exogenous hydrogen sulfide mitigates the fatty liver in

obese mice through improving lipid metabolism and antioxidant potential. *Med Gas*

Res. 2015;5(1). doi:10.1186/s13618-014-0022-y

23. Mani S, Li H, Untereiner A, et al. Decreased endogenous production of hydrogen

sulfide accelerates atherosclerosis. *Circulation*. 2013;127(25):2523-2534.

doi:10.1161/CIRCULATIONAHA.113.002208

24. Distrutti E, Mencarelli A, Santucci L, et al. The methionine connection: Homocysteine

and hydrogen sulfide exert opposite effects on hepatic microcirculation in rats.

Hepatology. 2008;47(2):659-667. doi:10.1002/hep.22037

25. Fiorucci S, Antonelli E, Mencarelli A, et al. The third gas: H2S regulates perfusion

pressure in both the isolated and perfused normal rat liver and in cirrhosis.

Hepatology. 2005;42(3):539-548. doi:10.1002/hep.20817

26. Sestito S, Nesi G, Pi R, Macchia M, Rapposelli S. Hydrogen Sulfide: A Worthwhile Tool

in the Design of New Multitarget Drugs. *Front Chem*. 2017;5.

doi:10.3389/fchem.2017.00072

27.   Whiteman M, Le Trionnaire S, Chopra M, Fox B, Whatmore J. Emerging role of

hydrogen sulfide in health and disease: critical appraisal of biomarkers and

pharmacological tools. *Clin Sci*. 2011;121(11):459-488. doi:10.1042/cs20110267

28.   Libiad M, Motl N, Akey DL, et al. Thiosulfate Sulfurtransferase like Domain Containing

1 Protein Interacts with Thioredoxin. *J Biol Chem*. January 2018:jbc.RA117.000826.

doi:10.1074/jbc.RA117.000826

29.   Reiffenstein R. Toxicology Of Hydrogen Sulfide. *Annu Rev Pharmacol Toxicol*.

1992;32(1):109-134. doi:10.1146/annurev.pharmtox.32.1.109

30.   Tiranti V, Viscomi C, Hildebrandt T, et al. Loss of ETHE1, a mitochondrial dioxygenase,

causes fatal sulfide toxicity in ethylmalonic encephalopathy. *Nat Med*.

2009;15(2):200-205. doi:10.1038/nm.1907

31.   Hildebrandt TM, Grieshaber MK. Three enzymatic activities catalyze the oxidation of

sulfide to thiosulfate in mammalian and invertebrate mitochondria. *FEBS J*.

2008;275(13):3352-3361. doi:10.1111/j.1742-4658.2008.06482.x

32.   Norris EJ, Culberson CR, Narasimhan S, Clemens MG. The liver as a central regulator

of hydrogen sulfide. *Shock*. 2011. doi:10.1097/SHK.0b013e3182252ee7

33.   Jackson MR, Melideo SL, Jorns MS. Human sulfide:Quinone oxidoreductase catalyzes

the first step in hydrogen sulfide metabolism and produces a sulfane sulfur

metabolite. *Biochemistry*. 2012. doi:10.1021/bi300778t

34.   Libiad M, Yadav PK, Vitvitsky V, Martinov M, Banerjee R. Organization of the human

mitochondrial hydrogen sulfide oxidation pathway. *J Biol Chem*. 2014.

doi:10.1074/jbc.M114.602664

35.   Morton NM, Beltram J, Carter RN, et al. Genetic identification of thiosulfate

sulfurtransferase as an adipocyte-expressed antidiabetic target in mice selected for

leanness. *Nat Med*. 2016;22(7). doi:10.1038/nm.4115

36.   Gero, Domokos Torregrossa R, Perry A, Waters A, et al. The novel mitochondria-

targeted hydrogen sulfide (H2S) donors AP123 and AP39 protect against

hyperglycemic injury in microvascular endothelial cells in vitro. *Pharmacol Res*.

2016;113:186-198.

https://reader.elsevier.com/reader/sd/pii/S1043661816306314?token=3623BE15023

42420421E3956873F1E65FD3FCBDE1D675D7396DB26A…1/13.

37.   Karwi QG, Bice JS, Baxter GF. Pre- and postconditioning the heart with hydrogen

sulfide (H2S) against ischemia/reperfusion injury in vivo: a systematic review and

meta-analysis. *Basic Res Cardiol*. 2018;113(1). doi:10.1007/s00395-017-0664-8

38.   Mason TM. The role of factors that regulate the synthesis and secretion of very-low-

density lipoprotein by hepatocytes. *Crit Rev Clin Lab Sci*. 1998;35(6):461-487.

doi:10.1080/10408369891234246

39.   Vitvitsky V, Yadav PK, Kurthen A, Banerjee R. Sulfide oxidation by a noncanonical

pathway in red blood cells generates thiosulfate and polysulfides. *J Biol Chem*.

2015;290(13):8310-8320. doi:10.1074/jbc.M115.639831

40.   Vitvitsky V, Yadav PK, An S, Seravalli J, Cho US, Banerjee R. Structural and mechanistic

insights into hemoglobincatalyzed hydrogen sulfide oxidation and the fate of

polysulfide products. *J Biol Chem*. 2017. doi:10.1074/jbc.M117.774943

41.   Banerjee R, Chiku T, Kabil O, Libiad M, Motl N, Yadav PK. *Assay Methods for H2S*

Biogenesis and Catabolism Enzymes. Vol 554. Academic Press Inc.; 2015:189-200.

doi:10.1016/bs.mie.2014.11.016

42.   Singha S, Kim D, Moon H, et al. Toward a selective, sensitive, fast-responsive, and

611   biocompatible two-photon probe for hydrogen sulfide in live cells. *Anal Chem*.

612   2015;87(2):1188-1195. doi:10.1021/ac503806w

613 43. Arndt S, Baeza-Garza CD, Logan A, et al. Assessment of H 2 S in vivo using the newly

614   developed mitochondria-targeted mass spectrometry probe MitoA. *J Biol Chem*.

615   2017;292(19):7761-7773. doi:10.1074/jbc.M117.784678

616 44. Krishnan N, Fu C, Pappin DJ, Tonks NK. H2S-Induced Sulfhydration of the Phosphatase

617   PTP1B and Its Role in the Endoplasmic Reticulum Stress Response. *Sci Signal*. 2011.

618   doi:10.1126/scisignal.2002329

619 45. Kabil O, Motl N, Banerjee R. H2S and its role in redox signaling. *Biochim Biophys Acta -*

620   *Proteins Proteomics*. 2014;1844(8):1355-1366. doi:10.1016/j.bbapap.2014.01.002

621 46. Koike S, Nishimoto S, Ogasawara Y. Cysteine persulfides and polysulfides produced by

622   exchange reactions with H2S protect SH-SY5Y cells from methylglyoxal-induced

623   toxicity through Nrf2 activation. *Redox Biol*. 2017;12:530-539.

624   doi:10.1016/j.redox.2017.03.020

625 47. Nagahara N. Catalytic Site Cysteines of Thiol Enzyme: Sulfurtransferases. *J Amino*

626   *Acids*. 2011. doi:10.4061/2011/709404

627 48. Batista TM, Garcia-Martin R, Cai W, et al. Multi-dimensional Transcriptional

628   Remodeling by Physiological Insulin In Vivo. *Cell Rep*. 2019;26(12):3429-3443.e3.

629   doi:10.1016/j.celrep.2019.02.081

630 49. Sutherland C, O'Brien R, Granner D. *Insulin Action Gene Regulation. In: Madame*

631   *Curies Bioscience Database [Internet].* Austin(TX); 2013.

632   https://www.ncbi.nlm.gov/books/NBK6471.

633 50. Xie L, Gu Y, Wen M, et al. Hydrogen sulfide induces Keap1 S-sulfhydration and

634   suppresses diabetes-accelerated atherosclerosis via Nrf2 activation. *Diabetes*.

635   2016;65(10):3171-3184. doi:10.2337/db16-0020

636 51. Yang G, Zhao K, Ju Y, et al. Hydrogen sulfide protects against cellular senescence via s-

637   sulfhydration of keap1 and activation of Nrf2. *Antioxidants Redox Signal*.

638   2013;18(15):1906-1919. doi:10.1089/ars.2012.4645

639 52. Módis K, Coletta C, Erdélyi K, Papapetropoulos A, Szabo C. Intramitochondrial

640   hydrogen sulfide production by 3-mercaptopyruvate sulfurtransferase maintains

641   mitochondrial electron flow and supports cellular bioenergetics. *FASEB J*. 2013.

642   doi:10.1096/fj.12-216507

643 53. Szabo C, Ransy C, Módis K, et al. Regulation of mitochondrial bioenergetic function by

644   hydrogen sulfide. Part I. Biochemical and physiological mechanisms. *Br J Pharmacol*.

645   2014. doi:10.1111/bph.12369

646 54. Nagahara N. Multiple role of 3-mercaptopyruvate sulfurtransferase: antioxidative

647   function, H2S and polysulfide production and possible SOx production. *Br J*

648   *Pharmacol*. 2018;175(4):577-589. doi:10.1111/bph.14100

649 55. Kimura Y, Koike S, Shibuya N, Lefer D, Ogasawara Y, Kimura H. 3-Mercaptopyruvate

650   sulfurtransferase produces potential redox regulators cysteine- and glutathione-

651   persulfide (Cys-SSH and GSSH) together with signaling molecules H_2S_2, H_2S_3 and

652   H_2S. *Sci Rep*. 2017;7(1):1-14. doi:10.1038/s41598-017-11004-7

653 56. Nagahara N, Tanaka M, Tanaka Y, Ito T. Novel Characterization of Antioxidant

654   Enzyme, 3-Mercaptopyruvate Sulfurtransferase-Knockout Mice: Overexpression of

655   the Evolutionarily-Related Enzyme Rhodanese. *Antioxidants*. 2019;8(5):116.

656   doi:10.3390/antiox8050116

657 57. Rui L. Energy metabolism in the liver. *Compr Physiol*. 2014;4(1):177-197.

658   doi:10.1002/cphy.c130024

58. Qian K, Zhong S, Xie K, Yu D, Yang R, Gong DW. Hepatic ALT isoenzymes are elevated

in gluconeogenic conditions including diabetes and suppressed by insulin at the

protein level. *Diabetes Metab Res Rev*. 2015;31(6):562-571. doi:10.1002/dmrr.2655

59. Sookoian S, Castaño GO, Scian R, et al. Serum aminotransferases in nonalcoholic fatty

liver disease are a signature of liver metabolic perturbations at the amino acid and

Krebs cycle level. *Am J Clin Nutr*. 2016;103(2):422-434. doi:10.3945/ajcn.115.118695

60. Ghosh Dastidar S, Jagatheesan G, Haberzettl P, et al. Glutathione S -transferase P

deficiency induces glucose intolerance via JNK-dependent enhancement of hepatic

gluconeogenesis . *Am J Physiol Metab*. 2018;315(5):E1005-E1018.

doi:10.1152/ajpendo.00345.2017

61. Slocum SL, Skoko JJ, Wakabayashi N, et al. Keap1/Nrf2 pathway activation leads to a

repressed hepatic gluconeogenic and lipogenic program in mice on a high-fat diet.

Arch Biochem Biophys. 2016;591:57-65. doi:10.1016/j.abb.2015.11.040

62. Meakin PJ, Chowdhry S, Sharma RS, et al. Susceptibility of Nrf2-Null Mice to

Steatohepatitis and Cirrhosis upon Consumption of a High-Fat Diet Is Associated with

Oxidative Stress, Perturbation of the Unfolded Protein Response, and Disturbance in

the Expression of Metabolic Enzymes but Not with Insulin Resistance. *Mol Cell Biol*.

2014;34(17):3305-3320. doi:10.1128/mcb.00677-14

63. Zhang YKJ, Wu KC, Liu J, Klaassen CD. Nrf2 deficiency improves glucose tolerance in

mice fed a high-fat diet. *Toxicol Appl Pharmacol*. 2012;264(3):305-314.

doi:10.1016/j.taap.2012.09.014

64. Tolwani RJ, Hamm DA, Tian L, et al. Medium-chain acyl-CoA dehydrogenase deficiency

in gene-targeted mice. *PLoS Genet*. 2005;1(2):0205-0212.

doi:10.1371/journal.pgen.0010023

65.   Onkenhout W, Venizelos V, Van Der Poet PFH, Van Den Heuvel MPM, Poorthuis

BJHM. *Identification and Quantification of Intermediates of Unsaturated Fatty Acid*

Metabolism in Plasma of Patients with Fatty Acid Oxidation Disorders. Vol 10.; 1995.

66.   Hildebrandt TM, Di Meo I, Zeviani M, Viscomi C, Braun H. Proteome adaptations in

Ethe1-deficient mice indicate a role in lipid catabolism and cytoskeleton organization

via post-translational protein modifications. *Biosci Rep*. 2013;33(4):575-584.

doi:10.1042/bsr20130051

67.   Bonomi F, Pagani S, Cerletti P, Cannella C. *Rhodanese-Mediated Sulfur Transfer to*

Succinate Dehydrogenase. Vol 72.; 1977.

68.   Pagani S, Galante YM. *INTERACTION OF RHODANESE WITH MITOCHONDRIAL NADH*

DEHYDROGENASE. Vol 742.; 1983.

69.   Nandi DL, Westley J. Reduced thioredoxin as a sulfur-acceptor substrate for

rhodanese. *Int J Biochem Cell Biol*. 1998. doi:10.1016/S1357-2725(98)00050-8

70.   Nandi DL, Horowitz PM, Westley J. Rhodanese as a thioredoxin oxidase. *Int J Biochem*

Cell Biol. 2000;32(4):465-473. doi:10.1016/S1357-2725(99)00035-7

71.   Smirnov A, Comte C, Mager-Heckel AM, et al. Mitochondrial enzyme rhodanese is

essential for 5 S ribosomal RNA import into human mitochondria. *J Biol Chem*.

2010;285(40):30792-30803. doi:10.1074/jbc.M110.151183

72.   Uyeda K, Repa JJ. Carbohydrate response element binding protein, ChREBP, a

transcription factor coupling hepatic glucose utilization and lipid synthesis. *Cell*

Metab. 2006;4(2):107-110. doi:10.1016/j.cmet.2006.06.008

73.   Kim MS, Krawczyk SA, Doridot L, et al. ChREBP regulates fructose-induced glucose

production independently of insulin signaling. *J Clin Invest*. 2016;126(11):4372-4386.

doi:10.1172/JCI81993

74.   Xue R, Hao D-D, Sun J-P, et al. Hydrogen Sulfide Treatment Promotes Glucose Uptake

by Increasing Insulin Receptor Sensitivity and Ameliorates Kidney Lesions in Type 2

Diabetes. *Antioxid Redox Signal*. 2013;19(1):5-23. doi:10.1089/ars.2012.5024

75.   Feng X, Chen Y, Zhao J, Tang C, Jiang Z, Geng B. Hydrogen sulfide from adipose tissue

is a novel insulin resistance regulator. *Biochem Biophys Res Commun*. 2009.

doi:10.1016/j.bbrc.2009.01.059

76.   Geng B, Cai B, Liao F, et al. Increase or Decrease Hydrogen Sulfide Exert Opposite

Lipolysis, but Reduce Global Insulin Resistance in High Fatty Diet Induced Obese Mice.

PLoS One. 2013;8(9). doi:10.1371/journal.pone.0073892

77.   Lutchmansingh FK, Hsu JW, Bennett FI, et al. Glutathione metabolism in type 2

diabetes and its relationship with microvascular complications and glycemia. *PLoS*

One. 2018;13(6). doi:10.1371/journal.pone.0198626

78.   Jain SK, Micinski D, Huning L, Kahlon G, Bass PF, Levine SN. Vitamin D and L-cysteine

levels correlate positively with GSH and negatively with insulin resistance levels in the

blood of type 2 diabetic patients. *Eur J Clin Nutr*. 2014;68(10):1148-1153.

doi:10.1038/ejcn.2014.114

79.   Zhang L, Wang Y, Li Y, et al. Hydrogen sulfide (H2S)-releasing compounds: Therapeutic

potential in cardiovascular diseases. *Front Pharmacol*. 2018;9(SEP).

doi:10.3389/fphar.2018.01066

80.   Szabó G, Veres G, Radovits T, et al. Cardioprotective effects of hydrogen sulfide. In:

Nitric Oxide - Biology and Chemistry. Vol 25. ; 2011:201-210.

doi:10.1016/j.niox.2010.11.001

81.   Miller RA, Buehner G, Chang Y, Harper JM, Sigler R, Smith-Wheelock M. Methionine-

deficient diet extends mouse lifespan, slows immune and lens aging, alters glucose,

T4, IGF-I and insulin levels, and increases hepatocyte MIF levels and stress resistance.

Aging Cell. 2005. doi:10.1111/j.1474-9726.2005.00152.x

82.   Lee BC, Kaya A, Gladyshev VN. Methionine restriction and life-span control. *Ann N Y*

Acad Sci. 2016. doi:10.1111/nyas.12973

83.   Shimokawa I, Komatsu T, Hayashi N, et al. The life-extending effect of dietary

restriction requires Foxo3 in mice. *Aging Cell*. 2015. doi:10.1111/acel.12340

84.   Hine C, Kim HJ, Zhu Y, et al. Hypothalamic-Pituitary Axis Regulates Hydrogen Sulfide

Production. *Cell Metab*. 2017. doi:10.1016/j.cmet.2017.05.003

85.   Hine C, Zhu Y, Hollenberg AN, Mitchell JR. Dietary and Endocrine Regulation of

Endogenous Hydrogen Sulfide Production: Implications for Longevity. *Antioxid Redox*

Signal. 2018;28(16):1483-1502. doi:10.1089/ars.2017.7434

86.   Libiad M, Sriraman A, Banerjee R. Polymorphic variants of human rhodanese exhibit

differences in thermal stability and sulfur transfer kinetics. *J Biol Chem*. 2015.

doi:10.1074/jbc.M115.675694

87.   Billaut-Laden I, Allorge D, Crunelle-Thibaut A, et al. Evidence for a functional genetic

polymorphism of the human thiosulfate sulfurtransferase (Rhodanese), a cyanide and

H2S detoxification enzyme. *Toxicology*. 2006;225(1):1-11.

doi:10.1016/j.tox.2006.04.054

88.   Morris AA, Page RL, Baumgartner LJ, et al. Thiocyanate Accumulation in Critically Ill

Patients Receiving Nitroprusside Infusions. *J Intensive Care Med*. 2017;32(9):547-553.

doi:10.1177/0885066616657004

89.   Bromley J, Hughes BGM, Leong DCS, Buckley NA. Life-threatening interaction

between complementary medicines: Cyanide toxicity following ingestion of

amygdalin and vitamin C. *Ann Pharmacother*. 2005;39(9):1566-1569.

doi:10.1345/aph.1E634

90.   O'brien B, Quigg C, Leong T. *Severe Cyanide Toxicity from "Vitamin Supplements."* Vol

12. Lippincott Williams & Wilkins; 2005.

http://www.cancer.gov/cancerinfo/pdq/cam.laetrile.

91.   Curry SC, Carlton MW, Raschke RA. Prevention of fetal and maternal cyanide toxicity

from nitroprusside with coinfusion of sodium thiosulfate in gravid ewes. *Anesth*

Analg. 1997;84(5):1121-1126. doi:10.1213/00000539-199705000-00031

92.   Petrova F. Hydrogen cyanide and cyanides : human health aspects. 2004:67.

93.   Vinnakota C V., Peetha NS, Perrizo MG, et al. Comparison of cyanide exposure

markers in the biofluids of smokers and non-smokers. *Biomarkers*. 2012;17(7):625-

633. doi:10.3109/1354750X.2012.709880

94.   Kashala-Abotnes E, Okitundu D, Mumba D, Boivin MJ, Tylleskär T, Tshala-Katumbay D.

Konzo: a distinct neurological disease associated with food (cassava) cyanogenic

poisoning. *Brain Res Bull*. 2019;145:87-91. doi:10.1016/j.brainresbull.2018.07.001

95.   Swerdlow DI, Preiss D, Kuchenbaecker KB, et al. HMG-coenzyme A reductase

inhibition, type 2 diabetes, and bodyweight: Evidence from genetic analysis and

randomised trials. *Lancet*. 2015;385(9965):351-361. doi:10.1016/S0140-

6736(14)61183-1

Tables

Table 1. Sulfur species in blood, urine, tissue and cells

	C57Bl/6J	*Tst*$^{-/-}$	*Tst*$^{-/-}$/6J ratio	Significance
A Trunk Blood (Micromolar)				
MBB-S (Sulfide)	2.28 +/- 0.43	22.18 +/- 0.85	9.73	****
MBB-SSO3 (Thiosulfate)	n.d.	6.25 +/- 3.17	n.c.	ns
B Trunk Plasma (Micromolar)				
MBB-S (Sulfide)	1.88 +/- 0.64	24.50 +/- 2.02	13.03	****
MBB-SSO3 (Thiosulfate)	3.99 +/- 0.99	80.29 +/- 13.6	20.12	**
MBB-GSH (reduced Glutathione)	48.0 +/- 1.15	86.25 +/- 6.27	1.80	***
C Urine (Micromoles/creatine/24hrs)				
MBB-SSO3 (Thiosulfate)	4.99 +/- 2.6	2374 +/- 319	475.75	****
D Inferior Vena Cava (Micromolar)				
MBB-S (Sulfide)	1.22 +/- 0.20	3.58 +/- 0.87	2.93	ns (0.08)
MBB-SSO3 (Thiosulfate)	6.58 +/- 4.51	88.3 +/- 13.0	13.42	*
E Liver (µmoles/kg wet liver)				
MBB-S (Sulfide)	13 +/- 1	17 +/- 3	1.31	ns
MBB-SSO3 (Thiosulfate)	4 +/- 1	15 +/- 7	3.75	ns
DNFB-GSH (reduced Glutathione)	6470 +/- 380	6850 +/- 30	1.04	ns
DNFB-Cysteine (Cysteine)	82 +/- 13	67 +/- 11	0.82	ns
F Sulfide P3 fluorescence (A510nm/protein)				
Hepatocyte	7.22 +/- 1.00	7.89 +/- 0.80	1.09	ns
G Mitochondrial sulfide (MitoA)				
Liver	0.78 +/- 0.16	1.14 +/- 0.45	1.46	ns
* P < 0.05, ** P < 0.01, *** P < 0.001 **** P < 0.0001				

Table 1. *Tst* deletion results in altered sulfur metabolites in blood and liver. (**A**) Sulfide dibimane, thiosulfate-MBB, measured by fluorescence detection following HPLC, from whole blood taken from trunk blood of ND-fed C57Bl/6J (n = 4) and *Tst*$^{-/-}$ (n = 4) mice. (**B**) Sulfide dibimane, thiosulfate-MBB, and rGSH-MBB, measured by fluorescence detection following HPLC, from EDTA-Plasma of ND-fed C57Bl/6J (n = 4) and *Tst*$^{-/-}$ (n = 4) mice. (**C**) Thiosulfate-MBB corrected for creatinine from 24 hour urine samples, taken from ND-fed C57Bl6/J (n = 4) and *Tst*$^{-/-}$ (n = 5) mice. (**D**) Sulfide dibimane, and thiosulfate-MBB, from whole blood taken from the inferior vena cava downstream of the hepatic vein of ND-fed C57Bl/6J (n = 3) and *Tst*$^{-/-}$ (n = 3) mice. (**E**) Sulfide dibimane, thiosulfate-MBB, rGSH-MBB, and cysteine-MBB from whole liver (n=4/genotype) of ND-fed C57Bl/6J (n = 4) and *Tst*$^{-/-}$ (n = 4) mice. (**F**) Fluorescence from cultured hepatocytes following incubation with P3 (sulfide reactive probe) from ND-fed C57Bl/6J (n = 4) and *Tst*$^{-/-}$ (n = 4) mice. (**G**) Ratio of Mito N/MitoA from the liver of ND-fed C57Bl/6J (n = 5) and *Tst*$^{-/-}$ (n = 5) mice. Data are represented as mean ±SEM. Significance was calculated using unpaired two-tailed student's t-test. * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001

Table 2. GO Terms persulfidation rate ($Tst^{-/-}$ vs C57Bl/6J liver, ND-fed)

GO-ID	Name	Direction ($Tst^{-/-}$ vs 6J)	Genes
GO terms Identified by log fold change			
0050660	FAD binding	Decreased	12
0008168	Methyltransferase activity	Decreased	9
0016741	Transferase activity, transferring one-carbon groups	Decreased	9
0008565	Protein transporter activity	Decreased	8
0008238	Exopeptidase activity	Decreased	7
0005777	Peroxisome	Decreased	7
0042579	Microbody	Decreased	7
0003995	Acyl-CoA dehydrogenase activity	Decreased	6
0008483	Transaminase activity	Decreased	6
0016769	Transferase activity, transferring nitrogenous groups	Decreased	6
0008757	S-adenosylmethionine-dependent methyltransferase activity	Decreased	6
0016655	Oxidoreductase activity, acting on NADH/NADPH, quinone	Decreased	5
0004177	Aminopeptidase activity	Decreased	5
0000059	Protein import into nucleus, docking	Decreased	3
0005643	Nuclear pore	Decreased	3
0031965	Nuclear membrane	Decreased	3
0044453	Nuclear membrane part	Decreased	3
0046930	Pore complex	Decreased	3
0015629	Actin cytoskeleton	Decreased	3
0016652	Oxidoreductase activity, NADH/NADPH, NAD/NADP acceptor	Decreased	3
0050662	Coenzyme binding	Increased	5
0016651	Oxidoreductase activity, NADH/NADPH,	Increased	5
0003954	NADH dehydrogenase activity	Increased	4
0008137	NADH dehydrogenase (ubiquinone) activity	Increased	4
0050136	NADH dehydrogenase (quinone) activity	Increased	4
0006739	NADP metabolism	Increased	3
0006769	Nicotinamide metabolism	Increased	3
0006733	Oxidoreduction coenzyme metabolism	Increased	3

Table 2. Tst deletion results in differential persulfidation rate of liver proteins. Significant GO terms represented by peptides with different persulfidation rates in the ND-fed $Tst^{-/-}$ liver relative to C57Bl/6J. '**Direction**' indicates whether the persulfidation is decreased or increased in $Tst^{-/-}$ relative to C57Bl/6J. '**Genes**' indicates the number of genes in the $Tst^{-/-}$ that represent the changes driving the GO term.

Table 3. KEGG pathways ($Tst^{-/-}$ vs C57Bl/6J liver, ND-fed)

Entry	Name	Genes	Significance
A. Increased in ND $Tst^{-/-}$ liver			
00250	Alanine, aspartate and glutamate metabolism	6	***
00260	Glycine, serine and threonine metabolism	5	**
00270	Cysteine and methionine metabolism	4	*
04122	Sulfur relay system	2	*
B. Reduced in ND $Tst^{-/-}$ liver			
00980	Metabolism of xenobiotics by cytochrome P450	12	****
00982	Drug metabolism – cytochrome P450	12	****
05204	Chemical carcinogenesis	12	****
00480	Glutathione metabolism	8	***
00040	Pentose and glucoronate interconversions	5	**
04142	Lysosome	6	**
04390	Hippo signalling pathway	4	**
00500	Starch and sucrose metabolism	5	**
05215	Prostate cancer	3	**
04024	cAMP signalling pathway	4	*
04141	Protein processing in ER	9	*
05211	Renal cell carcinoma	3	*
00830	Retinol metabolism	6	*
00053	Ascorbate and aldarate metabolism	4	*
00860	Porphyrin and chlorophyll metabolism	4	*
04722	Neurotrophin signalling pathway	3	*
04670	Leukocyte transendothelial migration	4	*
04010	MAPK signalling pathway	4	*
04720	Long-term potentiation	2	*
04914	Progesterone-mediated oocyte maturation	2	*
04062	Chemokine signalling pathway	3	*
04110	Cell cycle	3	*
04015	Rap1 signaling pathway	4	*
00983	Drug metabolism - other enzymes	5	*
04918	Thyroid hormone synthesis	3	*
04612	Antigen processing and presentation	3	*
05203	Viral carcinogenesis	5	*

* P < 0.05, ** P < 0.01, *** P < 0.001 **** P < 0.0001

808 **Table 3. Tst deletion results in differential hepatic protein abundance of KEGG pathways.** (**A**)
809 Significant KEGG pathway terms represented by proteins that are more abundant in the ND-fed Tst^{-}
810 $^{/-}$ liver compared with ND-fed C57Bl/6J. (**B**) Significant KEGG pathway terms represented by proteins
811 that are less abundant in the ND-fed $Tst^{-/-}$ liver compared with ND-fed C57Bl/6J. '**Genes**' indicates
812 the number of genes in the $Tst^{-/-}$ that represent the changes driving the KEGG pathway.

Table 4. GO Terms: Congruence of peptide abundance and persulfidation

GO-ID	Name	Persulfidation ($Tst^{-/-}$ vs 6J)	Abundance ($Tst^{-/-}$ vs 6J)
0008483	Transaminase activity	Decreased	Increased
0016769	Transferase activity, transferring nitrogenous	Decreased	Increased
0003995	Acyl-CoA dehydrogenase activity	Decreased	Decreased
0005777	Peroxisome	Decreased	Decreased
0042579	Microbody	Decreased	Decreased

816
817 **Table 4. The $Tst^{-/-}$ liver proteome and persulfidome converges on lipid oxidation pathways.** GO
818 terms that are significantly regulated following comparison of ND-fed $Tst^{-/-}$ relative to C57Bl/6J, at
819 *both* the level of cysteine persulfidation and protein abundance.

Figures and Legends

Figure 1. *Tst* deletion results in impaired glucose and lipid metabolism. (**A**) Plasma glucose over 120 minutes, following pyruvate (i.p., 1.5mg/g) administration in overnight fasted C57Bl/6J (black line, n = 9) and $Tst^{-/-}$ (red line, n = 8) normal diet-fed mice. (**B**) Extinction of NADH measured by absorbance at 340nm, coupled to PEPCK activity from liver homogenates taken from C57Bl/6J (white bar, n = 6) and $Tst^{-/-}$ (red bar, n = 6) normal diet-fed mice. (**C**) Production of ^{13}C (M+2) acetyl-CoA and ^{13}C (M+3) aspartate generated after a 1 hour pulse of 1mM 3-carbon labelled ^{13}C (M+3) pyruvate in ^{12}C pyruvate free media, expressed as a percentage of the total amount of detected metabolite, in primary hepatocytes from C57Bl/6J (white bars, n = 6) and $Tst^{-/-}$ (red bars, n = 5) normal diet-fed mice. (**D**) Blood glucose during the pre-clamp phase of the hyperinsulinemic, euglycemic clamp from C57Bl/6J (black lines), and $Tst^{-/-}$ (red lines) fed a control (ND, solid lines, n = 3, 6) or high fat diet (HFD, broken lines, n= 6, 7). (**E**) Mean integrated radioactive glucose (inversely related to whole body glucose uptake) during a hyperinsulinemic, euglycaemic clamp from normal diet fed C57Bl/6J control (white, n = 3), and $Tst^{-/-}$ (red, n = 6) mice. (**F**) Plasma glucose expressed as % of baseline glucose, over 120 minutes following insulin (i.p., 1mU/g) administration in 4 hour fasted C57Bl/6J (black line, n = 8) and $Tst^{-/-}$ (red line, n = 7) normal diet-fed mice. (**G**) HPLC quantified total and VLDL plasma triglyceride in 4 hour fasted C57Bl/6J (white bar, n = 6), and $Tst^{-/-}$ (red bar, n = 6) normal diet-fed mice. (**H**) Representative light microscopic images of fixed liver stained with Oil-Red O from normal diet-fed (ND) or high fat diet-fed (HFD) C57Bl/6J and $Tst^{-/-}$ mice. (**I**) Analysis of the area of red staining (Oil Red O) after thresholding, using Image J, from normal diet-fed (no pattern, n = 3-4/genotype) or high fat diet-fed (hatched pattern, n = 4-5/genotype) C57Bl/6J (white bars) and $Tst^{-/-}$ (red bars) mice. Data are represented as mean ±SEM. Significance was calculated using repeated measures ANOVA (**A**,**F**) 2-WAY ANOVA (**C**,**I**), 3-WAY repeated measures ANOVA (**D**) or unpaired two-tailed student's t-test (**B**,**E**,F,**G**) * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001. For (**C**) the 2-WAY ANOVA revealed no overall effect of genotype, however a significant interaction between metabolite and genotype was found. The * on the histogram represents this interaction. For (**D**) significant effects of time (****), diet (*) and genotype (*) were found. For (**F**) the analysis was performed on absolute glucose values and demonstrated a significant effect of time (****) and an interaction between time and genotype (*). T-tests revealed that the decrement of glucose from baseline at 30 and 60 minutes after insulin was greater in the $Tst^{-/-}$ (*). For (**I**) no main genotype effect was found, but a significant effect of diet (***), and an interaction (*) were found. Post Hoc analysis using Sidaks' multiple comparison test show an effect of diet on the 6J controls (***), whereas no effect of diet is found on the $Tst^{-/-}$.

881
882
883

**Figure 2. *Tst* deletion results in increased hepatic sulfur excretion and a reduction of protein
persulfidation.** (**A**) Cysteine concentrations (MBB-HPLC) in the media incubated with primary
hepatocytes in the presence (hatched pattern) or absence (no pattern) of 1mM methionine, from
C57Bl/6J (white bars, n = 4/treatment) and $Tst^{-/-}$ (red bars, n = 4/treatment) mice. (**B**) Glutathione
concentrations (MBB-HPLC) in the media incubated with primary hepatocytes in the presence
(hatched pattern) or absence (no pattern) of 1mM methionine, from C57Bl/6J (white bars, n =
4/treatment) and $Tst^{-/-}$ (red bars, n = 4/treatment) mice. (**C**) Pie chart depicting the proportion of
liver peptides that are significantly higher (82 peptides, purple space) or lower (311 peptides, yellow
space), in their persulfidation rate in the $Tst^{-/-}$ (n = 3) relative to C57Bl/6J (n = 3) mice. (**D**) Total DTT-
released cysteine-persulfidated liver protein as measured by REVERT total protein stain following
western blotting, normalised to the total input protein of the sample from $Tst^{-/-}$ (red bar, n = 4) and
C57Bl/6J (white bar, n = 4) mice. Data with error bars are represented as mean ±SEM. Significance was
calculated using 2-WAY ANOVA (**A**, **B**) or student's t-test (**D**), * P < 0.05, ** P < 0.01. For (**A**) and (**B**)
the 2-WAY ANOVA reveals a main effect of genotype, indicated by * or ** on the histogram. A
significant effect of methionine was also found for both (**A**) and (**B**) not indicated on the histogram.
For (**C**) peptides were selected as being significant at a P-diff of 0.95 or greater.

A Proteome overlap of *Tst*⁻/⁻ with high fat feeding

B Impact of high fat feeding on proteome

Protein enrichment by organelle

C

☐ Number of down-regulated proteins ☐ Number of up-regulated proteins

Figure 3. *Tst* deletion engenders a high fat feeding–like hepatic proteome with a distinct organellar signature. (**A**) Venn diagram representing the number of proteins significantly different (at P < 0.01) between normal diet-fed *Tst*⁻/⁻ and C57Bl/6J (Red circle), and the number of regulated proteins between high fat fed and normal diet-fed C57Bl/6J (Green circle). The overlap (brown) represents those proteins regulated in the same direction by both comparisons (n= 4/genotype). (**B**) Number of proteins significantly different (at p < 0.01) between 58% high fat and normal diet in either C57Bl/6J (white bar), or *Tst*⁻/⁻ mice (Red bar), (n= 4/genotype). (**C**) Pie charts depicting the proportion of individual liver proteins that are upregulated (Blue space) compared to downregulated (yellow space) after GO term categorisation according to subcellular location. Upper row; normal diet-fed *Tst*⁻/⁻ relative to normal diet-fed C57Bl/6J. Lower row, high fat-fed C57Bl/6J relative to normal diet-fed C57Bl/6J.

Figure 4. *Tst* deletion results in increased hepatocyte respiration but impaired medium–chain fat respiration. (**A**) Electron microscope images of liver, visualising mitochondria from normal diet-fed C57Bl/6J (n = 4) or *Tst*⁻/⁻ (n = 4) mice. (**B**) Seahorse trace representing the mean oxygen consumption rate (OCR), normalised to protein, by hepatocytes from normal diet-fed C57Bl/6J (n = 6) or *Tst*⁻/⁻ (n = 6) mice during a mitochondrial stress test. (**C**) Respiratory OCR linked to ATP production (oligomycin sensitive) by hepatocytes from normal diet-fed C57Bl/6J (n = 6) or *Tst*⁻/⁻ (n = 6) mice, calculated from Figure 4B. (**D**) Respiratory OCR relating to proton leak (oligomycin insensitive) by hepatocytes from normal diet-fed C57Bl/6J (n = 6) or *Tst*⁻/⁻ (n = 6) mice, calculated from Figure 4B. (**E**) Reduction of maximal uncoupled respiration following inhibition of long chain fatty acid mitochondrial import using etomoxir (8 μM), from normal diet-fed C57Bl/6J (n = 4) or *Tst*⁻/⁻ (n = 4) mice. (**F**) Stimulation of maximal uncoupled respiration following addition of medium chain fatty acid octanoate (250 μM), from normal diet-fed C57Bl/6J (n = 4) or *Tst*⁻/⁻ (n = 4) mice. Data are represented as mean ±SEM. Significance was calculated using an unpaired two tailed, student's t-test (**C, D, E, F**), * P < 0.05.