1 Title

2 **The hepatic compensatory response to elevated systemic sulfide**
3 **impairs medium chain fat oxidation and promotes diabetes.**

4 Authors

Roderick N. Carter[1], Matthew T.G. Gibbins[1,a], Martin E. Barrios-Llerena[1,b], Stephen E. Wilkie[1,2],
Peter L. Freddolino[3], Marouane Libiad[3,c], Victor Vitvitsky[3], Barry Emerson[1,d], Thierry Le
Bihan[4,e], Madara Brice[1], Huizhong Su[5,f], Scott Denham[1], Natalie Z.M. Homer[1], Clare Mc
Fadden[1,g], Anne Tailleux[6], Nourdine Faresse[7,h], Thierry Sulpice[7], Francois Briand[7], Tom
Gillingwater[8], Kyo Han Ahn[9], Subhankar Singha[9,i], Claire McMaster[10], Richard C. Hartley[10], Bart
Staels[6], Gillian Gray[1], Andrew J. Finch[5,j], Colin Selman[2], Ruma Banerjee[3], Nicholas M. Morton[1]*

[1]University/British Heart Foundation Centre for Cardiovascular Science, University of Edinburgh, Queen's
Medical Research Institute, Edinburgh, EH16 4TJ, United Kingdom
[2]Glasgow Ageing Research Network (GARNER), Institute of Biodiversity, Animal Health and Comparative
Medicine, University of Glasgow, Glasgow, G12 8QQ, United Kingdom
[3]The Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, MI 48109, USA
[4]SynthSys - Systems and Synthethic Biology, Edinburgh, EH9 3JD, United Kingdom
[5]Cancer Research UK Edinburgh Centre, MRC Intitute of Genetics & Molecular Medicine, University of
Edinburgh, Western General Hospital, Edinburgh, EH4 2XR, United Kingdom
[6]Université de Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U101-EGID, F-59000, Lille, France
[7]Physiogenex S.A.S, Prologue Biotech, 516 rue Pierre et Marie Curie, 31670 Labège, France
[8]College of Medicine & Veterinary Medicine, University of Edinburgh, Old Medical School (Anatomy), Teviot
Place, Edinburgh, EH8 9AG, United Kingdom
[9]Department of Chemistry, POSTECH, 77 Cheongam-Ro, Nam-Gu, Pohang, Gyungbuk, 37673, South Korea
[10]School of Chemistry, Joseph Black Building, University of Glasgow, Glasgow, G12 8QQ, United Kingdom

Current affiliations where different from the place where work for the manuscript was carried out;
[a]Cambridge Biomedical Campus, Royal Papworth Hospital NHS Foundation Trust, Cardiology department,
Cambridge, CB2 0AY, United Kingdom
[b]The International Clinical Research Centre, St. Anne's University Hospital, Brno, 656 91, Czech Republic
[c]Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, 91198,
Gif-sur-Yvette cedex, France
[d]BD Research Centre Ireland, Co. Limerick, Castletroy, Ireland
[e]Rapid Novor Inc, 44 Gaukel St, Kitchener, ON N2G 4P3, Canada
[f]Wellcome Trust Centre for Mitochondrial Research, Newcastle University, NE2 4HH, United Kingdom
[g]Springer Nature Campus, London, N1 9FN, United Kingdom
[h]D.I.V.A-expertise, 1 place Pierre Potier, 31100 Toulouse, France
[i]Institute of Advanced Studies and Research, JIS University, Kolkata 700091, India
[j]Cancer Research UK Barts Centre, London, EC1M 6BQ, United Kingdom

*Lead Contact details: Nicholas M Morton
Molecular Metabolism Group, University/British Heart Foundation Centre for Cardiovascular Science,
University of Edinburgh, Queen's Medical Research Institute, Edinburgh, EH16 4TJ, United Kingdom.
nik.morton@ed.ac.uk

Competing interests statement,

The authors have declared that no conflict of interest exists.

Abstract

Impaired hepatic glucose and lipid metabolism are hallmarks of type–2 diabetes. Increased sulfide production from cysteine, or sulfide–donor compounds, may beneficially regulate hepatic metabolism. Disposal of sulfide through the sulfide oxidation pathway (SOP) is critical for maintaining sulfide within a safe physiological range. We show that mice lacking the liver–enriched mitochondrial SOP enzyme thiosulfate sulfur–transferase ($Tst^{-/-}$ mice) exhibit high circulating sulfide, increased gluconeogenesis, hypertriglyceridemia and fatty liver, despite whole–body insulin–sensitisation. Unexpectedly, hepatic sulfide levels were normal in $Tst^{-/-}$ mice, a result of homeostatic induction of mitochondrial sulfide disposal and glutathione excretion associated with net suppression of protein persulfidation and nuclear respiratory factor–2 target proteins. Proteomic and persulfidomic profiling converged on hepatic lipid metabolism, revealing a deficit in medium–chain fatty acid oxidation in $Tst^{-/-}$ mice. In conclusion our findings have implications for sulfide-donor strategies in the context of liver function and metabolic disease.

Introduction

The prevalence of Type 2 diabetes (T2D) continues to soar in parallel with that of obesity[1]. Increased hepatic glucose production and aberrant hepatic lipid metabolism are cardinal features of T2D[2,3]. Dysregulation of hepatic nutrient metabolism in T2D is a promising area for therapeutic intervention because it precipitates the more severe liver pathologies that manifest along the spectrum of non-alcoholic fatty liver disease (NAFLD), steatosis, steatohepatitis and hepatocellular carcinoma[4].

Hydrogen sulfide (hereafter referred to as sulfide), an endogenously produced gaseous signalling molecule[5–8], has recently emerged as a modulator of nutrient metabolism[9–12]. Enzymatic sulfide production from sulfur amino acids is catalysed by cystathionine beta–synthase; CBS, and cystathionine gamma lyase; CTH[13,14] and by 3-mercaptopyruvate sulfur transferase; MPST[15–17]. Thioredoxin-mediated reduction of cysteine persulfides on proteins also regulates free sulfide and cysteine persulfide levels[18]. Endogenously produced and exogenously administered sulfide specifically influences hepatic glucose and lipid metabolism[19,20]. Thus, *in vitro,* treatment of murine hepatocytes with NaHS, or overexpression of rat *Cth* in HepG2 liver cells, increased glucose production through increased gluconeogenesis and reduced glycogen storage[21]. Conversely, glucose production was lower in hepatocytes from *Cth* gene knockout mice (*Cth*$^{-/-}$ mice) that exhibit low sulfide production[21]. Elevation of sulfide with NaHS administration *in vivo* reduced cholesterol and triglyceride accumulation in the liver of high fat diet (HFD)–fed mice[22]. In contrast, inter-cross of sulfide production–deficient *Cth*$^{-/-}$ mice with the hyperlipidemic *Apoe*$^{-/-}$ mouse strain (*Cth*$^{-/-}$*Apoe*$^{-/-}$ mice) produced a phenotype of elevated plasma cholesterol following exposure to an atherogenic diet[23]. Consistent with their higher cholesterol, *Cth*$^{-/-}$*Apoe*$^{-/-}$ mice developed fatty streak lesions earlier than *Apoe*$^{-/-}$ mice, and this effect was reversed

by NaHS administration[23]. Sulfide may also indirectly impact hepatic nutrient metabolism

through its effect on hepatic artery vasorelaxation and thus liver perfusion[24,25]. The

apparently beneficial effects of sulfide administration in multiple disease indications has led

to a major drive towards development of targeted H_2S–donor molecules as a therapeutic

approach[26,27]. However, an often-overlooked aspect of net sulfide exposure, key to the

efficacy of therapeutic H_2S–donors, is that it is regulated through its oxidative disposal. Thus,

endogenous sulfide exposure is actively limited to prevent mitochondrial respiratory

toxicity[28–30]. Sulfide is rapidly oxidized[31,32] through the mitochondrial sulfide oxidation

pathway (SOP), consisting of sulfide quinone oxidoreductase (SQOR), persulfide dioxygenase

(ETHE1/PDO) and thiosulfate sulfurtransferase (TST, also known as rhodanese)[31,33,34]. The

liver is highly abundant in SOP enzymes, and is a major organ of whole body sulfide disposal[32].

Mice lacking the *Ethe1* gene (*Ethe1*$^{-/-}$ mice) die of fatal sulfide toxicity[30], consistent with its

critical role in sulfide oxidation and the severe pathological consequences of unchecked

sulfide build-up in tissues. However, the importance of mitochondrial TST in the SOP *in vivo*

remains obscure. In contrast to *Ethe1*$^{-/-}$ mice, *Tst*$^{-/-}$ mice were grossly normal despite

exhibiting substantially elevated blood sulfide levels, as implied by qualitative measures[35].

This revealed an important yet distinct role for TST in the SOP for the first time *in vivo*.

Nevertheless, *Tst*$^{-/-}$ mice showed an apparently diabetogenic impairment of glucose

tolerance[35], consistent with the concept that increased sulfide promotes hepatic glucose

production[21]. As *Tst* deficiency represents a model of chronic but viable sulfide elevation,

determining the molecular mechanisms driving their aberrant metabolic profile can provide

important insights into the optimal range for therapeutic sulfide exposure, particularly in light

of the current interest in developing mitochondrially–targeted sulfide–donors[36,37]. To this end

we sought to define the impact of *Tst* deficiency on the underlying molecular pathways that

impact hepatic metabolism.

Results

Tst$^{-/-}$ mice exhibit increased hepatic gluconeogenesis and dyslipidaemia despite peripheral insulin sensitisation. TST *mRNA* expression is highest in the liver (e.g. it is ~8-fold higher than in white adipose tissue http://biogps.org/#goto=genereport&id=22117). We therefore hypothesised that liver TST deficiency was the principal driver of the impaired glucose tolerance observed in *Tst$^{-/-}$* mice[35]. We first tested this by challenging the mice with pyruvate, which is converted to glucose in the liver. *Tst$^{-/-}$* mice exhibited higher glucose production than C57BL/6J controls in response to pyruvate (Figure 1A). We next tested phosphoenolpyruvate carboxykinase (PEPCK) activity, the rate-determining enzyme in *de-novo* hepatic glucose synthesis, and found it was higher in liver homogenates from *Tst$^{-/-}$* mice (Figure 1B). Next, we traced the metabolite incorporation of $^{13}C_3$-pyruvate by isolated *Tst$^{-/-}$* hepatocytes, which showed a significant reduction in $^{13}C_2$-acetyl-CoA and a significant increase in $^{13}C_3$-aspartate, consistent with skewing of the tricarboxylic acid cycle towards gluconeogenic precursors (Figure 1C). We also assessed whether *Tst* deficiency is associated with impaired lipid metabolism, another hallmark of diabetes. Fast protein liquid chromatography analysis of triglyceride levels and their lipoprotein distribution revealed significantly higher total plasma triglycerides in *Tst$^{-/-}$* mice (Figure 1D). The higher triglyceride was selectively associated with an increased VLDL triglyceride fraction (Figure 1D), consistent with a dominant liver-driven impairment in lipid metabolism[38]. Total and distinct lipoprotein fraction plasma cholesterol levels were similar between genotypes (Figure S1A and S1B), suggestive of a triglyceride–selective effect of *Tst* deficiency on hepatic lipid efflux. HFD significantly increased liver lipid content of C57BL/6J mice but did not further increase the lipid levels in the liver of *Tst$^{-/-}$* mice (Figure 1E-1F). Liver glycogen, a marker of long–term

carbohydrate storage was comparable between $Tst^{-/-}$ and C57BL/6J control mice (Figure

S1C). Steady–state phosphorylation levels of hepatic insulin–responsive proteins AKT and the

cellular energy–sensing protein AMPK were comparable between $Tst^{-/-}$ mice and C57BL/6J

littermates (Figure S1D).

Despite unchanged steady–state markers of hepatic insulin sensitivity, impaired glucose

tolerance suggested that whole body, and usually hepatic, insulin–resistance was present. To

test this, we performed an insulin tolerance test. Unexpectedly, the glucose decrement in

response to insulin was greater in $Tst^{-/-}$ mice (Figure 1G), demonstrating a net increase in

dynamic whole–body insulin sensitivity despite increased hepatic glucose output. We

confirmed increased whole–body insulin sensitivity under short–term steady–state

conditions using euglycaemic hyperinsulinaemic clamps. Blood glucose tended to be higher

in the $Tst^{-/-}$ mice during baseline conditions (Figure 1H). During the clamp, when blood

glucose levels were maintained constant, the glucose infusion rate was comparable between

genotypes (Table S1). However, an overall increase in glucose uptake (integral glucose) by

tissues in the $Tst^{-/-}$ mice was apparent (Figure 1I, Table S1), supporting increased peripheral

insulin–sensitivity concurrent with higher endogenous glucose production. After HFD feeding,

$Tst^{-/-}$ mice exhibited relatively lower fasting insulin and higher hepatic glucose output across

baseline pre-clamp conditions (Table S2). However, the metabolic phenotype of $Tst^{-/-}$ mice

largely converged on that of the C57BL/6J mice during the hyperinsulinemic clamp (Table S2).

TST deficiency elicits compensatory hepatic sulfide disposal mechanisms that drive reduced

global protein persulfidation. A role for TST in the disposal of sulfide was suggested by its

participation in the SOP[31,34] and supported *in vivo* by the qualitatively higher blood sulfide of

$Tst^{-/-}$ mice[35]. Here, we quantified circulating sulfide, showing approximately 10–fold

elevation in the blood and plasma of $Tst^{-/-}$ mice (Table 1A and 1B). Thiosulfate, an oxidised

metabolite of sulfide [39,40] and a TST substrate[41], was approximately 20–fold higher in plasma

(Table 1B), and profoundly higher (450–fold) in urine (Table 1C) of $Tst^{-/-}$ mice compared to

C57Bl/6J mice. Reduced glutathione (GSH) levels were ~2–fold higher in the plasma of $Tst^{-/-}$

mice (Table 1B). To determine any direct hepatic contribution to the elevated systemic sulfide

in vivo, whole blood was sampled from the inferior vena cava (IVC) (Table 1D). IVC sulfide

levels tended to be higher in the $Tst^{-/-}$ mice, but the magnitude of the increase (~3-fold) did

not parallel that in trunk blood (~10-fold), suggesting liver was not a major source of the

elevated circulating sulfide. Surprisingly, liver sulfide, thiosulfate, cysteine and GSH levels

were similar between $Tst^{-/-}$ and C57BL/6J mice (Table 1E). Further, cultured hepatocytes

from $Tst^{-/-}$ and C57BL/6J mice exhibited similar intracellular sulfide levels, as estimated

using P3, a sulfide–selective fluorescent probe[42] (Table 1F). Mitochondrial sulfide levels in

liver reported by MitoA/MitoN[43] were similarly unchanged between genotypes (Table 1G).

The apparently unaltered hepatic steady-state sulfide levels, despite higher circulating

sulfide, suggested a profound homeostatic mechanism was invoked in the liver of $Tst^{-/-}$

mice. We assessed respiratory sulfide disposal (antimycin–sensitive) and found this was

markedly increased in hepatocytes from $Tst^{-/-}$ mice, whereas antimycin–insensitive sulfide

disposal was relatively reduced compared to hepatocytes from C57BL/6J mice (Table S3).

Isolated liver mitochondria from $Tst^{-/-}$ hepatocytes also exhibited a higher sulfide disposal

rate (Table S3). In addition, cysteine and GSH was excreted at higher levels from $Tst^{-/-}$

hepatocytes under basal conditions and after stimulation of sulfur amino acid metabolism by

addition of methionine (Figure 2A and 2B). Consistent with higher GSH turnover, hepatocytes

from $Tst^{-/-}$ mice showed resistance to exogenous H_2O_2–mediated mitochondrial ROS

production (Figure S2). We next determined the global hepatic protein persulfidation profile,

the major post–translational modification mediated by sulfide[44–46]. Mass spectrometry

analysis of maleimide–labelled liver peptides revealed a greater abundance of peptides with

a lower persulfidation level (under-persulfidated) in the liver of $Tst^{-/-}$ mice (Figure 2C). We

confirmed this using semi-quantitative western–blot analysis on pulled down maleimide–

labelled proteins (Figure 2D). Gene Ontology (GO) analysis of proteins exhibiting altered

persulfidation levels revealed 28 terms different between $Tst^{-/-}$ and C57BL/6J mice by log–

fold change (Table 2). Major under-persulfidated terms (20) included "*FAD-binding, methyl*

transferase, peroxisome, acyl-CoA dehydrogenase activity and transaminase" while the less

abundant (8) over-persulfidated terms were dominated by terms related to "*Nicotinamide*

metabolism".

The hepatic proteome of Tst$^{-/-}$ mice reveals a distinct molecular signature of altered sulfur

and mitochondrial nutrient metabolism. To gain molecular insight into the mechanisms

underlying the apparently diabetogenic liver phenotype in ND-fed $Tst^{-/-}$ mice, we performed

a comparative iTRAQ proteomic analysis of whole liver with ND-fed C57BL/6J mice. Kyoto

Encyclopedia of Genes and Genomes (KEGG) analysis (at significance threshold $P < 0.05$)

revealed 4 up–regulated pathways in liver of $Tst^{-/-}$ mice related to amino acid metabolism,

including sulfur amino acids, and sulfur metabolism (Table 3A). GO analysis (at threshold $P <$

0.05) revealed 95 significantly up–regulated terms in liver of $Tst^{-/-}$ mice (Table S4A). Of the

top 14 up–regulated terms, 7 referred to amino acid metabolism and 1 referred to the

organellar term '*mitochondrion*'. KEGG pathway analysis revealed 27 down–regulated

pathways in the liver of $Tst^{-/-}$ mice (Table 3B) including Phase 1 and 2 detoxification

pathways (Cytochrome P450s, Glutathione and Glucuronidation) and '*Lysosome*', '*Protein*

processing in the Endoplasmic reticulum' organellar terms. 213 GO terms were significantly

down–regulated in $Tst^{-/-}$ mice (Table S4B). Among the most significant down–regulated

terms were phase 2 detoxification pathways such as '*glutathione binding'* and '*glutathione*

transferase activity' and 'endoplasmic reticulum'. We validated a representative subset of protein changes (Figures S3A), noting broadly consistent direction of change in the key pathways. At the single protein level, the most robust change we observed was increased MPST protein in whole liver (Figure S3A) and mitochondrial sub-fractions (Figure S3B). This change was remarkable as mRNA levels for *Mpst* were lower in *Tst*$^{-/-}$ mice (Figure S3C), likely as a result of loss of proximal *Mpst* promoter function; *Mpst* is a paralog of *Tst*[47] juxtaposed approximately 1kb from *Tst*. Given the evidence for glucose and lipid metabolic changes in the *Tst*$^{-/-}$ mice, we undertook a further focussed comparison of canonical proteins in these pathways (Table S5). Four GO terms were enriched in liver from *Tst*$^{-/-}$ mice, all of which were down–regulated; *'Lipid metabolic process'*, *'fatty acid beta-oxidation'*, *'Acyl-CoA dehydrogenase activity'* and *'Acyl-CoA hydrolase activity'* (Table S5).

Hepatic protein expression in Tst$^{-/-}$ *mice is consistent with lower NRF2 activation*. We performed a transcription factor binding site (TFBS) enrichment analysis in the promoters of proteins that are up–regulated in the liver of *Tst*$^{-/-}$ mice to look for potential hub transcriptional drivers of the proteome profile (Figure S4A). This revealed a statistically significant under-representation of TFBS for the sulfide–responsive[48,49] NRF2 transcription factor (Figure S4A). To explore whether there was evidence for reduced NRF2 activation in the *Tst*$^{-/-}$ liver, we looked at the relative expression of 47 NRF2–regulated proteins in the proteome of ND-fed *Tst*$^{-/-}$ and ND-fed C57Bl/6J mice. While 37 of these target proteins were similar in expression between the genotypes, 10 NRF2 target proteins were lower in *Tst*$^{-/-}$ liver (Figure S4B) and none of the 47 NRF2 target proteins were increased in *Tst*$^{-/-}$ liver.

Proteome of TST deficiency versus HFD response in C57BL/6J mice reveals distinct regulation of lipid metabolism and detoxification pathways. We examined mechanistic commonalities

between the diabetogenic hepatic phenotype of $Tst^{-/-}$ mice and that induced by the diabetogenic HFD–feeding regimen in C57BL/6J mice. Note, ND-fed $Tst^{-/-}$ mice were in a pre-existing diabetogenic state (Figure 1) that does not further worsen with HFD (Figure 1E-F, Table S1B), suggesting gross phenotypic convergence of the two genotypes after HFD. We compared the identity and direction of change of the 188 proteins differentially expressed in ND-fed $Tst^{-/-}$ mice (versus ND-fed C57BL/6J mice; Figure 3A) to proteins that were differentially expressed in response to HFD in C57BL/6J mice (432 proteins; Figure 3A). There was a striking 67% overlap in individual proteins (126) in this comparison (Figure 3A). When we analysed these two protein signatures for directionally shared pathways, one KEGG pathway *'Glycine, serine and threonine metabolism'* was up–regulated (Table S6A) and 12 KEGG pathways were down–regulated, including *'drug metabolism'* and *'endoplasmic reticulum'* (Table S6B). Consistent with a pre-existing HFD–like proteome, the proteomic response in the liver of $Tst^{-/-}$ mice to HFD was muted, relative to that observed in C57BL6J mice (106 proteins, a 4–fold lower response; Figure 3B). We then considered contrasting, rather than congruent, proteomic responses arising from TST deficiency versus HFD responses in C57BL/6J mice to illuminate potential novel pathways underlying the otherwise functionally similar overall diabetogenic hepatic $Tst^{-/-}$ phenotype. 5 KEGG pathways (Table S7A) and 4 GO terms (Table S7B) were regulated in the opposite direction in this comparison. Strikingly, the GO terms were all related to lipid metabolism, which were up–regulated in the HFD response, but down–regulated with TST deficiency (Table S7A-B). An organelle–focussed protein analysis showed shared up-regulation of mitochondrial and endoplasmic reticulum pathways between TST deficiency (Figure 3C upper row) and C57BL/6J HFD-responses (Figure 3C lower row), but a striking discordance in peroxisomal protein pathways (up-regulated by

HFD, down-regulated TST deficiency) and nuclear proteins (down-regulated by HFD, up-

regulated TST deficiency; Figure 3C).

The Tst⁻/⁻ liver proteome and persulfidome converge on transamination and lipid oxidation

pathways. To assess whether conservation of changes at protein and post-translational

modification levels can illuminate key regulatory hubs driving the hepatic phenotype we ran

a congruence analysis of the proteome and persulfidome. We found that GO terms referring

to amino acid, lipid metabolism and peroxisome were significantly regulated at both the level

of protein abundance and persulfidation status (Table 4).

Tst⁻/⁻ hepatocytes exhibit elevated mitochondrial respiration and a defect in medium–chain

fatty acid oxidation. Enhanced respiratory sulfide disposal was found from *Tst⁻/⁻*

hepatocytes, and enrichment of mitochondrial proteins was suggested from the liver

proteome of the *Tst⁻/⁻* mice (Table S4). We therefore sought to determine if TST deficiency

affected respiratory function and substrate utilisation of the hepatocyte. Analysis of electron

micrographs prepared from the liver of ND-fed *Tst⁻/⁻* mice and C57BL/6J controls showed

morphologically normal mitochondria (Figure 4A). Basal respiration, comprising ATP–linked

and leak respiration, was significantly higher in hepatocytes from *Tst⁻/⁻* mice (Figure 4B–4D).

Maximal hepatocyte respiratory capacity and non–respiratory oxygen consumption was

similar between genotypes (Figures S5A and S5B). In line with phenotypic convergence

following HFD, hepatocyte respiration was comparable between genotypes from HFD-fed

mice (Figures S5C-S5H). A unique feature of the liver from *Tst⁻/⁻* mice that contrasted with

HFD response in C57BL/6J mice was a decrease in proteins and persulfidation levels of

proteins in lipid oxidation pathways. We therefore investigated hepatocyte respiration of

lipids. Using a low pyruvate (100 μM) medium to reveal respiratory dependency on other

substrates, we showed that CPT1A–mediated mitochondrial oxidation of endogenous long

chain fatty acids (LCFA; etomoxir-inhibited) was similar between genotypes (Figure 4E). Next,

we by–passed CPT1A–mediated LCFA transfer, and revealed a marked deficit in respiration

stimulated by the medium chain fatty acid octanoate in hepatocytes from $Tst^{-/-}$ mice (Figure

4F). A similar experiment adding back pyruvate revealed comparable stimulation of

respiration between genotypes (Figure S5I). In amino acid free media, combined glutamine–,

aspartate– and alanine–stimulated hepatocyte respiration was comparable between

genotypes (Figure S5J).

Discussion

Elevated TST expression in adipose tissue was recently identified as a genetic mechanism driving metabolically protective leanness in mice[35]. Conversely, mice with genetic deletion of *Tst* (*Tst*$^{-/-}$ mice) exhibited impaired glucose tolerance relative to C57BL/6N controls[35]. A key observation from the *Tst*$^{-/-}$ mouse was qualitative evidence for elevated blood sulfide levels[35], which implicated the mitochondrial sulfurtransferase in the breakdown of sulfide for the first time *in vivo*. However, the mechanistic link between *Tst* deficiency and impaired glucose tolerance remained unclear. The *Tst*$^{-/-}$ mice had a subtle adipose tissue phenotype, with comparable fat accumulation upon high fat feeding, suggesting alternate organs as a potential source of impaired glucose homeostasis. Moreover, TST is most highly abundant in the liver, consistent with its role in detoxification of cyanide and related molecules. Indeed, hepatic TST mRNA levels are ~8-fold higher than adipose tissue, with the difference in enzyme activity likely substantially exceeding this, per gram of tissue. Based upon the reported effects of sulfide to increase hepatic glucose production[20,21] and modulate lipid metabolism[22,23], we hypothesised that TST deficiency predominantly drives a diabetogenic state through increased hepatic sulfide exposure. Consistent with this, we found evidence for increased gluconeogenesis, steatosis, elevated plasma VLDL triglycerides and, quantitatively, a 10-fold higher circulating blood sulfide level in *Tst*$^{-/-}$ mice. Unexpectedly, and in spite of the markedly increased circulating sulfide levels, steady-state sulfide level was normal in the liver of *Tst*$^{-/-}$ mice. Moreover, we found evidence for multiple mechanisms for increased, not decreased, hepatic sulfide disposal, reduced downstream sulfide signalling and novel molecular explanations for the apparently diabetogenic phenotype. Our data are consistent with the liver being a major site for sulfide disposal rather than production[32]. Moreover, our

data suggest that the liver of $Tst^{-/-}$ mice has over-compensated in its response to maximise

hepatic sulfide removal, thus engendering detrimental metabolic consequences indirectly.

This response is achieved through a combination of distinct compartmentalised cellular

responses including increased respiratory sulfide disposal and export of cysteine and GSH and

is further evidenced by the reduction of global hepatic protein persulfidation and depletion

of sulfide-responsive[49] NRF2 target proteins. Additionally, up-regulation of translation and

recruitment of MPST to the mitochondria of $Tst^{-/-}$ mice is observed. This response, in face

of reduced transcription of $Mpst$ due to $Mpst$ promoter disruption by the Tst null allele,

suggests a powerful post-transcriptional cellular sulfide sensing mechanism. Moreover, if

MPST is compensating for TST–mediated sulfide disposal in this context, this suggests a

subversion of normal MPST function away from sulfide production[50,51 52,53]. Intriguingly, TST

expression levels were elevated in the liver of $Mpst^{-/-}$ mice, providing further support for an

exquisite reciprocal compensatory mechanism between these two enzymes[54].

The unexpected finding of normal hepatic sulfide levels in the $Tst^{-/-}$ mice led us to discover

that the metabolic phenotype we observed was driven by the very mechanisms invoked to

maintain sulfide within a normal range rather than sulfide excess *per se*. Several observations

were consistent with this. For example, the major amino acid pathways increased in the liver

proteome of $Tst^{-/-}$ mice were transaminases involved in metabolism of GSH that support

increased export of sulfur equivalents as GSH (and cysteine). These same transaminases

support gluconeogenesis by redirecting Krebs cycle intermediates[55–57]. In addition,

glutathione-S-transferases (GST) that inhibit gluconeogenesis[58] were lower in the liver of $Tst^{-/-}$

mice. Further, NRF2 activation, which represses gluconeogenesis[59] appears lower in liver

of $Tst^{-/-}$ mice. The involvement of NRF2 in the $Tst^{-/-}$ liver phenotype is further supported by

the phenotype of $Nrf2^{-/-}$ mice that also exhibited steatohepatitis, in the absence of insulin

resistance[60]. However, we note that NRF2 signalling can be complex and dependent upon

dietary context; $Nrf2^{-/-}$ mice showed improved glucose tolerance after a high–fat diet[61]

suggesting any contribution of a NRF2 signalling deficit in the liver of the $Tst^{-/-}$ mice changes

upon high fat feeding. Beyond gluconeogenesis, we also showed that the hepatocytes of $Tst^{-/-}$

mice exhibited defective lipid metabolism. Specifically, medium–chain fatty acid (MCFA)

oxidation was impaired, associated with selective reduction of protein levels and

persulfidation levels of lipid catabolic enzymes. This represents a novel mechanism linking

altered sulfide metabolism to lipid oxidation, hepatic lipid accumulation and dyslipidaemia.

Consistent with impaired MCFA oxidation defects as a driver, at least in part, of the phenotype

of $Tst^{-/-}$ mice, steatosis is observed in medium–chain acyl-CoA dehydrogenase $(Mcad)^{-/-}$

mice[62] and dyslipidemia is found in MCADD deficient humans[63]. The data we present adds to

a growing understanding of the link between sulfide regulating genes and nutrient

metabolism that has hitherto focussed on the enzymes of sulfide production. Specifically, we

provide support for the importance of the sulfide oxidising pathway as a regulator of cellular

sulfide exposure. Unexpectedly, the data reveal novel cellular mechanisms that are engaged

to homeostatically regulate sulfide disposal and that can impact upon cell energetics and

nutrient metabolism.

Our findings may have implications for potentially unexpected side effects of sulfide donor

therapeutics. In normal mice, *in vivo* sulfide administration for 4 weeks post HFD partially

reversed hepatic lipid accumulation invoked by chronic (16 weeks) HFD[22]. No evidence was

provided for whether sulfide disposal mechanisms were altered[22]. This efficacious sub-

chronic sulfide administration regimen contrasts with our genetic model of chronic sulfide

elevation as a driver of dysregulated metabolism and NAFLD. Clearly, the normal mice in the

Na_2S administration studies had a fully functional SOP, suggesting the presence of TST is

required to achieve the beneficial metabolic effects of Na_2S administration. This is also

consistent with the apparently low sulfide signalling status (evidenced by lower

persulfidation, NRF2 target protein abundance) in the liver of the $Tst^{-/-}$ mice. The benefits

of elevated sulfide cannot be realised, perhaps because the major mediator of those effects

is missing and the alternate mechanisms invoked do not fully compensate (e.g. MPST) or

actively drive aberrant nutrient metabolism.   Comparable studies of glucose and lipid

metabolism after manipulation of other sulfide regulating genes are limited. However in a

contrasting model of reduced sulfide production ($Cth^{-/-}$ mice), plasma triglycerides were

lowered[23], consistent and opposite to what we observed with the $Tst^{-/-}$ mice. The hepatic

sulfide disposal status of the $Cth^{-/-}$ mouse model is unknown, but our findings predict a

suppression of the SOP to spare the limited endogenous sulfide produced. Intriguingly they

also predict a knock-on effect on nutrient homeostasis due to reduced metabolic demand of

the TST/SOP axis. A more direct model informing on the effects of impairment of the sulfide

disposal pathway is deficiency of the key mitochondrial SOP enzyme ETHE1. $Ethe1^{-/-}$ mice

suffer fatal sulfide toxicity[30] and therefore comparable metabolic studies are lacking.

However, one notable observation is that $Ethe1^{-/-}$ mice have an apparently 10-fold higher

liver sulfide exposure than control mice[30], in contrast to the normalised hepatic sulfide levels

of $Tst^{-/-}$ mice. Circulating sulfide levels were not reported for comparison, but the

presumably relatively lower systemic sulfide levels of $Tst^{-/-}$ mice appear to have permitted

an effective homeostatic sulfide disposal response in the liver to avoid toxicity, albeit with a

metabolic cost. Consequently, the liver of $Tst^{-/-}$ mice has a distinct functional and proteomic

profile to that of the $Ethe1^{-/-}$ mice. For example, in the liver of $Tst^{-/-}$ and $Ethe1^{-/-}$ mice[64],

proteins of the glutathione–S–transferase Mu type (GSTM) and peroxiredoxin (PRDX) families

were altered, but sometimes in the opposite direction or with alteration of distinct protein

sub-classes. A notable difference is also observed in amino acid metabolism. The liver of

Ethe1$^{-/-}$ mice exhibited increased expression of enzymes of branched chain amino acid

metabolism[64], distinct from the predominantly glutathione–related amino acid pathways that

are increased in liver of *Tst*$^{-/-}$ mice. Beyond sulfide, TST may also have distinct cellular roles

that affect metabolism such as mitoribosomal synthesis, ROS attenuation and modulation of

mitochondrial iron-sulfur clusters[65–69].

Given the pro-diabetogenic liver phenotype in *Tst*$^{-/-}$ mice, its was surprising that insulin

signalling in the liver appeared normal, and peripheral insulin sensitivity was increased. There

are precedents for increased hepatic glucose production independent of insulin resistance, as

found in the *Nrf2*$^{-/-}$ mice[60] and as driven by the transcription factor ChREBP[70,71]. There is

also evidence to support insulin–sensitising effects of sulfide administration *in vivo* in mice

and rats[72–74], consistent with a sulfide–mediated insulin–sensitisation of non-hepatic tissues

in *Tst*$^{-/-}$ mice. Higher circulating GSH in *Tst*$^{-/-}$ mice may also promote peripheral insulin–

sensitisation[75,76]. Clearly, the net balance of glucose production from the liver and its

peripheral disposal remains abnormal in *Tst*$^{-/-}$ mice. Indeed, the baseline metabolic

phenotype of *Tst*$^{-/-}$ mice resembles in many ways that of a normal mouse fed a HFD. Thus

we expected to gain insight into conserved pro-diabetogenic mechanisms from the overlap

between the liver proteome of *Tst*$^{-/-}$ mice and that of HFD-fed C57BL/6J mice. However,

upon closer inspection this was not entirely the case, with some lipid metabolism and

peroxisomal proteins changing in the opposite direction to those observed in HFD–fed

C57BL/6J mice. Unlike a HFD state, which is associated with dominant hepatic insulin

resistance, the increased hepatic glucose production in ND-fed *Tst*$^{-/-}$ mice appears to be a

result of their over-compensated sulfide-removal response.

Sulfide donor therapeutics were proposed as a clinical strategy for improving cardiovascular

health[27,77,78]. Elevated endogenous sulfide was also implicated in the beneficial metabolic

effects of caloric restriction[11,79–83]. Our results suggest that chronic sulfide elevation may have

unintended detrimental consequences, driving liver glucose production and fat accumulation

to undesirable levels. This caveat may be fortunately limited to cases where SOP proteins are

compromised through rare genetic effects – such as TST variants[84,85]. More broadly, a number

of drugs or supplements are known to increase cyanide, which may dominantly inhibit TST

activity and result in secondary sulfide overexposure. These include nitroprusside[86] and

amygdalin[87,88]. Indeed the TST metabolite thiosulfate is commonly co-administered with

nitroprusside to prevent cyanide toxicity[89]. Furthermore, dietary and environmental exposure

to cyanogenic compounds[90], e.g. smoking[91] or cyanogenic diets[92] may interfere with normal

TST function and could lead to increased sensitivity to sulfide therapeutics. In contrast, we

have shown that administration of the TST substrate thiosulfate can ameliorate diabetes[35]

further underlining the potential utility of targeting the SOP in metabolic disease. As with all

therapeutic strategies, a careful cost-benefit analysis is required. A comparable case of

relevance are the statins, one of the most potent and widely used drugs to prevent

atherosclerosis, which also carry a higher risk for diabetes[93]. The full impact of TST

manipulation on opposing metabolic pathways requires further study. Our current study

sheds light on the underlying hepatic mechanisms invoked for sulfide disposal that are

relevant to current sulfide–donor strategines and may inform on routes to reduce their

potential metabolic side-effects.

Methods

The methods for all data and work presented in this manuscript is provided as supplementary information.

Author Contributions

N.M.M. and R.N.C conceived experiments. R.N.C., M.T.G.G, M. B.-L., A. M.-C., M.L., V.V., B.E., T. LeB., M.B., S.H., S.D., N.H., C. Mc F., A.T., N. F., T.G., performed experiments. R.N.C., M. B.-L., P.F., V.V., T. LeB., N.H., T.S., F.B., T.G., R.C.H., B.S., G.G., A.J.F., C.S., R.B. and N.M.M. analysed and interpreted data and commented on the manuscript. K.H.A., S.S. generated reagents. C.McM. and R.C.H. generated reagents. R.N.C. and N.M.M. wrote the manuscript.

Acknowledgements

This work was funded by a Wellcome Trust New Investigator Award to NMM (100981/Z/13/Z) and Diabetes UK grant (17/0005697). RCH was funded by a Medical Research Council Discovery Award (MC-PC-15076). We would like to thank Professor Ken Olsen and Eric DeLeon (Notre Dame) for advice establishing sulfide measurement by probe.

References

1. World Health Organisation. Global report on diabetes.
   2016;https://www.who.int/publications-detail/global-report-on-diabetes

2. Lewis GF, Carpentier A, Adeli K, Giacca A. Disordered fat storage and mobilization in
   the pathogenesis of insulin resistance and type 2 diabetes. *Endocr Rev*.
   2002;23(2):201-229. doi:10.1210/edrv.23.2.0461

3. Consoli A, Nurjhan N, Capani F, Gerich J. Predominant role of gluconeogenesis in
   increased hepatic glucose production in NIDDM. *Diabetes*. 1989;38(5):550-557.
   doi:10.2337/diab.38.5.550

4. Caron S, et al. Transcriptional activation of apolipoprotein CIII expression by glucose
   may contribute to diabetic dyslipidemia. *Arterioscler Thromb Vasc Biol*.
   2011;31(3):513-519. doi:10.1161/ATVBAHA.110.220723

5. Abe K, Kimura H. The possible role of hydrogen sulfide as an endogenous
   neuromodulator. *J Neurosci*. 1996;16(3):1066-1071.
   doi:10.1523/JNEUROSCI.16-03-01066

6. Wang R. Physiological Implications of Hydrogen Sulfide: A Whiff Exploration That
   Blossomed. *Physiol Rev*. 2012;92(2):791-896. doi:10.1152/physrev.00017.2011

7. Mishanina T V., Libiad M, Banerjee R. Biogenesis of reactive sulfur species for
   signaling by hydrogen sulfide oxidation pathways. *Nat Chem Biol*.
   2015;11(7):457-464. doi:10.1038/nchembio.1834

8. Filipovic MR, Zivanovic J, Alvarez B, Banerjee R. Chemical Biology of H_2S Signaling
   through Persulfidation. *Chem Rev*. 2018;118(3):1253-1337.
   doi:10.1021/acs.chemrev.7b00205

9. Desai KM, Chang T, Untereiner A, Wu L. Hydrogen sulfide and the metabolic

syndrome. *Expert Rev Clin Pharmacol*. 2011;4(1):63-73. doi:10.1586/ecp.10.133

10.   Szabo C. Roles of Hydrogen Sulfide in the Pathogenesis of Diabetes Mellitus and Its

Complications. *Antioxid Redox Signal*. 2011;17(1):68-80. doi:10.1089/ars.2011.4451

11.   Hine C, et al. Endogenous hydrogen sulfide production is essential for dietary

restriction benefits. *Cell*. 2015;160(1-2):132-44. doi:10.1016/j.cell.2014.11.048

12.   Carter RN, Morton NM. Cysteine and hydrogen sulphide in the regulation of

metabolism: Insights from genetics and pharmacology. *J Pathol*. 2016;238(2):321-332.

doi:10.1002/path.4659

13.   Chen X, Jhee KH, Kruger WD. Production of the neuromodulator H2S by

cystathionine beta-synthase via the condensation of cysteine and homocysteine. *J*

Biol Chem. 2004;279(50):52082-6. doi:10.1074/jbc.C400481200

14.   Singh S, Padovani D, Leslie RA, Chiku T, Banerjee R. Relative contributions of

cystathionine beta-synthase and gamma-cystathionase to H2S biogenesis via

alternative trans-sulfuration reactions. *J Biol Chem*. 2009;284(33):22457-66.

doi:10.1074/jbc.M109.010868

15.   Shibuya N, et al. 3-Mercaptopyruvate sulfurtransferase produces hydrogen sulfide

and bound sulfane sulfur in the brain. *Antioxid Redox Signal*. 2009;11(4):703-14.

doi:10.1089/ars.2008.2253

16.   Mikami Y, Shibuya N, Kimura Y, Nagahara N, Ogasawara Y, Kimura H. Thioredoxin and

dihydrolipoic acid are required for 3-mercaptopyruvate sulfurtransferase to produce

hydrogen sulfide. *Biochem J*. 2011;439(3):479-85. doi:10.1042/BJ20110841

17.   Yadav PK, Yamada K, Chiku T, Koutmos M, Banerjee R. Structure and kinetic analysis

of H2S production by human mercaptopyruvate sulfurtransferase. *J Biol Chem*.

2013;288(27):20002-13. doi:10.1074/jbc.M113.466177

18.   Wedmann R, et al. Improved tag-switch method reveals that thioredoxin acts as

509   depersulfidase and controls the intracellular levels of protein persulfidation. *Chem*

510   *Sci*. 2016;7(5):3414-3426. doi:10.1039/c5sc04818d

511   19.   Mani S, Cao W, Wu L, Wang R. Hydrogen sulfide and the liver. *Nitric Oxide - Biol*

512   *Chem*. 2014;312(3):H406-H414. doi:10.1016/j.niox.2014.02.006

513   20.   Pichette J, Gagnon J. Implications of Hydrogen Sulfide in Glucose Regulation: How

514   H2S Can Alter Glucose Homeostasis through Metabolic Hormones. *Oxid Med Cell*

515   *Longev*. 2016;2016:3285074. doi:10.1155/2016/3285074

516   21.   Zhang L, Yang G, Untereiner A, Ju Y, Wu L, Wang R. Hydrogen sulfide impairs glucose

517   utilization and increases gluconeogenesis in hepatocytes. *Endocrinology*.

518   2013;154(1):114-126. doi:10.1210/en.2012-1658

519   22.   Wu D, et al. Exogenous hydrogen sulfide mitigates the fatty liver in obese mice

520   through improving lipid metabolism and antioxidant potential. *Med Gas Res*.

521   2015;5(1):1-8. doi:10.1186/s13618-014-0022-y

522   23.   Mani S, et al. Decreased endogenous production of hydrogen sulfide accelerates

523   atherosclerosis. *Circulation*. 2013;127(25):2523-2534.

524   doi:10.1161/CIRCULATIONAHA.113.002208

525   24.   Distrutti E, et al. The methionine connection: Homocysteine and hydrogen sulfide

526   exert opposite effects on hepatic microcirculation in rats. *Hepatology*.

527   2008;47(2):659-667. doi:10.1002/hep.22037

528   25.   Fiorucci S, et al. The third gas: H2S regulates perfusion pressure in both the isolated

529   and perfused normal rat liver and in cirrhosis. *Hepatology*. 2005;42(3):539-548.

530   doi:10.1002/hep.20817

26.   Sestito S, Nesi G, Pi R, Macchia M, Rapposelli S. Hydrogen Sulfide: A Worthwhile Tool

in the Design of New Multitarget Drugs. *Front Chem*. 2017;5:72.

doi:10.3389/fchem.2017.00072

27. Whiteman M, Le Trionnaire S, Chopra M, Fox B, Whatmore J. Emerging role of

hydrogen sulfide in health and disease: critical appraisal of biomarkers and

pharmacological tools. *Clin Sci*. 2011;121(11):459-488. doi:10.1042/cs20110267

28. Libiad M, et al. Thiosulfate Sulfurtransferase like Domain Containing 1 Protein

Interacts with Thioredoxin. *J Biol Chem*. January 2018;293(8):2675-2686.

doi:10.1074/jbc.RA117.000826

29. Reiffenstein R. Toxicology Of Hydrogen Sulfide. *Annu Rev Pharmacol Toxicol*.

1992;32(1):109-134. doi:10.1146/annurev.pharmtox.32.1.109

30. Tiranti V, et al. Loss of ETHE1, a mitochondrial dioxygenase, causes fatal sulfide

toxicity in ethylmalonic encephalopathy. *Nat Med*. 2009;15(2):200-205.

doi:10.1038/nm.1907

31. Hildebrandt TM, Grieshaber MK. Three enzymatic activities catalyze the oxidation of

sulfide to thiosulfate in mammalian and invertebrate mitochondria. *FEBS J*.

2008;275(13):3352-3361. doi:10.1111/j.1742-4658.2008.06482.x

32. Norris EJ, Culberson CR, Narasimhan S, Clemens MG. The liver as a central regulator

of hydrogen sulfide. *Shock*. 2011;36(3):242-50. doi:10.1097/SHK.0b013e3182252ee7

33. Jackson MR, Melideo SL, Jorns MS. Human sulfide:Quinone oxidoreductase catalyzes

the first step in hydrogen sulfide metabolism and produces a sulfane sulfur

metabolite. *Biochemistry*. 2012;51(34):6804-15. doi:10.1021/bi300778t

34. Libiad M, Yadav PK, Vitvitsky V, Martinov M, Banerjee R. Organization of the human

mitochondrial hydrogen sulfide oxidation pathway. *J Biol Chem*.

2014;289(45):30901-10. doi:10.1074/jbc.M114.602664

35.   Morton NM, et al. Genetic identification of thiosulfate sulfurtransferase as an

adipocyte-expressed antidiabetic target in mice selected for leanness. *Nat Med*.

2016;22(7):771-9. doi:10.1038/nm.4115

36.   Gero D, et al. The novel mitochondria-targeted hydrogen sulfide (H2S) donors AP123

and AP39 protect against hyperglycemic injury in microvascular endothelial cells in

vitro. *Pharmacol Res*. 2016;113:186-198. doi: 10.1016/j.phrs.2016.08.019

37.   Karwi QG, Bice JS, Baxter GF. Pre- and postconditioning the heart with hydrogen

sulfide (H2S) against ischemia/reperfusion injury in vivo: a systematic review and

meta-analysis. *Basic Res Cardiol*. 2018;113(1):6. doi:10.1007/s00395-017-0664-8

38.   Mason TM. The role of factors that regulate the synthesis and secretion of very-low-

density lipoprotein by hepatocytes. *Crit Rev Clin Lab Sci*. 1998;35(6):461-487.

doi:10.1080/10408369891234246

39.   Vitvitsky V, Yadav PK, Kurthen A, Banerjee R. Sulfide oxidation by a noncanonical

pathway in red blood cells generates thiosulfate and polysulfides. *J Biol Chem*.

2015;290(13):8310-8320. doi:10.1074/jbc.M115.639831

40.   Vitvitsky V, Yadav PK, An S, Seravalli J, Cho US, Banerjee R. Structural and mechanistic

insights into hemoglobincatalyzed hydrogen sulfide oxidation and the fate of

polysulfide products. *J Biol Chem*. 2017;292(13):5584-5592.

doi:10.1074/jbc.M117.774943

41.   Banerjee R, Chiku T, Kabil O, Libiad M, Motl N, Yadav PK. *Assay Methods for H2S*

Biogenesis and Catabolism Enzymes. Vol 554. Academic Press Inc.; 2015:189-200.

doi:10.1016/bs.mie.2014.11.016

42.   Singha S, et al. Toward a selective, sensitive, fast-responsive, and biocompatible two-

photon probe for hydrogen sulfide in live cells. *Anal Chem*. 2015;87(2):1188-1195.

doi:10.1021/ac503806w

43.   Arndt S, et al. Assessment of H2S in vivo using the newly developed mitochondria-

targeted mass spectrometry probe MitoA. *J Biol Chem*. 2017;292(19):7761-7773.

doi:10.1074/jbc.M117.784678

44.   Krishnan N, Fu C, Pappin DJ, Tonks NK. H2S-Induced Sulfhydration of the Phosphatase

PTP1B and Its Role in the Endoplasmic Reticulum Stress Response. *Sci Signal*.

2011;4:ra86. doi:10.1126/scisignal.2002329

45.   Kabil O, Motl N, Banerjee R. H2S and its role in redox signaling. *Biochim Biophys Acta -*

Proteins Proteomics. 2014;1844(8):1355-1366. doi:10.1016/j.bbapap.2014.01.002

46.   Koike S, Nishimoto S, Ogasawara Y. Cysteine persulfides and polysulfides produced by

exchange reactions with H2S protect SH-SY5Y cells from methylglyoxal-induced

toxicity through Nrf2 activation. *Redox Biol*. 2017;12:530-539.

doi:10.1016/j.redox.2017.03.020

47.   Nagahara N. Catalytic site cysteines of thiol enzyme: eulfurtransferases. *J Amino*

Acids. 2011;2011:709404. doi:10.4061/2011/709404

48.   Xie L, et al. Hydrogen sulfide induces Keap1 S-sulfhydration and suppresses diabetes-

accelerated atherosclerosis via Nrf2 activation. *Diabetes*. 2016;65(10):3171-3184.

doi:10.2337/db16-0020

49.   Yang G, et al. Hydrogen sulfide protects against cellular senescence via s-

sulfhydration of keap1 and activation of Nrf2. *Antioxidants Redox Signal*.

2013;18(15):1906-1919. doi:10.1089/ars.2012.4645

50.   Módis K, Coletta C, Erdélyi K, Papapetropoulos A, Szabo C. Intramitochondrial

hydrogen sulfide production by 3-mercaptopyruvate sulfurtransferase maintains

mitochondrial electron flow and supports cellular bioenergetics. *FASEB J*.

2013;27(2):601-11. doi:10.1096/fj.12-216507

51. Szabo C, et al. Regulation of mitochondrial bioenergetic function by hydrogen sulfide.

Part I. Biochemical and physiological mechanisms. *Br J Pharmacol*.

2014;171(8):2099-122. doi:10.1111/bph.12369

52. Nagahara N. Multiple role of 3-mercaptopyruvate sulfurtransferase: antioxidative

function, H2S and polysulfide production and possible SOx production. *Br J*

Pharmacol. 2018;175(4):577-589. doi:10.1111/bph.14100

53. Kimura Y, Koike S, Shibuya N, Lefer D, Ogasawara Y, Kimura H. 3-Mercaptopyruvate

sulfurtransferase produces potential redox regulators cysteine- and glutathione-

persulfide (Cys-SSH and GSSH) together with signaling molecules H2S2, H2S3 and H2S.

Sci Rep. 2017;7(1):1-14. doi:10.1038/s41598-017-11004-7

54. Nagahara N, Tanaka M, Tanaka Y, Ito T. Novel Characterization of Antioxidant

Enzyme, 3-Mercaptopyruvate Sulfurtransferase-Knockout Mice: Overexpression of

the Evolutionarily-Related Enzyme Rhodanese. *Antioxidants*. 2019;8(5):116.

doi:10.3390/antiox8050116

55. Rui L. Energy metabolism in the liver. *Compr Physiol*. 2014;4(1):177-197.

doi:10.1002/cphy.c130024

56. Qian K, Zhong S, Xie K, Yu D, Yang R, Gong DW. Hepatic ALT isoenzymes are elevated

in gluconeogenic conditions including diabetes and suppressed by insulin at the

protein level. *Diabetes Metab Res Rev*. 2015;31(6):562-571. doi:10.1002/dmrr.2655

57. Sookoian S, et al. Serum aminotransferases in nonalcoholic fatty liver disease are a

signature of liver metabolic perturbations at the amino acid and Krebs cycle level. *Am*

J Clin Nutr. 2016;103(2):422-434. doi:10.3945/ajcn.115.118695

58. Ghosh Dastidar S, et al. Glutathione S -transferase P deficiency induces glucose

intolerance via JNK-dependent enhancement of hepatic gluconeogenesis. *Am J*

Physiol Metab. 2018;315(5):E1005-E1018. doi:10.1152/ajpendo.00345.2017

59.   Slocum SL, et al. Keap1/Nrf2 pathway activation leads to a repressed hepatic

gluconeogenic and lipogenic program in mice on a high-fat diet. *Arch Biochem*

Biophys. 2016;591:57-65. doi:10.1016/j.abb.2015.11.040

60.   Meakin PJ, et al. Susceptibility of Nrf2-Null Mice to Steatohepatitis and Cirrhosis upon

Consumption of a High-Fat Diet Is Associated with Oxidative Stress, Perturbation of

the Unfolded Protein Response, and Disturbance in the Expression of Metabolic

Enzymes but Not with Insulin Resistance. *Mol Cell Biol*. 2014;34(17):3305-3320.

doi:10.1128/mcb.00677-14

61.   Zhang YKJ, Wu KC, Liu J, Klaassen CD. Nrf2 deficiency improves glucose tolerance in

mice fed a high-fat diet. *Toxicol Appl Pharmacol*. 2012;264(3):305-314.

doi:10.1016/j.taap.2012.09.014

62.   Tolwani RJ, et al. Medium-chain acyl-CoA dehydrogenase deficiency in gene-targeted

mice. *PLoS Genet*. 2005;1(2):0205-0212. doi:10.1371/journal.pgen.0010023

63.   Onkenhout W, Venizelos V, Van Der Poet PFH, Van Den Heuvel MPM, Poorthuis BJ.

Identification and quantification of intermediates of unsaturated fatty acid

metabolism in plasma of patients with fatty acid oxidation disorders. *Clin Chem*.

1995;41(10):1467-74. PMID: 7586519

64.   Hildebrandt TM, Di Meo I, Zeviani M, Viscomi C, Braun H. Proteome adaptations in

Ethe1-deficient mice indicate a role in lipid catabolism and cytoskeleton organization

via post-translational protein modifications. *Biosci Rep*. 2013;33(4):575-584.

doi:10.1042/bsr20130051

65.   Bonomi F, Pagani S, Cerletti P, Cannella C. Rhodanese-Mediated Sulfur Transfer to

Succinate Dehydrogenase. *Eur J Biochem*. 1977;72(1):17-24.

doi:10.1111/j.1432-1033.1977.tb11219.x

66.   Pagani S, Galante YM. Interaction of rhodanese with mitochondrial NADH

dehydrogenase. *Biochim Biophys Acta*. 1983;742(2):278-84.

doi:10.1016/0167-4838(83)90312-6

67.   Nandi DL, Westley J. Reduced thioredoxin as a sulfur-acceptor substrate for

rhodanese. *Int J Biochem Cell Biol*. 1998;30(9):973-7.

doi:10.1016/S1357-2725(98)00050-8

68.   Nandi DL, Horowitz PM, Westley J. Rhodanese as a thioredoxin oxidase. *Int J Biochem*

Cell Biol. 2000;32(4):465-473. doi:10.1016/S1357-2725(99)00035-7

69.   Smirnov A, et al. Mitochondrial enzyme rhodanese is essential for 5S ribosomal RNA

import into human mitochondria. *J Biol Chem*. 2010;285(40):30792-30803.

doi:10.1074/jbc.M110.151183

70.   Uyeda K, Repa JJ. Carbohydrate response element binding protein, ChREBP, a

transcription factor coupling hepatic glucose utilization and lipid synthesis. *Cell*

Metab. 2006;4(2):107-110. doi:10.1016/j.cmet.2006.06.008

71.   Kim MS, et al. ChREBP regulates fructose-induced glucose production independently

of insulin signaling. *J Clin Invest*. 2016;126(11):4372-4386. doi:10.1172/JCI81993

72.   Xue R, et al. Hydrogen Sulfide Treatment Promotes Glucose Uptake by Increasing

Insulin Receptor Sensitivity and Ameliorates Kidney Lesions in Type 2 Diabetes.

Antioxid Redox Signal. 2013;19(1):5-23. doi:10.1089/ars.2012.5024

73.   Feng X, Chen Y, Zhao J, Tang C, Jiang Z, Geng B. Hydrogen sulfide from adipose tissue

is a novel insulin resistance regulator. *Biochem Biophys Res Commun*.

2009;380(1):153-9. doi:10.1016/j.bbrc.2009.01.059

74.   Geng B, et al. Increase or Decrease Hydrogen Sulfide Exert Opposite Lipolysis, but

677   Reduce Global Insulin Resistance in High Fatty Diet Induced Obese Mice. *PLoS One*.

678   2013;8(9):e73892. doi:10.1371/journal.pone.0073892

679   75.   Lutchmansingh FK, et al. Glutathione metabolism in type 2 diabetes and its

680   relationship with microvascular complications and glycemia. *PLoS One*.

681   2018;13(6):e0198626. doi:10.1371/journal.pone.0198626

682   76.   Jain SK, Micinski D, Huning L, Kahlon G, Bass PF, Levine SN. Vitamin D and L-cysteine

683   levels correlate positively with GSH and negatively with insulin resistance levels in the

684   blood of type 2 diabetic patients. *Eur J Clin Nutr*. 2014;68(10):1148-1153.

685   doi:10.1038/ejcn.2014.114

686   77.   Zhang L, et al. Hydrogen sulfide (H2S)-releasing compounds: Therapeutic potential in

687   cardiovascular diseases. *Front Pharmacol*. 2018;9:1066.

688   doi:10.3389/fphar.2018.01066

689   78.   Szabó G, et al. Cardioprotective effects of hydrogen sulfide. *Nitric Oxide.*

690   2011;25(2):201-210. doi:10.1016/j.niox.2010.11.001

691   79.   Miller RA, Buehner G, Chang Y, Harper JM, Sigler R, Smith-Wheelock M. Methionine-

692   deficient diet extends mouse lifespan, slows immune and lens aging, alters glucose,

693   T4, IGF-I and insulin levels, and increases hepatocyte MIF levels and stress resistance.

694   *Aging Cell*. 2005;4(3):119-25. doi:10.1111/j.1474-9726.2005.00152.x

695   80.   Lee BC, Kaya A, Gladyshev VN. Methionine restriction and life-span control. *Ann N Y*

696   *Acad Sci*. 2016;1363:116-24. doi:10.1111/nyas.12973

697   81.   Shimokawa I, et al. The life-extending effect of dietary restriction requires Foxo3 in

698   mice. *Aging Cell*. 2015;14(4):707-9 doi:10.1111/acel.12340

82.   Hine C, et al. Hypothalamic-pituitary axis regulates hydrogen sulfide production. *Cell*

Metab. 2017;25(6):1320-1333 doi:10.1016/j.cmet.2017.05.003

83.   Hine C, Zhu Y, Hollenberg AN, Mitchell JR. Dietary and endocrine regulation of

endogenous hydrogen sulfide production: implications for longevity. *Antioxid Redox*

Signal. 2018;28(16):1483-1502. doi:10.1089/ars.2017.7434

84.   Libiad M, Sriraman A, Banerjee R. Polymorphic variants of human rhodanese exhibit

differences in thermal stability and sulfur transfer kinetics. *J Biol Chem*.

2015;290(39):23579-88 doi:10.1074/jbc.M115.675694

85.   Billaut-Laden I, et al. Evidence for a functional genetic polymorphism of the human

thiosulfate sulfurtransferase (Rhodanese), a cyanide and H2S detoxification enzyme.

Toxicology. 2006;225(1):1-11. doi:10.1016/j.tox.2006.04.054

86.   Morris AA, et al. Thiocyanate Accumulation in Critically Ill Patients Receiving

Nitroprusside Infusions. *J Intensive Care Med*. 2017;32(9):547-553.

doi:10.1177/0885066616657004

87.   Bromley J, Hughes BGM, Leong DCS, Buckley NA. Life-threatening interaction

between complementary medicines: Cyanide toxicity following ingestion of

amygdalin and vitamin C. *Ann Pharmacother*. 2005;39(9):1566-1569.

doi:10.1345/aph.1E634

88.   O'Brien B, Quigg C, Leong T. Severe Cyanide Toxicity from "Vitamin Supplements." *Eur*

J Emerg Med. 2005;12(5):257-8. doi:10.1097/00063110-200510000-00014

89.   Curry SC, Carlton MW, Raschke RA. Prevention of fetal and maternal cyanide toxicity

from nitroprusside with coinfusion of sodium thiosulfate in gravid ewes. *Anesth*

Analg. 1997;84(5):1121-1126. doi:10.1213/00000539-199705000-00031

90.   Petrova F. Hydrogen cyanide and cyanides : human health aspects. 2004;CICAD;61

https://www.who.int/ipcs/publications/cicad/en/cicad61.pdf?ua=1

91. Vinnakota C V, et al. Comparison of cyanide exposure markers in the biofluids of

smokers and non-smokers. *Biomarkers*. 2012;17(7):625-633.

doi:10.3109/1354750X.2012.709880

92. Kashala-Abotnes E, Okitundu D, Mumba D, Boivin MJ, Tylleskär T, Tshala-Katumbay D.

Konzo: a distinct neurological disease associated with food (cassava) cyanogenic

poisoning. *Brain Res Bull*. 2019;145:87-91. doi:10.1016/j.brainresbull.2018.07.001

93. Swerdlow DI, et al. HMG-coenzyme A reductase inhibition, type 2 diabetes, and

bodyweight: Evidence from genetic analysis and randomised trials. *Lancet*.

2015;385(9965):351-361. doi:10.1016/S0140-6736(14)61183-1

Figures and Legends

Figure 1. *Tst* deletion results in impaired glucose and lipid metabolism. (A) Plasma glucose over 120
minutes, following pyruvate (i.p., 1.5mg/g) administration in overnight fasted C57Bl/6J (black line, n =
9) and $Tst^{-/-}$ (red line, n = 8) normal diet-fed mice. **(B)** Extinction of NADH measured by absorbance
at 340nm, coupled to PEPCK activity from liver homogenates taken from C57Bl/6J (white bar, n = 6)
and $Tst^{-/-}$ (red bar, n = 6) normal diet-fed mice. **(C)** Production of ^{13}C (M+2) acetyl-CoA and ^{13}C (M+3)
aspartate generated from 3-carbon labelled ^{13}C (M+3) pyruvate, expressed as a percentage of the total
amount of detected metabolite, in primary hepatocytes from C57Bl/6J (white bars, n = 6) and $Tst^{-/-}$
(red bars, n = 5) normal diet-fed mice. **(D)** HPLC quantified total and VLDL plasma triglyceride in 4 hour
fasted C57Bl/6J (white bar, n = 6), and $Tst^{-/-}$ (red bar, n = 6) normal diet-fed mice. **(E)** Representative
light microscopic images of fixed liver stained with Oil-Red O from normal diet-fed (ND) or high fat
diet-fed (HFD) C57Bl/6J and $Tst^{-/-}$ mice. **(F)** Analysis of the area of red staining (Oil Red O) after
thresholding, using Image J, from normal diet-fed (no pattern, n = 3-4/genotype) or high fat diet-fed
(hatched pattern, n = 4-5/genotype) C57Bl/6J (white bars) and $Tst^{-/-}$ (red bars) mice. **(G)** Plasma
glucose over 120 minutes following insulin (i.p., 1mU/g) administration in 4 hour fasted C57Bl/6J
(black line, n = 8) and $Tst^{-/-}$(red line, n = 7) normal diet-fed mice. Inset; Reduction in plasma glucose
from baseline levels 60 minutes after the injection of insulin. **(H)** Blood glucose during the pre-clamp
phase of the hyperinsulinemic, euglycaemic clamp from C57Bl/6J (white, n = 3), and $Tst^{-/-}$ (red, n =
6) normal diet-fed mice. Significance is reported from area under curve analysis. **(I)** Mean integrated
radioactive glucose (inversely related to whole body glucose uptake) during a hyperinsulinemic,
euglycaemic clamp from normal diet fed C57Bl/6J control (white, n = 3), and $Tst^{-/-}$ (red, n = 6) mice.
Data are represented as mean ±SEM. Significance was calculated using repeated measures ANOVA
(A,G) 2-WAY ANOVA **(C,F)** or unpaired two-tailed student's t-test **(B,D,H,I)** * P < 0.05, ** P < 0.01. For
(C) the 2-WAY ANOVA revealed no overall effect of genotype, however a significant interaction
between metabolite and genotype was found. The * on the histogram represents this interaction. For
(F) no main genotype effect was found, but a significant effect of diet, and an interaction were found.
Post Hoc analysis using Sidaks' multiple comparison test show an effect of diet on the 6J controls (***),
whereas no effect of diet is found on the $Tst^{-/-}$.

Figure 2. *Tst* deletion results in increased hepatic sulfur excretion and a reduction of protein persulfidation. (**A**) Cysteine concentrations (MBB-HPLC) in the media incubated with primary hepatocytes in the presence (hatched pattern) or absence (no pattern) of 1mM methionine, from C57Bl/6J (white bars, n = 4/treatment) and *Tst*$^{-/-}$ (red bars, n = 4/treatment) mice. (**B**) Glutathione concentrations (MBB-HPLC) in the media incubated with primary hepatocytes in the presence (hatched pattern) or absence (no pattern) of 1mM methionine, from C57Bl/6J (white bars, n = 4/treatment) and *Tst*$^{-/-}$ (red bars, n = 4/treatment) mice. (**C**) Pie chart depicting the proportion of liver peptides that are significantly higher (82 peptides, purple space) or lower (311 peptides, yellow space), in their persulfidation rate in the *Tst*$^{-/-}$ (n = 3) relative to C57Bl/6J (n = 3) mice. (**D**) Total DTT-released cysteine-persulfidated liver protein as measured by REVERT total protein stain following western blotting, normalised to the total input protein of the sample from *Tst*$^{-/-}$ (red bar, n = 4) and C57Bl/6J (white bar, n = 4) mice. Data with error bars are represented as mean ±SEM. Significance was calculated using 2-WAY ANOVA (**A**, **B**) or student's t-test (**D**), * P < 0.05, ** P < 0.01. For (**A**) and (**B**) the 2-WAY ANOVA reveals a main effect of genotype, indicated by * or ** on the histogram. A significant effect of methionine was also found for both (**A**) and (**B**) not indicated on the histogram. For (**C**) peptides were selected as being significant at a P-diff of 0.95 or greater.

Number of down-regulated proteins   Number of up-regulated proteins

Figure 3. *Tst* deletion engenders a high fat feeding–like hepatic proteome with a distinct organellar signature. (**A**) Venn diagram representing the number of proteins significantly different (at P < 0.01) between normal diet-fed *Tst*$^{-/-}$ and C57Bl/6J (Red circle), and the number of regulated proteins between high fat fed and normal diet-fed C57Bl/6J (Green circle). The overlap (brown) represents those proteins regulated in the same direction by both comparisons (n= 4/genotype). (**B**) Number of proteins significantly different (at p < 0.01) between 58% high fat and normal diet in either C57Bl/6J (white bar), or *Tst*$^{-/-}$ mice (Red bar), (n= 4/genotype). (**C**) Pie charts depicting the proportion of individual liver proteins that are upregulated (Blue space) compared to downregulated (yellow space) after GO term categorisation according to subcellular location. Upper row; normal diet-fed *Tst*$^{-/-}$ relative to normal diet-fed C57Bl/6J. Lower row, high fat-fed C57Bl/6J relative to normal diet-fed C57Bl/6J.

Figure 4. *Tst* deletion results in increased hepatocyte respiration but impaired medium–chain fat respiration. (**A**) Electron microscope images of liver, visualising mitochondria from normal diet-fed C57Bl/6J (n = 4) or *Tst*$^{-/-}$ (n = 4) mice. (**B**) Seahorse trace representing the mean oxygen consumption rate (OCR), normalised to protein, by hepatocytes from normal diet-fed C57Bl/6J (n = 6) or *Tst*$^{-/-}$ (n = 6) mice during a mitochondrial stress test. (**C**) Respiratory OCR linked to ATP production (oligomycin sensitive) by hepatocytes from normal diet-fed C57Bl/6J (n = 6) or *Tst*$^{-/-}$ (n = 6) mice, calculated from Figure 4B. (**D**) Respiratory OCR relating to proton leak (oligomycin insensitive) by hepatocytes from normal diet-fed C57Bl/6J (n = 6) or *Tst*$^{-/-}$ (n = 6) mice, calculated from Figure 4B. (**E**) Reduction of maximal uncoupled respiration following inhibition of long chain fatty acid mitochondrial import using etomoxir (8 µM), from normal diet-fed C57Bl/6J (n = 4) or *Tst*$^{-/-}$ (n = 4) mice. (**F**) Stimulation of maximal uncoupled respiration following addition of medium chain fatty acid octanoate (250 µM), from normal diet-fed C57Bl/6J (n = 4) or *Tst*$^{-/-}$ (n = 4) mice. Data are represented as mean ±SEM. Significance was calculated using an unpaired two tailed, student's t-test (**C, D, E, F**), * P < 0.05.

Tables

Table 1. Sulfur species in blood, urine, tissue and cells

	C57Bl/6J	$Tst^{-/-}$	$Tst^{-/-}$/6J ratio	Significance
A Trunk Blood (Micromolar)				
MBB-S (Sulfide)	2.28 +/- 0.43	22.18 +/- 0.85	9.73	****
MBB-SSO3 (Thiosulfate)	n.d.	6.25 +/- 3.17	n.c.	ns
B Trunk Plasma (Micromolar)				
MBB-S (Sulfide)	1.88 +/- 0.64	24.50 +/- 2.02	13.03	****
MBB-SSO3 (Thiosulfate)	3.99 +/- 0.99	80.29 +/- 13.6	20.12	**
MBB-GSH (reduced Glutathione)	48.0 +/- 1.15	86.25 +/- 6.27	1.80	***
C Urine (Micromoles/creatine/24hrs)				
MBB-SSO3 (Thiosulfate)	4.99 +/- 2.6	2374 +/- 319	475.75	****
D Inferior Vena Cava (Micromolar)				
MBB-S (Sulfide)	1.22 +/- 0.20	3.58 +/- 0.87	2.93	ns (0.08)
MBB-SSO3 (Thiosulfate)	6.58 +/- 4.51	88.3 +/- 13.0	13.42	*
E Liver (µmoles/kg wet liver)				
MBB-S (Sulfide)	13 +/- 1	17 +/- 3	1.31	ns
MBB-SSO3 (Thiosulfate)	4 +/- 1	15 +/- 7	3.75	ns
DNFB-GSH (reduced Glutathione)	6470 +/- 380	6850 +/- 30	1.04	ns
DNFB-Cysteine (Cysteine)	82 +/- 13	67 +/- 11	0.82	ns
F Sulfide P3 fluorescence (A510nm/protein)				
Hepatocyte	7.22 +/- 1.00	7.89 +/- 0.80	1.09	ns
G Mitochondrial sulfide (MitoA)				
Liver	0.78 +/- 0.16	1.14 +/- 0.45	1.46	ns

* P < 0.05, ** P < 0.01, *** P < 0.001 **** P < 0.0001

Table 1. *Tst* Deletion Results in Altered Sulfur Metabolites in Blood and Liver. (**A**) Sulfide dibimane, thiosulfate-MBB, measured by fluorescence detection following HPLC, from whole blood taken from trunk blood of ND-fed C57Bl/6J (n = 4) and $Tst^{-/-}$ (n = 4) mice. (**B**) Sulfide dibimane, thiosulfate-MBB, and rGSH-MBB, measured by fluorescence detection following HPLC, from EDTA-Plasma of ND-fed C57Bl/6J (n = 4) and $Tst^{-/-}$ (n = 4) mice. (**C**) Thiosulfate-MBB corrected for creatinine from 24 hour urine samples, taken from ND-fed C57Bl6/J (n = 4) and $Tst^{-/-}$ (n = 5) mice. (**D**) Sulfide dibimane, and thiosulfate-MBB, from whole blood taken from the inferior vena cava downstream of the hepatic vein of ND-fed C57Bl/6J (n = 3) and $Tst^{-/-}$ (n = 3) mice. (**E**) Sulfide dibimane, thiosulfate-MBB, rGSH-MBB, and cysteine-MBB from whole liver (n=4/genotype) of ND-fed C57Bl/6J (n = 4) and $Tst^{-/-}$ (n = 4) mice. (**F**) Fluorescence from cultured hepatocytes following incubation with P3 (sulfide reactive probe) from ND-fed C57Bl/6J (n = 4) and $Tst^{-/-}$ (n = 4) mice. (**G**) Ratio of Mito N/MitoA from the liver of ND-fed C57Bl/6J (n = 5) and $Tst^{-/-}$ (n = 5) mice. Data are represented as mean ±SEM. Significance was calculated using unpaired two-tailed student's t-test. * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001

Table 2. GO Terms Persulfidation rate ($Tst^{-/-}$ vs C57Bl/6J liver, ND-fed)

GO-ID	Name	Direction ($Tst^{-/-}$ vs 6J)	Genes
GO terms Identified by log fold change			
0050660	FAD binding	Decreased	12
0008168	Methyltransferase activity	Decreased	9
0016741	Transferase activity, transferring one-carbon groups	Decreased	9
0008565	Protein transporter activity	Decreased	8
0008238	Exopeptidase activity	Decreased	7
0005777	Peroxisome	Decreased	7
0042579	Microbody	Decreased	7
0003995	Acyl-CoA dehydrogenase activity	Decreased	6
0008483	Transaminase activity	Decreased	6
0016769	Transferase activity, transferring nitrogenous groups	Decreased	6
0008757	S-adenosylmethionine-dependent methyltransferase activity	Decreased	6
0016655	Oxidoreductase activity, acting on NADH/NADPH, quinone	Decreased	5
0004177	Aminopeptidase activity	Decreased	5
0000059	Protein import into nucleus, docking	Decreased	3
0005643	Nuclear pore	Decreased	3
0031965	Nuclear membrane	Decreased	3
0044453	Nuclear membrane part	Decreased	3
0046930	Pore complex	Decreased	3
0015629	Actin cytoskeleton	Decreased	3
0016652	Oxidoreductase activity, NADH/NADPH, NAD/NADP acceptor	Decreased	3
0050662	Coenzyme binding	Increased	5
0016651	Oxidoreductase activity, NADH/NADPH,	Increased	5
0003954	NADH dehydrogenase activity	Increased	4
0008137	NADH dehydrogenase (ubiquinone) activity	Increased	4
0050136	NADH dehydrogenase (quinone) activity	Increased	4
0006739	NADP metabolism	Increased	3
0006769	Nicotinamide metabolism	Increased	3
0006733	Oxidoreduction coenzyme metabolism	Increased	3

905 **Table 2. Tst Deletion Results in Differential Persulfidation Rate of Liver Proteins**. Significant GO terms
906 represented by peptides with different persulfidation rates in the ND-fed $Tst^{-/-}$ liver relative to
907 C57Bl/6J. '**Direction**' indicates whether the persulfidation is decreased or increased in $Tst^{-/-}$ relative
908 to C57Bl/6J. '**Genes**' indicates the number of genes in the $Tst^{-/-}$ that represent the changes driving
909 the GO term.

Table 3. KEGG pathways ($Tst^{-/-}$ vs C57Bl/6J liver, ND-fed)

Entry	Name	Genes	Significance
A. Increased in ND $Tst^{-/-}$ liver			
00250	Alanine, Aspartate and glutamate metabolism	6	***
00260	Glycine, serine and threonine metabolism	5	**
00270	Cysteine and methionine metabolism	4	*
04122	Sulfur relay system	2	*
B. Reduced in ND $Tst^{-/-}$ liver			
00980	Metabolism of xenobiotics by cytochrome P450	12	****
00982	Drug metabolism – cytochrome P450	12	****
05204	Chemical carcinogenesis	12	****
00480	Glutathione metabolism	8	***
00040	Pentose and glucoronate interconversions	5	**
04142	Lysosome	6	**
04390	Hippo signalling pathway	4	**
00500	Starch and sucrose metabolism	5	**
05215	Prostate cancer	3	**
04024	cAMP signalling pathway	4	*
04141	Protein processing in ER	9	*
05211	Renal cell carcinoma	3	*
00830	Retinol metabolism	6	*
00053	Ascorbate and aldarate metabolism	4	*
00860	Porphyrin and chlorophyll metabolism	4	*
04722	Neurotrophin signalling pathway	3	*
04670	Leukocyte transendothelial migration	4	*
04010	MAPK signalling pathway	4	*
04720	Long-term potentiation	2	*
04914	Progesterone-mediated oocyte maturation	2	*
04062	Chemokine signalling pathway	3	*
04110	Cell cycle	3	*
04015	Rap1 signaling pathway	4	*
00983	Drug metabolism - other enzymes	5	*
04918	Thyroid hormone synthesis	3	*
04612	Antigen processing and presentation	3	*
05203	Viral Carcinogenesis	5	*
* P < 0.05, ** P < 0.01, *** P < 0.001 **** P < 0.0001			

916 **Table 3. Tst Deletion Results in Differential Hepatic Protein Abundance of KEGG Pathways.** (**A**)
917 Significant KEGG pathway terms represented by proteins that are more abundant in the ND-fed Tst^{-}
918 $^{/-}$ liver compared with ND-fed C57Bl/6J. (**B**) Significant KEGG pathway terms represented by proteins
919 that are less abundant in the ND-fed $Tst^{-/-}$ liver compared with ND-fed C57Bl/6J. '**Genes**' indicates
920 the number of genes in the $Tst^{-/-}$ that represent the changes driving the KEGG pathway.

Table 4. GO Terms Congruence of Abundance and Persulfidation

GO-ID	Name	Persulfidation ($Tst^{-/-}$ vs 6J)	Abundance ($Tst^{-/-}$ vs 6J)
0008483	Transaminase activity	Decreased	Increased
0016769	Transferase activity, transferring nitrogenous	Decreased	Increased
0003995	Acyl-CoA dehydrogenase activity	Decreased	Decreased
0005777	Peroxisome	Decreased	Decreased
0042579	Microbody	Decreased	Decreased

925 **Table 4. The $Tst^{-/-}$ liver proteome and persulfidome converges on lipid oxidation pathways.** GO
926 terms that are significantly regulated following comparison of ND-fed $Tst^{-/-}$ relative to C57Bl/6J, at *both* the
927 level of cysteine persulfidation and protein abundance.