Direct readout of heterochromatic H3K9me3 regulates DNMT1-mediated maintenance DNA methylation

Wendan Ren[1,12], Huitao Fan[2,3,12], Sara A Grimm[4], Yiran Guo[2,3], Jae Jin Kim[5], Linhui Li[1], Christopher James Petell[2,3], Xiao-Feng Tan[1], Zhi-Min Zhang[1,†], John P. Coan[6], Jiekai Yin[7], Linfeng Gao[7], Ling Cai[2,3], Brittany Detrick[1], Burak Çetin[8], Dinshaw J. Patel[9], Yinsheng Wang[7,10], Qiang Cui[11], Brian D. Strahl[2,3], Or Gozani[6], Kyle M. Miller[5], Seán E. O'Leary[1,8], Paul A. Wade[4], Gang Greg Wang[2,3,*], Jikui Song[1,7,*]

[1]Department of Biochemistry, University of California, Riverside, CA 92521, USA

[2]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27599, USA

[3]Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27599, USA

[4]Division of Intramural Research, Epigenetics and Stem Cell Biology Laboratory, National Institute of Environmental Health Sciences, Research Triangle Park, Durham, NC 27709, USA

[5]Department of Molecular Biosciences, LIVESTRONG Cancer Institute of the Dell Medical School, Institute for Cellular and Molecular Biology, University of Texas at Austin, Austin, TX 78712, USA

[6]Department of Biology, Stanford University, Stanford, CA 94305, USA

[7]Environmental Toxicology Graduate Program, University of California, Riverside, CA 92521, USA

[8]Cell, Molecular, and Developmental Biology Graduate Program, University of California, Riverside, Riverside, CA 92521.

[9]Structural Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA.

[10]Department of Chemistry, University of California, Riverside, CA 92521, USA

[11]Departments of Chemistry, Physics and Biomedical Engineering, Boston University, Boston, MA 02215, USA.

[†]Current address: School of pharmacy, Jinan University, 601 Huangpu Avenue West, Guangzhou 510632, China

[12]These authors contributed equally to this work

*Correspondence: greg_wang@med.unc.edu; jikui.song@ucr.edu

ABSTRACT (124 words)

In mammals, repressive histone modifications such as trimethylation of histone H3 Lys9 (H3K9me3), frequently coexist with DNA methylation, producing a more stable and silenced chromatin state. However, it remains elusive how these epigenetic modifications crosstalk. Here, through structural and biochemical characterizations, we identified the replication foci targeting sequence (RFTS) domain of maintenance DNA methyltransferase DNMT1, a module known to bind the ubiquitylated H3 (H3Ub), as a specific reader for H3K9me3/H3Ub, with the recognition mode distinct from the typical trimethyl-lysine reader. Disruption of the interaction between RFTS and the H3K9me3Ub affects the localization of DNMT1 in ES cells and profoundly impairs the global DNA methylation and genomic stability. Together, this study reveals a previously unappreciated pathway through which H3K9me3 directly regulates DNMT1-mediated maintenance DNA methylation.

INTRODUCTION

DNA methylation is an evolutionarily conserved epigenetic mechanism that critically influences chromatin structure and function (1,2). In mammals, DNA methylation predominantly occurs at the C-5 position of cytosine within the CpG dinucleotide context, which regulates the silencing of retrotransposons (3), allele-specific genomic imprinting (4), X-chromosome inactivation (5), and tissue-specific gene expression that underlies cell fate commitment (6). During DNA replication, DNA methylation is stably propagated by DNA methyltransferase 1 (DNMT1), which replicates the DNA methylation patterns from parental DNA strands to the newly synthesized strands in a replication-dependent manner (7,8). Faithful propagation of DNA methylation patterns is essential for clonal transmission of epigenetic landscapes between cell generations.

DNMT1 is a multi-domain protein, comprised of a large N-terminal regulatory region and a C-terminal methyltransferase (MTase) domain, linked via a conserved (GK)n dipeptide repeat (Figure 1A). The regulatory region contains a replication-foci-targeting sequence (RFTS), a CXXC zinc finger domain, and a pair of bromo-adjacent homology (BAH) domains (9-12). Recent structural and biochemical evidence has revealed that both the RFTS and CXXC domains regulate the activity of DNMT1 through autoinhibitory mechanisms: the RFTS domain directly interacts with the MTase domain to inhibit DNA binding (11-15), whereas the CXXC domain specifically recognizes unmethylated CpG nucleotides, which in turn blocks the *de novo* methylation activity of DNMT1 (9). These N-terminal domains-mediated allosteric regulations, together with an inherent enzymatic preference of the MTase domain for hemimethylated CpG sites (10), shape the enzymatic specificity of DNMT1 in maintaining DNA methylation.

The crosstalk between DNA methylation and other gene silencing mechanisms, such as histone H3K9 methylation, is widely observed across evolution, ensuring proper chromatin assembly and loci-specific gene suppression (16,17). The H3K9 methylation-dependent DNA methylation in *Neurospora* and *Arabidopsis* has been elucidated, which is attributed to a direct or indirect H3K9me2/H3K9me3 readout mechanism of DNA methyltransferases (18,19). However, the mechanism by which H3K9 methylation is translated into DNA methylation in mammals remains far from being understood. Nevertheless, it has been established that DNMT1-mediated maintenance DNA methylation is critically regulated by Ubiquitin-like, containing PHD and RING Finger domains, 1 (UHRF1) (20,21). During S phase, UHRF1 is recruited to replicating heterochromatin through its association with both hemi-methylated CpG DNA and H3K9me3 (22), where it stochastically catalyzes the mono-ubiquitylation of histone H3 at lysine 14 (H3K14Ub), lysine 18 (H3K18Ub) and/or lysine 23 (H3K23Ub) (23-26). The DNMT1 RFTS domain recognizes all these modifications, with a preference for the two-mono-ubiquitin marks (i.e. H3K18Ub/H3K23Ub), leading to allosteric stimulation of DNMT1 (26). The structure of DNMT1 RFTS domain in complex with H3K18Ub/K23Ub revealed that the H3K18Ub/K23Ub binding leads to structural rearrangement of RFTS and its dissociation with the C-terminal linker, thereby facilitating the conformation transition of DNMT1 from an autoinhibitory state to an active one (26,27). These observations suggest a UHRF1-bridged link between H3K9me3 and DNA methylation. However, whether H3K9me3 directly interacts with DNMT1 to regulate maintenance DNA methylation remains unknown.

To determine how H3K9me3 regulates DNMT1-mediated maintenance DNA methylation, we examined the histone binding activity of DNMT1 RFTS domain and its relationship to the chromatin association and enzymatic activity of DNMT1. Through ITC and *in vitro* enzymatic assay, we identified that the RFTS domain binds specifically to H3K9me3, and the combinatorial readout of H3K9me3 and H3 ubiquitylation (H3Ub) by RFTS leads to synergistic enzymatic stimulation of DNMT1. Furthermore, we determined the crystal structure of bovine RFTS domain complexed with H3K9me3 peptide and two ubiquitins, providing the molecular basis for the H3K9me3 recognition. In addition, our cellular and genomic methylation analysis demonstrated that impairment of the RFTS-H3K9me3Ub recognition led to reduced co-localization of DNMT1 with H3K9me3, a global loss of DNA methylation patterns and genome instability in ES cells. Together, this study provides a mechanism by which H3K9me3 directly impacts DNMT1-mediated maintenance DNA methylation in mammalian cells.

RESULTS

The RFTS domain of DNMT1 is an H3K9me3Ub reader module

Recent studies have indicated that UHRF1-mediated recognition of H3K9me3 and hemi-methylated CpG DNA stimulates its E3 ubiquitin ligase activity on histone H3 (25,28), thus providing a linkage between H3K9me3 and H3 ubiquitylation during UHRF1/DNMT1-mediated DNA methylation. This observation prompted us to ask whether or not H3K9me3 directly influences the interaction between the RFTS domain of DNMT1 and ubiquitylated H3. To this end, we performed Isothermal Titration Calorimetry (ITC) assays using the purified human DNMT1 RFTS domain (hDNMT1$_{RFTS}$) and histone H3 peptides (residues 1-22, H3$_{1-22}$), either unmodified or with H3K9me3 modification (H3$_{1-22}$K9me3) (Supplementary Table S1). Titration of hDNMT1$_{RFTS}$ with H3$_{1-22}$ gives a dissociation constant (K_d) of 6.4 μM (Figure 1B). In comparison, titration of hDNMT1$_{RFTS}$ with H3$_{1-22}$K9me3 gives a K_d of 1.3 μM, suggesting that hDNMT1$_{RFTS}$ has a ~5-fold binding preference for H3K9me3 over unmodified H3 tails (Figure 1C). Furthermore, we probed the interaction of hDNMT1$_{RFTS}$ with the H3$_{1-22}$ peptide acetylated at K9 (H3$_{1-22}$K9Ac) and other histone tri-methylations (H3K4me3, H3K27me3, H3K36me3 and H4K20me3) and observed a much weaker binding (Supplementary Figure S1A-S1E), supporting a specific recognition of H3K9me3 by hDNMT1$_{RFTS}$.

Next, we asked whether H3K9me3 can enhance the interaction between H3K18Ub/K23Ub and the RFTS domain in the context of full-length DNMT1. Following a previously established approach (26,29), we installed dual ubiquitin marks onto a K18C/K23C mutant form of H3$_{1-24}$ (H3$_{1-24}$Ub2) or H3$_{1-24}$K9me3 (H3$_{1-24}$K9me3Ub2) peptides through dichloroacetone (DCA) linkage (29), followed by ITC binding assay against a hDNMT1 fragment (residues 351-1600, hDNMT$_{351-1600}$) spanning from the RFTS domain to the C-terminal MTase domain (Figure 1A). We found that hDNMT$_{351-1600}$ binds to the H3$_{1-24}$K9me3Ub2 peptide with a K_d of 17 nM (Figure 1E), exhibiting a ~5-fold binding preference over the H3$_{1-24}$Ub2 peptide (K_d = 91 nM) (Figure 1D). These data support that, together with H3Ub, H3K9me3 provides a signal for tethering DNMT1 onto chromatin via its RFTS domain.

The H3K9me3 readout boosts the stimulation effect of H3Ub on DNMT1 activity

In light of a previous study showing that recognition of H3Ub by the DNMT1 RFTS
domain results in enhanced enzymatic activity of DNMT1 (26), we have further
evaluated the effect of H3K9me3 on the enzymatic stimulation of hDNMT1. Here, we
found that, consistent with the previous observation (26), incubation of hDNMT1$_{351-1600}$
with equimolar H3K18Ub and H3Ub2 led to enhanced hDNMT1$_{351-1600}$ activity by 1.3
and 14 fold, respectively (Figure 1F). In contrast, incubation with equimolar
H3K9me3/K18Ub and H3K9me3Ub2 promoted the enzymatic activity of hDNMT1$_{351-1600}$
by 2.2 and 36 fold, respectively. Additionally, although incubation of hDNMT1$_{351-1600}$ with
equimolar H3K9me0 or H3K9me3 did not lead to appreciable changes in DNMT1
activity (Figure 1F), we observed a significant increase in activity of hDNMT1$_{351-1600}$
when incubated with H3K9me3 at higher peptide concentrations, relative to H3K9me0
peptide control (Supplementary Figure S1F). Together, these results show a specific
recognition of H3K9me3 by hDNMT1$_{RFTS}$ and suggest that the combinatorial readout of
H3K9me3 and H3Ub leads to a synergistic stimulation effect on the enzymatic activity of
DNMT1.

The structure of RFTS domain in complex with H3K9me3 peptide and two-mono ubiquitin

To gain a molecular understanding of the RFTS-H3K9me3 recognition, we next
crystallized the complex of the RFTS domain from bovine DNMT1 (bDNMT1$_{RFTS}$) with
both the H3$_{1-24}$K9me3/K18C/K23C peptide and G76C-mutated ubiquitin, and solved the
structure at 3.0 Å resolution (Figure 2A and Supplementary Table S2). The structure
reveals a bDNMT1$_{RFTS}$-H3K9me3-Ub2 complex, with clearly traced H3 peptide from
Arg2 to Leu20 (Figure 2B). The bDNMT1$_{RFTS}$ domain folds into a two-lobe architecture,
with an N-lobe harboring a zinc finger and a C-lobe dominated by a helical bundle
(Figure 2A), resembling that of hDNMT1$_{RFTS}$ (12,15,26,27). Like the previous reported
hDNMT1$_{RFTS}$-H3K18Ub/K23Ub complex (26) , the H3K9me3 peptide traverses across
the surface of the C-lobe and N-lobe of bDNMT1$_{RFTS}$ domain and engages extensive
intermolecular interactions, with the side chain of H3K9me3 residue embraced by a
surface groove at the interface between bDNMT1$_{RFTS}$ and one of the ubiquitin
molecules (Figure 2B and 2C). The two ubiquitin molecules are mostly positioned at the
N-lobe of the RFTS domain, separated by the ubiquitin recognition loop (URL; residues
386-398), and interact with the RFTS domain and H3 in a fashion similar to what was
previously observed for the hDNTM1$_{RFTS}$-H3K18Ub/K23Ub complex (Figure 2E-2G)
(26). Of particular note, residue G76C of one ubiquitin and H3 K18C are positioned in
a distance that allows disulfide bond formation, while the C-terminus of the other
ubiquitin is near to the side chain of H3K14 (Figure 2D), reminiscent of the close
proximity of these moieties in the hDNTM1$_{RFTS}$-H3K18Ub/K23Ub complex. Note that in
the hDNMT1$_{RFTS}$-H3K18Ub/K23Ub complex, the H3 peptide was conjugated with two
G76C-mutated ubiquitins through disulfide linkages before mixing with hDNMT1$_{RFTS}$,
whereas in the current complex, no covalent linkage was involved between the
H3K9me3 peptide and the ubiquitins prior to the complex assembly. In this regard, the
similar positioning of H3 and ubiquitins between the two complexes, crystallized under
different conditions, reinforces the notion that H3 and ubiquitin molecules are able to
engage DNMT1$_{RFTS}$ via independent inter-molecular interactions (26). These results
also indicate that H3K9me3 and H3Ub2 may act synergistically or independently for
associating DNMT1 onto chromatin.

Strikingly, in our structure, we found that both DNMT1 and H3Ub contribute to H3K9me3 engagement. First, H3K9me3 stacks its side chain against the indole ring of bDNMT1$_{RFTS}$ W463, with the quaternary ammonium group interacting with the side chain carboxylates of bDNMT1$_{RFTS}$ E461 and E483 through electrostatic attractions (Figure 3A). In addition, residues R72-G75 of one of the bound ubiquitins make both main-chain and side-chain van der Waals contacts with H3K9me3 from an opposite direction, leading to formation of an H3K9me3-binding pocket (Figure 3A). Such an H3K9me3-binding site of DNMT1$_{RFTS}$ is distinct from the multi-walled aromatic cage typical for a trimethyl-lysine (Kme3) reader (30), but is reminiscent of the interaction between the PHD finger domain of E3 ubiquitin ligase TRIM33 and H3K9me3 (Figure 3B) (31). Through superimposition with the previously determined hDNTM1$_{RFTS}$-H3K18Ub/K23Ub structure (Figure 3C) (26), we found that those two structures overlap very well, with a root-mean-square deviation (RMSD) of 0.6 Å over 384 aligned Cα atoms. However, in comparison to the H3K9me0 binding, association of H3K9me3 with bDNMT1$_{RFTS}$ W463 leads to a slight inward shift of the indole ring (Figure 3C), resulting in an increase in buried surface area of the H3K9-engaging pocket by 8-9 Å^2.

Sequence analysis of the DNMT1 RFTS domains reveals that both the H3K9me3- and ubiquitin-binding sites are highly conserved across evolution (Figure 3D). Among these, a di-tryptophan motif (W462 and W463 in bDNMT1) is unvaried from zebrafish to human (Figure 3D), in line with their important roles in binding to H3K9me3 and surrounding H3 tail residues (Figure 2C and 3A).

Mutational analysis of DNMT1-H3K9me3Ub2 binding

To test the above structural observations, we selected hDNMT1 W464 and W465, corresponding to bDNMT1 W462 and W463, respectively, for mutagenesis. Using Electrophoretic Mobility Shift Assays (EMSA), we found that the mutation of hDNMT1 W465 into alanine (W465A) leads to a significant reduction in the H3K9me3Ub2-binding affinity of hDNMT1$_{RFTS}$ and hDNMT1$_{351-1600}$ (Figure 4A and 4B), while the W464A/W465A mutation reduced the hDNMT1$_{RFTS}$-H3K9me3Ub2 binding even further (Figure 4A) and led to nearly undetectable binding between hDNMT1$_{351-1600}$ and H3K9me3Ub2 (Figure 4B). Likewise, the ITC assays indicated that the W465A mutation reduced the H3K9me0- and H3K9me3-binding affinities of hDNMT1$_{RFTS}$ by ~5 and ~17 fold, respectively (Figure 4C and 4D); introduction of a W464A/W465A mutation largely abolished the hDNMT1$_{RFTS}$-H3K9me3 binding (Figure 4E). Furthermore, we performed *in vitro* DNA methylation assays to evaluate the effect of their mutations on the enzymatic activity of DNMT1. Unlike wild-type hDNMT1$_{351-1600}$, which shows substantial increase in methylation efficiency in the presence of H3K9me3K18Ub and H3K9me3Ub2 (Figure 1F) as well as an enzymatic preference for H3K9me3 over H3K9me0 (Supplementary Figure S1F), introduction of the W465A or W464A/W465A mutation into hDNMT1$_{351-1600}$ greatly dampened the methylation-stimulating effects of H3K9me3, H3K9me3K18Ub or H3K9me3Ub2 (Figure 4F and Supplementary Figure S2A-S2B). These data support important roles of the W464 and W465 residues in mediating the DNMT1-H3K9me3Ub interaction and the consequent enzymatic activation of DNMT1.

Given that H3Ub is reportedly a transient mark in cells (32), we have focused on examining the requirement of the DNMT1$_{RFTS}$ domain for H3K9me3 binding in cells. To this end, we used the Dnmt1-knockout mouse embryonic stem cells (1KO-ESC) (33) and generated multiple 1KO-ESC lines with comparable, stable expression of exogenous DNMT1, either wild-type (DNMT1^{WT}) or an H3K9me3-binding-defective mutant (DNMT1^{W465A} or DNMT1$^{W464A/W465A}$) (Figure 4G). By co-immunoprecipitation (CoIP), we found that both the W464A and W464A/W465A mutations interfere with efficient binding of DNMT1 to H3K9me3, but not H4K20me3, in cells (Figure 4H). Furthermore, in the ES cells synchronized at S phase, confocal immunofluorescence (IF) microscopy showed punctate nuclear foci of DNMT1^{WT} that overlap with the H3K9me3-marked, DAPI-dense regions of chromatin (Figure 4I), whereas the RFTS-mutant DNMT1 proteins show a more diffuse distribution in the nucleus and lose their co-localization with H3K9me3 (Figure 4I). Together, these data support that the histone-engaging activity of the RFTS domain is critical for both the enzymatic stimulation and chromatin occupancy of DNMT1.

The RFTS domain of DNMT1 is crucial for global DNA methylation in cells

Next, we sought to examine the role of the DNMT1 RFTS domain in maintenance DNA methylation in cells. First, we used liquid chromatography-mass spectrometry (LC-MS) to quantify global levels of methylated cytosine (5mC) in 1KO-ESC cells reconstituted with DNMT1^{WT} or the RFTS-defective mutant. As expected, stable transduction of DNMT1^{WT} led to an increase in overall 5mC level (Figure 5A, WT vs. 1KO). However, such an increase was found compromised by the W465A or W464A/W465A mutation, with the latter exhibiting a more severe DNA methylation defect, lacking significant methylation-stimulating effect relative to mock (Figure 5A). Further, we carried out genome-wide methylation profiling with enhanced reduced representation bisulfite sequencing (eRRBS). Our eRRBS data showed the desired bisulfite conversion rates (Supplementary Figure S3A, 99.79-99.83%; and Supplementary Table S3), with at least 5-fold coverage for ~5-8 million of CpG sites in all samples. Relative to 1KO-ESCs reconstituted with DNMT1^{WT}, those with DNMT1^{W465A} or DNMT1$^{W464A/W465A}$ showed a marked decrease in overall CpG methylation, with a complete loss of most heavily methylated CpG sites that are typically seen at heterochromatic repetitive elements (34) (Figure 5B, red; and Supplementary Figure S3B and S3C). In particular, there is a significantly decreased level of CpG methylation at the H3K9me3-decorated genomic regions in cells reconstituted with either DNMT1^{W465A} or DNMT1$^{W464A/W465A}$, relative to DNMT1^{WT} controls (Figure 5C), as demonstrated by sub-telomeric regions located in the chromosomes 1 and X (Figure 5D and Supplementary Figure S3D). Again, these cellular assays show that the double RFTS mutant (DNMT1$^{W464A/W465A}$) produced more severe CpG methylation defects, relative to DNMT1^{W465A} (Figure 5A-5D and Supplementary Figure S3A-S3D), which is consistent with our *in vitro* biochemical and enzymatic observations. Collectively, we have demonstrated an essential role of the histone-engaging RFTS domain in maintenance of CpG methylation at the H3K9me3-associated heterochromatin in cells.

Maintenance of proper DNA methylation levels in cells is crucial for genome stability (35). To investigate the role of DNMT1-mediated DNA methylation in genome

stabilization, we further challenged 1KO-ESC cells, reconstituted with WT or mutant DNMT1, with ionizing radiation (IR) treatment. By using the neutral comet assay (Figure 5E), a surrogate method for scoring DNA double-strand break (DSB) lesions, we found that loss of DNMT1 rendered ESC cells a hyper-sensitivity to IR treatment, reflecting a possible change in chromatin structure or impairment in DSB repair (Figure 5E and 5F), a phenotype that can be rescued by complementation with WT DNMT1. Whereas cells with the RFTS single mutant (DNMT1^{W465A}) and double mutant (DNMT1$^{W464A/W465A}$) exhibited modest and severe impairment of IR resistance (Figure 5E and 5F), respectively, confirming the role of DNMT1-mediated DNA methylation in genomic stability maintenance.

Together, the above results strongly indicated that the recognition of H3K9 trimethylation by DNMT1 is important for maintenance DNA methylation, and also genome stability and radiation resistance of ES cells.

Discussion

The crosstalk between H3K9 methylation and DNA methylation, two of the major epigenetic silencing mechanisms, critically influences gene silencing and heterochromatin formation (16,17). For instance, previous studies have demonstrated that Suv39h-mediated H3K9-trimethylation promotes the enrichment of DNA methylation at major satellite repeats of pericentromeric heterochromatin (36), which is essential for maintaining heterochromatic assembly and genome stability (37). Whereas the mechanism by which H3K9 methylation and DNA methylation crosstalk has been established in fungi and plants (18,19), how H3K9me3 is translated into DNA methylation in mammals remains elusive. Through a set of structural, biochemical and cellular analyses, this study shows that the RFTS domain of DNMT1 specifically recognizes H3K9me3, in conjunction with the previously identified H3Ub mark. The combinatorial readout of H3K9me3/H3Ub not only leads to a synergistic stimulation effect on DNMT1 activity, but also regulates the genome targeting of DNMT1. This study therefore establishes a direct link between histone H3K9me3 modification and DNMT1-mediated maintenance DNA methylation, which profoundly influences the global DNA methylation patterns and genome stability.

Previous studies from others and us demonstrated that DNMT1 assumes autoinhibitory conformations either in the DNA-free state (11,12) or in the presence of unmethylated CpG DNA (9), in which the autoinhibitory linker located between the CXXC and BAH1 domains serves as a key inhibition-enforcing element in both regulations (9,11). This study reveals that the recognition between the RFTS domain and H3K9me3 strengthens the RFTS interaction with histone H3 tails that carry either one- or two-mono-ubiquitin mark, directly contributing to the relief of the autoinhibition of DNMT1. The specific recognition of H3K9me3 by the RFTS domains of DNMT1 presumably helps transduce the H3K9me3 signal into DNA methylation, thereby ensuring the epigenetic fidelity of DNA methylation in heterochromatin domains.

The RFTS-H3K9me3 interaction reinforces the previously identified UHRF1-H3K9me3 axis to H3K9me3-directed, replication-dependent DNMT1 methylation.

UHRF1 harbors a tandem TUDOR domain that recognizes H3K9me3 (38-43) and a RING finger domain that mediates H3 ubiquitylation for DNMT1 targeting (23,24,26), thereby serving as a platform for the functional crosstalk between H3K9me3 and DNA methylation (44). However, disruption of the interaction between H3K9me3 and UHRF1 via the TUDOR domain mutation only leads to a modest (~10%) reduction of DNA methylation. In this regard, the direct readout of H3K9me3Ub by the RFTS domain of DNMT1 provides a redundant mechanism in transducing H3K9me3 into maintenance DNA methylation. Note that the RFTS-H3K9me3Ub readout does not involves the discrimination of the methylation state of DNA substrates, therefore providing a mechanism in supporting the region-specific methylation maintenance by DNMT1, as opposed to site-specific methylation maintenance (45,46). Consistently, impairment of this interaction in cells compromises the DNMT1-mediated CpG methylation, leading to an aberrant landscape of DNA methylation and defects in maintenance of genome stability. This targeting-coupled allosteric stimulation mechanism is reminiscent of the role of histone H3K4me0 in DNMT3A-mediated *de novo* DNA methylation, in which the specific recognition of the DNMT3A ADD domain with H3K4me0 allosterically stimulates its enzymatic activity, thereby providing a mechanism of locus-specific DNA methylation establishment (47).

The DNMT1 RFTS domain adds to the reader modules that offer interpretation of specific histone modifications (30), but deviates from the typical Kme3 readout that depends on aromatic or other hydrophobic residues (30): the RFTS domain presents a single tryptophan to stack against H3K9me3, unlike the typical Kme3 readout involving a hydrophobic cage composed of multiple aromatic residues (30). These observations highlight the evolutionary divergence of the histone modification-binding mechanisms.

Materials and Methods

Plasmids

The plasmid that contains DNMT1 was purchased from Addgene (cat # 24952). The DNMT1 cDNA was fused to an N-terminal 3xFlag tag by PCR, followed by subcloning into the pPyCAGIP vector (48) (kind gift of I. Chambers). DNMT1 point mutation was generated by a QuikChange II XL Site-Directed Mutagenesis Kit (Agilent), with the residue numerations based on the isoform 1 of DNMT1 that contains 1616 amino acids. For domain analysis of the DNMT1 RFTS-H3K9me3 interactions, DNA encoding the human DNMT1 RFTS domain (residues 351-597, $hDNMT1_{RFTS}$) or the bovine DNMT1 RFTS domain (residues 349-594, $bDNMT1_{RFTS}$) was cloned into a modified pRSF-Duet vector preceded by an N-terminal His_6-SUMO tag and ULP1 (ubiquitin-like protease 1) cleavage site. For analysis of $hDNMT1_{351-1600}$ methylation activity, the hDNMT1 construct was inserted into an in-house expression vector as a His_6-MBP-tagged form. All plasmid sequences were verified by sequencing before use.

Protein purification

The plasmids were transformed into BL21(DE3) RIL cells (Novagen Inc). When the cell density reached an optical density at 600 nm (OD_{600}) of 0.6, the protein expression was induced by 0.1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) at 16 °C overnight.

The cells were harvested, and subsequently resuspended and lysed in a buffer containing 50 mM Tris-HCl (pH 7.5), 25 mM imidazole, 1 M NaCl, 0.5 mM DTT and 1 mM PMSF. The His$_6$-SUMO-tagged hDNMT1$_{RFTS}$ and bDNMT1$_{RFTS}$ proteins were first purified using a nickel column with elution buffer containing 25 mM Tris-HCl (pH 8.0), 100 mM NaCl and 300 mM imidazole. The eluted protein was incubated with ULP1 on ice for cleavage of the His$_6$-SUMO tag, followed by purification of the tag-free protein by anion exchange chromatography on a HiTrap Q XL column (GE Healthcare) and nickel affinity chromatography. The protein sample was finally purified on size-exclusion chromatography on a HiLoad 16/600 Superdex 75 pg column (GE Healthcare), pre-equilibrated with buffer (20 mM Tris, pH 7.5, 50 mM NaCl, 5 mM DTT). The His$_6$-MBP tagged DNMT1 proteins were first purified by Ni^{2+} chromatography, followed by ion-exchange chromatography on a Heparin HP (GE Healthcare) or Q HP column (GE Healthcare), removal of His$_6$-MBP tag by TEV protease cleavage, a second round of nickel affinity chromatography, and size-exclusion chromatography on a Superdex 200 16/600 column (GE Healthcare). DNMT1 mutants were introduced by site-directed mutagenesis and purified as that described for wild-type protein. All purified protein samples were stored at −80 °C before use.

Chemical modifications of histones

To prepare the unbiquitylated histones, the His$_6$-SUMO-Ub(G76C) and each synthesized histone peptide (H3$_{1-25}$K18C, H3$_{1-25}$K9me3K18C, H3$_{1-24}$K18CK23C or H3$_{1-24}$K9me3K18CK23C, each containing an additional tryptophan at the C-terminus), dissolved in buffer (250 mM Tris-HCl (pH 8.6), 8 M urea, 5 mM TCEP), were mixed in a 4:1 molar ratio, and incubated at room temperature for 30 min. The crosslinker 1,3-dichloroacetone, dissolved in N,N'-dimethylformamide, was added to the reaction mixture with the amount equal to one-half of the total sulfhydryl groups. After 2 hr.-incubation on ice, the reaction was stopped by 5 mM β-ME. Purification of H3$_{1-25}$K18Ub, H3$_{1-25}$K9me3K18Ub, H3$_{1-24}$Ub2 and H3$_{1-24}$K9me3Ub2 were achieved through cation-exchange chromatography on a mono S column (GE Healthcare).

Crystallization and structure determination

To crystallize the bDNMT1$_{RFTS}$-H3K9me3-Ub2 complex, bDNMT1$_{RFTS}$ was mixed with H3$_{1-24}$K9me3K18CK23C peptide, containing an additional tryptophan at the C-terminus, and ubiquitin in a 1:1:2 molar ratio. The complex was incubated on ice for 30 min before crystallization. The crystals were generated in a buffer containing 0.1 M citric acid (pH 3.5), 28% PEG8000 at 4 °C, using the hanging-drop vapor diffusion method. The crystals were soaked in the crystallization buffer supplemented with 20-25% (v/v) glycerol as cryo-protectant before flash frozen in liquid nitrogen. The X-ray diffraction data were collected on the beamline BL5.0.1 at Advanced Light Source (ALS), Lawrence Berkeley National Laboratory. The data were indexed, integrated and scaled by HKL2000 program (49) or XDS(50). The structure was solved by molecular replacement using PHASER(51) with the RFTS domain in the DNMT1 structure (PDB 4WXX) as searching model. Iterative cycles of model rebuilding and refinement were carried out using COOT (52) and PHENIX (53), respectively. Data collection and structure refinement statistics were summarized in Supplementary Table S2.

ITC binding assay

ITC measurements were performed using a MicroCal iTC200 instrument (GE Healthcare). Synthesized $H3_{1-22}$, $H3_{1-22}K9me3$, $H3_{1-22}K9Ac$, $H3_{1-15}K4me3$, $H3_{21-33}K27me3$, $H3_{31-43}K36me3$ and $H4_{14-25}K20me3$ peptides each contain a C-terminal tyrosine for spectroscopic measurement. To measure the bindings between $hDNMT1_{RFTS}$ and the peptides, 1 mM peptide was titrated over 0.1 mM $hDNMT1_{RFTS}$ in buffer containing 20mM Tris-HCl (pH 7.5), 100 mM NaCl, 10 μM $ZnCl_2$ and 1 mM DTT at 20 °C. To measure the bindings between $hDNMT1_{351-1600}$ and H3Ub2 or H3K9me3Ub2 peptides, 0.12 mM peptide was titrated over 12 μM $hDNMT1_{351-1600}$ sample in buffer containing 20 mM Tris-HCl (pH 7.5),100 mM NaCl and 1 mM DTT at 5 °C. Analyses of all data were performed with MicroCal Origin software, fitted with single-site binding mode. The ITC parameters were summarized in Supplementary Table S1.

DNA methylation kinetics assay

The DNA methylation assays were performed as previously described (9) with modifications. Synthesized $H3_{1-25}$, $H3_{1-25}K9me3$, $H3_{1-25}K18Ub$, $H3_{1-25}K9me3K18Ub$, $H3_{1-24}Ub2$ and $H3_{1-24}K9me3Ub2$ peptides, each with a C-terminal tryptophan, were used for evaluation of the RFTS-mediated enzymatic stimulation of $hDNMT1_{351-1600}$. Each reaction mixture contains 0.1 μM $hDNMT1_{351-1600}$, wild type or mutants, 0.5 mM S-adenosyl-L-[methyl^{3}H] methionine (SAM) (Perkin Elmer), 0.4 μM $(GT^{m}C)_{12}/(GAC)_{12}$ hemimethylated DNA duplex, and various amount of histone peptides in 50 mM Tris-HCl (pH 8.0), 7 mM β-ME, 5% glycerol, 100 μg/mL BSA and 100 mM NaCl, unless indicated otherwise. The reaction mixture was incubated at 37 °C for 20 min, before quenched by 2 mM cold SAM. Eight μL of the reaction mixture was applied onto DEAE filtermat (Perkin Elmer), sequentially washed with 0.2 M ammonium bicarbonate (twice), water and ethanol. The filter paper was then air dried and soaked in ScintiVerse cocktail (Thermo fisher). The activity was measured by Beckman LS6500 scintillation counter.

Electrophoretic mobility shift assay

To measure the $H3_{1-24}K9me3Ub2$-binding affinity, 1 μM $hDNMT1_{351-1600}$ or $hDNMT1_{RFTS}$ protein was incubated with various amount of $H3_{1-24}K9me3Ub2$ peptide in 10 μL binding buffer (20 mM Tris–HCl (pH 7.5), 100 mM NaCl, 1 mM DTT and 5% glycerol) at 4 °C for 1 hr. The sample mixture was resolved in 4-10% native gel using 0.5X TG buffer at 4 °C under 100 V for 3.5 hr. The gel image was visualized by coomassie blue staining.

To measure the DNA-binding affinity of $hDNM1_{351-1600}$, 0.1 μM 26-base pair DNA duplex containing one central hemimethylated CpG site (upper strand: 5′-ACACCAAGCCTGMGGAGGCTCACGGA-3′, M = 5-methylcytosine; lower strand: 5′-TCCGTGAGCCTCCGCAGGCTTGGTGT -3′) was mixed with 0, 1, 2 or 5 ìM $hDNMT1_{351-1600}$, wild type or mutants, in the presence or absence of the $H3_{1-24}K9me3Ub2$ peptide, in buffer containing 20 mM Tris–HCl (pH 7.5), 50 mM NaCl, 1 mM DTT and 5% glycerol at 4 °C for 1 hr, before resolved in a 6% TBE native gel. The protein-DNA complex was visualized by SYBR green staining.

Cell lines and tissue culture

Dnmt1-knockout mouse embryonic stem cells (1KO-ESCs; a gift from Dr. M. Okano) were cultivated as previously described on gelatin-coated dishes in the base ESC

culture medium supplemented with leukemia inhibitory factor(54). 1KO-ESCs were transfected by Lipofectamine 2000 (Invitrogen) with the pPyCAGIP empty vector or that carrying WT or mutant DNMT1. Forty-eight hours post-transfection, the transduced ES cells were selected out in culture medium with 1μg/mL puromycin for over two weeks. The pooled stable-expression cell lines and independent single-cell-derived clonal lines were first established, followed by further characterizations such as immunoblotting of DNMT1.

Antibodies and Western blotting

Antibodies used for immunoblotting include α-Flag (Sigma; M2), H4K20me3 (Abcam ab9053), H3K9me3 (Abcam ab8898), general histone H3 (Abcam ab1791) and α-Tubulin (Sigma). Whole cell protein lysates were prepared by NP40 lysis buffer (50mM Tris-HCl pH 8.0, 150mM NaCl, 1% NP-40) followed by brief sonication and centrifugation. After mixing with loading buffer and boiling for 5 minutes, the same amount of extracted samples were loaded into SDS-PAGE gels for immunoblotting analysis as previously described (54,55).

Quantification of 5-methyl-2′-deoxycytidine (5mC) in genomic DNA

Enzymatic digestion of cellular DNA and LC-MS/MS measurements of the levels of 5-mdC in the resulting nucleoside mixture was carried out as described previously (54).

Confocal immunofluorescence (IF)

The immunofluoresence was carried out as described before (56). In brief, 3xFlag-tagged DNMT1 transduced cells were fixed with 4% paraformaldehyde and 10-min permeabilization in 0.1% Triton X-100. After one-hour block in PBS with 2.5% of BSA, cells were stained with primary antibodies, followed by staining with the Alexa-488 or Alexa-594 conjugated secondary antibodies. The primary antibodies used include M2 anti-FLAG (Sigma) and H3K9me3 (Abcam). Fluorescence was detected in a FV1000 confocal microscope (UNC Imaging Core).

Co-immunoprecipitation (CoIP)

Cells were lysed in NP40 lysis buffer that contains 50mM Tris-HCl (pH 8.0), 150mM NaCl, 1% of NP-40, 1mM DTT and a complete protease inhibitor (Roche) as described (56,57). Antibodies H4K20me3 (Abcam ab9053), H3K9me3 (Abcam ab8898) and general histone H3 (Abcam ab1791) conjugated with protein A/G beads (Millipore) or anti-Flag M2-conjugated agarose beads (Sigma) were incubated with the lysates overnight at 4 °C. The beads were then washed 3–6 times with cell lysis buffer, and the bound proteins were eluted in SDS buffer and analyzed by western blotting.

Preparation of enhanced reduced representation bisulfite sequencing (eRRBS) libraries

Construction of eRRBS libraries was performed as described before (54). Briefly, 1 microgram of total genomic DNA (gDNA) was added with 0.1% of unmethylated lambda DNA (Promega), followed by one-hour digestion with three enzymes (MspI, MseI and BfaI) at 37 degree. The purified digested gDNA was then subjected to end repair, A-tailing and ligation to NEBNext Methylated Adaptors (NEBNext DNA Library Prep Kit), followed by purification using AMPure beads. Bisulfite conversion and library

construction were carried out as before using the EpiMark Bisulfite Conversion Kit (NEB cat# E3318) according to manufacturer's specifications. The generated multiplexed eRRBS libraries were subjected to deep sequencing in an Illumina HiSeq 4000 platform with a paired end PE150 cycle (carried out by UNC HTSF Genomic Core).

eRRBS data processing

General quality control checks were performed with FastQC v0.11.2 (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). The last 5 bases were clipped from the 3' end of every read due to questionable base quality in this region, followed by filtration of the sequences to retain only those with average base quality scores of more than 20. Examination of the 5' ends of the sequenced reads indicated that 70%-90% (average 83.0%) were consistent with exact matches to the expected restriction enzyme sites (i.e. MspI, BfaI, and MseI). Approximately 85% of both ends were consistent with the expected enzyme sites. Adapter sequence was trimmed from the 3' end of reads via Cutadapt v1.2.1 (parameters -a AGATCGGAAGAG -O 5 -q 0 -f fastq; https://doi.org/10.14806/ej.17.1.200). Reads less than 30nt after adapter-trimming were discarded. Filtered and trimmed datasets were aligned via Bismark v0.18.1 (parameters -X 1000 --non_bs_mm)(58), using Bowtie v1.2(59) as the underlying alignment tool. The reference genome index contained the genome sequence of enterobacteria phage λ (NC_001416.1) in addition to the mm10 reference assembly (GRCm38). For all mapped read pairs, the first 4 bases at the 5' end of read1 and the first 2 bases at the 5' end of read2 were clipped due to positional methylation bias, as determined from QC plots generated with the 'bismark_methylation_extractor' tool (Bismark v0.18.1). To avoid bias in quantification of methylation status, any redundant mapped bases due to overlapping read ends from the same read pair were trimmed. Read pairs in which either read end had 3 more or methylated cytosines in non-CpG context were assumed to have escaped bisulfite conversion and were discarded. Finally, mapped read pairs were separated by genome (mm10 or phage λ). Read pairs mapped to phage λ were used as a QC assessment to confirm that the observed bisulfite conversion rate was >99%. Read pairs mapped to the mm10 reference genome were used for downstream analysis.

eRRBS data analysis

Although the eRRBS data does carry stranded information, data from the plus and minus strands have been collapsed in this analysis. Weighted methylation scores were calculated as described by Schultz et al(60) at query regions of interest, such as H3K9me3 and H4K20me3 peaks. Depth tracks for the eRRBS data at all genomic loci were first generated in bedGraph format by BEDTools v2.24.0(61) 'genomecov', then converted to bigWig format by bedGraphToBigWig (UCSC utility script; http://hgdownload.soe.ucsc.edu/admin/exe/). Percent methylation tracks for the eRRBS data at CpG sites in non-cytosine context were generated by first constructing WIG tracks reporting %methylation (count of methylated C bases / total C bases) per site, followed by conversion to bigWig format by wigToBigWig (UCSC utility script).

Neutral comet assay

Neutral comet assays were performed using the CometAssay Reagent Kit (Trevigen) according to the manufacturer's instruction. Cells were plated and treated with ionizing

radiation (IR; 5Gy). After 2hr post-IR treatment, cells were harvested and mixed with LMAgarose (Trevigen). The mixtures were placed on glass slides (Trevigen). Cells were lysed with lysis solution (Trevigen) at 4 °C for 1h. The slides were washed with TBE (90mM Tris borate (pH 8.3) to removal of remain lysis solution and subject to electrophoresis at 40V for 40 min in TBE buffer. Samples were fixed with 70% ethanol for 10 min. The DNA was stained with SYBR-green (Invitrogen) for 15 min in RT. The images were taken by fluorescence microscopy and tail moments were calculated for at least 100 cells for each sample by Image J (v 1.48). Tail moment (TM) reflects both the tail length (TL) and the fraction of DNA in the comet tail (TM = %DNA in tail × TL/100).

Statistics

The two-tailed Student t-tests were performed to compare distributions between different groups. And the p value lower than 0.01 was considered to be statistically significant.

Accession codes. Coordinates and structure factors for the bDNMT1$_{RFTS}$-H3K9me3-Ub2 complex have been deposited in the Protein Data Bank under accession codes xxxx. The eRRBS data have been deposited in Gene Expression Omnibus (GEO) under accession code xxxxxxx.

Acknowledgments

We would like to thank staff members at the Advanced Light Source (ALS), Lawrence Berkeley National Laboratory for access to X-ray beamlines. We are also grateful for professional support of UNC facilities including Genomics Core, which are partly supported by the UNC Cancer Center Core Support Grant P30-CA016086. This work was supported by NIH grants (1R35GM119721 to J.S, 5R21ES025392 to Y.W., RO1GM079641 to O.G., 1R35GM126900 to B.D.S., 1R01CA215284 and 1R01CA211336 to G.G.W., and CA198279 and CA201268 to K.M.M.), NIH T32 Training Fellowship for Integrated Training in Cancer Model Systems (T32CA009156) and an American Cancer Society Postdoctoral Fellowship (PF-19-027-01–DMC) to C.J.P. and grants from When Everyone Survives (WES) Leukemia Research Foundation (to G.G. W.) and Gabrielle's Angel Foundation for Cancer Research (to G.G.W.). This work was also supported in part by the Intramural Research Program of the National Institute of Environmental Health Sciences, NIH (ES101965 to PAW). J.P.C. is supported in part by T32 CA09302. G.G.W. and K.M.M. are American Cancer Society (ACS) Research Scholar, C.J.P. is an ACS postdoctoral fellow, and G.G.W. is a Leukemia & Lymphoma Society (LLS) Scholar.

Author Contributions

W.R., H.F., S.A.G., Y.G., J.J.K., L.L., C.J.P., X-F. T., Z-M. Z., J.P.C, J.Y., L.G., S.L., L.C., B.D., Y.W., Q.C., and S.O. performed experiments. Y.W., B.S., O.G., K.M.M., S.O., P.A.W., G.G.W. and J.S. organized the study. D.J.P., O.G., G.G.W. and J.S. conceived the study, G.G.W. and J.S wrote the manuscript with input from all the authors.

References

1. Law, J.A. and Jacobsen, S.E. (2010) Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nat Rev Genet*, **11**, 204-220.
2. Smith, Z.D. and Meissner, A. (2013) DNA methylation: roles in mammalian development. *Nat Rev Genet*, **14**, 204-220.
3. Walsh, C.P., Chaillet, J.R. and Bestor, T.H. (1998) Transcription of IAP endogenous retroviruses is constrained by cytosine methylation. *Nat Genet*, **20**, 116-117.
4. Li, E., Beard, C. and Jaenisch, R. (1993) Role for DNA methylation in genomic imprinting. *Nature*, **366**, 362-365.
5. Panning, B. and Jaenisch, R. (1998) RNA and the epigenetic regulation of X chromosome inactivation. *Cell*, **93**, 305-308.
6. Yagi, S., Hirabayashi, K., Sato, S., Li, W., Takahashi, Y., Hirakawa, T., Wu, G., Hattori, N., Ohgane, J., Tanaka, S. *et al.* (2008) DNA methylation profile of tissue-dependent and differentially methylated regions (T-DMRs) in mouse promoter regions demonstrating tissue-specific gene expression. *Genome research*, **18**, 1969-1978.
7. Goll, M.G. and Bestor, T.H. (2005) Eukaryotic cytosine methyltransferases. *Annu Rev Biochem*, **74**, 481-514.
8. Jeltsch, A. (2006) On the enzymatic properties of Dnmt1: specificity, processivity, mechanism of linear diffusion and allosteric regulation of the enzyme. *Epigenetics*, **1**, 63-66.
9. Song, J., Rechkoblit, O., Bestor, T.H. and Patel, D.J. (2011) Structure of DNMT1-DNA complex reveals a role for autoinhibition in maintenance DNA methylation. *Science*, **331**, 1036-1040.
10. Song, J., Teplova, M., Ishibe-Murakami, S. and Patel, D.J. (2012) Structure-based mechanistic insights into DNMT1-mediated maintenance DNA methylation. *Science*, **335**, 709-712.
11. Takeshita, K., Suetake, I., Yamashita, E., Suga, M., Narita, H., Nakagawa, A. and Tajima, S. (2011) Structural insight into maintenance methylation by mouse DNA methyltransferase 1 (Dnmt1). *Proc Natl Acad Sci U S A*, **108**, 9055-9059.
12. Zhang, Z.M., Liu, S., Lin, K., Luo, Y., Perry, J.J., Wang, Y. and Song, J. (2015) Crystal Structure of Human DNA Methyltransferase 1. *J Mol Biol*, **427**, 2520-2531.
13. Bashtrykov, P., Rajavelu, A., Hackner, B., Ragozin, S., Carell, T. and Jeltsch, A. (2014) Targeted mutagenesis results in an activation of DNA methyltransferase 1 and confirms an autoinhibitory role of its RFTS domain. *Chembiochem : a European journal of chemical biology*, **15**, 743-748.
14. Berkyurek, A.C., Suetake, I., Arita, K., Takeshita, K., Nakagawa, A., Shirakawa, M. and Tajima, S. (2014) The DNA methyltransferase Dnmt1 directly interacts with the SET and RING finger-associated (SRA) domain of the multifunctional protein Uhrf1 to facilitate accession of the catalytic center to hemi-methylated DNA. *J Biol Chem*, **289**, 379-386.
15. Syeda, F., Fagan, R.L., Wean, M., Avvakumov, G.V., Walker, J.R., Xue, S., Dhe-Paganon, S. and Brenner, C. (2011) The replication focus targeting sequence (RFTS) domain is a DNA-competitive inhibitor of Dnmt1. *J Biol Chem*, **286**, 15344-15351.
16. Cheng, X. and Blumenthal, R.M. (2010) Coordinated chromatin control: structural and functional linkage of DNA and histone methylation. *Biochemistry*, **49**, 2999-3008.
17. Du, J., Johnson, L.M., Jacobsen, S.E. and Patel, D.J. (2015) DNA methylation pathways and their crosstalk with histone methylation. *Nature reviews. Molecular cell biology*, **16**, 519-532.
18. Du, J., Zhong, X., Bernatavichute, Y.V., Stroud, H., Feng, S., Caro, E., Vashisht, A.A., Terragni, J., Chin, H.G., Tu, A. *et al.* (2012) Dual binding of chromomethylase domains to H3K9me2-containing nucleosomes directs DNA methylation in plants. *Cell*, **151**, 167-180.
19. Freitag, M., Hickey, P.C., Khlafallah, T.K., Read, N.D. and Selker, E.U. (2004) HP1 is essential for DNA methylation in neurospora. *Mol Cell*, **13**, 427-434.

20. Bostick, M., Kim, J.K., Esteve, P.O., Clark, A., Pradhan, S. and Jacobsen, S.E. (2007) UHRF1 plays a role in maintaining DNA methylation in mammalian cells. *Science*, **317**, 1760-1764.

21. Sharif, J., Muto, M., Takebayashi, S., Suetake, I., Iwamatsu, A., Endo, T.A., Shinga, J., Mizutani-Koseki, Y., Toyoda, T., Okamura, K. *et al.* (2007) The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. *Nature*, **450**, 908-912.

22. Liu, X., Gao, Q., Li, P., Zhao, Q., Zhang, J., Li, J., Koseki, H. and Wong, J. (2013) UHRF1 targets DNMT1 for DNA methylation through cooperative binding of hemi-methylated DNA and methylated H3K9. *Nature communications*, **4**, 1563.

23. Nishiyama, A., Yamaguchi, L., Sharif, J., Johmura, Y., Kawamura, T., Nakanishi, K., Shimamura, S., Arita, K., Kodama, T., Ishikawa, F. *et al.* (2013) Uhrf1-dependent H3K23 ubiquitylation couples maintenance DNA methylation and replication. *Nature*, **502**, 249-253.

24. Qin, W., Wolf, P., Liu, N., Link, S., Smets, M., La Mastra, F., Forne, I., Pichler, G., Horl, D., Fellinger, K. *et al.* (2015) DNA methylation requires a DNMT1 ubiquitin interacting motif (UIM) and histone ubiquitination. *Cell research*.

25. Harrison, J.S., Cornett, E.M., Goldfarb, D., DaRosa, P.A., Li, Z.M., Yan, F., Dickson, B.M., Guo, A.H., Cantu, D.V., Kaustov, L. *et al.* (2016) Hemi-methylated DNA regulates DNA methylation inheritance through allosteric activation of H3 ubiquitylation by UHRF1. *eLife*, **5**.

26. Ishiyama, S., Nishiyama, A., Saeki, Y., Moritsugu, K., Morimoto, D., Yamaguchi, L., Arai, N., Matsumura, R., Kawakami, T., Mishima, Y. *et al.* (2017) Structure of the Dnmt1 Reader Module Complexed with a Unique Two-Mono-Ubiquitin Mark on Histone H3 Reveals the Basis for DNA Methylation Maintenance. *Mol Cell*, **68**, 350-360 e357.

27. Li, T., Wang, L., Du, Y., Xie, S., Yang, X., Lian, F., Zhou, Z. and Qian, C. (2018) Structural and mechanistic insights into UHRF1-mediated DNMT1 activation in the maintenance DNA methylation. *Nucleic Acids Res*, **46**, 3218-3231.

28. Fang, J., Cheng, J., Wang, J., Zhang, Q., Liu, M., Gong, R., Wang, P., Zhang, X., Feng, Y., Lan, W. *et al.* (2016) Hemi-methylated DNA opens a closed conformation of UHRF1 to facilitate its histone recognition. *Nature communications*, **7**, 11197.

29. Morgan, M.T., Haj-Yahya, M., Ringel, A.E., Bandi, P., Brik, A. and Wolberger, C. (2016) Structural basis for histone H2B deubiquitination by the SAGA DUB module. *Science*, **351**, 725-728.

30. Taverna, S.D., Li, H., Ruthenburg, A.J., Allis, C.D. and Patel, D.J. (2007) How chromatin-binding modules interpret histone modifications: lessons from professional pocket pickers. *Nat Struct Mol Biol*, **14**, 1025-1040.

31. Xi, Q., Wang, Z., Zaromytidou, A.I., Zhang, X.H., Chow-Tsang, L.F., Liu, J.X., Kim, H., Barlas, A., Manova-Todorova, K., Kaartinen, V. *et al.* (2011) A poised chromatin platform for TGF-beta access to master regulators. *Cell*, **147**, 1511-1524.

32. Yamaguchi, L., Nishiyama, A., Misaki, T., Johmura, Y., Ueda, J., Arita, K., Nagao, K., Obuse, C. and Nakanishi, M. (2017) Usp7-dependent histone H3 deubiquitylation regulates maintenance of DNA methylation. *Scientific reports*, **7**, 55.

33. Tsumura, A., Hayakawa, T., Kumaki, Y., Takebayashi, S., Sakaue, M., Matsuoka, C., Shimotohno, K., Ishikawa, F., Li, E., Ueda, H.R. *et al.* (2006) Maintenance of self-renewal ability of mouse embryonic stem cells in the absence of DNA methyltransferases Dnmt1, Dnmt3a and Dnmt3b. *Genes to cells : devoted to molecular & cellular mechanisms*, **11**, 805-814.

34. Saksouk, N., Simboeck, E. and Dejardin, J. (2015) Constitutive heterochromatin formation and transcription in mammals. *Epigenetics & chromatin*, **8**, 3.

35. Robertson, K.D. (2005) DNA methylation and human disease. *Nat Rev Genet*, **6**, 597-610.

36. Lehnertz, B., Ueda, Y., Derijck, A.A., Braunschweig, U., Perez-Burgos, L., Kubicek, S., Chen, T., Li, E., Jenuwein, T. and Peters, A.H. (2003) Suv39h-mediated histone H3 lysine 9 methylation

directs DNA methylation to major satellite repeats at pericentric heterochromatin. *Curr Biol*, **13**, 1192-1200.

37. Peters, A.H., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schofer, C., Weipoltshammer, K., Pagani, M., Lachner, M., Kohlmaier, A. *et al.* (2001) Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin and genome stability. *Cell*, **107**, 323-337.

38. Arita, K., Isogai, S., Oda, T., Unoki, M., Sugita, K., Sekiyama, N., Kuwata, K., Hamamoto, R., Tochio, H., Sato, M. *et al.* (2012) Recognition of modification status on a histone H3 tail by linked histone reader modules of the epigenetic regulator UHRF1. *Proc Natl Acad Sci U S A*, **109**, 12950-12955.

39. Cheng, J., Yang, Y., Fang, J., Xiao, J., Zhu, T., Chen, F., Wang, P., Li, Z., Yang, H. and Xu, Y. (2013) Structural insight into coordinated recognition of trimethylated histone H3 lysine 9 (H3K9me3) by the plant homeodomain (PHD) and tandem tudor domain (TTD) of UHRF1 (ubiquitin-like, containing PHD and RING finger domains, 1) protein. *J Biol Chem*, **288**, 1329-1339.

40. Karagianni, P., Amazit, L., Qin, J. and Wong, J. (2008) ICBP90, a novel methyl K9 H3 binding protein linking protein ubiquitination with heterochromatin formation. *Mol Cell Biol*, **28**, 705-717.

41. Rothbart, S.B., Dickson, B.M., Ong, M.S., Krajewski, K., Houliston, S., Kireev, D.B., Arrowsmith, C.H. and Strahl, B.D. (2013) Multivalent histone engagement by the linked tandem Tudor and PHD domains of UHRF1 is required for the epigenetic inheritance of DNA methylation. *Genes Dev*, **27**, 1288-1298.

42. Rothbart, S.B., Krajewski, K., Nady, N., Tempel, W., Xue, S., Badeaux, A.I., Barsyte-Lovejoy, D., Martinez, J.Y., Bedford, M.T., Fuchs, S.M. *et al.* (2012) Association of UHRF1 with methylated H3K9 directs the maintenance of DNA methylation. *Nat Struct Mol Biol*, **19**, 1155-1160.

43. Xie, S., Jakoncic, J. and Qian, C. (2012) UHRF1 double tudor domain and the adjacent PHD finger act together to recognize K9me3-containing histone H3 tail. *J Mol Biol*, **415**, 318-328.

44. Hashimoto, H., Horton, J.R., Zhang, X. and Cheng, X. (2009) UHRF1, a modular multi-domain protein, regulates replication-coupled crosstalk between DNA methylation and histone modifications. *Epigenetics*, **4**, 8-14.

45. Jones, P.A. and Liang, G. (2009) Rethinking how DNA methylation patterns are maintained. *Nat Rev Genet*, **10**, 805-811.

46. Jurkowska, R.Z., Jurkowski, T.P. and Jeltsch, A. (2011) Structure and function of mammalian DNA methyltransferases. *Chembiochem : a European journal of chemical biology*, **12**, 206-222.

47. Guo, X., Wang, L., Li, J., Ding, Z., Xiao, J., Yin, X., He, S., Shi, P., Dong, L., Li, G. *et al.* (2014) Structural insight into autoinhibition and histone H3-induced activation of DNMT3A. *Nature*.

48. Chambers, I., Colby, D., Robertson, M., Nichols, J., Lee, S., Tweedie, S. and Smith, A. (2003) Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell*, **113**, 643-655.

49. Otwinowski, Z. and Minor, W. (1997) Processing of X-ray diffraction data collected in oscillation mode. *Methods in enzymology*, **276**, 307-326.

50. Kabsch, W. (2010) Xds. *Acta Crystallogr D Biol Crystallogr*, **66**, 125-132.

51. McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C. and Read, R.J. (2007) Phaser crystallographic software. *J Appl Crystallogr*, **40**, 658-674.

52. Emsley, P. and Cowtan, K. (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr*, **60**, 2126-2132.

53. Adams, P.D., Grosse-Kunstleve, R.W., Hung, L.W., Ioerger, T.R., McCoy, A.J., Moriarty, N.W., Read, R.J., Sacchettini, J.C., Sauter, N.K. and Terwilliger, T.C. (2002) PHENIX: building new software for automated crystallographic structure determination. *Acta Crystallogr D Biol Crystallogr*, **58**, 1948-1954.

54. Zhang, Z.M., Lu, R., Wang, P., Yu, Y., Chen, D., Gao, L., Liu, S., Ji, D., Rothbart, S.B., Wang, Y. *et al.* (2018) Structural basis for DNMT3A-mediated de novo DNA methylation. *Nature*, **554**, 387-391.

55. Lu, R., Wang, P., Parton, T., Zhou, Y., Chrysovergis, K., Rockowitz, S., Chen, W.Y., Abdel-Wahab, O., Wade, P.A., Zheng, D. *et al.* (2016) Epigenetic Perturbations by Arg882-Mutated DNMT3A Potentiate Aberrant Stem Cell Gene-Expression Program and Acute Leukemia Development. *Cancer Cell*, **30**, 92-107.

56. Cai, L., Rothbart, S.B., Lu, R., Xu, B., Chen, W.Y., Tripathy, A., Rockowitz, S., Zheng, D., Patel, D.J., Allis, C.D. *et al.* (2013) An H3K36 Methylation-Engaging Tudor Motif of Polycomb-like Proteins Mediates PRC2 Complex Targeting. *Mol Cell*, **49**, 571-582.

57. Xu, B., On, D.M., Ma, A., Parton, T., Konze, K.D., Pattenden, S.G., Allison, D.F., Cai, L., Rockowitz, S., Liu, S. *et al.* (2015) Selective inhibition of EZH2 and EZH1 enzymatic activity by a small molecule suppresses MLL-rearranged leukemia. *Blood*, **125**, 346-357.

58. Krueger, F. and Andrews, S.R. (2011) Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics*, **27**, 1571-1572.

59. Langmead, B., Trapnell, C., Pop, M. and Salzberg, S.L. (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome biology*, **10**, R25.

60. Schultz, M.D., Schmitz, R.J. and Ecker, J.R. (2012) 'Leveling' the playing field for analyses of single-base resolution DNA methylomes. *Trends in genetics : TIG*, **28**, 583-585.

61. Quinlan, A.R. and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*, **26**, 841-842.

Figure legends

Figure 1. Specific interaction between the DNMT1 RFTS domain and H3K9me3Ub.

(**A**) Domain architecture of human DNMT1 (hDNMT1), with individual domains delimited by residues numbers. (**B,C**) ITC binding assays of hDNMT1$_{RFTS}$ over H3$_{1-22}$ (**B**) and H3$_{1-22}$K9me3 (**C**) peptides. (**D,E**) ITC binding assays of hDNMT1$_{351-1600}$ over H3$_{1-24}$Ub2 (**D**) and H3$_{1-24}$K9me3Ub2 (**E**) peptides. (**F**) *In vitro* DNA methylation assays for hDNMT1$_{351-1600}$ in the absence or presence of H3 peptides with the indicated modification. Mean and s.d. were derived from three independent measurements. n.s., not significant; **$p < 0.01$; ***$p < 0.001$.

Figure 2. Structural details for the bDNMT1$_{RFTS}$ domain in complex with H3K9me3K18C/K23C and ubiquitin.

(**A**) Crystal structure of bovine DNMT1$_{RFTS}$ (light green) in complex with H3$_{1-24}$K9me3/K18C/K23C peptide (yellow) and G76C-mutated ubiquitin (magenta and salmon). The zinc ions are shown as purple spheres. (**B**) Crystal structure of bovine DNMT1$_{RFTS}$ (light green) in complex with H3$_{1-24}$K9me3/K18C/K23C peptide (yellow) and G76C-mutated ubiquitin (purple and salmon). The Fo-Fc omit map (blue) of the H3 peptide was contoured at the 1.5 σ level. (**C**) Close-up views of the intermolecular interactions between bDNMT1 RFTS (light green) and the H3$_{1-24}$K9me3K18CK23C peptide (yellow stick). The two ubiquitin molecules are colored in salmon and magenta, respectively. Hydrogen bonds are shown as dashed lines. The zinc ion is shown as purple sphere. (**D**) Close-up views of the close proximity between the H3$_{1-24}$K9me3K18CK23C peptide (yellow stick) and the ubiquitin molecules (salmon and magenta). (**E**) Close-up views of the intermolecular interactions between URL of

bDNMT1 RFTS (light green) and two ubiquitin molecules (magenta and salmon). (**F,G**) Close-up views of the intermolecular interactions between the conserved I44 patch of two ubiquitin molecules (magenta and salmon) and bDNMT1 RFTS (light green).

Figure 3. Structure of the H3K9me3-binding pocket.

(**A**) Close-up view of the H3K9me3-binding pocket with residues from bDNMT1 RFTS, H3K9me3 and ubiquitin molecules shown as light green, yellow and salmon sticks. (**B**) Close-up view of the interaction between the residue W889 of the TRIM33 PHD domain and H3K9me3 (PDB 3U5N). (**C**) Structural superposition of bDNMT1$_{RFTS}$-H3K9me3-Ub2 and hDNMT1$_{RFTS}$-H3Ub2 complexes (PDB 5WVO), with the H3K9me3-binding sites shown in the expanded view. (**D**) Sequence alignment of the DNMT1 RFTS domain from human (hDNMT1), bovine (bDNMT1), mouse (mDNMT1), Xenopus Laevis (xDNMT1) and Zebrafish (zDNMT1). Strictly conserved residues are colored in white and shaded in red. Similar residues are colored in red. The secondary structures corresponding to hDNMT1 RFTS are marked on top. The residues for interaction with H3K9me3 peptide are marked by blue arrows. The residues for interacting with ubiquitin molecules are marked by dark arrows. The H3 K9me3-interacting residues are marked by red arrows.

Figure 4. Biochemical and cellular analysis of the RFTS-H3K9me3Ub2 interaction.

(**A**) EMSA analysis of the interaction between hDNMT1$_{351-597}$, either wild type (WT), W465A or W464A/W465A, and the H3$_{1-24}$K9me3Ub2 peptide. (**B**) EMSA analysis of the interaction between hDNMT1$_{351-1600}$, either wild type (WT), W465A or W464A/W465A, and the H3$_{1-24}$K9me3Ub2 peptide. (**C,D**) ITC binding assays of hDNMT1$_{RFTS}$ W465A mutant over H3$_{1-22}$K9m3 (**C**) and H3$_{1-22}$ (**D**) peptides. (**E**) ITC binding assays of hDNMT1$_{RFTS}$ W464A/W465A mutant over H3$_{1-22}$K9me3 peptide. (**F**) Immunoblots of the indicated Flag-tagged DNMT1 after stable reconstitution into the independently derived 1KO-ESC lines. (**G**) CoIP (upper panels) detecting association of the indicated Flag-tagged DNMT1 with H3K9me3 or H4K20me3. Bottom panels are immunoblots of input. (**H**) Representative confocal immunofluorescence images revealing localization of the indicated DNMT1 (Flag-tagged, red), H3K9me3 (green) and chromatin (stained by DAPI, blue) in the 1KO-ESC stable expression lines synchronized at S phase. Scale bar, 5 micrometers. (**I**) DNA methylation activities of hDNMT1$_{351-1600}$, either WT, W465A or W464A/W465A, in the absence of presence of H3$_{1-24}$K9me3Ub2 peptide. Mean and s.d. were derived from three independent measurements. (**, $p < 0.01$, ***, $p < 0.001$, n.s. not significant, Student's *t*-test)

Figure 5. The role of RFTS mutant in cellular CpG methylation pattern and IR response.

(**A**) LC-MS analysis of global 5-methyl-2-deoxycytidine (5-mdC) levels in 1KO-ESC lines after stable transduction of empty vector or the indicated DNMT1 (n = 3-6 biological replicates). Data are mean±s.d. **, $p < 0.01$; ****, $p < 0.0001$. (**B**) Bar plots showing the CpG methylation levels in 1KO-ESC lines with stable expression of the indicated DNMT1. (**C**) Box plot shows weighted methylation levels of CpGs located within H3K9me3 peaks in 1KO-ESC lines with stable expression of the indicated DNMT1. Scores are calculated after aggregating data from three replicated samples per group. Only H3K9me3 peaks with the mapped reads are included (n = 103,603 for WT,

101,791 for W465, and 102,892 for W464A/W465A). In the plot, the box depicts the 25th to 75th percentiles, with the band in the box representing the median. (**D**) Representative IGV view shows CpG methylation at an H3K9me3-marked genomic region located in the chromosome X among three replicated 1KO-ESC lines with stable expression of the indicated DNMT1. Cytosines covered by at least 5 reads according to eRRBS data are shown, with each site designated by a vertical line. (**E,F**) Neutral comet assays revealing DNA breaks (DNA breaks quantified in panel **F**) after ionizing radiation (IR) treatment of 1KO-ESC cells reconstituted with vector control or the indicated DNMT1. Box-and whisker plots in panel F depict 25-75% in the box, whiskers are 10-90%, and median is indicated. Data represent the mean ± S.E.M. from >100 cells (N = 3 biologically independent replicates).

Figure 1

Figure 2

Figure 3

Figure 4

A

hDNMT1_{RFTS}

	WT				W465A				W464A/W465A			

H3K9me3Ub2 (μM)   0   0.5   1   2   0   0.5   1   2   0   0.5   1   2

Complex–

Free hDNMT1_{RFTS}–

B

hDNMT1₃₅₁₋₁₆₀₀

	WT				W465A				W464A/W465A			

H3K9me3Ub2 (μM)   0   0.5   1   2   0   0.5   1   2   0   0.5   1   2

Complex –
Free hDNMT1 –

C

Time (min)
0   10   20   30   40   50   60   70

μcal/sec

kcal mol⁻¹ of injectant

W465A
hDNMT1_{RFTS}
vs
H3K9me3

K_d = 22 μM

Molar Ratio

D

Time (min)
0   10   20   30   40   50   60   70

μcal/sec

kcal mol⁻¹ of injectant

W465A
hDNMT1_{RFTS}
vs
H3K9me0

K_d = 30 μM

Molar Ratio

E

Time (min)
0   10   20   30   40   50   60   70

μcal/sec

kcal mol⁻¹ of injectant

W464A/W465A
hDNMT1_{RFTS}
vs
H3K9me3

Molar Ratio

F

Radioactivity(x10³ c.p.m.)

☐ Peptide free
▨ H3K9me3Ub2

***

**

n.s.

WT   W465A   W464A/W465A

G

	vector	WT			W465A			W464A/W465A		
		1	2	3	1	2	3	1	2	3

KD

150–

50–

H

	vector	WT	W465A	vector	WT	W464A/W465A	

Flag IP

Flag-DNMT1
H3K9me3
H4K20me3

Input

Flag-DNMT1
H3K9me3
H4K20me3
General H3

I

	Flag-DNMT1	H3K9me3	DAPI	Merged
WT				
W464A / W465A				
W465A				

Figure 5

Supplementary Information for
Direct readout of heterochromatic H3K9me3 regulates DNMT1-mediated maintenance DNA methylation

Wendan Ren[1,12], Huitao Fan[2,3,12], Sara A Grimm[4], Yiran Guo[2,3], Jae Jin Kim[5], Linhui Li[1], Christopher James Petell[2,3], Xiao-Feng Tan[1], Zhi-Min Zhang[1,†], John P. Coan[6], Jiekai Yin[7], Linfeng Gao[7], Ling Cai[2,3], Brittany Detrick[1], Burak Çetin[8], Dinshaw J. Patel[9], Yinsheng Wang[7,10], Qiang Cui[11], Brian D. Strahl[2,3], Or Gozani[6], Kyle M. Miller[5], Seán E. O'Leary[1,8], Paul A. Wade[4], Gang Greg Wang[2,3,*], Jikui Song[1,7,*]

[1]Department of Biochemistry, University of California, Riverside, CA 92521, USA

[2]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27599, USA

[3]Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27599, USA

[4]Division of Intramural Research, Epigenetics and Stem Cell Biology Laboratory, National Institute of Environmental Health Sciences, Research Triangle Park, Durham, NC 27709, USA

[5]Department of Molecular Biosciences, LIVESTRONG Cancer Institute of the Dell Medical School, Institute for Cellular and Molecular Biology, University of Texas at Austin, Austin, TX 78712, USA

[6]Department of Biology, Stanford University, Stanford, CA 94305, USA

[7]Environmental Toxicology Graduate Program, University of California, Riverside, CA 92521, USA

[8]Cell, Molecular, and Developmental Biology Graduate Program, University of California, Riverside, Riverside, CA 92521.

[9]Structural Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA.

[10]Department of Chemistry, University of California, Riverside, CA 92521, USA

[11]Departments of Chemistry, Physics and Biomedical Engineering, Boston University, Boston, MA 02215, USA.

[†]Current address: School of pharmacy, Jinan University, 601 Huangpu Avenue West, Guangzhou 510632, China

[12]These authors contributed equally to this work

*Correspondence: greg_wang@med.unc.edu; jikui.song@ucr.edu

Supplementary Figure S1. Biochemical analysis of the interaction between DNMT1 RFTS domain and histone peptides.

(**A-E**) ITC binding curves of hDNMT1$_{RFTS}$ with the H3$_{1-22}$K9Ac peptide (**A**), H3$_{1-15}$K4me3 peptide (**B**), H3$_{21-33}$K27me3 peptide (**C**), H3$_{31-41}$K36me3 peptide (**D**) and H4$_{11-25}$K20me3 peptide (**E**). (**F**) DNA methylation activity of hDNMT1$_{351-1600}$ in the presence of 100-fold molar excess of H3K9me0 or H3K9me3 peptides. Mean and s.d. were derived from three independent measurements. (***, $p < 0.001$, Student's t-test).

Supplementary Figure S2. *In vitro* **DNA methylation assays of hDNMT1**$_{351\text{-}1600}$ **mutants with H3 peptide.**

(**A**) DNA methylation activity of W465A- or W464A/W465A-mutated hDNMT1$_{351\text{-}1600}$ in the presence of H3K9me0 or H3K9me3 peptides. (**B**) DNA methylation activity of W465A- or W464A/W465A-mutated hDNMT1$_{351\text{-}1600}$ in the absence or presence of the H3K9me3Ub. Mean and s.d. were derived from three independent measurements. (n.s. not significant).

Supplementary Figure S3. eRRBS profilings of DNA methylation in 1KO-ESC cells reconstituted with either WT or RFTS-mutated DNMT1.

(**A**) The rates of bisulfite conversion, labeled on top of columns, for all cytosines in each 1KO-ESC cell sample with expression of the indicated DNMT1, as determined by the unmethylated lambda DNA used as spike-in controls. (**B**) Distribution of absolute methylation levels for CpG sites with >5 coverage among 1KO-ESC lines with stable expression of the indicated DNMT1, as detected by eRRBS. (**C**) Violin plots showing distribution of absolute methylation levels for CpG sites with >5 coverage at all CpG sites or those within the repeated genomic sequences among the indicated cells samples. White dots are the median and box lines are the first and third quartile of the data. (**D**) Representative IGV views of CpG methylations at an H3K9me3 (red, bottom)

marked genomic region located in the chromosome 1 among three replicated 1KO-ESC lines with stable expression of the indicated DNMT1. Cytosines covered by at least 5 reads according to eRRBS data are shown, with each site designated by a vertical line.

Supplementary Table S1. Summary of ITC binding parameters.

Protein	Peptide	K_d (µM)	N value
hDNMT1$_{RFTS}$, WT	H3(1-22)[#]	6.4 ± 0.8	0.74±0.05
hDNMT1$_{RFTS}$, WT	H3(1-22)K9me3[#]	1.3 ± 0.1	0.77± 0.1
hDNMT1$_{351-1600}$, WT	H3(1-24)Ub2[#]	91 ± 20	0.91± 0
hDNMT1$_{351-1600}$, WT	H3(1-24)K9me3Ub2[#]	17 ± 0.3	0.96±0.1
hDNMT1$_{RFTS}$, WT	H3(1-22)K9Ac	61 ± 14	0.95±0.1
hDNMT1$_{RFTS}$, WT	H3(1-15)K4me3	284 ± 32	1.0*
hDNMT1$_{RFTS}$, WT	H3(21-33)K27me3	NDB	
hDNMT1$_{RFTS}$, WT	H3(31-43)K36me3	562 ±77	1.0*
hDNMT1$_{RFTS}$, WT	H4(14-25)K20me3	339 ±11	1.0*
hDNMT1$_{RFTS}$, W465A	H3(1-22)K9me3	22 ± 0.7	1.1±0.04
hDNMT1$_{RFTS}$, W465A	H3(1-22)	30 ± 6	1.0±0.2
hDNMT1$_{RFTS}$, W464AW465A	H3(1-22)K9me3	NDB	

NDB, no detectable binding. *The N value was set manually. [#]The mean value and S.D. were derived from two-independent measurements.

Supplementary Table S3. Summary of eRRBS data analysis.

Supplementary Table S2. Data collection and refinement statistics.

	bDNMT1 RFTS – Ubiquitin – H3K9me3 (PDB: XXXX)
Data collection	
Space group	P 2_1 $2_1$2
Cell dimensions	
a, b, c (Å)	70.3, 196.4, 67.7
α, β, γ (°)	90, 90, 90
Resolution (Å)	49.11-3.01(3.12-3.01)[a]
R_{merge}	0.126(1.023)
$I/\sigma(I)$	12.3(1.3)
$CC_{1/2}$	0.998(0.583)
Completeness (%)	99.1(92.3)
Redundancy	9.8(7.9)
Refinement	
No. reflections	19,094
R_{work} / R_{free}	0.212/0.275
No. atoms	
Protein	6184
Zn^{2+}	2
Water	48
B factors (Å^2)	
Protein	85.1
Zn^{2+}	82.0
Water	67.5
r.m.s deviations	
Bond lengths (Å)	0.003
Bond angles (°)	0.588

[a]Values in parentheses are for highest-resolution shell.