Antibody repertoire and gene expression dynamics of diverse human B cell states during affinity maturation.

Hamish W King[1,2] *, Nara Orban[3], John C Riches[4,5], Andrew J Clear[4], Gary Warnes[6], Sarah A Teichmann[2,7], Louisa K James[1] * §

[1] Centre for Immunobiology, Blizard Institute, Queen Mary University of London, London E1 2AT, UK

[2] Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SA, UK

[3] Barts Health Ear, Nose and Throat Service, The Royal London Hospital, London E1 1BB, UK

[4] Centre for Haemato-Oncology, Barts Cancer Institute, Queen Mary University of London, London EC1M 6BQ, UK

[5] The Francis Crick Institute, London NW1 1AT, UK

[6] Flow Cytometry Core Facility, Blizard Institute, Queen Mary University of London, London E1 2AT, UK

[7] Theory of Condensed Matter, Cavendish Laboratory, Department of Physics, University of Cambridge, Cambridge CB3 0EH, UK

§ Lead contact

* To whom correspondence can be addressed: h.king@qmul.ac.uk, drhamishking@gmail.com and louisa.james@qmul.ac.uk

Abstract

In response to antigen challenge, B cells clonally expand, undergo selection and differentiate to produce mature B cell subsets and high affinity antibodies. However, the interplay between dynamic B cell states and their antibody-based selection is challenging to decipher in primary human tissue. We have applied an integrated analysis of bulk and single-cell antibody repertoires paired with single-cell transcriptomics of human B cells undergoing affinity maturation. We define unique gene expression and antibody repertoires of known and novel B cell states, including a pre-germinal centre state primed to undergo class switch recombination. We dissect antibody class-dependent gene expression of germinal centre and memory B cells to find that class switching prior to germinal centre entry dictates the capacity of B cells to undergo antibody-based selection and differentiate. Together, our analyses provide unprecedented resolution into the gene expression and selection dynamics that shape B cell-mediated immunity.

Introduction

Effective immunity depends on the ability of B cells to evolve a functional antibody repertoire in response to antigen challenge. Following antigen encounter, activated B cells either differentiate into short-lived plasma cells or following cognate interaction with antigen-specific T cells can form germinal centres (GCs) within secondary lymphoid tissues, such as the spleen, peripheral lymph nodes and tonsils (Cyster and Allen, 2019). These GCs are transient structures in which B cells undergo iterative cycles of clonal expansion and somatic hypermutation in the variable regions of their immunoglobulin heavy and light chain genes followed by affinity-based selection for clones with high antigen-specificity. This highly dynamic process occurs in spatially and transcriptionally distinct dark and light zones (DZ and LZ) under the regulation of a network of specialised T follicular helper cells, follicular dendritic cells and

macrophages (Mesin et al., 2016). B cells can differentiate and exit the GC reaction either as antibody-secreting plasmablasts committed to the plasma cell lineage or memory B cells, which are long-lived quiescent cells capable of being reactivated upon secondary exposure to the antigen (Suan et al., 2017b). The effector functions of antibodies expressed by B cells are broadly determined by antibody class (IgM, IgD, IgG, IgA, IgE) and more precisely by isotype or subclass (IgG1-4, IgA1-2), specified by the constant domain genes in the immunoglobulin heavy chain (IgH) locus. Although all naïve B cells express IgM and IgD, during maturation they may undergo class switch recombination, which involves the deletional recombination of IgM and IgD constant domain genes and expression of a different downstream constant domain gene (IgG1-4, IgA1-2 or IgE) (Stavnezer and Schrader, 2014). The combined outcomes of B cell differentiation, antigen affinity maturation and class switch recombination ultimately shape the antibody repertoire and the B cell-mediated immune response more broadly.

During their maturation in germinal centres, B cells express their antibody immunoglobulin genes as part of a membrane-bound complex termed the B cell receptor (BCR). Antigen-binding and downstream signalling of the BCR is a primary determinant of GC B cell survival and even differentiation through differential expression of key transcription factors regulated by BCR-mediated signalling (Kwak et al., 2019, Shlomchik et al., 2019). Studies have shown that BCR activation thresholds and downstream signalling can differ as a result of isotype-specific differences in the extracellular, transmembrane and intracellular domains of immunoglobulin proteins forming the BCR (Martin and Goodnow, 2002, Engels et al., 2014, Xu et al., 2014b). Thus, as well as shaping the effector functions of the subsequent antibody repertoire, class switch recombination contributes towards B cell survival or fate specification within the GC reaction. However, resolving the combined contribution of somatic hypermutation, maturation and class switching in the polyclonal context of primary human lymphoid tissues remains an enormous challenge for the field (Mesin et al., 2016). Furthermore, while it has long been held that class switch recombination occurs exclusively within the GC, as this is where the highest AICDA expression is detected, other work has demonstrated that class switch recombination often occurs prior to formation of the GC response (Roco et al., 2019, Toellner et al., 1996, Pape et al., 2003). This has raised questions about our current understanding of the cellular states and dynamics during human B cell maturation *in vivo* and demands a systematic and unbiased approach to better define how the human antibody repertoire is shaped through somatic hypermutation, class switch recombination and differentiation into different B cell fates.

We have applied an integrated strategy of bulk and single-cell antibody repertoire analysis paired with single-cell transcriptomics of human B cells from a model secondary lymphoid tissue, tonsils. We compare and contrast the antibody repertoires of major B cell subsets to reveal unique class switch hierarchies of memory B cells and plasmablasts. We then discover and define novel transcriptional B cell states during the GC response using single-cell RNA-seq. In particular, we reveal the unique gene expression of a B cell state primed to undergo class switch recombination before entering the GC. By leveraging the single-cell resolution of our datasets, we deconvolve the contribution of somatic

hypermutation and antibody class to gene expression patterns linked with altered BCR signalling, B cell
maturation and fate decisions within the GC. Finally, we define diverse memory B cell states within
secondary lymphoid tissue and explore the impact of class switch recombination on their functional
potential. Our analyses reveal a striking importance for class switch recombination in shaping B cell
fate and maturation in the GC and memory B cell fate that reframes our understanding of antibody-
based selection and B cell differentiation.

Results

Subset- and subclass-specific antibody class switch recombination landscapes.

To begin to untangle the antibody-based selection of different B cell fates, we characterised the antibody
repertoires of four broadly defined B cell subsets from the human tonsil; naïve, germinal centre (GC) B
cells, memory B cells (MBCs) and plasmablasts, in addition to total CD19+ B cells (Figure 1A). We
applied a subclass-specific and quantitative unique molecular identifier-based repertoire sequencing
protocol for IgH VDJ sequences (Horns et al., 2016). Naïve B cells were comprised of more than 95%
unswitched and unmutated IgM and IgD sequences, while GC and MBC samples consisted of both
switched and unswitched IgH sequences with elevated somatic hypermutation (Figure 1B-C).
Plasmablasts were nearly all switched and highly mutated. Consistent with the low abundance of IgE+
B cells in human tonsils, we detected only a single IgE sequence, preventing us from drawing any
conclusions about IgE-expressing B cells.

Higher IgH somatic hypermutation frequencies within the GC are typically a reflection of higher affinity
BCRs and are proposed to bias GC B cells towards the plasmablast cell fate rather than the MBC fate
(Shinnakasu et al., 2016, Suan et al., 2017a). In keeping with this, plasmablast-derived antibody
repertoires generally contained higher somatic hypermutation frequencies than those of MBCs at a bulk
level (Figure 1C). However, by resolving for antibody subclass, we found that while MBC-derived IgM
and IgD sequences were consistently less mutated than plasmablast-derived IgM and IgD, the somatic
hypermutation levels for switched isotypes were broadly similar between different B cell subsets (Figure
1D). Comparison of the clonal diversity of subclass-specific MBCs and plasmablasts revealed
unswitched and IgA+ MBCs were less clonally expanded (as evidenced by higher diversity) than
plasmablasts of the same isotype, while IgG+ MBCs and plasmablasts appeared to have clonally
expanded to similar degrees (Figure 1E). This is not likely explained by differences in their somatic
hypermutation frequencies (Figure 1D), but instead may reflect differential selection or differentiation
processes linked with specific class switch recombination outcomes.

Figure 1. Subclass- and subset-specific features of human tonsillar B cell repertoires.

A) Cell sorting strategy to isolate naïve, germinal centre (GC), memory B cells (MBC) and plasmablasts (PB) from live CD19+ human tonsillar B cells for antibody repertoire analysis. Representative of *n*=8.

B) Mean antibody subclass frequencies within each B cell subset across donors.

C) Mean frequencies of antibody somatic hypermutation levels (None, Low, High) within each B cell subset across donors.

D) Somatic hypermutation frequencies for subclass-specific antibody sequences within each B cell subset. Violin plots show all unique sequences per subset, while points represent mean for each donor.

E) Mean clonal diversity scores per donor of subclass- and subset-specific B cell clones.

F) Mean switched antibody subclass frequencies within each B cell subset across donors.

G) Mean subclass frequencies for expanded clones spanning different B cell populations (x axis). For each class of clone, subset-specific members are examined (see groups at top).

H) Observed/expected frequencies for isotype pairs detected in reconstructed phylogenies of clonally-related sequences. Antibody subclasses are ordered according to the IgH locus. nd denotes not detected.

I) Same as in H), except phylogenies are restricted to subset-specific sequences.

We next examined whether there were differences in antibody subclass frequencies or the manner in

which subclasses might have arisen in our broadly defined B cell populations. Intriguingly, we observed

that as well as an increased propensity to retain IgM expression (Figure 1B), MBCs were 3.3- or 7.3-

fold more likely than plasmablasts to express IgA1 or IgA2 respectively, while plasmablasts were

significantly more likely to express IgG1 (Figure 1F). These enrichments were linked with specific B cell

fates even for expanded clones spanning different B cell populations (Figure 1G). Finally, to explore

how these unique class switch patterns might have arisen, we reconstructed phylogenies for 28,845

expanded B cell clones and calculated the likelihood that specific class switch recombination events

were observed compared to that expected by chance (Figure 1H-I). Clonal lineages within the MBC

pool exhibited greater likelihoods for switching of isotype pairs located close to each other in linear

space along the IgH locus, compared to plasmablast clones which demonstrated a more eclectic pattern

of class-switch likelihoods (Figure 1I). Of note, both the antibody subclass frequencies and

reconstructed class switch hierarchies of MBCs closely resembled those of GC cells, consistent with

models that propose a stochastic exit of MBCs from the GC (Duffy et al., 2012, Good-Jacobson and

Shlomchik, 2010). Together, these analyses reveal that the antibody-based selection mechanisms of

two major mature B cell subsets (MBCs and plasmablasts) exhibit important differences related

primarily to their propensity to have undergone class switch recombination earlier in their maturation.

Single-cell atlas of tonsillar immune cells defines B cell states during affinity maturation.

To better understand the antibody-based selection mechanisms shaping the maturation of different B

cell subsets, we performed single-cell RNA-seq (scRNA-seq) paired with single-cell B cell repertoire

VDJ sequencing (scVDJ-seq) for unsorted tonsillar immune cells from the same samples used for our

bulk B cell repertoire analyses (Figure 2A-C). After stringent quality control, we retained the

transcriptomes of 32,607 immune cells (n=7; median of 3142 and mean of 4658 cells per donor) from

which we identified 30 transcriptionally distinct cell types (Figure 2D-F). Although our primary focus was

understanding B cell maturation, given the importance of other immune cell populations in this process

we also annotated varied T cell and non-lymphoid populations, including naïve and/or central memory

(NCM) CD4+ and CD8+ T cells, follicular helper (TfH) T cells, follicular regulatory T (Tfr) cells, regulatory

T cells (Treg), cytotoxic CD8+ T cells, innate lymphoid cells (ILCs), natural killer (NK) cells, follicular

dendritic cell (FDC), plasmacytoid-derived dendritic cell (pDCs), classical dendritic cell (cDC1), and

several macrophage (MAC) clusters (Figures 2D-E and S1) that will act as a valuable resource for those

studying immune cell dynamics in human secondary lymphoid tissues.

We next characterised tonsillar B cell states using both their unique gene expression and scVDJ-

derived antibody repertoire features. To improve the power and accuracy of our scVDJ analyses we

applied a novel strategy of integrating single-cell and bulk repertoires to benefit from the deeper

sampling of IgH sequences from bulk repertoires to enhance genotype correction, identification of

clonally-related sequences and quantitation of somatic hypermutation levels for scVDJ-derived
sequences (Figure 2G). We defined gene expression signatures for 12 distinct B cell types or states
(Figure 2H) and complemented marker gene-based annotation with antibody isotype frequencies
(Figure 2I), somatic hypermutation levels (Figure 2J), clonal diversity (Figure 2K-L) and relationships
with other B cell subsets (Figure 2M-N). We identified all major stages of B cell maturation, including
naïve, activated, GC (including both LZ and non-proliferating DZ cells), MBCs, tissue-resident FCRL4+
MBCs, plasmablasts and a cycling population consisting mostly of DZ GC cells (Figure 2H). Whilst
apoptotic cells normally comprise a sizeable proportion of GC B cells these were not retained in our
analysis as they failed to generate sufficiently high quality transcriptomic data. Our analysis of B cell
states also identified a "preGC" B cell state expressing unmutated IgM and IgD that transcriptionally
shared markers with both naïve and LZ GC populations, but had yet to acquire features consistent with
B cell maturation in the GC such as *CD27* and *CD38* expression, hypermutated antibody genes or
clonal expansion (see Figure 3 for further details). We also annotated a population of class switched
and hypermutated GC B cells that amongst other transcriptionally unique features (*FRZB*, *BTNL9*,
FGFR1) express low to intermediate levels of the plasmablast-specific transcription factors *PRDM1* and
XBP1 (Figure 2H), suggesting that these cells may be a pre-plasmablast (prePB) state within the GC.
Finally, we discover a transcriptionally distinct and clonally-expanded IgM+ B cell population in the GC
with elevated expression of genes associated with inhibitory BCR signalling, such as *FCRL2*, *FCRL3*,
SAMSN1, and *SIGLEC10* that we have labelled as FCRL2/3^{high} GC B cells. Some of these cells were
part of large expanded GC-derived clones that also contained MBCs or plasmablasts (Figure 2M-N),
indicating that this cell state arises as part of productive GC reactions and is unlikely to be derived from
a separate B cell lineage.

Crucially, all annotated B cell states were observed at reproducible frequencies across patients,
regardless of their history of tonsillitis (Figure 2O), although exactly how some of these transcriptional
states relate to other human B cell populations will require further work. We also note that we have
identified a greater diversity of B cell fates in our paediatric tonsil samples (typically <10 years old) than
previous single-cell studies of other human lymphoid tissues from adult donors (typically >40 years old)
(Madissoon et al., 2019, James et al., 2020), highlighting the importance of profiling immunologically
active tissues to understand B cell maturation. Together, our uniquely comprehensive transcriptomic
and repertoire analyses of a model secondary lymphoid organ have allowed us to create a detailed
overview of human B cell maturation that will allow us to interrogate gene expression and antibody
repertoire dynamics before, during and after maturation in the GC.

Figure 2. A single-cell atlas of human tonsillar immune cells to understand B cell maturation.

A) Schematic of tonsillar immune cell isolation followed by single-cell profiling of gene expression and antibody sequences.

B) UMAP projection of tonsillar immune scRNA-seq data (32,607 cells; 7 donors) annotated as B, T or non-lymphoid cells.

C) Same as in B), with cells annotated for whether a high quality immunoglobulin scVDJ sequence was assembled.

....continued on following page...

D) UMAP projection of T cell populations in the tonsillar immune scRNA-seq data, including CD4$^+$ naïve or central memory
(CD4$^+$ NCM), CD4$^+$, T follicular helper (TfH), T follicular regulatory (TfH), T regulatory (Treg), CD8$^+$ naïve or central
memory (CD8$^+$ NCM), CD8$^+$ cytotoxic, TIM3$^+$ CD4/CD8 double-negative (TIM3$^+$ DN) and cycling T cells, in addition to
innate lymphoid cells (ILC) and natural killer (NK) cells.
E) UMAP projection of non-lymphoid cell populations in the tonsillar immune scRNA-seq data, including
monocyte/macrophage precursors (Precursors), macrophages (MAC1, MAC2, MAC3), conventional dendritic cell 1
(cDC1), plasmacytoid-derived dendritic cell (pDC) and follicular dendritic cell (FDC) subsets.
F) UMAP projection of B cell populations in the tonsillar immune scRNA-seq data, including naïve, activated, pre-germinal
centre (preGC), light zone GC (LZ GC), GC, dark zone (DZ GC), cycling, FCRL2/3^{high} GC, pre-plasmablasts (prePB),
plasmablasts, MBC and FCRL4$^+$ MBC subsets.
G) Schematic of scVDJ and bulk repertoire integration, including genotype inference and correction, identification of clonally-
related sequences and quantitation of somatic hypermutation.
H) Mean expression of key marker genes used to define B cell scRNA-seq clusters. Frequency of cells for which each gene
is detected is denoted by size of the dots.
I) Relative scVDJ-derived antibody subclass frequencies in different B cell states.
J) Somatic hypermutation frequencies of scVDJ-derived antibody genes in different B cell states.
K) Clonal diversity scores of B cell clones identified in scRNA-seq dataset. Error bars denote ±SD.
L) Number of members per clonotype in different B cell states from integrated repertoire analysis.
M) Co-occurrence of expanded clones across B cell states. Numbers reflect a binary detection event.
N) Clonal relationships between scRNA-seq-defined B cell states and repertoires of sorted B cell subsets.
O) Relative frequencies of different B cell states separated by clinical indication for tonsillectomy. OSA; obstructive sleep
apnoea (n=2), RT; recurrent tonsillitis (n=3), RT+OSA (n=2).
219

Dynamic gene expression during human B cell activation and GC formation.

Initially in a B cell response, B cells acquire antigen, either in soluble form or displayed on the surface
of follicular antigen-presenting cells (APCs), which results in their activation and migration from the
follicle to the T cell zone where they can form or participate in GC reactions. However, reconstructing
the events during this important process in primary human tissues is extremely challenging. We
therefore used our single-cell profiling of human B cell maturation to explore the dynamics of early B
cell activation and GC formation. We first identified an activated B cell state with elevated expression
of known activation marker genes (Activated; *CD69, CD83*, *JUN*) (Figures 2 and 3A) and which
demonstrate the highest frequency of predicted cell-cell communication with APCs (FDCs, other
dendritic cells, macrophages) (Figure 3B). This activated B cell state appeared capable of coordinating
help from both APCs and T cells through IL6 signalling (Arkatkar et al., 2017, Ise et al., 2018) and/or
ICAM1-ITGAL1 (LFA-1) interactions (Zaretsky et al., 2017) (Figure 3C). Many of these same predicted
cell-cell interactions were also detected in preGC B cells (Figure 3C), suggesting that these might reflect
a transitional cell state between antigen dependent-activation and GC entry or formation. Indeed, we
found that preGC B cells exhibit a strong directionality towards the light zone of the GC using RNA
velocity inference (Figure 3D). We next reconstructed a pseudotemporal trajectory of naïve, activated,
preGC and LZ GC B cells, revealing a continuum of gene expression from early activation events to
bona fide GC B cells and allowing us to identify dynamic expression of key signalling molecules and
transcription factors (Figures 3E-F and S2A). Crucially, these gene expression changes through
pseudotime were well correlated with an experimentally-derived time course of *in vitro*-stimulated B
cells (Figure 3G; Shinohara et al., 2014), strongly supporting our pseudotemporal ordering of defined
B cell states during B cell activation and GC entry. This roadmap of B cell activation may allow future
improvement to *in vitro* B cell culture protocols to better model B cell activation dynamics *in vivo*.

Figure 3. Reconstruction of *in vivo* B cell activation reveals class switch recombination in a preGC state.

A) Mean expression of key marker genes for activated B cells in scRNA-seq.

B) Frequency of significant predicted ligand-receptor pair interactions between major B cell states and different T cell populations and antigen-presenting cells (APCs) performed using CellPhoneDB.

C) Selected ligand–receptor interactions between B cell subsets and APCs, CD4+ T cells and TfH cells with CellPhoneDB.

D) Grid-based visualisation of tonsillar B cell RNA velocities. Arrow size conveys strength of predicted directionality.

E) Relative frequencies of B cell types in a velocity-based pseudotime reconstruction of B cell activation and GC formation.

F) Heatmap depicting dynamic gene expression across velocity-based pseudotime reconstruction in E).

G) Smoothed anti (α)-IgM-treatment gene signature scores (±95% CI) across velocity-based pseudotime.

H) Smoothed expression of class switch recombination genes differentially expressed through velocity-based pseudotime.

I) Smoothed class switch recombination gene signature score through velocity-based pseudotime.

J) Smoothed expression of I promoter germline transcripts for IgM (I_μ), IgA (I_α) and IgG (I_γ) through velocity-based pseudotime. $I_{\alpha+\gamma}$ denotes the sum of I_α and I_γ expression.

K) Relative antibody subclass frequencies across velocity-based pseudotime.

L) Relative frequencies of cells with high class switch recombination signature scores in different B cell states for each donor ($n = 7$). p values denote results of Student's T test.

M) Mean expression of genes implicated in class switch recombination for B cell subsets.

One surprising result from our reconstruction of human B cell activation and GC entry was the discovery that several genes associated with class switch recombination were most highly expressed prior to GC entry, with a specific enrichment of these genes in the preGC B cell population (Figure 3H-I). This

included *APEX1* expression, of which its translated product APE1 is required for class switch

recombination to occur in a dose-dependent manner (Masani et al., 2013, Xu et al., 2014a) and is

expressed almost exclusively by non-GC B cells (Figure S2B; Roco et al., 2019). We also found that

preGC B cells had the highest expression of IgH germline transcripts (GLTs) (Figure 3J), which

preceded switching from IgM/IgD to other isotypes (Figure 3K), in fitting with a recent study describing

GLT transcription prior to GC formation in mouse models (Roco et al., 2019). These findings contrast

with the prevailing view that class switch recombination occurs in the DZ of the GC, where AICDA

expression is highest. However, we detected little enrichment of our class switch recombination

signature in non-cycling DZ GC B cells (Figure 3L), and although there was an enrichment of class

switch recombination genes in cycling B cell populations, this likely reflects their involvement in cell

cycle-linked DNA recombination and repair. Our analysis also highlights many other genes previously

linked with class switch recombination (Figure 3M), including those capable of binding to switch region

sequences within the IgH locus (*NME2*, *NCL*, *DDX21*), interacting with the class switch recombination

machinery (*NPM1*, *SERBP1*) or regulating *AICDA*/AICDA transcript/protein stability (*mir155HG,*

HSP90AB1) (Borggrefe et al., 1998, Hanakahi et al., 1997, Mondal et al., 2016, Orthwein et al., 2010,

Shinozaki et al., 2006, McRae et al., 2017, Zheng et al., 2019). Notably, the microRNA gene *miR155HG*,

formerly called BIC (B-cell Integration Cluster), was up-regulated in preGC B cells and has been shown

to be essential for B cells to form GCs and undergo class switch recombination in mice (Thai et al.,

2007, Vigorito et al., 2007). Furthermore, the transcription factor genes *BATF*, *IRF4* and *BHLHE40* are

enriched in the preGC B cell state, of which BATF and IRF4 are known to regulate GC formation in a B

cell-intrinsic manner (Morman et al., 2018, Willis et al., 2014). Intriguingly, BHLHE40 is capable of

binding to the major regulatory regions α_1 RR and α_2 RR of the IgH locus (Figure S2C-D), implicating

this poorly understood transcription factor in the regulation of class switch recombination prior to GC

formation. Together, these analyses strongly support a model where class switch recombination occurs

primarily before formation of the GC response (Roco et al., 2019, Toellner et al., 1996, Pape et al.,

2003) and our detailed gene expression analyses define the cellular state involved.

Antibody-based selection of B cell fate in the germinal centre

Class switch recombination before entry into the GC has the potential to dramatically influence the

antibody-based selection of B cells within the subsequent GC reaction as a consequence of differential

signalling through the membrane-bound immunoglobulin BCR. We therefore turned to dissect the gene

expression dynamics linked with antibody-based selection and fate specification of B cells within the

GC reaction.

During affinity maturation and selection in the GC, B cells cycle between physically distinct light and

dark zones. While we clearly identified LZ and DZ B cell populations in our scRNA-seq dataset, we also

found that many GC B cells existed in a continuum between these two states (Figure 4A-B) similar to

previous reports (Milpied et al., 2018), with the exception of the FCRL2/3[high] and prePB clusters which

existed as transcriptionally distinct states (Figures 2H and 4C). In addition to unique gene expression

patterns, these two sub-populations of GC B cells also exhibit unique patterns of class switching, with

prePB B cells in the GC more likely to express class-switched isotypes and FCRL2/3[high] GC B cells

more likely to retain expression of IgM and IgD (Figures 2I and 4D). Furthermore, cells clonally related

to FCRL2/3[high] GC B cells were almost exclusively IgM[+] and had rarely undergone class-switching

(Figure 4E). These observations suggested that the outcome of class switch recombination may be

linked with specific gene expression programs of GC B cells.

To address this, we first needed to examine the contribution of antibody maturation state to GC B cell

gene expression programs, given lower average somatic hypermutation frequencies of IgM[+] B cells

compared to B cells expressing switched isotypes (Figure 1D). We leveraged our paired single-cell VDJ

and transcriptomic datasets to stratify all non-cycling GC B cells (excluding prePB and FCRL2/3[high]

populations) based on their IgH somatic hypermutation frequencies as a proxy for affinity (Figure 4F).

Strikingly, GC B cells with high or low somatic hypermutation were significantly enriched with gene sets

derived from experimentally determined high- or low-affinity antigen-binding B cells respectively

(Shinnakasu et al., 2016), higher expression of the B cell maturation marker CD27 and larger clone

sizes (Figure 4F), reflecting increased expansion and maturation based on BCR affinity. We found that

the direct comparison of GC B cells expressing different antibody classes was confounded by gene

expression linked with high and low somatic hypermutation frequencies (Figure 4G), consistent with

high and low affinity binding events differentially regulating GC B cells (Shinnakasu et al., 2016). To

overcome this, we examined GC B cells with matched somatic hypermutation levels expressing

different antibody classes (IgM/D, IgG or IgA) at single-cell resolution. This revealed differential

expression of genes involved directly in cell survival, BCR signalling, antigen presentation, immune

responses and metabolism, as well as other pathways more indirectly linked with BCR activity

(RhoA/RhoGDI, Integrin, NFAT, eIF2), between unswitched and switched GC B cells (Tybulewicz and

Henderson, 2009, Mielke et al., 2011, Arana et al., 2008, Scharenberg et al., 2007) (Figure 4H-J). These

differences may be linked with differential exposure to T cell-derived cytokines such as IL4, TGFB,

IFNG and CD40LG, or signalling through different toll-like receptors (TLR) (Figure 4I). Intriguingly,

several genes involved with GC confinement or regulating B cell niche homing were up-regulated in

IgG[+] and IgA[+] GC B cells, such as genes required for CXCL12-mediated migration to GCs (*LCP1* and

MYO1E (Dubovsky et al., 2013, Girón-Pérez et al., 2020)) and the GC confinement receptor *P2RY8*

(Muppidi et al., 2014).

Figure 4. Influence of antibody class on B cell fate and function within the germinal centre.

A) Diffusion-based graph visualisation and pseudotemporal ordering of GC B cell scRNA-seq populations.

B) Single-cell gene expression heatmap of major GC B cell states ordered by pseudotime within each cluster.

C) Expression of key marker genes for prePB and FCRL2/3high GC B cells.

D) Relative frequency of GC B cells that have undergone class switch recombination (n = 6). "GC" is the combination of LZ GC, GC and DZ GC clusters in (A). Error bars denote SEM.

....continued on following page...

E)   Percentage of members within each GC B cell clone that express IgM or IgD for clones that contain FCRL2/3[high] or prePB
cells. Clonal families include sequences from both scVDJ and bulk repertoires.
F)   High and low affinity gene signature scores for GC B cells (LZ GC, GC and DZ GC clusters) grouped by antibody somatic
hypermutation (SHM) frequency (n = 2045 cells). Also shown are *CD27* expression levels and integrated clone sizes.
G)   Scatterplot comparing log enrichment of genes in class-switched vs unswitched (x axis) and high vs low affinity/SHM (y
axis) GC B cells. Colour denotes statistical significance and c*or* denotes Pearson's Correlation coefficient.
H)   Heatmap of significantly enriched gene ontologies for genes enriched in class-specific high affinity/SHM GC B cells.
I)   Same as in (H), but for gene pathways downstream of cytokines (upper panel) and toll-like receptors (TLRs).
J)   Mean expression of genes enriched in class-specific high affinity/SHM GC B cells grouped by predicted functions.
K)   Mean expression of transcription factor genes enriched in class-specific high affinity/SHM GC B cells.
L)   Single-cell gene expression of transcription factors in class-specific high affinity/SHM GC B cells.
352

Although most gene expression differences were comparable between IgG and IgA and we identified

few significant or meaningful differences for subclass-specific B cells (Figure S3), one interesting

example of class-specific gene expression was the enrichment of *CLEC16A* in IgA+ GC B cells given

that this gene is associated with a selective IgA immunodeficiency (Ferreira et al., 2010). Finally, to try

and understand the upstream regulation of these class-specific gene expression networks we examined

the expression of transcription factors within class-specific GC B cells (Figure 4K-L). We found that

IgM+ B cells express lower levels of transcription factors like *BCL6*, *XBP1* and *ID3* known to regulate

the ability of B cells to remain in the GC or differentiate but higher levels of the transcription factor

BACH2 that represses plasma cell differentiation (Todd et al., 2009, Gloury et al., 2016, Huang et al.,

2014, Shinnakasu et al., 2016). We also found differential expression of other transcription factors such

as *LMO2*, *TOX*, *BCL11A* and *CUX1*, suggesting that they may play a role in the unique transcriptional

wiring of switched and unswitched B cells within the GC. Our single-cell resolution of the GC B cell

response has allowed us to uncouple antibody affinity and class and to dissect the differential

contributions of these two critical arms of the B cell repertoire in shaping B cell fate and function in the

GC. Importantly, these differential gene expression patterns suggest varying abilities of switched and

unswitched B cells to survive and reside in the GC and establish that one of the dominant influences in

shaping antibody-based selection in the GC is whether a B cell has undergone class switch

recombination.

Diverse memory B cell states and activation dynamics in secondary lymphoid tissue.

Maturation state and antibody class appeared to both impact gene expression dynamics within the GC,

presumably through membrane-specific isoforms of immunoglobulin as part of the BCR. Upon GC exit

and differentiation, plasmablasts lose membrane-bound immunoglobulin expression and instead start

to secrete large amounts of soluble antibody. In contrast MBCs retain BCR expression of which the

antibody isotype may influence the phenotypic properties of different MBC subsets (Engels and

Wienands, 2018).

We therefore sought to determine whether antibody class expression by MBCs might be linked with

different functional abilities and to better define the heterogeneity within the MBC pool in secondary

lymphoid tissue. A significant proportion of memory B cells are unswitched (Figure 1B), so to examine

potential differential gene expression across class switched memory B cells subsets we generated
paired single-cell transcriptomics and VDJ repertoires for IgD-depleted or IgM/IgD-depleted MBCs from
the same tonsillar immune cell preparations analysed previously. Dataset integration and quality control
provided 21,595 high-quality MBC single-cell transcriptomes that we annotated with 11 clusters
reflecting different MBC subsets and states (Figure 5A-C), all of which lacked marker gene expression
for naïve or GC B cells (Figure S4A). In addition to tissue-resident FCRL4$^+$ MBCs previously identified
(Figure 2; Ehrhardt et al., 2005), we annotated two rare *CR2*/CD21^{low} MBC subsets resembling
populations described elsewhere (Lau et al., 2017, Thorarinsdottir et al., 2016) (Figures 5A,C and S4B).
However, the majority of MBC diversity within human tonsils appeared to reflect differences in cellular
state or signalling activity rather than distinct cell types, such as different activation states (Activated 1
and Activated 2), heat shock protein (HSP)-related gene activity (HSP-response), and an IFN-
responsive MBC state (IFN-response) (Figures 5C and S4C). We also identified a preGC MBC
population with similar gene expression to naïve preGC cells and an enrichment for class switch
recombination genes (Figures 5C and S4D-E), as well as an FCRL2/3^{high} MBC state similar to the
FCRL2/3^{high} GC population (Figures 5C and S4D), suggesting that these may be widely shared
functional states spanning multiple B cell fates.

We next considered whether class-switched and unswitched MBCs exhibit different gene expression
networks that might reflect unique functional abilities (Dogan et al., 2009). We found little evidence that
antibody class contributed towards the likelihood of an MBC to exist in a given state, with the exception
of FCRL2/3^{high} MBCs which, similar to FCRL2/3^{high} GC B cells, were enriched for IgM$^+$ cells (Figures 5B
and S4F). Intriguingly, key marker genes associated with this state were broadly up-regulated in IgM$^+$
cells across all MBC clusters (Figure S4G), suggesting a close relationship between the expression of
these genes and IgM expression. We therefore compared gene expression of switched and unswitched
MBCs of equivalent somatic hypermutation levels as a proxy for affinity (similar to our analysis of GC B
cells in Figure 4), as MBCs expressing switched isotypes tended to have higher somatic hypermutation
frequencies than unswitched MBCs (Figure S4H). We found widespread differences between
unswitched and switched MBCs that were independent of MBC subset or state (Figure 5D), indicating
a specific transcriptional wiring of switched and unswitched MBCs that might reflect altered abilities to
signal to other immune cell populations, proliferate, survive, differentiate, migrate and respond to
challenges (Figure 5E). In particular, we noted elevated expression in unswitched MBCs of genes
known to regulate B cell migration within lymphoid tissues and genes with the potential to signal to and
activate other immune cell types, including *IL6*, *CLEC2B*, *CR2* (CD21), *CKLF*, *S1PR1*, *CCR6*, and
SEMA7A (Figure 5F; Arkatkar et al., 2017, Suzuki et al., 2008, Elgueta et al., 2015, Cinamon et al.,
2004, Han et al., 2001, Welte et al., 2006). This could all contribute to the increased capacity of IgM$^+$
MBCs to re-initiate GC reactions as part of a recall memory response (Dogan et al., 2009, Lutz et al.,
2015, Seifert et al., 2015).

Figure 5. Diverse memory B cell states and antibody class-dependent gene expression.

A) Clustering and UMAP visualisation of 21,595 memory B cell (MBC) single-cell transcriptomes. Identified cell populations include multiple basal and activated states, MBCs enriched for heat shock protein (HSP)-response and interferon (IFN)-response genes, preGC MBCs, FCRL2/3high MBCs, tissue-resident FCRL4+ MBCs and two CD21low populations.

B) scVDJ-derived antibody isotypes of single MBC transcriptomes with high quality VDJ sequences (*n*=15,531 cells).

C) Mean expression of top marker genes for MBC states.

D) Average log fold change (logFC) of genes significantly enriched in switched or unswitched MBC states with similarly high affinity (based on SHM frequency). CD21low clusters had too few cells and were excluded.

E) Gene ontologies for genes significantly enriched in switched or unswitched MBCs with high affinity/SHM, in any cell state.

F) Average logFC of selected immunologically-relevant genes significantly enriched in switched or unswitched MBC populations with high affinity/SHM.

G) Single-cell scores for preGC and class switch recombination (CSR) signature gene sets in switched and unswitched preGC MBC with high affinity/SHM.

H) Single-cell gene expression of key class switched recombination genes in switched and unswitched preGC MBC with high affinity/SHM.

However, IgM+ MBCs were also enriched for many genes proposed to regulate or inhibit B cell activation, such as *FCRLA*, *FCRL2*, *FCRL3*, *CBLB*, *CD72* and *SIGLEC10* (Wu and Bondada, 2009, Sohn et al., 2003, Meyer et al., 2018, Kochi et al., 2009, Shabani et al., 2014), which may reflect a fine regulatory balance controlling their activation threshold. Perhaps underpinning this, unswitched MBCs also expressed higher levels of the transcription factor genes *POU2F2* (OCT2) and *FOXP1* than class-switched MBCs (Figure 5F), which coordinate the capacity of B cells to respond normally to antigen

receptor signals and directly repress key regulators of plasma cell differentiation respectively (van
Keimpema et al., 2015, Corcoran et al., 2014). This is consistent with switched IgG+ MBCs being more
likely to differentiate into plasma cells, while unswitched IgM+ MBCs are more likely to re-enter or form
secondary GC responses to gain higher affinity (Dogan et al., 2009, Lutz et al., 2015, Seifert et al.,
2015). Indeed, we found that unswitched preGC MBCs exhibited significantly higher expression of many
genes linked with the preGC state, including those associated with class switch recombination of their
antibody genes (Figure 5G-H). This indicates that unswitched MBCs are more primed to undergo class
switching for the first time as they re-enter the GC reaction, which will have an important impact on their
subsequent selection dynamics in the GC.

Discussion

Antibody responses are the foundation of effective immune memory and our ability to manipulate them
through vaccination has contributed significantly to the success of modern medicine. While it is known
that high affinity class-switched antibodies are generated during GC reactions, the complexity and
dynamic nature of this response has presented a significant challenge for those seeking to understand
how human B cell-mediated immunity is derived. By combining bulk antibody repertoire analysis with
single-cell transcriptomics we have generated a detailed resource of the human GC response in a
model secondary lymphoid tissue. This allowed us to define gene expression signatures of known and
novel B cell states, most notably a population primed to undergo class switch recombination before
entering the GC reaction. Whether a B cell undergoes class switch recombination at this stage then
influences their capacity to undergo antibody-based selection within the GC and secondary activation
as MBCs.

Although first histologically observed over a century ago, many questions remain about how B cells
enter, experience and exit the GC reaction (Mesin et al., 2016, Shlomchik et al., 2019, Cyster and Allen,
2019). The practical challenges of sequentially sampling patient lymphoid tissues during an immune
response make this especially true for understanding the most dynamic aspects of human B cell
maturation. In particular, understanding the early events that facilitate GC entry by human B cells could
provide new targets for adjuvants during vaccination or other immunotherapies. We have used
pseudotemporal ordering to map the gene expression dynamics of both the early stages of B cell
activation that correspond to antigen-dependent signalling through the BCR and the subsequent
transition to a transcriptionally distinct preGC state, the latter of which is presumably under the
regulation of cognate antigen-specific T helper cells. Our discovery that this preGC state is primed to
undergo class switch recombination supports mounting evidence that class switching occurs before the
classical GC response (Toellner et al., 1996, Roco et al., 2019, Pape et al., 2003) and has profound
implications for understanding antibody-based selection dynamics in the GC.

Within the GC, B cell survival and selection is dependent on antigen binding to the BCR and its
downstream signalling pathways (Kwak et al., 2019). By using single-cell transcriptomics paired with
BCR sequence analysis we have been able to uncouple antibody class, affinity and B cell phenotype
at single-cell resolution. We show that whether a B cell has undergone class switch recombination
during GC entry is a major determinant of that B cell's capacity to expand, acquire high antigen affinity
and differentiate into plasmablasts or MBCs. IgG$^+$ and IgA$^+$ GC B cells have gene expression patterns
consistent with increased BCR signalling and a greater capacity to remain within the GC, acquire T cell
help and undergo somatic hypermutation to increase their affinity than GC B cells that have retained
IgM expression. If class switch recombination does indeed occur predominantly prior to GC entry, as
we and others suggest, these data support a model whereby the ability of a B cell to acquire high affinity
is primarily dictated by the outcome of a specific class switch recombination "checkpoint" at the preGC
stage. This would explain our observation that switched MBCs have comparable somatic hypermutation
frequencies to switched plasmablasts, in contrast to the prevailing paradigm that higher affinity GC B
cells preferentially differentiate towards the plasmablast fate whereas lower affinity clones seed the
memory compartment (Suan et al., 2017b, Phan et al., 2006, Shinnakasu et al., 2016). The differences
in affinity between these populations may instead be explained by the likelihood of whether they
retained IgM expression prior to entering affinity maturation in the GC. This class switch recombination
"checkpoint" is also relevant during the secondary activation of MBCs, where IgM$^+$ MBCs appear more
likely to undergo class switching compared to IgG$^+$ or IgA$^+$ MBCs, which may be important to provide a
higher affinity secondary response through more prolonged GC maturation as well as more diverse
effector functions from class-switched antibodies. Finally, our discovery of a conserved "unswitched"
signature of elevated FCRL2 and FCRL3 expression (amongst other genes) across different B cell
states, raises interesting questions about how these genes might regulate B cell function and immune
responses more broadly.

Although the direct mechanisms shaping these gene expression differences between class-specific B
cells remain to be elucidated, variations in the immunoglobulin tail tyrosine domain, linker flexibility or
glycosylation sites between IgM and other antibody isotypes may all contribute to differential BCR
signalling that could shape the behaviour of class-specific B cells (Martin and Goodnow, 2002, Engels
et al., 2014, Xu et al., 2014b). Of note, we did not identify many major or meaningful differences in gene
expression between IgG$^+$ and IgA$^+$ B cells, or between subclass-specific B cells, which may reflect the
need to increase the power of future studies to identify potentially subtle gene expression differences
between the less abundant isotypes. Unfortunately, we did not identify any IgE$^+$ GC B cells, which are
very rare in human tonsils, likely as they have been described to rapidly exit GCs due to IgE-specific
BCR signalling (Haniuda et al., 2016). Given their relative availability, human tonsils are a useful tissue
with which to examine the GC response. It will be of interest for future studies to compare class-specific
gene expression differences in other tissues and contexts to determine the contribution of the local
cellular environment on B cell maturation (James et al., 2020, Smillie et al., 2019). Similarly, recent

technical advances allowing the simultaneous readout of antigen specificity, antibody sequences and
gene expression (Setliff et al., 2019) make it possible to examine how different types of antigens may
be linked with different B cell phenotypes. Finally, our profiling of human MBCs in a secondary lymphoid
tissue revealed diverse states reflecting different activation, signalling and functional potential. Given
an emerging appreciation for heterogeneity within both human and mouse MBC populations (Good-
Jacobson and Shlomchik, 2010), our single-cell characterisation of different MBC states will act as a
valuable resource to interrogate the potential relevance for such diverse populations in mediating
humoral immunity.

Together, our integrated analyses of antibody repertoires and gene expression of human B cell states
during affinity maturation highlight how the outcome of class switch recombination is a major
determinant of B cell fate and function. More broadly, our detailed annotation of diverse B cell states
provides a new and uniquely detailed framework through which to view B cell-mediated immune
responses in the context of both health and B cell-related pathologies such as allergy, multiple sclerosis,
rheumatoid arthritis and lymphoma.

528 **Methods**

529 Human ethics, tissue collection and sorting of B cell subsets

Routine tonsillectomy patients at the Royal London Hospital aged between 3 and 14 were consented
for tissue collection with approval from North West - Greater Manchester East Research Ethics
Committee under REC reference 17/NW/0664. Following removal of dead tissue and clotted blood,
each palatine tonsil was bisected and processed separately as follows, with all repertoire and single-
cell analyses performed on a single bisected sample. Tonsillar tissue was dissected manually into
approximately 2-3 mm pieces prior to homogenisation with the gentleMACS^{TM} Dissociator using C tubes
and two rounds of the Multi_C_01_01 setting in 8 mL RPMI + 10% fetal calf serum (FCS). Dissociated
cells were then passed through a 70 µM filter prior to isolation of mononuclear lymphocytes using Ficoll-
Paque^{TM} gradients. Isolated mononuclear cells were then washed in RPMI + 10% FCS before cell
counting and viability determination using Trypan Blue staining. Cells to be used for 10X single-cell
transcriptomics were processed immediately, while cells for bulk repertoire sequencing were either
stored overnight at 4°C or cryopreserved in FCS with 10% DMSO at -70°C.

For bulk B cell repertoires, we labelled 1-1.3×10^8 tonsillar lymphocytes per donor for fluorescence-
activated cell sorting (FACS). Briefly, cells were washed and stained with Zombie NIR™ Fixable
Viability Kit (BioLegend) to label dead cells, followed by washing with FACS buffer (PBS + 0.5 % BSA
+ 2 mM EDTA) and incubation with human FcR Blocking Reagent (Miltentyi Biotec). Cells were then
stained with CD19-APC (clone HIB19; BioLegend), CD38-PE-Cy7 (clone HB-7; BioLegend), CD27-
Pacific Blue^{TM} (clone O323; BioLegend), IgD-PerCP-Cy5.5 (clone IA6-2; BioLegend), and IgM-FITC

(clone MHM-88; BioLegend). For bulk B cell repertoires, two aliquots of 250,000 cells for the following

populations were sorted using a BD FACSAria™ IIIu: total B cells (live CD19$^+$), naïve B cells (live CD19$^+$

IgD$^+$ CD38$^-$), germinal centre B cells (live CD19$^+$ IgD$^-$ CD38$^+$), memory B cells (live CD19$^+$ IgD$^-$ CD38$^-$),

and plasmablasts (live CD19$^+$ IgD$^-$ CD38^{++}). Gates were set using fluorescence minus one (FMO)

controls. Sorted B cell samples were processed immediately for RNA extraction. For single-cell RNA-

seq, 50,000-200,000 class-switched memory B cells (live CD19$^+$ IgD$^-$ CD38$^-$ IgM$^+$ (n=2) or IgM$^-$ (n=4))

were sorted.

Bulk VDJ repertoire library preparation and sequencing

RNA was isolated from sorted B cell aliquots using the RNAqueous™-Micro Total RNA Isolation Kit

(ThermoScientific) supplemented with β-mercaptoethanol according to manufacturer's protocol. RNA

was stored long-term at -80°C or processed immediately to generate bulk repertoire sequencing

libraries of immunoglobulin heavy chains (IgH) as previously described (Horns et al., 2016), with minor

changes. Briefly, 50 to 100 ng RNA from sorted B cell subsets were annealed to a pooled set of five

isotype-specific IgH constant region primers containing unique molecular identifiers (UMIs) of either 10

or 12 nucleotides at 72°C for 5 minutes before being immediately placed on ice for 2 minutes. We then

performed first-strand cDNA synthesis using SuperScript IV reverse transcriptase (ThermoFisher

Scientific) with recommended reagent concentrations and the following cycling conditions in a

thermocycler: 105°C lid; 55°C 10 minutes; 80°C 10 minutes; 4°C hold. Second-strand cDNA synthesis

was performed using Phusion® High-Fidelity DNA Polymerase (NEB) and six IgH variable region

primers containing 10 or 12 nucleotide UMIs with the following cycling conditions: 105°C lid; 98°C 4

minutes; 52°C 1 minutes; 72°C 5 minutes; 4°C hold. Double-stranded cDNA was then purified using

(0.6X) Ampure XP beads (Beckman Coulter) before amplification with Illumina adapter-containing

primers (Nextera i7 indices) and NEBNext Ultra II Q5 Master Mix (NEB) as follows: 105°C lid; 98°C 30

seconds; (98°C 10 seconds, 72°C 50 seconds) × 22 to 28 cycles; 72°C 2 minutes; 4°C hold. Amplified

libraries were purified using (0.6X) Ampure XP beads and quantified by Qubit™ dsDNA HS Assay Kit

prior to multiplexing. Libraries were sequenced with PhiX spike-in using paired-end 301 bp reads on

the Illumina MiSeq platform.

10x Genomics Chromium single-cell transcriptomics and VDJ library preparation, sequencing and raw data processing

Total tonsillar immune cells (n=7) or FACS-enriched memory B cells (n=6) were loaded according to

the manufacturer's protocol for either for the Chromium single-cell 3' kit (v2; n = 1) or 5' gene expression

(v1; n = 6) to attain between 2000-5000 cells per well. Library preparation for both gene expression and

VDJ (BCR) was performed according to the manufacturer's protocol prior to sequencing on the Illumina

NextSeq 500 platform with 26/8/134 bp or 155/8/155 bp read configurations respectively. BaseCall files

were subsequently used to generate library-specific FASTQ files with cellranger mkfastq (v3.0.0) prior

to running cellranger count (v3.0.0) with the GRCh38 (release 92) reference to produce cell barcode-

gene expression matrices using default settings. For single-cell VDJ datasets, cellranger vdj (v3.0.0)

was run using the refdata-cellranger-vdj-GRCh38-alts-ensembl-2.0.0 reference from 10x Genomics

using default settings. Poor quality contigs that either did not map to immunoglobulin chains or were

designated incomplete by cellranger were discarded. We further filtered IgH contigs as to whether they

had sufficient coverage of constant regions to ensure accurate isotype assignment between closely

related subclasses using MaskPrimers.py (pRESTO v0.5.10; Vander Heiden et al., 2014).

Quality control and sequence assembly of bulk B cell repertoires

Raw sequencing data from bulk VDJ libraries were processed to generate UMI-collapsed consensus

VDJ sequences using pRESTO (v0.5.10; Vander Heiden et al., 2014). Paired-end sequencing reads

with mean Phred quality scores less than 25 were removed, and remaining sequences were annotated

and trimmed for PCR primer and UMI sequences. UMI barcodes were then filtered by length and the

presence of ambiguous nucleotides, prior to UMI alignment using MUSCLE (v3.8.31; Edgar, 2004).

UMI consensus sequences were then generated, with a minimum of three reads per UMI required, prior

to assembly of paired-end UMI consensus sequences into a single VDJ contig and annotation of

constant region isotype using MaskPrimers.py align (v0.5.10; Vander Heiden et al., 2014). Duplicate

VDJ sequences within each subset were then collapsed using CollapseSeq.py (v0.5.10; Vander Heiden

et al., 2014) before VDJ gene assignment and functional annotation using AssignGenes.py (ChangeO

v0.4.5; Gupta et al., 2015) and IgBLAST (v1.12.0; Ye et al., 2013).

Identification of clonally-related sequences, genotype inference and calculation of IgH

mutation frequencies.

Following initial quality control, all single-cell VDJ sequences were combined together with bulk BCR

repertoire sequences from the same donor for subsequent processing. IgH sequences were annotated

using IgBLAST (v1.12.0; Ye et al., 2013) and assigned isotype classes using AssignGenes.py prior to

correction of ambiguous V gene assignments using TIgGER (v0.3.1; Gupta et al., 2015, Gadala-Maria

et al., 2015). Clonally-related IgH sequences were identified using DefineClones.py (ChangeO v0.4.5;

Gupta et al., 2015) with a nearest neighbour distance threshold of 0.0818, as determined by the mean

99% confidence interval of all 8 donors with distToNearest (Shazam v0.1.11; Gupta et al., 2015).

CreateGermlines.py (ChangeO v0.4.5) was then used to infer germline sequences for each clonal

family and observedMutations (Shazam v0.1.11) was used to calculate somatic hypermutation

frequencies for each IgH sequence. Sequences with mutation frequencies greater than 0.02 were

annotated as "High" mutation levels, those between 0 and 0.02 as "Low" mutation levels and 0 as
"None". For bulk BCR repertoire analysis in Figure 1, single-cell VDJ sequences were excluded and
analysed, providing ~1.5 million high-confidence and unique IgH sequences, with a median of 14 UMIs
per sequence, a median of 28,918 unique sequences per donor per subset and approximately 96-99%
of these sequences appeared to be functional.

Single-cell VDJ analysis was performed broadly as described previously (James et al., 2020). Briefly,
the number of quality filtered and annotated IgH, IgK or IgL were determined per unique cell barcode
prior to integration with single-cell gene expression objects. If more than one contig per chain was
identified, metadata for that cell was ascribed as "Multi". IgH diversity analyses were performed using
the rarefyDiversity and testDiversity of Alakazam (v0.2.11; Gupta et al., 2015). To assess clonal
relationships between cell types, co-occurrence of expanded clone members between cell types was
reported as a binary event for each clone that contained a member within two different cell types in
either single-cell or bulk repertoires. For comparisons of somatic hypermutation and isotype frequencies
between subsets we used Wilcoxon Rank Sum Signed test while Student's t test was used to compare
mean clonal diversity scores.

Data quality control, processing and annotation of single-cell RNA-seq.

Gene expression count matrices from cellranger were used to calculate percentage mitochondrial
expression per cell barcode prior to mitochondrial genes being removed from gene expression matrices.
Similarly, the V, D and J gene counts for each immunoglobulin and T cell receptor were summed to
calculate an overall expression before individual genes were removed from gene expression matrices.
Counts of individual IgH constant region genes were also summed together (IgG1-4, and IgA1-A2) and
removed from gene expression matrices. Modified gene-by-cell matrices were then used to create
Seurat objects for each sample using Seurat (v3.0.3; Butler et al., 2018, Stuart et al., 2019), removing
genes expressed in fewer than 3 cells. Cell barcodes with >1000 and <60000 UMIs and >500 and
<7000 genes detected were removed, as were cell barcodes with >30% mitochondrial reads. Individual
samples were then log transformed, normalised by a factor of 10000 prior to predicting cell cycle phases
using the CellCycleScoring command and then identifying the 3000 most variable genes within each
sample using the "vst" method. We then performed a preliminary integration of all unsorted immune
cells or all sorted memory B cell datasets together using FindIntegrationAnchors and IntegrateData
(3000 genes) before regressing out cell cycle scores and mitochondrial gene expression, performing
principle component analysis (PCA) and preliminary clustering and cell type annotation. One cluster
was identified to be enriched with predicted doublets based on the results from DoubletFinder (v2.0.1;
McGinnis et al., 2019) and scrublet (v0.2; Wolock et al., 2019), and a small number of cell barcodes
with co-expression of B/T/non-lymphoid markers were manually removed by filtering on UMAP
coordinates. Following the removal of poor quality cell barcodes from gene expression matrices based

on these preliminary analyses of the unsorted immune cell and sorted memory B cell libraries, we then

integrated all normalised count matrices together using the unsorted immune cell count matrices as a

reference with 4000 highly variable genes before scaling the integrated data and regressing cell cycle

and mitochondrial gene expression, running PCA and identifying broad cell type lineages (B cell, T cell

and non-lymphoid cells) using a broad resolution for clustering. These lineages were then separated

for more detailed cell state annotation by recomputing the PCA (RunPCA), nearest neighbour graph

(FindNeighbors) and unbiased clustering (FindClusters). Uniform Manifold Approximation and

Projection (UMAP) was then used to visualise both integrated and lineage-specific datasets. B cells

were annotated with scVDJ metadata from the integrated repertoire analysis detailed above.

664   Differential gene expression and signature enrichment analysis.

Gene expression markers for different clusters of unsorted B cells, T cells, non-lymphoid cells and

sorted MBCs were identified using FindAllMarkers from Seurat with default settings, including Wilcoxon

test and Bonferroni *p* value correction (v3.0.3; Butler et al., 2018, Stuart et al., 2019). Differential gene

expression for antibody class-specific or somatic hypermutation frequency for GC B cells or class-

switched MBCs was performed using FindAllMarkers with Benjamini-Hochberg false discovery rate

(FDR) correction. Genes were deemed significantly different if FDR < 0.05, average log fold change >

0.1 and the gene was detected in >20% of cells in that group. Gene ontology analyses for high

affinity/SHM class-specific GC B cells were performed with Ingenuity Pathway Analysis (Qiagen)

software using avg_logFC values of all genes significantly enriched in at least one class. For IFN-

response MBC cluster gene ontology enrichment was performed with all significant gene markers for

this cluster in Metascape (Zhou et al., 2019) using default settings, as were all genes significantly

enriched or depleted in switched or unswitched in at least one MBC cluster.

Enrichment of gene set signatures for single cells was calculated using AUCell (v1.5.5; Aibar et al.,

2017). For class switch recombination, a manually curated shortlist of genes was determined for genes

linked with CSR that were reliably detected in our sparse scRNA-seq datasets (Stavnezer and

Schrader, 2014). Gene signatures of high and low affinity mouse LZ GC B cells (Shinnakasu et al.,

2016) were obtained by quantifying RNA-seq transcript counts against GRCm38 transcriptome build

using Salmon (v1.0.0; Patro et al., 2017), collapsing protein-coding transcripts into a single gene count

using tximport (v1.10.1; Soneson et al., 2016), identifying significant gene expression differences

between the two groups using DESeq2 (v1.22.2; Love et al., 2014) with a threshold of log fold change

> 1.5 and *padj* < 0.05 and converting mouse gene IDs to human using bioMart (v2.38.0; Durinck et al.,

2005, Durinck et al., 2009). For IgM stimulation gene sets, Geo2R (Barrett et al., 2012) was used to

analyse microarray data from a timecourse of wild type splenic mouse B cells stimulated with 10 μg/ml

of anti-IgM (Shinohara et al., 2014) and identify genes enriched following α-IgM treatment compared to

control untreated cells with FDR < 0.05 and a fold change > 2. To calculate preGC and FCRL2/3[high]

signature scores in MBC subsets, the top 50 most significantly enriched gene markers per cluster in
the unsorted B cell subset analysis were used with AUCell. Unless indicated otherwise, Wilcoxon
Ranked Signed Sum test was used to test for significant differences.

Prediction of cell-cell communication using CellPhoneDB.

To evaluate potential cell-cell communication, we used CellPhoneDB (v2.0.6; Vento-Tormo et al., 2018,
Efremova et al., 2020) to examine the expression of ligand-receptor pairs between different scRNA-seq
clusters. Briefly, we exported raw gene count matrices from Seurat, converted gene IDs to Ensembl
IDs using bioMart (v2.38.0; Durinck et al., 2005). We re-annotated all non-lymphoid cell type clusters
as antigen-presenting cells (APCs), naïve and effector T cell groups by CD4 or CD8 expression, Treg
and Tfr as "Treg" and rare GC subsets (prePB and FCRL2/3^{high}) as "GC" and exported cell type
metadata for use with raw count data using the "statistical_analysis" command of CellPhoneDB with
database v2.0.0. The number of unique significant ligand-receptor co-expression pairs (putative
interactions; p value < 0.05) between each cell type was then counted and visualised as a heatmap,
while exemplar interacting pairs were visualised by calculating mean average expression level of gene
1 in cell type 1 and gene 2 in cell type 2 are indicated by colour and p values indicated by circle size.

RNA velocity and pseudotemporal ordering.

To calculate single-cell velocities we first quantified spliced and unspliced transcripts for the filtered
barcodes output from cellranger using velocyto (v0.17.10; La Manno et al., 2018). Loom files were then
combined using loompy (v2.0.17) before reformatting cell barcode names to be compatible with Seurat
objects and merging with a Scanpy (v1.4; Wolf et al., 2018) object containing the raw gene expression
matrix of high quality annotated single B cell transcriptomes (see above) using scVelo (v0.1.23; Bergen
et al., 2019). scVelo was then used to pre-process, filter and normalise velocyto-derived counts with
default settings prior to computation of the first- and second-order moments (scv.pp.moments) and
subsequent velocity estimation using a dynamical model (scv.tl.recover_dynamics and scv.tl.velocity).
Velocities were then projected and visualised onto UMAP embeddings at a grid level using an inverted
transition matrix obtained from scv.tl.transition_matrix prior to scv.tl.velocity_embedding, basis='umap'.

For pseudotemporal ordering of the B cell activation and GC entry trajectory, a partition-based graph
abstraction (PAGA) was performed for Naïve, Activated, preGC and LZ GC B cell clusters (Scanpy;
tl.paga) before computing connectivity of single cells using a diffusion map (Scanpy; tl.diffmap).
Velocity-based pseudotime reconstruction was performed using default settings for scVelo commands
tl.recover_latent_time and tl.velocity_pseudotime, although the pseudotemporal order was reversed to
place naïve cluster at pseudotime = 0. Dynamic gene expression changes were examined by using
tl.rank_velocity_genes (scVelo) to sub-cluster original cell type annotations (resolution = 1) based on

RNA velocity and the top 200 genes per sub-cluster were reported before filtering out ribosomal genes
and collapsing to unique genes. Genes were then clustered through pseudotime for heatmap
visualisation with smoothed expression scores in scVelo. Quantitation of individual gene expression or
AUCell-derived signature scores of single-cells across pseudotime was performed using smoothed
normalised counts with geom_smooth() including 95% confidence intervals. For pseudotemporal
analysis of GC B cell subsets, a similar approach was taken for the LZ GC, GC, DZ GC, FCRL2/3^{high}
GC and prePB clusters, except that diffusion-based pseudotime was calculated with Scanpy (tl.dpt)
independent of RNA velocity measurements. Visualisation of a custom list of top GC subset marker
genes for LZ GC, GC and DZ GC clusters was performed using pl.paga.path heatmap with Scanpy.

Immunohistochemistry

Tonsil biopsies were cut from formalin fixed paraffin-embedded blocks then deparaffinized in xylene
and rehydrated through a series of ethanols to water. Endogenous peroxidase was blocked with 3%
hydrogen peroxide. Heat-mediated antigen retrieval was performed using a commercial citrate-based
unmasking buffer (Vector Labs) at 120°C using a pressure cooker. Sections (3μm thick) were then
incubated for 40 minutes at RT with 1:1000 dilution of anti-APE1 (HPA002564; Sigma). Detection of
primary antibody was performed using the Super-sensitive–Polymer HRP system (Biogenex) and
staining visualized using purple chromogen VIP (Vector Labs) and hematoxylin as a nuclear
counterstain. Slides were then scanned (Pannoramic 250 Flash) before being left to soak in xylene to
de-coverslip. Once the coverslips were removed, slides were rehydrated through ethanol to water. De-
staining and stripping of the primary antibodies and the heat-labile VIP chromogen was achieved using
a subsequent round of heat-mediated antigen retrieval as per the first round of staining. The second
primary antibody CD20 (M0755; Dako) was incubated for 40 minutes at RT at a dilution of 1:500,
followed by detection, visualization and scanning as before. Negative controls were performed by
treating sequential sections as above but without the second primary antibody (CD20) to confirm
complete antibody and signal stripping. Images were prepared using CaseViewer (3DHistTech).

Quantitation of IgH germline transcripts.

To quantify expression of IgH germline transcripts (GLT), all mapped reads to the IgH locus
(chr14:105540180-105879151) were extracted from cellranger-derived bam files. DropEst (v0.8.6;
Petukhov et al., 2018) was then used to count reads against a custom GTF containing coordinates for
I promoter GLT sequences (Sideras et al., 1989, Fujieda et al., 1996), annotation of membrane-specific
IgH exons and IgH switch regions identified by enrichment of the WRCY motif using HOMER2 (v4.9.1;
Heinz et al., 2010). Counts were then read into Seurat without filtering for log10-normalisation and
scaling. Due to mapping ambiguities between subclass-specific regions, subclass-specific counts were
summed together.

Accession codes and data availability

Raw sequencing and processed data files for single-cell RNA sequencing, single-cell VDJ sequencing, and bulk B cell repertoires are available at ArrayExpress (accession numbers: E-MTAB-8999, E-MTAB-9003 and E-MTAB-9005). RNA-seq and microarray data for high affinity and α-IgM stimulation gene expression signatures were obtained from GSE73729 and GSE41176 respectively. DNase-seq and ChIP-seq datasets from ENCODE were visualised using UCSC Genome Browser (Kent et al., 2002, Rosenbloom et al., 2012).

Acknowledgements

This research was supported by funding from the Wellcome Trust to L. K. J. (208961/Z/17/Z), S. A. T. (206194) and J. C. R. (110020/Z/15/Z). H. W. K. was funded by a Sir Henry Wellcome PostDoctoral Fellowship (213555/Z/18/Z). We would like to thank the Bart's and the London Genome Centre at QMUL for library sequencing support. We also thank members of the James and Teichmann labs for their help and support, especially Mirjana Efremova for her help with CellPhoneDB. Finally, we are grateful to Neil McCarthy, Kylie James, Kerstin Meyer, Lou Herman and Jo Spencer for their help reviewing the manuscript.

Author contributions

H. W. K. initiated the project, designed and performed experiments, analysed data and wrote the manuscript. N. O. carried out tonsillectomy tissue collection. J. C. R. and A. J. C. designed and performed immunohistochemistry experiments. G. W. assisted in FACS sorting. S. A. T. supervised and interpreted data analysis. L. K. J. initiated the project, designed and supervised experiments and data analysis, interpreted data and wrote the manuscript.

Conflict of Interest Statement

In the past three years, S.A.T has worked as a consultant for Genentech, Biogen and Roche, and is a remunerated member of the Foresite Labs Scientific Advisory Board.

References

Aibar, S., González-Blas, C. B., Moerman, T., Huynh-Thu, V. A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J.-C., Geurts, P., Aerts, J*., et al.* 2017. SCENIC: single-cell regulatory network inference and clustering. *Nat Methods,* 14**,** 1083-1086.

Arana, E., Harwood, N. E. & Batista, F. D. 2008. Regulation of integrin activation through the B-cell receptor. *J Cell Sci,* 121**,** 2279-2286.

Arkatkar, T., Du, S. W., Jacobs, H. M., Dam, E. M., Hou, B., Buckner, J. H., Rawlings, D. J. & Jackson, S. W. 2017. B cell–
derived IL-6 initiates spontaneous germinal center formation during systemic autoimmunity. *J Exp Med,* 214**,** 3207-3217.
Barrett, T., Wilhite, S. E., Ledoux, P., Evangelista, C., Kim, I. F., Tomashevsky, M., Marshall, K. A., Phillippy, K. H., Sherman,
P. M., Holko, M*., et al.* 2012. NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids Res,* 41**,**
D991-D995.
Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. 2019. Generalizing RNA velocity to transient cell states through
dynamical modeling. *bioRxiv***,** 820936; doi:10.1101/820936.
Borggrefe, T., Wabl, M., Akhmedov, A. T. & Jessberger, R. 1998. A B-cell-specific DNA Recombination Complex. *J Biol Chem,*
273**,** 17025-17035.
Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. 2018. Integrating single-cell transcriptomic data across different
conditions, technologies, and species. *Nat Biotechnol,* 36**,** 411-420.
Cinamon, G., Matloubian, M., Lesneski, M. J., Xu, Y., Low, C., Lu, T., Proia, R. L. & Cyster, J. G. 2004. Sphingosine 1-
phosphate receptor 1 promotes B cell localization in the splenic marginal zone. *Nat Immunol,* 5**,** 713-20.
Corcoran, L., Emslie, D., Kratina, T., Shi, W., Hirsch, S., Taubenheim, N. & Chevrier, S. 2014. Oct2 and Obf1 as Facilitators
of B:T Cell Collaboration during a Humoral Immune Response. *Front Immunol,* 5**,** 108.
Cyster, J. G. & Allen, C. D. C. 2019. B Cell Responses: Cell Interaction Dynamics and Decisions. *Cell,* 177**,** 524-540.
Dogan, I., Bertocci, B., Vilmont, V., Delbos, F., Megret, J., Storck, S., Reynaud, C. A. & Weill, J. C. 2009. Multiple layers of B
cell memory with different effector functions. *Nat Immunol,* 10**,** 1292-9.
Dubovsky, J. A., Chappell, D. L., Harrington, B. K., Agrawal, K., Andritsos, L. A., Flynn, J. M., Jones, J. A., Paulaitis, M. E.,
Bolon, B., Johnson, A. J*., et al.* 2013. Lymphocyte cytosolic protein 1 is a chronic lymphocytic leukemia membrane-
associated antigen critical to niche homing. *Blood,* 122**,** 3308-3316.
Duffy, K. R., Wellard, C. J., Markham, J. F., Zhou, J. H., Holmberg, R., Hawkins, E. D., Hasbold, J., Dowling, M. R. & Hodgkin,
P. D. 2012. Activation-induced B cell fates are selected by intracellular stochastic competition. *Science,* 335**,** 338-41.
Durinck, S., Moreau, Y., Kasprzyk, A., Davis, S., De Moor, B., Brazma, A. & Huber, W. 2005. BioMart and Bioconductor: a
powerful link between biological databases and microarray data analysis. *Bioinformatics,* 21**,** 3439-40.
Durinck, S., Spellman, P. T., Birney, E. & Huber, W. 2009. Mapping identifiers for the integration of genomic datasets with the
R/Bioconductor package biomaRt. *Nat Protoc,* 4**,** 1184-91.
Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res,* 32**,**
1792-7.
Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. 2020. CellPhoneDB: inferring cell–cell communication
from combined expression of multi-subunit ligand–receptor complexes. *Nat Protoc,* 15**,** 1484–1506.
Ehrhardt, G. R., Hsu, J. T., Gartland, L., Leu, C. M., Zhang, S., Davis, R. S. & Cooper, M. D. 2005. Expression of the
immunoregulatory molecule FcRH4 defines a distinctive tissue-based population of memory B cells. *J Exp Med,* 202**,**
783-91.
Elgueta, R., Marks, E., Nowak, E., Menezes, S., Benson, M., Raman, V. S., Ortiz, C., O'connell, S., Hess, H., Lord, G. M*., et*
al. 2015. CCR6-dependent positioning of memory B cells is essential for their ability to mount a recall response to antigen.
J Immunol, 194**,** 505-513.
Engels, N., Konig, L. M., Schulze, W., Radtke, D., Vanshylla, K., Lutz, J., Winkler, T. H., Nitschke, L. & Wienands, J. 2014.
The immunoglobulin tail tyrosine motif upgrades memory-type BCRs by incorporating a Grb2-Btk signalling module. *Nat*
Commun, 5**,** 5456.
Engels, N. & Wienands, J. 2018. Memory control by the B cell antigen receptor. *Immunol Rev,* 283**,** 150-160.
Ferreira, R. C., Pan-Hammarström, Q., Graham, R. R., Gateva, V., Fontán, G., Lee, A. T., Ortmann, W., Urcelay, E.,
Fernández-Arquero, M., Núñez, C*., et al.* 2010. Association of IFIH1 and other autoimmunity risk alleles with selective
IgA deficiency. *Nat Genet,* 42**,** 777-780.
Fujieda, S., Lin, Y. Q., Saxon, A. & Zhang, K. 1996. Multiple types of chimeric germ-line Ig heavy chain transcripts in human
B cells: evidence for trans-splicing of human Ig RNA. *J Immunol,* 157**,** 3450-3459.
Gadala-Maria, D., Yaari, G., Uduman, M. & Kleinstein, S. H. 2015. Automated analysis of high-throughput B-cell sequencing
data reveals a high frequency of novel immunoglobulin V gene segment alleles. *PNAS,* 112**,** E862-70.
Girón-Pérez, D. A., Vadillo, E., Schnoor, M. & Santos-Argumedo, L. 2020. Myo1e modulates the recruitment of activated B
cells to inguinal lymph nodes. *J Cell Sci,* 133**,** jcs235275.
Gloury, R., Zotos, D., Zuidscherwoude, M., Masson, F., Liao, Y., Hasbold, J., Corcoran, L. M., Hodgkin, P. D., Belz, G. T., Shi,
W*., et al.* 2016. Dynamic changes in Id3 and E-protein activity orchestrate germinal center and plasma cell development.
J Exp Med, 213**,** 1095-1111.
Good-Jacobson, K. L. & Shlomchik, M. J. 2010. Plasticity and Heterogeneity in the Generation of Memory B Cells and Long-
Lived Plasma Cells: The Influence of Germinal Center Interactions and Dynamics. *J Immunol,* 185**,** 3117-3125.
Gupta, N. T., Vander Heiden, J. A., Uduman, M., Gadala-Maria, D., Yaari, G. & Kleinstein, S. H. 2015. Change-O: a toolkit for
analyzing large-scale B cell immunoglobulin repertoire sequencing data. *Bioinformatics,* 31**,** 3356-3358.
Han, W., Lou, Y., Tang, J., Zhang, Y., Chen, Y., Li, Y., Gu, W., Huang, J., Gui, L., Tang, Y*., et al.* 2001. Molecular cloning and
characterization of chemokine-like factor 1 (CKLF1), a novel human cytokine with unique structure and potential
chemotactic activity. *Biochem J,* 357**,** 127-35.
Hanakahi, L. A., Dempsey, L. A., Li, M.-J. & Maizels, N. 1997. Nucleolin is one component of the B cell-specific transcription
factor and switch region binding protein, LR1. *PNAS,* 94**,** 3605-3610.
Haniuda, K., Fukao, S., Kodama, T., Hasegawa, H. & Kitamura, D. 2016. Autonomous membrane IgE signaling prevents IgE-
memory formation. *Nat Immunol,* 17**,** 1109-17.
Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H. & Glass, C. K. 2010.
Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage
and B cell identities. *Mol Cell,* 38**,** 576-89.

Horns, F., Vollmers, C., Croote, D., Mackey, S. F., Swan, G. E., Dekker, C. L., Davis, M. M. & Quake, S. R. 2016. Lineage
tracing of human B cells reveals the in vivo landscape of human antibody class switching. *eLife,* 5**,** e16578.
Huang, C., Geng, H., Boss, I., Wang, L. & Melnick, A. 2014. Cooperative transcriptional repression by BCL6 and BACH2 in
germinal center B-cell differentiation. *Blood,* 123**,** 1012-20.
Ise, W., Fujii, K., Shiroguchi, K., Ito, A., Kometani, K., Takeda, K., Kawakami, E., Yamashita, K., Suzuki, K., Okada, T.*, et al.*
2018. T Follicular Helper Cell-Germinal Center B Cell Interaction Strength Regulates Entry into Plasma Cell or Recycling
Germinal Center Cell Fate. *Immunity,* 48**,** 702-715.e4.
James, K. R., Gomes, T., Elmentaite, R., Kumar, N., Gulliver, E. L., King, H. W., Stares, M. D., Bareham, B. R., Ferdinand, J.
R., Petrova, V. N.*, et al.* 2020. Distinct microbial and immune niches of the human colon. *Nat Immunol,* 21**,** 343-353.
Kent, W. J., Sugnet, C. W., Furey, T. S., Roskin, K. M., Pringle, T. H., Zahler, A. M., Haussler & David 2002. The Human
Genome Browser at UCSC. *Genome Res,* 12**,** 996-1006.
Kochi, Y., Myouzen, K., Yamada, R., Suzuki, A., Kurosaki, T., Nakamura, Y. & Yamamoto, K. 2009. FCRL3, an autoimmune
susceptibility gene, has inhibitory potential on B-cell receptor-mediated signaling. *J Immunol,* 183**,** 5502-10.
Kwak, K., Akkaya, M. & Pierce, S. K. 2019. B cell signaling in context. *Nat Immunol,* 20**,** 963-969.
La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K., Kastriti, M. E., Lönnerberg,
P., Furlan, A.*, et al.* 2018. RNA velocity of single cells. *Nature,* 560**,** 494-498.
Lau, D., Lan, L. Y.-L., Andrews, S. F., Henry, C., Rojas, K. T., Neu, K. E., Huang, M., Huang, Y., Dekosky, B., Palm, A.-K. E.*,*
et al. 2017. Low CD21 expression defines a population of recent germinal center graduates primed for plasma cell
differentiation. *Sci Immunol,* 2**,** eaai8153.
Love, M. I., Huber, W. & Anders, S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.
Genome Biol, 15**,** 550.
Lutz, J., Dittmann, K., Bosl, M. R., Winkler, T. H., Wienands, J. & Engels, N. 2015. Reactivation of IgG-switched memory B
cells by BCR-intrinsic signal amplification promotes IgG antibody production. *Nat Commun,* 6**,** 8575.
Madissoon, E., Wilbrey-Clark, A., Miragaia, R. J., Saeb-Parsy, K., Mahbubani, K. T., Georgakopoulos, N., Harding, P.,
Polanski, K., Huang, N., Nowicki-Osuch, K.*, et al.* 2019. scRNA-seq assessment of the human lung, spleen, and
esophagus tissue stability after cold preservation. *Genome Biol,* 21**,** 1.
Martin, S. W. & Goodnow, C. C. 2002. Burst-enhancing role of the IgG membrane tail as a molecular determinant of memory.
Nat Immunol, 3**,** 182-8.
Masani, S., Han, L. & Yu, K. 2013. Apurinic/Apyrimidinic Endonuclease 1 Is the Essential Nuclease during Immunoglobulin
Class Switch Recombination. *Mol Cell Biol,* 33**,** 1468-1473.
Mcginnis, C. S., Murrow, L. M. & Gartner, Z. J. 2019. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data
Using Artificial Nearest Neighbors. *Cell Systems,* 8**,** 329-337.e4.
Mcrae, E. K. S., Booy, E. P., Moya-Torres, A., Ezzati, P., Stetefeld, J. & Mckenna, S. A. 2017. Human DDX21 binds and
unwinds RNA guanine quadruplexes. *Nucleic Acids Res,* 45**,** 6656-6668.
Mesin, L., Ersching, J. & Victora, G. D. 2016. Germinal Center B Cell Dynamics. *Immunity,* 45**,** 471-482.
Meyer, S. J., Linder, A. T., Brandl, C. & Nitschke, L. 2018. B Cell Siglecs-News on Signaling and Its Interplay With Ligand
Binding. *Front Immunol,* 9**,** 2820-2820.
Mielke, N., Schwarzer, R., Calkhoven, C. F., Kaufman, R. J., Dörken, B., Leutz, A. & Jundt, F. 2011. Eukaryotic initiation factor
2alpha phosphorylation is required for B-cell maturation and function in mice. *Haematologica,* 96**,** 1261-1268.
Milpied, P., Cervera-Marzal, I., Mollichella, M.-L., Tesson, B., Brisou, G., Traverse-Glehen, A., Salles, G., Spinelli, L. & Nadel,
B. 2018. Human germinal center transcriptional programs are de-synchronized in B cell lymphoma. *Nat Immunol,* 19**,**
1013-1024.
Mondal, S., Begum, N. A., Hu, W. & Honjo, T. 2016. Functional requirements of AID's higher order structures and their
interaction with RNA-binding proteins. *PNAS,* 113**,** E1545-E1554.
Morman, R. E., Schweickert, P. G., Konieczny, S. F. & Taparowsky, E. J. 2018. BATF regulates the expression of Nfil3,
Wnt10a and miR155hg for efficient induction of antibody class switch recombination in mice. *Eur J Immunol,* 48**,** 1492-
1505.
Muppidi, J. R., Schmitz, R., Green, J. A., Xiao, W., Larsen, A. B., Braun, S. E., An, J., Xu, Y., Rosenwald, A., Ott, G.*, et al.*
2014. Loss of signalling via Galpha13 in germinal centre B-cell-derived lymphoma. *Nature,* 516**,** 254-8.
Orthwein, A., Patenaude, A.-M., Affar, E. B., Lamarre, A., Young, J. C. & Di Noia, J. M. 2010. Regulation of activation-induced
deaminase stability and antibody gene diversification by Hsp90. *J Exp Med,* 207**,** 2751-2765.
Pape, K. A., Kouskoff , V., Nemazee , D., Tang , H. L., Cyster , J. G., Tze , L. E., Hippen , K. L., Behrens , T. W. & Jenkins ,
M. K. 2003. Visualization of the Genesis and Fate of Isotype-switched B Cells during a Primary Immune Response. *J*
Exp Med, 197**,** 1677-1687.
Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. 2017. Salmon provides fast and bias-aware quantification of
transcript expression. *Nat Methods,* 14**,** 417-419.
Petukhov, V., Guo, J., Baryawno, N., Severe, N., Scadden, D. T., Samsonova, M. G. & Kharchenko, P. V. 2018. dropEst:
pipeline for accurate estimation of molecular counts in droplet-based single-cell RNA-seq experiments. *Genome Biol,* 19**,**
78.
Phan, T. G., Paus, D., Chan, T. D., Turner, M. L., Nutt, S. L., Basten, A. & Brink, R. 2006. High affinity germinal center B cells
are actively selected into the plasma cell compartment. *J Exp Med,* 203**,** 2419-24.
Roco, J. A., Mesin, L., Binder, S. C., Nefzger, C., Gonzalez-Figueroa, P., Canete, P. F., Ellyard, J., Shen, Q., Robert, P. A.,
Cappello, J.*, et al.* 2019. Class-Switch Recombination Occurs Infrequently in Germinal Centers. *Immunity,* 51**,** 337-
350.e7.
Rosenbloom, K. R., Sloan, C. A., Malladi, V. S., Dreszer, T. R., Learned, K., Kirkup, V. M., Wong, M. C., Maddren, M., Fang,
R., Heitner, S. G.*, et al.* 2012. ENCODE Data in the UCSC Genome Browser: year 5 update. *Nucleic Acids Res,* 41**,**
D56-D63.

Scharenberg, A. M., Humphries, L. A. & Rawlings, D. J. 2007. Calcium signalling and cell-fate choice in B cells. *Nat Rev*
Immunol, 7**,** 778-789.
Seifert, M., Przekopowitz, M., Taudien, S., Lollies, A., Ronge, V., Drees, B., Lindemann, M., Hillen, U., Engler, H., Singer, B.
B.*, et al.* 2015. Functional capacities of human IgM memory B cells in early inflammatory responses and secondary
germinal center reactions. *PNAS,* 112**,** E546-55.
Setliff, I., Shiakolas, A. R., Pilewski, K. A., Murji, A. A., Mapengo, R. E., Janowska, K., Richardson, S., Oosthuysen, C., Raju,
N., Ronsard, L.*, et al.* 2019. High-Throughput Mapping of B Cell Receptor Sequences to Antigen Specificity. *Cell,* 179**,**
1636-1646.e15.
Shabani, M., Bayat, A. A., Jeddi-Tehrani, M., Rabbani, H., Hojjat-Farsangi, M., Ulivieri, C., Amirghofran, Z., Baldari, C. T. &
Shokri, F. 2014. Ligation of human Fc receptor like-2 by monoclonal antibodies down-regulates B-cell receptor-mediated
signalling. *Immunology,* 143**,** 341-53.
Shinnakasu, R., Inoue, T., Kometani, K., Moriyama, S., Adachi, Y., Nakayama, M., Takahashi, Y., Fukuyama, H., Okada, T.
& Kurosaki, T. 2016. Regulated selection of germinal-center cells into the memory B cell compartment. *Nat Immunol,* 17**,**
861-869.
Shinohara, H., Behar, M., Inoue, K., Hiroshima, M., Yasuda, T., Nagashima, T., Kimura, S., Sanjo, H., Maeda, S., Yumoto, N.*,*
et al. 2014. Positive Feedback Within a Kinase Signaling Complex Functions as a Switch Mechanism for NF-κB
Activation. *Science,* 344**,** 760-764.
Shinozaki, F., Minami, M., Chiba, T., Suzuki, M., Yoshimatsu, K., Ichikawa, Y., Terasawa, K., Emori, Y., Matsumoto, K.,
Kurosaki, T.*, et al.* 2006. Depletion of hsp90beta induces multiple defects in B cell receptor signaling. *J Biol Chem,* 281**,**
16361-9.
Shlomchik, M. J., Luo, W. & Weisel, F. 2019. Linking signaling and selection in the germinal center. *Immunol Rev,* 288**,** 49-63.
Sideras, P., Mizuta, T. R., Kanamori, H., Suzuki, N., Okamoto, M., Kuze, K., Ohno, H., Doi, S., Fukuhara, S., Hassan, M. S.*,*
et al. 1989. Production of sterile transcripts of C gamma genes in an IgM-producing human neoplastic B cell line that
switches to IgG-producing cells. *Int Immunol,* 1**,** 631-42.
Smillie, C. S., Biton, M., Ordovas-Montanes, J., Sullivan, K. M., Burgin, G., Graham, D. B., Herbst, R. H., Rogel, N., Slyper,
M., Waldman, J.*, et al.* 2019. Intra- and Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis. *Cell,* 178**,**
714-730.e22.
Sohn, H. W., Gu, H. & Pierce, S. K. 2003. Cbl-b negatively regulates B cell antigen receptor signaling in mature B cells through
ubiquitination of the tyrosine kinase Syk. *J Exp Med,* 197**,** 1511-1524.
Soneson, C., Love, M. & Robinson, M. 2016. Differential analyses for RNA-seq: transcript-level estimates improve gene-level
inferences [version 2; peer review: 2 approved]. *F1000Research,* 4.
Stavnezer, J. & Schrader, C. E. 2014. IgH Chain Class Switch Recombination: Mechanism and Regulation. *J Immunol,* 193**,**
5370-5378.
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., 3rd, Hao, Y., Stoeckius, M., Smibert, P. &
Satija, R. 2019. Comprehensive Integration of Single-Cell Data. *Cell,* 177**,** 1888-1902.e21.
Suan, D., Kräutler, N. J., Maag, J. L. V., Butt, D., Bourne, K., Hermes, J. R., Avery, D. T., Young, C., Statham, A., Elliott, M.*,*
et al. 2017a. CCR6 Defines Memory B Cell Precursors in Mouse and Human Germinal Centers, Revealing Light-Zone
Location and Predominant Low Antigen Affinity. *Immunity,* 47**,** 1142-1153.e4.
Suan, D., Sundling, C. & Brink, R. 2017b. Plasma cell and memory B cell differentiation from the germinal center. *Curr Opin*
Immunol, 45**,** 97-102.
Suzuki, K., Kumanogoh, A. & Kikutani, H. 2008. Semaphorins and their receptors in immune cell interactions. *Nat Immunol,*
9**,** 17-23.
Thai, T.-H., Calado, D. P., Casola, S., Ansel, K. M., Xiao, C., Xue, Y., Murphy, A., Frendewey, D., Valenzuela, D., Kutok, J.
L.*, et al.* 2007. Regulation of the Germinal Center Response by MicroRNA-155. *Science,* 316**,** 604-608.
Thorarinsdottir, K., Camponeschi, A., Cavallini, N., Grimsholm, O., Jacobsson, L., Gjertsson, I. & Mårtensson, I. L. 2016.
CD21(-/low) B cells in human blood are memory cells. *Clin Exp Immunol,* 185**,** 252-262.
Todd, D. J., Mcheyzer-Williams, L. J., Kowal, C., Lee, A. H., Volpe, B. T., Diamond, B., Mcheyzer-Williams, M. G. & Glimcher,
L. H. 2009. XBP1 governs late events in plasma cell differentiation and is not required for antigen-specific memory B cell
development. *J Exp Med,* 206**,** 2151-9.
Toellner, K. M., Gulbranson-Judge, A., Taylor, D. R., Sze, D. M. & Maclennan, I. C. 1996. Immunoglobulin switch transcript
production in vivo related to the site and time of antigen-specific B cell activation. *J Exp Med,* 183**,** 2303-12.
Tybulewicz, V. L. J. & Henderson, R. B. 2009. Rho family GTPases and their regulators in lymphocytes. *Nat Rev Immunol,* 9**,**
630-644.
Van Keimpema, M., Grüneberg, L. J., Mokry, M., Van Boxtel, R., Van Zelm, M. C., Coffer, P., Pals, S. T. & Spaargaren, M.
2015. The forkhead transcription factor FOXP1 represses human plasma cell differentiation. *Blood,* 126**,** 2098-2109.
Vander Heiden, J. A., Yaari, G., Uduman, M., Stern, J. N. H., O'connor, K. C., Hafler, D. A., Vigneault, F. & Kleinstein, S. H.
2014. pRESTO: a toolkit for processing high-throughput sequencing raw reads of lymphocyte receptor repertoires.
Bioinformatics, 30**,** 1930-1932.
Vento-Tormo, R., Efremova, M., Botting, R. A., Turco, M. Y., Vento-Tormo, M., Meyer, K. B., Park, J.-E., Stephenson, E.,
Polański, K., Goncalves, A.*, et al.* 2018. Single-cell reconstruction of the early maternal–fetal interface in humans. *Nature,*
563**,** 347-353.
Vigorito, E., Perks, K. L., Abreu-Goodger, C., Bunting, S., Xiang, Z., Kohlhaas, S., Das, P. P., Miska, E. A., Rodriguez, A.,
Bradley, A.*, et al.* 2007. microRNA-155 Regulates the Generation of Immunoglobulin Class-Switched Plasma Cells.
Immunity, 27**,** 847-859.
Welte, S., Kuttruff, S., Waldhauer, I. & Steinle, A. 2006. Mutual activation of natural killer cells and monocytes mediated by
NKp80-AICL interaction. *Nat Immunol,* 7**,** 1334-1342.

Willis, S. N., Good-Jacobson, K. L., Curtis, J., Light, A., Tellier, J., Shi, W., Smyth, G. K., Tarlinton, D. M., Belz, G. T., Corcoran,
L. M., *et al.* 2014. Transcription factor IRF4 regulates germinal center cell formation through a B cell-intrinsic mechanism.
J Immunol, 192, 3200-6.
Wolf, F. A., Angerer, P. & Theis, F. J. 2018. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol,*
19, 15.
Wolock, S. L., Lopez, R. & Klein, A. M. 2019. Scrublet: Computational Identification of Cell Doublets in Single-Cell
Transcriptomic Data. *Cell Systems,* 8, 281-291.e9.
Wu, H. J. & Bondada, S. 2009. CD72, a coreceptor with both positive and negative effects on B lymphocyte development and
function. *J Clin Immunol,* 29, 12-21.
Xu, J., Husain, A., Hu, W., Honjo, T. & Kobayashi, M. 2014a. APE1 is dispensable for S-region cleavage but required for its
repair in class switch recombination. *PNAS,* 111, 17242-17247.
Xu, Y., Xu, L., Zhao, M., Xu, C., Fan, Y., Pierce, S. K. & Liu, W. 2014b. No receptor stands alone: IgG B-cell receptor intrinsic
and extrinsic mechanisms contribute to antibody memory. *Cell Research,* 24, 651-664.
Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. 2013. IgBLAST: an immunoglobulin variable domain sequence analysis tool.
Nucleic Acids Res, 41, W34-W40.
Zaretsky, I., Atrakchi, O., Mazor, R. D., Stoler-Barak, L., Biram, A., Feigelson, S. W., Gitlin, A. D., Engelhardt, B. & Shulman,
Z. 2017. ICAMs support B cell interactions with T follicular helper cells and promote clonal selection. *J Exp Med,* 214,
3435-3448.
Zheng, S., Kusnadi, A., Choi, J. E., Vuong, B. Q., Rhodes, D. & Chaudhuri, J. 2019. NME proteins regulate class switch
recombination. *FEBS Lett,* 593, 80-87.
Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., Benner, C. & Chanda, S. K. 2019. Metascape
provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun,* 10, 1523.

1018

Supplementary Figures

Figure S1

Figure S1. Annotation of non-B cell populations in the human tonsils.

A) Mean expression of key marker genes used to define T cell scRNA-seq clusters, including CD4+ naïve or central memory (CD4+ NCM), CD4+, T follicular helper (TfH), T follicular regulatory (TfH), T regulatory (Treg), CD8+ naïve or central memory (CD8+ NCM), CD8+ cytotoxic (CD8+ CTL), TIM3+ CD4/CD8 double-negative (TIM3+ DN) and cycling T cells, in addition to innate lymphoid cells (ILC) and natural killer (NK) cells. Frequency of cells for which each gene is detected is denoted by size of the dots.

B) Relative frequencies of different T cell subsets separated by clinical indication for tonsillectomy. OSA; obstructive sleep apnoea (*n* = 2), RT; recurrent tonsillitis (*n* = 3), RT+OSA (*n* = 2).

C) Mean expression of key marker genes used to define non-lymphoid cell scRNA-seq clusters, including monocyte/macrophages precursor (Precursors), macrophage (MAC1, MAC2, MAC3), conventional dendritic cell 1 (cDC1), plasmacytoid-derived dendritic cell (pDC) and follicular dendritic cell (FDC) subsets. Frequency of cells for which each gene is detected is denoted by size of the dots.

D) Relative frequencies of different non-lymphoid cell subsets separated by clinical indication for tonsillectomy.

Figure S2. Dynamic gene expression during class switch recombination and transcription factor binding at the immunoglobulin locus.

A) Smoothed gene expression for example cell surface receptor or cytokine (top row) and transcription factor (bottom row) genes that are differentially regulated through velocity-based pseudotime of B cell activation and GC entry.

B) Immunohistochemistry of CD20 (B cell marker) and APE1 (*APEX1*) in two paediatric human tonsils reveals depleted expression of APE1 in germinal centres (GCs) compared to the follicular zone.

C) Schematic of the human immunoglobulin heavy chain (IgH) locus, with intergenic (I) promoters, switch regions, germline transcripts and regulatory regions (Eµ, α₁ RR, α₂ RR).

D) Open chromatin (DNase-seq) and ChIP-seq from ENCODE consortium at Eµ, α₁ RR, α₂ RR, and a control neighbouring locus (*CRIP1 / C14orf80*) for EBV-transformed B lymphocyte cell line GM12878 and control non-B lymphocyte cell lines (K562, A549, HEPG2).

1046
Figure S3. Class- and subclass-specific gene expression analyses.
A) Pseudobulk heatmaps of average expression per donor of differentially expressed genes between class-specific GC B
      cells with similar affinity (based on SHM frequency).
B) Same as in A), but for subclass-specific gene expression analyses.

1051
Figure S4. Characterisation of memory B cell states identified by scRNA-seq.
A) Single-cell expression of memory B cell (*TNFRSF1B*), naïve/undifferentiated (*TCL1A*) and germinal centre (*CD38*)
markers across all B cell subsets (left) and sorted memory B cell subsets (right).
B) Single-cell expression of key marker genes differentially expressed by CD21^{low} MBC populations.
C) Top gene ontologies for significantly enriched genes in the IFN-response MBC cluster.
D) Single-cell AUCell-derived scores for top 50 marker genes of the naïve preGC B cells and FCRL2/3^{high} GC B cells in MBC
subsets. * denotes *p* value < 0.001.
E) Single-cell AUCell-derived scores for class switch recombination gene set in MBC subsets. * denotes *p* value < 0.001.
F) Relative frequencies of scVDJ-derived antibody subclass expression within different MBC scRNA-seq populations.
G) Single-cell expression of key marker genes of the FCRL2/3^{high} B cell states between switched and unswitched MBCs.
H) Somatic hypermutation frequencies of scVDJ-derived antibody genes between switched and unswitched B cells within
different MBC populations.

33