A transient role of primary cilia in controlling direct versus indirect neurogenesis in the developing cerebral cortex

Kerstin Hasenpusch-Theil[1], Christine Laclef[2,3], Matt Colligan[1], Eamon Fitzgerald[1], Katherine Howe[1], Emily Carroll[1], Shaun R. Abrams[4], Jeremy F. Reiter[4,5], Sylvie Schneider-Maunoury[2], and Thomas Theil[1,6]

[1]Centre for Discovery Brain Sciences, Hugh Robson Building, University of Edinburgh, Edinburgh, EH8 9XD, United Kingdom

[2] Sorbonne Université, CNRS UMR7622, INSERM U1156, Institut de Biologie Paris Seine (IBPS) - Developmental Biology Unit, 75005, Paris, France

[3]Present address: Sorbonne Université, Institut du Fer à Moulin INSERM U1270, Paris, France

[4]Department of Biochemistry and Biophysics, University of California, San Francisco, CA 94143

[5]Chan Zuckerberg Biohub, San Francisco, CA94158, USA Chan Zuckerberg Biohub, San Francisco, CA 94158, USA

[6]Author for correspondence (email: thomas.theil@ed.ac.uk)

Running title: Inpp5e and direct/indirect neurogenesis

Key words: *Inpp5e*; *Tctn2*; primary cilia; neurogenesis; cortex

ABSTRACT

During the development of the cerebral cortex, neurons are generated directly from radial glial cells or indirectly via basal progenitors. The balance between these division modes determines the number and types of neurons formed in the cortex thereby affecting cortical functioning. Here, we investigate the role of primary cilia in this process. We show that a mutation in the ciliary gene *Inpp5e* leads to a transient increase in direct neurogenesis and subsequently to an overproduction of layer V neurons in newborn mice. Loss of *Inpp5e* also affects ciliary structure coinciding with increased Akt and mTOR signalling and reduced Gli3 repressor levels. Genetically re-storing Gli3 repressor rescues the decreased indirect neurogenesis in *Inpp5e* mutants. Overall, our analyses reveal how primary cilia determine neuronal subtype composition of the cortex by controlling direct vs indirect neurogenesis. These findings have implications for understanding cortical malformations in ciliopathies with *INPP5E* mutations.

INTRODUCTION

Building a functional cerebral cortex which confers humans with their unique cognitive capabilities requires controlling the proliferation of neural progenitor cells and the timing and modes of neurogenic cell divisions. Varying the timing and modes of neurogenesis affects neuronal numbers and subtype composition of the cortex (Florio & Huttner, 2014). In the developing murine cortex, radial glial cells (RGCs) represent the major neural stem cell type. Residing in the ventricular zone, they express Pax6 and undergo interkinetic nuclear migration dividing at the ventricular surface (Götz, Stoykova, & Gruss, 1998; Warren et al., 1999). Initially, RGCs go through rounds of symmetric proliferative divisions to produce two RGCs increasing the progenitor pool but switch to asymmetric divisions at the beginning of cortical neurogenesis. RGCs generate neurons in two ways, either directly or indirectly via the production of basal progenitors (BPs) that settle in the subventricular zone (SVZ) and express the Tbr2 transcription factor (Englund et al., 2005). In the mouse, the majority of BPs divide once to produce two neurons whereas the remainders undergo one additional round of symmetric proliferative division before differentiating into two neurons (Haubensak, Attardo, Denk, & Huttner, 2004; Miyata et al., 2004; Noctor, Martinez-Cerdeno, Ivic, & Kriegstein, 2004). In this way, BPs increase neuron output per radial glial cell and have therefore been implicated in the evolutionary expansion of the mammalian cerebral cortex (Martinez-Cerdeno, Noctor, & Kriegstein, 2006). Thus, the balance between direct and indirect neurogenesis is an important factor in generating appropriate neuron numbers and types.

The mechanisms that fine tune this balance and thereby adjust the numbers and types of neurons produced in the cortex have only recently been investigated. Mitotic spindle orientation (Postiglione et al., 2011) and endoplasmic reticulum (ER) stress (Gladwyn-Ng et al., 2018; Laguesse et al., 2015) are contributing factors to control the generation of basal progenitors. In addition, levels of Slit/Robo and Notch/Delta signaling were shown to be evolutionarily conserved factors that determine the predominant mode of neurogenesis (Cardenas et al., 2018). Moreover, feedback signals from postmitotic neurons control the fate of radial glial daughter cells involving the release of Neurotrophin-3 and Fgf9 (Parthasarathy, Srivatsa, Nityanandam, & Tarabykin, 2014; Seuntjens et al., 2009) as well as the activation of a Notch-dependent signaling pathway (W. Wang et al., 2016). These studies highlight the importance of cell-cell signaling in controlling the cell lineage of cortical progenitors (Silva, Peyre, & Nguyen, 2019) and emphasize the necessity of studying the cellular mechanisms by which these signals control the decision by RGCs to undergo direct or indirect neurogenesis.

Given the importance of cell-cell signaling, it is likely that the primary cilium, a signaling hub in embryogenesis in general and in neural development in particular (Valente, Rosti, Gibbs, & Gleeson, 2014), plays key roles in determining the balance between direct versus indirect neurogenesis. The cilium is a subcellular protrusion that predominately emanates from the apical surface of radial glial cells projecting into the ventricular lumen. The phenotypes of several mouse

lines mutant for ciliary genes underline the importance of the primary cilium in forebrain development
but these mutants often suffer from severe patterning defects (Ashique et al., 2009; Besse et al.,
2011; Willaredt et al., 2008) which make studying ciliary roles in determining the lineage of cortical
progenitors difficult. To address such functions, we investigated corticogenesis in a mouse mutant
for the ciliary gene *Inpp5e*.

INPP5E is mutated in Joubert syndrome (JS) (Bielas et al., 2009; Jacoby et al., 2009), a
ciliopathy characterized by cerebellar defects in which a subset of patients also shows malformations
of the cerebral cortex including heterotopias, polymicrogyria and agenesis of the corpus callosum
(Valente et al., 2014). *Inpp5e* encodes Inositol polyphosphate 5 phosphatase E, an enzyme that is
localized in the ciliary membrane and that hydrolyses the phosphatidylinositol polyphosphates
$PI(4,5)P_2$ and $PI(3,4,5)P_3$ (Bielas et al., 2009; Jacoby et al., 2009). In this way, it controls the inositol
phosphate composition of the ciliary membrane and thereby regulates the activity of several
signaling pathways and cilia stability (Bielas et al., 2009; Chavez et al., 2015; Garcia-Gonzalo et al.,
2015; Jacoby et al., 2009; Plotnikova et al., 2015). In contrast to *Inpp5e*'s extensively studied
biochemical and cellular roles, little is known how these diverse functions are employed at the tissue
level to control RGC lineage.

Here, we show that loss of *Inpp5e* function results in an increase in neuron formation at the
expense of basal progenitor production in the E12.5 cortex and in an overproduction of Ctip2+ layer
V neurons in newborn mutants. Moreover, RGC cilia show unusual membranous structures and/or
abnormal numbers of microtubule doublets affecting the signaling capabilities of the cilium. The
levels of Gli3 repressor (Gli3R), a critical regulator of cortical stem cell development (Hasenpusch-
Theil et al., 2018; H. Wang, Ge, Uchida, Luu, & Ahn, 2011), is reduced and re-introducing Gli3R
rescues the decreased formation of basal progenitors. Taken together, these findings implicate the
primary cilium in controlling the decision of RGCs to either undergo direct neurogenesis or to form
basal progenitors, thereby governing the neuronal subtype composition of the cerebral cortex.
105

106   **RESULTS**

108   ***Inpp5e*^{Δ/Δ} embryos show mild telencephalic patterning defects**

109   Controlling the balance between direct and indirect neurogenesis in the developing cerebral

cortex is mediated by cell-cell signaling (Cardenas et al., 2018) and is hence likely to involve the

primary cilium. To investigate potential ciliary roles, we started characterizing cortical stem cell

development in embryos mutant for the *Inpp5e* gene which has a prominent role in ciliary signaling

and stability. Mutations in ciliary genes have previously been found to result in telencephalon

patterning defects, most notably in a ventralisation of the dorsal telencephalon and/or in defects at

the corticoseptal (CSB) and pallial/subpallial boundaries (PSPB) (Ashique et al., 2009; Besse et al.,

2011; Willaredt et al., 2008). Therefore, we first considered the possibility that such early patterning

defects may be present in *Inpp5e* mutant embryos and could affect cortical stem cell development.

In situ hybridization and immunofluorescence analyses of E12.5 control and *Inpp5e*^{Δ/Δ} embryos

revealed no obvious effect on the expression of dorsal and ventral telencephalic markers at the

corticoseptal boundary (SupFig. 1). In contrast, the pallial/subpallial boundary was not well defined

with a few scattered Pax6+ and *Dlx2* expressing cells on the wrong side of the boundary, i.e. in the

subpallium and pallium, respectively (SupFig. 1). Moreover, the hippocampal anlage appeared

smaller and disorganized with low level and diffuse expression of cortical hem markers (SupFig. 2),

consistent with known roles of Wnt/β-catenin and Bmp signaling in hippocampal development

(Galceran, Miyashita-Lin, Devaney, Rubenstein, & Grosschedl, 2000; Lee, Tole, Grove, & McMahon,

2000). In contrast, progenitors in the neocortical ventricular zone of *Inpp5e* mutant mice expressed

the progenitor markers *Emx1*, *Lhx2*, *Pax6* and *Ngn2*, though the levels of Pax6 protein expression

appeared reduced in the medial neocortex suggestive of a steeper lateral to medial Pax6 expression

gradient in mutant embryos. These expression patterns were maintained in E14.5 *Inpp5e*^{Δ/Δ} embryos

but revealed an area in the very caudal/dorsal telencephalon where the neocortex was folded

(SupFig. 3). At this level, folds were also present in the hippocampal anlage. Taken together, these

findings indicate that *Inpp5e* mutants have mild patterning defects affecting the integrity of the PSPB,

hippocampal development and the caudal-most neocortex while the rostral neocortex shows no

obvious malformation and can therefore be analysed for effects of the *Inpp5e* mutation on direct and

indirect neurogenesis.

***Inpp5e* controls direct vs indirect neurogenesis in the lateral neocortex**

138   Based upon these findings, we started analyzing the proliferation and differentiation of radial

glial cells in *Inpp5e*^{Δ/Δ} embryos in the rostrolateral and rostromedial neocortex to avoid the

regionalization defects described above. As a first step, we determined the proportion of radial glial

cells, basal progenitors and neurons in these regions in E12.5 embryos. Double

immunofluorescence for PCNA which labels all progenitor cells (Hall et al., 1990) and the radial glial

marker Pax6 did not reveal differences in the proportions of radial glial cells at both medial and lateral

levels (Fig. 1A-D). In contrast, the proportion of Tbr2+ basal progenitors was reduced laterally but

not medially (Fig. 1E-H). This decrease coincided with an increase in Tbr1+ neurons specifically in

the lateral neocortex (Fig. 1I-L). To determine whether these alterations are maintained at a later

developmental stage, we repeated this investigation in E14.5 embryos. This analysis revealed no

significant differences in the proportion of Pax6+ RGCs (Fig. 2A-D). Similarly, there was no alteration

in the proportion of Tbr2+ basal progenitors in lateral neocortex, however, their proportion was

reduced medially (Fig. 2E-H).

To label cortical projection neurons, we used double immunofluorescence for Tbr1 and Ctip2

which allowed us to distinguish between Ctip2+Tbr1+ and Ctip2+Tbr1- neurons. Quantifying these

subpopulations showed no effect on the formation of Ctip2+Tbr1+ neurons in $Inpp5e^{\Delta/\Delta}$ embryos. In

contrast, the proportion of Ctip2+Tbr1- neurons was reduced medially but increased in the lateral

neocortex (Fig. 2I-R). Taken together, these findings show that initially Tbr1+ and later Ctip2+Tbr1-

neurons were increasingly formed in the lateral neocortex of $Inpp5e^{\Delta/\Delta}$ embryos and that the

proportion of basal progenitors recovered after an initial down-regulation.

To address the defective cellular processes underlying these neurogenesis defects in $Inpp5e$

mutants, we first measured proliferation rates of cortical progenitors and performed double

immunofluorescence for PCNA and pHH3 which labels mitotic radial glial cells located at the

ventricular surface and dividing basal progenitors in abventricular positions (SupFig. 4). This analysis

revealed no statistically significant differences in the E12.5 and E14.5 lateral neocortex of control

and $Inpp5e^{\Delta/\Delta}$ embryos. The proportion of mitotic basal progenitors, however, was reduced in the

E12.5 medial neocortex (SupFig. 4).

The cell cycle represents another key regulator of neuronal differentiation and a mutation in

$Kif3a$ affects ciliogenesis and the cell cycle in the developing neocortex (Wilson, Wilson, Wang,

Wang, & McConnell, 2012). To investigate the possibility of altered cell cycle kinetics, we used a

BrdU/IdU double labelling strategy (Martynoga, Morrison, Price, & Mason, 2005; Nowakowski,

Lewin, & Miller, 1989) to determine S phase length and total cell cycle length in radial glial cells and

found no statistically significant changes in these parameters (SupFig. 5).

Finally, the increased neuron production could also be explained by an increase in direct

neurogenesis at the expense of basal progenitor cell formation. To test this possibility, we gave BrdU

to E11.5 pregnant mice 24h before dissecting the embryos. We then used BrdU immunostaining in

conjunction with Tbr1 and Tbr2 to identify the neurons and basal progenitors formed in the lateral

neocortex within the 24h time period. This analysis showed that the proportion of Tbr1+ neurons

compared to the total number of BrdU+ cells increased while the Tbr2+ proportion decreased in

$Inpp5e$ mutants (Fig. 3). Since the cell cycle of basal progenitors is longer than 24h (Arai et al.,

2011), the 24h interval used in our cell cycle exit experiment was too short for newly formed basal

progenitors to undergo one additional round of the cell cycle and as the BrdU label would have been

diluted with a further round of division, this analysis supports our hypothesis that direct neurogenesis

became more prevalent in *Inpp5e*$^{\Delta/\Delta}$ radial glial cells.

Cortical malformations in *Inpp5e*$^{\Delta/\Delta}$ embryos

Next, we investigated the consequences of this increase in direct neurogenesis on cortical

size and layer formation. Since *Inpp5e*$^{\Delta/\Delta}$ newborn pups die perinatally (Bielas et al., 2009), we

focused our analysis on E18.5 embryos. The mutant lacked obvious olfactory bulbs, as revealed by

whole mounts of control and mutant brains (SupFig. 6). To gain insights into the overall histology of

the mutant forebrain, we stained coronal sections with DAPI. This analysis showed that the mutant

cortex was thinner laterally but not medially with a more pronounced reduction of the thickness at

caudal levels (SupFig. 7). In addition, the hippocampus was malformed with a smaller dentate gyrus.

Investigating the expression of markers characteristic of the entire hippocampus (*Nrp2*; (Galceran

et al., 2000)), the CA1 field (Scip1; (Frantz, Bohner, Akers, & McConnell, 1994)) and the dentate

gyrus (Prox1; (Oliver et al., 1993)) showed that these hippocampal structures were present but were

severely reduced in size and disorganized in *Inpp5e*$^{\Delta/\Delta}$ embryos (SupFig. 8). In addition, the corpus

callosum, the major axon tract connecting the two cerebral hemispheres, was correctly formed but

was smaller. We confirmed this effect by staining callosal axons and surrounding glial cells that guide

these axons to the contralateral hemisphere with L1 and GFAP, respectively (SupFig. 9).

After characterizing the gross morphology of the *Inpp5e*$^{\Delta/\Delta}$ cortex, we next investigated

whether the increased neuron formation in E12.5 mutant embryos led to changes in the neuronal

subtype composition of the E18.5 cortex. To this end, we used immunofluorescence labelling for

Tbr1 and Ctip2 to analyse the formation of layer VI and V neurons, respectively, whereas Satb2

served as a layer II-IV marker (Fig. 4). Inspecting these immunostainings at low magnification

showed that Tbr1+, Ctip2+ and Satb2+ neurons occupied their correct relative laminar positions in

Inpp5e mutants (Fig. 4A-F) except for neuronal heterotopias which were present in all mutant brains,

though their number and position varied (Fig. 4D). These immunostainings also revealed a medial

shift in the position of the rhinal fissure, a sulcus that is conserved across mammalian species and

separates neocortex from the paleocortical piriform cortex (Ariens-Kapers, Huber, & Crosby, 1936).

This shift was more marked caudally and suggests a dramatic expansion of the *Inpp5e* mutant

piriform cortex at the expense of neocortex at caudal most levels (Fig. 4D-F). Using the Tbr1/Ctip2

and Satb2 stainings, we determined the proportions of deep and superficial layer neurons,

respectively. Because of the expanded piriform cortex in *Inpp5e* mutants, we limited this

investigation to the unaffected rostral neocortex. In the rostrolateral neocortex, we found the

proportion of Tbr1+ neurons to be reduced (Fig. 4G, H, M). This reduction coincided with an

increased proportion of Ctip2+ layer V neurons (Fig. 4I, J, N) while the Satb2 population was

unchanged (Fig. 4K, L, O). In contrast, the rostromedial neocortex did not show any differences (Fig.

4P-X). Thus, the increase in direct neurogenesis in the lateral neocortex during earlier development

concurs with a change in the proportions of E18.5 Tbr1+ and Ctip2+ deep layer neurons.

A mutation in the ciliary gene *Tctn2* leads to increased telencephalic neurogenesis

To start to unravel the mechanisms by which *Inpp5e* controls cortical stem cell development,

we first analysed whether the increased early neurogenesis is restricted to *Inpp5e*$^{\Delta/\Delta}$ mutants or is

observed in another mutant affecting cilia. To this end, we focused on the *TECTONIC 2* (*TCTN2*)

gene which is crucial for ciliary transition zone architecture (Shi et al., 2017) and which, like *INPP5E*,

is mutated in Joubert Syndrome (Garcia-Gonzalo et al., 2011). Interestingly, E12.5 *Tctn2*$^{-/-}$ mutant

embryos (Reiter & Skarnes, 2006) also showed an increased proportion of Tbr1+ projection neurons

and a concomitant decrease in Tbr2+ basal progenitors in the dorsolateral telencephalon (Fig. 5).

Due to embryonic lethality, however, we were not able to investigate the formation of cortical neurons

at later stages.

Ciliary defects in the forebrain of E12.5 *Inpp5e*$^{\Delta/\Delta}$ embryos

Our findings in the *Inpp5e* and *Tctn2* mutants suggested a role for cilia in cortical progenitor

cells to control early neurogenesis. Therefore, we examined the presence and the structure of

primary cilia in the developing forebrain of *Inpp5e*$^{\Delta/\Delta}$ embryos by immunofluorescence and electron

microscopy. We first analyzed the presence of the small GTPase Arl13b, enriched in ciliary

membranes, and of γ-Tubulin (γTub), a component of basal bodies (Caspary, Larkins, & Anderson,

2007). We found no major difference in the number, the apical localization or the size of cilia in

control and *Inpp5e*$^{\Delta/\Delta}$ neuroepithelial cells in the E12.5 telencephalon (Fig. 6A,B) or diencephalon

(data not shown).

To gain insights into the fine structure of these primary cilia we performed electron

microscopy analyses. Scanning electron microscopy (SEM) provided an observation of the cilia

protruding into the telencephalic ventricles. In control embryos, almost all radial glial cells had a

single, ~1 μm long primary cilium (Fig. 6C), as previously described (Besse et al., 2011). Some

Inpp5e$^{\Delta/\Delta}$ mutant cells also displayed an apparently normal cilium (Fig. 6D, E), whereas other cells

harbored abnormal cilia, either with a lateral blob (arrowhead in Fig. 6D) or as a short and bloated

cilium-like protrusion (arrows in Fig. 6D,E).

Transmission electron microscopy (TEM) confirmed the presence of abnormal cilia in

Inpp5e$^{\Delta/\Delta}$ embryos. Cilia were recognized by basal bodies anchored to the apical membrane in both

control and *Inpp5e*$^{\Delta/\Delta}$ radial glial cells (Fig. 6F-L, N, Q). However, in *Inpp5e*$^{\Delta/\Delta}$ cells, some cilia lacked

the axoneme and showed unusual membranous structures that resemble budding vesicles emerging

from the lateral surface of the cilium (Fig. 6G, K), internal vesicles (arrows in Fig. 6I, K, L), or

undulating peripheral membranes (Fig. 6I), indicating an *Inpp5e*-dependent defect in ciliary

membrane morphology. Transverse sections revealed the presence of cilia with apparently normal

9+0 axonemes, as well as cilia containing abnormal numbers of microtubule doublets in *Inpp5e*$^{\Delta/\Delta}$

embryos (Fig. 6O, P). To quantify these ciliary defects, we counted the number of normal versus

abnormal cilia on TEM images obtained from control and *Inpp5e*$^{\Delta/\Delta}$ embryos, and found an increase

in abnormal cilia in *Inpp5e*$^{\Delta/\Delta}$ compared to control embryos (Fig. 6R). Taken together, a significant

number of abnormal primary cilia were found at the apical end of E12.5 radial glial cells in the

forebrain of *Inpp5e*$^{\Delta/\Delta}$ embryos. These abnormalities are consistent with a role of *Inpp5e* in

maintaining cilia stability (Jacoby et al., 2009).

Receptor tyrosine kinase and mTOR signaling in *Inpp5e*$^{\Delta/\Delta}$ embryos

Given these defects in ciliary structure, we started to investigate cilia-controlled signaling

pathways, defects in which might underlie the increased direct neurogenesis in the *Inpp5e*$^{\Delta/\Delta}$ mutant

radial glial cells. Inpp5e hydrolyzes the 5-phosphate of PI(3,4,5)P$_3$ which serves as a second

messenger in receptor tyrosine kinase signaling. We therefore investigated the amount and

distribution of pErk1/2 as a read-out of receptor tyrosine kinase signaling. Western blots using

dissected dorsal telencephalon from E12.5 and E14.5 embryos revealed no difference in the ratio

between pErk1/2 and Erk1/2 (SupFig. 10). Moreover, pErk1/2 expression in the E12.5 forebrain

remains restricted to the LGE, the diencephalon and to meninges covering the telencephalon but

was not identified in cortical progenitors (SupFig. 10). In E14.5 embryos, we detected pErk1/2

immunoreactivity in cortical progenitors in a lateral to medial gradient covering the whole extent of

the neocortex. In contrast, high level pErk1/2 expression remained confined to the lateral neocortex

with little or no expression medially in *Inpp5e*$^{\Delta/\Delta}$ embryos (SupFig. 10). These findings suggest that

pErk signaling does not play a role in the early neurogenesis defect in *Inpp5e* mutants but may be

involved in the recovery of basal progenitor formation.

PI(3,4,5)P$_3$, a substrate of the Inpp5e phosphatase, is essential for the effective activation of

the serine threonine kinase Akt (Kisseleva, Cao, & Majerus, 2002; Plotnikova et al., 2015) raising

the possibility that loss of *Inpp5e* results in activated Akt signaling in the neocortex. Moreover,

phosphorylation of Akt at Serine[473] has been implicated in inhibiting cilia assembly and promoting

cilia disassembly (Mao et al., 2019). We therefore investigated the amount of phospho-Akt[S473] in

western blots on protein extracts from the E12.5 and E14.5 dorsal telencephalon. This analysis

revealed increased phosphorylation of Akt at Serine[473] at E12.5 but no significant change at E14.5

(SupFig. 11). Since Akt acts upstream of mTOR (Yu & Cui, 2016), we also analysed the activity of

the mTOR pathway and investigated the phosphorylation of the S6 ribosomal protein, at Serine[235/236]

and Serine[S240/244] (Ferrari, Bandi, Hofsteenge, Bussian, & Thomas, 1991). Interestingly, *Inpp5e*

mutants showed increased S6 phosphorylation at Serine[235/236] in E12.5 and E14.5 embryos whereas

an increase in phospho-S6[S240/244] remained restricted to E12.5 (SupFig. 11). Taken together, these

findings indicate that *Inpp5e* inactivation leads to increased activation of Akt and mTOR signaling in

the E12.5 dorsal telencephalon.

Next, we investigated a potential role of elevated mTOR signaling in controlling the balance

between direct and indirect neurogenesis and treated E11.5 pregnant mice with the mTORC1

inhibitor rapamycin (1mg/kg bodyweight). We harvested embryos 24h later and determined the

proportions of basal progenitors and neurons. This analysis revealed that administering rapamycin

did not rescue basal progenitor and neuron formation although the cortex of rapamycin treated

Inpp5e$^{\Delta/\Delta}$ embryos appeared elongated (SupFig. 11). Based on this experiment, augmented mTOR

signaling does not underlie the altered ratio of direct vs indirect neurogenesis in *Inpp5e*$^{\Delta/\Delta}$ embryos.

Restoring Gli3 repressor ratio rescues cortical malformations in *Inpp5e*$^{\Delta/\Delta}$ embryos

Primary cilia also play a crucial role in Shh signaling by controlling the proteolytic cleavage

of full length Gli3 (Gli3FL) into the Gli3 repressor form (Gli3R) in the absence of Shh and by

converting Gli3FL into the transcriptional activator Gli3A in the presence of Shh. Moreover, the dorsal

telencephalon predominately forms Gli3R (Fotaki, Yu, Zaki, Mason, & Price, 2006) and mice that

can only produce Gli3R have no obvious defect in cortical development (Besse et al., 2011; Böse,

Grotewold, & Rüther, 2002). In addition, we recently showed that Gli3 has a prominent role in radial

glial cells controlling the switch from symmetric proliferative to asymmetric neurogenic cell division

(Hasenpusch-Theil et al., 2018). Therefore, we hypothesized that alterations in Gli3 processing

caused by abnormal cilia function underlies the increased direct neurogenesis and the cortical

malformations in *Inpp5e*$^{\Delta/\Delta}$ embryos. In situ hybridization showed that *Gli3* mRNA expression might

be slightly reduced but the overall expression pattern in the telencephalon remains unaffected

(SupFig. 12). We next investigated the formation of Gli3FL and Gli3R in the E12.5 dorsal

telencephalon of *Inpp5e*$^{\Delta/\Delta}$ embryos using Western blots. This analysis revealed no change in the

levels of Gli3FL but a significant decrease inGli3R which resulted in a reduced Gli3R to Gli3FL ratio

in the mutant (Fig. 7A-D) suggesting that the *Inpp5e* mutation affects Gli3 processing.

The next set of experiments aimed to clarify a role for the reduced Gli3 processing. To this

end, we restored Gli3R levels by crossing *Inpp5e* mutants with *Gli3*$^{\Delta699/+}$ mice that can only produce

Gli3R in a cilia-independent manner (Besse et al., 2011; Böse et al., 2002). Overall inspection of

Inpp5e$^{\Delta/\Delta}$;*Gli3*$^{\Delta699/+}$ embryos revealed normal eye formation whereas *Inpp5e*$^{\Delta/\Delta}$ embryos either

completely lacked eyes or showed microphthalmia (Jacoby et al., 2009) (SupFig. 13). Moreover, the

overall morphology of the telencephalon is much improved in *Inpp5e*$^{\Delta/\Delta}$;*Gli3*$^{\Delta699/+}$ embryos as

compared to *Inpp5e*$^{\Delta/\Delta}$ embryos. In E18.5 *Inpp5e*$^{\Delta/\Delta}$ *Gli3*$^{\Delta699/+}$ mutants, the corpus callosum has a

thickness indistinguishable from that of control embryos (SupFig. 14). In E12.5 and E14.5

Inpp5e$^{\Delta/\Delta}$;*Gli3*$^{\Delta699/+}$ embryos, the neocortex lacks the undulations of the VZ present in *Inpp5e*$^{\Delta/\Delta}$

embryos (data not shown) and the morphology of the hippocampal anlage is more akin to that in

wild-type embryos but it is still smaller and less bulged (Fig. 7E, H, K, N).

We also determined the proportions of basal progenitors and Tbr1+ neurons at E12.5 which

were decreased and increased, respectively, in the lateral neocortex of *Inpp5e*$^{\Delta/\Delta}$ embryos.

Remarkably, there was no statistically significant difference between control and $Inpp5e^{\Delta/\Delta}$;$Gli3^{\Delta699/+}$

embryos (Fig. 7E-J) indicating that the neurogenesis phenotype of E12.5 $Inpp5e^{\Delta/\Delta}$ mutants is

rescued by a single copy of the $Gli3^{\Delta699}$ allele. The proportion of Tbr1+Ctip2+ neurons was not

affected in the medial neocortex of E14.5 $Inpp5e^{\Delta/\Delta}$;$Gli3^{\Delta699/+}$ embryos (Fig. 7K, M, N) either, however,

the proportion of Tbr1-Ctip2+ neurons was reduced (Fig. 7K, N, P). Similarly, the proportions of basal

progenitors in the E14.5 medial neocortex was reduced in $Inpp5e^{\Delta/\Delta}$ $Gli3^{\Delta699/+}$ embryos as in $Inpp5e^{\Delta/\Delta}$

embryos (Fig. 7L, O, S). As re-introducing a single $Gli3^{\Delta699}$ allele does not completely rescue the

$Inpp5e^{\Delta/\Delta}$ neurogenesis phenotype, we generated $Inpp5e^{\Delta/\Delta}$ embryos homozygous for $Gli3^{\Delta699}$.

Interestingly, the morphology of the telencephalon including the hippocampal formation was

indistinguishable between control and $Inpp5e^{\Delta/\Delta}$;$Gli3^{\Delta699/\Delta699}$ embryos (Fig. 7K, Q) and the proportions

of Tbr1-Ctip2+ and basal progenitors were not affected any longer (Fig. 7K, L. P-S). Taken together,

these findings indicate that re-introducing a single copy of the *Gli3R* allele into the *Inpp5e* mutant

background leads to a partial rescue of cortical neurogenesis in $Inpp5e^{\Delta/\Delta}$ embryos whereas two

copies are required for a full rescue.

DISCUSSION

Generating a functional cerebral cortex requires a finely tuned balance between direct and indirect neurogenesis to form subtypes of cortical projection neurons in appropriate numbers. Here, we show that the ciliary mouse mutants *Inpp5e* and *Tctn2* present with a transient increase in neurons forming directly from radial glia progenitors in the lateral neocortex at the expense of basal progenitor formation. This increase in neurogenesis results in augmented formation of Ctip2+ layer V neurons in the *Inpp5e* mutant cortex. Our studies also revealed that the *Inpp5e* mutation interfered with the stability of the RGC primary cilium and its signaling functions, leading to a reduction in the Gli3R levels. Since re-introducing Gli3R in an *Inpp5e* mutant background restored the decreased formation of normal proportions of basal progenitors and neurons, our findings implicate a novel role for primary cilia in controlling the signaling events that direct the decision of RGCs to undergo either direct or indirect neurogenesis.

Primary cilia affect the decision between direct and indirect neurogenesis

Radial glial cells in the developing mouse neocortex have the potential to undergo symmetric proliferative or asymmetric cell divisions with the latter division mode producing neurons in a direct manner or indirectly via basal progenitors. Balancing out these division modes is important not only to determine final neuronal output and cortical size but also the types of cortical projection neurons and, hence, subtype composition of the adult neocortex. In the E12.5 *Inpp5e* and *Tctn2* mouse mutants, we identified an increased formation of neurons in the lateral neocortex. Based on our cell cycle exit experiment additional neurons are formed from RGCs at the expense of basal progenitors. Given the cell cycle length of basal progenitors of >24 hours (Arai et al., 2011), it is unlikely that new born basal progenitors would have undergone an additional round of cell division to produce two neurons within the time frame of this experiment. Such an extra division would also have diluted the BrdU label. We therefore conclude that the *Inpp5e* mutation caused RGCs to preferentially produce neurons directly.

Interestingly, this increase in direct neurogenesis led to an increased proportion of Ctip2+ deep layer V neurons in the E18.5 neocortex but did not coincide with a reduced proportion of upper layer neurons. This effect could be explained in several mutually non-exclusive ways. First, neurons born at E12.5 initially express both Ctip2 and Tbr1 (Fig. 7) and later down-regulate Ctip2. *Inpp5e* could therefore affect the signaling that controls this downregulation. Secondly, the proportions of basal progenitors and neurons was normalized in E14.5 mutants. Since basal progenitors are a main source of upper layer neurons (Arnold et al., 2008; Vasistha et al., 2015), this normalization would account for the sufficient numbers of Satb2+ upper layer neurons. Newly formed projection neurons signal back to RGCS via Jag1, Fgf9 and Neurotrophin 3 (Parthasarathy et al., 2014; Seuntjens et al., 2009; W. Wang et al., 2016) to control the sequential production of deep and upper layer neurons and of glia (Silva et al., 2019). *Inpp5e* might affect these signals by controlling cilia stability and/or levels of PI(3,4,5)P_3 (Bielas et al., 2009; Jacoby et al., 2009) that acts as a second messenger in

receptor tyrosine kinase signaling. This possibility is supported by our observation that the extent of

the pErk expression domain in the neocortex is diminished in E14.5 $Inpp5e^{\Delta/\Delta}$ embryos. Regardless

of the exact mechanism, our findings suggest a novel, spatially and temporally restricted role for

Inpp5e in controlling the decision between direct and indirect neurogenesis. This function differs from

those described for other cilia mutants. Conditional inactivation of *Ift88* and *Kif3a* leads to a larger

cortex (Foerster et al., 2017; Wilson et al., 2012) with a modest increase in BP production in the

absence of a delay in neurogenesis (Foerster et al., 2017) while *Rpgrip1l* mutants have reduced

numbers of both basal progenitors and neurons (Postel, Karam, Pezeron, Schneider-Maunoury, &

Clement, 2019). These findings highlight the multiple and varied roles cilia play in cortical

development.

391 *Inpp5e* controls direct/indirect neurogenesis through Gli3 processing

Our study also shed lights into the mechanisms by which *Inpp5e* controls the decision

between direct/indirect neurogenesis. Most notably, the Gli3R level and Gli3R/Gli3FL ratio are

decreased in $Inpp5e^{\Delta/\Delta}$ embryos. While the *Inpp5e* mutation does not lead to an up-regulation of Shh

signaling in the dorsal telencephalon (Magnani et al., 2015), re-introducing a single or two copies of

Gli3R in an *Inpp5e* mutant background partially and fully restores the neurogenesis defects,

respectively. This rescue indicates that reduced levels of Gli3R rather than the reduction in the

Gli3R/Gli3FL ratio are responsible for the prevalence of direct neurogenesis in $Inpp5e^{\Delta/\Delta}$ embryos.

This idea is consistent with the findings that (i) $Gli3^{\Delta699/\Delta699}$ embryos that cannot produce Gli3FL and

Gli3A show no obvious phenotype in cortical development (Besse et al., 2011; Böse et al., 2002),

(ii) dorsal telencephalic patterning defects in $Gli3^{Xt/Xt}$ mutants are not rescued in $Shh^{-/-}/Gli3^{XtXt}$ double

mutants (Rallu et al., 2002; Rash & Grove, 2007), (iii) Shh promotes the generation of olfactory bulb

interneurons and cortical oligodendrocytes and neurogenesis in the subventricular zone by reducing

Gli3R rather than by promoting Gli activator function (Petrova, Garcia, & Joyner, 2013; H. Wang,

Kane, Lee, & Ahn, 2014; Zhang et al., 2020). In addition, there is also a dramatic rescue of eye

development and the rescue also extends to other malformations of the $Inpp5e^{\Delta/\Delta}$ forebrain, including

the corpus callosum, the hippocampus and the expansion of the piriform cortex, structures that are

also affected in *Gli3* null and hypomorphic mutants (Amaniti et al., 2015; Johnson, 1967; Magnani

et al., 2014; Theil, Alvarez-Bolado, Walter, & Rüther, 1999; Wiegering, Petzsch, Kohrer, Ruther, &

Gerhardt, 2019). Taken together, these findings support the idea that *Inpp5e* and the primary cilium

control key processes in cortical development by regulating the formation of Gli3R.

Our analyses support several mutually non-exclusive mechanisms how the *Inpp5e* mutation

impacts on Gli3 processing. First, our electron microscopy study revealed severe structural

abnormalities in large proportions of cilia. The Inpp5e phosphatase hydrolyses $PI(3,4,5)P_3$, which is

essential for the effective activation of the serine threonine kinase Akt (Kisseleva et al., 2002;

Plotnikova et al., 2015). Following PIP3 binding, Akt translocates to the membrane and becomes

phosphorylated at T308 by phosphoinositide dependent kinase-1 (Pdk1) and at S473 by mammalian

target of rapamycin complex (mTORC2) (Yu & Cui, 2016). Consistent with the loss of *Inpp5e* function

and a resulting increase in $PI(3,4,5)P_3$, our western blot analysis revealed elevated $pAkt^{S473}$ levels.

Increased phosphorylation at this site has been implicated in inhibiting cilia assembly and promoting

cilia disassembly (Mao et al., 2019) and could hence explain the structural defects of RGC *Inpp5e*$^{\Delta/\Delta}$

cilia. Secondly, *Inpp5e* could control Gli3 processing through its effect on the transition zone (TZ). It

is required for TZ molecular organisation (Dyson et al., 2017) and its substrate PI(4,5)P2 plays a

role in TZ maturation in Drosophila (Gupta, Fabian, & Brill, 2018). This model is further supported by

our finding that a mouse mutant for the TZ protein Tctn2 phenocopies the *Inpp5e*$^{\Delta/\Delta}$ neurogenesis

defect. In turn, several mouse mutants defective for TZ proteins show microphthalmia (Garcia-

Gonzalo et al., 2011; Sang et al., 2011; Yee et al., 2015). Tctn proteins are also required for Gli3

processing (Sang et al., 2011; Thomas et al., 2012; C. Wang, Li, Meng, & Wang, 2017) and the TZ

protein Rpgrip1l controls the activity of the proteasome at the basal body responsible for proteolytic

cleavage of Gli3 (Gerhardt et al., 2015). Taken together, these findings indicate that *Inpp5e* mutation

might affect the ability of RGCs to switch to indirect neurogenesis through defects in cilia stability

and/or the integrity of the ciliary transition zone.

Implications for Joubert Syndrome

In humans, hypomorphic *INPP5E* mutations contribute to Joubert Syndrome (JS), a

ciliopathy characterized by cerebellar malformations and concomitant ataxia and breathing

abnormalities. In addition, a subset of JS patients exhibit cortical abnormalities including

polymicrogyria, neuronal heterotopias and agenesis of the corpus callosum (Poretti, Huisman,

Scheer, & Boltshauser, 2011). Strikingly, the *Inpp5e* mouse mutant also shows several of these

abnormalities. In the caudal telencephalon, the otherwise lissencephalic cortex formed folds

reminiscent of the polymicrogyria in JS patients. In addition, the mutant formed leptomeningeal

heterotopias with 100% penetrance, but their number and location varied. Mutations in ciliary genes

were previously associated with heterotopia formation in humans and mice (Magnani et al., 2015;

Uzquiano et al., 2019). Mice carrying mutations in the *Eml1* gene encoding a microtubule-associated

protein show subcortical heterotopias due to a mispositioning of radial glial cells and impaired

primary cilia formation (Uzquiano et al., 2019). Finally, the corpus callosum is thinner but callosal

axons normally project to the contralateral cerebral hemisphere in *Inpp5e* mutants. This phenotype

is milder compared to that of other mouse mutants with altered cilia that show complete agenesis of

the corpus callosum with callosal axons forming Probst bundles (Benadiba et al., 2012; Laclef et al.,

2015; Putoux et al., 2019). Unlike these other ciliary mutants, the corticoseptal boundary which plays

a crucial role in positioning guidepost cells that control midline crossing of callosal axons (Magnani

et al., 2014) is not obviously affected in *Inpp5e*$^{\Delta/\Delta}$ embryos. Instead, the thinner corpus callosum is

likely to be the result of an expanded piriform cortex. Despite these mechanistic differences,

however, re-introducing Gli3R into the cilia mutant background restores callosal development in both

groups of mutants suggesting that cilia control two independent steps in corpus callosum formation

by regulating Gli3 processing. Thus, the *Inpp5e*$^{\Delta/\Delta}$ mutant recapitulates cortical abnormalities in JS

patients and starts to help unravelling the pathomechanisms underlying these defects.

MATERIAL & METHODS

Mice

All experimental work was carried out in accordance with the UK Animals (Scientific Procedures) Act
1986 and UK Home Office guidelines. All protocols were reviewed and approved by the named
veterinary surgeons of the College of Medicine and Veterinary Medicine, the University of Edinburgh,
prior to the commencement of experimental work. $Inpp5e^{\Delta/+}$ and $Gli3^{\Delta699/+}$ mouse lines have been
described previously (Böse et al., 2002; Jacoby et al., 2009). $Inpp5e^{\Delta/+}$ mice were interbred to
generate $Inpp5e^{\Delta/\Delta}$ embryos; exencephalic $Inpp5e^{\Delta/\Delta}$ embryos were excluded from the analyses.
Wild-type and $Inpp5e^{\Delta/+}$ litter mate embryos served as controls. $Inpp5e^{\Delta/\Delta};Gli3^{\Delta699/+}$ and
$Inpp5e^{\Delta/\Delta};Gli3^{\Delta699/\Delta699}$ embryos were obtained from inter-crosses of $Inpp5e^{\Delta/+};Gli3^{\Delta699/+}$ mice using
wild-type, $Inpp5e^{\Delta/+}$ and $Gli3^{\Delta699/+}$ embryos as controls. Embryonic (E) day 0.5 was assumed to start
at midday of the day of vaginal plug discovery. Transgenic animals and embryos were genotyped as
described (Böse et al., 2002; Jacoby et al., 2009). For each marker and each stage, 3-8 embryos
were analysed.

For measuring cell cycle lengths, pregnant females were intraperitoneally injected with a
single dose of IdU (10mg/ml) at E12.5, followed by an injection of BrdU (10mg/ml) 90 min later.
Embryos were harvested 30 min after the second injection. For cell cycle exit analyses, BrdU was
injected peritoneally into E11.5 pregnant females and embryos were harvested 24 hrs later.
Rapamycin was dissolved in 100% EtOH and diluted in vehicle containing 5% PEG400, 5% Tween80
immediately before use. E11.5 pregnant females received a single intraperitoneal injection of
Rapamycin at 1 mg/kg body weight or of vehicle and were killed 24 hrs later.

Immunohistochemistry and in situ hybridisation

For immunohistochemistry, embryos were fixed overnight in 4% paraformaldehyde, incubated in
30% sucrose at +4°C for 24h, embedded in 30% sucrose/OCT mixture (1:1) and frozen on dry ice.
Immunofluorescence staining was performed on 12 to 14 μm cryostat sections as described
previously (Theil, 2005) with antibodies against Arl13b (Neuromab 75-287; 1:1500), rabbit anti-BrdU
(1:50, Abcam #ab6326), mouse anti-BrdU/IdU (B44) (1:50, BD Biosciences #347580), rat anti-Ctip2
(1:1000, Abcam #18465), rabbit anti-pErk$^{T202/Y204}$ (1:1000, Cell Signaling Technology #9101), rabbit
anti-GFAP (1:1000, Dako #Z 0334), rat anti-L1, clone 324 (1:1000, Millipore #MAB5272), rabbit anti-
Pax6 (1:400, Biolegend #901301), mouse anti-PCNA (1:500, Abcam #29), rabbit anti-Prox1 (1:1000,
RELIA*Tech* #102-PA32). rabbit anti-pHH3 (1:100, Millipore #06-570), mouse anti-Satb2 (1:200,
Abcam #51502), rabbit anti-Tbr1 (1:400, Abcam #31940), rabbit anti-Tbr2 (1:1000, Abcam #23345)
and γTUB (Sigma T6557; 1:2000). Primary antibodies for immunohistochemistry were detected with
Alexa- or Cy2/3-conjugated fluorescent secondary antibodies. The Tbr1 signals were amplified using
biotinylated secondary IgG antibody (swine anti-rabbit IgG) (1:400, BD Biosciences) followed by

Alexa Fluor 488 or 568 Streptavidin (1:100, Invitrogen). For counter staining DAPI (1:2000, Life

Technologies) was used. Prox1 and pErk$^{T202/Y204}$ proteins were detected non-fluorescently using

biotinylated goat anti-rabbit IgG (1: 400, BD Biosciences) followed by avidin-HRP and DAB detection

(Vector labs) as described previously (Magnani et al., 2010).

In situ hybridisation on 12μm serial paraffin sections were performed as described previously (Theil,

2005) using antisense RNA probes for *Axin2* (Lustig et al., 2002), *Bmp4* (Jones, Lyons, & Hogan,

1991), *Dbx1* (Yun, Potter, & Rubenstein, 2001), *Dlx2* (Bulfone et al., 1993), *Emx1* (Simeone et al.,

1992), *Gli3* (Hui, Slusarski, Platt, Holmgren, & Joyner, 1994), *Lhx2* (Liem, Tremml, & Jessell, 1997),

Msx1 (Hill et al., 1989), *Ngn2* (Gradwohl, Fode, & Guillemot, 1996), *Nrp2* (Galceran et al., 2000),

Pax6 (Walther & Gruss, 1991), *Scip1* (Frantz et al., 1994), *Wnt2b* (Grove, Tole, Limon, Yip, &

Ragsdale, 1998).

Western blot

Protein was extracted from the dorsal telencephalon of E12.5 (3 tissues pooled per sample) and

E14.5 (single tissue per sample) wild-type and *Inpp5e*$^{\Delta/\Delta}$ embryos (n=4 samples per genotype) as

described previously (Magnani et al., 2010). For the detection of Gli3 10μg protein lysates were

subjected to gel electrophoresis on a 3-8% NuPAGE® Tris-Acetate gel (Life Technologies), and

protein was transferred to a Immobilon-FL membrane (Millipore), which was incubated with goat

anti-h/m Gli3 (1:500, R&D Systems #AF3690) and mouse anti-β-Actin antibody (1:15000, Abcam

#ab6276). After incubating with donkey anti-goat IgG IRDye680RD (1:15000, LI-COR Biosciences)

and donkey anti-mouse IgG IRDye800CW secondary antibodies (1:15000, Life Technologies),

signal was detected using LI-COR's Odyssey Infrared Imaging System with Odyssey Software.

Values for protein signal intensity were obtained using Image Studio Lite Version3.1. Gli3 repressor

and activator protein levels were compared between wild-type and mutant tissue using a paired t-

test.

For all other Western blot analyses 20μg protein lysates were loaded on 4–12% NuPAGE® Bis-Tris

gels (Life Technologies) and later transferred to an Immobilon–FL membrane. Membranes were

incubated with the following primary antibodies: rabbit anti-phospho-Akt^{S473} (1:1000, Cell Signaling

Technologies CST #9271), rabbit anti-Akt (1:1000, CST #9272), rabbit anti-phospho-Erk$^{T202/Y204}$

(1:1000, CST #9101), rabbit anti-Erk (L34F12) (CST # 4696), rabbit anti-ribosomal protein phospho-

S6$^{S235/S236}$ (1:1000, CST #2211), rabbit anti-ribosomal protein phospho-S6$^{S240/S244}$ (1:1000, CST

#2215) and mouse anti-ribosomal protein S6 (C-8) (1:1000, Santa Cruz Technologies #sc-74459).

For the detection of phosphorylated proteins goat anti-rabbit IRDye 680RD (1:15,000, LI-COR

Biosciences) and for total proteins goat anti-mouse or goat anti-rabbit IRDye 800CW (1:15,000, LI-

COR Biosciences) were used as secondary antibodies. The signals were detected via the Odyssey

Imaging System and further analysed using Image Studio Lite Version4.0. The ratios between

phosphorylated and total protein were compated between wild-type and mutant tissue using a paired

t-test.

Scanning and transmission electron microscopy

TEM and SEM image acquisition were performed in the Cochin Imaging Facility and on the IBPS EM Facility, respectively. For scanning electron microscopy, embryos were dissected in 1.22x PBS (pH 7.4) and fixed overnight with 2% glutaraldehyde in 0.61x PBS (pH 7.4) at 4°C. Heads were then sectioned to separate the dorsal and ventral parts of the telencephalon, exposing their ventricular surfaces. Head samples were washed several times in 1.22x PBS and postfixed for 15 minutes in 1.22x PBS containing 1% OsO4. Fixed samples were washed several times in ultrapure water, dehydrated with a graded series of ethanol and prepared for scanning electron microscopy using the critical point procedure (CPD7501, Polaron). Their surfaces were coated with a 20 nm gold layer using a gold spattering device (Scancoat Six, Edwards). Samples were observed under a Cambridge S260 scanning electron microscope at 10 keV.

For transmission electron microscopy tissues were fixed for 1 hour with 3% glutaraldehyde, post-fixed in 1.22x PBS containing 1% OsO4, then dehydrated with a graded ethanol series. After 10 minutes in a 1:2 mixture of propane:epoxy resin, tissues were embedded in gelatin capsules with freshly prepared epoxy resin and polymerized at 60°C for 24 hours. Sections (80 nm) obtained using an ultramicrotome (Reichert Ultracut S) were stained with uranyl acetate and Reynold's lead citrate and observed with a Philips CM10 transmission electron microscope.

Statistical Analyses

Data were analysed using GraphPadPrism 6 software with n=3-8 embryos for all analyses. Power analysis of pilot experiments informed minimum samples size. Mann Whitney tests were performed for immunohistochemical analyses in general. Cortical thickness was analysed using a two way ANOVA followed by Sidak's multiple comparisons test. Paired t-tests were performed for Western blots and a fisher's exact test was used to analyse the quantification of normal and abnormal cilia. The mTORC1 and Gli3 rescue experiments were evaluated with one way ANOVAS followed by Tukey's multiple comparisons test. A single asterisk indicates significance of $p<0.05$, two asterisks indicate significance of $p<0.01$ and three asterisks of $p<0.005$. Due to morphological changes blinding was not possible and scores were validated by a second independent observer.

ACKNOWLEDGEMENTS

We are grateful to Drs Thomas Becker, John Mason, Pleasantine Mill, and David Price for critical comments on the manuscript, Dr Christos Gkogkas for invaluable advice on mTOR signaling, and Stéphane Schurmans for the *Inpp5e*$^{\Delta/+}$ mouse line. We also thank Dr Michaël Trichet (electron microscopy platform of the IBPS-Sorbonne Universités Paris 6) and Dr Alain Schmitt (electron

microscopy platform of the Institut Cochin CNRS-UMR 8104) for their help with scanning and
transmission electron microscopy analyses, respectively. This work was supported by a grant from
the Biotechnology and Biological Sciences Research Council to TT (BB/P00122X/1) and NIH
R01GM095941 to JFR.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

Amaniti, E. M., Fu, C., Lewis, S., Saisana, M., Magnani, D., Mason, J. O., & Theil, T. (2015).
Expansion of the piriform cortex contributes to corticothalamic pathfinding defects in Gli3
conditional mutants. *Cereb Cortex, 25*(2), 460-471. doi:10.1093/cercor/bht244
Arai, Y., Pulvers, J. N., Haffner, C., Schilling, B., Nusslein, I., Calegari, F., & Huttner, W. B. (2011).
Neural stem and progenitor cells shorten S-phase on commitment to neuron production.
Nature communications, 2, 154. doi:10.1038/ncomms1155
Ariens-Kapers, C. U., Huber, G. C., & Crosby, E. C. (1936). The comparative anatomy of the
nervous system of the vertebrates including man. *New York Macmillan*.
Arnold, S. J., Huang, G. J., Cheung, A. F., Era, T., Nishikawa, S., Bikoff, E. K., . . . Groszer, M.
(2008). The T-box transcription factor Eomes/Tbr2 regulates neurogenesis in the cortical
subventricular zone. *Genes Dev, 22*(18), 2479-2484. doi:10.1101/gad.475408
Ashique, A. M., Choe, Y., Karlen, M., May, S. R., Phamluong, K., Solloway, M. J., . . . Peterson, A.
S. (2009). The Rfx4 transcription factor modulates Shh signaling by regional control of
ciliogenesis. *Sci Signal, 2*(95), ra70. doi:10.1126/scisignal.2000602
Benadiba, C., Magnani, D., Niquille, M., Morle, L., Valloton, D., Nawabi, H., . . . Durand, B. (2012).
The ciliogenic transcription factor RFX3 regulates early midline distribution of guidepost
neurons required for corpus callosum development. *PLoS genetics, 8*(3), e1002606.
doi:10.1371/journal.pgen.1002606
Besse, L., Neti, M., Anselme, I., Gerhardt, C., Ruther, U., Laclef, C., & Schneider-Maunoury, S.
(2011). Primary cilia control telencephalic patterning and morphogenesis via Gli3
proteolytic processing. *Development, 138*(10), 2079-2088. doi:10.1242/dev.059808
Bielas, S. L., Silhavy, J. L., Brancati, F., Kisseleva, M. V., Al-Gazali, L., Sztriha, L., . . . Gleeson, J.
G. (2009). Mutations in INPP5E, encoding inositol polyphosphate-5-phosphatase E, link
phosphatidyl inositol signaling to the ciliopathies. *Nat Genet, 41*(9), 1032-1036.
doi:10.1038/ng.423
Böse, J., Grotewold, L., & Rüther, U. (2002). Pallister-Hall syndrome phenotype in mice mutant for
Gli3. *Hum Mol Genet, 11*(9), 1129-1135.
Bulfone, A., Puelles, L., Porteus, M. H., Frohman, M. A., Martin, G. R., & Rubenstein, J. L. (1993).
Spatially restricted expression of Dlx-1, Dlx-2 (Tes-1), Gbx-2, and Wnt-3 in the embryonic
day 12.5 mouse forebrain defines potential transverse and longitudinal segmental
boundaries. *J Neurosci, 13*(7), 3155-3172.
Cardenas, A., Villalba, A., de Juan Romero, C., Pico, E., Kyrousi, C., Tzika, A. C., . . . Borrell, V.
(2018). Evolution of Cortical Neurogenesis in Amniotes Controlled by Robo Signaling
Levels. *Cell, 174*(3), 590-606 e521. doi:10.1016/j.cell.2018.06.007
Caspary, T., Larkins, C. E., & Anderson, K. V. (2007). The graded response to Sonic Hedgehog
depends on cilia architecture. *Developmental cell, 12*(5), 767-778.
doi:10.1016/j.devcel.2007.03.004
Chavez, M., Ena, S., Van Sande, J., de Kerchove d'Exaerde, A., Schurmans, S., & Schiffmann, S.
N. (2015). Modulation of Ciliary Phosphoinositide Content Regulates Trafficking and Sonic
Hedgehog Signaling Output. *Developmental cell, 34*(3), 338-350.
doi:10.1016/j.devcel.2015.06.016
Dyson, J. M., Conduit, S. E., Feeney, S. J., Hakim, S., DiTommaso, T., Fulcher, A. J., . . . Mitchell,
C. A. (2017). INPP5E regulates phosphoinositide-dependent cilia transition zone function. *J
Cell Biol, 216*(1), 247-263. doi:10.1083/jcb.201511055
Englund, C., Fink, A., Lau, C., Pham, D., Daza, R. A., Bulfone, A., . . . Hevner, R. F. (2005). Pax6,
Tbr2, and Tbr1 are expressed sequentially by radial glia, intermediate progenitor cells, and
postmitotic neurons in developing neocortex. *J Neurosci, 25*(1), 247-251.
Ferrari, S., Bandi, H. R., Hofsteenge, J., Bussian, B. M., & Thomas, G. (1991). Mitogen-activated
70K S6 kinase. Identification of in vitro 40 S ribosomal S6 phosphorylation sites. *J Biol
Chem, 266*(33), 22770-22775.
Florio, M., & Huttner, W. B. (2014). Neural progenitors, neurogenesis and the evolution of the
neocortex. *Development, 141*(11), 2182-2194. doi:10.1242/dev.090571

Foerster, P., Daclin, M., Asm, S., Faucourt, M., Boletta, A., Genovesio, A., & Spassky, N. (2017).
mTORC1 signaling and primary cilia are required for brain ventricle morphogenesis.
Development, 144(2), 201-210. doi:10.1242/dev.138271
Fotaki, V., Yu, T., Zaki, P. A., Mason, J. O., & Price, D. J. (2006). Abnormal positioning of
diencephalic cell types in neocortical tissue in the dorsal telencephalon of mice lacking
functional Gli3. *J Neurosci, 26*(36), 9282-9292.
Frantz, G. D., Bohner, A. P., Akers, R. M., & McConnell, S. K. (1994). Regulation of the POU
domain gene SCIP during cerebral cortical development. *J Neurosci, 14*(2), 472-485.
Galceran, J., Miyashita-Lin, E. M., Devaney, E., Rubenstein, J. L., & Grosschedl, R. (2000).
Hippocampus development and generation of dentate gyrus granule cells is regulated by
LEF1. *Development, 127*(3), 469-482.
Garcia-Gonzalo, F. R., Corbit, K. C., Sirerol-Piquer, M. S., Ramaswami, G., Otto, E. A., Noriega, T.
R., . . . Reiter, J. F. (2011). A transition zone complex regulates mammalian ciliogenesis
and ciliary membrane composition. *Nat Genet, 43*(8), 776-784. doi:10.1038/ng.891
Garcia-Gonzalo, F. R., Phua, S. C., Roberson, E. C., Garcia, G., 3rd, Abedin, M., Schurmans, S., .
. . Reiter, J. F. (2015). Phosphoinositides Regulate Ciliary Protein Trafficking to Modulate
Hedgehog Signaling. *Developmental cell, 34*(4), 400-409. doi:10.1016/j.devcel.2015.08.001
Gerhardt, C., Lier, J. M., Burmuhl, S., Struchtrup, A., Deutschmann, K., Vetter, M., . . . Ruther, U.
(2015). The transition zone protein Rpgrip1l regulates proteasomal activity at the primary
cilium. *J Cell Biol, 210*(1), 115-133. doi:10.1083/jcb.201408060
Gladwyn-Ng, I., Cordon-Barris, L., Alfano, C., Creppe, C., Couderc, T., Morelli, G., . . . Nguyen, L.
(2018). Stress-induced unfolded protein response contributes to Zika virus-associated
microcephaly. *Nature neuroscience, 21*(1), 63-71. doi:10.1038/s41593-017-0038-4
Götz, M., Stoykova, A., & Gruss, P. (1998). Pax6 controls radial glia differentiation in the cerebral
cortex. *Neuron, 21*(5), 1031-1044.
Gradwohl, G., Fode, C., & Guillemot, F. (1996). Restricted expression of a novel murine atonal-
related bHLH protein in undifferentiated neural precursors. *Dev Biol, 180*(1), 227-241.
Grove, E. A., Tole, S., Limon, J., Yip, L., & Ragsdale, C. W. (1998). The hem of the embryonic
cerebral cortex is defined by the expression of multiple Wnt genes and is compromised in
Gli3-deficient mice. *Development, 125*(12), 2315-2325.
Gupta, A., Fabian, L., & Brill, J. A. (2018). Phosphatidylinositol 4,5-bisphosphate regulates cilium
transition zone maturation in Drosophila melanogaster. *J Cell Sci, 131*(16).
doi:10.1242/jcs.218297
Hall, P. A., Levison, D. A., Woods, A. L., Yu, C. C., Kellock, D. B., Watkins, J. A., . . . et al. (1990).
Proliferating cell nuclear antigen (PCNA) immunolocalization in paraffin sections: an index
of cell proliferation with evidence of deregulated expression in some neoplasms. *J Pathol,*
162(4), 285-294. doi:10.1002/path.1711620403
Hasenpusch-Theil, K., West, S., Kelman, A., Kozic, Z., Horrocks, S., McMahon, A. P., . . . Theil, T.
(2018). Gli3 controls the onset of cortical neurogenesis by regulating the radial glial cell
cycle through Cdk6 expression. *Development, 145*(17). doi:10.1242/dev.163147
Haubensak, W., Attardo, A., Denk, W., & Huttner, W. B. (2004). Neurons arise in the basal
neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc*
Natl Acad Sci U S A, 101(9), 3196-3201.
Hill, R. E., Jones, P. F., Rees, A. R., Sime, C. M., Justice, M. J., Copeland, N. G., . . . Davidson, D.
R. (1989). A new family of mouse homeo box-containing genes: molecular structure,
chromosomal location, and developmental expression of Hox-7.1. *Genes Dev, 3*(1), 26-37.
Hui, C. C., Slusarski, D., Platt, K. A., Holmgren, R., & Joyner, A. L. (1994). Expression of three
mouse homologs of the Drosophila segment polarity gene cubitus interruptus, Gli, Gli-2,
and Gli-3, in ectoderm- and mesoderm-derived tissues suggests multiple roles during
postimplantation development. *Dev Biol, 162*(2), 402-413.
Jacoby, M., Cox, J. J., Gayral, S., Hampshire, D. J., Ayub, M., Blockmans, M., . . . Schurmans, S.
(2009). INPP5E mutations cause primary cilium signaling defects, ciliary instability and
ciliopathies in human and mouse. *Nat Genet, 41*(9), 1027-1031.
Johnson, D. R. (1967). Extra-toes: a new mutant gene causing multiple abnormalities in the
mouse. *J Embryol Exp Morphol, 17*(3), 543-581.

Jones, C. M., Lyons, K. M., & Hogan, B. L. (1991). Involvement of Bone Morphogenetic Protein-4 (BMP-4) and Vgr-1 in morphogenesis and neurogenesis in the mouse. *Development, 111*(2), 531-542.

Kisseleva, M. V., Cao, L., & Majerus, P. W. (2002). Phosphoinositide-specific inositol polyphosphate 5-phosphatase IV inhibits Akt/protein kinase B phosphorylation and leads to apoptotic cell death. *J Biol Chem, 277*(8), 6266-6272. doi:10.1074/jbc.M105969200

Laclef, C., Anselme, I., Besse, L., Catala, M., Palmyre, A., Baas, D., . . . Schneider-Maunoury, S. (2015). The role of primary cilia in corpus callosum formation is mediated by production of the Gli3 repressor. *Hum Mol Genet, 24*(17), 4997-5014. doi:10.1093/hmg/ddv221

Laguesse, S., Creppe, C., Nedialkova, D. D., Prevot, P. P., Borgs, L., Huysseune, S., . . . Nguyen, L. (2015). A Dynamic Unfolded Protein Response Contributes to the Control of Cortical Neurogenesis. *Developmental cell, 35*(5), 553-567. doi:10.1016/j.devcel.2015.11.005

Lee, S. M., Tole, S., Grove, E., & McMahon, A. P. (2000). A local Wnt-3a signal is required for development of the mammalian hippocampus. *Development, 127*(3), 457-467.

Liem, K. F., Jr., Tremml, G., & Jessell, T. M. (1997). A role for the roof plate and its resident TGFbeta-related proteins in neuronal patterning in the dorsal spinal cord. *Cell, 91*(1), 127-138.

Lustig, B., Jerchow, B., Sachs, M., Weiler, S., Pietsch, T., Karsten, U., . . . Behrens, J. (2002). Negative Feedback Loop of Wnt Signaling through Upregulation of Conductin/Axin2 in Colorectal and Liver Tumors. *Mol. Cell. Biol., 22*(4), 1184-1193.

Magnani, D., Hasenpusch-Theil, K., Benadiba, C., Yu, T., Basson, M. A., Price, D. J., . . . Theil, T. (2014). Gli3 controls corpus callosum formation by positioning midline guideposts during telencephalic patterning. *Cerebral cortex, 24*(1), 186-198. doi:10.1093/cercor/bhs303

Magnani, D., Hasenpusch-Theil, K., Jacobs, E. C., Campagnoni, A. T., Price, D. J., & Theil, T. (2010). The Gli3 hypomorphic mutation Pdn causes selective impairment in the growth, patterning, and axon guidance capability of the lateral ganglionic eminence. *J Neurosci, 30*(41), 13883-13894. doi:30/41/13883 [pii]
10.1523/JNEUROSCI.3650-10.2010

Magnani, D., Morle, L., Hasenpusch-Theil, K., Paschaki, M., Jacoby, M., Schurmans, S., . . . Theil, T. (2015). The ciliogenic transcription factor Rfx3 is required for the formation of the thalamocortical tract by regulating the patterning of prethalamus and ventral telencephalon. *Hum Mol Genet, 24*(9), 2578-2593. doi:10.1093/hmg/ddv021

Mao, H., Tang, Z., Li, H., Sun, B., Tan, M., Fan, S., . . . Sun, Y. (2019). Neddylation inhibitor MLN4924 suppresses cilia formation by modulating AKT1. *Protein Cell*. doi:10.1007/s13238-019-0614-3

Martinez-Cerdeno, V., Noctor, S. C., & Kriegstein, A. R. (2006). The role of intermediate progenitor cells in the evolutionary expansion of the cerebral cortex. *Cereb Cortex, 16 Suppl 1*, i152-161. doi:10.1093/cercor/bhk017

Martynoga, B., Morrison, H., Price, D. J., & Mason, J. O. (2005). Foxg1 is required for specification of ventral telencephalon and region-specific regulation of dorsal telencephalic precursor proliferation and apoptosis. *Dev Biol, 283*(1), 113-127.

Miyata, T., Kawaguchi, A., Saito, K., Kawano, M., Muto, T., & Ogawa, M. (2004). Asymmetric production of surface-dividing and non-surface-dividing cortical progenitor cells. *Development, 131*(13), 3133-3145. doi:10.1242/dev.01173

Noctor, S. C., Martinez-Cerdeno, V., Ivic, L., & Kriegstein, A. R. (2004). Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nature neuroscience, 7*(2), 136-144. doi:10.1038/nn1172

Nowakowski, R. S., Lewin, S. B., & Miller, M. W. (1989). Bromodeoxyuridine immunohistochemical determination of the lengths of the cell cycle and the DNA-synthetic phase for an anatomically defined population. *J Neurocytol, 18*(3), 311-318.

Oliver, G., Sosa-Pineda, B., Geisendorf, S., Spana, E. P., Doe, C. Q., & Gruss, P. (1993). Prox 1, a prospero-related homeobox gene expressed during mouse development. *Mech Dev, 44*(1), 3-16. doi:10.1016/0925-4773(93)90012-m

Parthasarathy, S., Srivatsa, S., Nityanandam, A., & Tarabykin, V. (2014). Ntf3 acts downstream of Sip1 in cortical postmitotic neurons to control progenitor cell fate through feedback signaling. *Development, 141*(17), 3324-3330. doi:10.1242/dev.114173

Petrova, R., Garcia, A. D., & Joyner, A. L. (2013). Titration of GLI3 repressor activity by sonic hedgehog signaling is critical for maintaining multiple adult neural stem cell and astrocyte functions. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 33*(44), 17490-17505. doi:10.1523/JNEUROSCI.2042-13.2013

Plotnikova, O. V., Seo, S., Cottle, D. L., Conduit, S., Hakim, S., Dyson, J. M., . . . Smyth, I. M. (2015). INPP5E interacts with AURKA, linking phosphoinositide signaling to primary cilium stability. *J Cell Sci, 128*(2), 364-372. doi:10.1242/jcs.161323

Poretti, A., Huisman, T. A., Scheer, I., & Boltshauser, E. (2011). Joubert syndrome and related disorders: spectrum of neuroimaging findings in 75 patients. *AJNR Am J Neuroradiol, 32*(8), 1459-1463. doi:10.3174/ajnr.A2517

Postel, M., Karam, A., Pezeron, G., Schneider-Maunoury, S., & Clement, F. (2019). A multiscale mathematical model of cell dynamics during neurogenesis in the mouse cerebral cortex. *BMC Bioinformatics, 20*(1), 470. doi:10.1186/s12859-019-3018-8

Postiglione, M. P., Juschke, C., Xie, Y., Haas, G. A., Charalambous, C., & Knoblich, J. A. (2011). Mouse inscuteable induces apical-basal spindle orientation to facilitate intermediate progenitor generation in the developing neocortex. *Neuron, 72*(2), 269-284. doi:10.1016/j.neuron.2011.09.022

Putoux, A., Baas, D., Paschaki, M., Morle, L., Maire, C., Attie-Bitach, T., . . . Durand, B. (2019). Altered GLI3 and FGF8 signaling underlies acrocallosal syndrome phenotypes in Kif7 depleted mice. *Hum Mol Genet, 28*(6), 877-887. doi:10.1093/hmg/ddy392

Rallu, M., Machold, R., Gaiano, N., Corbin, J. G., McMahon, A. P., & Fishell, G. (2002). Dorsoventral patterning is established in the telencephalon of mutants lacking both Gli3 and Hedgehog signaling. *Development, 129*(21), 4963-4974.

Rash, B. G., & Grove, E. A. (2007). Patterning the dorsal telencephalon: a role for sonic hedgehog? *J Neurosci, 27*(43), 11595-11603.

Reiter, J. F., & Skarnes, W. C. (2006). Tectonic, a novel regulator of the Hedgehog pathway required for both activation and inhibition. *Genes Dev, 20*(1), 22-27. doi:10.1101/gad.1363606

Sang, L., Miller, J. J., Corbit, K. C., Giles, R. H., Brauer, M. J., Otto, E. A., . . . Jackson, P. K. (2011). Mapping the NPHP-JBTS-MKS protein network reveals ciliopathy disease genes and pathways. *Cell, 145*(4), 513-528. doi:10.1016/j.cell.2011.04.019

Seuntjens, E., Nityanandam, A., Miquelajauregui, A., Debruyn, J., Stryjewska, A., Goebbels, S., . . . Tarabykin, V. (2009). Sip1 regulates sequential fate decisions by feedback signaling from postmitotic neurons to progenitors. *Nature neuroscience, 12*(11), 1373-1380. doi:10.1038/nn.2409

Shi, X., Garcia, G., 3rd, Van De Weghe, J. C., McGorty, R., Pazour, G. J., Doherty, D., . . . Reiter, J. F. (2017). Super-resolution microscopy reveals that disruption of ciliary transition-zone architecture causes Joubert syndrome. *Nat Cell Biol, 19*(10), 1178-1188. doi:10.1038/ncb3599

Silva, C. G., Peyre, E., & Nguyen, L. (2019). Cell migration promotes dynamic cellular interactions to control cerebral cortex morphogenesis. *Nat Rev Neurosci, 20*(6), 318-329. doi:10.1038/s41583-019-0148-y

Simeone, A., Gulisano, M., Acampora, D., Stornaiuolo, A., Rambaldi, M., & Boncinelli, E. (1992). Two vertebrate homeobox genes related to the Drosophila empty spiracles gene are expressed in the embryonic cerebral cortex. *EMBO J, 11*(7), 2541-2550.

Theil, T. (2005). Gli3 is required for the specification and differentiation of preplate neurons. *Dev Biol, 286*(2), 559-571.

Theil, T., Alvarez-Bolado, G., Walter, A., & Rüther, U. (1999). Gli3 is required for Emx gene expression during dorsal telencephalon development. *Development, 126*(16), 3561-3571.

Thomas, S., Legendre, M., Saunier, S., Bessieres, B., Alby, C., Bonniere, M., . . . Attie-Bitach, T. (2012). TCTN3 mutations cause Mohr-Majewski syndrome. *Am J Hum Genet, 91*(2), 372-378. doi:10.1016/j.ajhg.2012.06.017

Uzquiano, A., Cifuentes-Diaz, C., Jabali, A., Romero, D. M., Houllier, A., Dingli, F., . . . Francis, F. (2019). Mutations in the Heterotopia Gene Eml1/EML1 Severely Disrupt the Formation of Primary Cilia. *Cell Rep, 28*(6), 1596-1611 e1510. doi:10.1016/j.celrep.2019.06.096

Valente, E. M., Rosti, R. O., Gibbs, E., & Gleeson, J. G. (2014). Primary cilia in
neurodevelopmental disorders. *Nat Rev Neurol, 10*(1), 27-36.
doi:10.1038/nrneurol.2013.247
Vasistha, N. A., Garcia-Moreno, F., Arora, S., Cheung, A. F., Arnold, S. J., Robertson, E. J., &
Molnar, Z. (2015). Cortical and Clonal Contribution of Tbr2 Expressing Progenitors in the
Developing Mouse Brain. *Cereb Cortex, 25*(10), 3290-3302. doi:10.1093/cercor/bhu125
Walther, C., & Gruss, P. (1991). Pax-6, a murine paired box gene, is expressed in the developing
CNS. *Development, 113*(4), 1435-1449.
Wang, C., Li, J., Meng, Q., & Wang, B. (2017). Three Tctn proteins are functionally conserved in
the regulation of neural tube patterning and Gli3 processing but not ciliogenesis and
Hedgehog signaling in the mouse. *Dev Biol, 430*(1), 156-165.
doi:10.1016/j.ydbio.2017.08.003
Wang, H., Ge, G., Uchida, Y., Luu, B., & Ahn, S. (2011). Gli3 is required for maintenance and fate
specification of cortical progenitors. *J Neurosci, 31*(17), 6440-6448. doi:31/17/6440 [pii]
10.1523/JNEUROSCI.4892-10.2011
Wang, H., Kane, A. W., Lee, C., & Ahn, S. (2014). Gli3 repressor controls cell fates and cell
adhesion for proper establishment of neurogenic niche. *Cell Rep, 8*(4), 1093-1104.
doi:10.1016/j.celrep.2014.07.006
Wang, W., Jossin, Y., Chai, G., Lien, W. H., Tissir, F., & Goffinet, A. M. (2016). Feedback
regulation of apical progenitor fate by immature neurons through Wnt7-Celsr3-Fzd3
signalling. *Nature communications, 7*, 10936. doi:10.1038/ncomms10936
Warren, N., Caric, D., Pratt, T., Clausen, J. A., Asavaritikrai, P., Mason, J. O., . . . Price, D. J.
(1999). The transcription factor, Pax6, is required for cell proliferation and differentiation in
the developing cerebral cortex. *Cereb Cortex, 9*(6), 627-635.
Wiegering, A., Petzsch, P., Kohrer, K., Ruther, U., & Gerhardt, C. (2019). GLI3 repressor but not
GLI3 activator is essential for mouse eye patterning and morphogenesis. *Dev Biol, 450*(2),
141-154. doi:10.1016/j.ydbio.2019.02.018
Willaredt, M. A., Hasenpusch-Theil, K., Gardner, H. A., Kitanovic, I., Hirschfeld-Warneken, V. C.,
Gojak, C. P., . . . Tucker, K. L. (2008). A crucial role for primary cilia in cortical
morphogenesis. *J Neurosci, 28*(48), 12887-12900.
Wilson, S. L., Wilson, J. P., Wang, C., Wang, B., & McConnell, S. K. (2012). Primary cilia and Gli3
activity regulate cerebral cortical size. *Dev Neurobiol, 72*(9), 1196-1212.
doi:10.1002/dneu.20985
Yee, L. E., Garcia-Gonzalo, F. R., Bowie, R. V., Li, C., Kennedy, J. K., Ashrafi, K., . . . Reiter, J. F.
(2015). Conserved Genetic Interactions between Ciliopathy Complexes Cooperatively
Support Ciliogenesis and Ciliary Signaling. *PLoS Genet, 11*(11), e1005627.
doi:10.1371/journal.pgen.1005627
Yu, J. S., & Cui, W. (2016). Proliferation, survival and metabolism: the role of PI3K/AKT/mTOR
signalling in pluripotency and cell fate determination. *Development, 143*(17), 3050-3060.
doi:10.1242/dev.137075
Yun, K., Potter, S., & Rubenstein, J. L. (2001). Gsh2 and Pax6 play complementary roles in
dorsoventral patterning of the mammalian telencephalon. *Development, 128*(2), 193-205.
Zhang, Y., Liu, G., Guo, T., Liang, X. G., Du, H., Yang, L., . . . Chen, B. (2020). Cortical Neural
Stem Cell Lineage Progression Is Regulated by Extrinsic Signaling Molecule Sonic
Hedgehog. *Cell Rep, 30*(13), 4490-4504 e4494. doi:10.1016/j.celrep.2020.03.027
847
848

Figure 1: Increased neuron formation in the dorsolateral telencephalon of E12.5 *Inpp5$^{\Delta/\Delta}$*
mutant embryos. (A-D) Pax6/PCNA double immunofluorescence staining revealed the proportion
of apical radial glial cells which remained unaltered in the mutant. The boxes in (A) indicate the
regions in the medial (m) and lateral (l) telencephalon at which cell counts were performed. (E-H)
Reduced proportions of basal progenitors in the lateral telencephalon as revealed by staining for
Tbr2 and PCNA. (I-L) Tbr1 immunostaining showed that the proportion of neurons is increased in
the lateral telencephalon. All statistical data are presented as means ± 95% confidence intervals
(CI); Mann Whitney tests; n = 4 except for (H) with n=5; * p < 0.05. Scale bar: 100μm. ctx: cortex;
LGE: lateral ganglionic eminence.

Figure 2: Proportions of radial glial cells, basal progenitors and neurons in the neocortex of E14.5 *Inpp5e*$^{\Delta/\Delta}$ embryos. (A-D) The proportion of radial glial cells remains unaffected by the *Inpp5e* mutation as revealed by Pax6/PCNA double immunofluorescence. The boxes in A indicate the regions in the medial (m) and lateral (l) telencephalon at which cell counts were performed. (E-H) Tbr2/PCNA double staining showed a reduced proportion of basal progenitors in the *Inpp5e*$^{\Delta/\Delta}$ medial but not lateral neocortex. (I-R) Neuron formation in the *Inpp5e*$^{\Delta/\Delta}$ mutant necocortex. The proportion of Tbr1$^+$Ctip2$^+$ neurons is not significantly altered (I-L) whereas the proportion of Tbr1$^-$Ctip2$^+$ neurons is decreased and increased in the medial and lateral neocortex, respectively (M-R). All statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; * p < 0.05. Scale bars: 100μm (A) and 50μm (M). ctx: cortex; LGE: lateral ganglionic eminence.

Figure 3: Increased neurogenesis at the expense of basal progenitor formation in E12.5
***Inpp5e*$^{\Delta/\Delta}$ mutants.** Immunohistochemistry on sections of E12.5 control (A, D) and *Inpp5e*$^{\Delta/\Delta}$
embryos (B, E) that were treated with BrdU 24 hours earlier. (A-C) Tbr2/BrdU double labelling
showed that less basal progenitors formed from the BrdU labelled progenitor cohort. (D-F) The
proportion of newly formed Tbr1^{+} neurons is increased. The arrows in D and E label Tbr1^{+}BrdU^{+}
cells. All statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney
tests; n = 4; * p < 0.05. Scale bar: 50μm.

Figure 4: Increased formation of layer V neurons in E18.5 *Inpp5e*^{Δ/Δ} mutants. (A-F) Coronal sections immunostained for the deep layer markers Tbr1 (layer VI) and Ctip2 (layer V) and for the upper layer marker Satb2 (layers II-IV); there is no obvious defect in layering in the *Inpp5e*$^{Δ/Δ}$ mutant except for the formation of a heterotopia (asterisk in D). At caudal levels, the cortex becomes thinner and the rhinal fissure is shifted medially as indicated by the arrows. (G-O) Formation of cortical neurons at rostrolateral levels. The proportion of Tbr1+layer VI neurons is decreased with a concomitant increase in Ctip2+layer V neurons. (P-X) Portion of cortical neurons at rostromedial levels. Immunolabeling with cortical layer markers revealed no significant difference. Note that due to the thinner cortex, the position of layer VI Tbr1+ (Q) and layer V Ctip2+ neurons (J, S) appears to be shifted to more superficial positions, however, the relative order of these layers remains unaffected. All statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; * $p < 0.05$. Scale bars: 500μm (A) and 100μm (G). CC: corpus callosum; ctx: cortex; sep: septum; str: striatum.

Figure 5: Increased generation of cortical neurons in E12.5 *Tctn2*[-/-] embryos. (A-C) Double

immunofluorescence for PCNA and Tbr2 revealed a significantly decreased proportion of basal

progenitors. (D-F) The portion of Tbr1[+] cortical neurons is increased. All statistical data are presented

as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; * $p < 0.05$. Scale bar: 50μm.

bv: blood vessel.

Figure 6: Ciliary defects in E12.5 *Inpp5e*^ΔΔ forebrain. (A-B) Immunohistochemistry for Arl13b and γ-Tubulin (γTUB) on E12.5 brain cryosections showed an accumulation of ciliary axonemes and basal bodies, respectively, at the apical border of radial glial cells facing the ventricules in both control (A) and *Inpp5e*^ΔΔ (B) embryos without any obvious difference. Scale bars: 10 μm.

(C-E) Scanning electron microscopy (SEM) on E12.5 control (C) and *Inpp5e*^ΔΔ (D, E) brains highlighted the presence of primary cilia projecting from the apical surface of radial glial cells in both control (A) and *Inpp5e*^ΔΔ (B) embryos. However, SEM also revealed the presence of abnormal cilia

in *Inpp5e*$^{\Delta/\Delta}$ embryos having a spherical shape (arrows in D and E) or aberrant lateral buddings

(arrowheads in D and E). Scale bars: 2 μm.

(F-L) Transmission electron microscopy (TEM) analysis on E12.5 brains showed longitudinal

sections of primary cilia in control (F) and *Inpp5e*$^{\Delta/\Delta}$ (G-L) embryos. In control primary cilia, the

axoneme appeared as an extension of the basal body (bb, black arrowheards) (F-H). In addition to

cilia with normal morphology, abnormal cilia were identified in *Inpp5e*$^{\Delta/\Delta}$ embryos thanks to the

presence of a basal body apparently correctly docked to the apical membrane. Abnormal cilia lacked

an axoneme (I, J, L) or showed unusual membranous structures, such as budding (G, K) or internal

(I, K, L) vesicles (arrows) or undulating peripheral membranes (I). Note that tight junctions (white

arrowheads in F and G) appeared normal in *Inpp5e*$^{\Delta/\Delta}$ (G) and control (F) embryos, suggesting that

apico-basal polarity of *Inpp5e*$^{\Delta/\Delta}$ radial glial cells was not compromised. Scale bars: 200 nm.

(M-Q) TEM images showing transverse sections of the axoneme (M, O, P) and the basal body (N,

Q) in control (M, N) and *Inpp5e*$^{\Delta/\Delta}$ (O-Q) embryos with no major difference in the basal bodies

between control (N) and *Inpp5e*$^{\Delta/\Delta}$ (Q) embryos. However, transverse section of primary cilia in

Inpp5e$^{\Delta/\Delta}$ brains revealed the presence of normal axonemes composed of 9 correctly organized

doublets of microtubules on some radial glial cells (O), while others harboured an abnormal axoneme

containing a lower number of microtubule doublets (P). Scale bars: 50 nm. (R) Graph showing the

number of normal versus abnormal cilia (cil.) found on TEM images from control (n=3) or *Inpp5e*$^{\Delta/\Delta}$

(n=3) embryos. cc: counted cilia.

Figure 7: Re-introducing the Gli3 repressor rescues the neurogenesis defect in *Inpp5e*
mutants. (A-D). Gli3 Western blot on E12.5 dorsal telencephalic tissue revealed the Gli3 full length
(FL) and repressor (R) forms (A). While Gli3FL levels are not affected (B), levels of Gli3R (C) and
the Gli3R/Gli3FL ratio (D) are decreased in *Inpp5e*$^{\Delta/\Delta}$ embryos. A paired t-test was used to evaluate
levels of Gli3FL and Gli3R and the Gli3R/Gli3FL ratio in four control/*Inpp5e*$^{\Delta/\Delta}$ embryo pairs derived
from four different litters. (E-S) Formation of basal progenitors and neurons in the neocortex of
Inpp5e$^{\Delta/\Delta}$;Gli3$^{\Delta699/+}$ and *Inpp5e*$^{\Delta/\Delta}$;Gli3$^{\Delta699/699}$ embryos. (E-N) In the lateral neocortex of E12.5
embryos, there is no significant difference in the proportions of Tbr1+ neurons (E-G) and basal
progenitor cells (H-J) between control and *Inpp5e*$^{\Delta/\Delta}$;Gli3$^{\Delta699/+}$ embryos. (K-S) In the E14.5 medial
neocortex, the proportion of Tbr1$^-$Ctip2$^+$ neurons (P) and basal progenitors (S) are significantly
reduced in *Inpp5e*$^{\Delta/\Delta}$;*Gli3*$^{\Delta699/+}$ embryos but not in *Inpp5e*$^{\Delta/\Delta}$;Gli3$^{\Delta699/699}$ embryos. All statistical data
except for the Western blot data (C, D) are presented as means ± 95% confidence intervals (CI);
Mann Whitney (n=5) (G, J) and ONE way ANOVA followed by Tukey's multiple comparison test (J,
M, P); * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Scale bars: 50μm (E, K, L), 100μm (A), and 250μm (F',
K', L'). ctx: cortex; MGE: medial ganglionic eminence; LGE: lateral ganglionic eminence.
949

Supplementary Figure 1: Formation of the telencephalic boundaries in *Inpp5e$^{\Delta/\Delta}$* embryos. (A-F) Formation of the corticoseptal boundary. Expression of the dorsal marker gene *Pax6* (A, B, D, E) and of the ventral marker gene *Dlx2* (C, F) remain restricted to the cortex and septum, respectively, with a sharp expression boundary between both tissues. (G-L) Formation of the pallial/subpallial boundary. While there is a sharp expression boundary between cortex and lateral ganglionic eminence (LGE) in wild-type embryos (G-I), scattered Pax6 and *Dlx2* expressing cells (arrows in J and K) are found in the mutant LGE and cortex, respectively, while the *Dbx1* expression domain characteristic of the ventral pallium (VP) is fuzzier (L). CGE: caudal ganglionic eminence; ctx: cortex; MGE: medial ganglionic eminence; sep: septum; th: thalamus. Scale bars: 200μm.

Supplementary Figure 2: Wnt/β-catenin and Bmp signalling in the dorsomedial telencephalon of E12.5 *Inpp5e^ΔΔ* embryos. (A, B, E, F) Reduced Wnt/β-catenin signalling in *Inpp5e^ΔΔ* embryos. (A, E) *Wnt2b* expression is confined to the cortical hem (h) while there are only few scattered Wnt2b expressing cells in the mutant. (B, F) Graded expression of the Wnt target gene *Axin2* in the dorsal midline is reduced in mutant embryos. (C, D, G, H) Roof plate (rp) expression of *Bmp4* and its target gene *Msx1* are reduced in *Inpp5e^ΔΔ* embryos. Scale bar: 100μm.

Supplementary Figure 3: Expression of cortical progenitor markers in *Inpp5e*$^{\Delta/\Delta}$ embryos. (A-J) Dorsal marker gene expression at patterning stages (E12.5). *Emx1*, *Lhx2*, *Pax6* and *Ngn2* are still expressed in the developing neocortex of *Inpp5e*$^{\Delta/\Delta}$ embryos though the *Lhx2* medial to lateral (B, C, G, H) and the *Pax6*/*Ngn2* lateral to medial (D, E, I, J) expression gradients are flatter. (K-T) Neocortical progenitors express *Lhx2*, *Pax6* and *Ngn2* in E14.5 *Inpp5e*$^{\Delta/\Delta}$ embryos. Note the folding of the neocortex at caudal levels (arrows in P, R and T). CGE: caudal ganglionic eminence; ctx: cortex; MGE: medial ganglionic eminence; LGE: lateral ganglionic eminence; pt: prethalamus; sep: septum; th: thalamus. Scale bars: 200μm.

Supplementary Figure 4: Proportion of mitotic progenitors in *Inpp5e*$^{\Delta/\Delta}$ embryos. (A-H) Proportions of mitotic progenitors in E12.5 control (A, E) and *Inpp5e*$^{\Delta/\Delta}$ embryos (B, F) as revealed by pHH3 (mitotic cells) and PCNA (all progenitor cells) double immunofluorescence. Note the reduction in mitotic basal progenitors in the *Inpp5e*$^{\Delta/\Delta}$ medial neocortex (A, B, D). (I-P) The proportions of apical and basal progenitors is not significantly different in E14.5 control and *Inpp5e*$^{\Delta/\Delta}$ embryos. In all panels, radial glia cells divide at the ventricular surface whereas mitotic basal progenitors locate in abventricular positions. All statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; * $p < 0.05$. Scale bar: 50μm.

Supplementary Figure 5: Cell cycle of cortical progenitors in E12.5 *Inpp5e$^{\Delta/\Delta}$* embryos. (A) Schematic illustrating the BrdU/IdU double labelling strategy to measure S phase (T$_S$) and total cell cycle length (T$_C$). 90 minutes after an initial IdU administration, pregnant females received an intraperitoneal BrdU injection. Embryos are harvested 30 minutes later. (B, C) Double immunofluorescence to detect IdU+ and BrdU+ progenitors. (D) Quantification showing no significant change in T$_S$ and T$_C$. Statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; * p < 0.05. Scale bar: 50μm.

Supplementary Figure 6: Whole mount preparations of E18.5 brains. (A) Control. (B) *Inpp5e*$^{\Delta/\Delta}$ brain. Note the absence of obvious protrusions of the olfactory bulbs (ob) in the mutant. Ctx: cortex. Scale bar: 1 mm.

Supplementary Figure 7: Forebrain malformations in E18.5 *Inpp5e*$^{\Delta/\Delta}$ embryos. (A-F) Coronal sections through the forebrain of E18.5 control (A-C) and *Inpp5e*$^{\Delta/\Delta}$ (D-F) embryos. The asterisk in (D) demarcates a heterotopia in the *Inpp5e* mutant. Note that the mutant lateral neocortex is thinner at intermediate and caudal levels (E, F). The lines in (A) indicate where cortical thickness was measured at medial (m) and lateral (l) levels. (G, H) Quantification of cortical thickness. CC: corpus callosum; ctx: cortex; hip: hippocampus; sep: septum; str: striatum; th: thalamus. Scale bar: 500µm. Statistical data are presented as means ± 95% confidence intervals (CI); Two way ANOVA followed by Sidak multiple comparisons test; n = 4; * p < 0.05; ** p < 0.005; *** p < 0.001.

Supplementary Figure 8: Hippocampus formation in *Inpp5e*$^{\Delta/\Delta}$ embryos. (A-F) Hippocampal marker gene expression in E18.5 control (A-C) and *Inpp5e*$^{\Delta/\Delta}$ embryos (D-F). Expression of *Nrp2* labels the whole hippocampal formation (A), while *Scip1* is expressed in CA1 and in the neocortex (B). Prox1 expression is confined to the dentate gyrus (DG) (C). In *Inpp5e*$^{\Delta/\Delta}$ embryos, these hippocampal markers are expressed but their expression domains are severely reduced or disorganized (D-F).

Supplementary Figure 9: Formation of the corpus callosum in E18.5 *Inpp5e*$^{\Delta/\Delta}$ embryos. (A, B) Coronal section through the telencephalon stained with L1 and GFAP to reveal the corpus callosum (CC) and glial cells, respectively. The corpus callosum is smaller in *Inpp5e*$^{\Delta/\Delta}$ embryos while the glial wedge (GW), the induseum griseum glia (IGG) and the midline zipper glia (MZG) occupy their correct position surrounding the corpus callosum. (C) Quantification of corpus callosum thickness. Statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; * p < 0.05. Scale bar: 250μm.

Supplementary Figure 10: Erk signalling in *Inpp5e*$^{\square/\square}$ embryos. (A, E) Western blot analysis on dorsal telencephalic tissue from E12.5 (A) and E14.5 (E) embryos using anti-phosphoErk$^{T202/Y204}$ and total-Erk antibodies. (B, F) Quantification of the Western blots. A paired t-test was used to evaluate relative pErk/Erk expression levels in four control/*Inpp5e*$^{\square/\square}$ embryo pairs derived from four different litters; statistical data are presented as means ± 95% confidence intervals (CI). (C, D, G, H) Immunohistochemistry for phosphoErk$^{T202/Y204}$ on coronal sections of the telencephalon of E12.5 (C, D) and E14.5 (G, H) embryos. ctx: cortex; MGE: medial ganglionic eminence; LGE: lateral ganglionic eminence; th: thalamus Scale bars: 100 μm.

Supplementary Figure 11: Akt and mTOR signalling in *Inpp5e*$^{\Delta/\Delta}$ embryos. (A, E) Western blot analyses on dorsal telencephalic tissue from E12.5 (A, C, E) and E14.5 (B, D, F) embryos with the indicated antibodies. (A, B) Increased ratio of pAkt^{S473}/total Akt in E12.5 *Inpp5e*$^{\Delta/\Delta}$ embryos but not at E14.5. (C, D) Phosphorylation of S6 at residues S235/S236 is increased in the dorsal telencephalon of E12.5 and E14.5 *Inpp5e*$^{\Delta/\Delta}$ embryos. (E, F) E12.5 *Inpp5e*$^{\Delta/\Delta}$ embryos show an increased phosphorylation of S6 at residues S240/S244. (G-P) *Inpp5e*$^{\Delta/\Delta}$ embryos treated with rapamycin show an augmented proportion of Tbr1+ neurons (G-J, O) and a reduced proportion of Tbr2+ basal progenitors (K-N, P). For quantification of the Western blots, paired t-test were used to evaluate expression levels of the phosphorylated versus unphosphorylated proteins in four control/*Inpp5e*$^{\Delta/\Delta}$ embryo pairs derived from four different litters. Immunofluorescence stainings were analysed by Mann Whitney tests. Statistical data are presented as means ± 95% confidence intervals (CI); n = 4; * $p < 0.05$; *** $p < 0.005$.

Supplementary Figure 12: *Gli3* mRNA expression in Inpp5e mutants. (A, B) *Gli3* in situ hybridization showing *Gli3* mRNA expression in the cortex (ctx) and lateral ganglionic eminence (LGE) of control (A) and *Inpp5e*$^{\Delta/\Delta}$ embryos (B).

Supplementary Figure 13: Rescue of eye development in *Inpp5e*^{Δ/Δ};*Gli3*^{Δ699/+} embryos. (A-C) Side views of the heads of E12.5 embryos with the indicated genotype. *Inpp5e*^{Δ/Δ} embryos lack the eye completely or only form a small remnant whereas eye formation is not affected in *Inpp5e*^{Δ/Δ};*Gli3*^{Δ699/+} embryos. Scale bar: 1mm

Supplementary Figure 14: Rescue of corpus callosum formation in *Inpp5e*^{Δ/Δ};*Gli3*^{Δ699/+} embryos. (A-B) Coronal section through the telencephalon stained with L1 and GFAP to reveal the corpus callosum (CC) and glial cells, respectively. There is no significant difference in the size of the corpus callosum between control and *Inpp5e*^{Δ/Δ};*Gli3*^{Δ699/+} embryos; the glial wedge (GW), the induseum griseum glia (IGG) and the midline zipper glia (MZG) are formed in their correct position. (C) Quantification of corpus callosum thickness. Statistical data are presented as means ± 95% confidence intervals (CI); Mann Whitney tests; n = 4; Scale bar: 250µm.