Reference-based QUantification Of gene

Dispensability (QUOD)

Katharina Sielemann[1,2], Bernd Weisshaar[1,*] and Boas Pucker[1,3]

[1]Genetics and Genomics of Plants, Center for Biotechnology (CeBiTec) & Faculty of

Biology, Bielefeld University, 33615 Bielefeld, Germany

[2]Graduate School DILS, Bielefeld Institute for Bioinformatics Infrastructure (BIBI),

Bielefeld University, 33615 Bielefeld, Germany

[3]Evolution and Diversity, Department of Plant Sciences, University of Cambridge,

Cambridge, UK

*Correspondence: bernd.weisshaar@uni-bielefeld.de

Email addresses:

kfrey@cebitec.uni-bielefeld.de

bpucker@cebitec.uni-bielefeld.de

bernd.weisshaar@uni-bielefeld.de

Abstract

Background

Dispensability of genes in a phylogenetic lineage, e.g. a species, genus, or higher-level clade, is gaining relevance as most genome sequencing projects move to a pangenome level. Most analyses classify genes as core genes, which are present in all investigated individual genomes, and dispensable genes, which only occur in a single or a few investigated genomes. The binary classification as 'core' or 'dispensable' is often based on arbitrary cutoffs of presence/absence in the analysed genomes. Even when extended to 'conditionally dispensable', this concept still requires the assignment of genes to distinct groups.

Results

Here, we present a new method which overcomes this distinct classification by quantifying gene dispensability and present a dedicated tool for reference-based QUantification Of gene Dispensability (QUOD). As a proof of concept, sequence data of 966 *Arabidopsis thaliana* accessions (Ath-966) were processed to calculate a gene-specific dispensability score for each gene based on normalised coverage in read mappings. We validated this score by comparison of highly conserved Benchmarking Universal Single Copy Orthologs (BUSCOs) to all other genes. The average scores of BUSCOs were significantly lower than the scores of non-BUSCOs. Analysis of variation demonstrated lower variation values between replicates of a single accession than between iteratively, randomly selected accessions from the whole dataset Ath-966. Functional investigations revealed defense and antimicrobial response genes among the genes with high-dispensability scores.

Conclusions

Instead of classifying a gene as core or dispensable, QUOD assigns a dispensability

score to each gene. Hence, QUOD facilitates the identification of candidate

dispensable genes, associated with high dispensability scores, which often underlie

lineage-specific adaptation to varying environmental conditions.

Keywords

pangenomics, genomics, dispensability, bioinformatics, bioinformatic tool,

presence/absence variations

Background

Genetic variation is not restricted to single nucleotide polymorphisms or small

insertions and deletions but extends also to (large) structural variations. These

structural variations include copy number variations (CNVs) and presence/absence

variations (PAVs), which can cause substantial variation of the gene content among

individual genomes (1,2). The comparative analysis of multiple genomes of the same

phylogenetic clade allows the identification of PAVs that are connected to phenotypic

traits. In the case of crop species, the identification of PAVs underlying specific

agronomic traits which only occur in a single or a few species is feasible (3–5). As more

highly contiguous genome sequences become available, pangenomes are suitable to

describe and investigate the gene set diversity of a biological clade, e.g. species,

genus or higher (6,7).

Genes of a pangenome are thought to be divided into a core and a dispensable gene

set, the latter is also often referred to as 'accessory' in the literature. Core genes occur

in all investigated genomes, whereas dispensable genes only occur in a single or a

few genomes (8). In eukaryotic pangenome studies, core and dispensable genes are

mostly identified based on sequence similarity e.g. using GET_HOMOLOGUES-EST

Markov clustering (9), OrthoMCL gene family clustering (10) or BLASTN (11).

Sometimes, a third category of 'conditionally dispensable' genes is invoked (12) or

genes might be classified as 'cloud', 'shell', 'soft-core' and 'core' (13) or even as 'core',

'softcore', 'dispensable' and 'private' (14). However, this distinct classification is not

based on the biological dispensability of genes and relies on one or multiple arbitrary

cutoffs. Some studies consider genes as 'core' if these genes occur in at least 90 % of

the investigated genomes (11); in other studies, only genes which are found in all

genomes are part of the core genome (10). In addition, dependency groups might

influence the dispensability of certain genes. The possibility that two genes might be

'replaced' by a specific number of other genes has to be considered. Some genes, of

e.g. a gene family, might be required in a specific proportion and therefore are only

conditionally dispensable (12). Further, assemblies of genomes or transcriptomes

might be incomplete leading to artificially missing genes (15). One way to circumvent

this is to rely only on high-quality reference genome sequences, thus avoiding

additional assemblies which are potential sources of errors.

Here, we present QUOD - a bioinformatic tool to quantify gene dispensability. An *A.*

thaliana dataset of about 1,000 accessions was used to calculate a per gene

dispensability score derived from the coverage of all genes in the given genomes. This

score was validated by comparison of scores of BUSCOs and the functional

investigation of genes with high-dispensability scores. Our tool is easy to use for all

kinds of plant species and datasets of different levels of phylogenetic resolution. QUOD

extends the distinct classification of genes as 'core' and 'dispensable' based on an

arbitrary threshold to a continuous dispensability score.

Methods

Selection and preprocessing of datasets

Genomic reads (FASTQ format) of the investigated genomes were retrieved from the

Sequence Read Archive (SRA) (16) via fastq-dump. BWA-MEM (v.0.7.13) (17) was

applied to map all genomic paired-end Illumina reads to the corresponding reference

genome sequence using default parameters as well as -m to discard secondary

alignments. For *A. thaliana*, all available 1,135 datasets (18) (Additional file 1) were

subjected to a mapping against the AthNd-1_v2c genome sequence (19). The resulting

BAM files of these mappings were subjected to QUOD.

Calculation of gene dispensability scores – QUOD

QUOD calculates a reference-based gene dispensability score for each structurally

annotated gene based on supplied mapping files (BAM) (one per investigated genome)

and a structural annotation of the reference sequence (GFF)

(https://github.com/ksielemann/QUOD). The tool is written in Python3 and consists of

six different components (Additional file 2). During the first part of the analysis, the read

coverage per position (I) as well as the read coverage per gene (II) are calculated. In

the next step, genomes with an average coverage below a given cutoff (default=10)

are discarded and excluded from further analyses (III). Finally, an input matrix is

constructed (IV) and a dispensability score is determined for each gene (V). QUOD

assigns high gene dispensability scores to more likely dispensable genes. Optionally,

the results can be visualized as a colored histogram and a box plot (VI).

The dispensability score (ds(g)) is calculated as follows (cov.=coverage):

$$\text{dispensability score (gene g)} = 1 \Bigg/ \left[\frac{\sum_{n=1}^{N} \left(\frac{\text{average cov. of gene g in genome n}}{\text{average cov. over all genes in genome n}} \right)}{\text{total number of genomes (N)}} \right]$$

Comprehension of the dispensability score composition

For further investigation of the score composition of selected genes of interest, the

script   'score_composition.py'   can   be   used

(https://github.com/ksielemann/QUOD/score_composition.py). As output, a table

including (I) the dispensability score, (II) the average coverage of all investigated

genome sequences, (III) the average coverage of the accessions with the highest and

(IV) lowest 10 % of all coverage values, respectively, (V) the number of accessions

with zero coverage and (VI) the coverage for each accession, separately, is provided.

Further, the coverage distribution for each gene can be visualized in a box plot.

Identification of plastid sequences

Genes of Ath-966 with high similarity to plastid sequences were flagged via BLASTp

(20) of the encoded peptides against all organelle peptide sequences obtained from

the National Center for Biotechnology Information (NCBI). As a control, the sequences

were also searched against themselves. Peptide sequences of Nd-1 with a score ratio

≥ 0.8 were considered plastid-like sequences when comparing BLAST hits against self-

hits (19).

Score comparison between contrasting gene sets

Genes structurally annotated in AthNd-1_v2c were classified with BUSCO v3 (21)

running in protein mode on the encoded peptide sequences using 'brassicales odb10'

(order level) as reference (22). For comparison, BUSCO was additionally executed

using 'chlorophyta odb10' (phylum level) and 'embryophyta odb10' (clade level) as

reference. BUSCOs include single-copy genes and universal genes which are present

in > 90% of all species in the reference dataset and are used to measure the

completeness of assemblies and annotations (21). The scores of BUSCO and non-

BUSCO genes were compared using matplotlib (23) for visualization (box plot) and a

Mann–Whitney U test implemented in the Python package dabest (24) for

determination of the significance

(https://github.com/ksielemann/QUOD/BUSCO_comparison.py). Further, a Levene's

test, implemented in the Python package SciPy (25), was calculated to test for equal

variances among BUSCO genes and non-BUSCO genes. The dispensability score of

non-BUSCO genes might deviate more from the mean as non-BUSCO genes might

be less conserved compared to BUSCO genes and might include multi-copy genes.

Note that for all analyses performed within this study, the score of the size 'infinity'

(detected for one gene) was set to the next highest score to enable calculations.

A list of Nd-1 transposable element (TE) genes, which are Nd-1 gene structures

overlapping with sequences annotated as TEs, was obtained from Pucker *et al.* (19).

First, the score distribution of TE and non-TE genes was determined using a Mann–

Whitney U test implemented in the Python package SciPy (25)

(https://github.com/ksielemann/QUOD/analyse_TE_genes_and_scores.py). Next, the

minimal distance of each gene to its closest TE gene was calculated after extracting

the gene positions from the Nd-1 annotation file. Mixed linear modelling was performed

using Statsmodels v0.12.0 (26) to determine the interaction between the distance to

the closest TE gene and the gene dispensability score

(https://github.com/ksielemann/QUOD/mixed_linear_effects.py).

Correlation of gene length and exon number with the dispensability score

Length and number of exons per gene were extracted from the Nd-1 annotation file.

Linear mixed modelling was performed for gene length, exon number and the gene

dispensability score for the whole dataset Ath-966 as well as for three large *A. thaliana*

gene families (TAPscan (27)), namely MYBs (28), AP2/EREBP (29) and WRKYs (30)

using   Statsmodels   v0.12.0   (26)

(https://github.com/ksielemann/QUOD/mixed_linear_effects.py).

Variation between replicates

A total of 14 genomic datasets of the *A. thaliana* accession Col-0 were received from

the SRA (Additional file 3) to assess the technical variation between replicates of the

same accession. Col-0 was selected for this analysis, because multiple independent

and high-quality datasets are only available for this accession. Each dataset was

mapped to the TAIR10 reference genome sequence using BWA-MEM because a Col-

0 read mapped against AthNd-1_v2c would result in multiple differences caused by

accession-specific differences. The mappings were then subjected to QUOD,

expecting a dispensability score close to one for each gene as there should be no

variability between datasets of the same accession. As the distributions are different

(Kolmogorov-Smirnov test, $p \approx 3e-27$) and the sample size (n) is high, the Levene's

test was selected to test for equal variances, regarding the gene dispensability scores.

The test was applied for (1) the dataset including replicates only and (2) iteratively

(100x), randomly chosen subsets (n=14) of Ath-966

(https://github.com/ksielemann/QUOD/variance_in_repl_test.py).

Functional annotation

194 All genes of the *A. thaliana* Nd-1 genome sequence were annotated via reciprocal best

195 blast hits (RBHs) and best BLAST hits against Araport11 (19). Functional enrichment

196 analyses (PANTHER protein classes and 'biological process' GO terms) were

197 performed using the PANTHER Classification System of the Gene Ontology (31).

Mapper comparison

To evaluate the impact of the read mapping, the results of different mappers were

compared. In addition to BWA-MEM (v.0.7.13; see above) (17), Bowtie2 (v2.4.1;

default parameters) (32) and STAR (v2.5.1b) (33) were selected for this analysis.

STAR parameters required alignments with a similarity of at least 95% over at least

90% of the read pair length. The average coverage values per gene were investigated

for correlation using the Spearman correlation coefficient implemented in the Python

package SciPy (25).

Data Availability

The tool QUOD (QUOD.py) can be downloaded from GitHub

(https://github.com/ksielemann/QUOD; http://doi.org/10.5281/zenodo.4066818). A

data set to test QUOD is available on 'PUB - Publications at Bielefeld University'

(http://doi.org/10.4119/unibi/2946079).

Results

In this study, a bioinformatic tool was developed to calculate a gene-specific

dispensability score based on the normalised coverage in a read mapping. QUOD

allows the quantification of dispensability by calculation of a single score for each gene

(Figure 1). The binary classification of gene dispensability can be compared to the

original method of mRNA detection by endpoint RT-PCR providing only qualitative

results (34–36) which was replaced by quantitative analyses like RNA-Seq.

Gene dispensability scores

The gene dispensability score would initially be dependent on the sequencing depth

per genome. By division of the average coverage of gene g in genome n (N = total

number of investigated genome sequences) by the average coverage over all genes

in genome n, the score is normalised for differences in the sequencing depth of the

investigated genomes. A high value indicates that a gene is likely to be missing in some

genomes and therefore more likely dispensable than a gene with a lower dispensability

score. Due to this quantification approach, this method is not based on an arbitrary

cutoff to determine the core genome and the dispensable genome of any given

pangenome dataset. An example: Using a cutoff of 'gene n occurs in at least 90 % of

all genomes' to be considered a 'core' gene (dark blue), genes 1,2,4 and 6 (dark grey)

would be considered 'dispensable' (Figure 1). However, considering the coverage

(right panel), it is not clear if e.g. gene 1 is truly biologically dispensable. QUOD does

not rely on any thresholds for the classification of genes into 'core' and 'dispensable',

but provides a score based on the normalised coverage in a read mapping. The genes

could theoretically be ranked as well using the percentage of presence/absence of a

gene in the investigated genomes. However, this alternative approach would still rely

on a threshold, e.g. the number of mapped reads for a gene to be considered present

in a genome. This threshold is avoided using the QUOD method.

Figure 1: Illustration of the QUOD method using a fictional dataset. On the left side,

genes are classified as 'core' (dark blue) or 'dispensable' (dark grey) according to a

cutoff. On the right side, gene dispensability is quantified according to a dispensability

score based on the normalised coverage in a read mapping (I-X: investigated

genomes). Coloring of genes (right side) indicates different dispensability scores.

Extremely rare genes, which are absent from most genomes but present in the

reference, can be easily detected using QUOD.

As a proof of concept, *A. thaliana* sequence reads of 1,135 accessions were mapped to the *A. thaliana* Nd-1 genome sequence. All accessions with less than 10-fold read coverage were discarded. The remaining sequencing dataset Ath-966 was analysed with QUOD to calculate a dispensability score for each gene (Figure 2). Genes with high dispensability scores, colored in pink, are considered to be likely dispensable, whereas genes with dispensability scores close to one (dark purple/dark blue) are considered to be core genes.

Figure 2: Distribution of the gene dispensability scores for Ath-966. A) Histogram colored according to the dispensability score. The x-axis represents the dispensability score and the y-axis shows the number of genes in each bin in logarithmic scale. B) Box plot representing the dispensability score (x-axis) of all genes (y-axis). The mean

is represented by the dashed, blue line, the other blue line represents the median of

the scores. C) Genome-wide distribution of genes with different dispensability scores

in *A. thaliana* Nd-1. The colored heatmap shows the respective gene dispensability

scores. There are low (blue) and high (pink) scoring genes clustered in repetitive

regions, including centromeric and telomeric areas. The x-axis represents the size (in

Mbp) of each pseudochromosome in the assembly. The black dots represent the

position of the centromeres of the five chromosomes in the AthNd1_v2c assembly (19).

Genome-wide distribution of the gene dispensability scores

Next, the genome-wide distribution of genes with specific gene dispensability scores

was investigated in *A. thaliana* (Figure 2C). A high plasticity between accessions,

which means a high number of genes with exceptionally high and low scores (pink and

blue), in the (peri-)centromeric regions is visible based on a heatmap (Figure 2C).

As high and low scoring genes cluster in repetitive regions (mainly centromeres), the

score distribution of TE genes was investigated (Additional file 4). Scores of TE genes

are evenly distributed across all dispensability scores. In total, the mean score of TE

genes (mean ds ≈ 1.501) is significantly higher when compared to non-TE genes

(mean ds ≈ 1.168) (Mann-Whitney U test, p ≈ 6E-8), which are more frequent across

scores close to one. Moreover, the minimal distance of each gene to its closest TE

gene and the dispensability scores revealed no relation (Additional file 4).

To test the hypothesis whether genes with higher dispensability scores/more likely

dispensable genes are shorter and whether introns accumulate in core genes, the

correlation of the gene dispensability score with gene length and exon number,

respectively, were determined for the Ath-966 and for three selected gene families

separately. However, no clear trend was detectable (Additional file 5).

Validation of the reliability

Validation of the reliability of the gene dispensability quantification was achieved by

comparison of BUSCOs and non-BUSCOs (Additional file 6). BUSCO genes show on

average slightly lower scores than non-BUSCO genes for all three reference datasets

($p < 0.001$, Mann-Whitney U test). Levene's test was used to test for equal variances.

The results show that the variances for all reference datasets differ significantly

between BUSCO and non-BUSCO genes ($p < 0.001$, Levene's test). Thus, the

deviation of the dispensability score from the respective mean is significantly higher for

non-BUSCO genes in comparison to BUSCO genes.

Further, functional annotation of BUSCO outliers, which are genes of the 'brassicales

odb10' BUSCO gene set with dispensability scores below 0.75 or above 1.25,

revealed, amongst others, several repeat proteins, transmembrane proteins, a 'stress

induced protein', and multiple hypothetical proteins (Additional file 7).

Genes with high and low gene dispensability scores were assessed in more detail.

Among genes with high dispensability scores, several significantly enriched PANTHER

protein classes were detected, e.g. defense/immunity and antimicrobial response

proteins, small GTPases and G-proteins (Figure 3A, left). For genes with dispensability

scores < 0.8, proteins of the extracellular matrix were significantly enriched (Figure 3A,

right). 'Biological process' GO term enrichment revealed several significantly enriched

terms associated with the regulation of cellular processes as well as associated with

response to stimuli among genes with dispensability scores > 2 (Figure 3B, left). Genes

with low dispensability scores show enrichment of primary metabolic processes (Figure

3B, right).

ds ≈ 10.1
≈ 90 % (870 accessions): coverage = 0

Resistance gene, mediates resistance against the bacterial pathogen *Pseudomonas syringae*

Figure 3: Closer investigation of genes with scores >2 (left) and genes with scores <

0.8 (right). A) Significantly enriched PANTHER protein classes (padj < 0.05). B)

Significantly enriched GO biological process terms (padj < 0.05). Abbreviations: p =

process, mp = metabolic process. C) Example for lineage specific adaptation. The

gene NdCCHr1.g3308 (Nd-1 annotation nomenclature) (ds ≈ 10.1) has zero coverage

in 870 accessions (≈ 90%) and is annotated as resistance gene mediating resistance

against the bacterial pathogen *Pseudomonas syringae*.

The function of the 100 genes with the highest gene dispensability scores was

examined in detail for Ath-966 (Additional file 8). Fourteen genes of Ath-966 are

annotated as "disease resistance proteins", whereas seven genes are annotated as

transposons/transposases. Four genes are described as hypothetical proteins and 24

genes have no functional annotation. In addition, an example for lineage specific

adaptation is provided (Figure 3C). The gene NdCCHr1.g3308 has a dispensability

score of approx. 10. For 870 accessions, which are approx. 90 % of Ath-966, no

coverage was detected. The gene is annotated as resistance gene mediating

resistance against the bacterial pathogen *Pseudomonas syringae*.

Next, the variation between replicates of the same accession (Col-0) was determined

(Additional file 9). The variation of the gene dispensability score distribution of the

replicate dataset (one accession) ($\sigma^2 \approx 0.0226$) is significantly lower than the variation

between all iteratively, randomly selected subsets of *A. thaliana* accessions ($\sigma^2 \approx$

0.0392) (Levene's test, $p \approx 4e\text{-}19$). The average coverage per gene using different

read mappers revealed strong correlations in all comparisons (Additional file 10). The

coverage correlations, calculated using Spearman correlation coefficient, between

BWA-MEM and bowtie2 (r ≈ 0.810, p ≈ 0.0), BWA-MEM and STAR (r ≈ 0.814, p ≈ 0.0)

as well as bowtie2 and STAR (r ≈ 0.760, p ≈ 0.0) are similar.

Discussion

QUOD was developed for the quantification of gene dispensability in plant pangenome

datasets. Multiple accessions of several plant species have been sequenced and pose

potential use cases for QUOD (Additional file 11). Dropping sequencing costs will lead

to an increasing availability of comprehensive sequence datasets which would permit

the application of QUOD. Additionally, QUOD is not restricted to plants, but could be

applied to other species (e.g. pig (37)). However, an accurate determination of gene

dispensability scores free of systematic biases might rely on a uniform selection of

genomes from the respective taxonomic group and on uniform read coverage of genes.

In addition, non-random fragmentation of DNA prior to sequencing (38) may cause

biases. The variation among replicates of the same accession (Col-0; σ^2 ≈ 0.0226)

might be attributed to technical biases, e.g. during sequencing library preparation. The

comparison of different read mappers revealed a significant correlation for the average

coverage per gene. Outlier samples, detected by the investigation of the average

coverage per gene using different read mappers, might indicate technical issues. Even

though the correlations are strong, the same tool with the same parameter settings

needs to be used for the read mapping of all compared datasets within one single

QUOD run.

Most genes show dispensability scores close to one as the majority of genes are

widespread across species. The aim of QUOD is mainly the identification of the

'outliers' and therefore the more dispensable genes, which are genes not present in all

genomes. These dispensable genes represent a smaller fraction of the genome than

the core genes. Genome level patterns are expected to be similar for all species.

Further, QUOD is not an alternative to PAV detection methods as groups of genes can

still always be defined using PAV methods, but QUOD provides a quantitative

measurement for this issue.

As already stated in the Introduction, genome assemblies might be incomplete leading

to artificially missing genes (15). One way to circumvent this is to rely on a high-quality

reference genome sequence, thus avoiding additional assemblies which are potential

sources of errors. Recently released telomere-to-telomere assemblies indicate that

these resources will be available for many plant species in the near future (39). Further,

the usage of QUOD with a synthetic reference derived from multiple assemblies is

possible and can be implemented in the future. A graph-based assembly of a

pangenome comprising multiple accessions is already feasible for bacteria (40–42).

However, for large plant genome sequences graph-based pangenome assembly is

computationally expensive and not yet robust for complex structural variants like

inversions(43). Even though there are still several shortcomings, like loss of the sample

information (44), improved methods might be available in the near future and could be

used for the improved quantification of gene dispensability.

Genome-wide distribution of the gene dispensability scores

The genome-wide distribution of all gene dispensability scores (not only BUSCO

genes) of the *A. thaliana* genomes reveals the origin of exceptionally low dispensability

scores (Figure 2). Low scoring genes, which are colored in light blue in Figure 2, might

be TEs and other repeat genes associated with collapsed sequences in the assembly.

An accurate determination of the dispensability scores of these genes might be

possible using ideal genome sequences without any collapsed regions and with

specific read mappings e.g. using high quality long reads. However, low scoring genes

could still be useful to determine amplified TEs and other repeat genes. Moreover, the

genome-wide distribution plot (Figure 2C) shows that high and low scoring genes

cluster in repetitive regions, like e.g. centromeres or telomeres. Very similar

sequences, e.g. members of a gene family or close paralogs, might cause read

mapping errors confounding biases in the dispensability scores of these genes.

Additionally, this can be explained by variation in the recombination rate (45) and active

TEs in these regions. It was previously proposed, that dispensable genes are likely

located closer to TEs which are important factors in genome evolution (9). However,

in the results of our study, TE genes are widely distributed across all dispensability

scores as TEs can occur with variable copy numbers in genomes leading to low scores

and can as well be dispensable. Other studies detected a high number of TEs in the

dispensable genome (46). However, it is possible that only certain TE families might

be truly dispensable. One limitation is the accurate assignment of reads to repetitive

sections of the reference sequence during the read mapping (15). Further, only a

fraction of transposons might be correctly assembled and annotated due to several

computational challenges in highly repetitive and peri-centromeric regions (47).

Therefore, a different strategy might be needed to accurately quantify dispensability of

TEs. A high quality annotation of transposons and a following exclusion of these genes

from the analysis or improved read mapping to the consensus sequence might improve

the results. Again, long reads could be an alternative solution to handle regions which

might be ambiguous in read mappings. Moreover, heterochromatin or genome-purging

mechanisms (48) could influence the gene dispensability scores in these regions.

Additionally some of the low scoring genes were flagged as plastid-like sequences as

original sequencing data from plants contain high amounts of reads originating from

plastid sequences (49,50). Biases due to this plastid read contamination inflate the

coverage of sequences with high similarity to plastid sequences, resulting in an

exceptionally low gene dispensability score.

Validation of the reliability

We validated the reliability of the gene dispensability score by showing that more

conserved BUSCO genes get significantly lower dispensability scores than non-

BUSCO genes (Figure 3A). Based on the distribution of the scores in the boxplot

(Figure 3), the difference between BUSCOs and non-BUSCOs appears small, even

though the difference is significant (U test, p ≈ 4E-113, brassicales reference). It is

important to note that non-BUSCO genes can be highly conserved. Consequently, the

difference is only visible at the group level. The difference in the dispensability scores

of BUSCOs and non-BUSCOs is low as expected, because conserved multiple-copy

genes are not included in the BUSCO gene set (21). Therefore, the variance of the

dispensability scores of non-BUSCO genes is significantly larger than the variance

among BUSCO genes: non-BUSCO genes comprise highly conserved multi-copy

genes as well as less conserved genes. Further, functional annotation of BUSCO

outliers revealed several repeat proteins and transmembrane proteins. Repeat

proteins might lead to read mapping errors and consequently artificial variations in

coverage and dispensability scores. Transmembrane proteins are thought to be

involved in biotic stress response and might not be essential for some accessions and

therefore dispensable (51). This could explain the absence in some genomes resulting

in high dispensability scores of these genes. Therefore, many important, lower-scoring

genes might lie outside of the BUSCO reference set.

Functional annotation of the 100 most likely dispensable genes revealed a high

number of uncharacterised proteins, disease resistance proteins as well as

transposons and transposases in the *A. thaliana* genomes. It is possible that these

genes are undergoing pseudogenization and have not been functionally annotated due

to the lack of a visible phenotype when mutated. TEs were detected in other studies

as contributors to large structural variations between species and individuals and

substantial part of the dispensable genome (46). Previous pangenome analyses also

revealed that the dispensable genome comprises functions like 'defense response',

'diseases resistance', 'flowering time' and 'adaptation to biotic and abiotic stress'

(9,11,13). Comparable results were detected for the enriched protein classes and

'biological process' GO terms (Figure 3A, Figure 3B), even though very general terms,

like 'protein class', give little evidence about the function of genes. Moreover, we

provide a specific example for lineage specific adaptation associated with a high

dispensability score (Figure 3C): a gene mediating resistance against the bacterial

pathogen *Pseudomonas syringae*. Therefore, in depth investigation of genes with high

dispensability scores can result in the identification and characterization of phenotypic

variation (52) and important agronomic traits (13). We envision several applications for

the gene dispensability score generated by QUOD: (1) more accurate prediction if a

gene is associated with a specific trait, (2) development of dependency gene networks,

and (3) improved modeling of the evolutionary value of genes.

Conclusions

459 QUOD (reference-based QUantification Of gene Dispensability) overcomes the

460 problem of labeling genes as 'core' or 'dispensable' through implementation of a

461 quantification approach. Instead of a distinct classification, QUOD provides a ranking

462 of all genes based on assigned gene-specific dispensability scores and therefore does

463 not rely on any thresholds.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The tool QUOD for the reference-based QUantification Of gene Dispensability

(QUOD.py) can be downloaded from GitHub (https://github.com/ksielemann/QUOD;

http://doi.org/10.5281/zenodo.4066818).

Competing interests

The authors declare that they have no competing interests.

Funding

KS is funded by Bielefeld University.

Authors' contributions

KS, BW and BP designed the study, performed the experiments, analysed the data,

and wrote the manuscript. All authors read and approved the final version of this

manuscript.

Acknowledgements

We thank members of Genetics and Genomics of Plants for discussion of preliminary

results. We are very grateful to Janik Sielemann and Nathanael Walker-Hale for helpful

comments on the manuscript. We acknowledge support for the Article Processing

Charge by the Deutsche Forschungsgemeinschaft and the Open Access Publication

Fund of Bielefeld University. We thank the CeBiTec Bioinformatic Resource Facility

team for great technical support.

References

1.   Springer NM, Ying K, Fu Y, Ji T, Yeh C-T, Jia Y, et al. Maize Inbreds Exhibit High

Levels of Copy Number Variation (CNV) and Presence/Absence Variation (PAV)

in Genome Content. Ecker JR, editor. PLoS Genet. 2009 Nov 20;5(11):e1000734.

2.   Scherer SW, Lee C, Birney E, Altshuler DM, Eichler EE, Carter NP, et al.

Challenges and standards in integrating surveys of structural variation. Nat Genet.

2007 Jul;39(S7):S7–15.

3.   Tao Y, Zhao X, Mace E, Henry R, Jordan D. Exploring and Exploiting Pan-

genomics for Crop Improvement. Molecular Plant. 2019 Feb;12(2):156–69.

4.   Lu F, Romay MC, Glaubitz JC, Bradbury PJ, Elshire RJ, Wang T, et al. High-
resolution genetic mapping of maize pan-genome sequence anchors. Nat
Commun. 2015 Nov;6(1):6914.

5.   Swanson-Wagner RA, Eichten SR, Kumari S, Tiffin P, Stein JC, Ware D, et al.
Pervasive gene content variation and copy number variation in maize and its
undomesticated progenitor. Genome Research. 2010 Dec 1;20(12):1689–99.

6.   Tettelin H, Masignani V, Cieslewicz MJ, Donati C, Medini D, Ward NL, et al.
Genome analysis of multiple pathogenic isolates of Streptococcus agalactiae:
Implications for the microbial 'pan-genome'. Proceedings of the National Academy
of Sciences. 2005 Sep 27;102(39):13950–5.

7.   Vernikos G, Medini D, Riley DR, Tettelin H. Ten years of pan-genome analyses.
Current Opinion in Microbiology. 2015 Feb;23:148–54.

8.   Golicz AA, Batley J, Edwards D. Towards plant pangenomics. Plant Biotechnol J.
2016 Apr;14(4):1099–105.

9.   Gordon SP, Contreras-Moreira B, Woods DP, Des Marais DL, Burgess D, Shu S,
et al. Extensive gene content variation in the Brachypodium distachyon pan-
genome correlates with population structure. Nat Commun. 2017 Dec;8(1):2184.

10. Li Y, Zhou G, Ma J, Jiang W, Jin L, Zhang Z, et al. De novo assembly of soybean
wild relatives for pan-genome analysis of diversity and agronomic traits. Nat
Biotechnol. 2014 Oct;32(10):1045–52.

11. Zhao Q, Feng Q, Lu H, Li Y, Wang A, Tian Q, et al. Pan-genome analysis
highlights the extent of genomic variation in cultivated and wild rice. Nat Genet.
2018 Feb;50(2):278–84.

12. Marroni F, Pinosio S, Morgante M. Structural variation and genome complexity: is

dispensable really dispensable? Current Opinion in Plant Biology. 2014

Apr;18:31–6.

13. Golicz AA, Bayer PE, Barker GC, Edger PP, Kim H, Martinez PA, et al. The

pangenome of an agronomically important crop plant Brassica oleracea. Nat

Commun. 2016 Dec;7(1):13390.

14. Liu Y, Du H, Li P, Shen Y, Peng H, Liu S, et al. Pan-Genome of Wild and Cultivated

Soybeans. Cell. 2020 Jul;182(1):162-176.e13.

15. Alkan C, Sajjadian S, Eichler EE. Limitations of next-generation genome

sequence assembly. Nat Methods. 2011 Jan;8(1):61–5.

16. Leinonen, Rasko and Sugawara, Hideaki and Shumway, Martin and International

Nucleotide Sequence Database Collaboration. The sequence read archive.

Nucleic Acids Research. 2010;39(suppl_1):D19–D21.

17. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv preprint arXiv:13033997. 2013;

18. Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, et

al. 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis

thaliana. Cell. 2016 Jul;166(2):481–91.

19. Pucker B, Holtgräwe D, Stadermann KB, Frey K, Huettel B, Reinhardt R, et al. A

chromosome-level sequence assembly reveals the structure of the Arabidopsis

thaliana Nd-1 genome and its gene set. Feltus FA, editor. PLoS ONE. 2019 May

21;14(5):e0216233.

20. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment
search tool. Journal of Molecular Biology. 1990 Oct;215(3):403–10.

21. Simão, Felipe A and Waterhouse, Robert M and Ioannidis, Panagiotis and
Kriventseva, Evgenia V and Zdobnov, Evgeny M. BUSCO: assessing genome
assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 2015;31(19):3210–3212.

22. Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis
P, et al. OrthoDB v9.1: cataloging evolutionary and functional annotations for
animal, fungal, plant, archaeal, bacterial and viral orthologs. Nucleic Acids Res.
2017 Jan 4;45(D1):D744–9.

23. Hunter JD. Matplotlib: A 2D Graphics Environment. Comput Sci Eng.
2007;9(3):90–5.

24. Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. Moving beyond P values:
data analysis with estimation graphics. Nat Methods. 2019 Jul;16(7):565–6.

25. Jones E, Oliphant T, Peterson P, others. SciPy: Open source scientific tools for
Python. 2001;

26. Seabold S, Perktold J. Statsmodels: Econometric and statistical modeling with
python. Proceedings of the 9th Python in Science Conference. 2010;57.

27. Wilhelmsson PKI, Mühlich C, Ullrich KK, Rensing SA. Comprehensive Genome-
Wide Classification Reveals That Many Plant-Specific Transcription Factors
Evolved in Streptophyte Algae. Genome Biology and Evolution. 2017 Dec
1;9(12):3384–97.

28. Stracke R, Werber M, Weisshaar B. The R2R3-MYB gene family in Arabidopsis

thaliana. Current Opinion in Plant Biology. 2001 Oct;4(5):447–56.

29. Feng J-X, Liu D, Pan Y, Gong W, Ma L-G, Luo J-C, et al. An annotation update

via cDNA sequence analysis and comprehensive profiling of developmental,

hormonal or environmental responsiveness of the Arabidopsis AP2/EREBP

transcription factor gene family. Plant Molecular Biology. 2005 Dec;59(6):853–68.

30. Eulgem T, Rushton PJ, Robatzek S, Somssich IE. The WRKY superfamily of plant

transcription factors. Trends in Plant Science. 2000 May;5(5):199–206.

31. Carbon, Seth, Mungall, Chris. Gene Ontology Data Archive [Internet]. Zenodo;

2018 [cited 2020 Sep 29]. Available from: https://zenodo.org/record/3980761

32. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat

Methods. 2012 Apr;9(4):357–9.

33. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:

ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15–21.

34. Wang AM, Doyle MV, Mark DF. Quantitation of mRNA by the polymerase chain

reaction. Proceedings of the National Academy of Sciences. 1989 Dec

1;86(24):9717–21.

35. Gilliland G, Perrin S, Blanchard K, Bunn HF. Analysis of cytokine mRNA and DNA:

detection and quantitation by competitive polymerase chain reaction. Proceedings

of the National Academy of Sciences. 1990;87(7):2725–9.

36. Chiang PW, Song WJ, Wu KY, Korenberg JR, Fogel EJ, Van Keuren ML, et al.

Use of a fluorescent-PCR reaction to detect genomic sequence copy number and

transcriptional abundance. Genome Research. 1996 Oct 1;6(10):1013–26.

37. Tian X, Li R, Fu W, Li Y, Wang X, Li M, et al. Building a sequence map of the pig

pan-genome from multiple de novo assemblies and Hi-C data. Sci China Life Sci.

2020 May;63(5):750–63.

38. Poptsova MS, Il'icheva IA, Nechipurenko DYu, Panchenko LA, Khodikov MV,

Oparina NY, et al. Non-random DNA fragmentation in next-generation

sequencing. Sci Rep. 2015 May;4(1):4532.

39. Miga KH, Koren S, Rhie A, Vollger MR, Gershman A, Bzikadze A, et al. Telomere-

to-telomere assembly of a complete human X chromosome. Nature. 2020 Sep

3;585(7823):79–84.

40. Gautreau G, Bazin A, Gachet M, Planel R, Burlot L, Dubois M, et al.

PPanGGOLiN: Depicting microbial diversity via a partitioned pangenome graph.

Ouzounis CA, editor. PLoS Comput Biol. 2020 Mar 19;16(3):e1007732.

41. Ding W, Baumdicker F, Neher RA. panX: pan-genome analysis and exploration.

Nucleic Acids Research. 2018 Jan 9;46(1):e5–e5.

42. Chan AP, Sutton G, DePew J, Krishnakumar R, Choi Y, Huang X-Z, et al. A novel

method of consensus pan-chromosome assembly and large-scale comparative

analysis reveal the highly flexible pan-genome of Acinetobacter baumannii.

Genome Biol. 2015 Dec;16(1):143.

43. Chen N-C, Solomon B, Mun T, Iyer S, Langmead B. Reducing reference bias

using multiple population reference genomes [Internet]. Bioinformatics; 2020 Mar

[cited   2020   Sep   29].   Available   from:

http://biorxiv.org/lookup/doi/10.1101/2020.03.03.975219

44. Li H, Feng X, Chu C. The design and construction of reference pangenome

graphs. arXiv:200306079 [q-bio] [Internet]. 2020 Mar 12 [cited 2020 Sep 29];

Available from: http://arxiv.org/abs/2003.06079

45. Nachman M. Variation in recombination rate across the genome: evidence and

implications. Current Opinion in Genetics & Development. 2002 Dec 1;12(6):657–

63.

46. Morgante M, Depaoli E, Radovic S. Transposable elements and the plant pan-

genomes. Current Opinion in Plant Biology. 2007 Apr;10(2):149–55.

47. Platt RN, Blanco-Berdugo L, Ray DA. Accurate Transposable Element Annotation

Is Vital When Analyzing New Genome Assemblies. Genome Biol Evol. 2016

Feb;8(2):403–10.

48. Lee S-I, Kim N-S. Transposable Elements and Genome Size Variations in Plants.

Genomics Inform. 2014;12(3):87.

49. Yu J, Hu S, Wang J, Wong GK-S, Li S, Liu B, et al. A draft sequence of the rice

genome (Oryza sativa L. ssp. indica). Science. 2002 Apr 5;296(5565):79–92.

50. Tang J, Xia H, Cao M, Zhang X, Zeng W, Hu S, et al. A Comparison of Rice

Chloroplast Genomes. Plant Physiol. 2004 May;135(1):412–20.

51. Dodds PN, Rathjen JP. Plant immunity: towards an integrated view of plant–

pathogen interactions. Nat Rev Genet. 2010 Aug;11(8):539–48.

52. Hirsch CN, Foerster JM, Johnson JM, Sekhon RS, Muttoni G, Vaillancourt B, et

al. Insights into the Maize Pan-Genome and Pan-Transcriptome. Plant Cell. 2014

Jan;26(1):121–35.

Additional files

Additional file 1 (.tsv): SRA IDs of datasets downloaded to conduct the QUOD analysis

of the *A. thaliana* genomes.

Additional file 2 (.pdf): Illustration of the different components of QUOD.

Additional file 3 (.tsv): SRA IDs of datasets downloaded to conduct the analysis of

replicates (Col-0).

Additional file 4 (.pdf): Distribution of scores of TE genes and non-TE genes and

correlation of the distance to the closest TE gene with the gene dispensability score of

the *A. thaliana* genomes.

Additional file 5 (.pdf): Correlation of gene length and exon number with the

dispensability scores of the *A. thaliana* genomes.

Additional file 6 (.pdf): Comparison of BUSCO analyses for 'chlorophyta', 'brassicales'

and 'embryophyta' as reference.

Additional file 7 (.tsv): Functional annotation of BUSCO outliers (using 'brassicales

odb10' as reference) with a dispensability score smaller than 0.75 or greater than 1.25.

Additional file 8 (.tsv): Functional annotation of the 100 most likely dispensable genes

of the *A. thaliana* genomes.

Additional file 9 (.pdf): Analysis of variance of the gene dispensability score calculated

for replicates of the *A. thaliana* Col-0 accession and iteratively, randomly chosen

subsets of the whole dataset Ath-966.

Additional file 10 (.pdf): Correlation of the average coverage per gene using three

different read mappers: BWA-MEM, bowtie2 and STAR.

Additional file 11 (.tsv): Examples of diploid species where multiple cultivars were

already sequenced.