Amino acids that are well tolerated in random peptides in *E. coli* are enriched in young animal but not young plant genes

Luke Kosinski[1], Nathan Aviles[2], Kevin Gomez[3], Joanna Masel[4]

[1] Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ 85721

[2] Graduate Interdisciplinary Program in Statistics, University of Arizona, Tucson, AZ 85721

[3] Graduate Interdisciplinary Program in Applied Math, University of Arizona, Tucson, AZ 85721

[4] Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721

Corresponding author: Joanna Masel, masel@email.arizona.edu

Abstract

Proteins' great potential for harm, via misfolding and aggregation, does not prevent them from being born *de novo* from non-coding DNA. To investigate how newborn proteins might "first, do no harm", we estimate fitnesses from an experiment that competed *Escherichia coli* lineages that each expressed a unique random peptide. A variety of peptide metrics significantly predict lineage fitness, but almost all this predictive power stems from simple amino acid composition. Low fitness is predicted by hydrophobic and positively charged residues, positive net charge, aggregation prone regions, and under-dispersed hydrophobic residues, while high fitness is predicted by disorder-promoting, negatively charged, small residues, and negative net charge. The same amino acids that predict high fitness in *E. coli* are enriched in young Pfams in animals, but not in plants. To modify Jacques Monod's famous quote, what makes peptides benign in *E.coli* also makes them benign in elephants, but not in eucalyptus.

Introduction

Proteins are the workhorses of the cell, but they are dangerous. The polypeptide backbone is the key structural feature of amyloids, putting all proteins at risk of forming insoluble aggregates (1). The risks of aggregation are exacerbated by the fact that most proteins are expressed at or just beyond their solubility limits (2). Despite these dangers, new protein-coding genes are nevertheless born *de novo* from essentially random sequences (3-5). To be beneficial enough for *de novo* birth, a peptide must first do no harm. Here we ask what properties are predictive of whether a peptide is harmful.

Neme et al. (6) competed over 2 million *Escherichia coli* lineages, each containing a plasmid designed to express a unique random peptide, and tracked lineage frequencies over four days using deep DNA sequencing. This study has been criticized for including too little evidence in support of the beneficial nature of the top candidates (7, 8). But these criticisms do not detract from using the dataset to identify statistical predictors of harm vs. relatively benign effect.

Conveniently, some properties, such as intrinsic structural disorder and aggregation propensity, can be predicted computationally from peptide sequences alone. Because insoluble proteins have been implicated in toxicity and disease (1), we hypothesize that peptides with high intrinsic structural disorder (9, 10) are least likely to be deleterious. Random sequences with high predicted disorder are well-tolerated *in vivo* (11). Existing mouse (12) and Drosophila (13) proteins, which are the product of evolution, are predicted from their amino acid sequences to be more disordered than what would be translated from intergenic controls.

Younger protein-coding sequences should be particularly constrained to first do no harm, as they have had little time to evolve more sophisticated harm-avoidance strategies (14). In support of the idea that high intrinsic structural disorder is an accessible way to avoid harm, young animal and fungal domains (15) and genes (15), and novel overprinted viral genes (16) have higher predicted disorder than their older counterparts.

High predicted intrinsic structural disorder in young genes is driven primarily by amino acid composition, with amino acid order playing a more minor role (12). Fortunately, the dataset of Neme et al. (6) is large enough to look at the frequencies of each amino acid as predictors, rather than assume that existing prediction programs such as IUPred (17, 18) or Waltz (19, 20) integrate amino acid composition (and more) in the best possible way. We can then test whether such programs have additional ability to predict peptide fitness, above and beyond the influence of amino acid content.

How each amino acid frequency predicts peptide harm may also be useful in resolving disputes about the technique of phylostratigraphy, which assigns ages to genes. Detecting homologs is more difficult for fast-evolving sequences, which may be erroneously scored as young (21-23). Disordered proteins tend to be fast evolving (24), casting doubt over findings that young genes are more disordered. If the amino acid preferences of higher fitness random peptides match those of purportedly young protein-coding sequences, this would be evidence that the *de novo* gene birth process, rather than homology detection bias alone, causes trends in protein properties as a function of apparent gene age.

Monod and Jacob (25) famously claimed that "anything found to be true of *E. coli* must also be true of Elephants." We test whether amino acids that are good for random peptides in *E. coli* are enriched in young animal and plant proteins from a recent phylostratigraphy dataset (15) of over 400 different

eukaryotic species. We also test whether published, sequence-based estimators of disorder, aggregation propensity, water solubility, hydrophobic residue clustering etc. predict fitness information not already predicted by amino acid frequencies.

Results

Estimating the fitness effects of random peptides

Assessing predictors of the fitness effects of random peptides requires those fitness effects to be measured accurately and precisely. Neme et al. (6) tracked lineage frequencies over four days. If the counts for days two, three, and four passed a test of statistical significance as all higher or all significantly lower than the count for day one, they categorized the peptide as beneficial or deleterious, respectively. Out of the 1082 peptides with at least five sequencing reads, they found that 277 were beneficial, while 436 were deleterious.

We use a custom maximum likelihood framework (see Materials & Methods) to quantitatively estimate "fitness" and its associated confidence interval. "Fitness" here refers to allele frequency changes over an entire cycle of population growth and dilution, rather than per generation. Our method accounts for increasing mean population fitness over the four days. In the process, we find statistical support for increases in frequency (i.e. fitness − 1.96 × standard error > 1) for 425 peptides (Dataset S1). Our methods are both more sensitive and more specific. We exclude 51 of the original 277, in which the counts for days 2-4 are elevated relative to day 1, satisfying Neme et al.'s (6) criteria for "increasing," but the trend for days 2-4 is downward. 492 peptides had statistical support for decreases in frequency, with support lost for 15 of the original 436 deleterious peptides.

Note that some peptides are pseudoreplicates (see Materials & Methods). We cluster these, and conduct statistics either with cluster as a random effect, or taking a weighted mean across cluster. There were 646 total clusters, of which there was statistical support for increases in frequency for the highest-weighted peptide in 127, and for decreases in 480.

Amino acid counts are the best predictor of random peptide fitness effects

We estimated peptide disorder using IUPred (17, 18), water solubility using CamSol (26), and aggregation propensity using Waltz (19). Fewer than 6% of the random peptides have a predicted transmembrane helix (Dataset S1) from TMHMM (27), so our choice of these predictors is guided by our assumption that the random peptides are predominantly located in the cytosol. Having a predicted transmembrane helix did not in itself predict random peptide fitness effects (P = 0.2, likelihood ratio test relative to mixed model with only the intercept as a fixed effect). In contrast, each of our cytosol-solubility-inspired metrics significantly predicted random peptide fitness (Fig. 1A − 1C), with effects in the predicted direction (more disorder and more solubility are good, more aggregation propensity is bad).

However, when the frequencies of each of the twenty amino acids were also used as predictors, IUPred and CamSol scores were no longer significant predictors of fitness (P = 0.1 and 0.8, respectively, likelihood ratio test), suggesting that their predictive power came merely from being metrics of amino

acid composition. Waltz remained a statistically significant, albeit weak, predictor of fitness in the expected direction (P = 0.03).

We next tried two predictors that focus exclusively on the order rather than the frequency of amino acids in a peptide. First, the contribution of amino acid order to ISD (calculated as ΔISD = original sequence ISD – scrambled sequence ISD), is higher in younger mouse genes (12). However, this was not the case for the very youngest mouse genes, and including ΔISD did not significantly improve our model (P = 0.7).

Second, previous work found that young proteins have their hydrophobic amino acids more clustered together along the primary sequence than expected by chance, while old proteins have them more interspersed (14, 15). This hydrophobic dispersion metric (28, 29), which is normalized to exclude any influence from amino acid composition, not only significantly predicted the fitness of our random peptides on its own (fig. 1D), but including it was also a significant but modest improvement to our model with amino acid composition and Waltz (P = 0.008, likelihood ratio test).

However, while young proteins in nature prefer high clustering (14, 15), we find that benign random peptides prefer low clustering. To explain this apparent contradiction, note that we predict primarily what makes a peptide benign, not what makes it functional. It may be that in order to be functional, some minimal folding and interaction is required, which benefits from some clustering of hydrophobic residues. Hydrophobic clustering, like hydrophobicity itself, may represent a trade-off between higher values being better for folding and hence function, and lower values being better for avoiding misfolding and aggregation (14, 30).

Net charge may be important for disordered sequences (31-33). Negative charge is thought to facilitate solubility (34-36). Positive charge may be harmful; positively charged antimicrobial peptides (37) and toxic nanoparticles (38) interact in destabilizing ways with negatively charged phospholipid membranes and DNA. Net charge includes interaction among amino acids, and so cannot be captured by a simple linear function of amino acids' effects on fitness. Net charge predicted fitness on its own (fig. 1E, P = 2 × 10^{-4}), with positive charge predicting low fitness, as expected. Net charge remained significant when included in our model with amino acid composition, Waltz, and clustering (P = 1 × 10^{-4}, likelihood ratio test). The model was not improved by separate terms for positive and for negative net charge.

Even with hydrophobic clustering, Waltz, and net charge included in the model, most of our predictive power still comes from amino acid composition. Our full model with all significant predictors was only a very modest improvement over a model with just amino acid composition (Fig. 2B vs 2A).

Young animal sequences are enriched for amino acids that increase fitness in random peptides

For ease of interpretation, we calculated the marginal effect of each amino acid on fitness (see supplementary text and Table S2, displayed in fig. 3, x-axis), instead of simply using its slope in a regression model. Isoleucine has the greatest magnitude effect, strongly predicting low fitness. The other four most hydrophobic amino acids (phenylalanine, valine, leucine, and methionine), as well as all of the positively charged amino acids (histidine, lysine, and arginine), also predict low fitness, but not as strongly as isoleucine. Predictors of high fitness were, in order of strongest to weakest: proline, alanine, serine, glycine, and aspartic acid. Proline is a very unusual amino acid that contributes to higher disorder

and low aggregation propensity (39). Alanine and glycine are somewhat hydrophobic but are also the two smallest amino acids, while serine and aspartic acid are hydrophilic, and aspartic acid is negatively charged.

Next we tested whether the amino acids that are relatively benign in random peptides expressed under laboratory conditions in *E. coli* are similar to the amino acids that are found in purportedly young protein-coding sequences in natural populations. We took the latter from James et al. (15) who quantified how amino acid frequencies depend on the age of the protein domain, across over 400 eukaryotic species. As predicted by the de novo gene birth interpretation of phylostratigraphy trends, amino acids that are good for random peptides are enriched among the youngest animal Pfams (fig. 3A).

This prediction was not, however, supported for trends among recent plant domains (fig. 3B) nor among ancient (fig. 3C) domains older than 2.1 billion years. Plant and ancient trends reflect a *de novo* gene birth process that enriches for the most abundant amino acids in their respective lineages, rather than for amino acids that promote intrinsic structural disorder (15). We find that intrinsic structural disorder still predicts harmlessness in *E. coli*, even though we do not find evidence it shaped *de novo* gene birth in its distant ancestors. We also note that intrinsic structural disorder does shape recent *de novo* gene birth in viruses (16).

The correlation in fig. 3A is nevertheless an important external validation that patterns observed using phylostratigraphy can be indicative of the gene birth process, at least in animals. As Monod suggested, what makes a peptide benign in *E. coli* does also make a peptide benign in elephants – with the caveat that this might not be the case in eucalyptus.

Determining which sequences are young is difficult, and may be subject to systematic error (homology detection bias) (21-23). Our ability to externally validate animal phylostratigraphy slopes against an independent dataset generated in *E. coli* is a powerful demonstration that these slopes represent more than mere bias. This conclusion is also supported by the fact that these slopes differ among taxa and are not correlated with amino acid changeability (15).

Fitness is better predicted by amino acid composition than by GC content

Long et al. (40) proposed that selection acts directly on GC content, perhaps due to the three hydrogen bonds of G-C pairs. Amino acids encoded by Gs and Cs tend to promote higher intrinsic structural disorder (10), making it difficult to distinguish between selection for high GC content and selection for disorder-promoting amino acids. To do this, we compare amino acids that always have G or C to those that always have A or T, at both the first and second nucleotide positions in the codon. If selection were for GC nucleotides, we would expect GC to predict high marginal amino acid fitness effects at both positions. But if results are dramatically different at the two positions, this would show that it is selection on amino acid content that drives GC as a correlated trait. Results are statistically significant in the predicted direction at the second position (fig. 4A, P = 0.002, weighted Welch's t-test), and in the predicted direction but not statistically significant at the first (fig. 4B, P > 0.1). The effect size of GC content on fitness could not be statistically distinguished between the first and second position (fig. 4C), with wide and hence inconclusive error bars.

However, amino acid composition is a far stronger predictor of lineage fitness than GC content. To produce easily interpretable R^2 values, we compare weighted fixed-effect regression models, collapsing each cluster of similar peptides (see Materials and Methods) into a single pseudo-datapoint with value

given by the weighted mean and weight given by the sum of weights, instead of modeling within-cluster variance with a random effect term. While GC content is a predictor, offering a significant improvement over an intercept-only model (P = 5 × 10^{-13}, likelihood ratio test for nested models, weighted adjusted R^2 = 0.07), it predicts considerably less than amino acid composition (P = 10^{-23}, weighted adjusted R^2 = 0.17). Adding GC content to the amino acid frequencies-only model offers a modest improvement (P = 0.001, weighted adjusted R^2 values improves from 0.17 to 0.19), while adding amino acid composition to a GC content only model offers a major improvement (P = 10^{-15}, weighted adjusted R^2 improves from 0.07 to 0.19). Adding GC content to a fixed effect analog of our full model (i.e. one that also includes waltz, clustering and net charge) also offers a statistically significant but very minor improvement (P = 0.003, weighted adjusted R^2 improves from about 0.19 to 0.20). These findings suggest that while there may be some direct selection on GC content, the strength of this selection pales in comparison to selection on amino acid composition.

Our fitness predictor is validated on random N-terminal tags

In order to validate our fitness predictor against an external dataset, we tested the ability of our amino-acid-only to predict the fitness effects of shorter random peptides present as N-terminal tags. Goodman et al. (41) tagged the N-prime end of green fluorescent protein (GFP) with 137 different short random sequences (11 amino acids long), to allow for monitoring of random peptide expression level via fluorescence. Frumkin et al. (42) measured the fitness effects of these random peptide-tagged GFPs in *E. coli* using FitSeq (43). For 89 of them, Frumkin et al. (42) were able to calculate a "fitness residual" based on the deviation from the fitness expected from the level of GFP expression. Our fitness predictions were significantly correlated with their fitness residuals (fig. 5, P = 0.004, Spearman's correlation coefficient = 0.31). These findings provide an external validation of the accuracy of our fitness predictor, even in cases where a random segment makes up only a small portion of a total protein.

Discussion

We found that amino acid composition is the primary predictor of the fitness effects of random peptides expressed in *E. coli*. Hydrophobic or positively charged amino acids predict low fitness, while amino acids that are small, negatively charged, and/or promote intrinsic structural disorder predict high fitness. Amino acids that are good for random peptides in *E. coli* are enriched in young proteins in animals, suggesting that harm is avoided in similar ways across diverse lineages. What is truly astonishing is that simply knowing how many of which amino acids are present in a random peptide explained 17% of the variance in our dataset, or 19% if other properties of the protein sequence were also included.

Most studies of random peptides have focused on finding peptides that have specific binding or function (44, 45, e.g. 46). Some were motivated as proof-of-concept that random peptides can exhibit properties of native proteins, such as folding (47-49) and being soluble (50). Others focus on how to increase the percentage of native-like random peptides, e.g. by showing that more hydrophilic random peptide libraries have a higher percentage of stable and soluble peptides (51). Our work has a different intent, identifying properties that make a peptide less likely to be harmful. Neme et al.'s (6) experiment was perfect for this purpose because it used a large library of peptides with diverse properties, and selected under relatively permissive conditions, simply for relative growth rates in the presence of peptide expression. In contrast, a study design such as that of Knopp et al. (52), who selected random peptides that rescue viability in the presence of antibiotics, is less useful for our purposes.

Having a higher proportion of random peptides do no harm is expected to increase the success rate of future screens for peptide with specific properties. Nucleotide sequences with high %GC content tend to have more benign fitness effects, suggesting that higher %GC should be used in future random peptide libraries. However, very high GC content will yield low complexity sequences, which our predictor has not been trained on. The marginal fitness effects of each amino acid might be different in this very different context.

While the library used by Neme et al. (6) was designed to have equal frequencies of each nucleotide in the random region, and thus 50% GC content, the over two million random peptides that had at least one sequencing read had a GC content of ~59% in their random portion. The mean GC content of the peptide clusters we analyzed (see Materials and Methods) was similar, at ~58%, with higher fitness peptides within this group having higher %GC, as discussed in the Results. This suggests that many lower GC content sequences were so harmful that lineages that carried them went extinct prior to detection via sequencing.

Long et al. (40) proposed that there is direct selection for high GC content, as evidenced in part by a preference for amino acids with G or C at the second position of codons, in excess of that predicted from mutation accumulation experiments. Our findings cannot exclude this hypothesis, but show stronger selection on amino acid composition, selection that is capable of driving increased GC content in coding regions as a correlated trait. In intergenic regions, elevated %GC is likely driven mostly by GC-biased gene conversion. However, if driven in part by selection, it is possibly not just on GC content, but also on peptides from non-coding regions translated by error (53, 54). Selection on translation errors is for example strong enough to shape non-coding sequences beyond stop codons in *Saccharomyces cerevisiae* (55).

The fact that IUPred and CamSol have predictive power only when amino acid composition is not controlled for suggest that both predictors are useful primarily as summary statistics of amino acid content. Hydrophobic clustering, Waltz-predicted aggregation, and net charge have only a small amount of predictive power above and beyond that of amino acid composition. Considering the minor predictive power of metrics like hydrophobic clustering, which are independent of amino acid composition by construction, amino acid composition seems to be the primary determinant of peptide fitness, with the effects of amino acid order being subtle.

While driven by amino acid composition, our findings still strongly support the hypothesis that peptides with low structural disorder tend to be harmful. Disorder-promoting amino acids may help a peptide remain soluble even if unfolded. The two smallest amino acids, alanine and glycine, also tend to be benign, perhaps because they are hydrophobic enough to promote some amount of folding but flexible enough to avoid too much hydrophobic residue exposure.

Plants have different trends in amino acid composition change over time than animals, with young genes seeming to prefer readily available amino acids, rather than amino acids that promote intrinsic structural disorder (15). Nevertheless, our finding of consistency between what is benign in *E. coli* and what is benign in animals suggests a deep concordance in what makes a peptide harmful between two apparently disparate branches of life. The forces that drive protein birth therefore appear to share a key similarity between bacteria and Animalia. To modify Monod's famous quote, what is harmful in *E. coli* is also harmful in elephants, but not necessarily in eucalyptus.

Methods

Data retrieval

Neme et al. [6] performed seven experiments where *E. coli* lineages, each with a plasmid containing a unique random peptide, were grown and tracked using deep DNA sequencing. We downloaded sequencing counts from Dryad at http://dx.doi.org/10.5061/dryad.6f356, and obtained amino acid and nucleotide sequences directly from Rafik Neme. Experiment 7 was by far the largest, containing all the peptides that the other six experiments classified as "increasing" or "decreasing," and more. We thus chose to focus on experiment 7 in our work. Experiment 7 consists of the numbers of reads of each random peptide sequence in 5 replicate populations of *E. coli* at 4 time points. We summed across all 5 replicates to obtain a total number of reads for each polypeptide at each time point. Following Neme et al. [6], we took the 1061 peptides out of over one million that had ≥5 reads across all 5 replicates of experiment 7. We further excluded the six peptides that, while meeting the criterion of ≥5 reads, had all of those reads at the same timepoint, leaving 1055 peptides for analysis.

Estimating lineage fitness from random peptide sequencing counts

The expected number of reads λ_{it} of peptide i at times $t=1,2,3,4$ was modeled as:

$$\lambda_{it} = N_t p_{i0} \prod_{k=1}^{t} \frac{\omega_i}{W_{k-1}},$$

where N_t is the observed total number of reads, p_{i0} is the initial frequency of peptide i at the beginning of the experiment (prior to the round of selection used to produce the first measured timepoint $t = 1$), $\frac{\omega_i}{W_t}$ is the fitness of bacteria with peptide i at time t (i.e. their propensity to contribute to the next time point), and W_k is population mean fitness at time k.

The likelihoods of observed peptide counts were estimated from this expectation and two different error models. A Poisson distribution, which captures sampling error alone, was used to generate our initial estimates of p_{io}, ω_i, and W_k (collectively yielding λ_{it}) because it is analytically tractable. Under a Poisson error function, the likelihood of observing n_{it} reads of peptide i at time t is

$$f_{Poiss}(n_{it}|\lambda_{it}) = \frac{\lambda_{it}^{n_{it}} e^{-\lambda_{it}}}{n_{it}!}.$$

To also capture variance inflation κ due to PCR amplification, we used a negative binomial distribution in the Polya form:

$$f_{NBP}\left(n_{it} \mid \lambda_{i,t}, \kappa\right) = \left(\frac{\Gamma\left(n_{it} + \frac{\lambda_{i,t}}{\kappa-1}\right)}{n_{it}! \, \Gamma\left(\frac{\lambda_{i,t}}{\kappa-1}\right)}\right) \left(\frac{1}{\kappa}\right)^{\frac{\lambda_{i,t}}{\kappa-1}} \left(1 - \frac{1}{\kappa}\right)^{n_{it}}$$

where $\Gamma(\,\cdot\,)$ is the gamma function. We used the initial estimates of p_{io}, ω_i, and W_k to numerically fit the negative binomial model. For the specifics of fitting the Poisson and negative binomial models, see

Supporting Information. Weights were calculated, for use in downstream linear models, from this likelihood inference procedure, as the inverse of Fisher information (see Supporting Information).

An existing software package for estimating lineage fitness from sequencing counts is Fit-Seq (43), which captures the amplication of PCR error through a more sophisticated distribution for the number of reads that is derived in the supplementary information of Levy et al. (56). However, Fit-Seq assumes that mean fitness is a simple average of all measured lineages' fitness, requiring all individuals to be tagged and measured. But Neme et al.'s (6) experiment included lineages carrying an empty plasmid, i.e. with the selectable marker but no random peptide. Worse, the proportion of cells with an empty vector can be presumed to increase over time. In the absence of a reliable way to quantify these (Diethard Tautz, personal communication), we instead consider mean population fitness over time to be a set of independent parameters to be fitted.

Clustering non-independent sequences

Upon visual inspection, we found that some peptide sequences were extremely similar, with only one or two amino acid differences; these data points will not contain independent information about the relationship between sequence and fitness. To account for non-independence, we clustered peptides by their Hamming distance, and either took only the peptide whose fitness had the highest weight within its cluster, or took weighted means within clusters, or included cluster in our regression models as a random effect term. Single-link clustering with Hamming distance cutoffs of 6 to 29 amino acids all produced an identical set of 646 clusters for our 1055 peptides. The largest cluster had 228 random peptides, and the second largest had only 13. The vast majorities of clusters contained only 1 sequence (Dataset S1). A few peptides had mutations in their non-random regions; these mutations were counted in our Hamming distance measurements.

Such similar sequences are highly unlikely to arise by chance if the peptides were truly random; $20^{50} \approx 10^{65}$ peptides are possible, far more than the $\sim 2 \times 10^6$ observed. Because we analyze only peptides with at least 5 reads, replicated sequencing error is an unlikely cause. We see the same nearly-identical sequences appearing in every experimental replicate, suggesting either that mutations occurred during Neme et al.'s (6) initial growth phase, or that the "random" peptides synthesized for the experiment are not entirely random. We note that construction of the "random" peptide library involved ligations of a smaller set of "seed" sequences (Rafik Neme, personal communication), introducing non-randomness at this stage.

Predictors of fitness

All peptides are exactly 65 amino acids long with 50 amino acids of random sequence, so there was no need to control for length.

GC content

Many amino acid sequences mapped to several possible nucleotide sequences, as part of the same problem of mutation or non-random construction discussed above. To calculate one GC content for each random peptide, we calculated a simple average of GC content across all the nucleotide sequences in the dataset that map to the peptide with the largest weight in the cluster.

To calculate GC content for the over two million peptides with at least one sequencing read, we took a simple average of the GC content from the random portion of the peptides.

Disorder

IUPred (17, 18) returns an ISD score between zero and one for each amino acid in a sequence, with higher scores indicating greater intrinsic disorder. To calculate an ISD score for each random peptide, we took the average of the scores for the whole sequence (i.e. including non-random parts). A square root transform produced an approximately linear relationship with the log transform of fitness (see Supporting Information for why a log transform of fitness was chosen). IUPred2 was used for all analyses except ΔISD, which used the original IUPred program – differences between the two are minimal (18).

Aggregation propensity

Waltz (19) returns an aggregation score for each amino acid in a sequence. At least five sequential amino acids with a score greater than or equal to five indicates an aggregation-prone region. We scored a peptide as aggregation-prone if it had at least one aggregation-prone region, including contributions from non-random regions.

Solubility

CamSol (26) returns a solubility score for each amino acid in a sequence, as well as a simple average of all scores for a sequence, which CamSol calls a "solubility profile." We used the solubility profile of the full sequences, including non-random regions.

Hydrophobic clustering

We used a Perl script developed by Scott Foy ("calculate_hydrophobic_dispersion.pl" at https://github.com/MaselLab/A-shift-in-aggregation-avoidance-strategy-marks-a-long-term-direction-to-protein-evolution/tree/master/script) to calculate Irback's hydrophobicity dispersion metric (29). This metric splits a sequence into blocks of equal length and calculates dispersion as the variance of the hydrophobicity per block by the mean hydrophobicity per block, normalized to account for differences in mean hydrophobicity and length. We used blocks of 6 amino acids, considered I, L, M, F, and V hydrophobic, and scored whole peptide sequences including non-random regions.

Amino acid composition

We scored amino acid composition by counting frequencies among the 50 amino acids in the random portion of each peptide.

Net charge

Expected net charge was calculated for the whole peptide, including non-random regions, using the "charge" function from the "Peptides" R package version 2.4.1 (57), which uses the Henderson-Hasselbalch equation (58). We used pKscale = "EMBOSS" and pH = 7.5, which is a typical pH for *E. coli* (59).

The values for all the above predictors for each peptide are listed in Dataset S1.

Statistics

All statistical tests were carried out in R (60), with figures generated using "ggplot2" (61). Weighted linear mixed models were implemented using the "lmer" function from the "lme4" package (62), with

cluster as a random effect. See Supporting Information for details, including justification of a log-transform for fitness. When R^2 values were needed, we instead averaged peptides within the same cluster into a combined datapoint, allowing us to avoid the use of random effect term, and took R^2 values from the base R "lm" function.

Code availability

All code is available on GitHub at https://github.com/MaselLab/RandomPeptides.

Acknowledgements

This work was supported by the John Templeton Foundation (39667, 60814) and the National Institutes of Health (GM-104040, T32GM-008659, T32GM-084905). We thank Rafik Neme and Diethard Tautz for sharing their data with us and for graciously answering all our questions regarding their analyses, Dvir Schirman and Tzachi Pilpel for sharing their data with us, Joe Watkins for helpful discussions about our likelihood estimation procedure, and Catherine Weibel for driving the GC content analysis forward.

Works cited

1.   Chiti F & Dobson CM (2017) Protein misfolding, amyloid formation, and human disease: A summary of progress over the last decade. *Annu. Rev. Biochem.* 86:27-68.

2.   Vecchi G*, et al.* (2020) Proteome-wide observation of the phenomenon of life on the edge of solubility. *Proc. Natl. Acad. Sci. U.S.A.* 117(2):1015-1020.

3.   Van Oss SB & Carvunis AR (2019) *De novo* gene birth. *PLoS Genet.* 15(5):e1008160.

4.   McLysaght A & Guerzoni D (2015) New genes from non-coding sequence: the role of de novo protein-coding genes in eukaryotic evolutionary innovation. *Phil. Trans. R. Soc. B.* 370(1678):20140332.

5.   Vakirlis N, Carvunis A-R, & McLysaght A (2020) Synteny-based analyses indicate that sequence divergence is not the main source of orphan genes. *eLife* 9:e53500.

6.   Neme R, Amador C, Yildirim B, McConnell E, & Tautz D (2017) Random sequences are an abundant source of bioactive RNAs or peptides. *Nat. Ecol. Evol.* 1:0127.

7.   Knopp M & Andersson DI (2018) No beneficial fitness effects of random peptides. *Nat. Ecol. Evol.* 2(7):1046-1047.

8.   Weisman CM & Eddy SR (2017) Gene Evolution: Getting Something from Nothing. *Current Biology* 27(13):R661-R663.

9.   Linding R, Schymkowitz J, Rousseau F, Diella F, & Serrano L (2004) A comparative study of the relationship between protein structure and β-aggregation in globular and intrinsically disordered proteins. *J. Mol. Biol.* 342(1):345-353.

10.   Angyan AF, Perczel A, & Gaspari Z (2012) Estimating intrinsic structural preferences of *de novo* emerging random-sequence proteins: Is aggregation the main bottleneck? *FEBS Lett.* 586(16):2468-2472.

11.   Tretyachenko V*, et al.* (2017) Random protein sequences can form defined secondary structures and are well-tolerated in vivo. *Scientific Reports* 7:15449.

12.   Wilson BA, Foy SG, Neme R, & Masel J (2017) Young genes are highly disordered as predicted by the preadaptation hypothesis of *de novo* gene birth. *Nat. Ecol. Evol.* 1(6):0146.

13.   Heames B, Schmitz J, & Bornberg-Bauer E (2020) A continuum of evolving de novo genes drives protein-coding novelty in Drosophila. *J. Mol. Evol.* 38:382-398.

14.   Foy SG, Wilson BA, Bertram J, Cordes MHJ, & Masel J (2019) A shift in aggregation avoidance strategy marks a long-term direction to protein evolution. *Genetics* 211:1345-1355.

15.   James JE*, et al.* (2020) Universal and taxon-specific trends in protein sequences as a function of age. *bioRxiv*:2020.2003.2026.010728.

16. Willis S & Masel J (2018) Gene birth contributes to structural disorder encoded by overlapping genes. *Genetics* 210(1):303-313.

17. Dosztányi Z, Csizmok V, Tompa P, & Simon I (2005) The pairwise energy content estimated from amino acid composition discriminates between folded and intrinsically unstructured proteins. *J. Mol. Biol.* 347(4):827-839.

18. Meszaros B, Erdos G, & Dosztanyi Z (2018) IUPred2A: context-dependent prediction of protein disorder as a function of redox state and protein binding. *Nucleic Acids Res.* 46(W1):W329-W337.

19. Maurer-Stroh S*, et al.* (2010) Exploring the sequence determinants of amyloid structure using position-specific scoring matrices. *Nature Methods* 7(3):237-U109.

20. Fernandez-Escamilla A-M, Rousseau F, Schymkowitz J, & Serrano L (2004) Prediction of sequence-dependent and mutational effects on the aggregation of peptides and proteins. *Nature biotechnology* 22(10):1302-1306.

21. Alba MM & Castresana J (2007) On homology searches by protein Blast and the characterization of the age of genes. *BMC Evol. Biol.* 7:53.

22. Moyers BA & Zhang JZ (2015) Phylostratigraphic Bias Creates Spurious Patterns of Genome Evolution. *Mol. Biol. Evol.* 32(1):258-267.

23. Moyers BA & Zhang JZ (2016) Evaluating Phylostratigraphic Evidence for Widespread De Novo Gene Birth in Genome Evolution. *Mol. Biol. Evol.* 33(5):1245-1256.

24. Chen SCC, Chuang TJ, & Li WH (2011) The Relationships Among MicroRNA Regulation, Intrinsically Disordered Regions, and Other Indicators of Protein Evolutionary Rate. *Mol. Biol. Evol.* 28(9):2513-2520.

25. Monod J & Jacob F (1961) General conclusions - teleonomic mechanisms in cellular metabolism, growth, and differentiation. *Cold Spring Harb. Symp. Quant. Biol.* 26:389-401.

26. Sormanni P, Aprile FA, & Vendruscolo M (2015) The CamSol Method of Rational Design of Protein Mutants with Enhanced Solubility. *J. Mol. Biol.* 427(2):478-490.

27. Krogh A, Larsson B, von Heijne G, & Sonnhammer ELL (2001) Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes. *J. Mol. Biol.* 305(3):567-580.

28. Irback A, Peterson C, & Potthast F (1996) Evidence for nonrandom hydrophobicity structures in protein chains. *Proc. Natl. Acad. Sci. U.S.A.* 93(18):9533-9538.

29. Irback A & Sandelin E (2000) On hydrophobicity correlations in protein chains. *Biophysical Journal* 79(5):2252-2258.

30. Bertram J & Masel J (2020) Evolution rapidly optimizes stability and aggregation in lattice proteins despite pervasive landscape valleys and mazes. *Genetics* 214(4):1047-1057.

31.   Zarin T, *et al.* (2019) Proteome-wide signatures of function in highly diverged intrinsically disordered regions. *eLife* 8:e46883.

32.   Zarin T, Tsai CN, Ba ANN, & Moses AM (2017) Selection maintains signaling function of a highly diverged intrinsically disordered region. *Proc. Natl. Acad. Sci. U.S.A.* 114(8):E1450-E1459.

33.   Mao AH, Crick SL, Vitalis A, Chicoine CL, & Pappu RV (2010) Net charge per residue modulates conformational ensembles of intrinsically disordered proteins. *Proc. Natl. Acad. Sci. U.S.A.* 107(18):8183-8188.

34.   Kramer RM, Shende VR, Motl N, Pace CN, & Scholtz JM (2012) Toward a molecular understanding of protein solubility: increased negative surface charge correlates with increased solubility. *Biophysical Journal* 102(8):1907-1915.

35.   Trevino SR, Scholtz JM, & Pace CN (2008) Measuring and increasing protein solubility. *Journal of Pharmaceutical Sciences* 97(10):4155-4166.

36.   Dubreuil B, Matalon O, & Levy ED (2019) Protein abundance biases the amino acid composition of disordered regions to minimize non-functional interactions. *J. Mol. Biol.* 431(24):4978-4992.

37.   Lei J, *et al.* (2019) The antimicrobial peptides and their potential clinical applications. *Am. J. Transl. Res.* 11(7):3919-3931.

38.   Frohlich E (2012) The role of surface charge in cellular uptake and cytotoxicity of medical nanoparticles. *Int. J. Nanomed.* 7:5577-5591.

39.   Theillet F-X, *et al.* (2013) The alphabet of intrinsic disorder: I. Act like a Pro: On the abundance and roles of proline residues in intrinsically disordered proteins. *Intrinsically Disord. Proteins* 1(1):e24360.

40.   Long H, *et al.* (2018) Evolutionary determinants of genome-wide nucleotide composition. *Nat. Ecol. Evol.* 2(2):237-240.

41.   Goodman DB, Church GM, & Kosuri S (2013) Causes and effects of N-terminal codon bias in bacterial genes. *Science* 342(6157):475-479.

42.   Frumkin I, *et al.* (2017) Gene Architectures that Minimize Cost of Gene Expression. *Mol. Cell* 65(1):142-153.

43.   Li F, Salit ML, & Levy SF (2018) Unbiased fitness estimation of pooled barcode or amplicon sequencing studies. *Cell Syst.* 7(5):521-525.

44.   Frulloni L, *et al.* (2009) Identification of a Novel Antibody Associated with Autoimmune Pancreatitis. *N. Engl. J. Med.* 361(22):2135-2142.

45.   Keefe AD & Szostak JW (2001) Functional proteins from a random-sequence library. *Nature* 410(6829):715-718.

46.   Kaiser CA, Preuss D, Grisafi P, & Botstein D (1987) Many random sequences functionally replace the secretion signal sequence of yeast invertase. *Science* 235(4786):312-317.

47.   Davidson AR & Sauer RT (1994) Folded proteins occur frequently in libraries of random amino-acid sequences. *Proc. Natl. Acad. Sci. U.S.A.* 91(6):2146-2150.

48.   Chiarabelli C*, et al.* (2006) Investigation of de novo totally random biosequences Part II: On the folding frequency in a totally random library of de novo proteins obtained by phage display. *Chem. Biodivers.* 3(8):840-859.

49.   LaBean TH, Butt TR, Kauffman SA, & Schultes EA (2011) Protein folding absent selection. *Genes* 2(3):608-626.

50.   Prijambada ID*, et al.* (1996) Solubility of artificial proteins with random sequences. *FEBS Lett.* 382(1-2):21-25.

51.   Davidson AR, Lumb KJ, & Sauer RT (1995) Cooperatively folded proteins in random sequence libraries. *Nat. Struct. Biol.* 2(10):856-864.

52.   Knopp M*, et al.* (2019) De novo emergence of peptides that confer antibiotic resistance. *mBio* 10(3):e00837-00819.

53.   Wilson BA & Masel J (2011) Putatively noncoding transcripts show extensive association with ribosomes. *Genome Biol. Evol.* 3:1245-1252.

54.   Rajon E & Masel J (2011) Evolution of molecular error rates and the consequences for evolvability. *Proc. Natl. Acad. Sci. U.S.A.* 108(3):1082-1087.

55.   Kosinski LJ & Masel J (2020) Readthrough errors purge cryptic sequences, facilitating the birth of coding sequence. *Mol. Biol. Evol.* Published online ahead of print:msaa046.

56.   Levy SF*, et al.* (2015) Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature* 519(7542):181-186.

57.   Osorio D, Rondon-Villarreal, & Torres R (2015) Peptides: A package for data mining of antimicrobial peptides. *R J.* 7(1):4-14.

58.   Moore DS (1985) Amino-acid and peptide net charges – a simple calculational procedure. *Biochem. Educ.* 13(1):10-11.

59.   Wilks JC & Slonczewski JL (2007) pH of the cytoplasm and periplasm of *Escherichia coli*: Rapid measurement by green fluorescent protein fluorimetry. *J. Bacteriol* 189(15):5601-5607.

60.   Team RDC (2015) R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing).

61.   Wickham H (2016) *ggplot2: Elegant graphics for data analysis* (Springer-Verlag, New York).

62.   Bates D, Maechler M, Bolker B, & Walker S (2015) Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67(1):1-48.

Fig. 1. **Many metrics, on their own, predict peptide fitness effects.** Lines come from weighted linear mixed regression models (see Materials and Methods), where a random effect term controls for pseudoreplication among similar peptides. Point area is proportional to the weight for each peptide's fitness estimate. Fitness is on a log scale. A) ISD (on a square root scale), B) CamSol, C) Waltz, D) hydrophobic clustering, and E) net charge each significantly predict peptide fitness on their own (P = 9 × 10^{-7}, 4 × 10^{-4}, 2 × 10^{-4}, 0.02, and 2 × 10^{-4}, respectively). Boxes in C) represent 25% and 75% quantiles, whiskers represent 9% and 91% quantiles of fitness. Weighted R^2 values for weighted linear regressions of fitness on the predicted fitness from the ISD, CamSol, Waltz, clustering, and net charge models are 0.048, 0.042, 0.0056, 0.018, and 0.0065, respectively.

Fig. 2. **Amino acid counts are the best predictor of random peptide fitness effects.** Amino acid composition is a strong predictor of polypeptide fitness effects (A), and adding hydrophobic clustering, net charge, and Waltz-predicted aggregation propensity to the model (B), while statistically supported, offers little improvement in predictive ability. Weighted R^2 = 0.17 and 0.19, respectively. Obtained R^2 required us to drop the random effect term, so each cluster of non-independent peptides is represented by a single datapoint, with fitness and predicted fitness taken as their weighted average for the cluster, and weight taken as the sum of weights within the cluster. Point area is proportional to these summed weights. Blue lines show these weighted linear regressions of fitness on predicted fitness; orange lines show y=x.

Fig. 3. **Purportedly young animal Pfams are enriched for amino acids that predict high fitness in random peptides.** The y-axis represents the change in the frequency of each amino acid per billion years (BY) for non-transmembrane Pfam domains (15). Change is in number of percentage points, not percentage increase or decrease. For example, a change in glutamic acid content from 5% to 6% is a change of 1 percentage point. The x-axis shows each amino acid's marginal effect on fitness, which is the change in $\log_2$ fitness when one amino acid of the focal type replaces one randomly chosen amino acid of a different type in a random peptide (see Supporting Information). X-axis tick labels are converted to fold change for ease of interpretation (e.g. $2^{0.1} = 1.07$). Fitness effects predict A) animal, but not B) plant, or C) ancient (older than 2.1 billion years) Pfam phylostratigraphy slopes. Correlation coefficients and P-values come from weighted Pearson correlations. Note that the P-value for animal phylostratigraphy slopes vs marginal effects survives a conservative Bonferroni correction (P = 0.003 < 0.05/3 = 0.017).

Figure 4. **Amino acids that are constrained to use Gs and Cs tend to have higher marginal effects on fitness than those constrained to use As and Ts**. The difference is significant for constraints at the second nucleotide position of a codon (A) P = 0.002, weighted Welch's t-test), but not at the first (B) P > 0.1). Point area is proportional to weight, which is calculated as 1 / s.e.(marginal fitness effect)^2, as described in Supporting Information. The y-axis is the same as the Fig. 3 x-axis. C) The mean advantage of amino acids constrained to use GC rather than constrained to use AT is not distinguishable in size between the first and second codon positions. Y-axis gives the difference in the two weighted means of marginal fitness effects from A) and B). Error bars represent 95% confidence intervals on the difference between the means (calculated as difference +/- t_{crit} × se), where t_{crit} ≈ 2.1 is the critical value of the t-statistic with the appropriate degrees of freedom. Weighted Welch's t-test statistic and the corresponding standard error of the difference in means were calculated using the "wtd.t.test" function from the "weights" R package, version 1.0.1.

Figure 5: **Fitness predictions trained on the random peptides of Neme et al. (6) also work for short random tags attached to the N-terminus of GFP**. Predicted fitness comes from our amino acid composition-only mixed model. "Fitness residuals" of N-terminal tags are from Frumkin et al. (2017), and represent the difference between the fitness of the construct and the expected fitness from expression level.