Random peptides rich in small and disorder-promoting amino acids are less likely to be harmful

Luke Kosinski*,††,[1], Nathan Aviles†,††, Kevin Gomez‡, Joanna Masel§

* Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ 85721

† Graduate Interdisciplinary Program in Statistics, University of Arizona, Tucson, AZ 85721

‡Graduate Interdisciplinary Program in Applied Math, University of Arizona, Tucson, AZ 85721

§ Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721

†† These authors contributed equally to this work.

[1]Present address: Critical Path Institute, Tucson, AZ 85718

ORCID IDs: 0000-0002-8146-5955 (L. K.); 0000-0001-9998-4406 (N. A.); 0000-0002-6356-0318 (K. G.); 0000-0002-7398-2127 (J. M.)

13   Running title: Amino acid effects on fitness

14   Corresponding author: Joanna Masel, Department of Ecology and Evolutionary Biology,

15   University of Arizona, 1041 E Lowell St., Tucson, AZ 85721. E-mail: masel@email.arizona.edu

16   +1(520)626-9888

Abstract

Proteins are the workhorses of the cell, yet they carry great potential for harm via misfolding and aggregation. Despite the dangers, proteins are sometimes born *de novo* from non-coding DNA. Proteins are more likely to be born from non-coding regions that produce peptides that do little to no harm when translated than from regions that produce harmful peptides. To investigate which newborn proteins are most likely to "first, do no harm", we estimate fitnesses from an experiment that competed *Escherichia coli* lineages that each expressed a unique random peptide. A variety of peptide metrics significantly predict lineage fitness, but this predictive power stems from simple amino acid frequencies rather than the ordering of amino acids. Amino acids that are smaller and that promote intrinsic structural disorder have more benign fitness effects. We validate that the amino acids that indicate benign effects in random peptides expressed in *E. coli* also do so in an independent dataset of random N-terminal tags in which it is possible to control for expression level. The same amino acids are also enriched in young animal proteins.

Keywords: Experimental evolution, evolvability, fitness estimation, preadaptation, de novo gene birth

Significance statement

Proteins are sometimes born de novo. In an experiment to reproduce this process in *Escherichia coli*, we were able to predict 15% of the variation in random peptide fitness effects from their amino acid frequencies. In contrast, which order the amino acids are in seems to make no difference, adding no predictive power on top of simple amino acid frequencies. Amino acids that are smaller and promote intrinsic structural disorder have more benign fitness effects.

Introduction

Proteins are the workhorses of the cell, but they are dangerous. For example, the polypeptide backbone is the key structural feature of amyloids, putting all proteins at risk of forming insoluble aggregates (Chiti and Dobson 2017), and most proteins are expressed at or just beyond their solubility limits (Vecchi, et al. 2020). Despite these dangers, new protein-coding genes are nevertheless born *de novo* from essentially random sequences (McLysaght and Guerzoni 2015; Van Oss and Carvunis 2019; Vakirlis, Carvunis, et al. 2020). To be beneficial enough for *de novo* birth, a random peptide must first do no serious harm, i.e. it must not be detrimental to the basic functioning of a cell. Here we quantify the degree to which, and the summary statistics via which, a random peptide's propensity for harm can be predicted.

Neme et al. (2017) competed over 2 million *Escherichia coli* lineages, each containing a plasmid designed to express a unique random peptide, and tracked lineage frequencies over four days using deep DNA sequencing. This study has been criticized for providing too little support for the beneficial nature of the top candidates (Weisman and Eddy 2017; Knopp and Andersson 2018). But these criticisms do not detract from using the dataset to identify statistical predictors of serious harm versus relatively benign effect. Neme et al. (2017) used a strong promoter, so evaluation is of tolerance to high expression. Some fitness differences might be due to variation in expression e.g. due to auto-downregulation at the RNA level (Knopp and Andersson 2018) - we will return to this point in the last portion of the Results. Here we pursue analyses based on the hypothesis that the properties of the peptides contribute to variation in fitness among lineages.

Conveniently, computational predictors from peptide sequences alone are available for

some properties, such as intrinsic structural disorder (ISD) and aggregation propensity. Because

insoluble proteins have been implicated in toxicity and disease (Chiti and Dobson 2017) and

peptides with high ISD are less prone to forming insoluble aggregates (Linding, et al. 2004;

Angyan, et al. 2012), we hypothesize that highly disordered peptides are least likely to be

strongly deleterious. Random sequences with high predicted disorder are well-tolerated *in vivo*

(Tretyachenko, et al. 2017). Existing mouse (Wilson, et al. 2017) and Drosophila (Heames, et al.

2020) proteins, which are the product of evolution, are predicted from their amino acid

sequences to be more disordered than what would be translated from intergenic controls.

Younger protein-coding sequences should be particularly constrained to first do no harm,

as they have had little time to evolve more sophisticated harm-avoidance strategies (Foy, et al.

2019). In support of the idea that high ISD is an accessible way to avoid harm, young animal and

fungal domains (James, et al. 2021) and genes (Wilson, et al. 2017; Foy, et al. 2019; James, et al.

2021), and novel overprinted viral genes (Willis and Masel 2018) have higher predicted disorder

than their older counterparts. Some studies have found that putative de novo protein candidates

in *Saccharomyces* yeasts have lower rather than higher ISD (Carvunis, et al. 2012; Basile, et al.

2017; Vakirlis, et al. 2018), but this could be an artifact of proportionately greater inclusion of

non-genes, i.e. those for which there is no selection to retain a full-length translated peptide

(Graur, et al. 2013), within the younger age classes. When Wilson et al. (2017) reanalyzed

Carvunis et al.'s (2012) "proto-genes" of different ages, using more rigorous criteria to exclude

non-genes from the data, the direction of the ISD trend was reversed. The same reversal of trend

following a quality filter was also found by Vakirlis et al. (2018).

Protein ISD is determined both by the overall frequencies of the amino acids, and by the

order in which those amino acids are arranged. Prior research on young genes has suggested that

high predicted ISD in that context is driven primarily by amino acid frequencies, with amino acid

order playing a more minor role (Wilson, et al. 2017). Here we ask the same question with

respect to the role that both amino acid frequencies and their ordering have on peptide fitness,

including through the promotion of ISD. Fortunately, the dataset of Neme et al. (2017) is large

enough to look at the frequencies of each amino acid as predictors, rather than assume that

existing prediction programs such as IUPred (Dosztányi, et al. 2005; Meszaros, et al. 2018) or

Tango (Fernandez-Escamilla, et al. 2004; Linding, et al. 2004; Rousseau, et al. 2006) integrate

all information about both amino acid frequencies and ordering in the best possible way. We can

then test whether the way such programs integrate information about amino acid order gives

them more ability to predict peptide fitness above and beyond the influence of amino acid

frequencies. In doing so, we can estimate the relative roles of amino acid frequencies versus

amino acid ordering in predicting fitness, as well as determine which amino acids have which

effects.

Here we investigate the degree to which amino acid frequencies and amino acid ordering

can predict the fitness effects of random peptides, and if so, which properties are most predictive.

We also investigate whether the properties that help random peptides avoid harm in *E. coli* are

also enriched in young eukaryotic proteins. With our work, we hope to further our understanding

of how peptides avoid harm.

Results

105   Estimating the fitness effects of random peptides

Neme et al. (2017) tracked lineage frequencies over four days, and categorized a peptide as

increasing or decreasing in frequency by comparing the DNA sequencing counts of day 4 to day

1 using DESeq2 (Love, et al. 2014). They report time information only in the form of days of the

experiment, not the potentially non-constant number of generations. Even with this limitation,

their approach fails to use the full richness of data on all 4 days, and applies a significance

threshold that discards quantitative information.

We therefore reanalyze the same data, instead using a custom maximum likelihood

framework (see Materials & Methods) to quantitatively estimate "fitness" and its associated

confidence interval / weight. "Fitness" here refers to allele frequency changes over an entire

cycle of population growth and dilution, rather than per generation. Our method classifies

peptides quantitatively rather than qualitatively. It accounts for the fact that mean population

fitness increases over the four days (see Materials and Methods). Our use of all available data

within an appropriate maximum likelihood framework should make our method more sensitive

and specific for identifying benign vs harmful peptides (see Supplementary Text).

Note that some peptides have such similar sequences that they should be considered

pseudoreplicates (see Materials & Methods). We therefore grouped sequences into clusters based

on sequence similarity (see Materials and Methods). There were 646 total clusters, of which

there was statistical support for increases in frequency for the highest-weighted peptide in 138

clusters, and for decreases in 488 clusters. Some of our statistics use cluster as a random effect

within a linear mixed model. To generate interpretable R^2 values, we instead use a fixed effects

model where we collapse each cluster into a single pseudo-datapoint with value given by the

weighted mean and weight given by the sum of weights.

Due to the low number of sequences in some clusters, the residuals of our mixed model

are subject to shrinkage (Savic and Karlsson 2009). We therefore present only results that remain

significant in a fixed effect model in which we use only the highest weight peptide per cluster.

These models are useful for ensuring our results are robust, but we do not use them as our

primary model because they discard information, and so lose power on external sources of

information (i.e. the correlations in Figs. 2, 3, and 6 are weaker in the fixed effects model).

Most predictive power stems from amino acid frequencies rather than amino acid

order

We estimated peptide disorder using several metrics that contain information both about amino

acid frequencies and about their order: IUPred as an estimate of intrinsic structural disorder

(Dosztányi, et al. 2005; Meszaros, et al. 2018), CamSol as an estimate of water solubility

(Sormanni, et al. 2015), and Tango as an estimate of general aggregation propensity (Fernandez-

Escamilla, et al. 2004; Linding, et al. 2004; Rousseau, et al. 2006). Fewer than 6% of the random

peptides have a predicted transmembrane helix (Dataset S1) from TMHMM (Krogh, et al. 2001),

so our choice of these predictors is guided by our assumption that the random peptides are

predominantly located in the cytosol. Having a predicted transmembrane helix did not in itself

predict random peptide fitness effects (P = 0.2, likelihood ratio test relative to mixed model with

only the intercept as a fixed effect). In contrast, each of our cytosol-solubility-inspired metrics

significantly predicted random peptide fitness (Fig. 1A – 1C), with effects in the predicted

direction (more disorder and more solubility are good, more aggregation propensity is bad).

Adjusted R^2 values for IUPred, CamSol, and Tango are 0.029, 0.029, 0.016, respectively.

Another aggregation predictor, Waltz (Maurer-Stroh, et al. 2010), that specializes in β

aggregates, was also in the predicted direction of aggregation being harmful, but did not meet

statistical significance (P = 0.07).

Metrics that include amino acid frequency + order information

Frequency information only

Fig. 1. **Many metrics predict peptide fitness effects, but most predictive power comes from amino acid frequencies.** Three metrics that combine information on both amino acid frequencies and amino acid order ((A) IUPred, (B) CamSol, and (C) Tango), and two that contain only amino acid frequency information ((D) 19 custom weights on amino acid frequencies and (E) independently estimated disorder propensities used as weights on amino acid frequencies), each significantly predict peptide fitness on their own (P = 7×10^{-4}, 0.003, 0.01, 5×10^{-6}, and 9×10^{-7}, respectively, likelihood ratio test in mixed model compared to intercept-only model). Each point (n = 646) shows a cluster of sequences with similar amino acid sequences (see Methods for more details), and the area displayed for each point is proportional to summed weights across that cluster. Blue lines are fixed-effect weighted linear regressions of cluster fitness on the x-axis predictor, where clusters are collapsed to a single pseudo-datapoint by their weighted average and weights are sums within each cluster. Metrics that include both frequency and order information fail to outperform frequency-only based metrics, as shown by regression slopes (blue lines) and adjusted R^2 values (top right of each figure panel). Adjusted R^2 is calculated as $R^2_{adj} = 1 - (1 - R^2)\frac{n-1}{n-p-1}$, where n is the number of data points and p is the number of degrees of freedom in the predictor. Note that in part D the predictor (model-predicted fitness) is a composite of 19 degrees of freedom that have all been trained on the dataset, so care should be taken in comparing its blue regression line to that of the other panels, each of which has a predictor with only one degree of freedom – this problem does not apply to comparisons of adjusted R^2 values. Seven clusters with fitness greater than 2 are not shown here for ease of visualization; a complete y-axis is shown in supplemental fig. 1. Log-transforming fitness would remove high fitness skew, but

creates systematic heteroscedasticity, and so was not done (supplemental fig. 2). The lack of
systematic heteroscedasticity can be seen here in the form of similar point size across fitness
values.

Next we asked whether these sophisticated metrics offer additional predictive power

beyond mere amino acid frequencies, in the light of prior work on young genes in which little

additional predictive power was found (Wilson, et al. 2017). To do this, we fit a model of fitness

predicted by amino acid frequencies, measured from counts of each amino acid in each peptide's

random region (Fig. 1D), and compared its performance to predictors that incorporate ordering

information (Figs. 1A-C). The amino acid frequency-only model was a significant predictor of

fitness (P = 2 × 10^{-6}, likelihood ratio test compared to an intercept-only mixed model). It is also

more biologically predictive than other metrics, with adjusted R^2 = 0.15 (adjusted to account for

the number of predictors used) being far greater than the values of 0.027, 0.029, and 0.016 found

in Figs 1A-1C. Another, non-adjusted, way to look at biological effect size is the far steeper blue

line in Fig. 1D than in Figs. 1A-1C. Statistically, when the frequencies of each of the twenty

amino acids are used as predictors (Fig. 1D), then IUPred, CamSol, and Tango drop out of the

model (P = 0.2 for each, likelihood ratio test in mixed model, see Supplemental Table S1),

suggesting that their predictive power in Figs. 1A-1C came largely from being metrics of amino

acid frequencies. These results are surprising: one might expect sophisticated metrics that

incorporate both amino acid frequencies and order information to offer more predictive power

and explain a greater range of fitness than simple amino acid frequencies, yet they fail to do so.

Our Fig. 1D model using the frequencies of the 20 amino acids involves 19 degrees of

freedom, while the other metrics we examine involve only one. This makes it inappropriate to

compare the slopes of the blue lines, although adjusted R^2 values can still be compared, and the

fact that the other metrics drop out of a combined model is also informative. We also

investigated a one degree of freedom model of amino acid frequencies, in which relative weights

were specified in advance by a disorder propensity metric that assigns each amino acid a score

based on how frequently it is found in known disordered versus ordered proteins (Theillet, et al.

2013). Average disorder scores over each peptide's random region significantly predicted

random peptide fitness effects in a linear mixed model (Fig. 1E, $P = 9 \times 10^{-7}$, likelihood ratio test

compared to an intercept-only model). The effect size on predicted fitness from the 10% to the

90% quantiles of disorder propensity is 0.50 to 0.72, and the adjusted R^2 for the disorder

propensity model is 0.047. For comparison to other predictors with a single degree of freedom,

the model that got the largest effect size from incorporating both amino acid frequency and order

information was IUPred with an effect size from 0.52 to 0.70, and CamSol had the highest

adjusted R^2 at 0.029. This superiority of the one degree of freedom disorder propensity model

further suggests that predictive power resides with amino acid frequencies, not order

information.

We next investigated a metric of ISD that is comprised of only order information. This

can be calculated as the excess IUPred score of the real peptide in comparison to the average

IUPred score of a set of hypothetical peptides in which the order of the amino acids has been

randomly scrambled; this metric was previously found to be elevated in younger mouse genes

(Wilson, et al. 2017). However, adding this ΔISD metric to our model with amino acid

frequencies as predictors did not significantly improve the model ($P = 0.2$). This further supports

our conclusion that amino acid ordering plays only a minor role compared to amino acid

frequencies in the fitness effects of the random peptides examined here.

Small and disorder-promoting amino acids predict benign fitness effects

Next we quantify the statistical effect of each of the 20 amino acids on fitness. Naively, we could take the associated slope coefficient in a multiple regression model, which represents the change in fitness when one amino acid is gained. But in a peptide of fixed length, one amino acid cannot be gained without another amino acid being lost. We therefore instead calculate the marginal fitness effect of each amino acid on fitness (see supplementary text and Table S2, displayed in fig. 2, y-axis), representing the effect of gaining that amino acid and losing a randomly selected alternative.

Amino acids with smaller volumes (Tsai, et al. 1999) and higher disorder propensities (Theillet, et al. 2013) tend to have higher marginal fitness effects (fig. 2A and 2B; P = 0.01 for both disorder propensity and volume, likelihood ratio test for dropping either term from a weighted regression of marginal effect on both volume and disorder propensity). Volume and disorder propensity together explain over half the weighted variation in marginal fitness effect (weighted adjusted $R^2 = 0.52$). Other properties of amino acids, such as stickiness (Levy, et al. 2012), relative solvent accessibility (Tien, et al. 2013), amino acid cost in *E. coli* (Akashi and Gojobori 2002), frequency in the *E. coli* proteome (either raw or the excess/deficiency expected from number of codons; see Materials and Methods), and isoelectric point (Liu, et al. 2004) did not provide significant explanatory power on top of disorder propensity and volume (all P > 0.1, likelihood ratio test).

Fig. 2. **Amino acids that are small and are associated with disorder promote higher fitness**.
The y-axis shows each amino acid's marginal effect on fitness, which is the change in fitness
when one amino acid of the focal type replaces one randomly chosen amino acid of a different
type in a random peptide (see Supporting Information). Error bars are +/- one standard error. P-
values and correlation coefficients come from weighted Pearson's correlations, where weights
for marginal effects are calculated as 1 / s.e. (marginal fitness effect)^2, and volume and disorder
propensity are unweighted.

Tryptophan is an outlier for amino acid effects on fitness, with a slightly positive effect
on fitness despite both its large volume and its underrepresentation in disordered regions (fig. 2).
Removing tryptophan from a weighted regression model of volume and disorder propensity
predicting marginal effect increases the weighted adjusted R^2 from 0.52 to 0.68. Tryptophan,
encoded only by UGG, is nearly 60% more common among peptides with at least 5 sequence

reads than we expect from the 58% GC content of our dataset. Together with the confidence

interval for its marginal fitness effect including 1, this provides further evidence that tryptophan

is not harmful, making it a distinct outlier, for reasons that are not clear to us.

Isoleucine also stands out, as even more harmful than expected by its large size and order

propensity. Isoleucine's harmful effects may be exacerbated by its role in amyloid formation. For

example, familial amyloid cardiomyopathy is most commonly caused by a valine to isoleucine

mutation (Jacobson, et al. 1997; Dubrey, et al. 2015), suggesting that isoleucine has potential to

form dangerous amyloids where other hydrophobic amino acids do not. Isoleucine, valine, and

leucine are all hydrophobic amino acids with a branched carbon, but only raised isoleucine levels

are associated with a higher risk of Alzheimer's disease (Larsson and Markus 2017), further

suggesting that isoleucine may be especially prone to amyloid formation.

Young animal sequences are enriched for amino acids that increase fitness in

random peptides

As discussed in the Introduction, young domains have higher predicted ISD than their

older counterparts. One hypothesis to explain this observation is that in order to be successfully

born *de novo*, a protein sequence is especially constrained to first do no harm (Wilson, et al.

2017). However, the "phylostratigraphy" approach of assigning ages to genes, on the basis of the

species range in which they have homologs, is contentious. Detecting homologs is more difficult

for fast-evolving sequences, which may be erroneously scored as young (Alba and Castresana

2007; Moyers and Zhang 2015, 2016). Disordered proteins tend to be fast evolving (Chen, et al.

2011), suggesting that highly disordered genes could be misclassified as young because of their

fast evolutionary rate. If the amino acid enrichments of higher fitness random peptides match the

amino acid enrichments of young genes, this would be evidence that the *de novo* gene birth

process, rather than homology detection bias alone, causes trends in protein properties as a

function of apparent gene age.

To test this, we took the slopes of amino acid frequencies with protein domain age from

James et al. (2021), as quantified across over 400 eukaryotic species. As predicted, amino acids

that are good for random peptides are enriched among the youngest animal Pfams (fig. 3A). This

prediction was not, however, supported for trends among recent plant domains (fig. 3B) nor

among ancient (fig. 3C) domains older than 2.1 billion years. Plant and ancient trends reflect a *de*

novo gene birth process that enriches for the most abundant amino acids in their respective

lineages, such as cysteine, rather than for amino acids that promote ISD (James, et al. 2021). It is

interesting that we find that ISD still predicts harmlessness in *E. coli*, even though we do not find

evidence it shaped *de novo* gene birth in its distant ancestors. We also note that ISD does shape

recent *de novo* gene birth in viruses (Willis and Masel 2018).

Fig. 3. **Purportedly young animal Pfams are enriched for amino acids that predict high**

fitness in random peptides. The y-axis represents how the frequency of each amino acid

depends on the age of the sequence in billion years (BY), estimated as a linear regression slope

for non-transmembrane Pfam domains (James, et al. 2021). Frequency is in number of

percentage points, e.g. a difference in glutamic acid content of 5% vs. 6% is a difference of one

percentage point. The x-axis shows each amino acid's marginal effect on fitness, which is the

change in fitness when one amino acid of the focal type replaces one randomly chosen amino

acid of a different type in a random peptide (see Supporting Information). Error bars are +/- one

standard error. Fitness effects predict A) animal, but not B) plant, or C) ancient (older than 2.1

billion years) Pfam phylostratigraphy slopes. Correlation coefficients and P-values come from

weighted Pearson correlations. Note that the P-value for animal phylostratigraphy slopes vs

marginal effects survives a conservative Bonferroni correction (P = 0.002 < 0.05/3 = 0.017).

Fitness is better predicted by amino acid frequencies than by GC content

Long et al. (2018) proposed that selection acts directly on GC content, perhaps due to the three

hydrogen bonds of G-C pairs. Amino acids encoded by Gs and Cs tend to promote higher ISD

(Angyan, et al. 2012), making it difficult to distinguish between selection for high GC content

and selection against harmful amino acids. To attempt to distinguish between the two, we

compare amino acids that always have G or C to those that always have A or T, at both the first

and second nucleotide positions in the codon. If selection were for GC nucleotides, we would

expect GC to predict high marginal amino acid fitness effects at both positions. But if results are

dramatically different at the two positions, this would show that it is selection on amino acid

content that drives GC as a correlated trait. Results are statistically significant in the predicted

direction at the second position (fig. 4A, P = 0.001, weighted Welch's t-test), and in the

predicted direction but not statistically significant at the first (fig. 4B, P = 0.2). The effect size of

GC content on fitness could not be statistically distinguished between the first and second

position (fig. 4C), with wide and hence inconclusive error bars.

Linear models are compatible with partially independent contributions of both amino acid

frequencies and GC content to harm avoidance. GC content, calculated from the random portion

of each peptide's sequence (for more details, see Materials and Methods), is a statistically

significant predictor of fitness by itself (P = 6×10^{-11}, likelihood ratio test for nested fixed-effect

models relative to intercept-only model). However, the weighted adjusted R^2 of 0.06 for GC

content is much lower than the weighted adjusted R^2 of 0.15 for full amino acid frequency

information, i.e., it explains less of the variation than amino acid frequencies. Adding GC

content to the amino acid frequencies-only model offers only a modest improvement (P = 0.003,

332 weighted adjusted R^2 values improves from 0.15 to 0.16), while adding amino acid frequencies

333 to a GC content only model offers a notably larger improvement ($P = 1 \times 10^{-11}$, weighted

334 adjusted R^2 improves from 0.06 to 0.16). These weighted adjusted R^2 values suggest that while

335 there may be some direct selection on GC content, the effect of amino acid frequencies appears

336 to be well beyond what can be explained by GC content.

340 Fig. 4. **Amino acids that are constrained to use Gs and Cs tend to have higher marginal**

341 **effects on fitness than those constrained to use As and Ts**. The difference is significant for

342 constraints at the second nucleotide position of a codon (A) ($P = 0.001$, weighted Welch's t-test),

343 but not at the first (B) ($P = 0.2$). Point area is proportional to weight, which is calculated as 1 /

344 s.e.(marginal fitness effect)^2, as described in Supporting Information. The y-axis is the same as

345 the fig. 2 y-axis and fig. 3 x-axis. C) The mean advantage of amino acids constrained to use GC

346 rather than constrained to use AT is not distinguishable in size between the first and second

347 codon positions. Y-axis gives the difference in the two weighted means of marginal fitness

348 effects from A) and B). Error bars represent 95% confidence intervals on the difference between

349 the means (calculated as difference +/- $t_{crit} \times$ se), where $t_{crit} \approx 2.1$ is the critical value of the t-

350 statistic with the appropriate degrees of freedom. Weighted Welch's t-test statistic and the

corresponding standard error of the difference in means were calculated using the "wtd.t.test"

function from the "weights" R package, version 1.0.1.

To further verify that GC content is not the primary driver of our results, we crudely

controlled for %GC by splitting our dataset into high (> 57.4%) and low ($\leq$ 57.4%) GC random

sequences and repeated the analysis of figure 1D for each subset. The 57.4% cutoff was the

median GC among the pseudo-datapoints corresponding to clusters. High and low %GC data

subsets produced nearly identical fits to each other (fig. 5) and to figure 1D. The adjusted R^2 is

0.11 for the high and 0.09 for the low GC content subsets, with the drop to be expected given the

restriction of range. If %GC were a major driver, we would expect an offset between the

regression lines in fig. 5. Instead, these results are compatible with amino acid frequencies being

the primary driver of results, with %GC being mostly just a correlated trait with little causal

explanatory power.

Fig. 5. **Amino acid frequencies predict fitness in the same way for peptides encoded by high vs. low GC content sequences.** We split data into high (> 57.4%) and low (≤ 57.4%) GC content and separately fit a model to each in which amino acid frequencies predicted peptide fitness, as in figure 1D. Statistical significance is best assessed with GC content as a quantitative rather than a binary predictor (described in text). For the full methodological details of this figure, please refer to the Figure 1 legend.

The same amino acids predict benign fitness effects in random N-terminal tags

The degree to which benign effects are due to low expression of a random peptide, vs. benign effects of the peptide once expressed, remains unclear. We therefore tested the ability of our amino-acid-frequencies-only model, trained on the data of Neme et al. (2017), to predict residual fitness effects in a dataset that controls for peptide expression level. Goodman et al. (2013) tagged the N-prime end of green fluorescent protein (GFP) with 137 different short random sequences (11 amino acids long), allowing random peptide expression level to be measured via

fluorescence. Frumkin et al. (2017) measured the fitness effects of these random peptide-tagged

GFPs in *E. coli* using FitSeq (Li, et al. 2018). For 89 of them, Frumkin et al. (2017) were able to

calculate a "fitness residual" based on the deviation from the fitness expected from the level of

GFP expression. Note that while this fitness residual controls for expression level, it still contains

the cost of inefficient expression in addition to the fitness effect of the peptide itself. Frumkin et

al. (2017) found that low fitness residuals were associated with hydrophobic and expensive-to-

synthesize amino acids. These findings are consistent with our own estimates of direct peptide

effects, as hydrophobic amino acids tend to be order-prone (Linding, et al. 2004; Angyan, et al.

2012), and amino acid volume is highly correlated with synthesis cost in *E. coli* (Pearson's

correlation coefficient = 0.85, P = 2×10^{-6}, cost for amino acid synthesis in *E. coli* taken from

(Akashi and Gojobori 2002)). Indeed, predicted fitness values for Frumkin et al.'s (2017) N-

terminal tags were significantly correlated with their actual fitness residuals (fig. 6). The

consistency between our results and the findings of Frumkin et al. (2017), who control for

peptide expression level, provides an external validation of our results and suggests that our

findings are unlikely to be due to differences in peptide expression levels alone.

Fig. 6. **Fitness predictions trained on the random peptides of Neme et al. (2017) also work**

for short random tags attached to the N-terminus of GFP. Predicted fitness comes from our

amino acid frequencies-only mixed model. "Fitness residuals" of N-terminal tags are from

Frumkin et al. (2017), and represent the difference between the fitness of the construct and the

expected fitness from expression level. $n = 89$.

Discussion

We found that, while many metrics of peptide properties have some ability to predict the fitness

effects of random peptides expressed in *E. coli*, most predictive power stems from amino acid

frequencies. Simply knowing how many of which amino acids are present in these random

peptides can account for 15% of the variance in fitness among lineages, and adding more

predictors to account for amino acid order fails to add more predictive power. This indicates both

the success of our statistical method for inferring fitness, and that mere amino acid frequencies

without amino acid order can be informative of peptide fitness effects. Amino acids that are

small and promote disorder predict high fitness in *E. coli*, and align with those that are enriched

in young protein domains in animals.

Most studies of random peptides have focused on finding peptides that have specific

binding or function (e.g. Kaiser, et al. 1987; Keefe and Szostak 2001; Frulloni, et al. 2009).

Some were motivated as proof-of-concept that random peptides can exhibit properties of native

proteins, such as folding (Davidson and Sauer 1994; Chiarabelli, et al. 2006; LaBean, et al. 2011)

and being soluble (Prijambada, et al. 1996). Others focus on how to increase the percentage of

native-like random peptides, e.g. by showing that more hydrophilic random peptide libraries

have a higher percentage of stable and soluble peptides (Davidson, et al. 1995). Our work has a

different intent, identifying properties that make a peptide less likely to be harmful. Neme et al.'s

(2017) experiment was suitable for this purpose because it used a large library of peptides with

diverse properties, competed lineages growing under permissive conditions, and measured

relative growth rates (i.e. fitness). In contrast, a study design such as that of Knopp et al. (2019),

who selected random peptides that rescue viability in the presence of antibiotics, is less suitable

for our purposes because so few peptides, including harm-avoiding peptides, are viable. Neme et

al.'s (Neme, et al. 2017) study was also convenient because all peptides were the same length –

65 amino acids with 50 amino acids of random sequence – allowing us to neglect length in our

analysis (see (Castro and Tautz 2021) for the analysis of shorter sequences).

Having a higher proportion of random peptides do no harm is expected to increase the

success rate of future screens for peptides with specific properties. Nucleotide sequences with

high %GC content tend to encode peptides with more benign fitness effects, suggesting that

higher %GC should be used in future random peptide libraries. However, very high GC content

will yield low complexity sequences, which our predictor has not been trained on. The marginal

fitness effects of each amino acid might be different in this very different context.

While the library used by Neme et al. (2017) was designed to have equal frequencies of

each nucleotide in the random region, and thus 50% GC content, the over two million random

peptides that had at least one sequencing read had a GC content of ~59% in their random portion.

The mean GC content of the peptide clusters we analyzed (see Materials and Methods) was

similar, at ~58%, with higher fitness peptides within this group having still higher %GC, as

discussed in the Results. The enrichment from 50% GC to ~59% GC might be because many

lower GC content sequences were so harmful that lineages that carried them went extinct prior to

detection via sequencing. Note that sequencing methods vary in their GC bias (Benjamini and

Speed 2012; Choudhari and Grigoriev 2017), making these absolute values less reliable, and

making it important to use internal controls. Our use of four time points acts to some degree as

such an internal control, except to the degree that bias changes with %GC over the course of the

experiment.

Long et al. (2018) proposed that there is direct selection for high GC content, as

evidenced in part by a preference for amino acids with G or C at the second position of codons,

in excess of that predicted from mutation accumulation experiments. Our findings cannot

completely exclude this hypothesis, but show stronger selection on amino acid frequencies,

selection that is capable of driving increased GC content in coding regions as a correlated trait.

In intergenic regions, elevated %GC is likely driven mostly by GC-biased gene conversion.

However, elevated GC content could also be due, at least in part, to selection on peptides from

non-coding regions translated by error (Rajon and Masel 2011; Wilson and Masel 2011).

Selection on translation errors is for example strong enough to shape non-coding sequences

beyond stop codons in *Saccharomyces cerevisiae* (Kosinski and Masel 2020).

Fitness effects in Neme et al. (2017) might not be directly caused by peptide properties

alone but instead by the effect of both nucleotide and peptide properties on expression (Knopp

and Andersson 2018), with lower expression being less harmful. For example, auto-

downregulation at the mRNA level can cause significant difference in expression among

peptides, despite identical promoters. However, the properties we find to be predictive, such as

disorder and amino acid size, are not *a priori* related to auto-downregulation of mRNA in wild-

type *E. coli*, making the latter an unlikely explanation for our findings.

Our findings are consistent with the hypothesis that peptides with low structural disorder

tend to be harmful. We find that this effect is mediated by amino acid frequencies, with no

additional contribution from amino acid ordering, at least none that could be picked up by use of

the program IUPred. Disorder-promoting amino acids may help a peptide remain soluble even if

unfolded. Small amino acids also tend to be benign, perhaps because they are hydrophobic

enough to promote some amount of folding but flexible enough to avoid too much hydrophobic

residue exposure.

Our findings suggest that the easiest way to avoid harm is through disorder and small

size, but do not rule out other strategies that rely on capacity for folding. Indeed, BCS4, a *de*

novo evolved protein in *Saccharomyces cerevisiae*, has a hydrophobic core and is capable of

folding (Bungard, et al. 2017). Vakirlis et al. (2020) found that de novo proteins can emerge as

transmembrane proteins, which need to be lipid soluble, presumably requiring different harm-

avoidance strategies than peptides that are located in the cytosol.

The correlation between the extent to which an amino acid is enriched in young animal

protein domains and its marginal fitness effect in random peptides in *E. coli* is intriguing, and

consistent with a body of literature that *de novo* gene birth favors protein disorder. What is more,

our ability to externally validate animal phylostratigraphy slopes against random peptides in *E.*

coli provides additional support that these slopes represent more than mere bias, in contrast to

suggests that all patterns are due to homology detection bias (Alba and Castresana 2007; Moyers

and Zhang 2015, 2016). That is, if phylostratigraphy trends were due to an artifact such as

homology detection bias, such an artifact would be unlikely to bias our random peptide analysis

in the same direction.

Plants have different trends in amino acid frequencies as a function of sequence age than

animals do, with young genes seeming to prefer readily available amino acids, rather than amino

acids that promote ISD (James, et al. 2021). This could be because: 1) plants are less susceptible

to harm from random peptides, 2) other properties, such as amino acid availability, drive the

emergence of *de novo* genes in plants, or 3) the plant data lack the resolution needed to identify a

correlation with the properties studied here. We do not have the ability to differentiate between

these three possibilities here.

Nevertheless, our finding of consistency between what is benign in *E. coli* and what is

benign in animals suggests the possibility of a deep concordance in what makes a peptide

harmful between two apparently disparate branches of life. The forces that drive protein birth

therefore appear to share a key similarity between bacteria and Animalia. Monod once suggested

that what is true in *E. coli* must also be true in elephants; our work suggests that this may apply

to the properties that tend to make peptides less harmful. To modify Monod's famous quote,

what is harmful in *E. coli* is also harmful in elephants, but not necessarily in eucalyptus.

A major idea in our understanding of proteins is that form – that is, the fold that is

determined by the exact sequence of amino acids – determines function and thus fitness.

However, for these random peptides in *E. coli*, the amino acid content but not the sequence in

which they occur is the main determinant of benign vs harmful effects. Random peptides likely

exist as a diverse ensemble of structural states, but the same is increasingly acknowledged to be

true of functional proteins. While the ordering of amino acids in functional proteins no doubt

plays a role, perhaps mere amino acid frequencies are also more important than once thought in

this context too, especially in structurally disordered protein regions.

Methods

Data retrieval

Neme et al. (2017) performed seven experiments where *E. coli* lineages, each with a plasmid

containing a unique random peptide, were grown and tracked using deep DNA sequencing. We

downloaded sequencing counts from Dryad at http://dx.doi.org/10.5061/dryad.6f356, and

obtained amino acid and nucleotide sequences directly from Rafik Neme. Experiment 7 was by

far the largest with over 4 million reads, more than five times larger than the 2nd largest

experiment and over 1.2 million reads more than all other experiments combined. Experiment 7

contained all the peptides that the other six experiments classified as "increasing" or

"decreasing," and more. Small datasets from these other six experiments yield limited

information because of the need to model changing mean fitness in a population, including not

just the tracked lineages but also cells with an empty vector (see Estimating lineage fitness from

random peptide sequencing counts section). We therefore chose to restrict our analysis to

experiment 7.

Experiment 7 consists of the numbers of reads of each random peptide sequence in 5

replicate populations of *E. coli* at 4 time points. In principle, these independent measurements

can simply be summed, allowing more precise fitness estimation than could be achieved from a

single replicate. To first assess reproducibility, we estimated fitness for each replicate in

isolation. Following Neme et al. (2017), we calculated fitness only when had ≥5 reads available,

excluding cases in which all reads were on the same day. We grouped sequences into "clusters"

to avoid pseudoreplication (see the "Clustering non-independent sequences" section below) and

calculated the weighted mean fitness for each cluster. The five-choose-two pairs of replicates

each shared between 612 and 624 clusters, with Pearson's correlation coefficients ranging from

0.76 to 0.90 for fitness estimates of the same peptide clusters assessed in different replicates. We

therefore summed across all five replicates to obtain a total number of reads for each polypeptide

at each time point, and used these sums.

After summing across the first four replicates of experiment 7, we once again calculated

fitness for peptides with ≥5 reads in our summed dataset, ending up with 1055 peptides out of

over one million, grouped into 646 clusters. The amino acid sequences analyzed, the count data

for each day and replicate of experiment 7, and the quantities we compute for the peptide, are

available in our Supplemental Dataset S1, and the raw reads are available at the European

Nucleotide Archive (ENA) under the project number PRJEB19640.

Neme et al. (2017) used this 5 read cutoff because it is not possible to infer fitness with

any reasonable resolution for individual peptides with fewer than five reads. The dramatic nature

of the data reduction from over a million peptides to only 646 clusters is unsurprising, firstly

because each initial unique peptide was present in only one copy, and secondly because most

peptides are likely deleterious. We note therefore that our analyzed subset of peptides with at

least five reads are certainly non-lethal, and likely less deleterious than the average random

peptide. Nonetheless, we achieved enough resolution to distinguish between more and less

harmful peptides, with remarkably large effect sizes considering the restricted fitness range.

Estimating lineage fitness from random peptide sequencing counts

The expected number of reads λ_{it} of peptide i at times t=1,2,3,4 was modeled as:

549
$$\lambda_{it} = N_t p_{i0} \prod_{k=1}^{t} \frac{\omega_i}{W_{k-1}},$$

where N_t is the observed total number of reads, p_{i0} is the initial frequency of peptide i at the

beginning of the experiment (prior to the round of selection used to produce the first measured

timepoint $t = 1$), $\frac{\omega_i}{W_t}$ is the fitness of bacteria with peptide i at time t (i.e. their propensity to

contribute to the next time point), and W_k is population mean fitness at time k, including bacteria

containing empty vectors for which we have no direct count data.

The likelihoods of observed peptide counts were estimated from this expectation and two

different error models. A Poisson distribution, which captures sampling error for count data

given independence of each read, was used to generate our initial estimates of p_{io}, ω_i, and W_k

(collectively yielding λ_{it}) because it is analytically tractable. Under a Poisson error function, the

likelihood of observing n_{it} reads of peptide i at time t is

560
$$f_{Poiss}(n_{it}|\lambda_{it}) = \frac{\lambda_{it}^{n_{it}} e^{-\lambda_{it}}}{n_{it}!}.$$

561   To relax the independence assumption to also capture variance inflation κ due to PCR

562   amplification, we used a negative binomial distribution in the Polya form:

563
$$f_{NBP}\left(n_{it}\,\middle|\,\lambda_{i,t},\kappa\right) = \left(\frac{\Gamma\left(n_{it} + \frac{\lambda_{i,t}}{\kappa - 1}\right)}{n_{it}!\,\Gamma\left(\frac{\lambda_{i,t}}{\kappa - 1}\right)}\right)\left(\frac{1}{\kappa}\right)^{\frac{\lambda_{i,t}}{\kappa - 1}}\left(1 - \frac{1}{\kappa}\right)^{n_{it}}$$

where $\Gamma(\cdot)$ is the gamma function. We used the initial estimates of p_{io}, ω_i, and W_k to

numerically fit the negative binomial model. For the specifics of fitting the Poisson and negative

binomial models, see Supporting Information. Weights were calculated, for use in downstream

linear models, from this likelihood inference procedure, as the inverse of Fisher information (see

Supporting Information). We use Fisher information to derive an estimate of the standard error

from the curvature of the likelihood surface.

An existing software package for estimating lineage fitness from sequencing counts is

Fit-Seq (Li, et al. 2018), which captures the amplification of PCR error through a more

sophisticated distribution for the number of reads that is derived in the supplementary

information of Levy et al. (2015). However, Fit-Seq assumes that mean fitness is a simple

average of all measured lineages' fitness, requiring all individuals to be tagged and measured.

But Neme et al.'s (2017) experiment included lineages carrying an empty plasmid, i.e. with the

selectable marker but no random peptide. Worse, the proportion of cells with an empty vector

can be presumed to increase over time. In the absence of a reliable way to directly quantify cells

with empty vectors, we instead consider mean population fitness over time to be a set of

independent parameters to be fitted.

Clustering non-independent sequences

Upon visual inspection, we found that some peptide sequences were extremely similar, with only one or two amino acid differences; these data points will not contain independent information about the relationship between sequence and fitness. To account for non-independence, we clustered peptides by their Hamming distance, and either took only the peptide whose fitness had the highest weight within its cluster, or took weighted means within clusters, or included cluster in our regression models as a random effect term. Single-link clustering with Hamming distance cutoffs of 6 to 29 amino acids all produced an identical set of 646 clusters for our 1051 peptides. The largest cluster had 228 random peptides, and the second largest had only 13. The vast majorities of clusters contained only 1 sequence (Dataset S1). A few peptides had mutations in their non-random regions; these mutations were counted in our Hamming distance measurements.

Such similar sequences are highly unlikely to arise by chance if the peptides were truly random; $20^{50} \approx 10^{65}$ peptides are possible, far more than the $\sim 2 \times 10^6$ observed. Because we analyze only peptides with at least 5 reads, replicated sequencing error is an unlikely cause. We see the same nearly-identical sequences appearing in every experimental replicate, suggesting either that mutations occurred during Neme et al.'s (2017) initial growth phase, or that the "random" peptides synthesized for the experiment are not entirely random. We note that construction of the "random" peptide library involved ligations of a smaller set of starting "seed" sequences, introducing non-randomness at this stage.

Predictors of fitness

601   Neme et al. (2017) discarded all peptides with premature stop codons while producing the

602 dataset we analyzed, so they are all exactly 65 amino acids long with 50 amino acids of random

603 sequence. We therefore do not need to control for length, but do need to account for the mean

604 fitness of a population that includes lineages expressing shorter peptides. An updated analysis

605 was recently released that explored peptides of shorter, varying length (Castro and Tautz 2021).

GC content

608 Many amino acid sequences mapped to several possible nucleotide sequences, as part of the

609 same problem of mutation or non-random construction discussed above. To calculate one GC

610 content for each random peptide, we calculated a simple average of GC content across all the

611 nucleotide sequences in the dataset that map to the peptide with the largest weight in the cluster,

612 using only the random portion of the sequence.

Disorder

Protein disorder was measured using IUPred2 (Dosztányi, et al. 2005; Meszaros, et al. 2018) for

amino acid sequences, and using disorder propensity (Theillet, et al. 2013) for individual amino

acids. IUPred2 returns an ISD score between zero and one for each amino acid in a sequence,

with higher scores indicating greater intrinsic disorder. To calculate an ISD score for each

random peptide, we took the average of the scores for the whole sequence (i.e. including non-

random parts). We used a square root transform because it produced a more linear relationship

with fitness than no transform. All measurements referring to ISD or IUPred used IUPred2

except ΔISD, which used the original IUPred program – differences between the two are

minimal (Meszaros, et al. 2018).

Disorder propensity gives each amino acid a score based on the frequency it is found in

disordered proteins relative to ordered proteins (Theillet, et al. 2013). The disorder propensity

score for a peptide was determined by averaging the disorder propensity scores for the amino

acids in the random region. When we use the disorder propensity metric, we explicitly refer to it

as "disorder propensity" and not as "ISD."

Aggregation propensity

Tango (Fernandez-Escamilla, et al. 2004; Linding, et al. 2004; Rousseau, et al. 2006) returns an

aggregation score for each amino acid in a sequence. At least five sequential amino acids with a

score greater than or equal to five indicates an aggregation-prone region. We scored peptide

aggregation propensity as the number of amino acids within regions scored as aggregation-prone,

including contributions from non-random regions.

Solubility

CamSol (Sormanni, et al. 2015) returns a solubility score for each amino acid in a sequence, as

well as a simple average of all scores for a sequence, which CamSol calls a "solubility profile."

We used the solubility profile of the full sequences, including non-random regions.

Amino acid frequencies

We counted frequencies among the 50 amino acids in the random portion of each peptide.

The values for all the above predictors for each peptide are listed in Dataset S1.

E. coli genome amino acid frequencies

Total amino acid frequencies in the *E. coli* genome were calculated from the K-12 reference

proteome on UniProt (Bateman, et al. 2015), found at

https://www.uniprot.org/proteomes/UP000000625. Excess/deficiency in frequency from that

expected by codon number was determined by calculating the expected frequencies from codon

number (e.g. 3/61 for isoleucine) and subtracting from the raw frequencies.

Statistics

All statistical tests were carried out in R version 3.6.3 (R Core Team 2019), with figures

generated using "ggplot2" (Wickham 2016). Weighted linear mixed models were implemented

using the "lmer" function from the "lme4" package (Bates, et al. 2015), with cluster as a random

effect. See Supporting Information for details, including justification of a log-transform for

fitness. When R^2 values were needed, we instead averaged peptides within the same cluster into

a combined datapoint, allowing us to avoid the use of a random effect term. We calculated

adjusted R^2 values using the base R "lm" function. Adjusted R^2 is a modification of R^2 to

penalize additional predictors, and is calculated using the formula:

$$R^2_{adj} = 1 - (1 - R^2)\frac{n-1}{n-p-1},$$

where n are the number of data points and p are the number of predictors. Raw P-values are

reported unless otherwise noted, i.e. without correction for multiple comparisons.

Data and code availability

All code and supplemental data files are available on GitHub at

https://github.com/MaselLab/RandomPeptides. The data originally released in conjunction with

662   the publication of Neme et al. (2017) can be found at Dryad

663   http://dx.doi.org/10.5061/dryad.6f356, and the raw reads are available at the European

664   Nucleotide Archive (ENA) under the project number PRJEB19640.

Acknowledgements

This work was supported by the John Templeton Foundation (39667, 60814) and the National

Institutes of Health (GM-104040, T32GM-008659, T32GM-084905). We thank Rafik Neme and

Diethard Tautz for sharing their data with us and for graciously answering all our questions

regarding their analyses, Dvir Schirman and Tzachi Pilpel for sharing their data with us, Joe

Watkins for helpful discussions about our likelihood estimation procedure, and Catherine Weibel

for driving the GC content analysis forward.

Works cited

Akashi H, Gojobori T. 2002. Metabolic efficiency and amino acid composition in the proteomes

of Escherichia coli and Bacillus subtilis. Proc. Natl. Acad. Sci. U.S.A. 99:3695-3700.

Alba MM, Castresana J. 2007. On homology searches by protein Blast and the characterization

of the age of genes. BMC Evol. Biol. 7:53.

Angyan AF, Perczel A, Gaspari Z. 2012. Estimating intrinsic structural preferences of *de novo*

emerging random-sequence proteins: Is aggregation the main bottleneck? FEBS Lett.

586:2468-2472.

Basile W, Sachenkova O, Light S, Elofsson A. 2017. High GC content causes orphan proteins to

be intrinsically disordered. PLoS Computational Biology 13:e1005375.

Bateman A, Martin MJ, O'Donovan C, Magrane M, Apweiler R, Alpi E, Antunes R, Arganiska J,

Bely B, Bingley M, et al. 2015. UniProt: a hub for protein information. Nucleic Acids

Res. 43:D204-D212.

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4.

J. Stat. Softw. 67:1-48.

Benjamini Y, Speed TP. 2012. Summarizing and correcting the GC content bias in high-

throughput sequencing. Nucleic Acids Res. 40:e72.

Bungard D, Copple JS, Yan J, Chhun JJ, Kumirov VK, Foy SG, Masel J, Wysocki VH, Cordes

MHJ. 2017. Foldability of a natural de novo evolved protein. Structure 25:1687-1696.

Carvunis A-R, Rolland T, Wapinski I, Calderwood MA, Yildirim MA, Simonis N, Charloteaux

B, Hidalgo CA, Barbette J, Santhanam B. 2012. Proto-genes and de novo gene birth.

Nature 487:370-374.

Castro JF, Tautz D. 2021. The Effects of Sequence Length and Composition of Random

Sequence Peptides on the Growth of E. coli Cells. Genes 12:1913.

Chen SCC, Chuang TJ, Li WH. 2011. The Relationships Among MicroRNA Regulation,

Intrinsically Disordered Regions, and Other Indicators of Protein Evolutionary Rate. Mol.

Biol. Evol. 28:2513-2520.

Chiarabelli C, Vrijbloed JW, De Lucrezia D, Thomas RM, Stano P, Polticelli F, Ottone T, Papa

E, Luisi PL. 2006. Investigation of de novo totally random biosequences Part II: On the

folding frequency in a totally random library of de novo proteins obtained by phage

display. Chem. Biodivers. 3:840-859.

Chiti F, Dobson CM. 2017. Protein misfolding, amyloid formation, and human disease: A

summary of progress over the last decade. Annu. Rev. Biochem. 86:27-68.

Choudhari S, Grigoriev A. 2017. Phylogenetic heatmaps highlight composition biases in

sequenced reads. Microorganisms 5:4.

Davidson AR, Lumb KJ, Sauer RT. 1995. Cooperatively folded proteins in random sequence

libraries. Nat. Struct. Biol. 2:856-864.

Davidson AR, Sauer RT. 1994. Folded proteins occur frequently in libraries of random amino-

acid sequences. Proc. Natl. Acad. Sci. U.S.A. 91:2146-2150.

Dosztányi Z, Csizmok V, Tompa P, Simon I. 2005. The pairwise energy content estimated from

amino acid composition discriminates between folded and intrinsically unstructured

proteins. J. Mol. Biol. 347:827-839.

Dubrey S, Ackermann E, Gillmore J. 2015. The transthyretin amyloidoses: advances in therapy.

Postgraduate Medical Journal 91:439-448.

Fernandez-Escamilla A-M, Rousseau F, Schymkowitz J, Serrano L. 2004. Prediction of

sequence-dependent and mutational effects on the aggregation of peptides and proteins.

Nature biotechnology 22:1302-1306.

Foy SG, Wilson BA, Bertram J, Cordes MHJ, Masel J. 2019. A shift in aggregation avoidance

strategy marks a long-term direction to protein evolution. Genetics 211:1345-1355.

Frulloni L, Lunardi C, Simone R, Dolcino M, Scattolini C, Falconi M, Benini L, Vantini I,

Corrocher R, Puccetti A. 2009. Identification of a Novel Antibody Associated with

Autoimmune Pancreatitis. N. Engl. J. Med. 361:2135-2142.

Frumkin I, Schirman D, Rotman A, Li F, Zahavi L, Mordret E, Asraf O, Wu S, Levy SF, Pilpel

Y. 2017. Gene Architectures that Minimize Cost of Gene Expression. Mol. Cell 65:142-

153.

Goodman DB, Church GM, Kosuri S. 2013. Causes and effects of N-terminal codon bias in

bacterial genes. Science 342:475-479.

Graur D, Zheng YC, Price N, Azevedo RBR, Zufall RA, Elhaik E. 2013. On the Immortality of

Television Sets: "Function" in the Human Genome According to the Evolution-Free

Gospel of ENCODE. Genome Biol. Evol. 5:578-590.

Heames B, Schmitz J, Bornberg-Bauer E. 2020. A continuum of evolving de novo genes drives

protein-coding novelty in Drosophila. J. Mol. Evol. 38:382-398.

Jacobson DR, Pastore RD, Yaghoubian R, Kane I, Gallo G, Buck FS, Buxbaum JN. 1997.

Variant-sequence transthyretin (isoleucine 122) in late-onset cardiac amyloidosis in black

Americans. N. Engl. J. Med. 336:466-473.

James JE, Willis SM, Nelson PG, Weibel C, Kosinski LJ, Masel J. 2021. Universal and taxon-

specific trends in protein sequences as a function of age. eLife 10:e57347.

Kaiser CA, Preuss D, Grisafi P, Botstein D. 1987. Many random sequences functionally replace

the secretion signal sequence of yeast invertase. Science 235:312-317.

Keefe AD, Szostak JW. 2001. Functional proteins from a random-sequence library. Nature

410:715-718.

Knopp M, Andersson DI. 2018. No beneficial fitness effects of random peptides. Nat. Ecol.

Evol. 2:1046-1047.

Knopp M, Gudmundsdottir JS, Nilsson T, Konig F, Warsi O, Rajer F, Adelroth P, Andersson DI.

2019. De novo emergence of peptides that confer antibiotic resistance. mBio 10:e00837-

00819.

Kosinski LJ, Masel J. 2020. Readthrough errors purge cryptic sequences, facilitating the birth of

coding sequence. Mol. Biol. Evol. 37:1761–1774.

Krogh A, Larsson B, von Heijne G, Sonnhammer ELL. 2001. Predicting transmembrane protein

topology with a hidden Markov model: Application to complete genomes. J. Mol. Biol.

305:567-580.

LaBean TH, Butt TR, Kauffman SA, Schultes EA. 2011. Protein folding absent selection. Genes

2:608-626.

Larsson SC, Markus HS. 2017. Branched-chain amino acids and Alzheimer's disease: a

Mendelian randomization analysis. Scientific Reports 7:13604.

Levy ED, De S, Teichmann SA. 2012. Cellular crowding imposes global constraints on the

chemistry and evolution of proteomes. Proc. Natl. Acad. Sci. U.S.A. 109:20461-20466.

Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G. 2015. Quantitative

evolutionary dynamics using high-resolution lineage tracking. Nature 519:181-186.

Li F, Salit ML, Levy SF. 2018. Unbiased fitness estimation of pooled barcode or amplicon

sequencing studies. Cell Syst. 7:521-525.

Linding R, Schymkowitz J, Rousseau F, Diella F, Serrano L. 2004. A comparative study of the

relationship between protein structure and β-aggregation in globular and intrinsically

disordered proteins. J. Mol. Biol. 342:345-353.

Liu HX, Zhang RS, Yao XJ, Liu MC, Hu ZD, Fan BT. 2004. Prediction of the isoelectric point

of an amino acid based on GA-PLS and SVMs. Journal of Chemical Information and

Computer Sciences 44:161-167.

Long H, Sung W, Kucukyildirim S, Williams E, Miller SF, Guo W, Patterson C, Gregory C,

Strauss C, Stone C, et al. 2018. Evolutionary determinants of genome-wide nucleotide

composition. Nat. Ecol. Evol. 2:237-240.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 15:550.

Maurer-Stroh S, Debulpaep M, Kuemmerer N, de la Paz ML, Martins IC, Reumers J, Morris KL,

Copland A, Serpell L, Serrano L, et al. 2010. Exploring the sequence determinants of

amyloid structure using position-specific scoring matrices. Nature Methods 7:237-242.

McLysaght A, Guerzoni D. 2015. New genes from non-coding sequence: the role of de novo

protein-coding genes in eukaryotic evolutionary innovation. Phil. Trans. R. Soc. B.

370:20140332.

Meszaros B, Erdos G, Dosztanyi Z. 2018. IUPred2A: context-dependent prediction of protein

disorder as a function of redox state and protein binding. Nucleic Acids Res. 46:W329-

W337.

Moyers BA, Zhang JZ. 2016. Evaluating Phylostratigraphic Evidence for Widespread De Novo

Gene Birth in Genome Evolution. Mol. Biol. Evol. 33:1245-1256.

Moyers BA, Zhang JZ. 2015. Phylostratigraphic Bias Creates Spurious Patterns of Genome

Evolution. Mol. Biol. Evol. 32:258-267.

Neme R, Amador C, Yildirim B, McConnell E, Tautz D. 2017. Random sequences are an

abundant source of bioactive RNAs or peptides. Nat. Ecol. Evol. 1:0127.

Prijambada ID, Yomo T, Tanaka F, Kawama T, Yamamoto K, Hasegawa A, Shima Y, Negoro S,

Urabe I. 1996. Solubility of artificial proteins with random sequences. FEBS Lett.

382:21-25.

R Core Team. 2019. R: A language and environment for statistical computing: R Foundation for

Statistical Computing.

Rajon E, Masel J. 2011. Evolution of molecular error rates and the consequences for

evolvability. Proc. Natl. Acad. Sci. U.S.A. 108:1082-1087.

Rousseau F, Schymkowitz J, Serrano L. 2006. Protein aggregation and amyloidosis: confusion of

the kinds? Current opinion in structural biology 16:118-126.

Savic RM, Karlsson MO. 2009. Importance of shrinkage in empirical bayes estimates for

diagnostics: problems and solutions. AAPS J. 11:558-569.

Sormanni P, Aprile FA, Vendruscolo M. 2015. The CamSol Method of Rational Design of

Protein Mutants with Enhanced Solubility. J. Mol. Biol. 427:478-490.

Theillet F-X, Kalmar L, Tompa P, Han K-H, Selenko P, Dunker AK, Daughdrill GW, Uversky

VN. 2013. The alphabet of intrinsic disorder: I. Act like a Pro: On the abundance and

roles of proline residues in intrinsically disordered proteins. Intrinsically Disord. Proteins

1:e24360.

Tien MZ, Meyer AG, Sydykova DK, Spielman SJ, Wilke CO. 2013. Maximum allowed solvent

accessibilites of residues in proteins. PLoS One 8:e80635.

Tretyachenko V, Vymetal J, Bednarova L, Kopecky V, Hofbauerova K, Jindrova H, Hubalek M,

Soucek R, Konvalinka J, Vondrasek J, et al. 2017. Random protein sequences can form

defined secondary structures and are well-tolerated in vivo. Scientific Reports 7:15449.

Tsai J, Taylor R, Chothia C, Gerstein M. 1999. The packing density in proteins: Standard radii

and volumes. J. Mol. Biol. 290:253-266.

Vakirlis N, Acar O, Hsu B, Coelho NC, Van Oss SB, Wacholder A, Medetgul-Ernar K, Bowman

RW, Hines CP, Iannotta J, et al. 2020. De novo emergence of adaptive membrane

proteins from thymine-rich genomic sequences. Nature Communications 11:781.

Vakirlis N, Carvunis A-R, McLysaght A. 2020. Synteny-based analyses indicate that sequence

divergence is not the main source of orphan genes. eLife 9:e53500.

Vakirlis N, Hebert AS, Opulente DA, Achaz G, Hittinger CT, Fischer G, Coon JJ, Lafontaine I.

2018. A Molecular Portrait of De Novo Genes in Yeasts. Mol. Biol. Evol. 35:631–645.

Van Oss SB, Carvunis AR. 2019. *De novo* gene birth. PLoS Genet. 15:e1008160.

Vecchi G, Sormanni P, Mannini B, Vandelli A, Tartaglia GG, Dobson CM, Hartl FU,

Vendruscolo M. 2020. Proteome-wide observation of the phenomenon of life on the edge

of solubility. Proc. Natl. Acad. Sci. U.S.A. 117:1015-1020.

Weisman CM, Eddy SR. 2017. Gene Evolution: Getting Something from Nothing. Current

Biology 27:R661-R663.

Wickham H. 2016. ggplot2: Elegant graphics for data analysis. New York: Springer-Verlag.

Willis S, Masel J. 2018. Gene birth contributes to structural disorder encoded by overlapping

genes. Genetics 210:303-313.

Wilson BA, Foy SG, Neme R, Masel J. 2017. Young genes are highly disordered as predicted by

the preadaptation hypothesis of *de novo* gene birth. Nat. Ecol. Evol. 1:0146.

Wilson BA, Masel J. 2011. Putatively noncoding transcripts show extensive association with

ribosomes. Genome Biol. Evol. 3:1245-1252.