Infection with CagA⁺ *Helicobacter pylori* induces epithelial to mesenchymal transition in human cholangiocytes

Prissadee Thanaphongdecha[1,2,3], Shannon E. Karinshak[1], Wannaporn Ittiprasert[1], Victoria H. Mann[1], Yaovalux Chamgramol[3], Chawalit Pairojkul[3], Sutas Suttiprapa[2], Banchob Sripa [2,3,*], Paul J. Brindley[1,*]

[1] Department of Microbiology, Immunology and Tropical Medicine, and Research Center for Neglected Tropical Diseases of Poverty, School of Medicine & Health Sciences, The George Washington University, Washington DC, 20037, USA

[2] Tropical Disease Research Laboratory, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand

[3] Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand

[*] Equal contribution

Correspondence: Paul J. Brindley, Department of Microbiology, Immunology and Tropical Medicine, and Research Center for Neglected Tropical Diseases of Poverty, School of Medicine & Health Sciences, The George Washington University, Washington DC, 20037, USA, email, pbrindley@gwu.edu; or Banchob Sripa, Tropical Disease Research Laboratory, Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand, email, banchob@kku.ac.th

Abstract

Recent reports suggest that the East Asian liver fluke *Opisthorchis viverrini*, which is implicated in pathogenesis of opisthorchiasis-associated cholangiocarcinoma, serves as a reservoir of *Helicobacter pylori*. The affected cholangiocytes that line the intrahepatic biliary tract are considered to be the cell of origin of this malignancy. Here, we investigated interactions among a human cholangiocyte cell line, H69, a CCA cell line CC-LP-1, CagA$^+$ve *Helicobacter pylori*, and *H. bilis* bacilli. Exposure to increasing numbers of *H. pylori* at 0, 1, 10,100 bacteria per H69 cell for 24 hrs induced phenotypic changes in the cultured cholangiocytes including profusion of thread-like filopodia and loss of cell-cell contact in a dose-dependent fashion. In parallel, changes in mRNA expression followed exposure to *H. pylori*, with increased expression of epithelial to mesenchymal transition (EMT) associated-factors including snail, slug, vimentin, matrix metalloprotease, zinc finger E-box-binding homeobox, and the cancer stem cell marker CD44. Transcription levels encoding the cell adhesion marker CD24 decreased. Analysis in real time using the xCELLigence system to quantify cell proliferation, migration and invasion revealed that exposure to 10 to 50 of *H. pylori* stimulated migration of H69 cells and CC-LP-1 cells, a line derived form of a human cholangiocarcinoma, and invasion through an extracellular matrix. In addition, 10 bacilli of CagA$^+$ve *H. pylori*, but not *H. bilis*, stimulated contact-independent colony establishment in soft agar. These findings support the hypothesis that infection with *H. pylori* can contribute to the malignant transformation of the biliary epithelium.

Keyword : *Helicobacter pylori*, cholangiocyte, epithelial-to-mesenchymal transition.

Introduction

Complex associations occur between trematodes and bacterial commensals, including species capable of causing human diseases such as non-typhoidal salmonellosis, leptospirosis and Sennetsu fevers (1-3). There is increasing evidence that the East Asian liver fluke *Opisthorchis viverrini* may serve as a reservoir of *Helicobacter* and implicate *Helicobacter* in pathogenesis of opisthorchiasis-associated cholangiocarcinoma (CCA) (4-8). The International Agency for Research on Cancer of the World Health Organization classifies infection with the liver flukes *O. viverrini* and *Clonorchis sinensis* and with *H. pylori* as Group 1 carcinogens (9). In northern and northeastern Thailand and Laos, infection with *O. viverrini* is the major risk for CCA (9-12). Given the elevated prevalence of CCA in this region where infection with liver fluke prevails, and given evidence of linkages between human infection by *Helicobacter* during opisthorchiasis, these two biological carcinogens together may orchestrate the pathogenesis of opisthorchiasis and bile duct cancer. The association of *Helicobacter* and its virulence factors, including during opisthorchiasis, may underlie biliary tract disease including CCA in liver fluke-endemic regions (13). Infection with species of *Helicobacter* causes hepatobiliary tract diseases that can resemble opisthorchiasis (8, 14, 15) (16, 17). Lesions ascribed to liver fluke infection, including cholangitis, biliary hyperplasia and metaplasia, periductal fibrosis and CCA, may be due in part to *Helicobacter*-associated hepatobiliary disease.

Commensalism involving *H. pylori* and *O. viverrini* may have evolved and may facilitate conveyance of the bacillus into the human biliary tract during the migration of the juvenile fluke following ingestion of the metacercaria in raw or undercooked cyprinid fish and establishment of infection (4, 18). Since we have shown previously that curved rods resembling *Helicobacter* occur in the alimentary tract of the liver fluke (4), and given the low pH of the gut of *O. viverrini*, the *H. pylori* rods or spores might be transported by the migrating parasite. The spiral shape of *H. pylori* attach to biliary cells, which internalize in similar fashion to their behavior on gastric epithelium (19, 20).

Transition of epithelial cells to mesenchymal cells during disease and epithelial to mesenchymal transition (EMT) during development and wound healing follows conserved EMT routes with well-characterized molecular and morphological hallmarks (21). These hallmarks are clearly displayed at malignant transformation of the epithelium of the gastric mucosa resulting from infection with *H. pylori* (22). Here, we investigated interactions between CagA$^+$ve *H. pylori*, the related species *H. bilis* (23), and human cholangiocytes including the H69 cholangiocyte cell line (24, 25) and the CCLP-1cholangiocarcinoma cell line (26). Infection with CagA$^+$ *H. pylori* induced epithelial to mesenchymal transition in human cholangiocytes.

Materials and Methods

Human cholangiocyte lines

The immortalized intrahepatic cholangiocyte cell line, H69 (24, 25), Cellosaurus (27) identifier RRID:CVCL_812,1, and the primary cholangiocarcinoma cell line, CC-LP-1 (26, 28) Cellosaurus RRID CVCL_0205, were obtained as described (26, 28-30). In brief, H69 cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) (Thermo Fisher Scientific, Inc.),

DMEM/F12 (Sigma, St Louis) supplemented with 10% fetal bovine serum (FBS; Invitrogen;
Thermo Fisher Scientific, Inc.), supplemented with adenine, insulin, epinephrine,
triiodothyronine/transferrin, hydrocortisone and epidermal growth factor and
penicillin/streptomycin (PS). CC-LP-1 cells were cultured in DMEM containing 10%FBS, L-
glutamine and antibiotics as above. All cells maintained in humidified 5% CO_2 in air at 37°C.
The cells were cultured to ~80% confluence before co-culture with bacteria.
145
146   ***Helicobacter pylori* and *Helicobacter bili*s**
147
Helicobacter pylori, ATCC 43504, isolated originally from human gastric antrum (31) and *H.*
bilis, Fox *et al.* ATCC 49314, originally isolated from intestines and liver of mice (32, 33), were
obtained from the American Type Culture Collection (ATCC) (Manassas, VA) and a gift from
Dr. Fox MIT and maintained on trypticase soy agar with 5% sheep blood for 72 to 96 hours (34)
(Becton Dickinson, Cockeysville, MD), incubated at 37°C with rocking under microaerobic
conditions with the BBL Campy Pack Plus system (Becton Dickinson). After 96 hours, each
bacterial species was harvested independently by scraping the colonies from the agar plate, the
cell scraping was resuspended in DMEM/F12 medium to an optical density at 600 nm (OD_{600}) of
1. 0 corresponding to approximately 1×10^8 colony forming units (cfu)/ml (35-37).
157
158   **Morphological assessment of cholangiocytes**
159
160   *H. pylori* was co-cultured with H69 cells at different multiplicities of infection (MOIs) of 0, 10
161   and 100 for 24 hours in serum-free media. After 24 hours, the effect of *H. pylori* to H69 cell
162   morphology were observed under an inverted microscope (Zeiss Axio Observer A.1) (Figure 1).
163
164   ***In vitro* wound healing assay**
165
Monitoring cell migration in a two-dimensional confluent monolayer in controlled *in vitro*
conditions allowed investigators to simulate and explore critical mechanisms of the impact of *H.*
pylori on the process of wound healing. Sheet migration (38, 39) was used in this study to
observe H69 cell migration after *H. pylori* exposure using differing MOI of 0, 10 or 100.
Briefly, H69 cells were grown in 6-well plates (Greiner, Monroe NC) overnight to allow cell
adherence. A linear thin scratch (wound) was created to produce a gap in the confluent cell
monolayer using a sterile 20 µl pipette tip. The gap was observed and measured for up to 26
hours. The gap width was analyzed using ImageJ software (NIH, USA). The wound closure rate
over time was calculated and compared with control group (H69 cells grown in the absence of *H.*
pylori). The statistical significance was analyzed by One-way ANOVA (Prism).
176
177   **Real time quantitative PCR (RT-qPCR) analysis**
178
Total RNA from H69 cells were extracted with RNAzol[RT] (Molecular Research Center, Inc.)
according to the manufacturer's instructions. The RNA quality and concentration were assessed
using the NanoDrop 1000 spectrophotometer (Thermo Fisher Scientific, MA). Total RNA (500
ng) was reversed-transcribed using iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA).
Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) analysis of mRNA
expression of six members of the EMT-associated genes including Vimentin, Snai1, Slug, Zeb,

Jam1 and MMP7; two cancer stem cells including CD44 and CD24 were undertaken on an
ABI7300 system (ABI) using SsoAdvance SYBR green mixture (Bio-Rad), normalized to
human GAPDH (Supplementary Table 1 provides the primer sequences). The relative fold-
change was determined by the ΔΔCt method (40). Three biological replicates of these assays
were carried out.
190
191 **Cell migration after *H. pylori* co-culture**
192
To monitor cell migration/invasion in real time, we used the xCELLigence Real-time cell
analyzer (RTCA) DP instrument (ACEA Biosciences, San Diego CA) equipped with a CIM-
plate 16 (ACEA Biosciences, San Diego CA), which is a 16-well system in which each well is
composed of upper and lower chambers separated by an 8-um microporous membrane.
Migration/invasion was measured as the relative impedance change (cell index) across
microelectronic sensors integrated into the bottom of the membrane.   H69 and CC-LP-1 cells
were infected with *H. pylori* ATCC45304 strain at MOIs of 0,1,10,50 and 100 respectively for 2
hours prior to the experiment, then harvested later using 0.25% trypsin-EDTA and washed with
medium. A total of 30,000 of H69 or CC-LP-1 cells were seeded into each well of the upper
chamber (UC) in serum-free media. Complete media containing 10 % serum was added the
lower chamber (LC) of each well. The CIM-plates was left in an incubator for 1 hour to allow
cell attachment, then monitored for migration/invasion over 96 hours taking a reading every 15
minutes. For quantification, the cell index (CI) was averaged from three independent
measurements at each time point (41).
207
208 **Invasion monitoring in cells exposed to *H. pylori***
209
The rate of cell invasion was monitored in real-time using the xCELLigence system CIM-plates.
Two hours prior to conducting the experiment H69 or CC-LP1 cells were infected with *H. pylori*
ATCC45304 strain at either MOI of 0 or 10 . The UC of the CIM plate was coated with 20%
solution of Matrigel[TM] (BD Biosciences). A total of 30,000 of H69 or CC-LP-1 cells were
seeded in each well of the UC in serum-free media.. H69 or CC-LP media was added to each
well of the LC. The CIM-plates placed at 37^{o} C in 5% CO_2 for 60 minutes to allow cell
attachment. The impedance value of each well was monitored by the xCELLigence system at
intervals of 20 minutes for 96 hours, at which point the experiment was terminated. Invasion was
expressed as a CI value.
219
220 **Assessment of cell proliferation**
221
Cell proliferation of H69 was assessed using the xCELLigence DP system (ACEA Biosciences,
San Diego, CA, USA).   H69 cells were fasted for 4 to 6 hours in 1:20 diluted medium that had
been diluted 1:20 with 50% DMEM and 50% DMEM/Ham's F12, containing 1% of PS, *i.e.*
0.5% FBS final concentration, as described (42, 43), then the cells were co-cultured of viable
bacilli of *H. bilis* for 120 min before starting the assay. H69 cells were harvested later using
0.25% trypsin-EDTA and washed with medium. Five thousand H69 cells were seeded on to each
well of the E-plate in H69 medium and cultured for 24 hours. The culture medium was removed,
and the cells were gently washed with 1×PBS. The PBS was replaced with serum diluted
medium. The cellular growth was monitored in real time with readings collected at intervals of

20 minutes for ≥60 hours. For quantification, CI was averaged from three independent
measurements at each time point.
233
234 **Colony forming assay in soft agar**
235
236 H69 cells (10,000 cells per well) were mixed with 0.3% low melting temperature agarose
237 (NuSieveGTG; Lonza, Walkersville, MD) in growth medium, plated on top of a solidified layer
238 of 0.6% agarose in growth medium, in 6-well or 20 mm cell culture plates, and fed every three
239 days (twice a week) with growth medium. After 28 days or until the colonies had established,
240 the colonies with diameter 50 μm were counted using phase-contrast microscopy (44).
241
242 **Statistical analysis**
243
Differences in cell elongation, cell scattering, wound healing and colony formation among test
and control groups were analysed by one-way ANOVA. Fold differences in mRNA levels, real
time cell migration and invasion through extracellular matrix were analysed by two-way
ANOVA followed by Dunnett's test for multiple comparisons. Analysis was accomplished using
GraphPad Prism version 7 software (GraphPad, San Diego, CA). A P value of ≤ 0.05 was
considered to be statistically significant.
250
251 **Results**
252
***H. pylori* induces epithelial to mesenchymal transition (EMT) phenotype in biliary cells**
H69 were directly exposed to increasing number of *H. pylori* at 0, 10 and 100 bacilli per H69 cell
in 6 wells. After 24 hours of infection, the epithelial morphology of H69 was changed into
mesenchymal phenotype characterized by cell elongation, protrusion of thread-like filopodia and
loss cell-cell contact in dose dependent fashion, which implicates increasing change in cell
motility. The morphological change of H69 was also supported by increasing L/W ratio of *H.*
pylori-infected cells, and H69 cells were more isolated after exposure to *H. pylori*, in dose-
dependent fashion (Figure 2).This phenotype change can resemble the phenomenon found in
AGS cell line after exposure to *H. pylori* called "Hummingbird phenotype"(Figure 1). On the
other side, H69 treated with 5 ng of TGF-beta were surprisingly increased cell aggregation and
nodule formation (Supplementary Figure 6).
264

265
266
Figure 1. CagA[+] *H. pylori* ATCC 43504 induced phenotypic changes in cultured
cholangiocytes. The photomicrographs showed cell morphology by light microscopy of H69
cells exposed to increasing numbers of *H. pylori* at 0, 10 and 100 bacteria per cholangiocyte,
respectively (left to right) (A). The appearance of cells had changed from epithelial to

mesenchymal phenotype including loss of cell-cell contact, elongated and spindle-shaped
morphology along with growth as individual cells after 24 hours of *H. pylori* infection in dose-
dependent manner. Scale bars, 50 μm (right), 20× magnification.
274
Figure 2. Quantification of epidermal to mesenchymal transition induced by cagA[+] *H.*
pylori. Cell elongation was measured by length-to-width ratio and isolate single cells were
counted to determine cell scattering (A, B). The number of elongated cells increased in a dose-
dependent fashion in response to infection by *H. pylori* (C). By contrast, the number of isolate
individual H69 cells indicated cell scattering was also significantly increased in dose-dependent
fashion (D). Data are presented as the mean values with standard errors from three replications.
Asterisks indicates statistically significant different of *H. pylori*-infected group compared to
normal group at 24 hours (*, $P < 0.05$; **, $P < 0.005$; ***, $P < 0.001$; $P < 0.0001$).

283
284
285

Expression of epithelial to mesenchymal transition (EMT) associated-factors induced by exposure to *H. pylori*

The dynamic change of EMT at transcription level was determined by quantitative real-time PCR of six EMT-related factors including mesenchymal marker (vimentin), transcription factors (SnaiI, Slug, Zeb1), adhesion molecules (Jam1) and proteolytic enzyme (MMP7). Two cancer stem cell markers, CD24 and CD44, also were included in this analysis. Levels of fold-difference for each of the six markers increased in a dose-dependent fashion. The regulatory transcriptional factor, Snail was remarkably up-regulated with highest level of fold difference: 88, 656 and 5,309 folds in MOI 10, 50 and 100 respectively followed by MMP7 expression: 15, 21, 44 folds in MOI 10, 50 and 100 respectively. Slug, Zeb1, Vimentin and Jam1 expression were up-regulated at several levels of bacterial infection. In addition, the cancer stem cell markers CD24 was down-regulated, whereas CD44 was up-regulated, displaying a CD44+ high/CD24+ low pattern following exposure to *H. pylori* (Figure 3)

Figure 3. Transcriptional change of EMT-related genes and cancer stem cell markers after *H. pylori* exposure. mRNA expression of 6 EMT-related genes and 2 cancer stem cell markers were determined with three biological replicates after 24 hours of infection. Expression of Snail, Slug, vimentin, Jam, MMP7 and CD44 were dose-dependently increased, whereas CD24 was dose-dependently decreased. The regulatory transcriptional factor, Snail was remarkably up-regulated with high level of fold difference; 88, 656 and 5,309 folds with MOI 10, 50 and 100 respectively. MMP7 expression was also remarkably up-regulated with every multiplicity of

infection; Slug, Zeb1, vimentin and Jam1 expression were up-regulated in specific amount of bacterial infection. The cancer stem cell markers, CD44 was significantly up-regulated with MOI 50 and 100, whereas CD24 was down-regulated. These results indicated that *H. pylori* parallelly induced EMT and stemness of Human cholangiocyte, H69. qPCR data were normalized to GAPDH expression for each sample.

Quantitative monitoring of cell migration and Matrigel invasion in real-time by xCELLigence approach

Scratch assays revealed two-dimensional migration of H69 cells. Wound healing of H69 was significantly increased to 19.47% in MOI of 10. Nevertheless, the gap closing was slowed down to 11.09% by MOI of 100 compared with healthy H69 at 14.47% (Supplementary Figure 7). Analysis in real time using the xCELLigence approach revealed that exposure to 10 to 50 *H. pylori* but not 100 *H. pylori* stimulated migration of H69 and the CC-LP1, cell line derived from a human cholangiocarcinoma, and invasion of Matrigel extracellular matrix by cells of the CC-LP-1 cholangiocarcinoma cell line (Figure 4).

Figure 4. Investigation of EMT-associated cell behavior by Real-time cell migration and Matrigel invasion analysis using xCELLigence system. The immortalized human cholangiocyte, H69 infected with *H. pylori* showed different level of cell migratory responses

depending on number of infections. Group of cells infected with MOI 10 revealed the highest
cell migration rate followed by MOI 50, whereas MOI 100 attenuated cell migration, compared
with non-infected control (A). Similarly, the cholangiocarcinoma cell line, CC-LP-1 showed the
highest cell migration rate when stimulated with MOI 10 followed by MOI 50. Moreover, CC-
LP-1 cells significantly migrated faster even with 1 multiplicity of infection(B). Matrigel
invasion compared between MOI 10 and non-infected cells, the Invasion rate of both H69 and
CC-LP1 cells were significantly increased when infected with (C,D). Cell migration and invasion
were monitored for 100 hours with chemoattractant in the lower chamber of CIM plate.
337
338 *H. pylori* **induces contact-independent colony formation in soft agar**
339
The tumorigenesis of infected H69 was determined by anchorage independent cell growth in soft
agar. The number of colony was counted, and the results revealed that 10 multiplicity of
infection can significantly increase the number of colony in soft agar, but not 50 or 100 MOI.
The size of colonies were also increased in all infected group, but significant only in MOI of 50
(Supplementary Figure 3). The other species of *Helicobacter*, *H. bilis* which naturally lives in
the bile duct surprisingly decreased the number of colony formation in soft agar (Figure 5). The
inhibition effect of *H. bilis* to biliary cell proliferation was supported by real-time cell
proliferation approach (Supplementary Figure 4).

348
349
Figure 5. Anchorage independent cell growth in soft agar revealed cell transformation of
H69. Single cell of and- Infected H69 had gradually grown into colony in soft agar. Colony
formation was shown and counted at day 30 (A). In-infected group, number of colonies were
significantly increased with a MOI of 10; no different with MOI of 50; decreased with MOI of
100, compared with non-infected control (B). On the other side, number of colonies in infected
group had dramatically decreased in dose-dependent fashion, which indicated inhibitory effect of
this bacteria toward anchorage independent cell growth or cell transformation of H69 (C). Three
biological replicates of this assay were undertaken.
358

Discussion

Numerous species of *Helicobacter* have been described (45) while *H. pylori* was the first bacterial pathogen confirmed to cause gastric disease including peptic ulcer, gastric lymphoma and gastric adenocarcinoma (31, 46-49). Bacilli of *H. pylori* occur in the stomach in at least half the human population. Transmission from mother to child and by other routes takes place. The human-*H. pylori* association may be beneficial in early life, including contributions to a healthy microbiome and reduced early-onset asthma (50, 51). Many strains of *H. pylori* have been characterized at the molecular level, and these are generally classified as either CagA positive or CagA negative. The major virulence factor, cytotoxin-associated gene A (CagA) is a ~140 kDa protein encoded on the cag pathogenicity island (PAI). Cag PAI also encodes a type 4 secretion channel through which the CagA virulence factor is introduced into the host cell upon attachment by the bacillus (22). CagA locates to the cell membrane, from where it is phosphylorated by the Src kinases. Downstream of these modifications, the activated CagA orchestrates changes in morphology of the infected epithelial cell, which loses cell polarity and becomes motile, exhibiting a diagnostic 'hummingbird-like' phenotype (52-54). In addition to the well-known association with gastric cancer, *H. pylori* has also been associated with hepatological disease including in the biliary tract (7, 8, 55). The related species, *H. hepaticus* and *H. bilis*, also have been associated with hepatobiliary disease (55-57).

Infection with the fish-borne liver flukes *Opisthorchis viverrini, Clonorchis sinensis* as well as the bacterial infection with *Helicobacter pylori* are all classified as Group 1 carcinogens by the International Agency for Research on Cancer of the World Health Organization (9). Opisthorchiasis is a major risk factor for cholangiocarcinoma in the northern and northeastern part of Thailand and Laos (9-12). In addition to gastric disease, infection with species of *Helicobacter* causes hepatobiliary tract diseases that can resemble opisthorchiasis (8, 14, 15) (16, 17). Liver fluke infection can induce lesions in biliary system including cholangitis, biliary hyperplasia and metaplasia, periductal fibrosis and CCA. These lesions derive not only from liver fluke infection but perhaps are in part the consequence of hepatobiliary tract infection with *H. pylori*. Given the high incidence rate of cholangiocarcinoma in this endemic area where the infection with *O. viverrini* prevails, and given evidence of this liver fluke as a reservoir host of *H. pylori*, these two biological carcinogens may together drive the pathogenesis of opisthorchiasis and cholangiocarcinoma in this region. In addition, the virulence factors of *H. pylori* may be a key risk factor that underlie biliary tract disease including CCA during opisthorchiasis (13). Furthermore, the CagA oncoprotein secreted by *H. pylori* into epithelial cells is noted for its variation at the SHP2 binding site and, based on the sequence variation, is subclassified into two main types - East-Asian CagA and Western CagA. East-Asian CagA shows stronger SHP2 binding and greater biological activity than Western CagA. It has been proposed that in East Asia, the circulation of *H. pylori* strains encoding active forms of CagA underlie the elevated incidence of gastric adenocarcinoma (22).

Here, we investigated interactions among a human cholangiocyte cell line, H69, a CCA cell line CC-LP-1, CagA$^+$ve *Helicobacter pylori*, and *H. bilis* bacilli.. Infection of H69 with CagA$^+$ *H. pylori* induced epithelial to mesenchymal transition phenotype in dose-dependent manner, which was characterized by cell elongation and scattering which implicate increasing change in cell motility. This phenotype of H69-infected cells was reminiscent of the modified appearance of the

405 AGS gastric carcinoma cell after exposure *H. pylori*, which has been termed the hummingbird
406 phenotype (58). In AGS cells, delivery of CagA by the type IV secretion mechanism from *H.*
407 *pylori* to the epithelial cell subverts the normal signaling leading to actin-dependent
408 morphological rearrangement and the hummingbird appearance. The hitherto uniform polygonal
409 shape becomes markedly elongated with terminal needle-like projections (59).
410
Earlier studies in AGS cells infected with *H. pylori* had also demonstrated that the hummingbird
phenotype was accompanied early on with transcription changes that reflected the epithelial to
mesenchymal transition, EMT. Similar transcriptional changes were seen in the *H. plyori*-
infected H69 cells. Specifically, upregulation of expression of Snai1, Slug, vimentin, Jam1, and
MMP7 expression in dose-dependent fashion strongly supported, at the expression level, a
dynamic change in cholangiocyte reflective of EMT. Moreover, the E-cadherin repressor, Snail
was markedly up-regulated, which indicated that this factor may be a key driver of epithelial to
mesenchymal transition in H69. CD44 expression was up-regulated in dose-dependent fashion,
whereas CD24 decreased, showing a CD44 high/CD24 low phenotype during infection. These
results indicated that *H. pylori* may not only induce EMT but also contributes to stemness and
malignant transformation of the cholangiocyte.
422
Realtime cell monitoring approach by xCELLigence revealed that the H69 and CC-LP-1 cells
showed higher cell migration when exposed to *H. pylori*. The fastest migratory rate was MOI of
10 followed by MOI of 50. By contrast, the rate decreased with 100 bacteria per cell, suggested
direct detrimental effect of the elevated levels of bacterial exposure. Ten bacilli per cell stimulate
both migration and invasion of the basement membrane model matrix by both the H69 and CC-
LP-1 cells, which was evidence of behavioral changes in these epithelial cells characteristic of
epidermal to mesenchymal transition. In addition, following exposure to *H. pylori*, H69 cells
showed anchorage independent cell growth in soft agar, an accepted in vitro phenotype of the
escape from anoikosis that is characteristic of metastasis (60). At a MOI of 10, there was a
significant increase in numbers of colonies of the H69 cell in soft agar. By contrast, not only
were increase in colonies also seen after exposure of H69 cells to *H. bilis* in the model of contact
independent cell growth, and also cell proliferation as quantified in E-plate on the xCELLIgene
platform, both numbers of colonies and cell growth decreased. In general, these outcome
indicated that *H. pylori* has the potential to induce carcinogenic changes in cholangiocytes.
437
H. pylori-specific DNA sequences has been detected in CCA tumors and also from lesions
diagnosed as cholecystitis/cholelithiasis in regions endemic for opisthorchiasis (8, 14).
Moreover, serological findings indicate infection with *H. pylori* in Thais at high risk for CCA
(15). Given that here we have observed that CagA+ *H. pylori* stimulated epithelial to
mesenchymal transition of the human cholangiocyte which in turn indicates the underlying
mechanism of tissue fibrosis(61-63) and cancer metastasis(64-66) , an explanation for why
infection with the liver fluke induces bile duct cancer (67) might now be clearer – involvement
by *H. pylori* and its virulence factors. *Helicobacter* likely passes from the stomach to the
duodenum and enters the biliary tree through the duodenal papilla and ampulla of Vater (19, 68).
447
In contrast to the malignancy-conducive focus of this relationship between the liver fluke *O.*
viverrini and *H. pylori*, co-infections with *H. pylori* and other *Helicobacter* species and other
helminths have generally described diminution in risk for *H. pylori*-associated gastric carcinoma

(69-71) For example, the concurrent infection on mice with an intestinal nematode modulates
inflammation, induces a Th2-polarizing cytokine phenotype with concurrent downmodulation of
Th1 responses and the gastric immune responses, and reduces *Helicobacter*-induced gastric
atrophy (72). To conclude, infection with CagA$^+$ *H. pylori* induced epithelial to mesenchymal
transition in human cholangiocytes, along with other changes strongly reflective of malignant
transformation. Further investigation of the relationship of *O. viverrini* and *H. pylori* within the
infected biliary tract clearly is warranted.
458
459
460 **Abbreviations** Cytotoxin-associated gene A (*cag*A), cholangiocarcinoma (CCA); CagA
461 multimerization motif (CM); EMT, epithelial to mesenchymal transition; ZEB, zinc finger E-
462 box-binding homeobox 1; VIM, vimentin; SNAII, MMP7 and cancer stem cell marker CD44;
463 CD24.
464
465 **Competing Financial Interests**
466
467 The authors declare there were no competing financial interests.
468
469 **Acknowledgments**
470

We thank Dr. James G. Fox for advice and cultures of *H. bilis* and related bacilli. P.T.
acknowledges support as a PhD research scholar from the Thailand Research Fund under the
Royal Golden Jubilee scholars program (grant number PHD/0013/2555); B.S. acknowledges
support from Thailand Research Fund Senior Research Scholar. This work was supported by the
National Health Security Office, Thailand, the Higher Education Research Promotion and
National Research University Project of Thailand, Office of the Higher Education Commission,
through the Health Cluster (SHeP-GMS), the Faculty of Medicine, Khon Kaen University,
Thailand (award number I56110), and the Thailand Research Fund under the TRF Senior
Research Scholar (RTA 5680006); the National Research Council of Thailand. The National
Institute of Allergy and Infectious Diseases (NIAID), Tropical Medicine Research Center award
number P50AI098639, The National Cancer Institute, award number R01CA164719, and the
United States Army Medical Research and Materiel Command (USAMRMC), contract number
W81XWH-12-C-0267 also provided support. The content is solely the responsibility of the
authors and does not necessarily represent the official views of the funders including
USAMRMC, NIAID, NCI or the NIH.

486
487 **Author contributions**
488
489 P.T., S.S. C.P., Y.C., B.S. and P.J.B. conceived and designed the study. P.T., S.K., V.M., W.I.
490 performed the experiments. J.F. provides H. bilis and experimental designs. P.T., W.I., B.S., J.F.
491 and PJB analyzed and interpreted the findings. P.T., B.S., W.I., S.S., and PJB wrote the
492 manuscript. All authors read and approved the final version of the paper.
493
494

Supporting information

Supplementary Figure 1. TGF-Beta promoted nodule formation. TGF-beta (EMT
stimulating agent) induced nodule formation in the human cholangiocyte cell line, H69. Cells
were cultured in the presence of 5 ng/ml TGF-beta in serum and hormone-free media for 24
hours.

504
505
506

507
508
509

**Supplementary Figure 2. Wound healing revealed cell migration of H69 cholangiocytes
infected with *Helicobacter pylori* in two dimensions.** To investigate the effect of on cell
migration, the scratch assay was performed in triplicate with different number of bacterial
infections. Wound closure was significantly increased to 19.47% with a MOI 10 of *H. pylori*.
Nevertheless, MOI 100 showed lower cell migration at 11.09%, compared with control at
14.47%. The photomicrograph at 0 and 26 hours, 5×magnification.

516
517
518 **Supplementary Figure 3. Size of colony in soft agar.** The colonies with diameter over 50 μm
519 were measure and calculated. The average size of colonies were 77.63, 91.62, 95.25 and 81.46
520 μm in MOI 0f 0, 10, 50 and 100, respectively.
521
522

523
524

Supplementary Figure 4. Real time cell proliferation of-infected H69 determined by
xCELLigence approach. The immortalized human cholangiocyte, H69 cells were infected with
increasing number of (ATCC 43879) at MOI 10, 50 and 100, respectively. Cell growth indicated
by cell index of all infected groups were dramatically decreased and lower than control after 40
hours. MOI 100 had the lowest cell proliferation followed by MOI 50 and MOI 10 respectively.
The results indicated cell growth inhibitory effect of toward H69. Cells proliferation assay were
performed in E-plate in minimal media.
532
533

Gene name	Gen Bank Number	Primer sequence (5'-3')	Product size (bp)
CD24	NM_013230.3	Forward CACCCAGCATCCTGCTAGAC	259
		Reverse GAGACCACGAAGAGACTGGC	
CD44	NM_00610.3	Forward GGGAGTCAAGAAGGTGGAGC	237
		Reverse CTGAGACTTGCTGGCCTCTC	
F11r/Jam1	NM_016946.5	Forward GTGCCTACTCGGGCTTTTCT	201
		Reverse GAGCTTGACCTTGACCTCCC	
Gapdh	NM_002046.5	Forward CCCATGTTCGTCATGGGTGT	366
		Reverse TTCTAGACGGCAGGTCAGGT	
MMP7	NM_002423.5	Forward GGAGCTCATGGGGACTCCTA	172
		Reverse GGCCAAGTTCATGAGTTGCAG	
Snai1	NM_005985.3	Forward AGCTCTCTGAGGCCAAGGAT	325
		Reverse GACATTCGGGAGAAGGTCCG	
Vimentin	NM_003380.5	Forward GCAGGAGGCAGAAGAATGGT	216
		Reverse GCAGCTTCAACGGCAAAGTT	
Zeb1	NM_001128128.2	Forward CGCAGTCTGGGTGTAATCGT	477
		Reverse CCATGCCCTGAGGAGAACTG	

534
535
536 **Supplementary Table 1. Nucleotide sequences of primers specific for EM- associated**
537 **human genes, which were used in |quantitative real-time PCR analysis.**

References

1.   McNulty SN, Tort JF, Rinaldi G, Fischer K, Rosa BA, Smircich P, et al. Genomes of Fasciola hepatica from the Americas Reveal Colonization with Neorickettsia Endobacteria Related to the Agents of Potomac Horse and Human Sennetsu Fevers. PLoS Genet. 2017;13(1):e1006537.

2.   Van CD, Doungchawee G, Suttiprapa S, Arimatsu Y, Kaewkes S, Sripa B. Association between Opisthorchis viverrini and Leptospira spp. infection in endemic Northeast Thailand. Parasitol Int. 2017;66(4):503-9.

3.   Gendrel D, Kombila M, Beaudoin-Leblevec G, Richard-Lenoble D. Nontyphoidal salmonellal septicemia in Gabonese children infected with Schistosoma intercalatum. Clin Infect Dis. 1994;18(1):103-5.

4.   Deenonpoe R, Chomvarin C, Pairojkul C, Chamgramol Y, Loukas A, Brindley PJ, et al. The carcinogenic liver fluke Opisthorchis viverrini is a reservoir for species of Helicobacter. Asian Pac J Cancer Prev. 2015;16(5):1751-8.

5.   Deenonpoe R, Mairiang E, Mairiang P, Pairojkul C, Chamgramol Y, Rinaldi G, et al. Elevated prevalence of Helicobacter species and virulence factors in opisthorchiasis and associated hepatobiliary disease. Sci Rep. 2017;7:42744.

6.   Sripa B, Deenonpoe R, Brindley PJ. Co-infections with liver fluke and Helicobacter species: A paradigm change in pathogenesis of opisthorchiasis and cholangiocarcinoma? Parasitol Int. 2017;66(4):383-9.

7.   Boonyanugomol W, Chomvarin C, Song JY, Kim KM, Kim JM, Cho MJ, et al. Effects of Helicobacter pylori gamma-glutamyltranspeptidase on apoptosis and inflammation in human biliary cells. Dig Dis Sci. 2012;57(10):2615-24.

8.   Boonyanugomol W, Chomvarin C, Sripa B, Bhudhisawasdi V, Khuntikeo N, Hahnvajanawong C, et al. Helicobacter pylori in Thai patients with cholangiocarcinoma and its association with biliary inflammation and proliferation. HPB (Oxford). 2012;14(3):177-84.

9.   Humans IWGotEoCRt. Biological agents. Volume 100 B. A review of human carcinogens. IARC Monogr Eval Carcinog Risks Hum. 2012;100(Pt B):1-441.

10.   Sungkasubun P, Siripongsakun S, Akkarachinorate K, Vidhyarkorn S, Worakitsitisatorn A, Sricharunrat T, et al. Ultrasound screening for cholangiocarcinoma could detect premalignant lesions and early-stage diseases with survival benefits: a population-based prospective study of 4,225 subjects in an endemic area. BMC Cancer. 2016;16:346.

11.   Khuntikeo N, Loilome W, Thinkhamrop B, Chamadol N, Yongvanit P. A Comprehensive Public Health Conceptual Framework and Strategy to Effectively Combat Cholangiocarcinoma in Thailand. PLoS Negl Trop Dis. 2016;10(1):e0004293.

12.   Aye Soukhathammavong P, Rajpho V, Phongluxa K, Vonghachack Y, Hattendorf J, Hongvanthong B, et al. Subtle to severe hepatobiliary morbidity in Opisthorchis viverrini endemic settings in southern Laos. Acta Trop. 2015;141(Pt B):303-9.

13.   Segura-Lopez FK, Guitron-Cantu A, Torres J. Association between Helicobacter spp. infections and hepatobiliary malignancies: a review. World J Gastroenterol. 2015;21(5):1414-23.

14.   Boonyanugomol W, Chomvarin C, Sripa B, Chau-In S, Pugkhem A, Namwat W, et al. Molecular analysis of Helicobacter pylori virulent-associated genes in hepatobiliary patients. HPB (Oxford). 2012;14(11):754-63.

15.   Pisani P, Whary MT, Nilsson I, Sriamporn S, Wadstrom T, Fox JG, et al. Cross-reactivity between immune responses to Helicobacter bilis and Helicobacter pylori in a population in

Thailand at high risk of developing cholangiocarcinoma. Clin Vaccine Immunol.
2008;15(9):1363-8.
16.   Lvova MN, Tangkawattana S, Balthaisong S, Katokhin AV, Mordvinov VA, Sripa B.
Comparative histopathology of Opisthorchis felineus and Opisthorchis viverrini in a hamster
model: an implication of high pathogenicity of the European liver fluke. Parasitol Int.
2012;61(1):167-72.
17.   Mairiang E, Laha T, Bethony JM, Thinkhamrop B, Kaewkes S, Sithithaworn P, et al.
Ultrasonography assessment of hepatobiliary abnormalities in 3359 subjects with Opisthorchis
viverrini infection in endemic areas of Thailand. Parasitol Int. 2012;61(1):208-11.
18.   Plieskatt JL, Deenonpoe R, Mulvenna JP, Krause L, Sripa B, Bethony JM, et al. Infection
with the carcinogenic liver fluke Opisthorchis viverrini modifies intestinal and biliary
microbiome. FASEB J. 2013;27(11):4572-84.
19.   Boonyanugomol W, Chomvarin C, Hahnvajanawong C, Sripa B, Kaparakis-Liaskos M,
Ferrero RL. Helicobacter pylori cag pathogenicity island (cagPAI) involved in bacterial
internalization and IL-8 induced responses via NOD1- and MyD88-dependent mechanisms in
human biliary epithelial cells. PLoS One. 2013;8(10):e77358.
20.   Hatakeyama M. Helicobacter pylori CagA and gastric cancer: a paradigm for hit-and-run
carcinogenesis. Cell Host Microbe. 2014;15(3):306-16.
21.   Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal
transition. Nat Rev Mol Cell Biol. 2014;15(3):178-96.
22.   Hatakeyama M. Oncogenic mechanisms of the Helicobacter pylori CagA protein. Nat
Rev Cancer. 2004;4(9):688-94.
23.   Fox JG. Helicobacter bilis: bacterial provocateur orchestrates host immune responses to
commensal flora in a model of inflammatory bowel disease. Gut. 2007;56(7):898-900.
24.   Grubman SA, Perrone RD, Lee DW, Murray SL, Rogers LC, Wolkoff LI, et al.
Regulation of intracellular pH by immortalized human intrahepatic biliary epithelial cell lines.
Am J Physiol. 1994;266(6 Pt 1):G1060-70.
25.   Park J, Gores GJ, Patel T. Lipopolysaccharide induces cholangiocyte proliferation via an
interleukin-6-mediated activation of p44/p42 mitogen-activated protein kinase. Hepatology.
1999;29(4):1037-43.
26.   Shimizu Y, Demetris AJ, Gollin SM, Storto PD, Bedford HM, Altarac S, et al. Two new
human cholangiocarcinoma cell lines and their cytogenetics and responses to growth factors,
hormones, cytokines or immunologic effector cells. Int J Cancer. 1992;52(2):252-60.
27.   Bairoch A. The Cellosaurus, a Cell-Line Knowledge Resource. J Biomol Tech.
2018;29(2):25-38.
28.   Han C, Demetris AJ, Michalopoulos GK, Zhan Q, Shelhamer JH, Wu T. PPARgamma
ligands inhibit cholangiocarcinoma cell growth through p53-dependent GADD45 and p21
pathway. Hepatology. 2003;38(1):167-77.
29.   Smout MJ, Sotillo J, Laha T, Papatpremsiri A, Rinaldi G, Pimenta RN, et al.
Carcinogenic Parasite Secretes Growth Factor That Accelerates Wound Healing and Potentially
Promotes Neoplasia. PLoS Pathog. 2015;11(10):e1005209.
30.   Arunsan P, Ittiprasert W, Smout MJ, Cochran CJ, Mann VH, Chaiyadet S, et al.
Programmed knockout mutation of liver fluke granulin attenuates virulence of infection-induced
hepatobiliary morbidity. Elife. 2019;8.
31.   Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with
gastritis and peptic ulceration. Lancet. 1984;1(8390):1311-5.

32.   Dewhirst FE, Fox JG, Mendes EN, Paster BJ, Gates CE, Kirkbride CA, et al. 'Flexispira
rappini' strains represent at least 10 Helicobacter taxa. Int J Syst Evol Microbiol. 2000;50 Pt
5:1781-7.
33.   Fox JG, Yan LL, Dewhirst FE, Paster BJ, Shames B, Murphy JC, et al. Helicobacter bilis
sp. nov., a novel Helicobacter species isolated from bile, livers, and intestines of aged, inbred
mice. J Clin Microbiol. 1995;33(2):445-54.
34.   Hachem CY, Clarridge JE, Evans DG, Graham DY. Comparison of agar based media for
primary isolation of Helicobacter pylori. J Clin Pathol. 1995;48(8):714-6.
35.   Jung SW, Thamphiwatana S, Zhang L, Obonyo M. Mechanism of antibacterial activity of
liposomal linolenic acid against Helicobacter pylori. PLoS One. 2015;10(3):e0116519.
36.   Lin SN, Ayada K, Zhao Y, Yokota K, Takenaka R, Okada H, et al. Helicobacter pylori
heat-shock protein 60 induces production of the pro-inflammatory cytokine IL8 in monocytic
cells. J Med Microbiol. 2005;54(Pt 3):225-33.
37.   Pan H, Zhang Y, He GX, Katagori N, Chen H. A comparison of conventional methods
for the quantification of bacterial cells after exposure to metal oxide nanoparticles. BMC
Microbiol. 2014;14:222.
38.   Bindschadler M, McGrath JL. Sheet migration by wounded monolayers as an emergent
property of single-cell dynamics. J Cell Sci. 2007;120(Pt 5):876-84.
39.   Liang CC, Park AY, Guan JL. In vitro scratch assay: a convenient and inexpensive
method for analysis of cell migration in vitro. Nat Protoc. 2007;2(2):329-33.
40.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8.
41.   Xie H, Lee L, Scicluna P, Kavak E, Larsson C, Sandberg R, et al. Novel functions and
targets of miR-944 in human cervical cancer cells. Int J Cancer. 2015;136(5):E230-41.
42.   Matchimakul P, Rinaldi G, Suttiprapa S, Mann VH, Popratiloff A, Laha T, et al.
Apoptosis of cholangiocytes modulated by thioredoxin of carcinogenic liver fluke. Int J Biochem
Cell Biol. 2015;65:72-80.
43.   Rinaldi G, Yan H, Nacif-Pimenta R, Matchimakul P, Bridger J, Mann VH, et al.
Cytometric analysis, genetic manipulation and antibiotic selection of the snail embryonic cell
line Bge from Biomphalaria glabrata, the intermediate host of Schistosoma mansoni. Int J
Parasitol. 2015;45(8):527-35.
44.   Horibata S, Vo TV, Subramanian V, Thompson PR, Coonrod SA. Utilization of the Soft
Agar Colony Formation Assay to Identify Inhibitors of Tumorigenicity in Breast Cancer Cells. J
Vis Exp. 2015(99):e52727.
45.   Flahou B, Rimbara E, Mori S, Haesebrouck F, Shibayama K. The Other Helicobacters.
Helicobacter. 2015;20 Suppl 1:62-7.
46.   Cover TL. Helicobacter pylori Diversity and Gastric Cancer Risk. MBio.
2016;7(1):e01869-15.
47.   Marshall BJ. The pathogenesis of non-ulcer dyspepsia. Med J Aust. 1985;143(7):319.
48.   Gaynor EC, Szymanski CM. The 30th anniversary of Campylobacter, Helicobacter, and
Related Organisms workshops-what have we learned in three decades? Front Cell Infect
Microbiol. 2012;2:20.
49.   Sheh A, Fox JG. The role of the gastrointestinal microbiome in Helicobacter pylori
pathogenesis. Gut Microbes. 2013;4(6):505-31.

50.   Kienesberger S, Cox LM, Livanos A, Zhang XS, Chung J, Perez-Perez GI, et al. Gastric
Helicobacter pylori Infection Affects Local and Distant Microbial Populations and Host
Responses. Cell Rep. 2016;14(6):1395-407.
51.   Cover TL, Blaser MJ. Helicobacter pylori in health and disease. Gastroenterology.
2009;136(6):1863-73.
52.   Saadat I, Higashi H, Obuse C, Umeda M, Murata-Kamiya N, Saito Y, et al. Helicobacter
pylori CagA targets PAR1/MARK kinase to disrupt epithelial cell polarity. Nature.
2007;447(7142):330-3.
53.   Hatakeyama M, Higashi H. Helicobacter pylori CagA: a new paradigm for bacterial
carcinogenesis. Cancer Sci. 2005;96(12):835-43.
54.   Yamaoka Y. Mechanisms of disease: Helicobacter pylori virulence factors. Nat Rev
Gastroenterol Hepatol. 2010;7(11):629-41.
55.   Murphy G, Michel A, Taylor PR, Albanes D, Weinstein SJ, Virtamo J, et al. Association
of seropositivity to Helicobacter species and biliary tract cancer in the ATBC study. Hepatology.
2014;60(6):1963-71.
56.   Mateos-Munoz B, Perez-de-la-Serna J, Ruiz-de-Leon A, Serrano-Falcon B, Casabona-
Frances S, Velasco-Cerrudo A, et al. Enterohepatic Helicobacter other than Helicobacter pylori.
Rev Esp Enferm Dig. 2013;105(8):477-84.
57.   Zhou D, Wang JD, Weng MZ, Zhang Y, Wang XF, Gong W, et al. Infections of
Helicobacter spp. in the biliary system are associated with biliary tract cancer: a meta-analysis.
Eur J Gastroenterol Hepatol. 2013;25(4):447-54.
58.   Wroblewski LE, Peek RM, Jr., Wilson KT. Helicobacter pylori and gastric cancer:
factors that modulate disease risk. Clin Microbiol Rev. 2010;23(4):713-39.
59.   Segal ED, Cha J, Lo J, Falkow S, Tompkins LS. Altered states: involvement of
phosphorylated CagA in the induction of host cellular growth changes by Helicobacter pylori.
Proc Natl Acad Sci U S A. 1999;96(25):14559-64.
60.   Simpson CD, Anyiwe K, Schimmer AD. Anoikis resistance and tumor metastasis. Cancer
Lett. 2008;272(2):177-85.
61.   Fragiadaki M, Mason RM. Epithelial-mesenchymal transition in renal fibrosis - evidence
for and against. Int J Exp Pathol. 2011;92(3):143-50.
62.   Rout-Pitt N, Farrow N, Parsons D, Donnelley M. Epithelial mesenchymal transition
(EMT): a universal process in lung diseases with implications for cystic fibrosis
pathophysiology. Respir Res. 2018;19(1):136.
63.   Stone RC, Pastar I, Ojeh N, Chen V, Liu S, Garzon KI, et al. Epithelial-mesenchymal
transition in tissue repair and fibrosis. Cell Tissue Res. 2016;365(3):495-506.
64.   Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. J Clin Invest.
2009;119(6):1420-8.
65.   Li L, Li W. Epithelial-mesenchymal transition in human cancer: comprehensive
reprogramming of metabolism, epigenetics, and differentiation. Pharmacol Ther. 2015;150:33-
46.
66.   Sciacovelli M, Frezza C. Metabolic reprogramming and epithelial-to-mesenchymal
transition in cancer. FEBS J. 2017;284(19):3132-44.
67.   Jurberg AD, Brindley PJ. Gene function in schistosomes: recent advances toward a cure.
Front Genet. 2015;6:144.
68.   Pellicano R, Menard A, Rizzetto M, Megraud F. Helicobacter species and liver diseases:
association or causation? Lancet Infect Dis. 2008;8(4):254-60.

69. Holcombe C. Helicobacter pylori: the African enigma. Gut. 1992;33(4):429-31.
70. Du Y, Agnew A, Ye XP, Robinson PA, Forman D, Crabtree JE. Helicobacter pylori and
Schistosoma japonicum co-infection in a Chinese population: helminth infection alters humoral
responses to H. pylori and serum pepsinogen I/II ratio. Microbes Infect. 2006;8(1):52-60.
71. Whary MT, Sundina N, Bravo LE, Correa P, Quinones F, Caro F, et al. Intestinal
helminthiasis in Colombian children promotes a Th2 response to Helicobacter pylori: possible
implications for gastric carcinogenesis. Cancer Epidemiol Biomarkers Prev. 2005;14(6):1464-9.
72. Fox JG, Beck P, Dangler CA, Whary MT, Wang TC, Shi HN, et al. Concurrent enteric
helminth infection modulates inflammation and gastric immune responses and reduces
helicobacter-induced gastric atrophy. Nat Med. 2000;6(5):536-42.
746