Infection with CagA+ *Helicobacter pylori* induces epithelial to mesenchymal transition in human cholangiocytes

Prissadee Thanaphongdecha1,2,3, Shannon E. Karinshak1, Wannaporn Ittiprasert1, Victoria H. Mann1, Yaovalux Chamgramol3, Chawalit Pairojkul3, James G. Fox4, Sutas Suttiprapa2, Banchob Sripa 2,3,*, Paul J. Brindley1,*

1 Department of Microbiology, Immunology and Tropical Medicine, and Research Center for Neglected Tropical Diseases of Poverty, School of Medicine & Health Sciences, The George Washington University, Washington DC, 20037, USA

2 Tropical Disease Research Laboratory, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand

3 Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand

4 Division of Comparative Medicine, Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, 02139, MA

* Equal contribution

Correspondence: Paul J. Brindley, Department of Microbiology, Immunology and Tropical Medicine, and Research Center for Neglected Tropical Diseases of Poverty, School of Medicine & Health Sciences, The George Washington University, Washington DC, 20037, USA, email, pbrindley@gwu.edu; or Banchob Sripa, Tropical Disease Research Laboratory, Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand, email, banchob@kku.ac.th

Abstract

Recent reports suggest that the East Asian liver fluke *Opisthorchis viverrini*, which is implicated in pathogenesis of opisthorchiasis-associated cholangiocarcinoma, serves as a reservoir of *Helicobacter pylori*. The affected cholangiocytes that line the intrahepatic biliary tract are considered to be the cell of origin of this malignancy. Here, we investigated interactions among a human cholangiocyte cell line, H69, a CCA cell line CC-LP-1, CagA$_+$ve *Helicobacter pylori*, and *H. bilis* bacilli. Exposure to increasing numbers of *H. pylori* at 0, 1, 10,100 bacteria per cholangiocyte cell induced phenotypic changes in the cultured cholangiocytes including profusion of thread-like filopodia and loss of cell-cell contact in a dose-dependent fashion. In parallel, changes in mRNA expression followed exposure to *H. pylori*, with increased expression of epithelial to mesenchymal transition (EMT) associated-factors including snail, slug, vimentin, matrix metalloprotease, zinc finger E-box-binding homeobox, and the cancer stem cell marker CD44. Transcription levels encoding the cell adhesion marker CD24 decreased. Analysis in real time using the xCELLigence system to quantify cell proliferation, migration and invasion revealed that exposure to ten to 50 of *H. pylori* stimulated migration of H69 cells and CC-LP-1 cells, a line derived form of a human cholangiocarcinoma, and invasion through an extracellular matrix. In addition, ten bacilli of CagA$_+$ve *H. pylori* stimulated contact-independent colony establishment in soft agar. These findings support the hypothesis that infection with *H. pylori* can contribute to the malignant transformation of the biliary epithelium.

Keyword : *Helicobacter pylori*, cholangiocyte, epithelial-to-mesenchymal transition.

Introduction

Complex associations occur between trematodes and bacterial commensals, including species capable of causing human diseases such as non-typhoidal salmonellosis, leptospirosis and Sennetsu fevers (1-3). There is increasing evidence that the East Asian liver fluke *Opisthorchis viverrini* may serve as a reservoir of *Helicobacter* and implicate *Helicobacter* in pathogenesis of opisthorchiasis-associated cholangiocarcinoma (CCA) (4-8). The International Agency for Research on Cancer of the World Health Organization classifies infection with the liver flukes *O. viverrini* and *Clonorchis sinensis* and with *H. pylori* as Group 1 carcinogens (9). In northern and northeastern Thailand and Laos, infection with *O. viverrini* is the major risk for CCA (9-12). Given the elevated prevalence of CCA in this region where infection with liver fluke prevails, and given evidence of linkages between human infection by *Helicobacter* during opisthorchiasis, these two biological carcinogens together may orchestrate the pathogenesis of opisthorchiasis and bile duct cancer. The association of *Helicobacter* and its virulence factors, including during opisthorchiasis, may underlie biliary tract disease including CCA in liver fluke-endemic regions (13). Infection with species of *Helicobacter* causes hepatobiliary tract diseases that can resemble opisthorchiasis (8, 14, 15) (16, 17). Lesions ascribed to liver fluke infection, including cholangitis, biliary hyperplasia and metaplasia, periductal fibrosis and CCA, may be due in part to *Helicobacter*-associated hepatobiliary disease.

Commensalism involving *H. pylori* and *O. viverrini* may have evolved and may facilitate conveyance of the bacillus into the human biliary tract during the migration of the juvenile fluke following ingestion of the metacercaria in raw or undercooked cyprinid fish and establishment of infection (4, 18). Since we have shown previously that curved rods resembling *Helicobacter* occur in the alimentary tract of the liver fluke (4), and given the low pH of the gut of *O. viverrini*, the *H. pylori* rods or spores might be transported by the migrating parasite. The spiral shape of *H. pylori* attach to biliary cells, which internalize in similar fashion to their behavior on gastric epithelium (19, 20).

Transition of epithelial cells to mesenchymal cells during disease and epithelial to mesenchymal transition (EMT) during development and wound healing follows conserved EMT routes with well-characterized molecular and morphological hallmarks (21). These hallmarks are clearly displayed at malignant transformation of the epithelium of the gastric mucosa resulting from infection with *H. pylori* (22). Here, we investigated interactions between CagA+ve *H. pylori*, the related species *H. bilis* (23), and human cholangiocytes including the H69 cholangiocyte cell line (24, 25) and the CCLP-1cholangiocarcinoma cell line (26). Infection with CagA+ *H. pylori* induced epithelial to mesenchymal transition in human cholangiocytes.

Materials and Methods

Cell lines of human cholangiocytes

The immortalized intrahepatic cholangiocyte cell line, H69 (24, 25), Cellosaurus (27) identifier RRID:CVCL_812,1, and the primary cholangiocarcinoma cell line, CC-LP-1 (26, 28) Cellosaurus RRID CVCL_0205, were obtained as described (26, 28-30). In brief, H69 cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) (Thermo Fisher Scientific, Inc.),

DMEM/F12 (Millipore-Sigma, St. Louis, MO) supplemented with 10% fetal bovine serum
(FBS)(Invitrogen; Thermo Fisher Scientific, Inc.), supplemented with adenine, insulin,
epinephrine, triiodothyronine/transferrin, hydrocortisone and epidermal growth factor and
penicillin/streptomycin (PS) (all from Millipore-Sigma). CC-LP-1 cells were cultured in DMEM
containing 10%FBS, L-glutamine and antibiotics, as above. Both cell lines were maintained in
humidified 5% CO_2 in air at 37°C. The cells were cultured to ~80% confluence before co-culture
with *H. pylori*.
111
112 ***Helicobacter pylori* and *Helicobacter bili*s**
113
Helicobacter pylori, ATCC 43504, isolated originally from human gastric antrum (31), and *H.*
bilis, Fox *et al.* ATCC 49314, originally isolated from intestines and liver of mice (32, 33), were
obtained from the American Type Culture Collection (ATCC) (Manassas, VA) and maintained
on trypticase soy agar with 5% sheep blood for 72 to 96 hours (34) (Becton Dickinson,
Cockeysville, MD), incubated at 37°C with gentle agitation under microaerobic conditions using
the BBL Campy Pack Plus system (Becton Dickinson). After 96 hours, each species of
Helicobacter species was harvested independently by scraping the colonies from the agar plate,
the cell scraping was resuspended in DMEM/F12 medium to an optical density at 600 nm
(OD_{600}) of 1.0 corresponding to ~1×10^8 colony forming units (cfu)/ml (35-37).
123
124 **Morphological assessment of cholangiocytes**
125
126 *H. pylori* was co-cultured with H69 cells at increasing multiplicities of infection (MOIs) of 0 (no
127 bacilli), 10 and 100 for 24 hours in serum-free media. After 24 hours, the effects of *H. pylori on*
128 H69 cell morphology were documented and images captured using an inverted microscope
129 (Zeiss Axio Observer A1, Jena, Germany).
130
131 **Cell scattering and elongation**
132
Cells were seeded at 5×10^5 cells/well in 6-well culture plates. One day later, the cells were fed
with serum- and hormone free medium in the presence of *H. pylori* at MOI of 0, 10 and 100
respectively and maintained for 24 more hours. At that interval, scattering of cells was
documented using a camera fitted to an inverted microscope (Zeiss Axio Observer A1, Jena,
Germany). Cell scattering was quantified by counting the total number of isolated single cells per
field in ten randomly selected images at 5× magnification (modified from (38)). To assess
elongation of cells, cells from ~20 randomly selected fields of view at ×20 magnification, with
two to seven cells per field, were documented as above. The length to width ratio of the isolated
cells was established using the ImageJ software (39).
142
143 ***In vitro* wound healing assay**
144
Monitoring cell migration in a two-dimensional confluent monolayer in controlled *in vitro*
conditions allowed investigators to simulate and explore critical mechanisms of the impact of *H.*
pylori on the process of wound healing. A sheet migration approach (40, 41) was used in this
study to monitor wound closure in vitro following exposure of the cholangiocytes to *H. pylori* at
MOI of 0, 10 and 100. H69 cells were grown in 6-well plates (Greiner Bio-One, Monroe NC)

overnight to allow cell adherence. The cell monolayer was wounded with a thin linear scratch in
the confluent cell monolayer, using a sterile 20 µl pipette tip, and the impact and wound closured
monitored and measured 26 hours later, as we have described previously (29, 30). The width of
the gap was measured (ImageJ software, NIH) and rate of wound closure rate compared with
control cells not exposed to *H. pylori*.
155
156   **Real time quantitative PCR (RT-qPCR)**
157
Total RNA from H69 cells were extracted with RNAzolʀᴛ (Molecular Research Center, Inc.,
Cincinnati, OH) according to the manufacturer's instructions. The RNA quality and
concentration were assessed using the NanoDrop 1000 spectrophotometer (Thermo Fisher
Scientific). Total RNA (500 ng) was reversed-transcribed using iScript cDNA Synthesis Kit
(Bio-Rad, Hercules, CA). Quantitative reverse transcription-polymerase chain reaction (qRT-
PCR) analysis of mRNA expression of six members of the EMT-associated genes including
Vimentin, Snai1, Slug, Zeb, Jam1 and MMP7; two cancer stem cells including CD44 and CD24
were undertaken on an ABI7300 system (ABI) using SsoAdvance SYBR green mixture (Bio-
Rad), normalized to human GAPDH (Supplementary Table 1 provides the primer sequences).
The relative fold-change was determined by the ΔΔCt method (42). Three biological replicates
of the assay were undertaken.
169
170   **Assessment of cell proliferation**
171
Cell proliferation of H69 was assessed using the xCELLigence Real-time cell analyzer (RTCA)
DP instrument (ACEA Biosciences, San Diego CA), as reported (29, 43, 44). H69 cells were
fasted for 4 to 6 hours in 1:20 diluted medium that had been diluted 1:20 with 50% DMEM and
50% DMEM/Ham's F12, containing 1% of PS, *i.e.* 0.5% FBS final concentration, as described
(45, 46), then the cells were co-cultured of viable bacilli of *H. bilis* for 120 min before starting
the assay. H69 cells were harvested later using 0.25% trypsin-EDTA and washed with medium.
Five thousand H69 cells were seeded on to each well of the E-plate in H69 medium and cultured
for 24 hours. The culture medium was removed, and the cells were gently washed with 1×PBS.
The PBS was replaced with serum diluted medium. The cellular growth was monitored in real
time with readings collected at intervals of 20 minutes for ≥ 60 hours. For quantification, CI was
averaged from three independent measurements at each time point.
183
184   **Cell migration and invasion in response to *H. pylori***
185
To monitor the rate of cell migration and/or invasion in real time, we used the xCELLigence DP
instrument (above) equipped with a CIM-plate 16 (ACEA Biosciences, San Diego CA), which is
a 16-well Boyden chamber-like system in which each well is composed of upper and lower
chambers separated by an 8-um microporous membrane (Ref). Migration/invasion was
measured as the relative impedance change (cell index) across microelectronic sensors integrated
into the bottom of the membrane separating the upper chamber (UC) and lower chamber (LC) of
the well.
193
To investigate migration, H69 and CC-LP-1 cells were infected with *H. pylori* ATCC45304
strain at MOIs of 0, 1, 10, 50 and 100 respectively for 2 hours before the start of assay, after

which the cells were harvested following treatment for 3 min with 0.25% trypsin-EDTA and
washed with medium. Subsequently, 30,000 H69 or CC-LP-1 cells were seeded into each UC
well in serum-free medium. Complete medium supplemented with 10 % serum was added to
each LC well. Next, equilibration of the loaded CIM-plate was accomplished by incubation for
60 min to allow cell attachment, after which monitoring for migration was commenced in real
time for a duration of ~100 hours with readings recorded in real time at 20 min intervals. For
quantification, the cell index (CI) was averaged from three independent measurements at each
time point (47).
204
For the invasion assay, two hours prior to conducting the assay, H69 or CC-LP-1 cells were
infected with *H. pylori* ATCC45304 strain at either MOI of 0 or 10. The UC of the CIM plate
was coated with a 20% solution of Matrigel™ (BD Biosciences). A total of 30,000 of H69 or
CC-LP-1 cells were seeded in each well of the UC in serum-free media. H69 or CC-LP-1
medium was added as appropriate to each well of the lower chamber (LC). The loaded CIM-
plate was maintained at 37_{o} C in 5% CO_2 for 60 minutes to facilitate attachment of the cells. The
impedance value of each well was recorded at intervals of 20 minutes for ~100 hours, at which
point the experiment was terminated. The CI value provided a measurement of the rate of
invasion.
214
215 **Colony formation in soft agar**
216
H69 cells were infected with *H. pylori* or *H. bilis* at MOI of 0, 10, 50 and 100 at the start of the
assay. For the soft agar assay, H69 cells at 10,000 cells per well were mixed with 0.3% low
melting temperature agarose (NuSieve GTG) (Lonza, Walkersville, MD) in growth medium, and
plated on top of a solidified layer of 0.6% agarose in growth medium, in wells of 6- or 12-well
culture plates (Greiner Bio-One). New growth medium was added every three or four days until
28 days or until the colonies had established, at which time the number of colonies of $\geq$ 50 µm
diameter was counted and documented using an inverted microscope (48).
224
225 **Statistical analysis**
226
Differences in cell elongation, cell scattering, wound healing and colony formation among test
and control groups were analysed by one-way ANOVA. Fold differences in mRNA levels, real
time cell migration and invasion through extracellular matrix were analysed by two-way
ANOVA followed by Dunnett's test for multiple comparisons. Analysis was accomplished using
GraphPad Prism version 7 software (GraphPad, San Diego, CA). A *P* value of $\leq$ 0.05 was
considered to be statistically significant.
233
234 **Results**
235
236 ***H. pylori* induces epithelial to mesenchymal transition (EMT) in cholangiocytes**
237
H69 were directly exposed to increasing number of *H. pylori* at 0, 10 and 100 bacilli per
cholangiocyte in 6-well plates. At 24 hours after addition of the bacilli, the normal epithelial cell
appearance in tissue culture had altered to an elongated phenotype characterized by terminal
thread-like filopodia and diminished cell-to-cell contacts (Figures 1,2). The extent of the

transformation was dose-dependent and, also, it indicated increased cell motility. The
morphological change of H69 was quantified by measurement of the length to width ratio of the
exposed cells. In a dose dependent fashion, the length: width ratio increased significantly at both
MOI of and MOI of 100 when compared with MOI of 0 ($P \leq 0.01$) (Figure 2C). In addition, the
numbers of isolated, single cells increased significantly in similar fashion to the increasing
length: width ratio of the *H. pylori*-exposed H69 cells (Figure 2D, $P \leq 0.05$ and ≤ 0.001 at MOI
10 and 100, respectively). This phenotype resembled the 'hummingbird phenotype' displayed by
the AGS gastric cell line following exposure to *H. pylori* (22). Moreover, H69 cells culture in
medium supplemented with TGF-β at 5 ng/ml displayed cell aggregation and nodule formation
(Figure S1).
252
253 **Expression of epithelial to mesenchymal transition (EMT) associated-factors induced by**
254 **exposure to *H. pylori***
255
The dynamic change to EMT at transcriptional level was investigated using quantitative real-
time PCR of total cellular RNAs for six EMT-related factors including mesenchymal marker
(vimentin), transcription factors (Snail, Slug, Zeb1), adhesion molecules (Jam1) and proteolytic
enzyme (MMP7) (49). The cancer stem cell markers, CD24 and CD44 (50) were included in this
analysis. Levels of the fold-difference in transcription for each of the six markers increased in a
dose-dependent fashion. The regulatory transcriptional factor, Snail was markedly up-regulated
with highest level of fold difference, 88-, 656- and 5,309-fold at MOI of 10, 50 and 100,
respectively, followed by MMP7, with 15-, 21-, and 44-fold at MOI of 10, 50 and 100.
Transcription of Slug, Zeb1, vimentin and Jam1 also was up-regulated during bacterial infection.
CD24 expression was down-regulated whereas CD44 was up-regulated, revealing a CD44+
high/CD24+ low pattern in cells exposed to *H. pylori* (Figure 3; $P \leq 0.05$ to ≤ 0.0001 for
individual EMT markers and/or MOI level as shown). A pattern of CD44+ high/CD24+ low
expression is a hallmark biomarker of cancer stem cell activity in gastric carcinoma (50).
269
270 **Exposure to *H. pylori* induces cholangiocytes to migrate, to invade extracellular matrix,**
271 **and to close wounds**
272
Analysis in real time using the CIM plates fitted to the xCELLigence system a revealed that
exposure to *H. pylori* at MOI of 10 to 50 *H. pylori* stimulated significantly more migration of the
H69 cholangiocytes from 24 to 96 hours from the start of real time monitoring. (Figure 4A; the
graph includes *P* values at representative time points and MOI levels). The effect was not seen
with MOI 100. In addition, a similar assay was undertaken using the CC-LP-1
cholangiocarcinoma cell line, but with a lower *H. pylori* MOI range. At MOI of 1, 10 and 50, *H.*
pylori stimulated significantly more migration of CC-LP-1 cells from 20 to 40 hours of the real
time monitoring (Figure 4B). Concerning invasion of extracellular matrix, both the CC-LP-1 and
H69 cells migrated and invaded the Matrigel layer in the CIM plate at significantly higher rates
than did the control cells, with significant differences evident from about 50 until 100 hours
when the assay was terminated (Figure 4C, 4D). Third, scratch assays revealed two-dimensional
migration of H69 cells over 24 hours. In vitro wound closure by H69 cells increased significantly
to 19.47% in MOI of 10 ($P \leq 0.05$) although an effect at MOI 100 was not apparent in
comparison with the control group (Figure S2).
287

288 ***H. pylori* induces anchorage-independent colony formation**
289
As a prospective biomarker of malignant transformation of the *Helicobacter*-infected
cholangiocytes, anchorage independent colony formation by the cells was determined in a
medium of soft agar. Following maintenance of the cultures for up to 28 days, counts of the
numbers of colonies was revealed that a MOI of 10 *H. pylori* had induced a significant increase
colony numbers of the H69 cells (Figure 5A, 5B). Significant differences from the control group
were not see at MOI 50 or 100. The size of colonies also increased in all groups exposed to *H.*
pylori although this was statistically significant only at the MOI of 50 (Figure S3; $P \le 0.05$). By
contrast to *H. pylori*, exposure of H69 cells to *H. bilis*, a microbe that naturally resides in the
biliary tract and intestines (33, 51), decreased the number of colony formation in soft agar
(Figure 5D). The ostensibly neutral or even inhibitory effect of *H. bilis* on H69 biliary cell
proliferation was mirrored by inapparent increase of cellular proliferation over several days when
monitored in real-time by xCELLigene using MOI of 0, 10, 50 and 100 *H. bilis* (Figure S4).
302
303 **Discussion**
304
Numerous species of *Helicobacter* have been described (52) while *H. pylori* was the first
bacterial pathogen confirmed to cause gastric disease including peptic ulcer, gastric lymphoma
and gastric adenocarcinoma (31, 53-56). Bacilli of *H. pylori* occur in the stomach in at least half
the human population. Transmission from mother to child and by other routes takes place. The
human-*H. pylori* association may be beneficial in early life, including contributions to a healthy
microbiome and reduced early-onset asthma (57, 58). Many strains of *H. pylori* have been
characterized at the molecular level, and these are generally classified as either CagA positive or
CagA negative. The major virulence factor, cytotoxin-associated gene A (CagA) is a ~140 kDa
protein encoded on the cag pathogenicity island (PAI). Cag PAI also encodes a type 4 secretion
channel through which the CagA virulence factor is introduced into the host cell upon attachment
by the bacillus (22). CagA locates to the cell membrane, from where it is phosphorylated by the
Src kinases. Downstream of these modifications, the activated CagA orchestrates changes in
morphology of the infected epithelial cell, which loses cell polarity and becomes motile,
exhibiting a diagnostic 'hummingbird-like' phenotype (59-61).
319
The CagA oncoprotein secreted by *H. pylori* into epithelial cells is noted for its variation at the
SHP2 binding site and, based on the sequence variation, is subclassified into two main types -
East-Asian CagA and Western CagA. East-Asian CagA shows stronger SHP2 binding and
greater biological activity than Western CagA. It has been proposed that in East Asia, the
circulation of *H. pylori* strains encoding active forms of CagA underlie the elevated incidence of
gastric adenocarcinoma (22).
326
In addition to the well-known association with gastric cancer, *H. pylori* has also been associated
with hepatological disease including in the biliary tract (7, 8, 62). The related species, *H.*
hepaticus and *H. bilis*, also have been associated with hepatobiliary disease (62-64). Infection
with the fish-borne liver flukes *Opisthorchis viverrini, Clonorchis sinensis* as well as the
bacterial infection with *Helicobacter pylori* are all classified as Group 1 carcinogens by the
International Agency for Research on Cancer of the World Health Organization (9).
Opisthorchiasis is a major risk factor for cholangiocarcinoma in the northern and northeastern

part of Thailand and Laos (9-12). In addition to gastric disease, infection with species of
Helicobacter causes hepatobiliary tract diseases that can resemble opisthorchiasis (8, 14, 15) (16,
17). Liver fluke infection can induce lesions in biliary system including cholangitis, biliary
hyperplasia and metaplasia, periductal fibrosis and CCA. These lesions derive not only from
liver fluke infection but perhaps are in part the consequence of hepatobiliary tract infection with
H. pylori. *Helicobacter* may transit from the stomach to the duodenum and enter the biliary tree
through the duodenal papilla and ampulla of Vater (19, 65), and indeed may be vectored there by
the liver fluke, O. viverrini ((4-6). *H. pylori*-specific DNA sequences have been detected in
CCA tumors and also from lesions diagnosed as cholecystitis/cholelithiasis in regions endemic
for opisthorchiasis (8, 14). Furthermore, serological findings have implicated indicate infection
with *H. pylori* as a risk for CCA in Thailand (15).

Here, we investigated interactions among a human cholangiocyte cell line, H69, a CCA cell line
CC-LP-1, CagA+ve *Helicobacter pylori*, and *H. bilis* bacilli. Infection of H69 with CagA+ *H.*
pylori induced epithelial to mesenchymal transition phenotype in dose-dependent manner, which
was characterized by cell elongation and scattering which implicate increasing change in cell
motility. This phenotype of H69-infected cells was reminiscent of the modified appearance of the
AGS gastric carcinoma cell after exposure *H. pylori*, which has been termed the hummingbird
phenotype (66). In AGS cells, delivery of CagA by the type IV secretion mechanism from *H.*
pylori to the epithelial cell subverts the normal signaling leading to actin-dependent
morphological rearrangement and the hummingbird appearance. The hitherto uniform polygonal
shape becomes markedly elongated with terminal needle-like projections (67).

Earlier studies in AGS cells infected with *H. pylori* had also demonstrated that the hummingbird
phenotype was accompanied early on with transcription changes that reflected the epithelial to
mesenchymal transition, EMT. Similar transcriptional changes were seen in the *H. pylori*-
infected H69 cells. Specifically, upregulation of expression of Snai1, Slug, vimentin, Jam1, and
MMP7 expression in dose-dependent fashion strongly supported, at the expression level, a
dynamic change in cholangiocyte reflective of EMT. Moreover, the E-cadherin repressor, Snail
was markedly up-regulated, which indicated that this factor may be a key driver of epithelial to
mesenchymal transition in H69. CD44 expression was up-regulated in dose-dependent fashion,
whereas CD24 decreased, showing a CD44 high/CD24 low phenotype during infection. These
results indicated that *H. pylori* may not only induce EMT but also contribute to stemness and
malignant transformation of the cholangiocyte.

Realtime cell monitoring approach by xCELLigence revealed that the H69 and CC-LP-1 cells
showed higher cell migration when exposed to *H. pylori*. The fastest migratory rate was evident
at MOI of 10, followed by MOI of 50. By contrast, the rate decreased with 100 bacteria per cell,
suggesting a direct detrimental effect of exposure to elevated levels of the bacilli. Ten bacilli per
cell stimulated both migration and invasion of the basement membrane model matrix by H69 and
by CC-LP-1 cells, a behavioral phenotype characteristic generally of epidermal to mesenchymal
transition. In addition, following exposure to *H. pylori*, H69 cells responded with an anchorage-
independent cell growth in soft agar, which is an in vitro phenotype indicative of the escape from
anoikosis and is characteristic of metastasis (68). The numbers of colonies of H69 cells in soft
agar were significantly increased at a *H. pylori* MOI of 10. By contrast, after exposure of H69
cells to *H. bilis* in the model of contact independent cell growth and also proliferation as

quantified on the xCELLigene platform, both numbers of colonies and cell growth decreased.
These responses indicated that CagA+ *H. pylori* has the potential to induce carcinogenic changes
in cholangiocytes in like fashion to its carcinogenicity for the gastric epithelia.
383
The findings presented here conform with a malignancy-conducive relationship between the liver
fluke *O. viverrini* and the bacilli *H. pylori*. By contrast, co-infections of *Helicobacter* species
including *H. pylori* and some other helminths have generally been associated with diminished
risk of *H. pylori*-associated gastric carcinoma (69-71). For example, the concurrent infection on
mice with an intestinal nematode modulates inflammation, induces a Th2-polarizing cytokine
phenotype with concurrent downmodulation of Th1 responses and the gastric immune responses,
and reduces *Helicobacter*-induced gastric atrophy (72). Nonetheless, given that CagA+ *H.
pylori* stimulated epithelial to mesenchymal transition of cholangiocytes which, in turn indicates
the underlying mechanism of tissue fibrosis (73-75) and cancer metastasis (76-78), an
explanation for why infection with the liver fluke induces bile duct cancer (79) might now be
clearer – involvement by *H. pylori* and its virulence factors. *Helicobacter* likely passes from the
stomach to the duodenum and enters the biliary tree through the duodenal papilla and ampulla of
Vater (19, 65). To conclude, infection with CagA₊ *H. pylori* induced epithelial to mesenchymal
transition in human cholangiocytes, along with other changes strongly reflective of malignant
transformation. Further investigation of the relationship of *O. viverrini* and *H. pylori* within the
infected biliary tract clearly is warranted.
400
401 **Abbreviations**   Cytotoxin-associated gene A (*cag*A), cholangiocarcinoma (CCA); CagA
402 multimerization motif (CM); EMT, epithelial to mesenchymal transition; ZEB, zinc finger E-
403 box-binding homeobox 1; VIM, vimentin; SNAII, MMP7 and cancer stem cell marker CD44;
404 CD24.
405
406 **Competing Financial Interests**
407
408 The authors declare there were no competing financial interests.
409
410 **Acknowledgments**
411

P.T. acknowledges support as a PhD research scholar from the Thailand Research Fund under
the Royal Golden Jubilee scholars program (grant number PHD/0013/2555); B.S. acknowledges
support from Thailand Research Fund Senior Research Scholar. This work was supported by the
National Health Security Office, Thailand, the Higher Education Research Promotion and
National Research University Project of Thailand, Office of the Higher Education Commission,
through the Health Cluster (SHeP-GMS), the Faculty of Medicine, Khon Kaen University,
Thailand (award number I56110), and the Thailand Research Fund under the TRF Senior
Research Scholar (RTA 5680006); the National Research Council of Thailand. The National
Institute of Allergy and Infectious Diseases (NIAID), Tropical Medicine Research Center award
number P50AI098639, The National Cancer Institute, award number R01CA164719, and the
United States Army Medical Research and Materiel Command (USAMRMC), contract number
W81XWH-12-C-0267 also provided support. The content is solely the responsibility of the
authors and does not necessarily represent the official views of the funders including
USAMRMC, NIAID, NCI or the NIH.

Author contributions

P.T., S.S., C.P., Y.C., J.F., B.S. and P.J.B. conceived and designed the study. P.T., S.K., V.M., W.I. performed the experiments. P.T., W.I., B.S. and PJB analyzed and interpreted the findings. P.T., B.S., W.I., S.S., and PJB wrote the manuscript. All authors read and approved the final version of the paper.

Figure legends

Figure 1. Exposure to CagA₊ *H. pylori* ATCC 43504 induced morphological alteration in cholangiocytes. The photomicrographs documented cell morphology of H69 cells exposed to increasing numbers of *H. pylori* at 0, 10 and 100 bacilli per cholangiocyte, respectively (left to right) (A). The appearance of cells had changed from an epithelial to a mesenchymal phenotype including loss of cell-cell contact, elongate and spindle-shape along with growth as individual cells at after 24 hours following *H. pylori* infection in dose-dependent manner. Scale bars, 5 μm (right), 20× magnification.

Figure 2. Quantification of cell elongation of cholangiocytes upon exposure to bacilli of CagA₊ *H. pylori* indicated epidermal to mesenchymal transition. The length-to-width ratio of single, isolated single cells were documented to determine cellular elongation and scattering (A, B). The number of elongated cells increased in a dose-dependent fashion in response to infection by *H. pylori* (C). By contrast, the number of isolated individual H69 cells indicated cell scattering was also significantly increased in dose-dependent fashion (D). Data are presented as the mean values with standard errors from three biological replicates. Asterisks indicates statistically significant different of *H. pylori*-infected group compared to normal group at 24 hours (*, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$).

Figure 3. Transcriptional change of EMT-related and cancer stem cell marker genes after *H. pylori* exposure. Messenger RNA expression of 6 EMT-related genes and 2 cancer stem cell markers were determined with three biological replicates after 24 hours of infection. Expression of Snail, Slug, vimentin, Jam, MMP7 and CD44 increased in a dose dependently fashion whereas CD24 transcription decreased. Expression of regulatory transcriptional factor, Snail was notably up-regulated by 88-, 656- and 5,309- fold at MOI 10, 50 and 100, respectively. MMP7 expression was markedly up-regulated at each MOI. Expression of each of Slug, Zeb1, vimentin and Jam1 up-regulated in dose-dependent manner. Transcription of the cancer stem cell marker CD44 was significantly up-regulated at MOI 50 and 100 whereas CD24 was down-regulated. Together, these changes indicated that *H. pylori* could induce both EMT and stemness in human cholangiocytes. The qPCR findings were normalized to the expression levels of GAPDH in each sample.

Figure 4. Exposing cholangiocytes to *H. pylori* induced migration and invasion through extracellular matrix. H69 cholangiocytes infected with *H. pylori* migrated through Matrigel, as monitored in real time over 100 hours using CIM plates fitted to a xCELLigence DP system. Cell migration and invasion were monitored for 100 hours with chemo-attractant in the lower chamber of CIM plate. H69 cells infected with MOI 10 revealed the highest cell migration rate

followed by MOI 50 whereas MOI 100 attenuated cell migration compared with non-infected
control (A). Similarly, the cholangiocarcinoma cell line, CC-LP-1 showed the highest cell
migration rate when stimulated with MOI 10 followed by MOI 50. Moreover, CC-LP-1 cells
significantly migrated faster even with at MOI of 1 (B). Invasion of the Matrigel extracellular
matrix compared between MOI 10 and non-infected cells; invasion rates for H69 and CC-LP-1
cells significantly increased when infected (C,D).
476
Figure 5. Anchorage-independent cell growth in soft agar revealed cellular transformation
of cholangiocytes by *H. pylori*. Representative micrographs of cholangiocyte colony appearance
at day 30 exposed H69 cells (panel A, left to right). At MOI of 10, the number of colonies
increased significantly ($P \leq 0.01$), but not at MOI of 50, and with a significant decrease at MOI
of 100, compared with the non-infected control cells (B). By contrast, exposure of H69 cells to
H. bilis resulted in markedly reduced numbers of colonies, in a dose-dependent fashion (C; $P \leq$
0.001). This indicated an inhibitory effect of *H. bilis* toward anchorage-independent cell growth
and/or cellular transformation of human cholangiocytes (C). Three biological replicates were
performed.

Supporting information

Figure S1. TGF-β promoted nodule formation. TGF-β, an EMT-stimulant, induced nodule
formation of H69 cholangiocytes. Cells were cultured in the presence of 5 ng/ml TGF-β in
serum- and hormone-free medium for 24 hours. Micrographs at 0 (left) and 24 (right) hours;
5×magnification.

Figure S2. Wound healing in a two-dimensional in vitro assay revealed cell migration of
H69 cholangiocytes infected with *H. pylori*. Wound closure was significantly increased to
19.47% at MOI 10 of *H. pylori* ($P \leq 0.05$). An increase in wound closure was not apparent at
MOI 100; 11.09% vs control, 14.47%. Micrographs at 0 (left) and 26 (right) hours;
5×magnification. the assays was performed in triplicate.
496
497 **Figure S3. Size of colonies of cholangiocytes in soft agar.** The colonies with diameter ≥ 50
498 µm were measured and counted;. mean diameters were 77.63, 91.62, 95.25 and 81.46 µm at MOI
499 of 0, 10, 50 and 100, respectively.
500
501 **Figure S4. Growth of *H. bilis* infected cholangiocytes as determined by the xCELLigence**
502 **approach.** H69 cells were infected with increasing number of H. bilis (ATCC 43879) at MOI of
503 0, 10, 50 and 100. Cell growth was inhibited when compared to the uninfected cells, at all MOIs.
504 The +proliferation assay was performed in E-plates in minimal (serum-depleted) medium (29).
505
506

Gene name	Gen Bank Number	Primer sequence (5'-3')	Product size (bp)
CD24	NM_013230.3	Forward CACCCAGCATCCTGCTAGAC	259
		Reverse GAGACCACGAAGAGACTGGC	
CD44	NM_00610.3	Forward GGGAGTCAAGAAGGTGGAGC	237
		Reverse CTGAGACTTGCTGGCCTCTC	
F11r/Jam1	NM_016946.5	Forward GTGCCTACTCGGGCTTTTCT	201
		Reverse GAGCTTGACCTTGACCTCCC	
Gapdh	NM_002046.5	Forward CCCATGTTCGTCATGGGTGT	366
		Reverse TTCTAGACGGCAGGTCAGGT	
MMP7	NM_002423.5	Forward GGAGCTCATGGGGACTCCTA	172
		Reverse GGCCAAGTTCATGAGTTGCAG	
Snai1	NM_005985.3	Forward AGCTCTCTGAGGCCAAGGAT	325
		Reverse GACATTCGGGAGAAGGTCCG	
Vimentin	NM_003380.5	Forward GCAGGAGGCAGAAGAATGGT	216
		Reverse GCAGCTTCAACGGCAAAGTT	
Zeb1	NM_001128128.2	Forward CGCAGTCTGGGTGTAATCGT	477
		Reverse CCATGCCCTGAGGAGAACTG	

507
508
509   **Table S1.** Nucleotide sequences of primers specific for EM- associated human genes, which
510   were used in |quantitative real-time PCR analysis.

References

1. McNulty SN, Tort JF, Rinaldi G, Fischer K, Rosa BA, Smircich P, et al. Genomes of Fasciola hepatica from the Americas Reveal Colonization with Neorickettsia Endobacteria Related to the Agents of Potomac Horse and Human Sennetsu Fevers. PLoS Genet. 2017;13(1):e1006537.

2. Van CD, Doungchawee G, Suttiprapa S, Arimatsu Y, Kaewkes S, Sripa B. Association between Opisthorchis viverrini and Leptospira spp. infection in endemic Northeast Thailand. Parasitol Int. 2017;66(4):503-9.

3. Gendrel D, Kombila M, Beaudoin-Leblevec G, Richard-Lenoble D. Nontyphoidal salmonellal septicemia in Gabonese children infected with Schistosoma intercalatum. Clin Infect Dis. 1994;18(1):103-5.

4. Deenonpoe R, Chomvarin C, Pairojkul C, Chamgramol Y, Loukas A, Brindley PJ, et al. The carcinogenic liver fluke Opisthorchis viverrini is a reservoir for species of Helicobacter. Asian Pac J Cancer Prev. 2015;16(5):1751-8.

5. Deenonpoe R, Mairiang E, Mairiang P, Pairojkul C, Chamgramol Y, Rinaldi G, et al. Elevated prevalence of Helicobacter species and virulence factors in opisthorchiasis and associated hepatobiliary disease. Sci Rep. 2017;7:42744.

6. Sripa B, Deenonpoe R, Brindley PJ. Co-infections with liver fluke and Helicobacter species: A paradigm change in pathogenesis of opisthorchiasis and cholangiocarcinoma? Parasitol Int. 2017;66(4):383-9.

7. Boonyanugomol W, Chomvarin C, Song JY, Kim KM, Kim JM, Cho MJ, et al. Effects of Helicobacter pylori gamma-glutamyltranspeptidase on apoptosis and inflammation in human biliary cells. Dig Dis Sci. 2012;57(10):2615-24.

8. Boonyanugomol W, Chomvarin C, Sripa B, Bhudhisawasdi V, Khuntikeo N, Hahnvajanawong C, et al. Helicobacter pylori in Thai patients with cholangiocarcinoma and its association with biliary inflammation and proliferation. HPB (Oxford). 2012;14(3):177-84.

9. Humans IWGotEoCRt. Biological agents. Volume 100 B. A review of human carcinogens. IARC Monogr Eval Carcinog Risks Hum. 2012;100(Pt B):1-441.

10. Sungkasubun P, Siripongsakun S, Akkarachinorate K, Vidhyarkorn S, Worakitsitisatorn A, Sricharunrat T, et al. Ultrasound screening for cholangiocarcinoma could detect premalignant lesions and early-stage diseases with survival benefits: a population-based prospective study of 4,225 subjects in an endemic area. BMC Cancer. 2016;16:346.

11. Khuntikeo N, Loilome W, Thinkhamrop B, Chamadol N, Yongvanit P. A Comprehensive Public Health Conceptual Framework and Strategy to Effectively Combat Cholangiocarcinoma in Thailand. PLoS Negl Trop Dis. 2016;10(1):e0004293.

12. Aye Soukhathammavong P, Rajpho V, Phongluxa K, Vonghachack Y, Hattendorf J, Hongvanthong B, et al. Subtle to severe hepatobiliary morbidity in Opisthorchis viverrini endemic settings in southern Laos. Acta Trop. 2015;141(Pt B):303-9.

13. Segura-Lopez FK, Guitron-Cantu A, Torres J. Association between Helicobacter spp. infections and hepatobiliary malignancies: a review. World J Gastroenterol. 2015;21(5):1414-23.

14. Boonyanugomol W, Chomvarin C, Sripa B, Chau-In S, Pugkhem A, Namwat W, et al. Molecular analysis of Helicobacter pylori virulent-associated genes in hepatobiliary patients. HPB (Oxford). 2012;14(11):754-63.

15. Pisani P, Whary MT, Nilsson I, Sriamporn S, Wadstrom T, Fox JG, et al. Cross-reactivity between immune responses to Helicobacter bilis and Helicobacter pylori in a population in

Thailand at high risk of developing cholangiocarcinoma. Clin Vaccine Immunol.
2008;15(9):1363-8.
16.   Lvova MN, Tangkawattana S, Balthaisong S, Katokhin AV, Mordvinov VA, Sripa B.
Comparative histopathology of Opisthorchis felineus and Opisthorchis viverrini in a hamster
model: an implication of high pathogenicity of the European liver fluke. Parasitol Int.
2012;61(1):167-72.
17.   Mairiang E, Laha T, Bethony JM, Thinkhamrop B, Kaewkes S, Sithithaworn P, et al.
Ultrasonography assessment of hepatobiliary abnormalities in 3359 subjects with Opisthorchis
viverrini infection in endemic areas of Thailand. Parasitol Int. 2012;61(1):208-11.
18.   Plieskatt JL, Deenonpoe R, Mulvenna JP, Krause L, Sripa B, Bethony JM, et al. Infection
with the carcinogenic liver fluke Opisthorchis viverrini modifies intestinal and biliary
microbiome. FASEB J. 2013;27(11):4572-84.
19.   Boonyanugomol W, Chomvarin C, Hahnvajanawong C, Sripa B, Kaparakis-Liaskos M,
Ferrero RL. Helicobacter pylori cag pathogenicity island (cagPAI) involved in bacterial
internalization and IL-8 induced responses via NOD1- and MyD88-dependent mechanisms in
human biliary epithelial cells. PLoS One. 2013;8(10):e77358.
20.   Hatakeyama M. Helicobacter pylori CagA and gastric cancer: a paradigm for hit-and-run
carcinogenesis. Cell Host Microbe. 2014;15(3):306-16.
21.   Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal
transition. Nat Rev Mol Cell Biol. 2014;15(3):178-96.
22.   Hatakeyama M. Oncogenic mechanisms of the Helicobacter pylori CagA protein. Nat
Rev Cancer. 2004;4(9):688-94.
23.   Fox JG. Helicobacter bilis: bacterial provocateur orchestrates host immune responses to
commensal flora in a model of inflammatory bowel disease. Gut. 2007;56(7):898-900.
24.   Grubman SA, Perrone RD, Lee DW, Murray SL, Rogers LC, Wolkoff LI, et al.
Regulation of intracellular pH by immortalized human intrahepatic biliary epithelial cell lines.
Am J Physiol. 1994;266(6 Pt 1):G1060-70.
25.   Park J, Gores GJ, Patel T. Lipopolysaccharide induces cholangiocyte proliferation via an
interleukin-6-mediated activation of p44/p42 mitogen-activated protein kinase. Hepatology.
1999;29(4):1037-43.
26.   Shimizu Y, Demetris AJ, Gollin SM, Storto PD, Bedford HM, Altarac S, et al. Two new
human cholangiocarcinoma cell lines and their cytogenetics and responses to growth factors,
hormones, cytokines or immunologic effector cells. Int J Cancer. 1992;52(2):252-60.
27.   Bairoch A. The Cellosaurus, a Cell-Line Knowledge Resource. J Biomol Tech.
2018;29(2):25-38.
28.   Han C, Demetris AJ, Michalopoulos GK, Zhan Q, Shelhamer JH, Wu T. PPARgamma
ligands inhibit cholangiocarcinoma cell growth through p53-dependent GADD45 and p21
pathway. Hepatology. 2003;38(1):167-77.
29.   Smout MJ, Sotillo J, Laha T, Papatpremsiri A, Rinaldi G, Pimenta RN, et al.
Carcinogenic Parasite Secretes Growth Factor That Accelerates Wound Healing and Potentially
Promotes Neoplasia. PLoS Pathog. 2015;11(10):e1005209.
30.   Arunsan P, Ittiprasert W, Smout MJ, Cochran CJ, Mann VH, Chaiyadet S, et al.
Programmed knockout mutation of liver fluke granulin attenuates virulence of infection-induced
hepatobiliary morbidity. Elife. 2019;8.
31.   Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with
gastritis and peptic ulceration. Lancet. 1984;1(8390):1311-5.

32.   Dewhirst FE, Fox JG, Mendes EN, Paster BJ, Gates CE, Kirkbride CA, et al. 'Flexispira
rappini' strains represent at least 10 Helicobacter taxa. Int J Syst Evol Microbiol. 2000;50 Pt
5:1781-7.
33.   Fox JG, Yan LL, Dewhirst FE, Paster BJ, Shames B, Murphy JC, et al. Helicobacter bilis
sp. nov., a novel Helicobacter species isolated from bile, livers, and intestines of aged, inbred
mice. J Clin Microbiol. 1995;33(2):445-54.
34.   Hachem CY, Clarridge JE, Evans DG, Graham DY. Comparison of agar based media for
primary isolation of Helicobacter pylori. J Clin Pathol. 1995;48(8):714-6.
35.   Jung SW, Thamphiwatana S, Zhang L, Obonyo M. Mechanism of antibacterial activity of
liposomal linolenic acid against Helicobacter pylori. PLoS One. 2015;10(3):e0116519.
36.   Lin SN, Ayada K, Zhao Y, Yokota K, Takenaka R, Okada H, et al. Helicobacter pylori
heat-shock protein 60 induces production of the pro-inflammatory cytokine IL8 in monocytic
cells. J Med Microbiol. 2005;54(Pt 3):225-33.
37.   Pan H, Zhang Y, He GX, Katagori N, Chen H. A comparison of conventional methods
for the quantification of bacterial cells after exposure to metal oxide nanoparticles. BMC
Microbiol. 2014;14:222.
38.   Masuda M, Kikuchi S, Maruyama T, Sakurai-Yageta M, Williams YN, Ghosh HP, et al.
Tumor suppressor in lung cancer (TSLC)1 suppresses epithelial cell scattering and
tubulogenesis. J Biol Chem. 2005;280(51):42164-71.
39.   Bourzac KM, Satkamp LA, Guillemin K. The Helicobacter pylori cag pathogenicity
island protein CagN is a bacterial membrane-associated protein that is processed at its C
terminus. Infect Immun. 2006;74(5):2537-43.
40.   Bindschadler M, McGrath JL. Sheet migration by wounded monolayers as an emergent
property of single-cell dynamics. J Cell Sci. 2007;120(Pt 5):876-84.
41.   Liang CC, Park AY, Guan JL. In vitro scratch assay: a convenient and inexpensive
method for analysis of cell migration in vitro. Nat Protoc. 2007;2(2):329-33.
42.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8.
43.   Arunsan P, Chaidee A, Cochran CJ, Mann VH, Tanno T, Kumkhaek C, et al. Liver fluke
granulin promotes extracellular vesicle-mediated crosstalk and cellular microenvironment
conducive to cholangiocarcinoma. Neoplasia. 2020;22(5):203-16.
44.   Ke N, Wang X, Xu X, Abassi YA. The xCELLigence system for real-time and label-free
monitoring of cell viability. Methods Mol Biol. 2011;740:33-43.
45.   Matchimakul P, Rinaldi G, Suttiprapa S, Mann VH, Popratiloff A, Laha T, et al.
Apoptosis of cholangiocytes modulated by thioredoxin of carcinogenic liver fluke. Int J Biochem
Cell Biol. 2015;65:72-80.
46.   Rinaldi G, Yan H, Nacif-Pimenta R, Matchimakul P, Bridger J, Mann VH, et al.
Cytometric analysis, genetic manipulation and antibiotic selection of the snail embryonic cell
line Bge from Biomphalaria glabrata, the intermediate host of Schistosoma mansoni. Int J
Parasitol. 2015;45(8):527-35.
47.   Xie H, Lee L, Scicluna P, Kavak E, Larsson C, Sandberg R, et al. Novel functions and
targets of miR-944 in human cervical cancer cells. Int J Cancer. 2015;136(5):E230-41.
48.   Horibata S, Vo TV, Subramanian V, Thompson PR, Coonrod SA. Utilization of the Soft
Agar Colony Formation Assay to Identify Inhibitors of Tumorigenicity in Breast Cancer Cells. J
Vis Exp. 2015(99):e52727.

49.   Zeisberg M, Neilson EG. Biomarkers for epithelial-mesenchymal transitions. J Clin
Invest. 2009;119(6):1429-37.
50.   Nguyen PH, Giraud J, Chambonnier L, Dubus P, Wittkop L, Belleannee G, et al.
Characterization of Biomarkers of Tumorigenic and Chemoresistant Cancer Stem Cells in
Human Gastric Carcinoma. Clin Cancer Res. 2017;23(6):1586-97.
51.   Shomer NH, Dangler CA, Schrenzel MD, Fox JG. Helicobacter bilis-induced
inflammatory bowel disease in scid mice with defined flora. Infect Immun. 1997;65(11):4858-
64.
52.   Flahou B, Rimbara E, Mori S, Haesebrouck F, Shibayama K. The Other Helicobacters.
Helicobacter. 2015;20 Suppl 1:62-7.
53.   Cover TL. Helicobacter pylori Diversity and Gastric Cancer Risk. MBio.
2016;7(1):e01869-15.
54.   Marshall BJ. The pathogenesis of non-ulcer dyspepsia. Med J Aust. 1985;143(7):319.
55.   Gaynor EC, Szymanski CM. The 30th anniversary of Campylobacter, Helicobacter, and
Related Organisms workshops-what have we learned in three decades? Front Cell Infect
Microbiol. 2012;2:20.
56.   Sheh A, Fox JG. The role of the gastrointestinal microbiome in Helicobacter pylori
pathogenesis. Gut Microbes. 2013;4(6):505-31.
57.   Kienesberger S, Cox LM, Livanos A, Zhang XS, Chung J, Perez-Perez GI, et al. Gastric
Helicobacter pylori Infection Affects Local and Distant Microbial Populations and Host
Responses. Cell Rep. 2016;14(6):1395-407.
58.   Cover TL, Blaser MJ. Helicobacter pylori in health and disease. Gastroenterology.
2009;136(6):1863-73.
59.   Saadat I, Higashi H, Obuse C, Umeda M, Murata-Kamiya N, Saito Y, et al. Helicobacter
pylori CagA targets PAR1/MARK kinase to disrupt epithelial cell polarity. Nature.
2007;447(7142):330-3.
60.   Hatakeyama M, Higashi H. Helicobacter pylori CagA: a new paradigm for bacterial
carcinogenesis. Cancer Sci. 2005;96(12):835-43.
61.   Yamaoka Y. Mechanisms of disease: Helicobacter pylori virulence factors. Nat Rev
Gastroenterol Hepatol. 2010;7(11):629-41.
62.   Murphy G, Michel A, Taylor PR, Albanes D, Weinstein SJ, Virtamo J, et al. Association
of seropositivity to Helicobacter species and biliary tract cancer in the ATBC study. Hepatology.
2014;60(6):1963-71.
63.   Mateos-Munoz B, Perez-de-la-Serna J, Ruiz-de-Leon A, Serrano-Falcon B, Casabona-
Frances S, Velasco-Cerrudo A, et al. Enterohepatic Helicobacter other than Helicobacter pylori.
Rev Esp Enferm Dig. 2013;105(8):477-84.
64.   Zhou D, Wang JD, Weng MZ, Zhang Y, Wang XF, Gong W, et al. Infections of
Helicobacter spp. in the biliary system are associated with biliary tract cancer: a meta-analysis.
Eur J Gastroenterol Hepatol. 2013;25(4):447-54.
65.   Pellicano R, Menard A, Rizzetto M, Megraud F. Helicobacter species and liver diseases:
association or causation? Lancet Infect Dis. 2008;8(4):254-60.
66.   Wroblewski LE, Peek RM, Jr., Wilson KT. Helicobacter pylori and gastric cancer:
factors that modulate disease risk. Clin Microbiol Rev. 2010;23(4):713-39.
67.   Segal ED, Cha J, Lo J, Falkow S, Tompkins LS. Altered states: involvement of
phosphorylated CagA in the induction of host cellular growth changes by Helicobacter pylori.
Proc Natl Acad Sci U S A. 1999;96(25):14559-64.

68. Simpson CD, Anyiwe K, Schimmer AD. Anoikis resistance and tumor metastasis. Cancer
Lett. 2008;272(2):177-85.
69. Holcombe C. Helicobacter pylori: the African enigma. Gut. 1992;33(4):429-31.
70. Du Y, Agnew A, Ye XP, Robinson PA, Forman D, Crabtree JE. Helicobacter pylori and
Schistosoma japonicum co-infection in a Chinese population: helminth infection alters humoral
responses to H. pylori and serum pepsinogen I/II ratio. Microbes Infect. 2006;8(1):52-60.
71. Whary MT, Sundina N, Bravo LE, Correa P, Quinones F, Caro F, et al. Intestinal
helminthiasis in Colombian children promotes a Th2 response to Helicobacter pylori: possible
implications for gastric carcinogenesis. Cancer Epidemiol Biomarkers Prev. 2005;14(6):1464-9.
72. Fox JG, Beck P, Dangler CA, Whary MT, Wang TC, Shi HN, et al. Concurrent enteric
helminth infection modulates inflammation and gastric immune responses and reduces
helicobacter-induced gastric atrophy. Nat Med. 2000;6(5):536-42.
73. Fragiadaki M, Mason RM. Epithelial-mesenchymal transition in renal fibrosis - evidence
for and against. Int J Exp Pathol. 2011;92(3):143-50.
74. Rout-Pitt N, Farrow N, Parsons D, Donnelley M. Epithelial mesenchymal transition
(EMT): a universal process in lung diseases with implications for cystic fibrosis
pathophysiology. Respir Res. 2018;19(1):136.
75. Stone RC, Pastar I, Ojeh N, Chen V, Liu S, Garzon KI, et al. Epithelial-mesenchymal
transition in tissue repair and fibrosis. Cell Tissue Res. 2016;365(3):495-506.
76. Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. J Clin Invest.
2009;119(6):1420-8.
77. Li L, Li W. Epithelial-mesenchymal transition in human cancer: comprehensive
reprogramming of metabolism, epigenetics, and differentiation. Pharmacol Ther. 2015;150:33-
46.
78. Sciacovelli M, Frezza C. Metabolic reprogramming and epithelial-to-mesenchymal
transition in cancer. FEBS J. 2017;284(19):3132-44.
79. Jurberg AD, Brindley PJ. Gene function in schistosomes: recent advances toward a cure.
Front Genet. 2015;6:144.
737

Figure 2

Figure axes: CI (y-axis) versus Time after addition of cells (hours) (x-axis).

Legend: Control; 10 H.bilis per cell; 50 H.bilis per cell; 100 H.bilis per cell