The impact of purifying and background selection on the inference of population history:

problems and prospects

Parul Johri[*, †], Kellen Riall[*], and Jeffrey D. Jensen[*, †]

[*]School of Life Sciences, Arizona State University, Tempe, AZ 85287

[†]Correspondence can be addressed to these authors

ABSTRACT

Current procedures for inferring population history are generally performed under the

assumption of complete neutrality - that is, neglecting both direct selection and the effects of

selection on linked sites. We here examine how the presence of purifying and background

selection may bias demographic inference by evaluating two commonly-used methods (MSMC

and *fastsimcoal2*), specifically studying how the underlying shape of the distribution of fitness

effects (DFE) and the fraction of directly selected sites interacts with demographic parameter

estimation. The results demonstrate that, even after masking functional genomic regions,

background selection effects may result in the mis-inference of population growth under models

of both constant population size as well as decline. This effect is amplified as the strength of

purifying selection and the density of directly selected sites increases. We propose a potential

solution to correct for this mis-inference. By treating the DFE as a nuisance parameter and

averaging across all potential realizations, we demonstrate that even directly selected sites may

be used to infer demographic histories with reasonable accuracy.

Keywords: demographic inference, background selection, MSMC, *fastsimcoal2*, approximate

Bayesian computation (ABC)

INTRODUCTION

The characterization of population history is a central goal of genomic analysis - with applications spanning from anthropological to agricultural to clinical (see review by Beichman *et al.* 2018). Furthermore, use of an appropriate underlying demographic model provides a necessary null for assessing the impact of selective effects across the genome (*e.g.*, Teshima *et al.* 2006; Thornton and Jensen 2007; Jensen *et al.* 2019). Multiple strategies have been proposed for performing demographic inference, utilizing expectations related to levels of variation, the site frequency spectrum, linkage disequilibrium, and within- and between-population relatedness (*e.g.*, Gutenkunst *et al.* 2009; Li and Durbin 2011; Lukic and Hey 2012; Harris and Nielsen 2013; Excoffier *et al.* 2013; Bhaskar *et al.* 2015; Sheehan and Song 2016; Ragsdale and Gutenkunst 2017; Steinrücken *et al.* 2019; Kelleher *et al.* 2019; Speidel *et al.* 2019). Although many methods perform well when evaluated under the stated assumption of neutrality, it is in fact difficult in practice to assure that the sites utilized for empirical analysis neither experience direct selection nor the effects of selection at linked sites. For example, inference is often performed using intergenic sites, or 4-fold degenerate and intronic sites in more compact genomes. While there is evidence for weak direct selection in all of these categories in multiple organisms (*e.g.*, Haddrill *et al.* 2005; Chamary and Hurst 2005; Andolfatto 2005; Lynch 2007; Zeng and Charlesworth 2010; Choi and Aquadro 2016; Jackson *et al.* 2017), it is also clear that such sites near or in coding regions will also experience background selection (BGS; Charlesworth *et al.* 1993; Charlesworth 2013). These BGS effects are known to affect the local underlying effective population size, and alter both levels and patterns of variation and linkage disequilibrium (Charlesworth *et al.* 1993; Kaiser and Charlesworth 2009; O'Fallon *et al.* 2010; Charlesworth 2013; Nicolaisen and Desai 2013; Ewing and Jensen 2016; Johri *et al.* 2020).

However, commonly-used approaches for performing demographic inference that assume complete neutrality, including *fastsimcoal2* (Excoffier *et al.* 2013) and MSMC/PSMC (Li and Durbin 2011; Schiffels and Durbin 2014), have yet to be thoroughly evaluated in light of this likely-to-be-violated model assumption. Yet it is not entirely uninvestigated. Rather than investigating existing software, Ewing and Jensen (2016) implemented an approximate Bayesian (ABC) approach to quantify the impact of unaccounted for BGS effects, demonstrating that weak purifying selection can generate a skew towards rare alleles that would be mis-interpreted as population growth. Under certain scenarios, this resulted in a many-fold mis-inference of

population size change. However, the density of directly selected sites and the shape of the

distribution of fitness effects (DFE) - parameters expected to be of great importance - have yet to

be considered. Spanning the range of these potential parameter values is also important for

understanding the expected impact on any given empirical application. As one example, the

proportion of the genome experiencing direct selection can vary greatly between species, with

estimates ranging from ~3-8% in humans, to ~12% in rice, to 37-53% in *D. melanogaster*, to 47-

68% in *S. cerevisiae* (Siepel *et al.* 2005; Liang *et al.* 2018). Further, many organisms have highly

compact genomes, with ~88% of the *E. coli* genome (Blattner *et al.* 1997) and effectively all of

many virus genomes (*e.g.*, >95% of the SARS-CoV-2 genome; Wu *et al.* 2020).

While such estimates allow us to approximate the effects of BGS in model organisms in

which recombination and mutation rates are well known, there are many non-model organisms in

which it is difficult to predict these effects. Moreover, while the genome-wide mean of *B*, a

widely-used measure of BGS effects that measures the level of variability relative to neutral

expectation, can range from ~0.45 in *D. melanogaster* to ~0.94 in humans (Charlesworth 2013;

though see Pouyet *et al.* 2018), existing demographic inference approaches are similarly applied

across organisms without a consideration of this important source of differences in levels of bias.

Here, we examine the effects of DFE shape and functional density on two common demographic

inference approaches - the multiple sequentially Markovian coalescent (MSMC) and

fastsimcoal2. Finally, we propose an extension within the approximate Bayesian computation

(ABC) framework to address this issue, treating the DFE as a nuisance parameter. Simulating

examples consistent with the structure of the *Drosophila* genome, we demonstrate greatly

improved demographic inference even when using directly selected sites alone.

METHODS

Simulations: Simulations were performed using SLiM 3.1 (Haller and Messer 2019) with 10

replicates per evolutionary scenario. For every replicate, 22 chromosomes of 150Mb each were

simulated, totaling ~3 Gb of information per individual genome (similar to the amount of

information in a human genome). Within each chromosome, 3 different types of regions were

simulated, representing non-coding intergenic, intronic, and exonic regions. Based on the NCBI

RefSeq human genome annotation downloaded from the UCSC genome browser

(http://genome.ucsc.edu/; Kent *et al.* 2002) for hg19, mean values of exon sizes and intron

numbers per gene were calculated. To represent mean values for the human genome (Lander *et*

al. 2001), each gene was comprised of 8 exons and 7 introns, and exon lengths were fixed at 350

bp. By varying the lengths of the intergenic and intronic regions, three different genomic

configurations with varying densities of functional elements were simulated and compared - with

5%, 10% and 20% of the genome being under direct selection - hereafter referred to as genome5,

genome10, and genome20, respectively. Genome5 was comprised of introns of 3000 bp and

intergenic sequence of 31000 bp, genome10 of introns of 1500 bp and intergenic sequence of

15750 bp, while genome20 was comprised of introns of 600 bp and intergenic sequence of 6300

bp. The total chromosome sizes of these genomes were approximately 150 Mb (150,018,599 bp,

150,029,949 bp, 150,003,699 bp in genome5, 10, and 20, respectively). In order to be

conservative and favorable with regards to the performance of existing demographic estimators,

intronic and intergenic regions were assumed to be neutral.

Recombination and mutation rates were assumed to be equal at 1×10^{-8} /site / generation.

Neither crossover interference (see discussion of Campos and Charlesworth 2019) nor gene

conversion were here considered. Exonic regions in the genomes experienced direct purifying

selection given by a discrete DFE comprised of 4 fixed classes: f_0 in which $0 \leq |2N_e s| < 1$ (*i.e.*,

effectively neutral), f_1 in which $1 \leq |2N_e s| < 10$ (*i.e.*, weakly deleterious), f_2 in which $10 \leq |2N_e s| <$

100 (*i.e.*, moderately deleterious), and f_3 in which $100 \leq |2N_e s| < 2N_e$ (*i.e.*, strongly deleterious),

where N_e is the effective population size and s is the decrease in fitness of the mutant

homozygote relative to wild-type. Within each bin, the distribution of s was assumed to be

uniform. All mutations were assumed to be semi-dominant. In all cases, the N_e corresponding to

the DFE refers to the ancestral effective population size. Six different types of DFE were

simulated: (1) a DFE skewed largely towards mildly deleterious mutations given by f_0=0.1,

f_1=0.7, f_2=0.1, f_3=0.1, (2) a DFE skewed towards moderately deleterious mutations, f_0=0.1,

f_1=0.1, f_2=0.7, f_3=0.1, (3) a DFE skewed towards strongly deleterious mutations f_0=0.1, f_1=0.1,

f_2=0.1, f_3=0.7, (4) a DFE with equal proportions of all mutations given by f_0=0.25, f_1=0.25,

f_2=0.25, f_3=0.25, (5) a DFE with equal proportions of neutral and strongly deleterious mutations

given by f_0=0.5, f_1=0.0, f_2=0.0, f_3=0.5, and (6) a DFE with a majority of neutral mutations and a

minority of strongly deleterious mutations given by f_0=0.7, f_1=0.0, f_2=0.0, f_3=0.3. In cases with

demographic non-equilibrium, the DFE shape was scaled with respect to the ancestral population

size.

Three different demographic models were tested for each of these DFEs: 1) demographic

equilibrium, 2) recent exponential 30-fold growth resembling that estimated for the human CEU

population (Gutenkunst *et al.* 2009), and 3) ~6-fold instantaneous decline, resembling the out-of-

Africa bottleneck in humans (Gutenkunst *et al.* 2009). The precise parameters of these three

scenarios are given in Supp. Table 1 and are also displayed in Figure 1a.

Running MSMC: In order to quantify the effect of purifying selection on demographic

inference, we used the entire chromosomes generated by SLiM to generate input files for

MSMC. For comparison, and to quantify the effect of BGS alone on demographic inference, we

masked the exonic regions to generate input files. For all parameters, MSMC was performed on

a single diploid genome, as those results were the most accurate (Supp Figure 1). Input files were

made using the script ms2multihetsep.py provided in the msmc-tools-Repository downloaded

from https://github.com/stschiff/msmc-tools. MSMC1 and 2 were run as follows:

msmc_1.1.0_linux64bit -t 5 -r 1.0 -o output_genomeID input_chr1.tab input_chr2.tab ...

input_chr22.tab

msmc2_linux64bit -r 1.0 -t 5 -o output_genomeID input_chr1.tab input_chr2.tab ...

input_chr22.tab

Running Fastsimcoal2: In order to minimize the effects of linkage disequilibrium, only SNPs

separated by 5 kb were used for inference, following Excoffier *et al.* (2013). Inference was also

performed by including all SNPs in order to assess the impact of violating the assumption of no

linkage disequilibrium on demographic inference. Site frequency spectra (SFS) were obtained for

both sets of SNPs for all 10 replicates of every combination of demographic history and DFE.

SNPs from all 22 chromosomes were pooled together to calculate the SFS. *Fastsimcoal2* was

used to fit each SFS to 4 distinct models: (a) equilibrium, which estimates only a single

population size parameter (N); (b) instantaneous size change (decline/growth), which fits 3

parameters - ancestral population size (N_{anc}), current population size (N_{cur}), and time of change

(T); (c) exponential size change (decline/growth), which also estimates 3 parameters - N_{anc}, N_{cur}

and T; and (d) an instantaneous bottleneck model with 3 parameters – N_{anc}, intensity, and time of

bottleneck. Model selection was performed as recommended by Excoffier *et al.* (2013). For each

demographic model, the maximum of maximum likelihoods from all replicates was used to

calculate the Akaike Information Criterion (AIC) = 2 × number of parameters – 2 ×

ln(likelihood). The relative likelihoods (Akaike's weight of evidence) in favor of the i^{th} model

was then calculated by:

$$w(i) = \frac{e^{-0.5\Delta_i}}{\sum_{j=1}^{4} e^{-0.5\Delta_j}}$$

where $\Delta_i = AIC_i - AIC_{min}$. The model with the highest relative likelihood was selected as the

best model, and the parameters estimated using that model were used to plot the final inferred

demography.

Performing inference by approximate Bayesian computation (ABC): ABC was performed

using the R package "abc" (Csilléry *et al.* 2010) and non-linear regression aided by a neural net

(used with default parameters as provided by the package) was used to correct for the

relationship between parameters and statistics. To infer posterior estimates, a tolerance of 0.1

was applied (*i.e.*, 10% of the total number of simulations were accepted by ABC to estimate the

posterior probability of each parameter). The weighted medians of the posterior estimates for

each parameter were used as point estimates. ABC inference was performed under two

conditions: (1) complete neutrality, and (2) in the presence of direct purifying selection. In both

cases only 2 parameters were inferred - ancestral (N_{anc}) and current (N_{cur}) population sizes.

However, in scenario 2, the shape of the DFE was also varied. Specifically, the parameters $f_0, f_1,$

$f_2,$ and f_3 were treated as nuisance parameters and were sampled such that $0 \leq f_i \leq 1,$ and $\Sigma_i f_i = 1,$

for $i = 0$ to 3. In addition, in order to limit the computational complexity involved in the ABC

framework, values of f_i were restricted to multiples of 0.05 (*i.e.*, $f_i \in \{0.0, 0.05, 0.10, \ldots, 0.95,$

$1.0\} \,\forall\, i$), which allowed us to sample 1,771 different DFE realizations. Simulations were

performed with functional genomic regions, and the demographic model was characterized by 1-

epoch changes in which the population either grows or declines exponentially from ancestral to

current size, beginning at a fixed time in the past. For the purpose of illustration, and for

comparison with the human-like parameter set above, parameters for ABC testing were selected

to resemble those of *D. melanogaster* African populations. Priors on ancestral and current

population sizes were drawn from a uniform distribution between 10^5-10^7 diploid individuals,

while the time of change was fixed at 10^6 ($\sim N_e$) generations. In order to simulate functional

regions, 94 single-exon genes, as described in Johri *et al*. (2020) and provided in

https://github.com/paruljohri/BGS_Demography_DFE/blob/master/DPGP3_data.zip, were

simulated with recombination rates specific to those exons (https://petrov.stanford.edu/cgi-

bin/recombination-rates_updateR5.pl) (Fiston-Lavier *et al.* 2010; Comeron *et al.* 2012).

Mutation rates were assumed to be fixed at 3×10^{-9} per site per generation (Keightley *et al.* 2009,

2014). All of these parameters were scaled by the factor 320 in order to decrease computational

time. The scaled population sizes thus ranged between ~300–30000 and were reported as scaled

values in the main text. One thousand replicate simulations were performed for every parameter

combination (N_{anc}, N_{cur}, f_0, f_1, f_2, f_3), 100 haploid genomes were randomly sampled without

replacement, and summary statistics were calculated using pylibseq 0.2.3 (Thornton 2003).

Means and variances (between replicates) of the following summary statistics were calculated

across the entire genomic region: nucleotide site diversity (π), Watterson's θ, Tajima's D, Fay

and Wu's H (both absolute and normalized), number of singletons, haplotype diversity, LD-

based statistics (r^2, D, D'), and divergence (*i.e.*, number of fixed mutations per site per

generation after the burn-in period). As opposed to the above examples, in this inference scheme

only exonic data (*i.e.*, directly selected sites) were utilized.

Data availability: The following data will be made publicly available at

https://github.com/paruljohri upon acceptance of the manuscript: (1) Scripts used to perform

simulations; (2) Input files used to run fastsimcoal2; (3) Scripts used for plotting; (4) Plotted

results of MSMC and *fastsimcoal2* for all models and scenarios tested in this work.

RESULTS and DISCUSSION

Effect of SNP number and genome size on inference under neutrality

The accuracy and performance of demographic inference was evaluated using two popular

methods, MSMC (Schiffels and Durbin 2014) and *fastsimcoal2* (Excoffier *et al.* 2013). In order

to assess performance, it was first necessary to determine how much genomic information is

required to make accurate inference when the assumptions of neutrality are met. Chromosomal

segments of varying sizes (1 Mb, 10 Mb, 50 Mb, 200 Mb, and 1 Gb) were simulated under
neutrality with 100 independent replicates each. Both *fastsimcoal2* and MSMC resulted in
incorrect inferences for all segments smaller than 1 Gb (Supp Figure 1, Supp Figure 2). In
addition, when 4 or 8 haploid genomes were used for inference, MSMC inferred a recent many-
fold growth for all segment sizes, but performed well when using 2 genomes for large segments
(Supp Figure 1). These results suggest exercising caution when performing inference on smaller
regions or genomes, and when using more than 1 diploid individual in MSMC. All further
analyses were restricted to characterizing demographic inference on data that roughly matched
the structure and size of the human genome - for every diploid individual, 22 chromosomes of
size 150 Mb each were simulated, which amounted to roughly 3 Gb of total sequence. Ten
independent replicates of each parameter combination were performed throughout and inference
was performed using 1 and 50 diploid individuals for MSMC and *fastsimcoal2,* respectively.
228
Effect of the strength of purifying selection on demographic inference
In order to test demographic inference in the presence of BGS, all 22 chromosomes were
simulated with exons of size 350 bp each, with varying sizes of introns and intergenic regions
(see Methods) to vary the fraction of the genome under selection. Because deleterious mutations
of differing severities result in background selection effects extending to variable distances,
demographic inference was evaluated for different possible DFE realizations - summarized by
four bins: $0 \leq |2N_e s| < 1$, $1 \leq |2N_e s| \leq 10$, $10 < |2N_e s| \leq 100$ and $100 < |2N_e s| \leq 2N_e$. Fitness effects
of mutations were uniformly distributed within each bin and the DFE shape was altered by
varying the proportion of mutations belonging to each fixed bin, given by f_0, f_1, f_2, and f_3,
respectively. Three highly skewed DFEs were initially used to assess the impact of the strength
of selection on demographic inference (with the remaining mutations equally distributed
amongst the other three classes): (1) a DFE in which 70% of mutations have weakly deleterious
fitness effects (*i.e.*, $f_1 = 0.7$), (2) a DFE in which 70% of mutations have moderately deleterious
fitness effects (*i.e.*, $f_2 = 0.7$), and (3) a DFE in which 70% of mutations have strongly deleterious
fitness effects (*i.e.*, $f_3 = 0.7$). In order to understand the effect of BGS, exonic sites were masked,
and only linked neutral intergenic sites were used for demographic inference. Three demographic
models were examined: (1) demographic equilibrium, (2) a 30-fold exponential growth,
mimicking the recent growth experienced by European human populations, and (3) ~6-fold

instantaneous decline, mimicking the out-of-Africa bottleneck in human populations (Figure 1a).
Although these models were parameterized using previous estimates of human demographic
history (Supp Table 1; Gutenkunst *et al.* 2009) for the purpose of illustration, these basic
demographic scenarios are applicable to many organisms.

Under demographic equilibrium, when 20% of the genome experiences direct selection
(which is masked) we observed an underestimate of the true population size, as well as the mis-
inference of recent population growth (Figure 1), a bias exacerbated in MSMC relative to
fastsimcoal2. Stronger growth is inferred when the effects of mutations are more deleterious.
Conversely, when the true demographic model is characterized by recent 30-fold growth,
demographic inference is accurate and performs equally well for both MSMC and *fastsimcoal2*,
with the exception of a slight under-estimation of the ancestral population size for all DFE types.
When the true model is population decline, weakly deleterious mutations alone did not affect
inference drastically and it was possible to recover the true model. However, moderately and
strongly deleterious mutations resulted in an underestimation of population sizes and an
inference of strong growth - to the extent that the decline model would be misinterpreted as a
growth model. We further tested the effect of BGS on demographic inference when changes in
population size were less severe, namely, when population growth and decline was only 2-fold -
with qualitatively similar results (Supp Figure 3).

Finally, given strong evidence that most organisms have a bi-modal DFE with a
significant proportion of strongly deleterious or lethal mutations (Sanjuán 2010; Kousathanas
and Keightley 2013; Jacquier *et al.* 2013; Bank *et al.* 2014), we investigated the effect of this
strongly deleterious class further. Thus, for comparison with the above, we simulated a rather
extreme case in which 30% or 50% of exonic mutations are strongly deleterious with fitness
effects uniformly sampled between $100 \leq |2N_{anc}s| \leq 2N_{anc}$, with the remaining mutations being
neutral. Similar to the above results, both equilibrium and decline models were falsely inferred as
growth, with an order of magnitude under-estimation of the true population size (Figure 2).

In sum, the neglect of BGS effects frequently results in the inference of population
growth, almost regardless of the true underlying model.

Effect of the density, and inclusion/exclusion, of directly selected sites on inference

Although we have shown that the presence of purifying selection biases demographic inference, the extent of mis-inference will necessarily depend on the fraction of the genome experiencing direct selection. We therefore compared models in which genomes contained 5%, 10%, and 20% functional regions. For this comparison, an equal proportion of mutations in each DFE bin were assumed (*i.e.*, $f_0 = f_1 = f_2 = f_3 = 0.25$). As before, when the true model is exponential growth, inference was unbiased, with a slight underestimation of ancestral population size when 20% of the genome is under selection (Figure 3). When the true model is population decline, it is inferred reasonably well if less than 10% of the genome experiences direct selection, and may be mis-inferred as growth with greater functional density, as shown in Figure 3. Similarly, the extent to which population size is under-estimated at demographic equilibrium increases with the fraction of the genome under selection. Finally, it is noteworthy that many such changes in population size that were falsely inferred were of a magnitude greater than 2-fold - suggesting the need for great caution when interpreting these plots from empirical data.

Importantly, the results presented do not significantly differ between inference performed while including directly selected sites (*i.e.*, no masking of functional regions) versus inference performed using linked neutral sites (*i.e.*, masking functional regions) as shown in Supp Figure 4. These results suggest that the exclusion of exonic sites, which is often assumed to provide a sufficiently neutral dataset to enable accurate demographic inference, is not necessarily a satisfactory solution unless gene density is low.

A potential solution: averaging across all possible DFEs

As shown above, demographic inference can be strongly affected by unaccounted for background selection effects, as well as by direct purifying selection. A potential solution is thus to correct for these effects when performing inference of population history. A promising first approach to estimating direct selection effects, DFE-alpha, takes a step-wise approach to infer demography using a presumed neutral class (synonymous sites); conditional on that demography, it then estimates parameters associated with the shape of the DFE (Keightley and Eyre-Walker 2007; Eyre-Walker and Keightley 2009; Schneider *et al.* 2011). However, this approach does not include the effects of selection at linked sites, which can result an over-estimate of population growth, as well as general mis-inference of the DFE (Johri *et al.* 2020).

Building on this idea, Johri *et al.* (2020) recently proposed an approach to include both

direct and background effects of purifying selection - simultaneously inferring the deleterious

DFE and demography. By utilizing the decay of BGS effects around functional regions, they

demonstrated high accuracy under the simple demographic models examined. Moreover, the

method makes no assumptions about the neutrality of synonymous sites, and can thus be used to

estimate selection acting on these sites, or in non-coding functional elements as well. However,

this computationally-intensive approach is specifically concerned with jointly inferring the DFE

with demography. As such, if an unbiased characterization of the population history is the sole

aim, such a procedure may be needlessly involved. We thus here examine the possibility of

rather treating the DFE as an unknown nuisance parameter, averaging across all possible DFE

shapes, in order to assess whether demographic inference may be improved simply by correcting

for these selection effects without inferring their underlying parameter values. This approach

thus utilizes functional (*i.e.*, directly selected) regions, a potential advantage in populations for

which only coding data may be available (*e.g.*, exome-capture data; see Jones and Good 2016),

or generally in organisms with high gene densities.

In order to illustrate this solution, a functional genomic element was simulated under

demographic equilibrium, 2-fold exponential population growth and 2-fold exponential

population decline with four different DFE shapes (as described previously, and shown in Figure

4). Inference was first performed assuming strict neutrality, and inferring a one-epoch size

change (thus estimating the ancestral (N_{anc}) and current population sizes (N_{cur})). As found in the

other inference approaches examined, population sizes were under-estimated and a false

inference of population growth was observed in almost all cases when selective effects are

ignored (Figure 4). Indeed, the reduction in diversity resulting from BGS (B) was found to differ

by demographic scenario, disproportionately amplifying mis-inference under certain models

(Figure 5). Next, the assumption of neutrality was relaxed, and mutations were simulated with

fitness effects characterized by a discrete DFE, with the same fitness classes given above ($f_0, f_1,$

f_2, f_3). Values for f_i were drawn from a uniform prior between 0 and 1 such that $\sum f_i = 1$. Note that

no assumptions were made about which sites in the genomic region were functionally important.

These directly selected sites were then used to infer demographic parameters. We found that by

varying the shape of the DFE, averaging across all realizations, and estimating only parameters

related to population history, highly accurate inference of modern and ancestral population size

is possible (Figure 4). These results demonstrate that even if the true DFE of a population is

unknown (as will always be the case empirically), it is possible to infer the demographic history

with reasonable accuracy by approximately accounting for these selective effects.

CONCLUSION

While commonly-used approaches to infer demography assume neutrality and

independence among segregating sites, empirically it is very difficult to verify those

assumptions. In addition, there is considerable evidence for the wide-spread effects of selection

on linked sites in many commonly studied organisms (Hernandez *et al.* 2011; Cutter and Payseur

2013; Elyashiv *et al.* 2016; Campos *et al.* 2017; Castellano *et al.* 2020). As such, we explored

how violations of this assumption may affect demographic inference, particularly with regard to

the underlying strength of purifying selection and the genomic density of directly selected sites.

Generally speaking, the neglect of these effects (*i.e.*, background selection) results in an

inference of population growth, with the severity of the growth model roughly scaling with

selection strength and density. Thus, when the true underlying model is in fact growth,

demographic mis-inference is not particularly severe; when the true underlying model is of

constant size or decline, the mis-inference can be extreme, with a many-fold under-estimation of

population size.

It is important to note that background selection effects extend to genomic distances in a

way that is positively related to the strength of purifying selection. For instance, strongly and

moderately deleterious mutations affect patterns of diversity at large genomic distances, whereas

mildly deleterious mutations primarily skew allele frequencies at adjacent sites. Thus, if

intergenic regions further away from exons are used to perform demographic inference, strongly

deleterious mutations are likely to predominantly bias inference. In contrast, if synonymous sites

are used to infer demographic history, mildly deleterious mutations will more predominantly

influence inference. Thus, as we here focused on relatively sparsely-coding genomes (resembling

human-like densities) and used intergenic sites to perform inference, moderately and strongly

deleterious mutations resulted in more severe demographic mis-inference. These results suggest

that demographic mis-inference will become stronger when such approaches are applied to

smaller, and more compact, genomes with higher gene densities.

Comparing the two methods investigated here, it appears that *fastsimcoal2* is less prone

to false fluctuations in population size caused by background selection. However, both methods

falsely infer growth under alternative models, which is exacerbated as the density of coding

regions increases. The correspondingly inferred times of population growth inferred by both

methods appear to be relatively randomly affected. Further, we observed little difference in

performance when using all SNPs relative to thinning SNPs to be separated by 5kb (Supp Figure

6), presumably because the models investigated do not generate strong LD. Overall, the degree

of mis-inference caused by a neglect of BGS is largely similar between the two methods.

However, it is noteworthy that even when all sites are strictly neutral, or only 5% of the

genome experiences direct selection, demographic equilibrium is mis-estimated as a series of

size changes within MSMC. The pattern of these erroneous size changes lend a characteristic

shape to the MSMC curve (*i.e.*, ancient decline and recent growth) which appears to resemble the

demographic history previously inferred for the Yoruban population (Schiffels and Durbin

2014), including the time at which changes in population size occurred (Supp Figure 5). Previous

work has demonstrated that this resulting model does not in fact well-explain the observed site-

frequency spectrum in the Yoruban population (Lapierre *et al.* 2017; Beichman *et al.* 2017). A

similar shape has also been inferred in the vervet subspecies (Warren *et al.* 2015; Figure 4), in

passenger pigeons (Hung *et al.* 2014; Figure 2), in elephants (Palkopoulou *et al.* 2018; Figure 4),

in Arabidopsis (Fulgione *et al.* 2018; Figure 3), and in grapevines (Zhou *et al.* 2017; Figure 2A).

Although under simulated neutrality the population size fluctuations are only ~1.2-fold, in most

empirical applications the fluctuations are of a somewhat larger magnitude (~ 2-fold in pigeons,

Arabidopsis, and grapevines). Nonetheless, this neutral, equilibrium performance of MSMC is

concerning, and adds to the other previously published cautions related to the interpretation of

MSMC results. For example, Mazet *et al.* (2016) and Chikhi *et al.* (2018) demonstrated that

under constant population size with hidden structure, inference may suggest false size change

(see also Orozco-terWengel 2016) . In addition, MSMC has been reported to falsely infer growth

prior to instantaneous bottlenecks (Bunnefeld *et al.* 2015). In addition, we observed that if

insufficient genomic data is utilized, or more than 2 haploid genomes are used to perform

inference, MSMC falsely infers recent growth of varying magnitudes (as previously observed by

Beichman *et al.* 2017).

In sum, we find that the effects of purifying and background selection result in similar
demographic mis-inference across approaches, and that masking functional sites does not yield
accurate parameter estimates. In order to side-step many of these difficulties, our proposed
approach of inferring demography by averaging selection effects across all possible DFE
realizations within an ABC framework appears to be highly promising. Utilizing only functional
regions, we found a great improvement in accuracy, without making any assumptions regarding
the true underlying shape of the DFE or the neutrality of particular classes of sites. As such, this
approach represents a more computationally-efficient avenue if only demographic parameters are
of interest, and ought to be particularly useful in the great majority of organisms in which
unlinked neutral sites either do not exist, or are difficult to identify and verify.
411
412

Acknowledgements

We would like to thank Brian Charlesworth and Susanne Pfeifer for helpful discussions related
to this project, and feedback on the manuscript. This research was conducted using resources
provided by Research Computing at Arizona State University
(http://www.researchcomputing.asu.edu) and the Open Science Grid, which is supported by the
National Science Foundation and the U.S. Department of Energy's Office of Science. This work
was funded by National Institutes of Health grant R01GM135899 to JDJ.

Figure 1: Inference of demography by MSMC (red lines; 10 replicates) and *fastsimcoal2* (blue lines; 10 replicates) with and without BGS, under demographic equilibrium (left column), 30-fold exponential growth (middle column), and ~6-fold instantaneous decline (right column). The true demographic model is depicted in black lines. (a) All genomic sites are strictly neutral. (b) Exonic sites experience purifying selection specified by a DFE comprised largely of weakly deleterious mutations ($f_0 = 0.1, f_1 = 0.7, f_2 = 0.1, f_3 = 0.1$). (c) Exonic sites experience purifying selection specified by a DFE comprised largely of moderately deleterious mutations ($f_0 = 0.1, f_1 = 0.1, f_2 = 0.7, f_3 = 0.1$). (d) Exonic sites experience purifying selection specified by a DFE comprised largely of strongly deleterious mutations ($f_0 = 0.1, f_1 = 0.1, f_2 = 0.1, f_3 = 0.7$). In this case, exons represent 20% of the genome. Here, exonic sites were masked/excluded when performing demographic inference - as such, this quantifies the effects of background selection alone.

Figure 2: Inference of demography by MSMC (red lines; 10 replicates) and *fastsimcoal2* (blue
lines; 10 replicates) in the presence of background selection generated by strongly deleterious
mutations. Here, directly selected sites comprised 20% of the genome and were masked when
performing demographic inference. (a) Exons experience purifying selection, with 30% of new
mutations being strongly deleterious, and the remaining being neutral. (b) Exons experience
purifying selection with 50% of new mutations being strongly deleterious, and the remaining
being neutral. The true demographic model is given as black lines.

Figure 3: Inference of demography by MSMC (red lines; 10 replicates) and *fastsimcoal2* (blue lines; 10 replicates) in the presence of background selection with varying proportions of the genome under selection, for demographic equilibrium (left column), exponential growth (middle column), and instantaneous decline (right column). Exonic sites were simulated with purifying selection given by the DFE: $f_0 = 0.25, f_1 = 0.25, f_2 = 0.25, f_3 = 025$ and were masked when performing inference. Directly selected sites comprise (a) 20% of the simulated genome, (b) 10% of the simulated genome, and (c) 5% of the simulated genome. The true demographic model is given by the black lines.

Figure 4: Comparison of inference of ancestral (N_{anc}) and current (N_{cur}) population sizes by
assuming neutrality *vs* by varying the DFE shape as a nuisance parameter, using an ABC
framework. Inference is shown for demographic equilibrium (left column), 2-fold exponential
growth (middle column), and 2-fold population decline (right column) for five separate DFE
shapes that define the extent of direct purifying selection acting on the genomic segment for
which demographic inference is performed: (a) neutrality ($f_0 = 1, f_1 = 0, f_2 = 0, f_3 = 0$), (b) weak
purifying selection ($f_0 = 0.1, f_1 = 0.7, f_2 = 0.1, f_3 = 0.1$), (c) moderately strong purifying selection
($f_0 = 0.1, f_1 = 0.1, f_2 = 0.7, f_3 = 0.1$), (d) strong purifying selection ($f_0 = 0.1, f_1 = 0.1, f_2 = 0.1, f_3 = 0.7$), and (e) a DFE in which all classes of mutations are equally present ($f_0 = f_1 = f_2 = f_3 = 0.25$).
In each, the horizontal lines give the true value (black for N_{anc}; and gray for N_{cur}) and the box-
plots give the estimated values. Black and gray boxes represent estimates when assuming
neutrality, while red boxes represent estimates when the DFE is treated as a nuisance parameter.

475
476
477

478
479
Figure 5: Reduction in nucleotide diversity (B) due to background selection under demographic
equilibrium (black bars), population growth (gray bars), and population decline (white bars),
under varying DFEs:
DFE 1: $f_0 = 0.1, f_1 = 0.7, f_2 = 0.1, f_3 = 0.1$;
DFE 2: $f_0 = 0.1, f_1 = 0.1, f_2 = 0.7, f_3 = 0.1$;
DFE 3: $f_0 = 0.1, f_1 = 0.1, f_2 = 0.1, f_3 = 0.7$;
DFE 4: $f_0 = f_1 = f_2 = f_3 = 0.25$;
DFE 5: $f_0 = 0.5, f_1 = 0.0, f_2 = 0.0, f_3 = 0.5$;
DFE 6: $f_0 = 0.7, f_1 = 0.0, f_2 = 0.0, f_3 = 0.3$.
Here, exonic sites comprise ~10% of the genome, roughly mimicking the density of the human
genome. BGS effects differ strongly by demographic model. Note that here $B = \pi(D) / \pi_0(D)$,
where $\pi(D)$ is the nucleotide diversity in neutral regions when exons experience purifying
selection under demographic model D (*i.e.*, with BGS), while $\pi_0(D)$ is the nucleotide diversity
under unlinked neutrality for the same demographic model D (*i.e.*, without BGS).

REFERENCES

Andolfatto P., 2005 Adaptive evolution of non-coding DNA in *Drosophila*. Nature 437: 1149–1152. https://doi.org/10.1038/nature04107

Bank C., G. B. Ewing, A. Ferrer-Admettla, M. Foll, and J. D. Jensen, 2014 Thinking too positive? Revisiting current methods of population genetic selection inference. Trends Genet. 30: 540–546. https://doi.org/10.1016/j.tig.2014.09.010

Beichman A. C., T. N. Phung, and K. E. Lohmueller, 2017 Comparison of single genome and allele frequency data reveals discordant demographic histories. G3 (Bethesda) 7: 3605–3620. https://doi.org/10.1534/g3.117.300259

Beichman A. C., E. Huerta-Sanchez, and K. E. Lohmueller, 2018 Using genomic data to infer historic population dynamics of nonmodel organisms. Annu. Rev. Ecol. Evol. Syst. 49: 433–456. https://doi.org/10.1146/annurev-ecolsys-110617-062431

Bhaskar A., Y. X. R. Wang, and Y. S. Song, 2015 Efficient inference of population size histories and locus-specific mutation rates from large-sample genomic variation data. Genome Res. 25: 268–279. https://doi.org/10.1101/gr.178756.114

Blattner F. R., G. Plunkett, C. A. Bloch, N. T. Perna, V. Burland, *et al.*, 1997 The complete genome sequence of *Escherichia coli* K-12. Science 277: 1453–1462. https://doi.org/10.1126/science.277.5331.1453

Bunnefeld L., L. A. F. Frantz, and K. Lohse, 2015 Inferring bottlenecks from genome-wide samples of short sequence blocks. Genetics 201: 1157–1169. https://doi.org/10.1534/genetics.115.179861

Campos J. L., L. Zhao, and B. Charlesworth, 2017 Estimating the parameters of background selection and selective sweeps in *Drosophila* in the presence of gene conversion. Proc. Natl. Acad. Sci. U.S.A. 114: E4762–E4771. https://doi.org/10.1073/pnas.1619434114

Campos J. L., and B. Charlesworth, 2019 The effects on neutral variability of recurrent selective sweeps and background selection. Genetics 212: 287–303. https://doi.org/10.1534/genetics.119.301951

Castellano D., A. Eyre-Walker, and K. Munch, 2020 Impact of mutation rate and selection at linked sites on DNA variation across the genomes of humans and other Homininae. Genome Biol. Evol. 12: 3550–3561. https://doi.org/10.1093/gbe/evz215

Chamary J., and L. D. Hurst, 2005 Evidence for selection on synonymous mutations affecting stability of mRNA secondary structure in mammals. Genome Biol. 6: R75. https://doi.org/10.1186/gb-2005-6-9-r75

Charlesworth B., M. T. Morgan, and D. Charlesworth, 1993 The effect of deleterious mutations on neutral molecular variation. Genetics 134: 1289–1303.

Charlesworth B., 2013 Background selection 20 years on. The Wilhelmine E. Key 2012
invitational lecture. J. Hered. 104: 161–171. https://doi.org/10.1093/jhered/ess136

Chikhi L., W. Rodríguez, S. Grusea, P. Santos, S. Boitard, *et al.*, 2018 The IICR (inverse
instantaneous coalescence rate) as a summary of genomic diversity: insights into
demographic inference and model choice. Heredity 120: 13–24.
https://doi.org/10.1038/s41437-017-0005-6

Choi J. Y., and C. F. Aquadro, 2016 Recent and long term selection across synonymous sites in
Drosophila ananassae. J. Mol. Evol. 83: 50–60. https://doi.org/10.1007/s00239-016-
9753-9

Comeron J. M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in
Drosophila melanogaster. PLoS Genet. 8: e1002905.
https://doi.org/10.1371/journal.pgen.1002905

Csilléry K., M. G. B. Blum, O. E. Gaggiotti, and O. François, 2010 Approximate Bayesian
Computation (ABC) in practice. Trends Ecol. Evol. 25: 410–418.
https://doi.org/10.1016/j.tree.2010.04.001

Cutter A. D., and B. A. Payseur, 2013 Genomic signatures of selection at linked sites: unifying
the disparity among species. Nat. Rev. Genet. 14: 262–274.
https://doi.org/10.1038/nrg3425

Elyashiv E., S. Sattath, T. T. Hu, A. Strutsovsky, G. McVicker, *et al.*, 2016 A genomic map of
the effects of linked selection in *Drosophila*. PLoS Genet. 12: e1006130.
https://doi.org/10.1371/journal.pgen.1006130

Ewing G. B., and J. D. Jensen, 2016 The consequences of not accounting for background
selection in demographic inference. Mol. Ecol. 25: 135–141.
https://doi.org/10.1111/mec.13390

Excoffier L., I. Dupanloup, E. Huerta-Sánchez, V. C. Sousa, and M. Foll, 2013 Robust
demographic inference from genomic and SNP data. PLoS Genet. 9: e1003905.
https://doi.org/10.1371/journal.pgen.1003905

Eyre-Walker A., and P. D. Keightley, 2009 Estimating the rate of adaptive molecular evolution
in the presence of slightly deleterious mutations and population size change. Mol. Biol.
Evol. 26: 2097–2108. https://doi.org/10.1093/molbev/msp119

Fiston-Lavier A.-S., N. D. Singh, M. Lipatov, and D. A. Petrov, 2010 *Drosophila melanogaster*
recombination rate calculator. Gene 463: 18–20.
https://doi.org/10.1016/j.gene.2010.04.015

Fulgione A., M. Koornneef, F. Roux, J. Hermisson, and A. M. Hancock, 2018 Madeiran
Arabidopsis thaliana reveals ancient long-range colonization and clarifies demography in
Eurasia. Mol. Biol. Evol. 35: 564–574. https://doi.org/10.1093/molbev/msx300

Gutenkunst R. N., R. D. Hernandez, S. H. Williamson, and C. D. Bustamante, 2009 Inferring the
581   joint demographic history of multiple populations from multidimensional SNP frequency
582   data. PLoS Genet. 5: e1000695. https://doi.org/10.1371/journal.pgen.1000695

583   Haddrill P. R., K. R. Thornton, B. Charlesworth, and P. Andolfatto, 2005 Multilocus patterns of
584   nucleotide variability and the demographic and selection history of *Drosophila*
585   *melanogaster* populations. Genome Res. 15: 790–799.
586   https://doi.org/10.1101/gr.3541005

587   Haller B. C., and P. W. Messer, 2019 SLiM 3: Forward genetic simulations beyond the Wright–
588   Fisher model. Mol. Biol. Evol. 36: 632–637. https://doi.org/10.1093/molbev/msy228

589   Harris K., and R. Nielsen, 2013 Inferring demographic history from a spectrum of shared
590   haplotype lengths. PLoS Genet. 9: e1003521.
591   https://doi.org/10.1371/journal.pgen.1003521

592   Hernandez R. D., J. L. Kelley, E. Elyashiv, S. C. Melton, A. Auton, *et al.*, 2011 Classic selective
593   sweeps were rare in recent human evolution. Science 331: 920–924.
594   https://doi.org/10.1126/science.1198878

595   Hung C.-M., P.-J. L. Shaner, R. M. Zink, W.-C. Liu, T.-C. Chu, *et al.*, 2014 Drastic population
596   fluctuations explain the rapid extinction of the passenger pigeon. Proc. Nat. Acad. Sci.
597   U.S.A. 111: 10636–10641. https://doi.org/10.1073/pnas.1401526111

598   Jackson B. C., J. L. Campos, P. R. Haddrill, B. Charlesworth, and K. Zeng, 2017 Variation in the
599   intensity of selection on codon bias over time causes contrasting patterns of base
600   composition evolution in *Drosophila*. Genome Biol. Evol. 9: 102–123.
601   https://doi.org/10.1093/gbe/evw291

602   Jacquier H., A. Birgy, H. L. Nagard, Y. Mechulam, E. Schmitt, *et al.*, 2013 Capturing the
603   mutational landscape of the beta-lactamase TEM-1. Proc. Natl. Acad. Sci. U.S.A. 110:
604   13067–13072. https://doi.org/10.1073/pnas.1215206110

605   Jensen J. D., B. A. Payseur, W. Stephan, C. F. Aquadro, M. Lynch, *et al.*, 2019 The importance
606   of the Neutral Theory in 1968 and 50 years on: A response to Kern and Hahn 2018.
607   Evolution 73: 111–114. https://doi.org/10.1111/evo.13650

608   Johri P., B. Charlesworth, and J. D. Jensen, 2020 Towards an evolutionarily appropriate null
609   model: jointly Inferring demography and purifying selection. Genetics.
610   https://doi.org/10.1534/genetics.119.303002

611   Jones M. R., and J. M. Good, 2016 Targeted capture in evolutionary and ecological genomics.
612   Mol. Ecol. 25: 185–202. https://doi.org/10.1111/mec.13304

613   Kaiser V. B., and B. Charlesworth, 2009 The effects of deleterious mutations on evolution in
614   non-recombining genomes. Trends Genet. 25: 9–12.
615   https://doi.org/10.1016/j.tig.2008.10.009

Keightley P. D., and A. Eyre-Walker, 2007 Joint inference of the distribution of fitness effects of
deleterious mutations and population demography based on nucleotide polymorphism
frequencies. Genetics 177: 2251–2261. https://doi.org/10.1534/genetics.107.080663

Keightley P. D., U. Trivedi, M. Thomson, F. Oliver, S. Kumar, *et al.*, 2009 Analysis of the
genome sequences of three *Drosophila melanogaster* spontaneous mutation accumulation
lines. Genome Res. 19: 1195–1201. https://doi.org/10.1101/gr.091231.109

Keightley P. D., R. W. Ness, D. L. Halligan, and P. R. Haddrill, 2014 Estimation of the
spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib
family. Genetics 196: 313–320. https://doi.org/10.1534/genetics.113.158758

Kelleher J., Y. Wong, A. W. Wohns, C. Fadil, P. K. Albers, *et al.*, 2019 Inferring whole-genome
histories in large population datasets. Nat. Genet. 51: 1330–1338.
https://doi.org/10.1038/s41588-019-0483-y

Kent W. J., C. W. Sugnet, T. S. Furey, K. M. Roskin, T. H. Pringle, *et al.*, 2002 The Human
Genome Browser at UCSC. Genome Res. 12: 996–1006.
https://doi.org/10.1101/gr.229102

Kousathanas A., and P. D. Keightley, 2013 A comparison of models to infer the distribution of
fitness effects of new mutations. Genetics 193: 1197–1208.
https://doi.org/10.1534/genetics.112.148023

Lander E. S., L. M. Linton, B. Birren, C. Nusbaum, M. C. Zody, *et al.*, 2001 Initial sequencing
and analysis of the human genome. Nature 409: 860–921.
https://doi.org/10.1038/35057062

Lapierre M., A. Lambert, and G. Achaz, 2017 Accuracy of demographic inferences from the site
frequency spectrum: the case of the Yoruba population. Genetics 206: 439–449.
https://doi.org/10.1534/genetics.116.192708

Li H., and R. Durbin, 2011 Inference of human population history from individual whole-
genome sequences. Nature 475: 493–496. https://doi.org/10.1038/nature10231

Liang P., H. S. A. Saqib, X. Zhang, L. Zhang, and H. Tang, 2018 Single-base resolution map of
evolutionary constraints and annotation of conserved elements across major grass
genomes. Genome Biol. Evol. 10: 473–488. https://doi.org/10.1093/gbe/evy006

Lukic S., and J. Hey, 2012 Demographic inference using spectral methods on SNP data, with an
analysis of the human out-of-Africa expansion. Genetics 192: 619–639.
https://doi.org/10.1534/genetics.112.141846

Lynch M., 2007 *The Origins of Genome Architecture*. Sinauer Associates, Sunderland,
Massachusetts.

Mazet O., W. Rodríguez, S. Grusea, S. Boitard, and L. Chikhi, 2016 On the importance of being
structured: instantaneous coalescence rates and human evolution--lessons for ancestral

population size inference? Heredity (Edinb) 116: 362–371.
https://doi.org/10.1038/hdy.2015.104

Nicolaisen L. E., and M. M. Desai, 2013 Distortions in genealogies due to purifying selection
and recombination. Genetics 195: 221–230. https://doi.org/10.1534/genetics.113.152983

O'Fallon B. D., J. Seger, and F. R. Adler, 2010 A continuous-state coalescent and the impact of
weak selection on the structure of gene genealogies. Mol. Biol. Evol. 27: 1162–1172.
https://doi.org/10.1093/molbev/msq006

Orozco-terWengel P., 2016 The devil is in the details: the effect of population structure on
demographic inference. Heredity (Edinb) 116: 349–350.
https://doi.org/10.1038/hdy.2016.9

Palkopoulou E., M. Lipson, S. Mallick, S. Nielsen, N. Rohland, *et al.*, 2018 A comprehensive
genomic history of extinct and living elephants. Proc. Nat. Acad. Sci. U.S.A. 115:
E2566–E2574. https://doi.org/10.1073/pnas.1720554115

Pouyet F., S. Aeschbacher, A. Thiéry, and L. Excoffier, 2018 Background selection and biased
gene conversion affect more than 95% of the human genome and bias demographic
inferences, (K. Veeramah, P. J. Wittkopp, and I. Gronau, Eds.). eLife 7: e36317.
https://doi.org/10.7554/eLife.36317

Ragsdale A. P., and R. N. Gutenkunst, 2017 Inferring demographic history using two-locus
statistics. Genetics 206: 1037–1048. https://doi.org/10.1534/genetics.117.201251

Sanjuán R., 2010 Mutational fitness effects in RNA and single-stranded DNA viruses: common
patterns revealed by site-directed mutagenesis studies. Philos. Trans. R. Soc. Lond., B,
Biol. Sci. 365: 1975–1982. https://doi.org/10.1098/rstb.2010.0063

Schiffels S., and R. Durbin, 2014 Inferring human population size and separation history from
multiple genome sequences. Nat. Genet. 46: 919–925. https://doi.org/10.1038/ng.3015

Schneider A., B. Charlesworth, A. Eyre-Walker, and P. D. Keightley, 2011 A method for
inferring the rate of occurrence and fitness effects of advantageous mutations. Genetics
189: 1427–1437. https://doi.org/10.1534/genetics.111.131730

Sheehan S., and Y. S. Song, 2016 Deep learning for population genetic inference. PLoS Comput.
Biol. 12: e1004845. https://doi.org/10.1371/journal.pcbi.1004845

Siepel A., G. Bejerano, J. S. Pedersen, A. S. Hinrichs, M. Hou, *et al.*, 2005 Evolutionarily
conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res. 15:
1034–1050. https://doi.org/10.1101/gr.3715005

Speidel L., M. Forest, S. Shi, and S. R. Myers, 2019 A method for genome-wide genealogy
estimation for thousands of samples. Nat. Genet. 51: 1321–1329.
https://doi.org/10.1038/s41588-019-0484-x

687 Steinrücken M., J. Kamm, J. P. Spence, and Y. S. Song, 2019 Inference of complex population
688  histories using whole-genome sequences from multiple populations. Proc. Natl. Acad.
689  Sci. U.S.A. 116: 17115–17120. https://doi.org/10.1073/pnas.1905060116

690 Teshima K. M., G. Coop, and M. Przeworski, 2006 How reliable are empirical genomic scans for
691  selective sweeps? Genome Res 16: 702–712. https://doi.org/10.1101/gr.5105206

692 Thornton K., 2003 Libsequence: a C++ class library for evolutionary genetic analysis.
693  Bioinformatics 19: 2325–2327. https://doi.org/10.1093/bioinformatics/btg316

694 Thornton K. R., and J. D. Jensen, 2007 Controlling the false-positive rate in multilocus genome
695  scans for selection. Genetics 175: 737–750. https://doi.org/10.1534/genetics.106.064642

696 Wang L., T. M. Beissinger, A. Lorant, C. Ross-Ibarra, J. Ross-Ibarra, *et al.*, 2017 The interplay
697  of demography and selection during maize domestication and expansion. Genome Biol.
698  18: 215. https://doi.org/10.1186/s13059-017-1346-4

699 Warren W. C., A. J. Jasinska, R. García-Pérez, H. Svardal, C. Tomlinson, *et al.*, 2015 The
700  genome of the vervet (*Chlorocebus aethiops sabaeus*). Genome Res. 25: 1921–1933.
701  https://doi.org/10.1101/gr.192922.115

702 Wu F., S. Zhao, B. Yu, Y.-M. Chen, W. Wang, *et al.*, 2020 A new coronavirus associated with
703  human respiratory disease in China. Nature 579: 265–269.
704  https://doi.org/10.1038/s41586-020-2008-3

705 Zeng K., and B. Charlesworth, 2010 Studying patterns of recent evolution at synonymous sites
706  and intronic sites in *Drosophila melanogaster*. J. Mol. Evol. 70: 116–128.
707  https://doi.org/10.1007/s00239-009-9314-6

708 Zhou Y., M. Massonnet, J. S. Sanjak, D. Cantu, and B. S. Gaut, 2017 Evolutionary genomics of
709  grape (*Vitis vinifera ssp. vinifera*) domestication. Proc. Nat. Acad. Sci. U.S.A. 114:
710  11715–11720. https://doi.org/10.1073/pnas.1709257114

Supp Figure 1: Performance of MSMC under neutrality when using 2, 4 and 8 individuals for inference and on varying chromosome sizes – (a) 1 Mb, (b) 10 Mb, (c) 50 Mb, (d) 200 Mb, (e) 1 Gb. Approximately 100 replicates of each are shown above. Grey lines indicate 0.5N and 2N. As shown, MSMC has a tendency to take a common shape, often falsely indicating recent population growth.

Supp Figure 2: Performance of *fastsimcoal2* under neutrality using 50 diploid individuals for inference when varying chromosome sizes – (a) 10 Mb, (b) 50 Mb, (c) 200 Mb, (d) 1 Gb. Approximately 100 replicates of each are shown above (blue lines) and the true model is shown in black.

Supp Figure 3: Inferred demography by MSMC (red lines) and *fastsimcoal2* (blue lines) in the presence of background selection, with the true DFE shown to the left of the panel, for 2-fold instantaneous decline (right column) and 2-fold exponential growth (left column). In this case, 20% of the genome experiences direct selection. The true demographic model is shown as black lines.

Supp Figure 4: Inference of demography by MSMC (red lines; 10 replicates) and *fastsimcoal2* (blue lines; 10 replicates) under demographic equilibrium (left column), 30-fold exponential growth (middle column), and ~6-fold instantaneous decline (right column) in the presence of direct purifying selection (*i.e.*, directly selected sites are not masked). The true demographic model is depicted in black lines. Exonic sites experience purifying selection specified by the following DFEs: (a) $f_0 = 0.1, f_1 = 0.7, f_2 = 0.1, f_3 = 0.1$, (b) $f_0 = 0.1, f_1 = 0.1, f_2 = 0.7, f_3 = 0.1$, (c) $f_0 = 0.1, f_1 = 0.1, f_2 = 0.1, f_3 = 0.7$, (d) $f_0 = 0.25, f_1 = 0.25, f_2 = 0.25, f_3 = 0.25$, (e) $f_0 = 0.7, f_1 = 0.0, f_2 = 0.0, f_3 = 0.3$, (f) $f_0 = 0.5, f_1 = 0.0, f_2 = 0.0, f_3 = 0.5$. In this case, 20% of the genome is composed of exons.

Supp Figure 5: Inference of demographic history by MSMC. Top panel / red line: simulations in which the true model is constant population size, and 50% of new mutations in exons are strongly deleterious with the remainder being neutral, where exons comprise of 5% of the genome. Bottom panel / green line: the empirical estimate of population history of the YRI population inferred with MSMC by Schiffels and Durbin (2014). The X-axis is in years (assuming a generation time of 30 years). Note that the Y-axes are on different scales, and the magnitude of change observed in the empirical data is considerably larger than that observed in the simulated data. Thus, this comparison is only meant to illustrate this common shape taken in MSMC plots (and see similar shapes in, for example, vervets (Warren *et al.* 2015; Figure 4) and passenger pigeons (Hung *et al.* 2014; Figure 2)).

Supp Figure 6: Demographic inference by *fastsimcoal2* when performed using all SNPs (right column) and when SNPs are thinned to be separated by 5 kb (left column). (a) Direct purifying selection (exonic sites are not masked) in 5% of the genome given by the DFE: $f_0 = f_1 = f_2 = f_3 = 0.25$. (b) Direct purifying selection (exonic sites are not masked) in 5% of the genome given by the DFE: $f_0 = 0.5, f_1 = 0.0, f_2 = 0.0, f_3 = 0.5$. (c) Direct purifying selection (exonic sites are not masked) in 10% of the genome given by the DFE: $f_0 = 0.1, f_1 = 0.7, f_2 = 0.1, f_3 = 0.1$. (d) Direct purifying selection (exonic sites are not masked) in 10% of the genome given by the DFE: $f_0 = 0.7, f_1 = 0.0, f_2 = 0.0, f_3 = 0.3$. (e) Background selection (*i.e.*, exonic sites are masked) in which 5% of the genome is exonic and given by the DFE: $f_0 = 0.5, f_1 = 0.0, f_2 = 0.0, f_3 = 0.5$. (f) Background selection (*i.e.*, exonic sites are masked) in which 20% of the genome is exonic and given by the DFE: $f_0 = 0.1, f_1 = 0.7, f_2 = 0.1, f_3 = 0.1$. (g) Background selection (*i.e.*, exonic sites are masked) in which 20% of the genome is exonic and given by the DFE: $f_0 = 0.1, f_1 = 0.1, f_2 = 0.7, f_3 = 0.1$. (h) Background selection (*i.e.*, exonic sites are masked) in which 20% of the genome is exonic and given by the DFE: $f_0 = f_1 = f_2 = f_3 = 0.25$.

Supp Table 1: Parameters underlying the human-like demographic models considered.

	Demographic models	Ancestral population size	Current population size	Time of change in generations
1	Equilibrium	10,000	10,000	NA
2	Exponential growth	1000	30,000	850
3	Instantaneous decline	12,300	2,100	4,750