<u>The Past, Present and Future of Elephant Landscapes in Asia</u>

Shermin de Silva[1,2*], Tiffany Wu[3,9], Philip Nyhus[3], Alison Thieme[1,10], Ashley Weaver[3], Josiah
Johnson[3], Jamie Wadey[4,11], Alexander Mossbrucker[5], Thinh Vu[6], Thy Neang[7], Becky Shu Chen[8],
Melissa Songer[1], Peter Leimgruber[1]

[1]Smithsonian Conservation Biology Institute, Conservation Ecology Center, Front Royal, VA,
USA

[2]Trunks & Leaves Inc., Newtonville, MA, USA

[3] Environmental Studies Program, Colby College, Waterville, ME, USA

[4] School of Environmental and Geographical Science, University of Nottingham Malaysia, Kuala
Lumpur, Malaysia

[5]Frankfurt Zoological Society, Jambi, Indonesia

[6]Department of Wildlife Management, Vietnam National University of Forestry, Hanoi, Vietnam

[7]Wild Earth Allies, Phnom Phenh, Cambodia

[8]Zoological Society of London, London, UK

[9]University of Michigan, Ann Arbor, MI, USA

[10]Maryland, College Park, MD, USA

[11]College of Science, Health, Engineering and Education, Murdoch University, Perth, WA,
Australia

*Corresponding author: Shermin@trunksnleaves.org; Trunks & Leaves Inc., 391 Walnut Street,
Unit 3, Newtonville MA 02460

Abstract

Habitat loss drives species' declines worldwide, but is seldom quantified over centennial

timescales. We constructed ecological niche models for Asian elephants based on land-use

change between 850-2015, and predictions under six different climate/socioeconomic scenarios

from 2015-2099. We find that over 64% of suitable natural habitat across diverse ecosystems

was lost over the past three centuries. Average patch size dropped 83% from approximately

99,000 km^2 to 16,000 km^2 and the area occupied by the largest patch decreased 83% from ~ 4

million km^2 (45% of area) to 54,000 km^2 (~7.5% of area). Over half of current elephant range

appears unsuitable. Habitat availability is predicted to decline further this century across all

scenarios. The most severe losses occur under RCP3.4-SSP4, representing mid-range emissions

but high regional inequities. We conclude that climate change mitigation measures must include

policies to ensure inter-regional socioeconomic equity to safeguard landscapes for elephants,

humans, and other species.

Keywords: Land-use change, ecological niche modelling, species distribution modelling, Asian

elephant, social-ecological systems, climate change scenarios

Running head: Elephant Landscapes

Introduction

Habitat loss and degradation are leading drivers of terrestrial biodiversity loss worldwide

[1–3]. An estimated three quarters of the Earth's land surface has been significantly altered by

human activities [4]. Historic reasons include conversion for cultivation and settlement, reflecting

both local and global socioeconomic drivers of land-use and land-cover (LULC) change [5,6].

Climate change is an additional contributor to species declines within the past century [4,6]. As a

result of these anthropogenic changes to climate and land-use, global forest extent has been

reduced by 32% relative to the pre-industrial period and ecological communities are estimated to

have lost over 20% of their biodiversity [4,6]. Yet it is not a given that efforts to address climate

change will also conserve species and habitats because these are distinct concerns [7]. Moreover,

climate change and LULC feed back on each other via economic systems as well as natural

processes such as carbon cycling [5]. These social-ecological systems impose inherent trade-offs

on efforts to mitigate climate change and conserve biodiversity [7,8]. Indeed, simultaneously

addressing both goals may be among the greatest and most urgent challenges to developing

sustainable societies [6].

Human activities can alter landscapes quickly, but long-term legacies may not be

immediately evident. Human-induced changes already restrict the ranges of many terrestrial

mammal species [9,10] but historical records on population abundance and distribution are often

limited for many taxa, complicating efforts to assess human impacts over longer periods. Though

LULC trends in recent decades may be inferred from satellite imagery [2,11], these fail to capture

processes operating over centuries. Nevertheless, longer historical perspectives are necessary to

appreciate the true magnitude of changes to threatened ecosystems. For example, historical

studies have influenced conservation policies related to remnant prairies in Oregon, wetlands in

Iowa, and forests in Germany, at times challenging traditional management practices [12].

However, a review of more than 200 historical ecology studies found that fewer than 12%

contained recommendations that "explicitly addressed ongoing or projected climate change" [12].

Moreover, although many studies attempt to define global priority regions for conservation [7,13],

they do not account for future changes in climate, socioeconomic conditions, and land-use. There

is a need for better integration of historical perspectives and present-day assessments with

forward-looking strategies and policies.

Land-use practices and climate change will continue to alter ecosystem structure and in

the future, resulting in changes to the availability of habitat for many species [1]. Wide-ranging

species cannot be conserved solely within existing or planned protected area networks [1]. The

survival of these species will depend not only on reducing proximate threats from illegal hunting

and sufficient protection of extant range, but on preparing for possible range shifts and

associated challenges under scenarios of global change. Typically, this is achieved through

ecological niche modelling (c.f. species distribution modelling) in which present-day species

occurrence data, together with environmental covariates, are used to infer possible occurrence or

suitable habitat at a different area or time [14–16]. Often, climatic variables form the basis of

predictive studies[1] but LULC features more directly constrain habitat availability. Moreover,

social and economic systems can be expected to evolve along with physical changes in the

climate and biosphere [17]. Therefore, biodiversity conservation and land-use planning efforts must

take into consideration not only possible global emissions outcomes, but also the corresponding

LULC and socioeconomic pathways taken to reach those outcomes.

Asia, Earth's most densely populated yet biodiverse continent, contains several hotspots

of species decline and threatened megafauna [9]. We model historic range in suitable habitat over

the past 300 years and possible shifts over the next 100 years for a widely-distributed endangered

mammal, the Asian elephant (*Elephas maximus*). Elephants are ecosystem engineers that

uniquely influence the structure of natural ecosystems [18] and also exemplify mutual challenges

for people and wildlife at frontiers of land-use change that manifest as human-elephant conflict

[19]. Occupying ecosystems ranging from rainforests to grasslands, this habitat generalist acts as a

surrogate for diverse ecoregions rather than particular biomes. This range is thought to represent

only a subset of past distributions, reflecting long-term contraction (Figure S1) [20]. However,

published depictions of past range are based on limited historical and anecdotal accounts

whereas elephants were probably never distributed uniformly over the area.

We base our predictions on the Land-Use Harmonization 2 (henceforth LUH2) data

products [21]. The LUH2 datasets provide historical reconstructions of land-use and management

between 850-2015, at annual increments. They also contain modelled scenarios from 2015-2100

representing combinations of a subset of Representative Concentration Pathways (RCPs)[22] and

Shared Socioeconomic Pathways (SSPs)[17]. These datasets therefore link historic and future land-

use patterns with a consistent set of variables, and pair the socioeconomic narratives with

possible emissions outcomes (Table 1; supplementary text section 1). We used a subset of LUH2

variables to model changes in habitat for elephants over multiple centuries (Table S1).

Our aims are to: (1) quantify historic trends in suitable natural habitat and fragmentation

over the 13 range countries throughout Asia; (2) characterize the suitability of present-day range

and identify other areas containing potential habitat; and (3) predict how the extent and

fragmentation of natural habitat may change under the six different RCP-SSP future scenarios.

Under the simple assumption that climate trends are the critical determinant of habitat loss for

this species, one would expect a decrease in suitable habitat with increasing RCP values. If

socioeconomic considerations have more influence on land-use trends, there would be no such

pattern. We highlight areas of concern and opportunity among the different regions, alongside

the effect of climatic vs. socioeconomic pathways on elephant landscapes.

Results

Historical change

Primary forest was less important for the model than elevation, forested and non-forested

secondary lands, croplands, pastures and wood harvest activities (Table S2). The amount

'suitable' natural habitat was relatively stable until the 1700s but afterwards declined by

approximately 64% (over 38% within current range; Figures 1 & S2, Table 2). Average patch

size fell by 83%, from 99,000 km^2 to 16,000 km^2 and amount of area occupied by the largest

patch (LPI) also decreased 83% from 4 million km^2 (~45% of total area) to a little over 54,000

km^2 (~7.5% of total area). The landscape contagion index nearly doubled (Figure 1). Mainland

China, India, Bangladesh, Thailand, Vietnam and Sumatra each lost more than half their suitable

range (Table 2). China lost 94% and India lost 86%. Bhutan, Nepal and Sri Lanka lost more

suitable habitat inside than outside current elephant range (Table 2). Trends in Lao PDR and

Malaysia showed a net gain, but not necessarily in areas within the current range (Table 2 and

Figure S2). Borneo appears to have experienced habitat restructuring rather than decline.

Animations of the changes between 1700 and 2015 are provided in supplementary videos

SV1and SV2.

Present-day suitability vs. distribution

Just over 48% of the current range was classified as suitable for elephants by 2015

(Figures S2 & S3, Table S3). India has the largest proportion, but only about a third was

classified as suitable by the year 2015. Sri Lanka and Malaysian Borneo appear to have

populations that are more than twice what would be expected relative to their share of current

range, with around 63% of the range in Sri Lanka and 95% of that in Borneo qualifying as

suitable (Table S3). On the other hand, Lao PDR, Thailand and Myanmar have much lower

populations than expected based on their share of current range, despite approximately 79%,

60% and 51% of these ranges respectively being suitable. Most remaining range in Vietnam

(98%) and Indonesian Borneo (100%) is suitable but extremely small, together accounting for

just 0.27% of the total.

Future scenarios

Suitable habitat is predicted to decline further by 2100 across all scenarios, but patterns

vary by country and scenario (Figures 2, S4-S6 and supplementary video SV3). Most countries

are predicted to lose habitat across the majority of scenarios, with India predicted to lose the

greatest in absolute and relative terms (Figures 2 & S4). However, peninsular Malaysia may gain

habitat within the current range for all cases. The Bornean portion of the range shows potential

for increase in three of six scenarios. Yunnan province, China, exhibits potential habitat gains in

five out of six scenarios, three of these more than doubling (Figures 2 & S4).

Scenarios representing lower RCPs are not consistently associated with better outcomes

for these landscapes, even within countries (Figure 2 & 3, Figures S5 & S6, Table 3). Scenarios

2.6/1 and 8.5/5, with the lowest and highest amounts of radiative forcing, respectively, show

similar overall losses (2.6/1 yields -24.24% and 8.5/5 yields -22.41%). Likewise, scenarios 4.5/2

and 7/3 have nearly identical changes in extent and fragmentation (Table 3, Figures 3 & S7).

Strikingly, SSP4, which is associated with two different possible RCPs (3.4/4 and 6.0/4), shows

greater losses in the lower RCP scenario (-83.26%) than in the higher RCP scenario (-19.29%),

with 3.4/4 being the worst outcome in terms of both habitat loss and fragmentation among all six

scenarios (Table 3, Figures 3 & S7). It consistently shows a severe decrease in the amount of

suitable habitat accompanied by reductions in patch size and increased contagion (Figure 3 and

supplementary video SV3). Other scenarios do not appreciably differ in terms of changes to

metrics of habitat fragmentation and no single one appears to be better than others. Despite the

overall losses, all other scenarios allow for the possibility that while decreases occur within this

decade, habitats can be stabilized or partially recovered thereafter (Figures 3 & S7).

Discussion

Habitat loss due to anthropogenic activities is a global concern for numerous taxa [2,3].

While several studies have characterized losses of particular biomes in more recent decades [2,11],

fewer have examined long-term implications of land-use trends across diverse ecoregions. We

present the first quantitative assessment of the impact of long-term land-use and global change

scenarios on habitat availability for an endangered keystone species at a range-wide scale,

encompassing important regional biodiversity. We discuss the processes that may drive observed

results at country-level, then present overall outcomes in relation to possible climate change and

socioeconomic trajectories.

We find that after several centuries of relative stability, nearly two-thirds of habitat

within the 13 range countries declined within just the past 300 years. Less than half of the current

range is classified as being suitable (14% of the potential habitat; Tables 2 & S3). Patch size

reduction and fragmentation was far greater than is evident from recent decades alone. While it is

possible that some of these trends are due to modelling error, our results closely match a review

of 4,018 terrestrial mammals showing that on average 48.6% of a species' range could be

classified as "suitable" on the basis of their preferred habitats [3]. This likely reflects a lag between

land-use changes and population responses. Elephants may no longer be able to disperse into

some areas, nor persist where there has been heavy offtake in the past [23,24]. Conversely, in the

absence of overharvest, they may survive for some time in suboptimal habitat even if failing to

reproduce [25]. Previous studies suggest that habitat is being lost even inside protected areas [26].

Given that over half of the current range seems unsuitable, it may be instructive to relate levels

of human-elephant conflict to our results, in addition to more fine-scaled characterizations of

changes in habitat suitability. Moreover, our results reinforce that attempts to limit elephants to

protected areas or extant ranges is impractical.

Trends in South Asia are largely driven by India and Sri Lanka, which contain the largest

remaining wild populations (Table S3). Both countries were transformed by colonial road-

building, logging, and plantations, during which elephants and other wildlife were eradicated

from higher elevations and lowland rainforests [27]. Those remaining in India now roam

fragmented and largely unprotected landscapes. A study in central Assam found that HEC rates

dramatically increased in the 1980s, corresponding to a drop in forest cover below 30-40% [28].

Likewise, another study focused around the Nilgiri Biosphere reserve found that deforestation

was associated with increases in negative incidents with elephants [29]. Current range in Sri Lanka

more closely matches areas classified as suitable habitat in the year 1700 than 2015 ([30] and this

paper, Figure 1). A substantial portion includes agricultural mosaics with substantial human

activity [30], consistent with the result that anthropogenic activities and land-use types are more

influential on the model than primary forests (Table S2). Yet extirpations may not occur for

several generations due to the slow pace of demographic processes [31]. Present-day development

policies largely continue failing to consider the likelihood of human-elephant conflict at planning

stages, followed by efforts to drive out elephants from development zones. Governmental

subsidies and irrigation infrastructure promote cultivation of conflict-prone food crops such as

rice, fruits and vegetables in and around remaining habitat. Both India and Sri Lanka are

predicted to continue losing habitat in future scenarios (Figures 2 & S4), which will likely

exacerbate conflict and drive elephant population declines, unless these trends are reversed by re-

aligning development agendas with stated commitments to biodiversity conservation.

The movements of elephants and people convert ecological events into social, economic

and political issues. Range shifts by elephants can introduce challenges for human communities

that have little experience with elephants. For instance, while Bangladesh and Nepal can

potentially gain habitat under certain scenarios, Bhutan is predicted to lose habitat under all

scenarios. Yet Bhutan has a constitutional mandate to maintain at least 60% of its forest cover.

All three countries share borders with India, where habitat losses are predicted and thus could

experience influx of elephants. Conversely, the political displacement of people can also

drastically alter habitat. In 2018 there was rapid, large-scale disruption of an elephant corridor at

Cox Bazar between Bangladesh and Myanmar with the settlement of Rohingya refugees[32]. Such

events might more likely under narratives SSP3 and SSP4. Future changes in political conditions

as much as suitable habitat have important implications for trans-boundary management plans,

wherein governments should recognize linked ecological and sociopolitical concerns.

In Southeast Asia, the disappearance of highly suitable habitat in what is now central

Thailand is particularly striking (Figure 1), much of it occurring between 1950-1990

(supplementary videos SV1 and SV2). This therefore reflects not only colonial era timber

extraction and land-use conversions, but also more the recent "Green Revolution" of industrial

agriculture. Although expanses of forest remain in Thailand and Myanmar, both have lower

estimated elephant populations than expected based on their share of current range (Table S3,

Figure S4). This might be driven by habitat quality [24] or high rates of historic offtake for both the

timber and tourism industries [33,34]. As a result, captive elephants now outnumber wild elephants

in both countries [24,33]. Myanmar had been experiencing high rates of poaching for skins [35]. Yet

both countries are pursuing re-introduction efforts. Given the negative trends across all future

scenarios, such programs might consider where suitable habitats are likely to shift to in the future

(Figures S5 & S6).

Two of the most threatened elephant populations are found in Sumatra and Vietnam.

Sumatra is expected to suffer losses even under low-emission scenarios. Although 98% of the

current range in Vietnam is classified as suitable (Table S3), the extent of this range is extremely

small and all future scenarios show losses of potential habitat outside the current range. Both

cases require concerted efforts to recover habitat and re-connect isolated wildlife populations

through ecosystem-level management.

Malaysia stands out as a region for potential gains both for mainland and Bornean

populations, with potential connectivity[36]. In both cases, this may be at least in part because

selectively logged or secondary forests have more open canopies than primary forests,

encouraging the growth of understory forage. Our results support observations that secondary

vegetation is more influential in predicting habitat use than primary forest (Table S2) [37,38].

Alternatively, the model may be failing to distinguish plantation landscapes (oil palm, coffee,

tea, rubber etc.) from natural forests. Because LUH2 variables explicitly differentiate primary

forest from other types, the latter explanation is unlikely, but one might question how these

categories were defined (see study limitations, below). Yunnan province in China may also offer

opportunities for population recovery and re-wilding, if there is sufficient political desire as well

as capacity to safeguard against hunting and trafficking. Most importantly, range expansion in all

cases can only be possible alongside management programs designed to also safeguard human

communities and livelihoods.

Surprisingly, low-emissions scenarios do not necessarily yield better outcomes than

higher-emissions scenarios. This is highlighted by the similarity between scenarios 2.6/1 and

8.5/5, together with the contrast between scenarios 3.4/4 and 6.0/4 (Table 3). It is especially

noteworthy that 3.4/4 resulted in distinctly worse habitat loss and fragmentation patterns than all

others (Table 3, Figures 3 & S7). Policies achieving lower emissions targets strictly via advances

in developed countries and/or through heavier reliance on biofuels, which already contribute

heavily to habitat loss, will further endanger at-risk ecosystems. Southeast Asia contains regions

with some of the greatest overlaps in carbon stocks and biodiversity but they are also among the

most threatened by fuel sources such as oil palm[7,9]. Scenario 3.4/4 represents a case in which

heightened regional inequities result in "a failure of current policies for energy access, leading to

continued and increased use of biomass in the households of developing countries" [39]. This

underscores the complexity of the relationship between land-use practices and climate change [6],

where climate trends cannot be divorced from societal concerns. While it is undoubtedly

important to stabilize the climate by limiting global emissions, the pathway to doing so must

manage Earth's social-ecological system in a manner that does not further undermine vulnerable

species, ecosystems, or human populations.[8]

Study limitations

Habitat suitability models typically rely on relating the species of interest (i.e.

occurrence, behavior) to ecological covariates. They exclude at least two important

considerations. First, habitat characteristics offer a limited view of which areas may support a

particular species in the absence of demographic data. Locations that animals may perceive as

attractive but promote unsustainable rates of mortality are known as ecological traps.[40] By

limiting ourselves to "natural" landscapes, we have chosen to be conservative, excluding the

possibility that elephants might potentially flourish in some human-dominated landscapes. This

decision is underpinned by the second set of considerations: human perceptions and behavior. A

species cannot flourish on a landscape that is otherwise ideal if people exclude or suppress it.

Given the extent of loss in natural habitats in and around extant range, human-modified

landscapes (c.f. "working landscapes") could play a pivotal role for elephants as well as other

wildlife in the future as well as re-colonized potential range, if conflicts are minimized [19,41].

Conversely, although we sampled occurrences from "natural" landscapes, including

protected areas in which human activities are now limited, historically these biodiverse

landscapes may have also been shaped by people. Traditional cultivation and diverse

agroforestry practices may have been part of an unrecognized anthropological history [42].

Communities with legacies of sustainable resource management have been displaced under

unjust colonial land-use practices, including through the creation of protected areas[43].

Nevertheless, concepts such as "protected area" and "land ownership" are baked into the

scenarios themselves, e.g. by assuming probability of land-use conversion to be contingent on

protected status [17,44]. These paradigms leave little room to envision other varied and socially just

alternative (e.g., communal) forms of land management.

The datasets underlying our results (the LUH2 variables) are derived from a many other

complex models rather than direct measurements and observations, each with assumptions and

limitations. The numerous interacting components preclude directly relating results to specific

features. Indeed, the scenario developers themselves refrain from commenting on 'motivating

forces,' especially with regard to mid-range outcomes[17]. At the same time, many modelling

choices entail simplification for practicality. For instance, the SSPs as a set incorporate

feedbacks from climate change on biological but not economic systems [45]. One could easily

imagine such feedbacks (e.g., costly natural disasters). The longer the time horizon of a model,

the greater the uncertainty and potential for error propagation. This applies equally to the niche

model as much as its underlying predictors. Conditions in the past or future may represent

variable combinations that do not currently occur, thus transfers between time points may

involve extrapolation. For all these reasons, our outcomes are only examples of possible worlds

at relatively coarse spatial resolution over long time periods rather than precise depictions of

reality. Our intent is not to identify a particular "best" outcome around which policies can be

crafted, nor even to predict which specific sites are of conservation value at fine-scale (see also

[46]). Rather, because Asian elephants are ecologically important flagship species for conservation

management and policy with respect to diverse ecoregions, we expect these findings to prompt

further discussions and studies to understand the impact of land-use trends for future planning,

climate change mitigation strategies, and sustainability initiatives at national and global levels.

Methods

Elephant occurrence

Elephant occurrence locations were initially compiled from the Global Biodiversity

Information Facility (https://www.gbif.org/), Movebank (https://www.movebank.org/) and published

literature [47–49] as well as data contributed by the authors based on direct sightings, data logged

via tracking devices, and camera traps. Records were first checked visually for errors (e.g.,

occurrences outside natural continental range) then refined to include locations representing

natural habitat where the species could conceivably flourish. We included logged forest because

secondary or regenerating forest can support elephants with potentially little conflict with

humans. We excluded monoculture plantations and agricultural areas, because even if such

human-dominated landscapes appear capable of supporting elephants and may even seem

preferred,[41] populations on the whole may be subject to chronic negative impacts due to

accidents, conflicts with people, pollutants and gradual anthropogenic degradation of

landscapes[31,50]. Moreover, we cannot know whether elephants would have in fact used human-

dominated landscapes in the past when more natural areas were readily available, whereas we

consider it safe to assume that elephants would have used the latter. Lastly, our goal was to

quantify changes to natural biomes representing a wide array of biodiversity, without

confounding these with other types of landscapes in which elephants might today persist.

To minimize sampling bias that could result in model overfitting, we subsampled data to

cover the full distribution as widely as possible while eliminating redundant points located

within any particular landscape. For instance, only one randomly selected data point per

individual, per population, was used from any tracking dataset. Outliers from this set were

removed using Cooks' distance to eliminate locations that could represent potential errors. The

final dataset consisted of 91 occurrence points spanning the years 1996-2015 which served as

training data (Figure S1), where all data other than from GBIF and the literature were contributed

by the authors or individuals listed in acknowledgments. QGIS and Google Earth Pro were used

to visualize and consolidate these data.

Predictor variables

The LUH2 data products were downloaded from the University of Maryland at

http://luh.umd.edu/data.shtml (LUHv2h "baseline" scenario released October 14th 2016 and

LUHv2f future scenarios released December 12th 2017). They contain three types of variables

gridded at 0.25° x 0.25° (approximately 30km^2 at the equator): state variables describing the

land-use of a grid cell for a given year, transition variables describing the changes in a grid cell

from one year to the next, and management variables that describe agricultural applications such

as irrigation and fertilizer use, totaling 46 variables. Of these, we selected 20 variables

corresponding to all three types (Table S1), which were expected to be relevant to elephant

habitat use. Using ArcGIS 10 (ESRI 2017) we extracted each variable for each year between

1700-2015 from the historical reconstructions constituting LUHv2h, as well as each year

between 2015-2099 for six different future scenarios from LUHv2f. We used 2099 rather than

2100 because variables representing land-use transitions, such as wood harvest rates, were not

available for 2100. Note that although wood harvest rates are considered "transition" variables,

in fact they represent the percentage of a pixel harvested within a given year, resulting in land-

use conversions between years. We separately obtained elevation from the SRTM Digital

Elevation Model (Table S5).

The RCP-SSP pairings of the LUH2 datasets (Table 1) do not include the entire space of

combinations because not all outcomes are either possible or relevant. Scenarios with strong

barriers to mitigation and adaptation cannot achieve low radiative forcing targets whereas those

lesser barriers are only of interest with respect to the lower range of targets (see section 1 of the

supplementary text) [39]. These scenarios therefore only represent a cross-section of futures that

may occur when specific socioeconomic conditions and policy decisions place differential

constraints on efforts to mitigate or adapt to climate change (see supplementary text, section 1),

not a factorial design.

Data analysis

We limited the geographic extent of all analyses to the 13 range countries in which

elephants are currently found. We used MAXENT, a maximum entropy algorithm[51], to model

habitat suitability. Raster files were binarized in ArcGIS into suitable and unsuitable habitat with

a pixel size of approximately 20 km^2 at a cutoff threshold (discussed below) for the years

between 850 and 2000 at 25-year intervals, plus 2015. While we recognize that the suitability of

any habitat patch for a species may in reality fall along a gradient, given the spatial and temporal

scale of the datasets used in this study, we feel such refinements would necessitate subjective

judgments that are biologically questionable. Binarization is also a practical requirement for

quantifying habitat loss and fragmentation. However, as there is no commonly accepted

threshold type [52], to ensure that the specific choice of threshold did not affect the observed trends

in suitable habitat, we initially used three possible thresholds: 0.237, representing 'maximum test

sensitivity plus specificity,' 0.284 corresponding to 'maximum training sensitivity plus

specificity,' and 0.331 representing '10th percentile training presence'. Unless otherwise stated,

for subsequent analyses we show only results using the threshold of 0.284, where everything

below this threshold was classified as 'unsuitable' while everything above it was classified as

'suitable'. The binarized maps were re-projected using the WGS84 datum and an Albers Equal

Area Conic projection.

To establish whether a model using the LUH2 variables yields reasonable predictions of

habitat suitability for elephants, we first compared the result of using LUH2 variables for the

year 2000 to one based on other features expected to directly influence present-day elephant

distributions, based on knowledge of the species' ecology [53–55]. These include climate, terrain,

land-cover, and human and livestock densities (Table S5). The details of this comparison and

results are provided in the Supplementary Information (Section 3, Figures S8 & S9). The LUH2

model was then applied to all prior and future years to generate predictions of habitat suitability.

Analyses were performed using the 'dismo' package in R (R Core Team 2017).

Polygons representing elephant range were digitized from Hedges et al. 2008 [57], limited

to the category identified as "active confirmed". We refer to the area within these polygons as

"current range," avoiding the term "occupied" to avoid confusion with occupancy modelling, and

areas outside as "potential range". We compared the total area of the habitat classified as suitable

within and outside the current elephant range for each range country as well as Borneo and

Sumatra, quantifying changes over time. We included the entirety of Borneo because both

genetic studies and geological history allow for the possibility that elephants could have been

natively distributed throughout the island [58,59], and there are no present-day physical barriers to

dispersal on the island. We ranked each region based on the proportion of the elephant

population found within the region as well as the percentage of current range within the region,

and calculated the ratio of these ranks (Table S3). Country-level analyses were compared for all

countries except China, Indonesia, and Malaysia. The extent of historic change was assessed for

all of mainland China, where elephants originally occurred, but analyses of future scenarios were

restricted to Yunnan province, where elephants are now confined. For Indonesia we

distinguished between Sumatran and Bornean populations, in recognition of the distinct

subspecies in these two regions. Likewise, for Malaysia we distinguished between Peninsular

and Bornean populations.

For all regions we calculated the historic change in suitable habitat by subtracting the

total area classified as suitable in the year 2015 from that of the year 1700, as major changes take

place within this period (Figure 1). Likewise, we calculated the net change in projected suitable

area between 2015 and 2099 across all six scenarios (Figure 2). We additionally calculated

fragmentation statistics (Table S4) for all LUH2-based predictions (historical as well as future

scenarios) with the program FRAGSTATS (v.4.2) [60]. These metrics characterize changes to the

spatial configuration of habitat in addition to their absolute extent. We used a 'no sampling'

strategy and set the search radius and threshold distance, respectively, to 61 km (approximately

three pixel lengths) based on the movement and dispersal capacity of elephants[61,62].

Acknowledgments

The authors wish to thank Kate Jenks and Megan Baker for providing some of the elephant

occurrence data. AM & SdS received funding from the U.S. Fish & Wildlife Asian Elephant

Conservation Fund supporting collection of elephant occurrence data in Sri Lanka and Sumatra.

Preparation of this manuscript was partially supported by a James Smithson Fellowship to SdS

and funding from the Section in Ecology, Behavior and Evolution at the University of California,

San Diego. TW thanks the Colby College Environmental Studies Program, the Ralph J. Bunche

Scholarship and Russell F. Cole Fellowship.

Author contributions

422 SdS and PL conceived of the study. SdS, PL, and PN guided analyses and study design. SdS,

423 TW, AW, AT, and JJ performed analyses. SdS and PN drafted the manuscript. All other authors

424 contributed data to the study, revised and approved the manuscript.

Competing interests

427 The authors declare no competing interests.

References

1. Jantz, S. M. *et al.* Future habitat loss and extinctions driven by land-use change in biodiversity hotspots under four scenarios of climate-change mitigation. *Conserv. Biol.* **29**, 1122–1131 (2015).

2. Watson, J. E. M. *et al.* Catastrophic declines in wilderness areas undermine global environment targets. *Curr. Biol.* **26**, 2929–2934 (2016).

3. Crooks, K. R. *et al.* Quantification of habitat fragmentation reveals extinction risk in terrestrial mammals. *Proc. Natl. Acad. Sci.* **114**, 7635–7640 (2017).

4. IPBES. *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.* (IPBES Secretariat, 2019).

5. Lambin, E. F., Geist, H. J. & Lepers, E. Dynamics of land-use and land-cover change in tropical regions. *Annu. Rev. Environ. Resour.* **28**, 205–241 (2003).

6. Díaz, S. *et al.* Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science.* **366**, (2019).

7. Soto-Navarro, C. *et al.* Mapping co-benefits for carbon storage and biodiversity to inform conservation policy and action. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, (2020).

8. O'Neill, D. W., Fanning, A. L., Lamb, W. F. & Steinberger, J. K. A good life for all within planetary boundaries. *Nat. Sustain.* **1**, 88–95 (2018).

9. Ripple, W. J. *et al.* Collapse of the world's largest herbivores. *Sci. Adv.* **1**, e1400103 (2015).

10. Tucker, M. A. *et al.* Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science.* **359**, 466–469 (2018).

11. Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change. *Science (80-.).* **342**, 850–853 (2013).

12. Beller, E. E., McClenachan, L., Zavaleta, E. S. & Larsen, L. G. Past forward: Recommendations from historical ecology for ecosystem management. *Glob. Ecol. Conserv.* **21**, (2020).

13. Strassburg, B. B. N., Iribarrem, A. & Beyer, H. L. Global priority areas for ecosystem restoration. *Nature* (2020). doi:10.1038/s41586-020-2784-9

14. Rondinini, C. *et al.* Global habitat suitability models of terrestrial mammals. *Philos. Trans. R. Soc. B Biol. Sci.* **366**, 2633–2641 (2011).

15. Soberón, J. & Townsend Peterson, A. Monitoring biodiversity loss with primary species-occurrence data: toward national-level indicators for the 2010 target of the Convention on Biological Diversity. *AMBIO A J. Hum. Environ.* **38**, 29–34 (2009).

16. Sánchez-Cordero, V., Illoldi-Rangel, P., Linaje, M., Sarkar, S. & Peterson, A. T. Deforestation and extant distributions of Mexican endemic mammals. *Biol. Conserv.* **126**,

465–473 (2005).

17.   O'Neill, B. C. *et al.* The roads ahead: Narratives for shared socioeconomic pathways
describing world futures in the 21st century. *Glob. Environ. Chang.* **42**, 169–180 (2015).

18.   Campos-Arceiz, A. & Blake, S. Megagardeners of the forest – the role of elephants in
seed dispersal. *Acta Oecologica* **37**, 542–553 (2011).

19.   Nyhus, P. J. *Human-Wildlife Conflict and Coexistence. Annual Review of Environment*
and Resources **41**, (2016).

20.   Olivier, R. Distribution and status of the Asian Elephant. *Oryx* **14**, 379–424 (1978).

21.   Hurtt, G. C. *et al.* Harmonization of Global Land Use Change and Management for the
Period 850-2100 (LUH2) for CMIP6. *Earth System Grid Federation* (2020).

22.   Moss, R. H. *et al.* The next generation of scenarios for climate change research and
assessment. *Nature* **463**, 747–756 (2010).

23.   Calabrese, A. *et al.* Conservation status of Asian elephants: the influence of habitat and
governance. *Biodivers. Conserv.* **26**, 2067–2081 (2017).

24.   Songer, M., Aung, M., Allendorf, T. D., Calabrese, J. M. & Leimgruber, P. Drivers of
change in Myanmar's wild elephant distribution. *Trop. Conserv. Sci.* **9**, (2016).

25.   de Silva, S. & Leimgruber, P. Demographic tipping points as early indicators of
vulnerability for slow-breeding megafaunal populations. *Front. Ecol. Evol.* **7**, 1–13
(2019).

26.   Neupane, D., Kwon, Y., Risch, T. S. & Johnson, R. L. Changes in habitat suitability over
a two decade period before and after Asian elephant recolonization. *Glob. Ecol. Conserv.*
22, e01023 (2020).

27.   Skinner, T. *Fifty years in Ceylon: an autobiography*. (Tisara Press, 1891).

28.   Chartier, L., Zimmermann, A. & Ladle, R. J. Habitat loss and human-elephant conflict in
Assam, India: does a critical threshold exist? *Oryx* **45**, 528–533 (2011).

29.   Puyravaud, J.-P., Gubbi, S., Poornesha, H. C. & Davidar, P. Deforestation increases
frequency of incidents with elephants (*Elephas maximus*). *Trop. Conserv. Sci.* **12**,
194008291986595 (2019).

30.   Fernando, P., de Silva, M. K. C. R., Jayasinghe, L. K. A., Janaka, H. K. & Pastorini, J.
First country-wide survey of the Endangered Asian elephant: towards better conservation
and management in Sri Lanka. *Oryx* 1–10 (2019). doi:10.1017/s0030605318001254

31.   de Silva, S. & Leimgruber, P. Demographic tipping points as early indicators of
vulnerability for slow-breeding megafaunal populations. *Front. Ecol. Evol.* **7**, (2019).

32.   Goldsmith, B. Elephants face 'time bomb' in Bangladesh land clash with Rohingya
refugees. (2019).

33.   Jackson, J., Childs, D. Z., Mar, K. U., Htut, W. & Lummaa, V. Long-term trends in wild-
capture and population dynamics point to an uncertain future for captive elephants.

Proceedings. Biol. Sci. **286**, 20182810 (2019).

34. Nijman, V. *An assessment of the live elephant trade in Thailand*. (2014).

35. Sampson, C. *et al.* New elephant crisis in Asia — Early warning signs from Myanmar.
PLoS One **13**, e0194113 (2018).

36. de la Torre, J. A. *et al.* Using elephant movements to assess landscape connectivity under
Peninsular Malaysia's central forest spine land use policy. *Conserv. Sci. Pract.* **1**, e133
(2019).

37. Evans, L. J., Goossens, B., Davies, A. B., Reynolds, G. & Asner, G. P. Natural and
anthropogenic drivers of Bornean elephant movement strategies. *Glob. Ecol. Conserv.* **22**,
e00906 (2020).

38. Rood, E., Ganie, A. A. & Nijman, V. Using presence-only modelling to predict Asian
elephant habitat use in a tropical forest landscape: Implications for conservation. *Divers.*
Distrib. **16**, 975–984 (2010).

39. Riahi, K. *et al.* The Shared Socioeconomic Pathways and their energy, land use, and
greenhouse gas emissions implications: An overview. *Glob. Environ. Chang.* **42**, 153–168
(2017).

40. Robertson, B. & Hutto, R. A framework for understanding ecological traps and an
evaluation of existing evidence. *Ecology* **87**, 1075–1085 (2006).

41. de la Torre, J. A. *et al.* There will be conflict – agricultural landscapes are prime, rather
than marginal, habitats for Asian elephants. *Anim. Conserv.* **24**, 720–732 (2021).

42. Cairns, M. *Shifting Cultivation Policies: Balancing Environmental and Social*
Sustainability. (2017).

43. Rangarajan, M. & Shahabuddin, G. Displacement and relocation from protected areas:
Towards a biological and historical synthesis. *Conserv. Soc.* **4**, 359–378 (2006).

44. Hasegawa, T., Fujimori, S., Ito, A., Takahashi, K. & Masui, T. Global land-use allocation
model linked to an integrated assessment model. *Sci. Total Environ.* **580**, 787–796 (2017).

45. Calvin, K. *et al.* The SSP4: A world of deepening inequality. *Glob. Environ. Chang.* **42**,
284–296 (2017).

46. Di Marco, M., Watson, J. E. M., Possingham, H. P. & Venter, O. Limitations and trade-
offs in the use of species distribution maps for protected area planning. *J. Appl. Ecol.* **54**,
402–411 (2017).

47. Hedges, S. *et al.* Distribution, status, and conservation needs of Asian elephants (*Elephas*
maximus) in Lampung Province, Sumatra, Indonesia. *Biol. Conserv.* **124**, 35–48 (2005).

48. Santiapillai, C. & Dissanayake, S. R. B. Observations on elephants in the Maduru Oya
National Park Sri Lanka (*Mammalia, Elephantidae*). *Gajah* **20**, 9–20 (2001).

49. Marasinghe, M. S. L. R. P., Dayawansa, N. D. K. & Silva, R. P. Area suitability prediction
for conserving elephants: An application of likelihood ratio prediction model. *Trop. Agric.*

Res. **25**, (2015).

50.   Goswami, V. R., Vasudev, D. & Oli, M. K. The importance of conflict-induced mortality
for conservation planning in areas of human–elephant co-occurrence. *Biol. Conserv.* **176**,
191–198 (2014).

51.   Hijmans, R. J. & Elith, J. Species distribution modeling with R. (2013).

52.   Bean, W. T., Stafford, R. & Brashares, J. S. The effects of small sample size and sample
bias on threshold selection and accuracy assessment of species distribution models.
Ecography (Cop.). **35**, 250–258 (2012).

53.   Fernando, P. & Leimgruber, P. Asian elephants and seasonally dry forests. in *The Ecology*
and Conservation of Seasonally Dry Forests in Asia (eds. McShea, W. J., Davies, S. J.,
Phumpakphan, N. & Pattanavibool, A.) 151–163 (Smithsonian Insitution Scholarly Press,
2011).

54.   Matawa, F., Murwira, A. & Schmidt, K. S. Explaining elephant (*Loxodonta africana*) and
buffalo (*Syncerus caffer*) spatial distribution in the Zambezi Valley using maximum
entropy modelling. *Ecol. Modell.* **242**, 189–197 (2012).

55.   Pastorini, J. *et al.* A preliminary study on the impact of changing shifting cultivation
practices on dry season forage for Asian elephants in Sri Lanka. *Trop. Conserv. Sci.* **6**,
770–780 (2013).

56.   Team, R. D. C. R: A language and environment for statistical computing. (2017).

57.   Hedges, S., Fisher, K. & Rose, R. *Range-wide mapping workshop for Asian elephants*
(Elephas maximus). (2008).

58.   Sharma, R. *et al.* Genetic analyses favour an ancient and natural origin of elephants on
Borneo. *Sci. Reports2* **8**, 1–11 (2018).

59.   Husson, L., Boucher, F. C., Sarr, A. C., Sepulchre, P. & Cahyarini, S. Y. Evidence of
Sundaland's subsidence requires revisiting its biogeography. *J. Biogeogr.* **47**, 843–853
(2020).

60.   McGarigal, K., Cushman, S. A. & Ene, E. FRAGSTATS v4: Spatial pattern analysis
program for categorical and continuous maps. (2012).

61.   Moßbrucker, A. M. Modeling the fate of Sumatran elephants in Bukit Tigapuluh
Indonesia: research needs & implications for population management. *J. For. Sci.* **10**, 5–
18 (2016).

62.   Fernando, P. *et al.* Ranging behavior of the Asian elephant in Sri Lanka. *Mamm. Biol. -*
Zeitschrift für Säugetierkd. **73**, 2–13 (2008).

Table 1 – Scenario summaries. Representative Concentration Pathway (RCP) number denotes
the radiative forcing target in W/m^2 achieved by a given pathway at stabilization by the year
2100. The Shared Socioeconomic Pathway (SSP) number denotes the specific narrative storyline
that describes the socioeconomic processes operating in the modelled hypothetical societies.
Here we describe features that are salient to land-use. See references [17,21,39] as well as in-table
references and supplementary text section 1 for additional details.

Scenario	Description
RCP 2.6- SSP 1:	Warming stays below 2°C by the end of the century, with climate policy implemented through carbon pricing. This sustainable development paradigm is characterized by the adoption of sustainable technologies including carbon capture and storage together with biofuel production and reforestation. The extent of protected areas globally achieves the 17% Aichi target. Global population levels off at mid-century with concurrent adoption of less resource-intensive and environmentally responsible lifestyles such as the consumption of less meat (thus requiring less pasture). Likewise, improvements in agricultural productivity and irrigation efficiency reduce the agricultural footprint. Economic development is high, along with strong governance and institutions at both national and international levels.
RCP 3.4- SSP 4:	Mid-range radiative forcing target in a world of deep regional inequalities both within and among countries. High-income countries enjoy technological progress including agricultural innovations improving efficiency, concurrent with greater demands for energy and food. The high carbon price results in ~1200 million ha of land being allocated to the production of bioenergy, driving an 80% increase in the extent of cropland between 2010-2100. Climate and environmental policies in middle/high-income regions yield increases in forest cover. However, low-income countries stagnate. Consumption is driven by population growth, while agricultural productivity remains low and there is a heavy reliance on traditional biofuels, resulting in higher rates of deforestation.
RCP 4.5- SSP 2:	The RCP target is achieved by shifts toward more sustainable energy sources, food systems with reduced carbon footprint, and expansion of forests following an initial decline. SSP 2 is an intermediate between SSP 1 and SSP 3, where population growth slows and levels off and most countries are politically stable but development progress is uneven within and among countries.
RCP 6- SSP 4:	An alternative to RCP3.4-SSP4 based on the same framework but assuming less stringent climate change mitigation policies. Crop and pasture land increase by 14% and 9% respectively by 2100, along with a 3% increase in forest cover globally. The latter is largely situated in middle- and high-income countries.
RCP 7- SSP 3:	SSP3 represents weak global institutions, nationalistic political and economic policies reduce international cooperation. Wealthier countries experience less population growth, others experience higher growth. Countries strive to meet their food and energy needs domestically. Slow technological change in the energy sector with continued reliance on fossil fuels. Land-use allocations are made through profit/productivity-maximizing decisions with unprotected and uncultivated areas more likely to be converted to other uses.
RCP 8.5- SSP 5:	A fossil fuel-driven world with rapid economic development accompanied by investments in human capital, promoting more equitable distribution of resources, stronger institutions and slower population growth.

Table 2 – Change in suitable habitat area by region from years 1700-2015. Areas are in km^2, ordered from ranges that experienced the greatest loss to those with the greatest gain in total suitable area. "Suitable current area" refers to the current range, i.e. areas known to still contain elephants by the year 2000, whereas "Suitable potential area" refers to area that is outside the current range, where it is unknown whether elephants were ever present. The amount of suitable area within current range is 14%.

	Suitable Current Area (km^2)			Suitable Potential Area (km^2)			Total Suitable Area (km^2)		
	1700	2015	% change	1700	2015	% change	1700	2015	% change
Mainland China	1,986	135	-93.2	1,119,258	65,054	-94.2	1,087,183	65,189	-94.2
India	216,207	82,793	-61.7	1,439,320	145,561	-89.9	1,655,527	228,354	-86.2
Bangladesh	6,322	1,770	-72.0	37,725	10,634	-71.9	44,046	12,405	-71.8
Thailand	40,449	31,303	-22.6	439,964	127,028	-71.2	480,413	158,331	-67.0
Vietnam	523	515	-1.4	196,259	80,923	-58.8	196,781	81,439	-58.6
Indonesia (Sumatra)	45,252	27,507	-39.2	317,636	123,278	-61.2	362,888	150,785	-58.5
Indonesia (Borneo)	826	928	12.3	428,709	282,061	-34.2	429,535	282,989	-34.1
Myanmar	32,026	36,591	14.3	289,533	181,179	-37.4	321,559	217,770	-32.3
Cambodia	7,904	12,508	58.3	147,795	102,867	-30.4	155,699	115,374	-25.9
Nepal	12,086	4,750	-60.7	44,333	37,905	-14.5	56,419	42,655	-24.4
Sri Lanka	31,654	22,603	-28.6	24,097	19,622	-18.6	55,750	42,225	-24.3
Bhutan	2,033	1,148	-43.6	5,560	5,126	-7.8	7,593	6,273	-17.4
Lao PDR	16,507	17,716	7.3	148,922	159,843	7.33	165,429	177,558	7.3
Malaysia (Peninsular)	7,796	10,682	37.0	95,029	105,418	10.9	102,825	116,100	12.9
Malaysia (Borneo)	7,216	12,007	66.4	100,023	161,316	61.3	107,239	173,323	61.6
Totals	428,787	262,956	-38.7	4,834,163	1,607,815	-66.8	5,228,886	1,870,770	-64.2

Table 3 – Change in suitable habitat across scenarios from 2015-2099. There is no clear association between emission levels or socioeconomic narratives and changes in habitat suitability.

RCP/SSP	Change in suitable area within current range (%)	Change suitable area outside current range (%)	Change in total suitable area (%)
2.6/1	-39.2	-21.8	-24.2
3.4/4	-89.5	-82.3	-83.3
4.5/2	-31.1	-9.3	-12.3
6.0/4	-36.0	-16.6	-19.3
7.0/3	-32.0	-9.8	-12.8
8.5/5	-46.0	-18.6	-22.4

Figure 1 – Loss of suitable habitat from 850 – 2015. Masked areas (Hainan island and part of Pakistan) have been excluded from analyses, and for visual clarity all of China is not shown. Shaded area (Borneo) is outside the currently known historic range. A) Habitat suitability predicted on the basis of elevation and the Land-use Harmonization (LUH) variables from the year 2000. B) Binarized map where 1 (yellow) indicates "suitable" areas with values above 0.284 and 0 (blue) indicates "unsuitable" areas. C) Changes in the extent and spatial configuration of suitable habitat, where each curve corresponds to the given threshold value (see methods). Total Suitable Area is the sum of all suitable habitat across the range. Area-Weighted Mean Patch Size is the weighted average of patch sizes. The Largest Patch Index is the percentage of total area occupied by the largest patch. The Landscape Contagion Index can be thought of as a measure of homogeneity, with higher probabilities representing fewer, more clumped patches. See Table S4 for complete list of fragmentation measures.

Figure 2 - Net change in suitable habitat from 2015-2099 by country and scenario (RCP/SSP, ordered from lower to higher radiative forcings). The term "current" indicates area falling within the current range. "Suitable" indicates pixel values above the threshold of 0.284, whereas "unsuitable" indicates values below it. Increase in the "suitable current" category signifies area within the present range that was converted from unsuitable to suitable, whereas decrease signifies the reverse. These do not imply actual range expansion or contraction. "Suitable potential range" represents change in area outside the current range.

Figure 3 – Landscape fragmentation indicators across different scenarios. No single scenario yields a consistently better outcome across all measures, however scenario 3.4/4 is consistently worse than all others.