Genome-wide comparisons and extrapolations of AU-rich elements in Plants with *Homo sapiens*

[1]Deepshikha Satish, [*]Anil Panwar

Affiliation(s) and address(es) of the author(s):

[1]Deepshikha Satish, International Center for Genetic Engineering and Biotechnology, New Delhi, India 110067.(deepshikha@icgeb.res.in, 0000-0002-1313-5495)

[*]Anil Panwar, Department of Molecular Biology, Biotechnology and Bioinformatics, CCS Haryana Agricultural University, Hisar, Haryana, India 125004. (kumarniki2003@gmail.com, 0000-0003-2071-7888)

Abstract

The Adenylate-Uridylate Rich elements (AREs) are adenine and uracil abundant sequences, established in ephemeral mRNAs, principally present in the 3' untranslated region (3` UTR). AREs are widely accepted as a cause of high turnover of the mRNAs containing them (Bakheet, 2001; Barreau, 2005; Shaw and Kamen, 1986). The mammalian ARE-mRNAs primarily translate into nuclear transcription factors, oncoproteins, cytokines, and G-protein-coupled receptors which indicates their indispensable role in the regulation of gene transcription during cell growth and differentiation, and the immune response (Chen and Shyu, 1995; Wilson et al., 1999). The present study is an attempt to comprehensively analyze the eloquent presence of AU rich elements in genome, transcriptome and 3`UTR of three important plant species (*Arabidopsis thaliana, Oryza sativa and Zea mays*) and compare the statistics of putative ARE motifs in plants with *H.sapiens.* Statistical analysis of genome-wide putative AU-rich elements revealed the explicit presence of ARE motifs in plants. This is the first study that analyses the presence of Adenylate-Uridylate Rich elements (AREs) in three different plants which can be further validated through experiments.

Keywords:
AU rich elements, Deadenylation, mRNA degradation, 3`UTR, Cancers, Immunity.

Introduction

The management and control of mRNA stability is of utmost essentiality for adjusting mRNA abundance and for restricting mRNA translation to prevent detrimental effects caused by overexpression of proteins (Caput et al., 1986). Adenylate-Uridylate Rich elements (AREs) are nucleotide sequences exuberant in adenine and uracil, present within the 3' untranslated region (UTR) of ephemeral mRNAs and are liable for high turnover of the mRNA containing them (Bakheet, 2001; Barreau, 2005; Shaw and Kamen, 1986). ARE containing mRNAs encode proteins that regulate either cell growth or the impedance of an organism to external factors like micro-organisms, inflammatory stimuli and environmental factors. Such genes require meticulous control of their temporal and spatial expression patterns which is accomplished, in addition to transcriptional control, by a regulation of the mRNA translation and therefore the stability of the mRNA a. In mammalian cells, ARE-mRNA principally codes for nuclear transcription factors, oncoproteins, cytokines, and G-protein-coupled receptors which specify their indispensable role in the modulation of immune reaction, gene transcription during cell growth and differentiation (Chen and Shyu, 1995; Wilson et al., 1999).

So far there is no universal consensus for ARE motifs but the scientific experiments illuminate a list of motifs which are predicted to promote mRNA degradation (Table1). Significant characteristics of ARE elements include the occurrence of several AUUUA pentamers (short distance from each other is preferred) and theoretically the pentamers are implanted within a broader U-rich or AU-rich context. Further, the minimal functional ARE sequence is the nonamer UUAUUUA(U/A)(U/A) in recurrent instances. The three broad classes of AREs according to Chen and co-workers can be recognized based on the subsequent sequence characteristics: class I—one or more AUUUA pentamers within a U-rich sequence; class II—tandem repeats of AUUUA; and class III—U-rich sequences lacking AUUUA motifs (Chen and Shyu, 1995; Wilusz et al., 2001). On the other hand, according to Bakheet's method, AREs are classified into five groups based on the number of continuous repeats of the core sequence of AUUUA. Groups 1, 2, 3 and 4 possess 5, 4, 3, and 2 continuous repeats of AUUUA, respectively, whereas group 5 AREs possess only one core pattern in the transcript (Bakheet, 2001).

Table 1: Significant AU rich elements motifs

Motifs	Nomenclature
ATTT	Tetramer
ATTTA	Pentamer
TATTTAT	Heptamer
AATTTAA	Heptamer
WATTTAW	Heptamer
WWATTTAWW	Nanomer
WTTTW	Pentamer
WWTTTWW	Heptamer
WWWTTTWWW	Nanomer
WWWWTTTWWWW	Undecamer
TTTGTTT	Heptamer
GTTTG	Pentamer
AWTAAA	Hexamer

These AREs belonging to different classes becomes functional only after interacting with AU-binding proteins (AUBPs) which recruit enzymatic machinery responsible for mRNA degradation in a process described as ARE-mediated decay (Barreau, 2005). Several different proteins (e.g. HuA, HuB, HuC, HuD, HuR) bind to these elements and stabilise the mRNA while others (AUF1, TTP, BRF1, TIA-1, TIAR, and KSRP) destabilize the mRNA, miRNAs may also bind to some of them. These proteins act as adapters that recognize the ARE as a signal and couple the bound mRNAs to the general mRNA destabilization/stabilization machinery. The activity of the AU-element binding proteins is regulated, usually by phosphorylation but also by subcellular localization. In the cytoplasm, AUBPs (like TTP) bind to ARE (AUUUA) in the mRNA. The binding of AUBPs recruits either Dcp, which promotes decapping of mRNA, or deadenylase, which removes the poly (A) tail. After exposure of terminal mRNA, exonucleases act to degrade the mRNA in either the 5′–3′ direction using Xrn1 or in the 3′–5′ direction using the exosome. The preferable mode of action is mediated by rapid deadenylation (poly(A) tail decay) and subsequent degradation of the mRNAs (Figure 1)

A significant study illustrate that the consistency of AREs is high among the most unstable mRNAs (half-life <2 h), while its existence decreases among mRNAs with enhancing half-life (Lam et al., 2001). Generally, AREs cause mRNAs to be degraded rapidly and thereby suppress gene expression at the post-transcriptional level. Thus, stabilization of the ARE-mRNAs in some cases can cause a prolonged response that may subsequently lead to a diseased state. Problems with mRNA stability have been identified in viral genomes, cancer cells, and various diseases. Accumlating evidences

indicate that many of these problems arise because of faulty ARE function. For instance, the *c-fos* gene produces a transcription factor that is activated in several cancers and it lacks the ARE elements. *c-myc* gene, also responsible for producing transcription factors found in several cancers, has also been reported to lack the ARE elements. The *Cox-2* gene catalyses the production of prostaglandins—it overexpresses in several cancers and is stabilized by the binding of CUGBP2 RNA-binding protein to ARE. The association between turnover of defective mRNA and diseases underscores the significance of a deep understanding of the ARE-mediated decay mechanism (Eberhardt et al., 2007; Espel, 2005; Khabar, 2005). Yet, a compilation of mRNAs and their AREs along with half-life is still missing in available databases.

Figure 1: ARE mediated decay of mRNA

AU rich elements are not only one of the most prominent indicator for RNA stability in the mammalian cells (Chen and Shyu, 1995) but, ARE-bearing mRNAs are wide-spread in species ranging from *Homo sapiens* to the *Saccharomyces cerevisiae* (Duttagupta et al., 2003). This widespread presence of AU rich elements and their crucial roles inspire us to extrapolate their cross-kingdom roles. Sporadic reports indicating the presence and role of AREs in plants kingdom are already available but no genome-wide study of ARE in multiple plants species is available yet. This

scientific void of knowledge motivates us to predict the genome-wide presence of different ARE motifs in three important plant species at genome, transcriptome and 3` UTR level and compare them with human AU rich elements statistics. To understand the significance of AREs, computational analysis was carried out to identify the prevalence of AREs in *Arabidopsis thaliana, Oryza sativa* and *Zea mays* mRNAs and compared with *Homo sapiens*. The present study might open a new avenue to further establish the relationship between ARE types, positions and the transcript half-life in the plant kingdom.

Material and methods:

Genomic Data and Transcriptomic data for the present study were retrieved from the NCBI server (https://www.ncbi.nlm.nih.gov/) while 3` Untranslated Region Data was retrieved from http://utrdb.ba.itb.cnr.it/. Complete genomes of *A. thaliana, O. sativa, Z. mays and H. sapiens* were taken as initiation point. An in-house Perl script was used to deduce a total number of 14 different ARE motifs (Figure 2). Further complete transcriptomic data belonging to the above three plant species and humans were used and the number of different ARE motifs were searched in this data as well. As most of the functional ARE sites reported till now belong to untranslated regions of the genome, hence we tried to estimate the number of different ARE motifs in a comprehensive pool of 3`UTRs belonging to four species under study. All the results obtained were analyzed and materialized using in-house shell scripts.

Figure 2 : Graphical representation of number of putative ARE motifs predicted in genomes, transcriptomes and 3`UTRs. Ara=Arabidopsis thaliana, Osa= O.sativa, Zma= Z.mays, Hum= H.sapiens. Numbers on the X-axis indicate sizes of respective nucleotide data.

Results and discussion:

Bioinformatics analysis predicted that as many as 7% (≈2000) of the mRNAs in the human transcriptome harbour AREs (Bakheet, 2001). However, very few of these AREs have actually been validated experimentally.

Table 2: Number of ARE motifs present in genomes of respective plant species and H.sapiens

Motifs	A. thaliana	O. sativa	Z. mays	Homo sapiens
ATTT	1310215	3265887	13788857	29851282
ATTTA	290586	720476	2871616	6479456
TATTTAT	35291	82237	372208	843462
AATTTAA	31894	83051	237269	626299
WATTTAW	124024	309717	1069736	2707257
WWATTTAWW	61751	146470	419257	1335749
WTTTW	1198621	2810838	11041178	27325857
WWTTTWW	464010	1145879	3531366	11472366
WWWTTTWWW	214694	548704	1277612	5273269
WWWWTTTWWWW	110086	271302	555394	2709335
WWWWWTTTWWWWW	60330	139527	247251	1527654
TTTGTTT	64233	102180	480650	1096963
GTTTG	183720	460821	2559491	3519206
AWTAAA	227579	528285	1951519	5305867

Majority of studies regarding AU rich elements uses animals as the target (Fan et al., 1997; He et al., 2012; Hitti et al., 2016; Spasic et al., 2012) while present study is the first of its kind where the magnitude of putative ARE sites are estimated in plants at the genome-wide scale.

Table 3: Number of ARE motifs present in transcriptomes of respective plant species and H.sapiens

Motifs	A. thaliana	O. sativa	Z. mays	Homo sapiens
ATTT	621490	251813	660984	3614954
ATTTA	105252	41320	113303	742011
TATTTAT	9971	4418	12667	84071
AATTTAA	7857	3122	7172	69770
WATTTAW	32771	14145	36793	292870
WWATTTAWW	12356	5001	12591	128664
WTTTW	496558	180175	476968	3153179
WWTTTWW	139497	55828	144431	1190921
WWWTTTWWW	50792	20978	50428	509720
WWWWTTTWWWW	21144	8388	18252	246544
WWWWWTTTWWWWW	9773	3690	7536	124387
TTTGTTT	45076	17609	43304	329354
GTTTG	149631	55170	162416	584026
AWTAAA	70352	22383	61958	615665

Hitherto, the presence of ARE motifs in 3`UTR only has been taken into cognisance by the various scientific studies but we made attempt to investigate the occurrence of ARE motifs at genomic and transcriptomic colosseum also. This comprehensive analysis made us to draw some interesting conclusions (Figure 2). The ARE motif occurrence in genomic data was anticipated to be directly correlated with the number of nucleotides. But corelation of number of ARE motifs at

transcriptomic and 3`UTR scale exhibited some very intriguing inclinations. For instance, the number of different ARE motifs in *A.thaliana* is not the exclusive reflection of the size of *A.thaliana* transcriptome size. Transcriptome size of *A.thaliana* (95.0 MB) is 33.50% smaller than *Z. Mays* (143 MB), regardless of this difference in size, number of ARE motifs found in *A.thaliana* are just 2% lesser than *Z.mays*. Further, this difference seems to be contributed by the overwhelming presence of "WTTTW" and "ATTT" motifs in the transcriptome of *A.thaliana* (Table 2, Table 3, Table 4, Figure 2). Further, this trend got reversed in 3`UTR data. Size of 3`UTR of *A.thaliana* is just 3.65% smaller than *Z.mays* while the number of putative ARE motifs are reported to be 46.1% higher in *A.thaliana* (Table 2, Table 3, Table 4, Figure 2).

Table 4: Number of ARE motifs present in 3`UTR of respective plant species and H.sapiens

Motif	A. thaliana	O. sativa	Z. mays	Homo sapiens
ATTT	84196	103982	49690	1313377
ATTTA	16111	18444	8759	269479
TATTTAT	2456	2380	1334	36417
AATTTAA	1289	1418	554	25487
WATTTAW	6783	7024	3357	115982
WWATTTAWW	2988	2747	1283	53239
WTTTW	86239	87007	43014	1251880
WWTTTWW	31031	30062	15043	503836
WWWTTTWWW	12931	12251	6104	224165
WWWWTTTWWWW	5829	5082	2331	113232
WWWWWTTTWWWWW	2756	2203	1009	58267
TTTGTTT	7649	4845	3098	62599
GTTTG	17445	20039	13765	200825
AWTAAA	11346	10371	6447	261558

The magnitudes of genomes, transcriptome and 3`UTR of all the three plant species were disproportionate and incomparable with *Homo sapiens*. For the purpose of accessing the general propensity of ARE motifs in all species at the genome, transcriptome and 3`UTR level we standardized the size of the respective genome, transcriptome and 3`UTR by converting them to 100 MB (one unit) (Figure 3). Following size standardization, ARE motifs at a different stages can be compared realistically.

Figure 3: General trend of total ARE motifs at genome, transcriptome and 3`UTR level. Size at each level is taken as 100 MB (1 Unit) in order to make the data of ARE motifs comparable.

The present study indicated differential presence of ARE motifs in different plant species. But strong presence of ARE motifs at each level in all species under study cannot be denied. The most important inference deduced is that, similar to *H.sapiens*, plants also have more number of ARE motifs at 3`UTR than in per unit transcriptome. This is a breakthrough observation, which indicates the high probability of occurrence of ARE motifs in plants. The comparable magnitude of various ARE motifs at the genome, transcriptome and 3`UTR level in plants with *H. sapiens* indicates their possible roles in plant cell functioning similar to *H.sapiens*.

Additionally *in-vitro, in-vivo* and computational interventions are needed to acertain if the plants also have significant role of AU rich elements similar to mammals. The preliminary study conducted here categorically indicated the wide swath presence of AU rich elements in plants. Most prominent ARE motif and their role ascertained through further experimentation will open new avenues in the field of plant immunity and plant cell metabolism studies.

Conclusion and future Prospects

The prevailing and significant ARE motifs are predicted computationally in three plant species and statistically proved to have a explicit presence in different plants species in the present genome-wide study. This is the first-ever study encompassing three different plants, showing the presence of AU rich elements at different genomic and transcriptomic levels. Existence of AU rich elements in plants could be used to study their role in plant immunity, stress tolerance, reproduction and productivity. The present study qualified to be an initiation point for further computational, statistical, in-vitro and in-vivo analysis.

Figure 4: Pipeline of proposed computational analysis of genome-wide ARE motifs

Figure 4 provided a concrete pipeline for comprehensive computational and statistical extrapolation. This pipeline suggests ARE cluster analysis for enhanced interpretation of ARE motif occurrence

and functioning. Understanding of localization and classification of ARE motifs is suggested to circumscribe 5`UTR and introns besides regions studied in the present case.

ARE sites proximity might also be studied to get a extensive insight about the functioning of AU rich element directed mRNA decay. Understanding of appropriate proximity for functional ARE motifs will be of immense interest because cellular machinery needs space to get loaded on ARE signals. Student's t-test, Two-way ANOVA test and Kolmogorov-Smimov test might be used for statistical analysis of elaborated results.

To get a comprehensible view regarding presence and significance of AU rich elements in plants study might be extended to encompass more number of plant species having complete genomes available and plants of immense economic importance e.g. *Triticum* aestivum, *Sorghum bicolour, Vitis vinifera, Medicago truncatula, Solanum tuberosum.*

References:

Bakheet, T., 2001. ARED: human AU-rich element-containing mRNA database reveals an unexpectedly diverse functional repertoire of encoded proteins. Nucleic Acids Research 29, 246–254. https://doi.org/10.1093/nar/29.1.246

Barreau, C., 2005. AU-rich elements and associated factors: are there unifying principles? Nucleic Acids Research 33, 7138–7150. https://doi.org/10.1093/nar/gki1012

Caput, D., Beutler, B., Hartog, K., Thayer, R., Brown-Shimer, S., Cerami, A., 1986. Identification of a common nucleotide sequence in the 3'-untranslated region of mRNA molecules specifying inflammatory mediators. Proceedings of the National Academy of Sciences 83, 1670–1674. https://doi.org/10.1073/pnas.83.6.1670

Chen, C.-Y.A., Shyu, A.-B., 1995. AU-rich elements: characterization and importance in mRNA degradation. Trends in Biochemical Sciences 20, 465–470. https://doi.org/10.1016/S0968-0004(00)89102-1

Duttagupta, R., Vasudevan, S., Wilusz, C.J., Peltz, S.W., 2003. A Yeast Homologue of Hsp70, Ssa1p, Regulates Turnover of the MFA2 Transcript through Its AU-Rich 3′ Untranslated Region. MCB 23, 2623–2632. https://doi.org/10.1128/MCB.23.8.2623-2632.2003

Eberhardt, W., Doller, A., Akool, E.-S., Pfeilschifter, J., 2007. Modulation of mRNA stability as a novel therapeutic approach. Pharmacology & Therapeutics 114, 56–73. https://doi.org/10.1016/j.pharmthera.2007.01.002

Espel, E., 2005. The role of the AU-rich elements of mRNAs in controlling translation. Seminars in Cell & Developmental Biology 16, 59–67. https://doi.org/10.1016/j.semcdb.2004.11.008

Fan, X.C., Myer, V.E., Steitz, J.A., 1997. AU-rich elements target small nuclear RNAs as well as mRNAs for rapid degradation. Genes & Development 11, 2557–2568. https://doi.org/10.1101/gad.11.19.2557

He, G., Sun, D., Ou, Z., Ding, A., 2012. The Protein Zfand5 Binds and Stabilizes mRNAs with AU-rich Elements in Their 3′-Untranslated Regions. J. Biol. Chem. 287, 24967–24977. https://doi.org/10.1074/jbc.M112.362020

Hitti, E., Bakheet, T., Al-Souhibani, N., Moghrabi, W., Al-Yahya, S., Al-Ghamdi, M., Al-Saif, M., Shoukri, M.M., Lánczky, A., Grépin, R., Győrffy, B., Pagès, G., Khabar, K.S.A., 2016. Systematic Analysis of AU-Rich Element Expression in Cancer Reveals Common Functional Clusters Regulated by Key RNA-Binding Proteins. Cancer Res 76, 4068–4080. https://doi.org/10.1158/0008-5472.CAN-15-3110

Khabar, K.S.A., 2005. The AU-Rich Transcriptome: More Than Interferons and Cytokines, and Its Role in Disease. Journal of Interferon & Cytokine Research 25, 1–10. https://doi.org/10.1089/jir.2005.25.1

Lam, L.T., Pickeral, O.K., Peng, A.C., Rosenwald, A., Hurt, E.M., Giltnane, J.M., Averett, L.M., Zhao, H., Davis, R.E., Sathyamoorthy, M., Wahl, L.M., Harris, E.D., Mikovits, J.A., Monks,

A.P., Hollingshead, M.G., Sausville, E.A., Staudt, L.M., 2001. Genomic-scale measurement of mRNA turnover and the mechanisms of action of the anti-cancer drug flavopiridol. Genome Biol 2, research0041.1. https://doi.org/10.1186/gb-2001-2-10-research0041

Shaw, G., Kamen, R., 1986. A conserved AU sequence from the 3′ untranslated region of GM-CSF mRNA mediates selective mRNA degradation. Cell 46, 659–667. https://doi.org/10.1016/0092-8674(86)90341-7

Spasic, M., Friedel, C.C., Schott, J., Kreth, J., Leppek, K., Hofmann, S., Ozgur, S., Stoecklin, G., 2012. Genome-Wide Assessment of AU-Rich Elements by the AREScore Algorithm. PLoS Genet 8, e1002433. https://doi.org/10.1371/journal.pgen.1002433

Wilson, G.M., Sun, Y., Lu, H., Brewer, G., 1999. Assembly of AUF1 Oligomers on U-rich RNA Targets by Sequential Dimer Association. J. Biol. Chem. 274, 33374–33381. https://doi.org/10.1074/jbc.274.47.33374

Wilusz, C.J., Wormington, M., Peltz, S.W., 2001. The cap-to-tail guide to mRNA turnover. Nat Rev Mol Cell Biol 2, 237–246. https://doi.org/10.1038/35067025