The TAT Protein Transduction Domain as an Intra-articular Drug-Delivery Technology

Sarah E. Mailhiot, Ph.D., sarahmailhiot@gmail.com[1]
Matthew A. Thompson, m.a.thompson14@comcast.net[2]
Akiko E. Eguchi, Ph.D., akieguchi@clin.medic.mie-u.ac.jp[3]
Sabrina E. Dinkel, B.S., sabrinadinkel@hotmail.com[2]
Martin K. Lotz, M.D., mlotz@scripps.edu[4]
*Steven F. Dowdy, Ph.D., SDOWDY@ucsd.edu[5]
*Ronald K. June II, Ph.D., rjune@montana.edu[2]

[1]University of Oulu, Oulu, Finland
[2]Montana State University, Bozeman, MT, USA
[3]Mie University, Tsu, Mie, Japan
[4]The Scripps Research Institute, La Jolla, CA, USA
[5]University of California, San Diego, La Jolla, CA, UsA

Word Count (including references): 2970 (Abstract through Acknowledgements)

Grant Support: NSF 1554708, NIH R01AR073964 and R37 AG059418

*Co-Corresponding Authors
Steven F. Dowdy, Ph.D.
Professor
Dept. of Cellular & Molecular Medicine
UCSD School of Medicine
9500 Gilman Drive, MC-0686
La Jolla, CA 92093-0686
SDOWDY@ucsd.edu

Ronald K. June II, Ph.D., Associate Professor
Montana State University, c/o MIE Department
220 Roberts Hall
Bozeman, MT 59717-3800
rjune@montana.edu
Office: 406.994.2203

49 **Abstract**
50

<u>Objective</u>

Intra-articular drug delivery holds great promise for the treatment of joint diseases such as

osteoarthritis. The objective of this study was to evaluate the TAT peptide transduction

domain (TAT-PTD) as a potential intra-articular drug delivery technology for synovial

joints.

<u>Design</u>

Experiments examined the ability of TAT conjugates to associate with primary

chondrocytes and alter cellular function both *in vitro* and *in vivo*. Further experiments

examined the ability of the TAT-PTD to bind to human osteoarthritic cartilage.

<u>Results</u>

The results show that the TAT-PTD associates with chondrocytes, is capable of delivering

siRNA for chondrocyte gene knockdown, and that the recombinant enzyme TAT-Cre is

capable of inducing *in vivo* genetic recombination within the knee joint in a reporter mouse

model. Lastly, binding studies show that osteoarthritic cartilage preferentially uptakes the

TAT-PTD from solution.

<u>Conclusions</u>

The results suggest that the TAT-PTD is a promising delivery strategy for intra-articular

therapeutics.

Keywords: cartilage; osteoarthritis; drug delivery; protein transduction; intra-articular

Introduction

Intra-articular drug delivery is important for targeting therapeutic compounds to the diverse cells and tissues involved in arthritis pathogenesis (*i.e.* cartilage, bone, and synovium). Joint vasculature includes large pores (~1-3 μm) in synovial blood vessels leading to fenestrated capillaries[4,17]. These, as well as lymphatic drainage[9,25], create challenges for drug residence times. Measurements show that most unmodified molecules do not possess sufficient joint residence time for effective pharmacologic intervention [8,15,21,22,32], demonstrating a need for new delivery strategies.

Previous studies have explored synovial joint drug delivery using nanoparticles, microspheres, and phase-transitioning polymers [6,13,19,24,28]. Diclofenac sodium delivered by poly(lactic-co-glycolic acid) (PLGA) microspheres to antigen-induced chronic arthritic rabbits formulated for experimentally optimized conditions for extended release found no significant difference between PLGA treatment groups and control [28]. Zilretta is an FDA-approved corticosteroid formulation for OA that achieves extended release through PLGA (poly lactic-co-glycolic acid) microspheres [18,20]. Elastin-like polypeptides (ELPs), a type of phase-transitioning polymers set to aggregate at 37°C following intra-articular injection, were found to have increased residency time for protein drug delivery during biodistribution tests in rats versus a soluble ELP control [6]. These are exciting technologies show partial success in animal models.

The TAT peptide transduction domain (TAT-PTD) is a small, positively charged peptide that is electrostatically attracted to the negatively charged cell surface [30] and induces uptake in all cell types by binding to an as yet unidentified surface protein(s) [12].

PTDs have enabled delivery of small and large (~150 kDa) macromolecules into cells and

have been tested in >2000 patients in over 25 clinical trials [29] and resulted in 2

statistically-significant Phase II trials, one published [11]. Previous studies suggest that TAT-

PTD transduction may be effective for modulating joint biology [16,23], and PTDs remain to

be systematically evaluated for joint disease applications.

Therefore, the objectives of this study were to first determine if the TAT-PTD could

transduce macromolecules into chondrocytes, and second, to assess the ability of the

TAT-PTD to bind to cartilage. Toward these goals, this study applied both TAT-

conjugated peptides and proteins in *in vitro* and *in vivo* models. The result show that the

TAT-PTD is capable of inducing transduction and binding to the osteoarthritis (OA)

cartilage matrix. Future studies may build on these results to apply the TAT-PTD toward

drug delivery in OA.

Materials and Methods

Primary Human Chondrocytes

This study utilized primary chondrocytes obtained with IRB approval and harvested

from human knee joint femoral and tibial cartilage with Outerbridge grades 1-4. Sources

of joint tissue included both healthy tissues obtained from tissue banks from patients

without history of arthritis or any macroscopic signs of joint degeneration and osteoarthritic

patients undergoing joint replacement surgery. Primary human chondrocytes (mean age:

64.6 years, age range: 36-94, 73% male and 27% female) were harvested from the joint

by dissecting the cartilage as described [1], and cultured in DMEM high glucose media,

10% fetal calf serum (FCS), 1% penicillin and streptomycin (PS), 1% Glutamine. First

passage cells were used in the experiments.

<u>Peptide and Protein Synthesis</u>

The TAT protein transduction domain (RKKRRQRRR) was used to facilitate

intracellular uptake of candidate molecules into primary human chondrocytes and mouse

knee joints. For cell association studies, the TAT PTD and a control PTD were labeled

with rhodamine and synthesized using solid-phase peptide synthesis with Fmoc protection

in 25 μM scale on a Symphony Quartet peptide synthesizer [14]. All peptides were cleaved

and deprotected using a standard protocol (95% trifluoroacetic acid, 1% water, 1%

triisopropylsilane). Crude peptides were precipitated using cold diethylether and purified

using prep-scale reversed-phase HPLC with a C18 column; purity was confirmed by

MALDI-TOF mass spectrometry using the matrix α-CHCA. Recombinant proteins were

purified as described previously [5,10]. Plasmids containing either TAT-Cre [31] or the TAT

PTD-DRBD (Protein-Transduction-Domain coupled to a Double-Stranded RNA Binding

Domain) [10] were transformed into BL21 Codon Plus *e. coli.* and expanded to ~1-2L scale.

Expression was induced via IPTG (isopropyl β-D-1 thiogalactopyranoside), and cells were

harvested after 2-3 hours. Multi-step purification proceed sequentially via affinity

chromatography, cation exchange, and anion exchange yielding purified proteins for *in*

vivo experiments [10]. Purified proteins were quantified via SDS-PAGE electrophoresis and

Coomassie staining against BSA standards. Activity was confirmed via *in vitro* assays in

established cell lines.

<u>Cell Association Experiments</u>

To assess the relevance of the TAT protein transduction domain as a drug delivery

tool for synovial joints, fluorescently labeled TAT peptides and siRNAs were used in cell

association experiments and assayed via flow cytometry and fluorescence microscopy as

follows. 50,000 cells per well were plated in 12 well plates. Two days later cells were

washed with HBSS and treated with control (GGSGGHHHHHHG) or TAT-peptides (1.25,

2.5, or 5 µM) for 1 hour or left untreated. Following treatment, cells were washed three

times with HBSS containing 0.5 mg/mL heparin sulfate, detached with accutase, and

resuspended in growth media for analysis. Cell-associated fluorescence was measured

via flow cytometry. Live cells were discriminated via forward and side scatter; cell

association was quantified using the mean fluorescence intensity. At each dosage, mean

fluorescence values between control and TAT-peptides were compared using paired

comparisons with an a priori significance level of 0.05. These studies were repeated for

chondrocytes from n = 6 donors. To evaluate the dose-dependent association of

fluorescently labeled TAT-peptides, primary human chondrocytes were cultured and

treated with either a rhodamine labeled TAT peptide or labeled control peptide (1-10µM)

for 10 minutes to 1 hour. The surface peptide was removed via HBSS washes and

accutase digestion prior to analysis by flow cytometry.

<u>In vitro siRNA Knockdown</u>

For a functional evaluation of the TAT PTD as a drug delivery method for synovial

joints, siRNA knockdown experiments were performed using the TAT PTD-DRBD [10].

siRNA was complexed with the PTD-DRBD at a 1:8 molar ratio, and cells were treated at

concentrations of 50-400 nM siRNA. Control cells were either untreated or treated by

cationic lipids (Lipofetamine 2000, Invitrogen) according to the manufacturer's

recommendations. For imaging, fluorescently-labeled siRNAs were used, and after 1 hour

of treatment cells were washed with PBS and then trypsin, followed by immediate

imaging. Otherwise, siRNAs targeting either GAPDH or FoxO1 were used. The off-target

control siRNA was designed against GFP [10]. siRNA effectiveness was determined via

qPCR (GAPDH normalized to β-macroglobulin) or Western blotting (FoxO1). Functional

siRNA-knockdown was examined via immunoblotting for FoxO1 and qPCR for GAPDH.

All antibodies were from AbCam, and siRNAs were validated sequences from Ambion

using previously described methodology [2].

<u>In vivo Genetic Recombination</u>

To assess the *in vivo* ability of TAT PTD fusions, to modulate synovial joint

biology, the protein TAT-Cre [31] was injected into the stifle joints of luciferase reporter mice

[26]. TAT-Cre was prepared as previously described. TAT-Cre was injected (10 μL of 60

μM) into the medial side of the left knees in the luciferase reporter mice (n=5). The

contralateral knees were injected with buffer as a control. After treatment, *in vivo*

luciferase imaging (IVIS Spectrum, Perkin Elmer, Waltham MA, USA) was performed

daily for 4 days following intraperitoneal injection of D-luciferin (150 mg/kg). After imaging

on day 5, a single mouse was euthanized and both knee joints were dissected to further

evaluate the tissue source of the luciferase signal. The remaining mice were imaged

monthly thereafter until 10 months following injection.

<u>Binding studies in Human OA Cartilage</u>

Explants of Kellgren-Lawrence Grade IV OA cartilage were obtained from knee

arthroplasty cases under IRB approval and equilibrated in sterile tissue culture.

Equilibrium peptide uptake was measured with previously published methods [7]. Briefly,

structurally intact 3 mm osteochondral cores (N=3) of human stage IV OA cartilage were

taken from patients undergoing knee replacements. For control studies, samples were

stored in phosphate buffered saline (PBS) with protease inhibitors for 24 hours prior to

experimentation. To investigate the effect of the negative fixed charge density in cartilage,

500 mM NaCl was added to the PBS to disrupt the negative fixed charge of the

proteoglycans. To investigate how bulk damage effects cartilage, samples were degraded

in 5 mM trypsin for 24 hours prior to uptake studies.

During uptake studies, samples were incubated in PBS plus protease inhibitors

and 100 mM labeled peptide, a control peptide or TAT, and graded amounts (150-400

mM) of unlabeled TAT for 48 hours. The control peptide is a peptide with the same fixed

solubility and size as TAT but without the functionality of TAT. The control is referred to as

control-TAMRA. The wet weight of each cartilage disk was measured. The fluorescence

excitation of the solution was measured at 560 nm, and calibration curves were created

based on known standards for both the TAT and control peptides. The sample was dried

to measure the dry weight. The dried sample was then degraded with proteinase K to

measure the uptake of peptide by fluorescence excitation. The uptake ratio was

calculated, defined as the concentration of labeled peptide inside the cartilage divided by

concentration of labeled peptide in the solution. We used linear regression to assess the

dependence of uptake ratio on peptide concentration.

Statistical Analysis

Analysis of variance (ANOVA) was used to detect statistical differences in

quantitative measures between experimental groups. Planned comparisons between

groups of interest were made using an *a priori* significance level of α = 0.05 with the

Bonferroni correction for multiple comparisons. Cell association experiments were

analyzed by performing a 2-factor ANOVA using the dependent variable of the mean live-

cell fluorescence in the rhodamine channel via flow cytometry. Independent variables

were peptide (TAT-PTD or control) and dosage, and paired comparison s were made

between TAT-PTD and control at each dosage. Results are plotted as normalized to the

average value achieved in the maximum group. GAPDH knockdown was assessed using

a 1-way ANOVA with groups of untreated, control-lipofectamine, siGAPDH-lipofectamine,

control-PTD-DRBD, and siGAPDH-PTD-DRBD. Bonferroni post-hoc tests were used to

infer specificity of knockdown comparting the siGAPDH groups to the untreated control

groups with an *a priori* significance level of α = 0.05. All data are expressed as mean ±

standard error of the mean. *In vivo* bioluminescence imaging experiments were assessed

by comparing the quantitative luminescence data for the buffer and TAT-Cre injected

knees over time using a 2-way ANOVA with planned comparisons between the buffer and

TAT-Cre injected knees at each timepoint. We used linear regression to assess the

dependence of uptake ratio on peptide concentration.

Results

TAT-PTD Associates with Primary Human Chondrocytes

As a first test of the TAT-PTD, fluorescent dye TAT conjugates were incubated

with primary chondrocytes *in vitro* followed by trypsinization, re-attachment and culture for

24 hours, and then either flow cytometry or fluorescent imaging. The TAT-PTD dye

resulted in significantly more cell-associated fluorescence than the control peptide in a

dose-dependent manner (Fig. 1A). The mean fluorescence intensity (MFI) for TAT-

treated cells was significantly greater than for the chondrocytes treated with the control

peptide at all doses (1.25-5 μM, all p $\leq$ 0.02). Cell association of TAT peptides was

observed in both male and female chondrocytes from donors ranging in age from 24 to

64, and from normal and OA donors (Fig. 2). Because the cells were trypsinized prior to

analysis, these data indicate that the TAT-PTD has strong potential for intracellular

delivery of both experimental and therapeutic molecules in primary human chondrocytes.

<u>Functional Delivery of siRNA using the TAT-DRBD</u>

To deliver siRNAs into monolayers of primary human chondrocytes in monolayer,

the siRNA was bound by a TAT-PTD-DRBD (Double Stranded RNAi Binding Domain)[10].

The TAT-PTD was able to efficiently deliver fluorescently-labeled siRNA to primary human

chondrocytes in monolayer. Fluorescence imaging (Fig. 3A) found diffuse fluorescence

using the PTD-DRBD whereas Lipofectamine resulted in punctuate staining. Significant

GAPDH mRNA knockdown was observed with both Lipofectamine (p=0.008) and the

PTD-DRBD (p<0.0001, Fig. 3B). In addition, protein-level knockdown of FoxO1 was

achieved with both delivery methods and resulted in concomitant decreases in GAPDH

(Fig. 3C). These data indicate that the TAT-fusion protein PTD-DRBD is capable of

delivering functional siRNAs for gene knockdown in primary articular chondrocytes.

<u>*In Vivo* Efficacy of TAT-Cre using Reporter Mice</u>

To test the *in vivo* efficacy of the TAT PTD, the recombinant enzyme TAT-Cre was

injected intra-articularly into bioluminescent luciferase reporter mice. TAT-Cre injection

resulted in significantly greater bioluminescence (p < 0.001) than buffer at all timepoints

(all p < 0.01). Note that previous studies established that Cre-only does not enter cells

[31].There was no main effect of time after injection on induced bioluminescence (p =

0.112). Bioluminescence was observed as rapidly as day 1 after injection and was

sustained for 196 days (Fig. 4). Bioluminescence was observed in both the femoral

condyles and the tibial plateau after injection of TAT-Cre.

<u>Human OA Cartilage Preferentially Uptakes the TAT-PTD</u>

To assess the transport properties of the TAT-PTD within human articular

cartilage, fluorescent TAT and control peptides were incubated with cartilage explants

(Fig. 5). The uptake ratio was not dependent on concentration of unlabeled TAT (slope: -

0.0006, 95% CI for uptake ratio: 1.67-3.11. n=48) or control peptide (slope: 0.0002, 95%

CI: 0.004-2.31. n=48) (Fig. 5A). Therefore, the uptake ratio was equivalent to the

partitioning coefficient [7]. The mean partitioning coefficient of the labeled TAT peptide was

2.39, and the 95% confidence interval did not include 1, indicating that TAT causes uptake

within human cartilage (Fig. 5A). The partitioning coefficient of TAT-PTD was reduced

when the cartilage disk was treated with trypsin (p=0.045, n=24, Figure 5), but not 500

mM NaCl (p=0.3, n=24).

Discussion

The objectives of this study were first to determine if the TAT-PTD could transduce

macromolecules into chondrocytes *in vitro*, and second to assess the ability of the TAT-

PTD to bind to cartilage and transduce cells that are embedded in the cartilage

extracellular matrix. The *in vitro* studies show that the TAT-PTD associated with primary

human chondrocytes and is capable of delivering functional siRNA for knockdown of gene

and protein expression by RNA interference. Using luciferase reporter mice, *in vivo*

injection of TAT-Cre resulted in induction of luciferase activity indicating that the TAT-PTD

can deliver functional enzymes directly to synovial tissues. Binding studies in human

osteoarthritic cartilage showed that the TAT-PTD preferentially binds to cartilage in a

trypsin-dependent manner, suggesting that the TAT-PTD may be capable of extending the

residence time of molecules injected into the joint.

The binding studies in human OA cartilage retrieved during joint replacement

found preferential binding of the TAT-PTD. To quantify uptake into cartilage, we defined

the uptake ratio as the ratio of the concentration of labeled peptide inside the cartilage to

the concentration of labeled peptide in the bath. Because the lower bound of the 95%

confidence interval for the uptake ratio of the TAT-PTD is greater than one, these data

show that the TAT-PTD increased uptake and partitioning of a fluorescent marker

TAMRA. Furthermore, the equilibrium distribution of TAT-PTD between cartilage and the

fluid indicated a partitioning coefficient greater than 1. Additionally, trypsin treatment

decreased uptake of TAT-PTD. This is notable because this is a common model for

disease conditions such as OA. Because these studies were performed in late-stage OA

cartilage, it was likely that the TAT-PTD may have even greater uptake and partitioning in

healthy and early-stage cartilage, although future studies are needed to confirm this.

These studies have important limitations. First, while the binding studies were

performed in PBS, it is unclear if the TAT-PTD will have similar transport properties within

synovial fluid of the joint. Thus, there is the possibility that delivery using the TAT-PTD

may be less efficient *in vivo*. The cell-association measurements cannot conclusively be

interpreted as evidence of chondrocyte TAT-PTD internalization. However, because

association measurements were performed after trypsinization these data, as well as the

siRNA knockdown studies, strongly suggest that the TAT-PTD is internalized by primary

human chondrocytes. While the positive results from the injection of TAT-Cre into reporter

mice suggest that *in vivo* delivery is possible, this study did not determine which joint cells

were transduced, and the efficacy of the TAT-PTD to deliver in human joints remains

unknown. Furthermore, although transduction was demonstrated in both male and female

chondrocytes, it is unclear if there are differences between male and female joint cells and

tissues. A final limitation of these studies is that all studies were examined at equilibrium.

The specific dynamics of TAT-PTD binding to articular cartilage remain unknown.

Addressing these limitations in future studies will enable more-precise modeling and

prediction of the dynamics of the TAT-PTD within synovial joints, and this information will

be valuable for assessing residence times of potential therapeutic compounds.

Previous work with AVIDIN, a small positively charged biotin binding protein,

shows that electrostatic interactions enhance uptake in intra-articular injections with higher

uptake of charged particles in the negatively charged cartilage as compared to

surrounding tissues [3]. Similarly, the TAT-PTD is positively charged and has the

advantage of also inducing intracellular uptake of its bound molecules [27]. Future work will

test the correlation between GAG content and TAT-PTD uptake with DMMB blue [7].

The data demonstrate that TAT-PTD can be used to deliver siRNA and protein to

modulate chondrocyte biology. Consistent knockdown of GAPDH was achieved using the

PTD-DRBD. TAT fusions penetrate articular cartilage and display prolonged joint

residence time. The TAT protein transduction domain facilitates peptide-uptake in primary

human chondrocytes from male and female donors with varying ages and OA grades.

Taken together, our data suggest that cytosolic delivery using the TAT-PTD has strong

potential for both basic and translational studies of synovial joint biology and osteoarthritis.

Figures
Figure 1

348
349
Figure 1 Dose-dependent association of TAT-PTDs with primary human chondrocytes.
Primary human chondrocytes were cultured and treated with either a rhodamine labeled
TAT peptide or labeled control peptide. Surface peptide was removed via HBSS washes
and accutase digestion prior to flow cytometry. Significantly more fluorescence was found
for all treatment dosages of TAT peptide compared to control peptide. Data from n=11
independent donors.
356

357
Figure 2 Cell association of TAT-PTDs in chondrocytes from a diverse array of donors.
Each panel shows a Safranin-O Fast Green stained section (top) and flow cytometry
histogram (bottom) from representative donors. For histograms, black population is
untreated cells, blue is cells treated with a control rhodamine-labeled peptide, and red is a
population treated with 5 µM rhodamine-labeled TAT peptide. (A) 25-year old male with
Mankin 1 deterioration. (B) 39-year old female with Mankin 2 deterioration. (C) 64-year
old male with Mankin 3 deterioration. (D) 64-year old female with Mankin 3 deterioration.
Cell association experiments performed on n=10 donors.
366

Figure 3

367
Figure 3
The TAT fusion protein PTD-DRBD delivers siRNA for RNA silencing in primary human
chondrocytes. (A) siRNA was complexed to the PTD-DRBD TAT delivery protein (PTD-
DRBD) or lipofectamine (L2K), and cells were treated as in Fig. 1A. Following PBS and
trypsin washes, cells were imaged via phase contrast and epifluorescence microscopy.
(B) Knockdown of GAPDH mRNA. Primary human chondrocytes were treated for 4
hours with siRNA designed against either GFP as a control (siGFP) or GAPDH
(siGAPDH) as a target gene. mRNA levels were quantified by qPCR. Both lipofectamine
and the PTD-DRBD resulted in significant knockdown of GAPDH mRNA. (C) Protein-level
knockdown of FoxO1 measured by immunoblot. Primary human chondrocytes treated as
in A with control siRNA (siGFP) or siRNA designed against FoxO1. Immunoblots
performed following lysis and SDS-PAGE after 48 hours. Knockdown experiments using
the TAT PTD-DRBD was performed on chondrocytes from n=10 donors with an siRNA
concentration of 400 nM. Legend: L2K = lipofectamine 2000. PTD-DRBD = protein
transduction domain coupled to a double-stranded RNA binding domain.
383

385
Figure 4 TAT-Cre induces genetic recombination in mouse knees. Cre-reporter mice
were injected with either TAT-Cre or buffer in the intra-articular space prior to *in vivo*
bioluminescence imaging. (A) Image of n=5 mice 4 days following injection. (B)
Representative dissected mouse knee. Top shows photograph. Bottom shows
bioluminescent image. (C) Quantification of TAT-Cre induced bioluminescence.
Luciferase activity was quantified within circular regions of interest overlaid on both the
control- and TAT-Cre- injected knees.
393

394
Figure 5
Transport studies of the TAT-PTD in human osteoarthritic articular cartilage. (A)
Equilibrium uptake ratio between cartilage samples and bath for the TAT-PTD across a
range of peptide concentrations in the bath. The uptake ratio of TAT-PTD within cartilage
showed a slope < 0.01, indicating that partitioning is not concentration dependent.
Therefore, the partitioning coefficient is the uptake ratio. (B) Both high-salt and trypsin
resulted in decreased partitioning coefficients compared to untreated cartilage. This
suggests that electrostatic interactions between cartilage GAGs and the TAT-PTD
mediate the uptake of TAT-PTDs into cartilage.
404

References

1.   Akagi R, Akatsu Y, Fisch KM, Alvarez-Garcia O, Teramura T, Muramatsu Y, et al.
Dysregulated circadian rhythm pathway in human osteoarthritis: NR1D1 and
BMAL1 suppression alters TGF-beta signaling in chondrocytes. *Osteoarthritis*
Cartilage. 2017;25(6):943-951.

2.   Akasaki Y, Alvarez-Garcia O, Saito M, Carames B, Iwamoto Y, Lotz MK. FoxO
transcription factors support oxidative stress resistance in human chondrocytes.
Arthritis Rheumatol. 2014;66(12):3349-3358.

3.   Bajpayee AG, Scheu M, Grodzinsky AJ, Porter RM. Electrostatic interactions
enable rapid penetration, enhanced uptake and retention of intra-articular injected
avidin in rat knee joints. *J Orthop Res.* 2014;32(8):1044-1051.

4.   Barland P, Novikoff AB, Hamerman D. Electron microscopy of the human synovial
membrane. *The Journal of cell biology.* 1962;14:207-220.

5.   Becker-Hapak M, Dowdy SF. Protein transduction: generation of full-length
transducible proteins using the TAT system. *Current protocols in cell biology /*
editorial board, Juan S Bonifacino [et al]. 2003;Chapter 20:Unit 20 22.

6.   Betre H, Liu W, Zalutsky MR, Chilkoti A, Kraus VB, Setton LA. A thermally
responsive biopolymer for intra-articular drug delivery. *Journal of controlled*
release : official journal of the Controlled Release Society. 2006;115(2):175-182.

7.   Byun S, Tortorella MD, Malfait AM, Fok K, Frank EH, Grodzinsky AJ. Transport
and equilibrium uptake of a peptide inhibitor of PACE4 into articular cartilage is
dominated by electrostatic interactions. *Arch Biochem Biophys.* 2010;499(1-2):32-
39.

8.   Chevalier X, Goupille P, Beaulieu AD, Burch FX, Bensen WG, Conrozier T, et al.
Intraarticular injection of anakinra in osteoarthritis of the knee: a multicenter,
randomized, double-blind, placebo-controlled study. *Arthritis and rheumatism.*
2009;61(3):344-352.

9.   Doan TN, Bernard FC, McKinney JM, Dixon JB, Willett NJ. Endothelin-1 inhibits
size dependent lymphatic clearance of PEG-based conjugates after intra-articular
injection into the rat knee. *Acta Biomater.* 2019;93:270-281.

10.   Eguchi A, Meade BR, Chang YC, Fredrickson CT, Willert K, Puri N, et al. Efficient
siRNA delivery into primary cells by a peptide transduction domain-dsRNA binding
domain fusion protein. *Nature biotechnology.* 2009;27(6):567-571.

11.   Glogau R, Blitzer A, Brandt F, Kane M, Monheit GD, Waugh JM. Results of a
randomized, double-blind, placebo-controlled study to evaluate the efficacy and
safety of a botulinum toxin type A topical gel for the treatment of moderate-to-

severe lateral canthal lines. *Journal of drugs in dermatology : JDD.* 2012;11(1):38-
45.

12. Gump JM, June RK, Dowdy SF. Revised role of glycosaminoglycans in TAT
protein transduction domain-mediated cellular transduction. *The Journal of*
biological chemistry. 2010;285(2):1500-1507.

13. Horisawa E, Hirota T, Kawazoe S, Yamada J, Yamamoto H, Takeuchi H, et al.
Prolonged anti-inflammatory action of DL-lactide/glycolide copolymer nanospheres
containing betamethasone sodium phosphate for an intra-articular delivery system
in antigen-induced arthritic rabbit. *Pharmaceutical research.* 2002;19(4):403-410.

14. June RK, Gogoi K, Eguchi A, Cui XS, Dowdy SF. Synthesis of a pH-sensitive
nitrilotriacetic linker to peptide transduction domains to enable intracellular delivery
of histidine imidazole ring-containing macromolecules. *Journal of the American*
Chemical Society. 2010;132(31):10680-10682.

15. Kaeding CC, Hill JA, Katz J, Benson L. Bupivacaine use after knee arthroscopy:
pharmacokinetics and pain control study. *Arthroscopy : the journal of arthroscopic*
& related surgery : official publication of the Arthroscopy Association of North
America and the International Arthroscopy Association. 1990;6(1):33-39.

16. Kim D, Jeon C, Kim JH, Kim MS, Yoon CH, Choi IS, et al. Cytoplasmic
transduction peptide (CTP): new approach for the delivery of biomolecules into
cytoplasm in vitro and in vivo. *Exp Cell Res.* 2006;312(8):1277-1288.

17. Knight AD, Levick JR. Morphometry of the ultrastructure of the blood-joint barrier in
the rabbit knee. *Quarterly journal of experimental physiology.* 1984;69(2):271-288.

18. Langworthy MJ, Conaghan PG, Ruane JJ, Kivitz AJ, Lufkin J, Cinar A, et al.
Efficacy of Triamcinolone Acetonide Extended-Release in Participants with
Unilateral Knee Osteoarthritis: A Post Hoc Analysis. *Adv Ther.* 2019;36(6):1398-
1411.

19. Lu Y, Zhang G, Sun D, Zhong Y. Preparation and evaluation of biodegradable
flubiprofen gelatin micro-spheres for intra-articular administration. *Journal of*
microencapsulation. 2007;24(6):515-524.

20. McAlindon TE, LaValley MP, Harvey WF, Price LL, Driban JB, Zhang M, et al.
Effect of Intra-articular Triamcinolone vs Saline on Knee Cartilage Volume and
Pain in Patients With Knee Osteoarthritis: A Randomized Clinical Trial. *JAMA.*
2017;317(19):1967-1975.

21. Neander G, Eriksson LO, Wallin-Boll E, Ersmark H, Grahnen A. Pharmacokinetics
of intraarticular indomethacin in patients with osteoarthritis. *European journal of*
clinical pharmacology. 1992;42(3):301-305.

22. Owen SG, Francis HW, Roberts MS. Disappearance kinetics of solutes from
synovial fluid after intra-articular injection. *British journal of clinical pharmacology.*
1994;38(4):349-355.

23. Ozaki D, Sudo K, Asoh S, Yamagata K, Ito H, Ohta S. Transduction of anti-
apoptotic proteins into chondrocytes in cartilage slice culture. *Biochem Biophys*
Res Commun. 2004;313(3):522-527.

24. Ratcliffe JH, Hunneyball IM, Wilson CG, Smith A, Davis SS. Albumin microspheres
for intra-articular drug delivery: investigation of their retention in normal and
arthritic knee joints of rabbits. *The Journal of pharmacy and pharmacology.*
1987;39(4):290-295.

25. Rodnan GP, Maclachlan MJ. The absorption of serum albumin and gamma
globulin from the knee joint of man and rabbit. *Arthritis and rheumatism.*
1960;3:152-157.

26. Safran M, Kim WY, Kung AL, Horner JW, DePinho RA, Kaelin WG, Jr. Mouse
reporter strain for noninvasive bioluminescent imaging of cells that have
undergone Cre-mediated recombination. *Molecular imaging.* 2003;2(4):297-302.

27. Schwarze SR, Ho A, Vocero-Akbani A, Dowdy SF. In Vivo Protein Transduction:
Delivery of a Biologically Active Protein into the Mouse. *Science.*
1999;285(5433):1569-1572.

28. Tuncay M, Calis S, Kas HS, Ercan MT, Peksoy I, Hincal AA. Diclofenac sodium
incorporated PLGA (50:50) microspheres: formulation considerations and in
vitro/in vivo evaluation. *International journal of pharmaceutics.* 2000;195(1-2):179-
188.

29. van den Berg A, Dowdy SF. Protein transduction domain delivery of therapeutic
macromolecules. *Current opinion in biotechnology.* 2011;22(6):888-893.

30. Varki, A., Cummings, R.D.M, Esko, J.D., Freeze, H.H., Stanley, P., Bertozzie,
C.R., Hart, G.W., and Entzler, M.E.. Essentials of Glycobiology, 2nd Edition. Cold
Spring Harbor, NY. Cold Spring Harbor Laboratory Press. 2009.

31. Wadia JS, Stan RV, Dowdy SF. Transducible TAT-HA fusogenic peptide enhances
escape of TAT-fusion proteins after lipid raft macropinocytosis. *Nature medicine.*
2004;10(3):310-315.

32. Wagner SC, Schweitzer ME, Weishaupt D. Temporal behavior of intraarticular
gadolinium. *Journal of computer assisted tomography.* 2001;25(5):661-670.

511