Pervasive generation of non-canonical subgenomic RNAs by SARS-CoV-2

Jason Nomburg[1,2,3], Matthew Meyerson[1,2,4,5*], James A. DeCaprio[1,3,5*]

[1]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston MA

[2]Broad Institute of MIT and Harvard, Cambridge, MA

[3]Harvard Program in Virology, Harvard University Graduate School of Arts and Sciences, Boston, MA

[4]Department of Genetics, Harvard Medical School, Boston, MA

[5]Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

*Correspondence to:

James A. DeCaprio

james_decaprio@dfci.harvard.edu

Matthew Meyerson

matthew_meyerson@dfci.harvard.edu

Abstract

Background: SARS-CoV-2, a positive-sense RNA virus in the family *Coronaviridae*, has caused a worldwide pandemic of coronavirus disease 2019 or COVID-19 Coronaviruses generate a tiered series of subgenomic RNAs (sgRNAs) through a process involving homology between transcriptional regulatory sequences (TRS) located after the leader sequence in the 5' UTR (the TRS-L) and TRS' located near the start of structural and accessory proteins (TRS-B) near the 3' end of the genome. In addition to the canonical sgRNAs generated by SARS-CoV-2, non-canonical sgRNAs (nc-sgRNAs) have been reported. However, the consistency of these nc-sgRNAs across viral isolates and infection conditions is unknown. The comprehensive definition of SARS-CoV-2 RNA products is a key step in understanding SARS-CoV-2 pathogenesis.

Methods: Here, we report an integrative analysis of eight independent SARS-CoV-2 transcriptomes generated using three sequencing strategies, five host systems, and seven viral isolates. Read-mapping to the SARS-CoV-2 genome was used to determine the 5' and 3' coordinates of all identified junctions in viral RNAs identified in these samples.

Results: Using junctional abundances, we show nc-sgRNAs make up as much as 33% of total sgRNAs *in vitro*, are largely consistent in abundance across independent transcriptomes, and increase in abundance over time during *in vitro* infection. By

assessing the homology between sequences flanking the 5' and 3' junction points, we

show that nc-sgRNAs are not associated with TRS-like homology. By incorporating read

coverage information, we find strong evidence for subgenomic RNAs that contain only

5' regions of ORF1a. Finally, we show that non-canonical junctions change the

landscape of viral open reading frames.

Conclusions: We identify canonical and non-canonical junctions in SARS-CoV-2

sgRNAs and show that these RNA products are consistently generated across many

independent viral isolates and sequencing approaches. These analyses highlight the

diverse transcriptional activity of SARS-CoV-2 and offer important insights into SARS-

CoV-2 biology.

Key Words

SARS-CoV-2, COVID-19, direct RNA sequencing, transcription

Background

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged from Wuhan, China in December 2019 and rapidly led to a world-wide pandemic, known as COVID-19 (1). The initial sequencing report found that this virus maintains 89.1% nucleotide identity to SARS-like coronaviruses found in bats in China (1).

Coronaviruses (CoV) including SARS-CoV-2 contain large, ~ 30kb, positive-sense, single-stranded RNA genomes with a unique genome organization structure. The 5' end of the genome contains two large open reading frames (ORFs), ORF1a and ORF1b. ORF1a produces a large polyprotein, while ribosomal slippage at an RNA pseudoknot structure and slippery sequence at the end of ORF1a occasionally leads to a frameshift and subsequent translation of a joint ORF1a and ORF1b polyprotein (2). ORF1a and ORF1ab polyproteins contain protease activity that can cleave the polyproteins into a variety of nonstructural proteins (Nsp) capable of facilitating viral transcription and replication and modulating host transcription and translation. Following ORF1b, there is an arrangement of ORFs encoding structural and accessory proteins that tend to vary in content and order across the CoV family. While structural proteins are incorporated into emergent virions, accessory proteins are thought to be dispensable for replication *in vitro* but to increase fitness *in vivo* (3-5).

Initial translation of ORF1ab can occur directly from incoming viral genomic RNA, generating the ORF1a and ORF1ab polyproteins and initiating infection. However, because the viral genome is so large, and because downstream ORFs are far from the

5' end of the genome, effective translation of these ORFs requires the generation of

subgenomic RNAs (sgRNAs) (6). To generate these sgRNAs, it is thought that CoV

undergoes a process of discontinuous extension during synthesis of negative-stranded

RNA templates. As the virus-encoded RNA-dependent RNA polymerase progresses

from the 3' end towards the 5' end of the viral genome, it encounters transcriptional

regulatory sequences (TRS) upstream of each major ORF (TRS-B). Here, the

polymerase can skip to a similar TRS that is 3' of a shared leader sequence within the

5' UTR of the CoV genome (TRS-L). These antisense sgRNAs are then transcribed,

resulting in a series of tiered sgRNAs containing the same 5' leader sequence (6).

Several studies have characterized SARS-CoV-2 sgRNAs using Nanopore direct RNA

sequencing (dRNAseq). In contrast to short-read sequencing, which can produce read

pairs of 150bp each, dRNAseq is capable of directly sequencing entire transcripts

including the full length of the viral genome (29903 bases). This allows dRNAseq to

identify RNA isoforms and variants that may be challenging to deconvolute using short-

read sequencing. Using nanopore sequencing, Taiaroa et al. identified eight major

sgRNAs, as well as a small number of non-canonical junctions between the 5' leader

sequence and downstream regions of the SARS-CoV-2 genome (7). Kim et al. identified

a total of 9 distinct subgenomic RNAs, as well as a series of non-canonical subgenomic

RNAs (nc-sgRNAs) that contain unexpected junctions between 5' and 3' sequences (8).

Using short-read DNA nanoball sequencing data, they confirmed the existence of

noncanonical junctions at various sites across the genome (8). Davidson et al. identified

a series of transcripts encoding the nucleocapsid (N) protein, but with distinct small

internal deletions, and validated this finding with proteomic evidence (9).

While these previous studies identified non-canonical RNA junctions and unexpected

RNA species in isolation, there were no integrative analyses of the SARS-CoV-2

junctions. It is unclear if identified non-canonical junctions are broadly representative of

SARS-CoV-2 biology or if they are a result of dataset- or isolate-specific artifacts. Here,

we report an integrative analysis of eight independent SARS-CoV-2 transcriptomes

generated using three sequencing strategies, five host systems, and seven viral isolates

to comprehensively characterize the landscape of canonical and non-canonical sgRNAs

generated by SARS-CoV-2.

Results

Analysis of three independent dRNAseq datasets reveals canonical and

noncanonical junctions.

To determine the major RNA species present in SARS-CoV-2-infected cells, we applied

a uniform computational approach to three independent dRNAseq datasets from

Taiaroa et al., Kim et al., and Davidson et al. (7, 10, 11). To assess sgRNA presence,

we identified junction-spanning reads following read-mapping against the SARS-CoV-2

reference genome.

Analysis of the 5' and 3' locations of junctions separated by at least 1000 bases

revealed that the majority of junctions consist of a 5' end originating in the 5' UTR and a

3' end at distinct sites in the 3' region of the genome (**Figure 1A-C**). These represent

the canonical junctions between the leader sequence and 3' sequences that generate

the major sgRNAs (6)**.** Plotting the counts of 3' junctions that contain a 5' end within the

first 100 bases revealed eight prevalent junction points along the 3' end of the genome

(**Figure 1D-F**), all of which are proximal to the TRS core sequence ACGAAC.

Interestingly, we observed that the 3' junction peak and TRS-B closest to ORF6 lands

151 bases within the M ORF. Based on these junction analyses and the locations of

identified TRS body sites, predicted sgRNAs and their most 5' gene or genes are shown

in **Figure 1G**. We predict 8 major sgRNAs based both on identification of the TRS core

sequence and presence of a dominant junction peak. In contrast to Kim et al. (8) but in

concordance with Taiaroa et al. (7) and Davidson et al. (9), we do not find evidence of a

prominent 3' junction point near ORF7B (**Figure 1D-F**).

Many SARS-CoV-2 junctions are non-canonical.

Similar to observations by Kim et al. (8), we identified an unexpectedly diffuse pattern of

noncanonical junctions across the genome (**Figure 1A-C**, black arcs), suggesting that

there are many unexpected RNA species. Had there been only canonical junctions, the

arcs on this plot would be exclusively between the first 100 bases and defined points on

the x-axis (**Figure 1A-F**).

Because the transcriptomes in these long-read sequencing studies (7, 8, 11) all consist

of dRNAseq from infected Vero cells, we sought to confirm this observation in

independent transcriptomes from other SARS-CoV-2 isolates, in other host systems,

and using other sequencing technologies. To this end, we assessed five additional

transcriptomes generated from Vero cells (12), A549 cells (13), Calu3 cells (14),

bronchial organoids (15), and ferret nasal washings (13) using Illumina PolyA or total

RNA sequencing. Detailed information regarding each sample is detailed in the

methods. Junction analysis in these transcriptomes likewise revealed pervasive non-

canonical junctions (**Supplementary Figure 1A-E**). Quantification of the relative

abundance of canonical and non-canonical junctions in these transcriptomes revealed

that up to 33% of junctions are non-canonical *in vitro* (**Figure 2A, Supplementary**

Figure 1F). The transcriptome from a ferret's nasal washing contained 51% non-

canonical junctions, indicating that non-canonical junctions are also produced *in vivo*.

We sought to assess if non-canonical junctions are driven by dataset-specific artifacts.

For example, it is possible that defective particles containing defective viral genomes

(DVGs), often with large internal deletions that increase their replicative fitness, could

accumulate during preparation of virion stocks (16) and generate the observed non-

canonical junctions during subsequent infections. If this was true, we would expect to

observe non-canonical junctions specific to a single virion preparation and accumulation

of these specific non-canonical junctions over the course of infection. To address this

possibility, we identified dataset-specific junctions by identifying junctions that are

differentially abundant across datasets (**Figure 2B**). This analysis revealed that the vast

majority of 5' and 3' non-canonical junctions are low abundance and are similarly

abundant across five independent SARS-CoV-2 sequencing studies (7-9, 12, 13)

(**Figure 2C, D**). This suggests that most non-canonical junctions are not driven by

dataset-specific artifacts.

However, we found that three 5' junction sites in the Finkel et al. transcriptome and

three 3' junctions in the Davidson et al. transcriptome were overrepresented (**Figure**

2C, D). Incorporating additional replicates of these transcriptomes and assessing the

percentage of non-canonical junctions at each site revealed that the Finkel et al.-

specific 5' junctions are highly elevated in both Finkel et al. replicates relative to other

transcriptomes (**Supplementary Figure 3A-C**). In contrast, the three Davidson et al.-

specific 3' junctions are present in other datasets but at slightly lower frequency

(**Supplementary Figure 3D, E**). The consistency of the Finkel et al.-specific junctions

across replicates and exclusion from other samples raises the possibility that these

junctions were generated from defective particles in the Finkel et al. virion preparation.

The proportion of non-canonical junctions increases over time *in vitro*.

Next, we assessed changes in junctions over time using the approach detailed in

Figure 3A using transcriptomes from early (4 or 5 hours post infection) and late (24

hours post infection) timepoints from Finkel et al. (12) and Emanual et al. (14). This

analysis revealed that there are large decreases in 5' junctions originating at position

64, and in 3' junctions ending in 28254 (proximal to the N TRS-B), with a smaller

decrease in 3' junctions ending at 26467 (proximal to the M TRS-B) (**Figure 3B,C**).

Notably, the Finkel et al.-specific 5' junction at 16971 increased in abundance over time.

In contrast, there were no specific large increases in percentage of junctions at other

positions (**Figure 3B,C**). Instead, there are small but widespread increases in non-

canonical junction sites throughout the SARS-CoV-2 genome (**Figure 3D, E**).

Quantification of the total percent changes of canonical and non-canonical junctions

revealed a large decrease in canonical N junctions and increase in non-canonical

junctions (**Figure 3F, G**). These results indicate that the abundance of non-canonical

junctions increases over time *in vitro*. It is unclear if a similar pattern occurs during

infection *in vivo*.

Non-canonical junctions are not associated with TRS-like homology.

While homology between the TRS-L and TRS-B is thought to be critical for generation

of canonical sgRNAs, the role of TRS-like homology in the generation of nc-sgRNAs is

unclear. To address this question, we determined the 30 bases surrounding the 5' and

3' sites of each junction and identified the longest continuous subsequence in common

between these sequence (**Figure 4A**). We found that junctions with a 3' end landing

within 15 nucleotides of a TRS-B overwhelmingly have junction homology associated

with variants of the TRS core sequence (**Figure 4B-D, blue**). In contrast, junctions

landing within ORFs and not proximal to a TRS-B do not have a prevalent homologous

sequence (**Figure 4B-D, red**). The exception are intra-M junctions, whose homology is

a consequence of the location of the ORF6 TRS within the M ORF.

While there is not a single strong motif driving non-canonical, intra-ORF junctions, it is

still possible that non-canonical junctions are associated with junction-specific homology

between the 5' and 3' junction sites. To address this possibility, we tallied the length of

homology between the 5' and 3' sites of canonical and non-canonical junctions. This

analysis revealed that canonical junctions have markedly longer homology lengths

(mean >10 bases) than non-canonical junctions (mean < 5 bases) (**Figure 4E-G**).

Indeed, the homology lengths of non-canonical junctions are only slightly longer than

homology lengths between 100,000 random pairs of SARS-CoV-2-derived 30 base

sequences (**Figure 4E-G**). These data indicate that non-canonical junctions are not

associated with TRS-like homology.

sgRNAs containing only a 5' region of ORF1a are produced in all studied *in vitro*

and *in vivo* infection systems.

Assessment of the three dRNAseq transcriptomes revealed evidence of sgRNAs that

contain only the 5' region of ORF1a (**Supplementary Figure 3A**). This is similar to an

observation previously reported by Kim et al. (8). Inspection of read coverage of ORF1a

in these transcriptomes confirmed that there is elevated 5' ORF1a coverage with

specific decreases in coverage concomitantly with sharp increases in junctions near

genome positions 1600 and 6000 (**Figure 5A-C**).

While there is clearly higher coverage of the 5' region of ORF1a in the three dRNAseq

transcriptomes, it remained unclear if these 5' ORF1a sgRNAs are a major percentage

of total ORF1a in the cell, or if this finding was due to a systematic bias favoring shorter

reads in nanopore sequencing. To address this question, we assessed ORF1a

coverage in four independent Illumina polyA-enriched RNAseq transcriptomes. Analysis

of these transcriptomes revealed a similar elevated 5' ORF1a coverage and consistent

drops in coverage around genome positions 1600 and 6000 (**Figure 5D-G**). Importantly,

one of these transcriptomes is from bronchial organoids (**Figure 5G**) and one

transcriptome is from the nasal washings of an infected ferret (**Figure 5F**), suggesting

that 5' ORF1a subgenomic RNAs can be produced *in vivo* and were not only due to

artifacts of highly replicative conditions *in vitro*. Furthermore, analysis of a transcriptome

generated from RNA that was not polyA-enriched revealed a consistent elevation in 5'

ORF1a coverage (**Figure 5H**), ruling out a systematic bias from polyA isolation. We

found that there is similar elevated 5' ORF1a coverage in the circulating human

coronavirus CoV 229E (17) (**Supplementary Figure 4**), supporting the existence of 5'

ORF1a RNAs in other coronaviruses. All together, these data support the hypothesis

that subgenomic RNAs containing just the 5' region of ORF1a are produced by SARS-

CoV-2.

Junctions have the potential to generate variant open reading frames.

To test the hypothesis that non-canonical junctions can result in variant open reading

frames (ORFs) in nc-sgRNAs, we assessed the ORF content of each read in the three

dRNAseq transcriptomes. Because dRNAseq can have an error rate of greater than

10% (18), we generated "coordinate-derived transcripts" by determining the start, end,

and junction coordinates of each raw transcript by mapping to SARS-CoV-2

(NC_045512.2) and extracting the associated genomic sequence. While this approach

is capable of overcoming sequencing errors and can resolve major RNA species, it is

not capable of resolving small or structurally complex RNA rearrangements. Because

the 5' end of nanopore reads may be truncated due to technical factors during

sequencing (19), we leveraged the assumption that functional sgRNAs should contain

the 5' leader sequence to identify reads generated from complete transcripts. While

excluding reads that don't contain the leader sequence may have reduced sensitivity, it

prevents identification of variant ORFs as a consequence of truncated reads.

ORFs were predicted directly from the coordinate-derived transcripts, and ORFs of

unexpected length were identified as "variant". Assessment of the relative abundance of

variant and canonical ORFs revealed evidence of variant ORFs of ORF1a/b, S, and M

(**Figure 6A**).

To characterize the identified M, S, and ORF1a ORF variants, we determined the stop

and start points of each ORF variant relative to their canonical counterparts (**Figure 6D-**

L). This analysis revealed that all theoretical M ORF variants were predicted to start at a

UUG start codon 42 amino acids before the end of the M gene and end at the canonical

stop codon (**Figure 6D, G, J**). M ORF variants make up between 7.9% and 19% of M

ORFs in the three dRNAseq datasets and are generated by the major junction point at

the ORF6 TRS sequence within the M reading frame (**Figure 1D-F**). If translated, M

ORF variants would lack the N-terminal transmembrane domains and contain regions

predicted to be present on the inside of the virion or in the cytosol. While the efficacy of

translation initiation from this start codon is unknown, it has an acceptable Kozak

consensus site with an A at the -3 position.

S ORF variants make up between 5.1% and 24.2% of all S ORFs identified in these

datasets (**Figure 6A-C**). The majority of S ORF variants are predicted from nc-sgRNAs

with 3' junctions landing within the S ORF (**Supplementary Figure 2D-F**), resulting in

N-terminal truncations with theoretical AUG and non-AUG start codons (**Figure 6E, H,**

K). If translated, these variants would lack the N-terminal signal sequence (20) and,

especially considering the variations in ORF start sites, it is unclear if these ORFs

encode functional isoforms.

Our identification of ORF1a variants is consistent with the observed patterns of 5'

ORF1a coverage (**Figure 5**). The peak end position of identified ORF1a variant ORFs is

around amino acid position 500 (**Figure 6F, I, L**), consistent with the observed decrease

in read coverage around genome position 1600. This position falls within the NSP2

amino acid sequence suggesting that, if translated, many ORF1a ORF variants would

consist of NSP1 and N-terminal fragments of NSP2. Interestingly, we found that

approximately 1/3 of ORF1a variants are in-frame fusions with downstream ORFs,

encoding mostly N (**Figure 6M-O**).

Discussion

While prior studies have identified nc-sgRNAs in SARS-CoV-2 (8, 11) and other

coronaviruses (17), there has been no comprehensive analysis of SARS-CoV-2 nc-

sgRNAs. We conducted an integrative analysis of eight independent SARS-CoV-2

transcriptomes to catalogue unexpected, non-canonical RNA species. We show that

SARS-CoV-2 produces nc-sgRNAs, that these nc-sgRNAs increase in abundance over

time, and that they are not associated with TRS-like homology. We highlight strong

evidence supporting the existence of nc-sgRNAs containing only the 5' region of ORF1a

and show that the presence of nc-sgRNAs can alter the landscape of viral ORFs.

We found that canonical sgRNAs are consistently produced across SARS-CoV-2

transcriptomes, and identify eight major sgRNAs encoding S, ORF3a, E, M, ORF6,

ORF7a/b, ORF8, and N. Each of these sgRNAs is supported by abundant 3' junctions

originating at a TRS-B just upstream the associated ORF. Similar to other reports, we

do not find evidence of a major series of sgRNAs encoding ORF10 (7, 8, 11). We also

do not find strong evidence of a major group of sgRNAs encoding ORF7B despite the

presence of a TRS-core-like sequence (A**A**GAAC) proximal to the ORF7B start.

Interestingly, up to 33% of sgRNAs from infected cells *in vitro* are non-canonical. One

potential source of nc-sgRNAs are DVGs introduced by defective particles during

infection. These can accumulate during the propagation of virus stocks. While

information on the passaging of viral stocks is only available for four transcriptomes, at

least five of the six SARS-CoV-2 transcriptomes generated from cell lines were

generated following a low MOI infection, which may reduce the impact of defective

particles (21, 22). Indeed, while there is some variation in the absolute abundance of

non-canonical junctions across transcriptomes that may reflect dataset-specific factors,

the relative abundance of non-canonical junctions at most genomic positions is

generally similar. One exception is that dataset-specific junctions are found in the Finkel

et al. transcriptomes. In particular, there is a strong 5' junction point at position 16971

that represents over 12% of non-canonical junctions in two Finkel et al. replicates but is

largely absent in other transcriptomes. A hypothesis that could explain this observation

is that there is a population of defective particles in the Finkel et al. virion preparation

that contain an internal deletion. This DVG, which is smaller than the full genome, may

have a replicative advantage during virus stock preparation (16). Indeed, the Finkel et

al.-specific junction at 16971 displays a relatively large increase in abundance over

time, in contrast to smaller and non-specific increases of non-canonical junctions at

other positions.

The consistent presence of non-canonical junctions across the SARS-CoV-2 genome in

independent transcriptomes suggests that the bulk of the observed nc-sgRNAs are not

dataset-specific but are generalizable to SARS-CoV-2 biology. Furthermore, their

nonspecific increase in abundance over time *in vitro* supports a model in which nc-

sgRNAs are generated nonspecifically and at higher frequencies later in infection. We

found that non-canonical junctions are not associated with TRS-like homology, raising

the possibility that there is another mechanism behind their formation. For example, it is

possible that factors such as RNA structure (23), accumulation of viral proteins, or

actions of the viral RdRp and associated cofactors (24) could influence nc-sgRNA

formation. Indeed, a recent study of the murine betacoronavirus MHV suggests that the

3'-to-5' exoribonuclease Nsp14 influences recombination site selection of the viral RdRp

(24). Additional studies are necessary to gain a mechanistic understanding of the

factors that influence nc-sgRNA generation and to understand the phenotypic effects of

nc-sgRNAs on SARS-CoV-2 pathogenesis. Notably, considering that defective viral

genomes in negative-sense RNA viruses have been associated with antiviral immunity,

dendritic cell maturation, and interferon production (5, 16, 25, 26) it will be important to

assess if SARS-CoV-2 nc-sgRNAs are associated with immune activity.

Junctions in viral RNA change the landscape of viral ORFs. Because the ORF6 TRS-B

is located 151 bases into the M ORF, sgRNAs created using this TRS-B contain a 3'

region of M. There is a UUG located 41 codons prior to the end of M that will now be

situated prior to the ORF6 on these sgRNAs. While UUG is a relatively weak start

codon, there is potential that this UUG could stimulate translation of a C-terminal M

fragment or regulate the expression of the downstream ORF6 as an upstream ORF

(27). Additional studies are needed to understand if this M ORF variant is translated and

to understand if its UUG start codon influences ORF6 translation.

In addition to this M ORF variant, we found that some nc-sgRNAs contain S and ORF1a

ORF variants. Many of the S ORF variants would lack the N-terminal signal sequence

(28) and, especially considering the variation in the hypothetical start positions of these

variants, it is unclear if they are translated. Notably, subgenomic RNAs encoding variant

S ORFs have been reported for SARS-CoV-1 (29).

In contrast, the ORF1a ORF variants have a peak stop position around ORF1a amino

acid position 500, which correlates well with an observed decrease in ORF1a 5'

coverage around genome position 1600. There is strong support for the existence of 5'

ORF1a-containing nc-sgRNAs across all eight transcriptomes studied here. Although

there isn't a strong peak of 5' junctions at specific positions in the 5' region of ORF1a,

the consistent decreases in coverage and concomitant increases in 5' junctions around

positions 1600 and 6000 are notable. It is possible that there are biological factors at

these positions that mediate these generalized inflection points but, unlike TRS-

mediated homology, don't concentrate junctions to a specific base. As posited by Kim et

al. (8), the existence of subgenomic RNAs with the 5' end of ORF1a raises the

possibility that these nc-sgRNAs may influence the relative abundance of Nsps, making

this observation an important target for future study.

Conclusions

In conclusion, we show that SARS-CoV-2 generates nc-sgRNAs *in vitro* and *in vivo*, and

that these nc-sgRNAs are largely consistent across independent transcriptomes. We

show that nc-sgRNAs are not associated with TRS-like homology, suggesting there may

be a homology-independent mechanism driving their formation. Finally, we show that

non-canonical junctions have the potential to influence the landscape of viral ORFs, and

that there is especially strong support for nc-sgRNAs that contain only a 5' portion of

ORF1a. Future studies are necessary to understand the mechanisms behind nc-sgRNA

formation, and to understand the influence of nc-sgRNAs on SARS-CoV-2

pathogenesis.

Materials and Methods

Sequencing data

Details on all studied transcriptomes are listed in the table below:

Source	Virus Isolate (MOI) (Details on stock generation)	Infection host	Timepoint	Accession or DOI	Technology
Taiaroa et al. (7)	Australia/VIC01/2020 (MOI: Unknown) (Passage unknown)	Vero cells	Unknown	SRR11350376	dRNAseq
Kim et al. (8)	Korea/KCDC02/2020 (MOI: 0.05) (4th passage. No plaque isolation)	Vero cells	24hpi	10.17605/OSF.IO/8F6N9	dRNAseq
Davidson et al. (9)	England/VE6-T/2020 (MOI: Unknown) (2nd passage. No plaque isolation).	Vero cells	Unknown	10.5281/zenodo.3722580	dRNAseq
Finkel et al. (12)	GISAID Acc. No. EPI_ISL_406862 (MOI: 0.2) (4th passage. No plaque isolation.)	Vero cells	5hpi	SRR11713354	Illumina PolyA
Finkel et al. (12)	GISAID Acc. No. EPI_ISL_406862 (MOI: 0.2) (4th passage. No plaque isolation.)	Vero cells	24hpi	SRR11713362	Illumina PolyA
Blanco-Melo et al. (13)	USA/WA1/2020 (MOI: 0.2) (Passage unknown)	A549-ACE2 cells	24hpi	SRR11517741	Illumina PolyA
Blanco-Melo et al. (13)	USA/WA1/2020 (5*10^4 PFU/ferret) (Passage unknown)	Ferret (Nasal washings)	3dpi	SRR11517855-SRR11517858	Illumina PolyA
Suzuki et al. (15)	SARS-CoV-2/Hu/DP/Kng/19-020 (5*10^4 PFU/100 organoids) (Passage unknown but **plaque isolated**)	Bronchial organoids	5dpi	SRR11811022	Illumina PolyA
Emanuel et al. (14)	Frankfurt strain AY310120.1 **OR** BetaCoV/Munich/BavPat1/2020|EPI_ISL_406862 (MOI: 0.33) (Passage unknown)	Calu3	4hpi	SRR11550047	Illumina Total RNA
Emanuel et al. (14)	Frankfurl strain AY310120.1 **OR** BetaCoV/Munich/BavPat1/2020|EPI_ISL_406862 (MOI: 0.33) (Passage unknown)	Calu3	24hpi	SRR11550045	Illumina Total RNA

Viehweger et al. (17)	CoV-229E (MOI: 3) (Passage unknown)	Huh7 cells	24hpi	ERR3460961	dRNAseq

Hpi: Hours post infection

Dpi: Days post infection

In the case of the Finkel et al. and Emanual et al. transcriptomes, the 24hpi samples

were used in all analyses except for the investigation of junction changes over time.

Determining junction coordinates.

Reads in all transcriptomes were mapped against the SARS-CoV-2 reference genome

(accession NC_045512.2) using minimap2 (30) and custom parameters detailed in

minimap_sars2.sh and inspired by Kim et al. (8), and unmapped reads were discarded.

The exceptions are the transcriptomes from Finkel et al. and Emanuel et al. – because

their read lengths were very short (60 and 72 bases in length respectively), mapping

was conducted with STAR (31) using settings detailed in star.sh.

The 5' and 3' junction positions of junction-spanning reads were determined with

generate_synthetic_transcripts.py. This script reads a BAM file and the SARS-CoV-2

genome fasta and, by parsing the BAM alignment CIGAR and start position, determines

the start, end, and junction coordinates of the alignment. Any minimap2-called deletion

or intron of at least 1000 bases in length was considered a junction. Junction

coordinates were then written to a tab-delimited output file for further analysis.

Analysis and visualization of junctions.

The 5' and 3' coordinates of each junction were processed in R (32), and figures

generated with ggplot2 (33). Gene-maps were generated using the gggenes R package

(34).

For the global distribution of reads in each transcriptome (**Figure 1A-C,**

Supplementary Figure 1A-F), specific 5'-3' junctions that occur at least twice are

represented as arcs. Each inverted peak represents a histogram of 5' or 3' junctions at

each genome position, with a bin size of 100 bases – this means that the magnitude of

each peak represents the total number of 5' or 3' junctions within each 100 base span.

Code: global_junctions.Rmd

The distribution of major 3' junctions (**Figure 1D-F**) was generated by plotting a

histogram of 3' junction positions that occur after position 21000 and that have a 5' end

prior to position 100. Bin-size is 20 bases. Code: canonical_3prime_junctions.Rmd

The percentage of non-canonical junctions in each transcriptome (**Figure 2A,**

Supplementary Figure 1G) was calculated by first assigning each individual 5'-3'

junction as canonical or non-canonical. A junction was considered to be canonical if it's

5' position was within 20 bases of the TRS-L core sequence ACGAAC (starting at

position 69), and it's 3' junction was within 15 bases of a TRS-B (defined by the

presence of the TRS core sequence). The percentage of total junctions that were

assigned as non-canonical was then determined. Code: global_junctions.Rmd

Identifying dataset-specific junctions (Figure 2).

The consistency of non-canonical junctions across transcriptomes (**Figure 2C, D**) was

determined separately for 5' and 3' junctions based on one replicate each from Taiaroa

et al., Kim et al., Davidson et al., Finkel et al., and Blanco-Melo et al. transcriptomes

generated from cells infected with SARS-CoV-2 *in vitro*. For each SARS-CoV-2 genome

position, the percentage of non-canonical junctions at that site was calculated for each

transcriptome (X). The mean (μ) and standard deviation (σ) of percentages at each site

across the five transcriptomes were then calculated. For each position in each

transcriptome, a z-score was calculated by subtracting the observed percentage by the

mean and dividing by the standard deviation ((X-μ)/σ). This process is visualized in

Figure 2B. Code: dataset_specific_junctions.Rmd

Change in junctions over time (**Figure 3**).

The change in junctions over time was determined based on datasets from Finkel et al.

(5hpi and 24hpi) and Emanuel et al. (4hpi and 24hpi). The percentage of junctions that

originate at each genome position was determined for each timepoint, and the absolute

difference between the two values was calculated. Notably, this analysis (**Figure 3B, C**)

focused on 5' and 3' counts at each position separately, such that a percentage of 5'

junctions and a percentage of 3' junctions at each position was determined. To

determine the aggregated change in each <u>class</u> of junctions (**Figure 3F,G**), each 5'-3'

junction was assigned to a class based on the location of its 5' and 3' position. If a

junction's 5' position was located within 20 bases of the TRS-L, and it's 3' position was

located within 15 bases of a TRS-B, it was assigned to the ORF closest to the TRS-B.

Otherwise, the junction was assigned as non-canonical. The percentage of total

junctions in each class at the early timepoint was subtracted from the class percentages

at the late timepoint for each dataset. Code: junctions_over_time.Rmd

Assessing homology near 5' and 3' positions of each junction (**Figure 4**).

The identity and length of homologous sequences flanking the 5' and 3' sites of each

junction were determined by identifying the 30 bases flanking each junction site. The

longest identical subsequence between these two 30 base sequences was then

collected. For **Figure 4B-D**, the most common homology sequence was plotted for each

group of junctions. Here, a junction was considered canonical if it has a 3' end within 15

bases of a canonical TRS-B or was assigned as "Within ORF" if it was otherwise within

an ORF. The exception were junctions with a 5' end within ORF1a – these were

assigned to ORF1a. The total distribution of homology lengths was also determined for

canonical and non-canonical junctions (**Figure 4E-G**). To determine the random

distribution of homology lengths, all 30 base sequences (30mers) present in the SARS-

CoV-2 genome were determined. The longest identical subsequence between 100,000

random pairs of 30mers that are derived from at least 1000 bases apart on the genome

was determined. Code: TRS_plotting.Rmd

Assessment of excess 5' ORF1a genome coverage (Figure 5).

Coverage at each position was assessed from SARS-CoV-2-mapped BAMs using the

depth subcommand of samtools (35). (Command:

samtools depth -aa -d0 file.bam > file.cov). Coverage and cumulative number of 5'

junctions (excluding junctions occurring before position 100) were plotted. Code:

ORF1a_coverage.Rmd

Analysis of ORFs on dRNAseq reads (Figure 6).

generate_synthetic_transcripts.py was used to generate "coordinate-derived transcripts"

from the SARS-CoV-2 mapped BAMs from the Taiaroa et al., Kim et al., and Davidson

et al. dRNAseq transcriptomes. This script reads a BAM file and the SARS-CoV-2

genome fasta and, by parsing the BAM alignment CIGAR and start position, determines

the start, end, and junction coordinates of the alignment. Coordinate-derived transcripts

were created by using these coordinates to retrieve the associated sequence from the

SARS-CoV-2 genome. Any minimap2-called deletion or intron of at least 1000 bases in

length was considered a junction. All coordinate-derived transcripts were then aligned

against a minimal leader sequence fragment (5' –

CTTCCCAGGTAACAAACCAACCAACTTTCGATCTCTTGTAGATCTGTTCTC – 3')

using blastn (36). This minimal sequence was used instead of the full leader sequence

because inspection of 5' coverage in the three long-read datasets revealed a significant

drop in coverage at the end of this sequence. Transcripts with an alignment of at least

45 nucleotides in length and a percent identity of at least 85%, and that contain the start

of this sequence within the first 50 nucleotides of the transcript, were kept. Transcripts

not containing this leader sequence were discarded.

ORFs were called directly from each transcript using Prodigal (37), using parameters

listed in prodigal.sh. ORFs were allowed to initiate from any nUG start codon, were not

required to be the most 5' ORF on each transcript and were only reported if they are at

least 30 codons in length. Each ORF was translated into amino acid sequence using

prodigal_to_orf_direct.py. If multiple Prodigal-predicted ORFs contained an overlapping

amino acid sequence, the longest ORF was output. Each ORF was output in fasta

format and labeled with transcript of origin and coordinates of the ORF on the transcript.

The amino acid sequence of each ORF was then mapped against canonical SARS-

CoV-2 proteins (from accession NC_045512.2) as well as Prodigal-predicted

unannotated ORFs from the SARS-CoV-2 genome using DIAMOND (38). DIAMOND

parameters are described in diamond.sh and include an E-value threshold of 10. During

downstream analysis, alignments were considered valid if the aligned portions

maintained at least 95% amino acid identity to the reference protein.

Various statistics of each ORF, including alignment information, the start position of the

ORF on the transcript, and fusion information, were generated from the DIAMOND

alignment file using parse_orf_assignments.py. An ORF was considered variant if it was

not the same length as the canonical protein, with a notable exception. Because there

are alternative start codons upstream of some annotated genes, and Prodigal would call

ORFs using these start codons if they are present on a transcript, ORFs were allowed

to have up to 20 amino acids of additional N-terminal amino acids while still being

classified as canonical if they end at the expected position. If an ORF had matches

against multiple SARS-CoV-2 proteins, it was given a primary assignment of the protein

with the longest alignment length.

Code availability

The reproducible computational pipeline used to generate all results is publicly available

at https://github.com/jnoms/virORF_direct.

List of abbreviations

CoV: Coronavirus

dRNAseq: Direct RNA sequencing

E: Envelope protein

M: Matrix protein

N: Nucleocapsid protein

Nsp: Nonstructural protein

ORF: Open reading frame

S: Spike protein

SARS-CoV-2: Systemic acute respiratory syndrome Coronavirus 2

TRS: Transcriptional regulatory sequence

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: Code is available in the virORF_direct repository,

https://github.com/jnoms/virORF_direct.

Competing interests

M.M. receives research support from Bayer, Ono, and Janssen, has patents licensed to

Bayer and Labcorp, and is a consultant for OrigiMed. J.A.D. received research support

from Constellation Pharmaceuticals and is a consultant for EMD Serono, Inc. and Merck

& Co. Inc.

Funding

This work was supported in part by the US Public Health Service grants R35CA232128

and P01CA203655 to J.A.D.

Authors' Contributions

J.N. analyzed the data; J.N., M.M., J.A.D. designed the study and wrote the manuscript.

Acknowledgements

The authors thank Taiaroa et al., Kim et al., Davidson et al., Blanco-Melo et al., Suzuki

et al., Finkel et al., and Emanuel et al. for collecting the RNA sequencing data and

making these data publicly available. Portions of this research were conducted by using

the O2 High Performance Compute Cluster, supported by the Research Computing

Group at Harvard Medical School.

Figure legends

Figure 1. SARS-CoV-2 generates a defined population of canonical subgenomic

RNAs.

A-C) For each location on the viral genome, a histogram of 5' and 3' junctions at that

position was calculated and plotted as an inverse peak. The histogram bin size is

100 bases, meaning each inverse peak represents the cumulative count of 5' or 3'

junctions occurring within that span. Curved lines represent the 5' and 3' locations of

junctions that occur at least twice. Red curves represent canonical junctions, black

curves represent non-canonical junctions. "Taiaroa" (7), "Kim" (8), and "Davidson"

(9) represent the three independent SARS-CoV-2 dRNAseq datasets investigated.

D-F) A histogram of 3' junctions past position 21000 with a 5' end before position

100 are plotted. Dashed lines indicate the start coordinates of annotated viral genes.

Bin-size is 20 bases.

G) Based on junction analysis, the predicted major species of virus-produced RNAs

are represented. The most 5' gene or genes on each subgenomic RNA is listed.

Figure 2. Non-canonical junctions are consistent across independent datasets.

A) Percentage of junctions that are non-canonical in five independent datasets.

Junctions were assigned as canonical if their 5' location was within 20 bases of

the TRS-L and their 3' location within 15 bases of a TRS-B, and otherwise

assigned as non-canonical. Taiaroa, Kim, and Davidson are dRNAseq, while

Finkel and Blanco-Melo are Illumina PolyA RNAseq.

B) Illustration of computational approach to determine the consistency of 5' and

3' junction points across independent datasets. The percentage of non-canonical

junctions at each genome position in each dataset was determined (X). The

mean (µ) and standard deviation (σ) of percentages at each position across the

five datasets was calculated. For each position in each dataset, the Z-score was

calculated as (X-µ)/σ (i.e. the number of standard deviations away from the

mean), and the percentages and Z-scores for each position in each dataset were

plotted.

C, D) Each point represents the percentage and Z-score of one position in one

dataset. For each position in the SARS-CoV-2 genome, the percentage and Z-

score of non-canonical junctions with a 5' end (D) or a 3' end (E) was determined

as described above for five independent datasets: Taiaroa, Kim, Davidson,

Finkel, and Blanco-Melo. Points with a percentage above 4% of non-canonical

junctions and a Z-score above 1 were highlighted.

Figure 3. Non-canonical junctions accumulate over time *in vitro*.

A) Illustration of computational approach used to determine change in junction

percentages over time. For each position in the SARS-CoV-2 genome and

separately for 5' and 3' junctions, the number of junctions at that position was

calculated. Using these numbers, the percentage of 5' or 3' junctions falling at

each position was calculated. The change in junction percentage at each position

is defined as the difference between the position's junction percentage at the late

and early timepoints, and this junction change was plotted for each position.

B-E) The change in junction percentage for 5' (orange) and 3' (blue) junctions

over time in the Finkel (B, D) and Emanuel (C, E) datasets was determined as

described above. Changes in the percentage of junctions Positions with at

change greater than 2.5% are annotated with text on the plot. Panels D and E

are zoomed in versions of B and C.

F, G) Junctions were assigned into groups based on their 5' and 3' junction

positions. If a junction had a 5' end within 20 bases of the TRS-L and within 15

bases of a TRS-B, it was considered a canonical junction belonging to the ORF

with a start closest to the TRS-B. Otherwise, it was considered non-canonical.

The percentage of junctions falling into each category was calculated for early

and late timepoints, and the difference between each category's percentage in

the late vs early timepoint was plotted.

Figure 4. Non-canonical junctions are not associated with TRS-like homology.

A) Illustration of computational approach used to assess homologous sequences

between the 5' and 3' junction points. For each of the three dRNAseq datasets,

the 30 bases flanking the 5' and 3' junction points were assessed, and the

longest homologous sequence between these two 30 base regions was

determined.

B-D) Homologous sequences present in the 15 bases on either side of the 5' and

3' ends of each junction. Junctions are classified by the location of their 3' end –

if this is within 15 bases of the canonical TRS site or if it falls within a gene it is

assigned accordingly. The only exception is ORF1a – junctions with a 5' end

originating in ORF1a are assigned to ORF1a. Labels represent the most

common homologous sequence between the ends of each junction. The core

TRS sequence ACGAAC is underlined.

E-G) The length of the longest homologous sequence for canonical (C) and non-

canonical (NC) are plotted. Here, junctions were considered canonical if they

have a 5' end within 20 nucleotides of the TRS-L and a 3' end within 15

nucleotides of a TRS-B. (R) represents random homology lengths. The points for

(R) were calculated by first extracting all possible 30 base sequences (30mers)

from the SARS-CoV-2 genome, and then assessing the length of the longest

homologous sequence between 100,000 random pairs of 30mers that are

separated by at least 1000 bases on the genome. Within each column, the

relative width of each band represents the relative abundance of junctions with

each homology length. The value at the top of each column is the mean

homology length.

Figure 5. Relative abundance of subgenomic RNAs that contain only the 5' end of

ORF1a.

A-H) Read coverage (black) and cumulative 5' junctions (red) are plotted for

eight independent datasets. The sequencing strategy and sample types are

annotated.

I) A schematic of a representative subgenomic RNA that consists of only the 5'

region of ORF1a. Pileups showing reads can be found in Supplementary Figure

3.

Figure 6. Junctions have the potential to generate variant open reading frames.

A-C) ORFs were predicted directly from transcript-derived reads for the three

dRNAseq datasets. Each ORF was aligned against the protein sequences of

canonical SARS-CoV-2 genes using the DIAMOND aligner. Variant ORFs were

defined as ORFs that were assigned to a canonical SARS-CoV-2 protein but had

an unexpected start or stop position, while perfectly-aligning ORFs were

considered canonical. The percentage of canonical and variant ORFs for each

protein is plotted.

D-L) Schematics of M, N, and ORF1ab are displayed with the approximate

location of predicted transmembrane domains labeled in red. A histogram of the

start and end sites of variant M, N, and ORF1ab ORFs are displayed. Start sites

of each variant ORF are on top, and end sites are on the bottom of each panel.

Percentages represent the percentage of each ORF that is variant. Histogram

bin-sizes: M, 1; N, 10 ; ORF1ab, 20. NTD: N-terminal domain. RBD: Receptor

binding domain. CD: Connector domain. NSP: Nonstructural protein.

M-O) The identity of ORF1a-fusion partners is plotted on the Y axis, with the

count of such fusions on the X axis. The top 10 fusion partners for each sample

are represented. Color indicates if the fusion partner is on the N and C terminus

of ORF1a, if the terminus is ambiguous, or if the fusion is a "self" fusion between

an upstream and downstream region of ORF1a. The percentage notes the

percent of ORFs assigned to ORF1a that have a fusion.

Supplementary Figure 1. There is evidence of non-canonical junctions in diverse

datasets.

A-E) For each location on the viral genome, a histogram of 5' and 3' junctions at

that position was calculated and plotted as an inverse peak. The histogram bin

size is 100 bases, meaning each inverse peak represents the cumulative count

of 5' or 3' junctions occurring within that span. Curved lines represent the 5' and

3' locations of junctions that occur at least twice. Red curves represent canonical

junctions, black curves represent non-canonical junctions.

F) The percentage of junctions that are canonical in the Suzuki (Bronchial

organoids) and Blanco-Melo (Ferret) datasets was determined. Junctions were

assigned as canonical if their 5' location was within 20 bases of the TRS-L and

their 3' location within 15 bases of a TRS-B, and otherwise assigned as non-

canonical.

Supplementary Figure 2. Dataset-specific junctions

A-E) The percentage of non-canonical junctions at each SARS-CoV-2 genome

position is plotted. Plots A-C show the percentage of 5' junctions at each

position, while D-E show the percentage of 3' junctions at each position. The

plotted samples include three replicates from Blanco-Melo et al., Davidson et al,

two replicates from Finkel et al, and the Kim and Taiaroa et al datasets.

Supplementary Figure 3. Subgenomic RNAs containing non-canonical junctions

are present in three independent dRNAseq datasets.

A-F) Subgenomic RNAs with 5' junctions occurring in ORF1a (A-C) or 3'

junctions occurring in S (D-F) are plotted for the three dRNAseq datasets. Each

row is a read, and regions in black indicate the read contains that sequence.

Supplementary Figure 4. There is elevated ORF1a coverage in CoV 229E.

A) Read coverage (black) and cumulative 5' junctions (red) are plotted for

dRNAseq data from CoV 229E-infected cells.

732   References

1.   Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. A new coronavirus
associated with human respiratory disease in China. Nature. 2020;579(7798):265-9.
2.   Plant EP, Dinman JD. The role of programmed-1 ribosomal frameshifting in
coronavirus propagation. Frontiers in bioscience: a journal and virtual library.
2008;13:4873.
3.   Narayanan K, Huang C, Makino S. SARS coronavirus accessory proteins. Virus
research. 2008;133(1):113-21.
4.   de Haan CA, Masters PS, Shen X, Weiss S, Rottier PJ. The group-specific
murine coronavirus genes are not essential, but their deletion, by reverse genetics, is
attenuating in the natural host. Virology. 2002;296(1):177-89.

5.   Yount B, Roberts RS, Sims AC, Deming D, Frieman MB, Sparks J, et al. Severe
acute respiratory syndrome coronavirus group-specific open reading frames encode
nonessential functions for replication in cell cultures and mice. Journal of virology.
2005;79(23):14909-22.
6.   Sawicki SG, Sawicki DL, Siddell SG. A contemporary view of coronavirus
transcription. Journal of virology. 2007;81(1):20-9.
7.   Taiaroa G, Rawlinson D, Featherstone L, Pitt M, Caly L, Druce J, et al. Direct
RNA sequencing and early evolution of SARS-CoV-2. bioRxiv.
2020:2020.03.05.976167.
8.   Kim D, Lee J-Y, Yang J-S, Kim JW, Kim VN, Chang H. The architecture of
SARS-CoV-2 transcriptome. Cell. 2020.
9.   Davidson AD, Williamson MK, Lewis S, Shoemark D, Carroll MW, Heesom KJ, et
al. Characterisation of the transcriptome and proteome of SARS-CoV-2 reveals a cell
passage induced in-frame deletion of the furin-like cleavage site from the spike
glycoprotein. Genome Medicine. 2020;12(1):1-15.
10.   Kim D, Lee J-Y, Yang J-S, Kim JW, Kim VN, Chang H. The architecture of
SARS-CoV-2 transcriptome. bioRxiv. 2020:2020.03.12.988865.
11.   Davidson AD, Williamson MK, Lewis S, Shoemark D, Carroll MW, Heesom K, et
al. Characterisation of the transcriptome and proteome of SARS-CoV-2 using direct
RNA sequencing and tandem mass spectrometry reveals evidence for a cell passage
induced in-frame deletion in the spike glycoprotein that removes the furin-like cleavage
site. bioRxiv. 2020:2020.03.22.002204.
12.   Finkel Y, Mizrahi O, Nachshon A, Weingarten-Gabbay S, Yahalom-Ronen Y,
Tamir H, et al. The coding capacity of SARS-CoV-2. bioRxiv. 2020:2020.05.07.082909.
13.   Blanco-Melo D, Nilsson-Payant BE, Liu W-C, Uhl S, Hoagland D, Møller R, et al.
Imbalanced host response to SARS-CoV-2 drives development of COVID-19. Cell.
2020.
14.   Emanuel W, Kirstin M, Vedran F, Asija D, Theresa GL, Roberto A, et al. Bulk and
single-cell gene expression profiling of SARS-CoV-2 infected human cell lines identifies
molecular targets for therapeutic intervention. bioRxiv. 2020:2020.05.05.079194.
15.   Suzuki T, Itoh Y, Sakai Y, Saito A, Okuzaki D, Motooka D, et al. Generation of
human bronchial organoids for SARS-CoV-2 research. bioRxiv.
2020:2020.05.25.115600.
16.   Genoyer E, López CB. The impact of defective viruses on infection and immunity.
Annual review of virology. 2019;6:547-66.
17.   Viehweger A, Krautwurst S, Lamkiewicz K, Madhugiri R, Ziebuhr J, Hölzer M, et
al. Direct RNA nanopore sequencing of full-length coronavirus genomes provides novel
insights into structural variants and enables modification analysis. Genome research.
2019;29(9):1545-54.
18.   Rang FJ, Kloosterman WP, de Ridder J. From squiggle to basepair:
computational approaches for improving nanopore sequencing read accuracy. Genome
biology. 2018;19(1):90.
19.   Workman RE, Tang AD, Tang PS, Jain M, Tyson JR, Razaghi R, et al. Nanopore
native RNA sequencing of a human poly (A) transcriptome. Nature Methods.
2019;16(12):1297-305.

20.   Bosch BJ, Martina BE, Van Der Zee R, Lepault J, Haijema BJ, Versluis C, et al.
Severe acute respiratory syndrome coronavirus (SARS-CoV) infection inhibition using
spike protein heptad repeat-derived peptides. Proceedings of the National Academy of
Sciences. 2004;101(22):8455-60.
21.   Thompson KAS, Yin J. Population dynamics of an RNA virus and its defective
interfering particles in passage cultures. Virology journal. 2010;7(1):257.
22.   Brooke CB. Biological activities of'noninfectious' influenza A virus particles.
Future virology. 2014;9(1):41-51.
23.   Lan TCT, Allan MF, Malsick LE, Khandwala S, Nyeo SSY, Bathe M, et al.
Structure of the full SARS-CoV-2 RNA genome in infected cells. bioRxiv.
2020:2020.06.29.178343.
24.   Gribble J, Pruijssers AJ, Agostini ML, Anderson-Daniels J, Chappell JD, Lu X, et
al. The coronavirus proofreading exoribonuclease mediates extensive viral
recombination. bioRxiv. 2020:2020.04.23.057786.
25.   Sun Y, Jain D, Koziol-White CJ, Genoyer E, Gilbert M, Tapia K, et al.
Immunostimulatory defective viral genomes from respiratory syncytial virus promote a
strong innate antiviral response during infection in mice and humans. PLoS Pathog.
2015;11(9):e1005122.
26.   Strahle L, Garcin D, Kolakofsky D. Sendai virus defective-interfering genomes
and the activation of interferon-beta. Virology. 2006;351(1):101-11.
27.   Barbosa C, Peixeiro I, Romão L. Gene expression regulation by upstream open
reading frames and human disease. PLoS Genet. 2013;9(8):e1003529.
28.   Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, et al. Cryo-
EM structure of the 2019-nCoV spike in the prefusion conformation. Science.
2020;367(6483):1260-3.
29.   Hussain S, Chen Y, Yang Y, Xu J, Peng Y, Wu Y, et al. Identification of novel
subgenomic RNAs and noncanonical transcription initiation signals of severe acute
respiratory syndrome coronavirus. Journal of virology. 2005;79(9):5288-95.
30.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34(18):3094-100.
31.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
32.   Team RC. R: A language and environment for statistical computing. 2013.
33.   Wickham H. ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics.
2011;3(2):180-5.
34.   Wilkins D. gggenes: draw gene arrow maps in 'ggplot2'. R package version 0.4.
0. 2019.
35.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence
alignment/map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
36.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.
BLAST+: architecture and applications. BMC bioinformatics. 2009;10(1):421.
37.   Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
bioinformatics. 2010;11(1):119.
38.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using
DIAMOND. Nature methods. 2015;12(1):59.

Figure 1

Figure 1. SARS-CoV-2 generates a defined population of canonical subgenomic RNAs.

A-C) For each location on the viral genome, a histogram of 5' and 3' junctions at that position was calculated and plotted as an inverse peak. The histogram bin size is 100 bases, meaning each inverse peak represents the cumulative count of 5' or 3' junctions occurring within that span. Curved lines represent the 5' and 3' locations of junctions that occur at least twice. Red curves represent canonical junctions, black curves represent non-canonical junctions. "Taiaroa", "Kim", and "Davidson" represent the three independent SARS-CoV-2 dRNAseq datasets investigated.

D-E) A histogram of 3' junctions past position 21000 with a 5' end before position 100 are plotted. Dashed lines indicate the start coordinates of annotated viral genes. Bin-size is 20 bases

G) Based on junction analysis, the predicted major species of virus-produced RNAs are represented. The most 5' gene or genes on each subgenomic RNA islisted.

Figure 2

Figure 2. Non-canonical junctions are consistent across independent datasets.

A) Percentage of junctions that are non-canonical in five independent datasets. Junctions were assigned as canonical if their 5' location was within 20 bases of the TRS-L and their 3' location within 15 bases of a TRS-B, and otherwise assigned as non-canonical. Taiaroa, Kim, and Davidson are dRNAseq, while Finkel and Blanco-Melo are Illumina PolyA RNAseq.

B) Description of computational approach to determine the consistency of 5' and 3' junction points across independent datasets. The percentage of non-canonical junctions at each genome position in each dataset was determined (X). The mean (μ) and standard deviation (σ) of percentages at each position across the five datasets was calculated. For each position in each dataset, the Z-score was calculated as $(X-\mu)/\sigma$ (i.e. the number of standard deviations away from the mean), and the percentages and Z-scores for each position in each dataset were plotted.

C, D) Each point represents the percentage and Z-score of one position in one dataset. For each position in the SARS-CoV-2 genome, the percentage and Z-score of non-canonical junctions with a 5' end (D) or a 3' end (E) was determined as described above for five independent datasets: Taiaroa, Kim, Davidson, Finkel, and Blanco-Melo. Points with a percentage above 4% of non-canonical junctions and a Z-score above 1 were highlighted.

Figure 3

Figure 3. Non-canonical junctions accumulate over time in vitro.

A) Description of computational approach used to determine change in junction percentages over time. For each position in the SARS-CoV-2 genome and separately for 5' and 3' junctions, the number of junctions at that position was calculated. Using these numbers, the percentage of 5' or 3' junctions falling at each position was calculated. The change in junction percentage at each position is defined as the difference between the position's junction percentage at the late and early timepoints, and this junction change was plotted for each position.

B-E) The change in junction percentage for 5' (orange) and 3' (blue) junctions over time in the Finkel (B, D) and Emanuel (C, E) datasets was determined as described above. Changes in the percentage of junctions Positions with at change greater than 2.5% are annotated with text on the plot. Figures D and E are zoomed in versions of A and B.

F, G) Junctions were assigned into groups based on their 5' and 3' junction positions. If a junction had a 5' end within 20 bases of the TRS-L and within 15 bases of a TRS-B, it was considered a canonical junction belonging to the ORF with a start closest to the TRS-B. Otherwise, it was considered non-canonical. The percentage of junctions falling into each category was calculated for early and late timepoints, and the difference between each category's percentage in the late vs early timepoint was plotted.

Figure 4. Non-canonical junctions are not associated with TRS-like homology.

A) Description of computational approach used to assess homologous sequences between the 5' and 3' junction points. For each of the three dRNAseq datasets, the 30 bases flanking the 5' and 3' junction points were assessed, and the longest homologous sequence between these two 30 base regions was determined.

B-D) Homologous sequences present in the 15 bases on either side of the 5' and 3' ends of each junction. Junctions are classified by the location of their 3' end – if this is within 15 bases of the canonical TRS site or if it falls within a gene it is assigned accordingly. The only exception is ORF1a – junctions with a 5' end originating in ORF1a are assigned to ORF1a. Labels represent the most common homologous sequence between the ends of each junction. The core TRS sequence ACGAAC is underlined.

E-G) The length of the longest homologous sequence for canonical (C) and non-canonical (NC) are plotted. Here, junctions were considered canonical if they have a 5' end within 20 nucleotides of the TRS-L and a 3' end within 15 nucleotides of a TRS-B. (R) represents random homology lengths. The points for (R) were calculated by first extracting all possible 30 base sequences (30mers) from the SARS-CoV-2 genome, and then assessing the length of the longest homologous sequence between 100,000 random pairs of 30mers that are separated by at least 1000 bases on the genome. Within each column, the relative width of each band represents the relative abundance of junctions with each homology length. The value at the top of each column is the mean homology length.

Figure 5

Figure 5. There is evidence of subgenomic RNAs that contain only the 5' end of ORF1a.
A-H) Read coverage (black) and cumulative 5' junctions (red) are plotted for eight independent datasets. The sequencing strategy and sample types are annotated.

I) A schematic of a representative subgenomic RNA that consists of only the 5' region of ORF1a. Pileups showing reads can be found in Supplementary Figure 3.

Figure 6. Junctions have the potential to generate variant open reading frames.

A-C) ORFs were predicted directly from transcript-derived reads for the three dRNAseq datasets, and each ORF was aligned against the protein sequences of canonical SARS-CoV-2 genes using the DIAMOND aligner. Variant ORFs were defined as ORFs that were assigned to a canonical SARS-CoV-2 protein but had an unexpected start or stop position, while perfectly-aligning ORFs were considered canonical. The percentage of canonical and variant ORFs for each protein is plotted.

D-L) Schematics of M, N, and ORF1ab are displayed with the approximate location of predicted transmembrane domains labeled in red. A histogram of the start and end sites of variant M, N, and ORF1ab ORFs are displayed. Start sites of each variant ORF are on top, and end sites are on the bottom of each panel. Percentages represent the percentage of each ORF that is variant. Histogram bin-sizes: M, 1; N, 10 ; ORF1ab, 20. NTD: N-terminal domain. RBD: Receptor binding domain. CD: Connector domain. NSP: Nonstructural protein.

M-O) The identity of ORF1a-fusion partners is plotted on the Y axis, with the count of such fusions on the X axis. The top 10 fusion partners for each sample are represented. Color indicates if the fusion partner is on the N and C terminus of ORF1a, if the terminus is ambiguous, or if the fusion is a "self" fusion between an upstream and downstream region of ORF1a. The percentage notes the percent of ORFs assigned to ORF1a that have a fusion.

Supplementary Figure 1

Supplementary Figure 1. There is evidence of non-canonical junctions in diverse datasets.

A-E) For each location on the viral genome, a histogram of 5' and 3' junctions at that position was calculated and plotted as an inverse peak. The histogram bin size is 100 bases, meaning each inverse peak represents the cumulative count of 5' or 3' junctions occurring within that span. Curved lines represent the 5' and 3' locations of junctions that occur at least twice. Red curves represent canonical junctions, black curves represent non-canonical junctions.

F) The percentage of junctions that are canonical in the Suzuki (Bronchial organoids) and Blanco-Melo (Ferret) datasets was determined. Junctions were assigned as canonical if their 5' location was within 20 bases of the TRS-L and their 3' location within 15 bases of a TRS-B, and otherwise assigned as non-canonical.

Supplementary Figure 2

Supplementary Figure 2. There are dataset-specific junctions in Finkel et al.

    A-E) The percentage of non-canonical junctions at each SARS-CoV-2 genome position is plotted. Plots A-C show the percentage of 5' junctions at each position, while D-E show the percentage of 3' junctions at each position. The plotted samples include three replicates from Blanco-Melo et al., Davidson et al, two replicate from Finkel et al, and the Kim and Taiaroa et al datasets.

Supplementary Figure 3

Supplementary Figure 3. Subgenomic RNAs containing non-canonical junctions are present in three independent dRNAseq datasets.

A-F) Subgenomic RNAs with 5' junctions occurring in ORF1a (A-C) or 3' junctions occurring in S (D-F) are plotted for the three dRNAseq datasets. Each row is a read, and regions in black indicate the read contains that sequence.

Supplementary Figure 4

A

Supplementary Figure 4. There is elevated ORF1A coverage in CoV 229E.
    A) Read coverage (black) and cumulative 5' junctions (red) are plotted for dRNAseq data from
    CoV 229E-infected cells.