Specificity of Gβ and γ subunits to the SNARE complex both at rest and after α[2a] adrenergic receptor stimulation

Yun Young Yim[1†], W. Hayes McDonald[2], Katherine M. Betke[1,5], Ali Kaya[1,6], Karren Hyde[1,7], Kevin Erreger[1], Ralf Gilsbach[3, 4], Lutz Hein[4], and Heidi E. Hamm[1*]

[1]Department of Pharmacology, Vanderbilt University, Nashville TN 37232-6600
†Currently at Nash Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY 10029
[2]Department of Biochemistry and Mass Spectrometry Research Center, Vanderbilt University, Nashville TN 37232-6600
[3]Fachbereich Medizin, Institute for Cardiovascular Physiology, Goethe University, Theodor-Stern-Kai 7, 60590, Frankfurt, Germany
[4]Institute of Experimental and Clinical Pharmacology and Toxicology, Faculty of Medicine, University of Freiburg, 79104, Freiburg, Germany
[5]Currently at Department of Anesthesiology and Perioperative Medicine, Oregon Health & Science University, Portland, Oregon, 97239
[6]Currently at Department of Chemistry and Chemical Biology, Cornell University Address, APS 9700 S. Cass Ave. Bldg. 436E Lemont, IL 60439
[7]Currently at Department of Cell and Developmental Biology, Vanderbilt University, Nashville TN 37232-6600
*Correspondence to heidi.hamm@vanderbilt.edu

Keywords: Gbeta, Ggamma, specificity, adrenergic receptor, SNARE, synaptic transmission, MRM

Summary

Specific Gβγ dimers interact with the SNARE complex following presynaptic α[2a]AR activation

in both adrenergic and non-adrenergic neurons.

Abstract

Though much is known about the various physiological functions of each GPCR and the specificity of Gα subunits, the specificity of G$\beta\gamma$ activated by a given GPCR and activating each effector in vivo is not known. Previously, we identified different Gβ and Gγ subunits interacting specifically with α_{2a}-adrenergic receptors (α_{2a}AR). In this study, we examined its in vivo specificity to the soluble NSF attachment proteins (SNARE) complex in adrenergic (auto-α_{2a}AR) and non-adrenergic (hetero-α_{2a}AR) neurons. We applied a quantitative targeted multiple reaction monitoring proteomic analysis of Gβ and Gγ subunits bound to the SNARE complex, and found only a subset of Gβ and Gγ bound. Without stimulation of auto-α_{2a}AR, Gβ_1 and Gγ_3 interacted with the SNARE complex. When auto-α_{2a}AR were activated, Gβ_1, Gβ_2, and Gγ_3 were found. Further understanding of in vivo G$\beta\gamma$ specificity to its effectors provides new insights into the multiplicity of genes for Gβ and Gγ.

Introduction

Numerous studies have demonstrated that different Gβ and Gγ subunits may selectively interact to form unique Gβγ dimers (*1-3*), which in turn target specific G protein-coupled receptors (GPCRs) and effectors of Gβγ (*1, 3-11*). Although many of these Gβγ-effector interactions and downstream signaling cascades have been well studied in vitro, factors controlling the specificity of Gβγ dimers associated with unique GPCRs or their effectors, as well as the in vivo expression abundance of Gβ and Gγ subunits or Gβγ dimers remain unclear. Only a few in vivo genetic deletion studies and knout-out animal studies have attempted to identify the specific roles of different Gβ and Gγ subunits(*11, 12*). With the recent development of targeted multiple reaction monitoring (MRM) proteomic method to detect each neuronal Gβ and Gγ subunits, our lab has showen the protein abundance and subcellular localization of each neuronal Gβ and Gγ subunits (*13*) and their specificity to α_{2a}-adrenergic receptors in adrenergic neurons (auto-α_{2a}AR) and non-adrenergic neurons (hetero-α_{2a}AR)(*14*). Only a subset of neuronal Gβ and Gγ subunits interact with auto-α_{2a}AR and hetero-α_{2a}AR. However, it remains unclear how G protein specificity plays out in α_{2a}AR-mediated effector interactions.

α_{2a}-adrenergic receptors (α_{2a}AR) are one of the best studied $G_{i/o}$ coupled GPCRs, given the availability of both knock out (α_{2a}AR KO) mice and transgenic mice with tagged receptors (FLAG-α_{2a}AR) (*15-19*). Briefly, the activation of α_{2a}AR releases GTP-$G\alpha_{i/o}$ and Gβγ, and these interact with various effectors to initiate the downstream signaling cascades(*11*). FLAG-α_{2a}AR transgenic mice express only auto-α_{2a}AR under the control of the dopamine-β-hydroxylase (Dbh) promoter while α_{2a}AR KO mice are the endogenous knockout of α_{2a}AR(*16*). Using these mice, auto-α_{2a}AR are found to be important for bradycardia and hypotension while α_{2a}AR in hetero-α_{2a}AR are involved in anesthetic sparing, hypothermia, analgesia, bradycardia, and hypotension

(*16*). However, it is unclear how G protein specificity, especially Gβγ, plays out in α_{2a}AR-mediated downstream signaling cascades and underlies such diverse physiological functions. Given the physiological importance of α_{2a}AR, and the different roles of auto-and hetero-α_{2a}AR, the signaling mechanisms of α_{2a}AR in both adrenergic and non-adrenergic neurons, especially in relation to the specificity of Gβγ, need to be further elucidated.

Soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE complex), Gβγ effector, is a key protein involved in synaptic transmission, essential for inter-neuronal communication in both the nervous and endocrine systems. The neuronal SNARE complex, comprised of SNAP25, syntaxin 1A, and synaptobrevin (VAMP) (*20-24*), brings the vesicle membrane and plasma membrane into close proximity, resulting in vesicle fusion at the plasma membrane (*25*). After membrane fusion, neurotransmitter is released into the synaptic junction, permitting communication between neurons encoding both external and internal sensory inputs, and promoting responsive behaviors in postsynaptic cells (*24*). Thus, it is crucial to understand how synaptic transmission is regulated in healthy subjects, and mechanisms of dysfunction in neurological diseases (*26-28*).

The activation of α_{2a}AR results in the inhibition of synaptic transmission via the Gβγ subunit in both adrenergic (auto-α_{2a}AR) and non-adrenergic neurons (hetero-α_{2a}AR) such as glutamatergic, GABAergic, serotonergic and dopaminergic neurons (*29-32*). In particular, numerous studies have reported that Gβγ subunits released from activated α_{2a}AR reduce presynaptic Ca^{2+} influx through the inhibition of voltage-gated calcium channels (VDCC) (*33-35*) and inhibit exocytosis through direct interaction with the SNARE complex (*36-38*). Furthermore, the direct interaction between Gβy and the SNARE complex upon α_{2a}AR activation is a primary mechanism to inhibit exocytosis of norepinephrine-containing vesicles in the central amygdala

(*38*). Delaney *et al*. show that inhibition of norepinephrine release, where high abundance of presynaptic α_{2a}AR are present, is mediated by a $G_{i/o}$-coupled receptor, and that G$\beta\gamma$ dimers released from active α_{2a}AR directly interact with the SNARE complex downstream of calcium entry to inhibit norepinephrine release (*38*). In addition, our knout-out study with SNAP25Δ3 mice no longer shows the inhibition of adrenergic signaling in the BNST, a key area for the integration of stress signals (*39*). The loss of 3 C-terminal residues in SNAP25 causes a loss of G$\beta\gamma$`s ability to regulate α_{2a}AR mediated inhibition of synaptic transmission. Although the G$\beta\gamma$–SNARE complex interaction has been well-characterized during α_{2a}AR-mediated inhibition of synaptic transmission, the specificity of G$\beta\gamma$ dimer isoforms for the SNARE complex is not known.

Together, we use various biochemical techniques and tools such as co-immunoprecipitation (coIP), quantitative MRM proteomic method, and transgenic animals to understand the specificities of Gβ and Gγ subunits to the SNARE complex. Here, we evaluated which G$\beta\gamma$ dimers released from epinephrine (epi)-activated α_{2a}AR interact with SNARE complex to regulate synaptic transmission. We discovered that only a subset of Gβ and Gγ subunits released from activated α_{2a}AR interact with the SNARE complex. Further understanding of α_{2a}AR-mediated G$\beta\gamma$ subunit specific interaction with the SNARE complex may lead to the development of new therapeutic targets that complement or contrast with currently available α_{2a}AR-targeted drugs.

Results

G$\beta\gamma$ - SNARE complex interaction. To study G$\beta\gamma$ specificity for the SNARE complex upon activation of α_{2a}AR, we used synaptosomes collected from the brains of wildtype, α_{2a}AR KO,

and FLAG-α_{2a}AR mice. Using a SNAP25 antibody (S9684) and DSP, a lipid-soluble thiol cleavable crosslinker, G$\beta\gamma$ and the SNARE complex were co-immunoprecipitated (coIP) and detected by Western blot (Fig. 1A). We showed that the SNAP25 (S9684) antibody selectively recognizes SNAP25 (fig.S1). To further validate that we were indeed coIPing the ternary SNARE complex, we detected Syntaxin1 in addition to SNAP25 from these coIP samples (fig. S2). The G$\beta\gamma$ and SNARE complex interaction was found in both unstimulated and stimulated coIP samples, but greater amounts were present in epi-stimulated coIP samples of wildtype and FLAG-α_{2a}-AR mice (Fig.1B to E). We know from a number of studies (*40, 41*) that there is a low affinity for interaction between G$\beta\gamma$ and the SNARE complex. However, the interaction is specific, as little non-specific binding was observed in IgG controls. Quantitative western blot analysis estimated approximately 1µg G$\beta\gamma$ was pulled down with SNAP25 per half brain (fig. S3). Together, these results suggest that this method is suitable to determine which G$\beta\gamma$ subunits associate with the SNARE complex in vivo.

Gβ_1, Gβ_2, Gβ_4, Gγ_2, Gγ_3, and Gγ_4 interact specifically with the SNARE complex upon epinephrine stimulation. To examine which neuronal Gβ and Gγ subunits interact with the SNARE complex upon activation of α_{2a}-AR, we applied a quantitative MRM method (*14*) to TCA precipitated and trypsin digested SNAP25 coIP samples of wildtype (WT), α_{2a}AR KO, and FLAG-α_{2a}AR mouse synaptosomes. Previously, only a subset of neuronal Gβ and Gγ subunits are selectively associated with α_{2a}AR(*14*). WT samples were used to detect the G$\beta\gamma$–SNARE complex interactions mediated by epi stimulation, including both auto- and hetero-α_{2a}AR, while α_{2a}AR KO samples were used to detect non α_{2a}AR-mediated interactions only. Lastly, FLAG-α_{2a}AR samples were used to detect presynaptic auto-α_{2a}AR-mediated G$\beta\gamma$-SNARE complex interactions. We used unstimulated WT samples (WT no epi), α_{2a}AR KO (α_{2a}AR KO no epi),

and FLAG-α_{2a}AR (FLAG-α_{2a}ARs no epi) as controls to detect basal G$\beta\gamma$–SNARE complex interactions. By comparing the amount of Gβ and Gγ subunits detected per mg of lysate between stimulated WT (WT +epi), α_{2a}AR KO (α_{2a}AR KO +epi), and FLAG-α_{2a}AR (FLAG-α_{2a}AR +epi), we detected not only auto-receptor mediated G$\beta\gamma$-SNARE complex interactions, but also quantify those mediated by non-α_{2a}AR and estimated heteroreceptor interactions.

To identify nonspecific interactions of Gβ and Gγ subunits, we used IgG alone controls (without primary antibody). We compared the amount of Gβ subunit detected per mg of lysate between WT IgG +epi and WT +epi samples and found Gβ_1, Gβ_2, and Gβ_4 significantly interacted with the SNARE complex upon α_{2a}AR stimulation (Fig. 2). As expected from previous studies (*13, 14*), Gβ_1 was detected at a significantly higher concentration than Gβ_2 and Gβ_4 subunits. Moreover, we did not identify interaction between Gβ_5 and the SNARE complex following α_{2a}AR activation.

Next, we identified the Gγ subunits that interact with the SNARE complex following receptor activation. We found enrichment of the Gγ_2, Gγ_3, and Gγ_4 subunits with the SNARE complex upon epi stimulation (Fig.3). Gγ_3 was the most abundant followed by Gγ_2 and Gγ_4, respectively. We detected a similar abundance of Gγ_2 and Gγ_4 associated with the SNARE complex. We were not able to detect an interaction between Gγ_{12} and the SNARE complex (Fig. 3). Although Gγ_{12} is specifically bound to α_{2a}AR (*14*), it may not interact with the SNARE complex but rather with other G$\beta\gamma$ effectors such as VDCCs (see Discussion). Even though we are not detecting specific G$\beta\gamma$ dimers in this study, according to our data (Fig. 2 and 3), we detect only a subset of possible G$\beta\gamma$ dimers and can postulate 9 different combinations of G$\beta\gamma$ dimers (G$\beta_1\gamma_2$, G$\beta_1\gamma_3$, G$\beta_1\gamma_4$, G$\beta_2\gamma_2$, G$\beta_2\gamma_3$, G$\beta_2\gamma_4$, G$\beta_4\gamma_2$, G$\beta_4\gamma_3$, and G$\beta_4\gamma_4$) which might interact

with the SNARE complex upon epi stimulation. These dimers may occur in the brain and play roles in the α_{2a}AR mediated G$\beta\gamma$-SNARE complex interactions. Based on the detected abundances of Gβ_1 and Gγ_3, we suggest that G$\beta_1\gamma_3$ is the most likely dimer to interact with the SNARE complex following receptor stimulation. Further biochemical analysis will be needed to validate the presence of these G$\beta\gamma$ dimers and their specificities for the SNARE complex upon α_{2a}AR activation in both adrenergic and non-adrenergic neurons.

Basal interaction of G$\beta\gamma$ and the SNARE complex. We showed in Fig 1B and C that G$\beta\gamma$ and the SNARE complex interact basally in unstimulated coIPed samples of wildtype and FLAG-α_{2a}-AR mice. To understand the basal G$\beta\gamma$–SNARE complex interaction in the absence of epi stimulation, we compared IgG control to no epi coIPed samples from α_{2a}AR KO and FLAG-α_{2a}AR synaptosomes using MRM mass spectrometry. α_{2a}AR KO samples were used to identify non-α_{2a}AR-mediated basal interactions while FLAG-α_{2a}AR samples were used to detect auto-α_{2a}AR-mediated G$\beta\gamma$-SNARE complex interactions. Here, we detected all neuronal Gβ subunits but found only Gβ_1 significantly interacting with the SNARE complex in both α_{2a}AR KO and FLAG-α_{2a}AR no epi conditions compared to that of the IgG no epi condition (Fig.4). This suggests that there is a basal interaction of Gβ_1 with the SNARE complex. The abundance of Gβ_1 between these two conditions was not different, suggesting that the basal interaction of Gβ_1 and the SNARE complex may not be α_{2a}AR mediated. Similarly, we detected Gγ_3 interaction with the SNARE complex in the unstimulated conditions of both genotypes compared to IgG control (Figure 5). Thus we suggest that the G$\beta_1\gamma_3$ dimer is a basal, non-α_{2a}AR mediated interaction. Various GPCRs have been implicated to mediate presynaptic inhibition through this mechanism; therefore, other endogenous GPCRs present in the synaptosomes could be causing

this effect, or alternatively, since no agonists of other GPCRs were present in the experiments, it may be a basal interaction with SNARE complex

Gβ_1, Gβ_2, and Gγ_3 interact specifically with the SNARE complex upon auto-α_{2a}ARs activation. As previously described, adrenergic auto-α_{2a}AR inhibits presynaptic neurotransmitter release through the interaction between G$\beta\gamma$ and the SNARE complex (*38*). To test whether auto-α_{2a}ARs and presynaptic hetero-α_{2a}AR utilize different G$\beta\gamma$ dimers upon activation, we coIPed samples from α_{2a}AR KO and FLAG-α_{2a}AR synaptosomes and again applied the quantitative MRM method to understand auto-α_{2a}ARs mediated G$\beta\gamma$-SNARE complex interactions. α_{2a}AR KO (no epi) and FLAG-α_{2a}AR (no epi) samples were used as controls to identify non-auto-α_{2a}AR mediated interactions, and α_{2a}AR KO +epi samples were used to detect non-α_{2a}AR-mediated G$\beta\gamma$-SNARE complex interactions. By comparing the result from α_{2a}AR KO, FLAG-α_{2a}AR, and wildtype, we can propose the specificity of Gβ and Gγ interactions with the SNARE complex mediated by hetero-α_{2a}AR.

Here, we were able to detect a subset of Gβ and Gγ subunit interactions with the SNARE complex upon auto-α_{2a}AR activation. When we compared the results from α_{2a}AR KO +epi and FLAG-α_{2a}AR +epi synaptosomes, we identified that Gβ_1 and Gβ_2, but not Gβ_4, enrich with the SNARE complex following auto-α_{2a}AR activation (Fig.6). We also detected significantly more Gβ_1 than Gβ_2 (Fig.6). Moreover, we detected a ~ 10-fold increased interaction of Gβ_1 and the SNARE complex upon epi stimulation compared to the basal state (compare Fig. 4 and 6).

Among the 3 Gγ subunits enriched with the SNARE complex upon stimulation of WT containing auto- and hetero-α_{2a}AR and non-α_{2a}AR (Fig. 3), the enrichment of Gγ_3 with the SNARE complex was detected following auto-α_{2a}AR activation (Fig. 7). Basally, without epi

stimulation, $G\gamma_3$ was found to interact with the SNARE complex (Fig 5). These findings suggest that $G\beta_1\gamma_3$ may be basally interacting with the SNARE complex and not α_{2a}ARs mediated (Fig. 4 and 5), whereas $G\beta_2\gamma_3$ may interact with the SNARE complex only after epi stimulation of auto-α_{2a}AR.

Discussion

Numerous in vitro studies have studied the effectors of $G\beta\gamma$ and their downstream signaling cascades(*1-4, 42*); however, only a few in vivo genetic deletion studies and knout-out animals studies have examined the specific roles of different $G\beta$ and $G\gamma$ subunits (*11, 12*). It remains unclear how particular $G\beta\gamma$ dimers form or how they localize to particular sites in vivo. In this paper, we addressed the in vivo specificity of $G\beta\gamma$ subunit interaction with the SNARE complex. We determined a surprising specificity in which $G\beta$ and γ subunits associate with the SNARE complex, including the interesting finding that some $G\beta\gamma$s interact with the SNARE complex regardless of receptor activation. Only a subset of neuronal $G\beta\gamma$ dimers—$G\beta_1$, $G\beta_2$, $G\beta_4$, $G\gamma_2$, $G\gamma_3$, and $G\gamma_4$—interact with the SNARE complex (Fig. 2 and 3) following epi stimulation in WT mice. Upon the activation of auto-α_{2a}AR, only $G\beta_1$, $G\beta_2$, and $G\gamma_3$ interacted with the SNARE complex (Fig. 6 and 7). From 9 possible $G\beta\gamma$ dimers, auto-α_{2a}AR may utilize only two $G\beta\gamma$ dimers, $G\beta_1\gamma_3$ and $G\beta_2\gamma_{3,}$ to inhibit neurotransmitter release through the $G\beta\gamma$-SNARE complex interaction, In addition, comparing WT and auto-α_{2a}AR samples (Fig.2 and 6), we found that $G\beta_4\gamma_2$ and $G\beta_4\gamma_4$ may be specifically utilized by presynaptic hetero-α_{2a}AR. Thus, this study provides a window into the specificity of $G\beta\gamma$ dimer function in vivo.

In both Western blot and quantitative MRM study in FLAG-α_{2a}AR animals, we detected $G\beta\gamma$-SNARE complex interactions in the basal state, without α_{2a}AR stimulation (Fig.1C, 4, and

5). Because we showed previously that the α_{2a}AR does not basally interact with G$\beta\gamma$ (*14*) and no basal G$\beta\gamma$ interaction to the SNARE complex was detected in the quantitative MRM study of wildtype animals (Fig. 2), we were surprised to see a basal interaction of G$\beta_1\gamma_3$ with the SNARE complex (Fig.4 and 5). With the limited tools, we cannot evaluate and understand the observed difference of basal interaction in wildtype and FLAG-α_{2a}AR animals However, the ~4-fold increase of abundance in Gβ and ~20-fold increase of abundance in Gγ observed in FLAG-α_{2a}AR animals raise an interacting hypothesis. The change in G$\beta\gamma$ associated to the SNARE complex may occur upon the activation of auto-α_{2a}AR. Further studies are necessary to understand the basal interaction of G$\beta_1\gamma_3$ with the SNARE complex, especially in the wildtype animal G$\beta_2\gamma_3$ appears only associated with the SNARE complex upon auto-α_{2a}AR activation.

Gβ_1 specifically interacts with the SNARE complex not only in the basal state and but also upon epi stimulation in FLAG-α_{2a}AR mice (Fig.4 and 6). The abundance of Gβ_1 interacting with the SNARE complex in WT samples (Fig. 2) was expected as Gβ_1 is the most abundant Gβ subunit in whole synaptosome as well as at both pre-and post-synaptic fractions(*13*). However, an increase of abundance in Gβ_1 (~10 fold) interacting with the SNARE complex upon auto-α_{2a}AR activation was not expected (Fig.6). This suggests that the basally bound Gβ_1 may have a unique role in basal modulation of exocytosis. So far, the role of this basal G$\beta\gamma$ association with the SNARE complex is not known, but it is well-known that basal exocytotic events (measured electrophysiologically as mini EPSCs) can be regulated by G$_{i/o}$ GPCR agonists(*43, 44*).

Combining this study of the G$\beta\gamma$ specificity to the SNARE complex with our previous studies of the presynaptic abundance specificity of G$\beta\gamma$ subunits for α_{2a}AR(*13, 14*), we hypothesize that the G$\beta\gamma$ subunit specificity may be encoded by an association of a particular

GPCR with a subset of Gβγs, –the abundance of each neuronal Gβ and Gγ subunit, and the distance to their effectors at the presynaptic terminal. In our previous receptor study comparing the protein abundance detected with α_{2a} receptors, we speculated that $G\gamma_2$, $G\gamma_3$, and $G\gamma_4$ were associated with auto-α_{2a}AR while $G\gamma_{12}$ is specific for hetero-α_{2a}AR (*14*). Given our current understanding with this study on the Gβ and Gγ specificity to the SNARE complex, we can more accurately show that $G\gamma_2$ and $G\gamma_4$ are involved in hetero-α_{2a}AR-mediated Gβγ-SNARE complex interaction, with no evidence for $G\gamma_{12}$ association to the SNARE complex. $G\gamma_{12}$ associated with hetero-α_{2a}AR may be associated with another Gβγ effector.

The Gγ subunits, more than Gβ, play key roles in Gβγ specificity. Previously, knock-down and knock-out studies showed that particular Gγ subunits have distinct physiological importance in biological systems (*11, 12*). To date, only 4 out of 12 Gγ subunits have been studied using knockdown or knock-out models. Mainly expressed in retina, $G\gamma_1$ is involved in photoreceptor degeneration. $G\gamma_2$ is involved in nociception and angiogenesis. $G\gamma_3$ is involved in increased susceptibility to seizures and resistance to diet induced obesity while $G\gamma_7$ affects the startle response(*4, 7*). Unique physiological phenotypes of each Gγ subunit suggest a great deal of specificity in Gβγ dimerization and signaling(*45-47*). The main molecular mechanisms behind binding selectivity between various Gβγs and their effectors are still unknown. However, ongoing investigations of interactions between each Gγ and t-SNARE are suggesting the importance of charged residues during this process(*48*).

Moreover, the fast rate of Gβγ-SNARE complex-mediated inhibition(*44*) suggests that Gβγ, the SNARE complex, and other synaptic proteins may be associated by scaffolding interactions prior to receptor activation, to facilitate Gβγ-SNARE complex interaction and

mediate Gβγ specificity. Previously, we identified two Gβγ binding sites on vesicle proximal and distal regions of SNAP25 (*41, 49, 50*), far enough apart to not both be occupied by a single Gβγ. However, the physiological and biochemical importance of these interaction sites still remains unclear. The two sites could underlie interaction via the SNARE complex with two different effectors (*51, 52*) or the amino terminal region may be a Gβγ binding (or "holding") site that can hand Gβγ off to the physiologically important carboxy-terminal interaction site (*50*). Using SNAP25Δ3 mice with a truncated C-terminus of SNAP25, we see a two-fold reduction in Gβγ-SNARE complex interaction, as well as an increase in stress-induced hyperthermia, defective spatial learning, impaired gait, supraspinal nociception (*39*), enhancement of long-term potentiation at Schaffer collateral-CA1 synapse (*53*), and resistance to diet-induced obesity(*54*). It will be interesting to determine the stoichiometry of Gβγ binding to the SNARE complexes, and whether identical Gβγ dimers bind to both amino- and carboxy-terminal sites of the SNARE complexes. Detailed quantitative analysis in combination with SNAP25 mutagenesis studies will be necessary to answer these questions.

The specificity of Gβγ dimers to the SNARE complex is not fully understood, but its importance and the relationship of GPCRs to this interaction have been well-studied. Hamid *et al*. demonstrated that $5HT_{1b}$ and $GABA_B$ receptors inhibit exocytotic release at the same synaptic terminal through different mechanisms (*43*), with $5HT_{1b}$ receptor-mediated inhibition of exocytosis using the Gβγ-SNARE complex interaction, while $GABA_B$ receptor-mediated inhibition works through Gβγ-VDCC interaction and decreased calcium entry. The authors suggested that this difference may depend on the N-terminal SNAP25 residues. The association of specific Gβγ subunits to GPCRs, such as $5HT_{1b}$ and $GABA_B$, and to the SNARE complex may

add a further layer of regulation of Gβγ-effector selectivity. Future studies to understand this specificity will clarify Gβγ isoform specificity important for both receptor and effector binding.

Materials and Methods

Animals. Adult, male FLAG-tagged alpha2a adrenergic receptors (α_{2a}-AR), α_{2a}-AR knock-out (KO), and wildtype mice (*55*) were decapitated and brain tissues were immediately homogenized to produce crude synaptosomes as described below. To minimize post-mortem differences, all tissues were processed in parallel. All animal handling and procedures were conducted in accordance with the Care and Use of Laboratory Animals of the National Institutes of Health and approved by the Vanderbilt Institutional Animal Care and Use Committee.

Drugs. Epinephrine (catalog E4642), prazosin (catalog P7791), and propranolol (catalog P0884) were purchased from Sigma-Aldrich.

Antibodies. For immunoprecipitation, rabbit anti-SNAP25 (Sigma, S9684) and rabbit ChromePure IgG (Jaskson Immuno Research, 011-030-003) were used. For the western blot analysis, mouse anti-SNAP25 (Santa Cruz, sc-376713, 1:500) and rabbit anti-Gβ (Santa Cruz, sc-378, 1:10,000 and 1:5000) were used. HRP-conjugated secondary antibodies were obtained from Perkin-Elmer and Jackson Immunoresearch and used at the following dilutions: goat anti-mouse (1:10,000), and mouse anti-rabbit light chain specific (1:7,500).

Synaptosome preparation, stimulation, and lysate protocol. Crude synaptosomes were isolated from mouse brain tissue, as described previously (*14, 56-58*) in the presence of 1mM ethylenediaminetetraacetic acid, 1uM propranolol, 1uM prazosin, and 2mM 3,3'-dithiobis [sulfosuccinimydlypropinate], the lipid soluble, thiol cleavable crosslinker (Pierce, 22585) and stimulated with 100µM epinephrine (epi). This mimics the local synaptic concentration of epi and it is a commonly used concentration in α_{2a}AR studies(*59-61*). As epi is not an α2aAR

selective agonist, we also used prazosin and propranolol to prevent off-target effects from non-α2aAR adrenergic receptors. Samples were frozen in liquid nitrogen and stored at -80°C.

Co-immunoprecipitation. Similar to the co-immunoprecipitation (coIP) of α_{2a}AR and G$\beta\gamma$ (*14*), we coIPed the SNARE complex and G$\beta\gamma$ from the precleared lysates (Fig.1A). Precleared lysates were aliquoted 400ul per tube. Instead of anti-HA or FLAG antibodies, they were incubated with an anti-SNAP25 coIP antibody. Because homozygous SNAP25 knock-out mice exhibit neonatal lethally(*62*) , we used rabbit IgG as a control.

Elution and trichloroacetic acid (TCA) precipitation of coIPs. Samples (approximately 11 coIPs per condition per genotype) were eluted using 100mM glycine (pH 2.5). 40µL of glycine was added to coIP beads and placed on a room temperature vortex shaker at medium speed for 8 mins and centrifuged at 5,000 x g for 2mins to pellet beads. Elutants were transferred to clean 1.5mL Eppendorf tubes. The elution was repeated a second time, and the elutants were pooled together. For western blot analysis, one tube per condition per genotype was saved separately Remaining elutants were pooled per genotype per condition and TCA precipitated to concentrate the Gβ and Gγ subunits for MRM (*14*). All samples were stored at -80°C freezer for the western blot or MRM analysis.

Immunoblot Analysis. The analysis was done as described previously (*14*). We used 15%, instead of 10% SDS-PAGE gel.

Heavy labeled peptide cocktail. A heavy labeled peptide cocktail was made as described previously (*13*).

Quantitative MRM of Gβ and Gγ subunits. As described previously (*14*), coIP samples containing Gβ and Gγ subunits were separated by SDS-PAGE, in-gel trypsin digested, and analyzed by scheduled multiple reaction monitoring (MRM) using a TSQ vantage triple

quadrupole mass spectrometer (Thermo Scientific) coupled directly to a nanoAcquity liquid chromatography system (Waters).

Statistical analysis. One-way analysis of variance (ANOVA) with a Tukey post hoc test was used to account for differences in protein abundance of Gβ and Gγ subunits and for the quantification of western blots (* $p<0.05$, ** $p<0.01$, *** $p<0.001$). All statistical tests were performed using GraphPad Prism v.9.0 for Windows, (GraphPad Software, La Jolla, California, USA, www.graphpad.com).

Supplementary Materials

Figure 1. Selectivity of SNAP25 IP Antibody

Figure 2. Western blot analysis of SNAP25 coIP.

Figure 3. Quantification of Gβ precipitated with SNAP25.

Reference

1.   A. V. Smrcka, G protein betagamma subunits: central mediators of G protein-coupled receptor signaling. *Cellular and molecular life sciences : CMLS* **65**, 2191-2214 (2008); published online EpubJul (10.1007/s00018-008-8006-5).
2.   G. J. Stephens, G-protein-coupled-receptor-mediated presynaptic inhibition in the cerebellum. *Trends Pharmacol Sci* **30**, 421-430 (2009); published online EpubAug (10.1016/j.tips.2009.05.008).
3.   J. D. Hildebrandt, Role of subunit diversity in signaling by heterotrimeric G proteins. *Biochemical Pharmacology* **54**, 325-339 (1997)Doi: 10.1016/s0006-2952(97)00269-4).
4.   A. N. Pronin, N. Gautam, Interaction between G-protein beta and gamma subunit types is selective. *Proceedings of the National Academy of Sciences of the United States of America* **89**, 6220-6224 (1992); published online EpubJul 1 (
5.   K. Yan, V. Kalyanaraman, N. Gautam, Differential ability to form the G protein betagamma complex among members of the beta and gamma subunit families. *The Journal of biological chemistry* **271**, 7141-7146 (1996); published online EpubMar 22 (
6.   J. D. Robishaw, C. H. Berlot, Translating G protein subunit diversity into functional specificity. *Curr Opin Cell Biol* **16**, 206-209 (2004); published online EpubApr (10.1016/j.ceb.2004.02.007).
7.   W. F. Schwindinger, K. S. Betz, K. E. Giger, A. Sabol, S. K. Bronson, J. D. Robishaw, Loss of G Protein γ7 Alters Behavior and Reduces Striatal αolf Level and cAMP Production. *Journal of Biological Chemistry* **278**, 6575-6579 (2003); published online EpubFebruary 21, 2003 (10.1074/jbc.M211132200).

8.   W. F. Schwindinger, K. E. Giger, K. S. Betz, A. M. Stauffer, E. M. Sunderlin, L. J. Sim-Selley, D. E. Selley, S. K. Bronson, J. D. Robishaw, Mice with Deficiency of G Protein g3 Are Lean and Have Seizures. *Mol. Cell. Biol.* **24**, 7758-7768 (2004); published online EpubSeptember 1, 2004 (10.1128/mcb.24.17.7758-7768.2004).

9.   W. F. Schwindinger, L. J. Mihalcik, K. E. Giger, K. S. Betz, A. M. Stauffer, J. Linden, D. Herve, J. D. Robishaw, Adenosine A2A receptor signaling and golf assembly show a specific requirement for the gamma7 subtype in the striatum. *The Journal of biological chemistry* **285**, 29787-29796 (2010); published online EpubSep 24 (10.1074/jbc.M110.142620).

10.   W. F. Schwindinger, U. L. Mirshahi, K. A. Baylor, K. M. Sheridan, A. M. Stauffer, S. Usefof, M. M. Stecker, T. Mirshahi, J. D. Robishaw, Synergistic roles for G-protein g3 and g7 subtypes in seizure susceptibility as revealed in double knockout mice. *Journal of Biological Chemistry* **287**, 7121-7133 (2011); published online EpubDecember 29, 2011 (10.1074/jbc.M111.308395).

11.   S. M. Khan, R. Sleno, S. Gora, P. Zylbergold, J. P. Laverdure, J. C. Labbe, G. J. Miller, T. E. Hebert, The expanding roles of Gbetagamma subunits in G protein-coupled receptor signaling and drug action. *Pharmacological reviews* **65**, 545-577 (2013); published online EpubApr (10.1124/pr.111.005603).

12.   N. Gautam, G. B. Downes, K. Yan, O. Kisselev, The G-protein betagamma complex. *Cell Signal* **10**, 447-455 (1998); published online EpubJul (

13.   Y. Y. Yim, W. H. McDonald, K. Hyde, O. Cruz-Rodriguez, J. J. G. Tesmer, H. E. Hamm, Quantitative Multiple-Reaction Monitoring Proteomic Analysis of Gbeta and Ggamma Subunits in C57Bl6/J Brain Synaptosomes. *Biochemistry* **56**, 5405-5416 (2017); published online EpubOct 10 (10.1021/acs.biochem.7b00433).

14.   Y. Y. Yim, K. M. Betke, W. H. McDonald, R. Gilsbach, Y. Chen, K. Hyde, Q. Wang, L. Hein, H. E. Hamm, The in vivo specificity of synaptic Gbeta and Ggamma subunits to the alpha2a adrenergic receptor at CNS synapses. *Sci Rep* **9**, 1718 (2019); published online EpubFeb 8 (10.1038/s41598-018-37222-1).

15.   R. J. Lu, Y. Li, Y. W. Zhang, Y. J. Chen, A. D. Shields, D. G. Winder, T. Angelotti, K. Jiao, L. E. Limbird, Y. Zhou, Q. Wang, Epitope-tagged Receptor Knock-in Mice Reveal That Differential Desensitization of alpha(2)-Adrenergic Responses Is because of Ligand-selective Internalization. *Journal of Biological Chemistry* **284**, 13233-13243 (2009); published online EpubMay 8 (DOI 10.1074/jbc.M807535200).

16.   R. Gilsbach, L. Hein, Are the pharmacology and physiology of alpha(2) adrenoceptors determined by alpha(2)-heteroreceptors and autoreceptors respectively? *British journal of pharmacology* **165**, 90-102 (2012); published online EpubJan (10.1111/j.1476-5381.2011.01533.x).

17.   T. Mizobe, [Adrenergic receptor and knockout mouse: 2). Alpha adrenergic receptor knockout mouse]. *Masui* **50**, 12-19 (2001); published online EpubJan (

18.   J. Lahdesmaki, J. Sallinen, E. MacDonald, B. K. Kobilka, V. Fagerholm, M. Scheinin, Behavioral and neurochemical characterization of alpha(2A)-adrenergic receptor knockout mice. *Neuroscience* **113**, 289-299 (2002).

19.   M. M. Bucheler, K. Hadamek, L. Hein, Two alpha(2)-adrenergic receptor subtypes, alpha(2A) and alpha(2C), inhibit transmitter release in the brain of gene-targeted mice. *Neuroscience* **109**, 819-826 (2002).

20.   R. G. Zhai, H. J. Bellen, Hauling t-SNAREs on the microtubule highway. *Nat Cell Biol* **6**, 918-919 (2004); published online EpubOct (10.1038/ncb1004-918).

21.   S. Schoch, E. D. Gundelfinger, Molecular organization of the presynaptic active zone. *Cell and tissue research* **326**, 379-391 (2006); published online EpubNov (10.1007/s00441-006-0244-y).

22.   T. C. Sudhof, The synaptic vesicle cycle. *Annu Rev Neurosci* **27**, 509-547 (2004)10.1146/annurev.neuro.26.041002.131412).

23.   A. G. Purves D, Fitzpatrick D, Hall WC, LaMantia AS, McNamara JO, White LE, *Neuroscience, 4th Edition*. (Sinauer Associates, Inc, 2008).

24.   T. C. Sudhof, The presynaptic active zone. *Neuron* **75**, 11-25 (2012); published online EpubJul 12 (10.1016/j.neuron.2012.06.012).

25.   N. P. Brose, AG. Sudhof, TC. Jahn, R., Synaptotagmin: a calcium sensor on the synaptic vesicle surface. *Science* **256**, 1021-1025 (1992); published online Epub1992 May 15 (

26.   A. F. Jeans, P. L. Oliver, R. Johnson, M. Capogna, J. Vikman, Z. Molnár, A. Babbs, C. J. Partridge, A. Salehi, M. Bengtsson, L. Eliasson, P. Rorsman, K. E. Davies, A dominant mutation in Snap25 causes impaired vesicle trafficking, sensorimotor gating, and ataxia in the blind-drunk mouse. *Proceedings of the National Academy of Sciences* **104**, 2431-2436 (2007); published online EpubFebruary 13, 2007 (10.1073/pnas.0610222104).

27.   B. Alten, Q. Zhou, O. H. Shin, L. Esquivies, P. Y. Lin, K. I. White, R. Sun, W. K. Chung, L. M. Monteggia, A. T. Brunger, E. T. Kavalali, Role of Aberrant Spontaneous Neurotransmission in SNAP25-Associated Encephalopathies. *Neuron* **109**, 59-72 e55 (2021); published online EpubJan 6 (10.1016/j.neuron.2020.10.012).

28.   A. Ramos-Miguel, C. L. Beasley, A. J. Dwork, J. J. Mann, G. Rosoklija, A. M. Barr, W. G. Honer, Increased SNARE Protein-Protein Interactions in Orbitofrontal and Anterior Cingulate Cortices in Schizophrenia. *Biol Psychiatry* **78**, 361-373 (2015); published online EpubSep 15 (10.1016/j.biopsych.2014.12.012).

29.   L. Hein, Adrenoceptors and signal transduction in neurons. *Cell and tissue research* **326**, 541-551 (2006); published online EpubNov (10.1007/s00441-006-0285-2).

30.   M. I. Forray, G. Bustos, K. Gysling, Noradrenaline inhibits glutamate release in the rat bed nucleus of the stria terminalis: in vivo microdialysis studies. *J Neurosci Res* **55**, 311-320 (1999); published online EpubFeb 01 (10.1002/(SICI)1097-4547(19990201)55:3<311::AID-JNR6>3.0.CO;2-E).

31.   M. J. Millan, F. Lejeune, A. Gobert, Reciprocal autoreceptor and heteroreceptor control of serotonergic, dopaminergic and noradrenergic transmission in the frontal cortex: relevance to the actions of antidepressant agents. *J Psychopharmacol* **14**, 114-138 (2000); published online EpubJun (10.1177/026988110001400202).

32.   R. Mongeau, M. Weiss, C. de Montigny, P. Blier, Effect of acute, short- and long-term milnacipran administration on rat locus coeruleus noradrenergic and dorsal raphe serotonergic neurons. *Neuropharmacology* **37**, 905-918 (1998); published online EpubJul (

33.   A. C. Dolphin, G protein modulation of voltage-gated calcium channels. *Pharmacological reviews* **55**, 607-627 (2003); published online EpubDec (

34.   S. R. Ikeda, Voltage-dependent modulation of N-type calcium channels by G-protein bg subunits. *Nature* **380**, 255-258 (1996); published online EpubMar 21 (10.1038/380255a0).

35.   S. R. Ikeda, K. Dunlap, Voltage-dependent modulation of N-type calcium channels: role of G protein subunits. *Adv Second Messenger Phosphoprotein Res* **33**, 131-151 (1999).

36.   R. Jahn, R. H. Scheller, SNAREs--engines for membrane fusion. *Nature reviews. Molecular cell biology* **7**, 631-643 (2006); published online EpubSep (10.1038/nrm2002).

37.   J. Malsam, S. Kreye, T. H. Sollner, Membrane fusion: SNAREs and regulation. *Cellular and molecular life sciences : CMLS* **65**, 2814-2832 (2008); published online EpubSep (10.1007/s00018-008-8352-3).

38.   A. J. Delaney, J. W. Crane, P. Sah, Noradrenaline modulates transmission at a central synapse by a presynaptic mechanism. *Neuron* **56**, 880-892 (2007); published online EpubDec 6 (DOI 10.1016/j.neuron.2007.10.022).

39.   Z. Zurawski, A. D. Thompson Gray, L. J. Brady, B. Page, E. Church, N. A. Harris, M. R. Dohn, Y. Y. Yim, K. Hyde, D. P. Mortlock, C. K. Jones, D. G. Winder, S. Alford, H. E. Hamm, Disabling the Gbetagamma-SNARE interaction disrupts GPCR-mediated presynaptic inhibition, leading to physiological and behavioral phenotypes. *Sci Signal* **12**, (2019); published online EpubFeb 19 (10.1126/scisignal.aat8595).

40.   Z. Zurawski, B. Page, M. C. Chicka, R. L. Brindley, C. A. Wells, A. M. Preininger, K. Hyde, J. A. Gilbert, O. Cruz-Rodriguez, K. P. M. Currie, E. R. Chapman, S. Alford, H. E. Hamm,

Gbetagamma directly modulates vesicle fusion by competing with synaptotagmin for binding to neuronal SNARE proteins embedded in membranes. *The Journal of biological chemistry* **292**, 12165-12177 (2017); published online EpubJul 21 (10.1074/jbc.M116.773523).

41.   E. J. Yoon, T. Gerachshenko, B. D. Spiegelberg, S. Alford, H. E. Hamm, Gbetagamma interferes with Ca2+-dependent binding of synaptotagmin to the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex. *Molecular pharmacology* **72**, 1210-1219 (2007); published online EpubNov (10.1124/mol.107.039446).

42.   J. Dingus, J. D. Hildebrandt, Synthesis and assembly of G protein betagamma dimers: comparison of in vitro and in vivo studies. *Sub-cellular biochemistry* **63**, 155-180 (2012)10.1007/978-94-007-4765-4_9).

43.   E. Hamid, E. Church, C. A. Wells, Z. Zurawski, H. E. Hamm, S. Alford, Modulation of neurotransmission by GPCRs is dependent upon the microarchitecture of the primed vesicle complex. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **34**, 260-274 (2014); published online EpubJan 1 (10.1523/JNEUROSCI.3633-12.2014).

44.   T. Gerachshenko, T. Blackmer, E. J. Yoon, C. Bartleson, H. E. Hamm, S. Alford, Gbetagamma acts at the C terminus of SNAP-25 to mediate presynaptic inhibition. *Nature neuroscience* **8**, 597-605 (2005); published online EpubMay (10.1038/nn1439).

45.   P. R. Albert, L. Robillard, G protein specificity: traffic direction required. *Cell Signal* **14**, 407-418 (2002); published online EpubMay (

46.   W. K. Lim, C. S. Myung, J. C. Garrison, R. R. Neubig, Receptor-G protein gamma specificity: gamma11 shows unique potency for A(1) adenosine and 5-HT(1A) receptors. *Biochemistry* **40**, 10532-10541 (2001); published online EpubSep 04 (

47.   M. A. Lindorfer, C.-S. Myung, Y. Savino, H. Yasuda, R. Khazan, J. C. Garrison, Differential Activity of the G Protein β5γ2 Subunit at Receptors and Effectors. *Journal of Biological Chemistry* **273**, 34429-34436 (1998); published online EpubDecember 18, 1998 (10.1074/jbc.273.51.34429).

48.   B. K. Mueller, A. I. Kaya, Z. Zurawski, Y. Y. Yim, J. Meiler, H. E. Hamm, Critical Residues of Gβγ for the interaction with the SNARE Complex. *bioRxiv*, (2020); published online Epub2020-01-01 00:00:00 (

49.   T. Blackmer, E. C. Larsen, C. Bartleson, J. A. Kowalchyk, E. J. Yoon, A. M. Preininger, S. Alford, H. E. Hamm, T. F. Martin, G protein betagamma directly regulates SNARE protein fusion machinery for secretory granule exocytosis. *Nature neuroscience* **8**, 421-425 (2005); published online EpubApr (10.1038/nn1423).

50.   C. A. Wells, Z. Zurawski, K. M. Betke, Y. Y. Yim, K. Hyde, S. Rodriguez, S. Alford, H. E. Hamm, Gbetagamma inhibits exocytosis via interaction with critical residues on soluble N-ethylmaleimide-sensitive factor attachment protein-25. *Molecular pharmacology* **82**, 1136-1149 (2012); published online EpubDec (10.1124/mol.112.080507).

51.   W. A. Catterall, A. P. Few, Calcium channel regulation and presynaptic plasticity. *Neuron* **59**, 882-901 (2008); published online EpubSep 25 (10.1016/j.neuron.2008.09.005).

52.   S. E. Jarvis, G. W. Zamponi, Masters or slaves? Vesicle release machinery and the regulation of presynaptic calcium channels. *Cell Calcium* **37**, 483-488 (2005); published online EpubMay (10.1016/j.ceca.2005.01.017).

53.   M. Irfan, Z. Zurawski, H. E. Hamm, C. Bark, P. K. Stanton, Disabling Gbetagamma-SNAP-25 interaction in gene-targeted mice results in enhancement of long-term potentiation at Schaffer collateral-CA1 synapses in the hippocampus. *Neuroreport* **30**, 695-699 (2019); published online EpubJul 3 (10.1097/WNR.0000000000001258).

54.   R. P. Ceddia, Z. Zurawski, A. T. Gray, F. Adegboye, F. Shi, D. Liu, O. P. McGuinness, S. Collins, H. E. Hamm, Impairment of the Gβγ-SNAP25 brake on exocytosis enhances insulin action, protects against diet-induced obesity, and promotes adipocyte browning. *bioRxiv*, (2020); published online Epub2020-01-01 00:00:00 (

55.   R. Gilsbach, C. Roser, N. Beetz, M. Brede, K. Hadamek, M. Haubold, J. Leemhuis, M. Philipp, J. Schneider, M. Urbanski, B. Szabo, D. Weinshenker, L. Hein, Genetic dissection of alpha2-adrenoceptor functions in adrenergic versus nonadrenergic cells. *Molecular pharmacology* **75**, 1160-1170 (2009); published online EpubMay (10.1124/mol.109.054544).

56.   E. G. Gray, V. P. Whittaker, The isolation of nerve endings from brain: an electron-microscopic study of cell fragments derived by homogenization and centrifugation. *Journal of anatomy* **96**, 79-88 (1962); published online EpubJan (

57.   V. P. Whittaker, I. A. Michaelson, R. J. Kirkland, The separation of synaptic vesicles from nerve-ending particles ('synaptosomes'). *The Biochemical journal* **90**, 293-303 (1964); published online EpubFeb (

58.   K. M. Betke, K. L. Rose, D. B. Friedman, A. J. Baucum, 2nd, K. Hyde, K. L. Schey, H. E. Hamm, Differential localization of G protein betagamma subunits. *Biochemistry* **53**, 2329-2343 (2014); published online EpubApr 15 (10.1021/bi500091p).

59.   A. E. Brady, Q. Wang, P. B. Allen, M. Rizzo, P. Greengard, L. E. Limbird, Alpha 2-adrenergic agonist enrichment of spinophilin at the cell surface involves beta gamma subunits of Gi proteins and is preferentially induced by the alpha 2A-subtype. *Molecular pharmacology* **67**, 1690-1696 (2005); published online EpubMay (10.1124/mol.104.005215).

60.   Q. Wang, J. Zhao, A. E. Brady, J. Feng, P. B. Allen, R. J. Lefkowitz, P. Greengard, L. E. Limbird, Spinophilin Blocks Arrestin Actions in Vitro and in Vivo at G Protein-Coupled Receptors. *Science* **304**, 1940-1944 (2004); published online EpubJune 25, 2004 (10.1126/science.1098274).

61.   Q. Wang, L. E. Limbird, Regulated interactions of the alpha 2A adrenergic receptor with spinophilin, 14-3-3zeta, and arrestin 3. *The Journal of biological chemistry* **277**, 50589-50596 (2002); published online EpubDec 27 (10.1074/jbc.M208503200).

62.   P. Washbourne, P. M. Thompson, M. Carta, E. T. Costa, J. R. Mathews, G. Lopez-Bendito, Z. Molnar, M. W. Becher, C. F. Valenzuela, L. D. Partridge, M. C. Wilson, Genetic ablation of the t-SNARE SNAP-25 distinguishes mechanisms of neuroexocytosis. *Nature neuroscience* **5**, 19-26 (2002); published online Epub01//print (http://www.nature.com/neuro/journal/v5/n1/suppinfo/nn783_S1.html).

Acknowledgments

We thank the proteomics core of the Mass Spectrometry research Center for advice and technical assistance and Drs. Boris Alten and Ege Kavalali for sharing SNAP25 WT and KO hippocampal culture with us. We would also like to thank Dr.Simon Alford for the illustration of summary figure. Lastly, we would like to thank Dr. Analisa Thompson Gray for reviewing and confirming the statistical tests we used for the analysis.

Funding

This work was supported by the National Institutes of Health (EY10291, MH101679, MH081917, and T32 GM07628).

Authors Contributions

Y.Y., W.H.M., K.M.B. , and H.E.H. participated in research design and preliminary study. Y.Y. and K.E. conducted experiments and data analysis. W.H.M. ran mass spectrometry and helped with data analysis. K.H., A.K., R.G., and L.H. contributed in mouse breeding and sampling. Y.Y. A.K., K.M.B. and H.E.H. wrote or contributed to the writing of the manuscript. All authors reviewed the results and approved the final version of the manuscript.

Competing Interests

All authors declare that they have no competing interests.

Data Availability

All data supporting the findings of this study are available from the lead author Heidi E. Hamm upon reasonable request.

Figure legends

Figure 1. Co-immunoprecipitation of the SNARE complex and Gβγ. Co-immunoprecipitation (coIP) experimental protocol (**A**), and representative western blot of coimmunoprecipitation of SNAP25 and Gβγ in wildtype (**B**) or FLAG-α_{2a}AR (**C**) mice following the stimulation of synaptosomes with (+, 100μM epi) or without epi (-). DSP, a lipid-soluble thiol cleavable crosslinker, was added to ensure the receptor and Gβγ remained intact during co-IP experiments. We detected Gβγ in the SNAP25 pulldown but not the IgG control pulldown regardless of receptor stimulation (**B** and **C**, red box). Quantified Gβ intensities were normalized to that of input and presented as relative fold change in wildtype (**D**) or FLAG-α_{2a}AR (**E**) mice (N=3). Data are presented as mean ± SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison. coIP: co-immunoprecipitated fraction; Sup: depleted supernatant

Figure 2. Gβ specificity to the SNARE complex upon α_{2a}AR activation. Quantification of Gβ subunits interacting with the SNARE complex in wildtype synaptosomes (N=8, unless otherwise noted). $G\beta_1$, $G\beta_2$, and $G\beta_4$ interact specifically with the SNARE complex upon α_{2a}AR activation with epi. Data are presented as mean ± SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison.

Figure 3. Gγ specificity to the SNARE complex upon α_{2a}AR activation. Quantification of Gγ subunits interacting with the SNARE complex in wildtype synaptosomes (N=8,unless otherwise noted). $G\gamma_2$, $G\gamma_3$, and $G\gamma_4$ may specifically interact with the SNARE complex upon α_{2a}AR activation. Data are presented as mean ± SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison.

Figure 4. Gβ subunits interacting basally with the SNARE complex. Quantification of Gβ subunits interacting with the SNARE complex basally (N=8, unless otherwise noted). $G\beta_1$, but not $G\beta_2$, specifically interacting with the SNARE complex in both α_{2a}AR KO and FLAG-α_{2a}AR no epi conditions compared to that of IgG no epi. Data are presented as mean ± SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison.

Figure 5. Gγ subunits interacting basally with the SNARE complex. Quantification of Gγ subunits interacting with the SNARE complex basally (N=8, unless otherwise noted). $G\gamma_3$ specifically interacted with the SNARE complex in both α_{2a}AR KO and FLAG-α_{2a}AR no epi conditions compared to that of IgG no epi. Data are presented as mean ± SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison.

Figure 6. Gβ specificity to the SNARE complex upon auto-α_{2a}ARs activation. Quantification of Gβ subunits interacting with the SNARE complex in α_{2a}AR KO and FLAG-α_{2a}AR synaptosomes (N=8, unless otherwise noted). $G\beta_1$ and $G\beta_2$ specifically interact with the SNARE

complex upon auto-α_{2a}AR activation. Data are presented as mean $\pm$ SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison.

Figure 7. Gγ subunit specificity to auto-α_{2a} adrenergic receptors. Quantification of Gγ subunits interacting with the SNARE complex in α_{2a}ARs KO and FLAG-α_{2a}AR synaptosomes (N=8, unless otherwise noted). Gγ_3 specifically interacts with SNARE upon auto-α_{2a}AR activation. Data are presented as mean $\pm$ SEM. One-way ANOVA* p<0.05, **p<0.01, *Post hoc* analysis by Tukey's multiple comparison.

Figure 8. **Summary Figure**. While the G$\beta\gamma$-SNARE complex interaction may exist basally (ex. G$\beta_1\gamma_3$), only a subset of G$\beta\gamma$ dimers made with Gβ_1, Gβ_2, Gβ_4, Gγ_2, Gγ_3, and Gγ_4 (ex. G$\beta_2\gamma_3$) specifically interact with the SNARE complex upon presynaptic α_{2a}AR activation in both adrenergic and non-adrenergic neurons.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.