1

The habenula clock influences prediction of danger

Adriana Basnakova[1,2], Ruey-Kuang Cheng[3], Joanne Chia Shu Ming[3], Giuseppe D'Agostino[3],

Suryadi[4], Germaine Tan Jia Hui[3], Sarah R. Langley[3] and *Suresh Jesuthasan[2,3]

[1]School of Biological Sciences, University of Manchester, UK.

[2]Institute of Molecular and Cell Biology, Singapore.

[3]Lee Kong Chian School of Medicine, Nanyang Technological University, 59 Nanyang

Drive, Singapore 636921.

[4]School of Physical and Mathematical Sciences, Nanyang Technological University

Singapore.

*corresponding author: sureshj@ntu.edu.sg

Abstract

The response of an animal to a sensory stimulus depends on the nature of the stimulus and on predictions mediated by spontaneous activity in neurons. Here, we ask how circadian variation in the prediction of danger, and thus the response to a potential threat, is controlled. We focus on the habenula, a mediator of threat response that functions by regulating neuromodulator release, and use zebrafish as the experimental system. Single cell transcriptomics indicates that multiple clock genes are expressed throughout the habenula, while quantitative in situ hybridization confirms that the clock oscillates. Two-photon calcium imaging indicates a circadian change in spontaneous activity of habenula neurons. To assess the role of this clock, a truncated *clocka* gene was specifically expressed in the habenula. This partially inhibited the clock, as shown by changes in *per3* expression as well as altered day-night variation in dopamine, serotonin and acetylcholine levels. Behaviourally, anxiety-like responses evoked by an alarm pheromone were reduced. Circadian effects of the pheromone were disrupted, such that responses in the day resembled those at night. Behaviours that are regulated by the pineal clock and not triggered by stressors were unaffected. These findings establish that the habenula clock regulates the expectation of danger, thus identifying one mechanism for circadian change in the response to a stressor.

Keywords: anxiety, circadian clock, habenula, spontaneous activity, predictive coding.

1. Introduction

Expectations play a major role in determining the response of a person or animal to a sensory stimulus or situation. In this framework, known as predictive coding, the brain is not a passive recipient of stimuli, but actively generates predictions that are updated by experience (Barrett, 2017; Friston, 2018). Thus, encountering a cue such as the smell or sound of a predator increases the expectation of danger, turning a normally innocuous stimulus into a threat. Abnormal expectations, or predictions, have been proposed to underlie stress-related conditions such as post-traumatic stress disorder. Predictions are mediated by spontaneous activity in the brain (Berkes et al., 2011), while updating of expectations is enabled by the release of neuromodulators. Dopamine and serotonin contribute to the comparison of current input with pre-existing expectations. When outcomes surpass predictions, dopamine release by neurons in the midbrain increases; when there is disappointment, dopamine release is

inhibited (Schultz, 1998). Serotonin release by the raphe provides an indication of the

magnitude of the prediction error (Matias et al., 2017). Acetylcholine appears to have a role

in setting the precision of predictions (Moran et al., 2013). Behaviour and emotional states

can thus be understood to be the result of genetically encoded programs that are continuously

modified by predictions, enabling the animal to respond optimally to perceived reality.

The habenula, which contains several subdomains, has emerged in recent years as an

important regulator of several neuromodulators involved in predictive coding. The medial

habenula in zebrafish (Hong et al., 2013), like that of mouse (Ren et al., 2011), includes

cholinergic neurons. Disruption of specific subdomains of the medial habenula leads to

anxiety-like behaviour (Agetsuma et al., 2010; Lee et al., 2010) or aversion (Hsu et al., 2014)

that is experience dependent. The lateral habenula regulates midbrain dopaminergic and

serotonergic neurons, and phasic activity occurs here when there is an unexpected

punishment or absence of reward (Matsumoto and Hikosaka, 2009). As an animal learns the

actions that prevent punishment, this activity lessens (Amo et al., 2014). In addition to

activity that is evoked by sensory stimuli or perceptions of reward, the habenula displays

spontaneous activity (Jetti et al., 2014; Kim and Chang, 2005). Excessive levels of burst

firing, which can be interpreted as constant negative prediction, has been detected in the

lateral habenula of animals with depressive behavior (Andalman et al., 2019; Yang et al.,

2018). To understand how mood and behaviour are shaped, it is thus important to not only

understand how the habenula responds to reward and sensory input, but to also investigate

how spontaneous activity is regulated.

One factor that affects the level of spontaneous activity in neurons is the molecular clock.

The molecular clock in most organisms consists of a transcriptional-translational feedback

loop with a period of ~ 24 hours (Takahashi, 2017). Core components of the clock in animals

include heterodimers of Clock and Arntl (Bmal), which enter the nucleus in the daytime.

Here, they drive transcription of *period* (*per*) and *cryptochrome* (*cry*) genes, which then

inhibit the Clock-Arntl heterodimer. Clock and Arntl also control transcription of channels

that affect resting membrane potential in neurons, driving this close to threshold and thus

increasing spontaneous firing rate in the daytime (Harvey et al., 2020). Components of the

clock machinery are found in nearly all cells, including those of the habenula (Baño-Otálora

and Piggins, 2017; Mendoza, 2017), and the habenula molecular clock continues cycling for

multiple days even in the absence of the main pacemaker of the brain, the suprachiasmatic

nucleus (Guilding et al., 2010; Salaberry et al., 2019). Spontaneous firing in the habenula of

rat (Zhao and Rusak, 2005) and mouse (Guilding et al., 2010; Sakhi et al., 2014a, 2014b)

varies across a circadian cycle, in both the lateral and medial habenula. The significance of

the habenula clock is unclear, however. Given the role of habenula activity in predictive

coding, we hypothesize that the clock causes a circadian change in behaviours that are

modified by recent experience. To test this, we use the zebrafish as the experimental system.

We examine anxiety-like behaviour, which is increased by expectation of danger.

2. Materials and Methods

2.1 Single cell data reanalysis

Count tables were downloaded from Gene Expression Omnibus, accession number

GSE105115. All processing downstream was done using R/BioConductor (R

4.0.2/Bioconductor 3.12) packages designed to handle single cell data using the

SingleCellExperiment class (v. 1.12.0) (Amezquita et al., 2020). A detailed explanation with

code and figures is provided as a GitHub repository at

https://github.com/langleylab/habenula_reanalysis. Briefly, gene-level count tables from the

SmartSeq2 experiment were merged retaining the information on the plate of origin. Genes

that were not detected in any barcode were removed. Barcodes were discarded according to

the following criteria: low expression, low number of genes detected, high percentage of

malat1 expression, high percentage of mitochondrial transcripts expression and high

percentage of ribosomal transcript expression. Thresholds for every criteria were set as 3

Mean Absolute Deviations (MADs), with the exception of mitochondrial % which was set at

4 MADs. Outlier calculations were done separately in each plate using the *isOutlier* (using

the *batch* argument) and *perCellQCMetrics* functions from the *scater* package (v. 1.18.6)

(McCarthy et al., 2017). These filters yielded 17673 genes and 1140 barcodes for

downstream analyses. Barcodes were then clustered using the *quickCluster* function from the

scran package (v. 1.18.5 (Lun et al., 2016)), blocking for the plate of origin. Per-cell size

factors were then determined by deconvolution using *scran*, and they were used to perform

multi-batch normalization (batching by plate) using the *multiBatchNorm* function from the

batchelor package (v. 1.6.2 (Haghverdi et al., 2018)). Highly variable genes were estimated

on the multi-batch normalized data blocking by plate using the *modelGeneVar* function from

the *scran* package, which estimates the mean-variance trend and decomposes variance for

each gene in technical and biological components according to the deviation from the trend.

Variances were combined across plates using the *combineVar* function from *scran*, and the

2000 genes with the highest biological variance were selected using the *getTopHVGs*

function from *scran*. These were used as an input to perform dimensionality reduction via

Principal Component Analysis as implemented in the *BiocSingular* package (v. 1.4.0, (Lun,

2020a)).

The first 50 principal components were then used as an input to perform batch (plate) effect

correction via the *fastMNN* function from *batchelor*, which determines a correction vector in

PCA space using mutual nearest neighbour pairs to estimate the amount and direction of

necessary correction. The dimensionality of the corrected space, i.e. the number of most

relevant dimensions for subsequent steps, was determined using the *maxLikGlobalDimEst*

function from the *intrinsicDimension* package (v. 1.2.0 (Johnsson et al., 2014)). The value for

the *k* parameter for this function (k = 30) was determined as the rounded square root of the

number of cells, and the estimation yielded 13 dimensions. Subsequently, the first 13

dimensions of the corrected space were used to run the t-stochastic neighbour embedding (t-

SNE) algorithm for data visualization, and to construct a Shared Nearest Neighbour (SNN)

graphs for clustering. Several SNN graphs were constructed at different resolutions, i.e. 10,

15, 30, 40 and 50 neighbours were considered for the graph construction; all edges were

weighted by the Jaccard similarity between the neighbours of each node. SNN graph

construction was performed using the *buildSNNGraph* function from *scran*. Then, the

Louvain community detection algorithm (as implemented in the *igraph* package (v. 1.2.6

(Csardi and Nepusz, 2006)) was used to identify clusters. Relationships between different

clustering results (obtained by using different neighbours in the SNN graph construction)

were inspected using a clustering tree drawn using *clustree* (v. 0.4.3 (Zappia and Oshlack,

2018)). Clusterings with 10 and 15 neighbours were further explored, as they yielded

numbers of clusters similar to the ones in the original publication. For both 10 and 15

neighbours, cluster separation was evaluated using three methods: approximate silhouette

width, neighborhood purity and pairwise modularity, as implemented in the *bluster* package

(v. 1.0.0 (Lun, 2020b)) in the *approxSilhouette*, *neighborPurity* and *pairwiseModularity*

functions. The 15 neighbour clustering showed slightly better separation so it was considered

for downstream analyses. In order to reconstruct the original cell type identities as identified

by the authors, the *AUCell* package (v. 1.12.0, (Aibar et al., 2017)) was used in conjunction
with the cluster marker genes as published by the authors, although these marker genes were
identified in the 10X dataset. Raw counts were ranked in each cell and the AUC for each
marker set was calculated; each cell was assigned the identity yielding the highest AUC value
using the 30% top ranking genes. AUCell is agnostic to the transcriptional space and to other
intrinsic properties of the dataset, which will be influenced by differences in filtering and
processing steps. Therefore the performance of AUCell was evaluated in this dataset
calculating the approximate silhouette width of the cell type assignment, and it was compared
to our clustering results by calculating pairwise Jaccard indices. Most identities (Hb1, Hb2,
Hb5, Hb6, Hb9, Hb1, Hb14, Hb15 and olfactory neurons) are reasonably well separated, but
only 4 (olfactory, Hb06, Hb03, and Hb08), are reproduced in our clustering with a Jaccard
index above 0.5 with any single cluster. This is in line with what the authors of the original
paper state regarding the poor performance of SmartSeq2 in terms of cell type recovery, and
is also to be expected when using cell type markers identified in the 10X dataset.
161
For the 10X datasets (larval) no filtering was performed as the distribution of QC metrics was
well within standard ranges, suggesting data were already filtered. Normalization, HVG
estimation and dimensionality reduction were performed in the same way as described above,
with the exception that there was no batch effect to account for and correct. An additional
doublet detection step was included using the *scDblFinder* package (v. 1.4.0, PMID
(Germain et al., 2020)), which flagged 172 barcodes as doublets and 4193 as singlets.
Doublets were not removed at this stage of the analysis. The intrinsic dimension estimation (k
= 66) yielded 12 dimensions to perform SNN graph clustering, which was performed using 5,
10, 15, 20, and 30 neighbours. In parallel, 36 dimensions were also considered as done by the
authors of the original publication. For the 12-dimensional space, 5 and 10 neighbours were
considered as they yielded numbers of clusters similar to those reported in the publication;
for the 36-dimensional space, 10 and 15 neighbours were considered. However, in both 12
and 36 dimensions a cluster was entirely made up of doublets, so doublets were removed and
the whole process (normalization, HVG selection and dimensionality reduction) was
repeated. Cluster separation metrics (approximate Silhouette and modularity) favoured
clustering with 10 neighbours (12 dimension) and 15 neighbours (36 dimensions). To
reproduce the original clustering results the AUCell package was used, and Jaccard indices
were used to match clusters to labels. In order to improve the label assignment, cells whose

top AUC label showed less than a 20% difference from the second best label were classified

as "ambiguous". Ambiguous labels were then reassigned using the closest centroid in the

approximate silhouette width calculation, thus making the labels conform more to the

transcriptional space. Improvements in the reassignment of these labels were measured by

comparing silhouette widths and the distribution maximum Jaccard indices for every cluster-

AUC label pair before and after reassignment. Both metrics showed an improvement in both

silhouette width and median best Jaccard index upon reassignment.

Adult zebrafish data was processed in the same above, accounting for batch effects (2

different captures). SNN graph construction with 4 neighbours in 16 dimensions was chosen

following the same pipeline and reasoning used in the larval dataset. AUCell was used to

determine identities using the markers provided by the authors of the original publication,

and the reassignment procedure described above was used to correct ambiguous labels.

2.2 Transgenic lines

TgBAC(gng8:GAL4)^{c426} (Hong et al., 2013) or T^{s1011t} (Scott and Baier, 2009) was used to

drive expression in the habenula. Calcium imaging was performed using

Tg(elavl3:GCaMP6f)^{a12200} and *Tg(UAS:GCaMP6s)^{sq205}*. *Tg(UAS:EGFP-2A-Δclocka-*

5×MYC)^{sq215}, abbreviated here as *Tg(UAS:EGFP-Δclocka)* was generated by injection of

pT2-UAS:EGFP-2A-Δclocka-5xMYC using Tol2 mediated transgenesis. All experiments

were carried out under guidelines approved by the IACUC of Biopolis (181408) and NTU

(A19014). The minimal number of animals was used.

2.3 Two-photon Imaging

2.3.1 Extended imaging

Based on the methods described by (Leung et al., 2019), non-anesthetized and non-paralyzed

larvae were mounted in 2% low melting agarose (LMA), in a glass-bottom dish (Matek). The

dish was filled with E3 medium supplemented with 1% Hepes and bubbled continuously with

carboxygen (95% oxygen). Imaging was performed with a Nikon A1R MP system, attached

to an upright FN1 microscope using a 25x 1.0 NA water dipping objective. Resonant

scanning was performed with 2x averaging, and 20 focal planes were collected, spaced 5 μm

apart. Signal was detected with a GaAsP detector. A piezo drive (Mad City Labs) was used

for fast focusing during imaging. 21 stacks were collected, at an interval of 1 second,

followed by a delay of 1 hour. The maximum imaging duration was 29 hours. After imaging,

larvae were checked for blood flow. They were then released from the agarose. Only data

from fish with normal blood flow and active swimming upon release were used for analysis.

To obtain a measure of calcium levels in the whole habenula, fluorescence intensities were

summed at each time point using Fiji. The intensity at each hour was averaged and z-scores

were plotted as a function of *zeitgeber* time.

218 *2.3.2 Short duration imaging*

6-day-old *Tg(elavl3:H2B-GCaMP6s)* or *Et(-1.5hsp70l:Gal4-VP16)^{s1011t}, Tg(UAS:GCaMP6f)*

larvae were immobilized with mivacurium, mounted in 2% low melting temperature agarose

and imaged with the two-photon system described above, using resonant scanning at a single

plan. Frame size was 512x256 pixels and 2x averaging was used, leading to a frame rate of

14.5 Hz. Segmentation of cells expressing cytoplasmic GCaMP6f was performed manually in

Fiji, while segmentation of cells expressing nuclear GCaMP6s was performed automatically

with Suite2p (Stringer and Pachitariu, 2019). Z-scores were obtained for each cell in a

recording, and cells were clustered using K-means clustering in Python. The classification

was performed using maximum likelihood and leave-one-out cross-validation. Specifically, at

each iteration we removed one dataset from the group and constructed empirical cluster

distributions for both categories (ZT3 and ZT15) from the remaining datasets. The removed

dataset was then assigned to the category whose empirical distribution maximized its

likelihood. This process was repeated independently until all datasets were classified.

232 2.4 Immunohistochemistry and imaging of EGFP-ΔCLK

The *Tg(gng8:GAL4,UAS:EGFP-Δclocka)* larvae at the age of 7 dpf were fixed in 4% PFA at

4ºC for ~22 h. For immunostaining of extracted brains, skin covering the brain was manually

removed and exposed brains were washed 3 times in PBST. Antigen retrieval was performed

by incubating the fish in 150 mM Tris-HCl (pH 9.0) for 5 min at room temperature in order

to equilibrate, and by subsequent heating of the samples at 70˚C for 15 min. Following

several washes and a blocking step, samples were incubated in the rabbit anti-GFP primary

antibody (Torrey Pines Biolabs Inc., #TP401), diluted 1:500, and the mouse anti-Myc

antibody (9E10; Santa Cruz Biotechnology), diluted 1:100, at 4ºC for a few days. The

secondary antibodies used were Alexa Fluor 488 goat anti-rabbit (ThermoFisher and Abcam),

diluted 1:1000, and Alexa Fluor 647 goat anti-mouse (ThermoFisher), diluted 1:500. The

immunostained brains were mounted in 2% LMA in PBS and imaged using a Zeiss LSM800

confocal microscope equipped with 40X 0.7 NA water immersion objective.

2.5 Screening and selection of *Tg(gng8:GAL4,UAS:EGFP-Δclocka)* fish

All of the *Tg(gng8:GAL4,UAS:EGFP-Δclocka)* fish used in the following experiments were

first screened for GFP fluorescence at 3 dpf using a Olympus MVX10 stereo fluorescence

microscope. Since the expression of EGFP-ΔCLK in the habenula was mosaic, only

individuals with expression in the majority of habenula cells were selected and used for

further experiments as ΔCLK-positive fish. Individuals with zero expression as well as larvae

exhibiting a minimal number of randomly labelled cells (less than ~15 cells per habenula)

were used as the control siblings (Figure S1). Following the behavioural experiments, larvae

were immobilized in tricaine methanesulfonate (MS-222; final concentration of 0.01 mg/ml;

Sigma), mounted in 2% LMA in E3 medium and the habenula-specific expression of EGFP

was assessed using a Zeiss LSM800 confocal microscope and 40X water dipping objective.

2.6 Quantification of DA and 5-HT levels

Three-month old female ΔCLK fish and their control siblings were immobilized using iced

water and euthanized by decapitation at ZT3 and ZT15. Whole brains, excluding the

olfactory bulb, were extracted in the ice-cold Ringer's solution (116 mM NaCl, 2.9 mM KCl,

1.8 mM $CaCl_2$, 5.0 mM HEPES, pH 7.2), snap-frozen in liquid nitrogen and stored in -80˚C

until further analysis. Supernatant was collected as described previously (Chatterjee and

Gerlai, 2009). Quantification of dopamine (DA) and serotonin (5-HT) was carried out by

LC/MS using the following analytical standards: dopamine hydrochloride (D2960000:

Sigma-Aldrich), dopamine-1,1,2,2-$d_4$hydrochloride (73483: Sigma-Aldrich) and serotonin

(14927: Sigma-Aldrich). Three biological replicates (i.e. measurements of different samples)

were performed.

2.7 Behaviour assays

2.7.1 Acoustic startle following exposure to Schreckstoff

Thirteen to sixteen day-old ΔCLK larvae and their non-expressing siblings were placed in

two transparent plastic tanks with dimensions of 80 x 54 x 33 (L x W x H in mm) and their

locomotor activity was recorded from above at 10 fps using a camera (acA2040-90µm USB

3.0; Basler). Each of the two tanks was filled with 20 mL of facility water and held 8 fish at

most. The water level in each tank was 5 mm in height to minimize vertical movement. The

alarm substance was freshly prepared from adult zebrafish each week (Mathuru et al., 2012),

and was tested for effectiveness on adults before use on larvae. An Arduino Grove Vibration

Motor was positioned in between the two plastic tanks, on the stage holding the tanks, to

provide the stimulus. The vibration was 1 sec in duration and was presented 3 times, at 1

minute intervals, starting five minutes before and after the delivery of the *Schreckstoff*. A

startle response was defined as a change in body orientation of more than 90 degrees within 5

sec in response to each vibration tone. The response (yes or no for each fish) was averaged

across 3 tones for each group to obtain a response percentage before and after the delivery of

the *Schreckstoff*. Both the vibration tones and the delivery of *Schreckstoff* were controlled by

an Arduino board synched with the video-recording Python codes. Experiments were

performed on 3 separate batches of larvae. A different batch of freshly prepared skin extract

was used for each experiment; the extract was verified for effectiveness by testing on adult

fish. Only extract that evoked freezing was used.

2.7.2 Locomotor activity

The *Tg(gng8:GAL4,UAS:EGFP-Δclocka)* larvae were raised under standard conditions. Each

experimental run involved animals from the same batch, originating from one pair of parents.

The larvae were screened and selected as described in the previous section. At the age of 6

dpf the larvae were placed individually in a 24-well plate filled with 1.5 ml of tank water per

well. The plate was then transferred into an incubator equipped with an IR camera (acA2040-

90μm USB 3.0: Basler), a white LED box and four IR LED bars (LBS2-00-080-3-IR850-

24V, 850 nm: TMS Lite). The camera and the LEDs were connected to a PC via a

microcontroller board (Arduino UNO SMD). At the onset of recording (ZT6 – 9), fish were

kept under ambient light, which was switched off at ZT14 and was kept off during the next

day for the monitoring of activity under DD. A customized script written in Python 2.7,

incorporating functions from the OpenCV library was used to control the Arduino

microcontroller and simultaneously video-track fish locomotor activity. The video was

captured using 576 x 864-pixel resolution at 10 frames per second and the background

subtraction method was implemented to extract the 'x' and 'y' coordinates of each fish

online. The position data was then converted to total distance (mm) travelled by each fish per

1 min time. The data conversion, further analysis and plotting were carried out using custom-

written Excel macros, Python 2.7 and Estimation Statistics (Ho et al., 2019).

2.7.3 Arousal threshold

Six days-old ΔCLK larvae and their control siblings were randomly placed in a 24-well plate and their locomotor activity was video-tracked as described in the previous section. In order to assess the arousal during sleep, vibration stimuli were applied through a speaker controlled, via an Arduino microcontroller board, by a modified version of the activity-tracking code in Python 2.7. Starting at ZT17, eighteen stimuli of nine different intensities, where 50% of computer audio output represented the highest intensity, were delivered for 200 ms (starting from 0%) in 12-16% increments every minute, first in a descending and then ascending order. The time interval between stimuli was determined based on previous findings that suggest a 30-second inter-stimulus gap is sufficient to avoid behavioural habituation (Burgess and Granato, 2007). The protocol was repeated at the commencement of every hour until ZT22, while the onset by ascending or descending order was altered each time. In total, 12 replicates (i.e. technical replicates) of each stimulus intensity were generated. A fish was scored as responding if it moved more than 7 pixels (~ 4 mm) following the stimulus delivery and the average percentage of responding fish was calculated for each of the stimulus intensities (Woods et al., 2014). The observed values were corrected for variations in the baseline locomotor activity, by first calculating the background probability of locomotion as a percentage of larvae which moved more than 7 pixels 5 s before the application of each stimulus. The average of 108 measures of the background probability of locomotion was then used to calculate the corrected probability of response at each of the stimulus intensities, implementing the following equation: corrected probability of response = observed value x [(observed value - background offset) / (max observed value - background offset)].

2.7.4 Novel tank assay

All experiments were conducted between ZT2 and ZT4 on 3-5 month-old adult zebrafish, as described elsewhere (Haghani et al., 2019). Fish were recorded in pairs in a darkened room with a black background, where tanks were illuminated from the top with a natural light LED bar. Novel tanks were made of glass measuring 20 cm (L) by 5 cm (W) by 12 cm (H) with opaque sides (to prevent fish from viewing each other), placed side by side. Fresh water from the facility was added to tanks up to the 10 mL mark to make a total volume of 1 L. The setup was placed within a curtained area to obscure the experimenter.

Prior to exposure to the novel tank, fish were handled carefully to ensure that handling stress

was minimized. Fish were gently netted in pairs from their home tanks in the facility to

smaller crossing tanks, where they were brought into the room with the behavioural setup.

Then, fish were netted into 100 mL beakers with ~30 mL of fish facility water and quickly

poured into the novel tank. The recording was started within 30 s of fish being in the

recording tanks. For exposure to *Schreckstoff,* fish were placed in 100 mL beakers containing

200 µL of heated skin extract (Chia et al., 2019) in 50 mL fish facility water for 2 minutes.

Fish were netted into another beaker containing fresh facility water before being poured into

the novel tank. Videos were acquired at 10 frames per second for 600 s (10 mins) using a

Basler Ace camera (acA1300–200 µm; 1,280 × 1,024). The position (x-y coordinates) of the

fish were tracked, and speed was binned by mm/s into a probability density curve.

For acute exposure to *Schreckstoff*, fish were first habituated in the glass tanks for 5-7 mins.

Then, fish were recorded in the tanks for 5 mins before 1 mL of 1:4 dilution of heated skin

extract was pipetted into the tanks. The recording continued for 5 mins after *Schreckstoff*

delivery for the alarm response to be observed. Two fish were tested at one time, each in a

separate tank, with one tank containing ΔCLK fish and the other a non-expressing sibling.

The tank used for a particular genotype was switched at every trial. Characteristic alarm

behaviours were quantified, such as time spent in the bottom third of the tank, number of

darting episodes (fish speed >5 SD above average swim speed for the first 5 mins), and

number of freezing episodes (movement of <5.5 mm per second). Tracking was automated to

prevent observer bias. Sample size was based on previous experiments (Chia et al., 2019).

2.7.5 Light/dark choice assay

All experiments were performed between ZT2 and ZT9 in a behavioural arena within

blackout curtains as described previously (Cheng et al., 2016). Four transparent plastic tanks

(dimensions: 43 mm W × 60 mm L × 30 mm H) filled with 30 mL fresh tank water served as

assay chambers. Opaque cardboard sheets were placed between the 4 tanks. Tanks were

placed on an Apple iPad screen and videos (9 fps) were recorded on a USB3.0 Basler camera

(Model# acA2040-90umNIR 1440 × 1080 pixels) with a long-pass filter (MIDOPT, LP830

830 nm) placed above. Two IR light bars (850 nm peak from TMS-lite) positioned next to the

4 tanks served as an IR light source. The light/dark compartments were created by white and

black rectangles (with 50% transparency level for the black) on a Microsoft PowerPoint slide

displayed on the iPad Air at highest brightness level. Sample size was based on previous

experiments(Cheng et al., 2016).

Larvae were exposed either to *Schreckstoff* or to control water (embryo medium) in a

treatment tank of the same dimensions with the same volume of tank water for 10 minutes.

Larvae were then gently pipetted from the treatment tank into two wash-out tanks

sequentially before being placed into the dark/light test chamber. Real-time tracking of the

fish was achieved by custom-written Python codes using the OpenCV library and recorded as

coordinates of the centroid of the fish across time. Larval behaviour was recorded for 10

minutes. Offline analysis was performed using custom-written Excel macros to determine the

position of the animals, the number of entries into each compartment, the percentage of total

time spent in each compartment, etc. A few fish (N=2 in both the Control group and

Gng8/ΔCLK group) were excluded from the analysis because they were entirely immobile or

because of tracking errors.

2.8 In situ hybridization

Hybridisation chain reaction with split initiator probes was conducted using the HCR v3.0

protocol for whole-mount zebrafish embryos and larvae from Molecular Instruments. Probe

hybridization buffer, amplification buffer, probe wash buffer, probe sets and HCR amplifiers

were obtained from Molecular Instruments.

Tg(gng8:GAL4, UAS:EGFP-Δclocka) larvae were maintained at 28°C under the standard 14-

10h LD cycle. For characterization of clock genes under DD, nacre and *Tg(gng8:GAL4,*

UAS:EGFP-Δclocka) larvae were maintained under standard LD conditions transferred to a

dark room at ZT14 the day before fixation.

All *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* larvae were screened for mosaic expression of

EGFP-ΔCLK at 5 dpf and EGFP-ΔCLK positive larvae were separated from non-expressing

siblings. All larvae were fixed at ZT3, ZT7, ZT11, ZT15, ZT19 and ZT23h with 4%

paraformaldehyde (PFA) in phosphate buffered saline (PBS) at 4°C in a 1.5mL eppendorf

tube (1 tube per time-point) and incubated at 4°C overnight. Fixations for DD larvae were

performed under dim light. For dehydration and permeabilization, larvae were washed 4 x 10

minutes and 1 x 50 minutes with 1mL of 100% methanol (MeOH) at room temperature, and

subsequently stored in -20°C for several days. For rehydration, larvae were washed 1 x 5

minutes with 1mL of 75% MeOH (in PBST), followed by 50%, and 25% MeOH (in PBST),

and subsequently 5 x 5 minutes washes with 100% PBST. Larvae were then treated with 1mL

of 30µg/mL proteinase K for 45 minutes at room temperature and washed twice with 1mL of

PBST without incubation. Postfixation of larvae was conducted by adding 1mL of 4% PFA

for 20 minutes at room temperature. The larvae were then washed 1 x 1 hour and 4 x 5

minutes with 1mL of PBST. For pre-hybridization, 4 larvae from each sample were

transferred to a 1.5 mL eppendorf tube and 500 µL of probe hybridization buffer was added

and incubated at 37°C for 30 minutes. For detection of clock genes, probe hybridisation

buffer was aspirated from the samples and probe solution (2 µL each of 1 µM *arntl1b* and

per3 qHCR probe sets or 3 µL of *cry1a* dHCR probe set in 500 µL of probe hybridisation

buffer at 37°C) was added and incubated at 37°C for ~22 hours. Samples were washed 4 x

15 minutes with 500 µL of probe wash buffer at 37°C, followed by 2 x 5 minutes washes

with 500 µL of 5X SSCT at room temperature for removal of excess probes. 5X SSCT was

aspirated from the samples and 500 µL of amplification buffer was added and incubated for

30 minutes at room temperature. For amplification of signal, amplification buffer was

aspirated from the samples and hairpin solution (10 µL each of snap-cooled HCR amplifier

hairpins h1 and h2 for all 4 probe sets in 500 µL of amplification buffer at room temperature)

was added and incubated in a dark box at room temperature for ~22 hours. Hairpin solution

was then removed by 2 x 5 minutes and 2 x 30 minutes washes and another 1 x 5 minutes

wash with 500µL of 5X SSCT at room temperature. Samples were lastly stored at 4°C in a

dark box before microscopy.

2.8.1 qHCR Imaging

425   4 larvae from each sample were mounted in 2% LMA in PBS in dorsal view and imaged

426   using a Zeiss LSM800 upright confocal microscope with 40 x water dipping objective at 0.5x

427   zoom. The acquisition settings were identical in all experiments. Z-stacks were collected at

428   512x512 resolution, giving a x-y pixel size of 0.62 µm, and a z-step of 0.73 µm. Sample size

429   was based on (Trivedi et al., 2018).

2.8.2 Image Analysis

To quantify gene expression in the entire habenula across different fish, segmentation of

habenula was performed manually using the Segmentation Editor in Fiji (Schindelin et al.,

2012). The integrated intensity of *per3* and *arntl1b* was measured using the 3D Objects

Counter (Bolte and Cordelières, 2006). The ratio was calculated to compensate for potential

tube-tube variation as well as change in intensity with imaging depth. Values were plotted

using GraphPad Prism.

To compare the level of *per3* in ΔCLK expressing and non-expressing cells, within-fish

analysis was performed on single planes. ΔCLK cells were segmented by thresholding the

GFP signal, while the habenula was manually segmented. Masks for control cells were

obtained by subtracting the ΔCLK mask from the habenula mask. ROIs were then created,

and mean intensity measured, based on methods described by (Choi et al., 2018), i.e. with a

mean filter of 3 and background correction. Statistical analysis was performed with

estimation statistics (Ho et al., 2019). 5000 reshuffles of the control and test labels were

performed to calculate each permutation P value.

2.8.3 dHCR imaging

Following label for *cry1a*, 3 larvae from each of 6 time-points were imaged with confocal

microscopy, using an Achroplan 63x water dipping objective and 2048 x 4093 image size,

giving a x-y pixel size of 0.0495 μm. z step was 0.78 μm. Sample size was based on Choi et

al, 2018. To analyse the images, the habenula in one plane ~ 10 μm from the surface was

manually segmented. Spots were identified in Fiji using a Laplacian transform with FeatureJ,

with the smoothing scale set at 5. The number of spots was counted using FindMaxima, with

the prominence > 10.0. Identical settings were used for acquisition and analysis of all dHCR

images.

3. Results

3.1 Multiple clock genes are expressed in the zebrafish habenula

Using in situ hybridization in brain sections with DIG-labelled probes, a number of clock

genes, such as *per1b* and *bmal* (*arntl1a*), have been shown to be expressed in the habenula of

adult zebrafish (Weger et al., 2013). To better characterize clock gene expression in this

structure, we analyzed single cell transcriptomes of the zebrafish habenula (Pandey et al.,

2018). A diversity of clock gene components, including *arntl1b*, *arntl2*, *per1a*, *per2*, *per3*,

cry1a, *cry1ab*, *cry1ba*, *cry1bb*, *clocka*, *clockb* and *nr1d1* were found (Figure 1). These

transcripts were isolated from *gng8*-expressing habenula neurons, which are found

throughout the habenula (Hong et al., 2013; Pandey et al., 2018). All clusters contained clock

genes, suggesting that the molecular clock is broadly distributed.

To examine the temporal dynamics of the clock in larval zebrafish, quantitative in situ

hybridization using hybridization chain reaction (HCR) (Choi et al., 2018) was carried out

on fish that were fixed at 6 different time points (Figure 2). With qHCR imaging, *arntl1b*

appeared to be expressed at a higher level in the habenula compared to the rest of the brain

(Figure 2A), with a stronger signal at approximately *zeitgeber* time ZT 15; expression at ZT

19 and ZT 23 was detected at the posterior margins. *per3* was detectable broadly in the brain,

including the entire habenula, with strongest expression at ZT 23 and ZT 3 (Figure 2B).

Signal intensity obtained with qHCR imaging provides a relative measure of transcript level

(Choi et al., 2018), and can vary with sample thickness and between tubes. To minimize error

from these potential confounds, we calculated the ratio of *arntl1b/per3* as a means of

comparing gene expression between animals. This ratio showed a circadian variation, with a

peak at ZT 11 - 15 (Figure 2D). To test whether clock gene cycling requires light, we

examined expression in fish grown in constant darkness. The ratio of *arntl1b*/*per3* expression

was maximal at CT 15 and minimal at CT3 (Figure 2E-G). We used single molecule imaging,

with dHCR imaging of *cry1a*, to further test whether the clock is cycling. Again, a circadian

change in expression level was detected in the habenula in constant darkness (Fig 2H, I), with

a peak at CT3 and a trough at CT15, similar to what has been reported in other tissues under

LD (Cavallari et al., 2011) or DD conditions (Kobayashi et al., 2000). These observations

confirm previous findings using clock reporters (Wang et al., 2020; Weger et al., 2013) that

the molecular clock is active in the zebrafish habenula.

3.2 The zebrafish habenula displays circadian variation in intracellular calcium

The presence of the molecular clock should lead to a number of circadian changes in a cell.

One such change is the level of cytoplasmic calcium (Pennartz et al., 2002). To examine

intracellular calcium levels in the zebrafish habenula, two-photon imaging was carried out in

fish held in constant darkness. Confocal microscopy was not used to avoid light-induced

disruption of the molecular clock by visible light (Tamai et al., 2007). By imaging a series of

z-stacks every hour, such that the entire habenula could be captured (Figure 3A-D), we

observed an increase in overall fluorescence intensity during the subjective day, followed by

a decrease during the subjective night (Figure 3E; 5 out of 6 fish imaged). This finding is

consistent with the hypothesis that the habenula contains an active circadian clock that has

the potential to drive changes in the intracellular calcium even in the absence of light, a

prominent circadian *zeitgeber*.

In the mouse, neurons in both medial (Sakhi et al., 2014a) and lateral (Sakhi et al., 2014b)

habenula display spontaneous neural activity that varies across a circadian cycle. To assess

whether this is also the case in the zebrafish habenula, we carried out calcium imaging at ~15

Hz at a single plane. We imaged at two different time points, starting at ZT3 and ZT15. The

z-score for fluorescence of each cell was calculated, and a vector was generated for the

activity of cells with high variance. A difference was observed between the means of this

vector at ZT3 and ZT15 (Figure 3F). To further characterize the activity, cells were classified

using K-means clustering. The cluster composition was distinct between day and night

(Figure 3G). To test the robustness of this apparent difference, we asked if a classifier could

correctly assign a random dataset to either ZT3 or ZT15. Indeed, 21 out of 26 datasets of the

cytoplasmic indicator and 10 out of 13 datasets with the nuclear indicator were correctly

assigned. These findings suggest that neural activity in zebrafish habenula is different at day

versus night.

3.3 Expression of a truncated *clock* gene in the habenula affects function

The expression of clock genes and the circadian change in intracellular calcium in the

habenula raises the possibility that an intrinsic clock influences habenula function. To test

this, we selectively expressed a truncated *clocka* gene (Dekens and Whitmore, 2008), referred

to as Δ*clocka*, in the habenula. This construct has been shown to act in a dominant manner to

inhibit the molecular clock in the zebrafish pineal (Livne et al., 2016). Δ*clocka* was expressed

in the habenula using the GAL4/UAS system with *TgBAC(gng8:GAL4)* (Hong et al., 2013)

as the driver. In addition to Δ*clocka*, the effector construct also included enhanced green

fluorescent protein (EGFP), which was separated from ΔClocka (ΔCLK) by a 2A peptide

linker, thus enabling the expression of two separate proteins. The presence of EGFP in the

habenula neurons of *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* fish (Figure 4A, B) indicates that

the construct is expressed appropriately and enables convenient identification of ΔCLK-

positive individuals (Livne et al., 2016). Labelling with an antibody against the Myc tag,

which is fused in frame at the C-terminus of ΔCLK, indicates that the truncated protein is

translated (Fig. 4C). Cells expressing ΔCLK would be expected to have a lower level of

Clock-regulated genes such as *per*, than non-expressing cells. To assess this, qHCR imaging

was performed on larvae with scattered expression of ΔCLK, and the level of *per3* in cells

with EGFP expression was compared with the ratio in non-expressing cells in the same

sample. As seen in Fig. 4D the level of *per3* differed between ΔCLK and control cells at all

time points, with a paired Cohen's D of 1.58 [95.0%CI 1.03, 4.22] at CT3, 0.525 [95.0%CI

0.337, 4.25] at CT7, -1.89 [95.0%CI -4.78, -0.734] at CT11, -1.57 [95.0%CI -3.38, -1.15] at

CT15, 0.872 [95.0%CI 0.636, 2.26] at CT19 and 0.637 [95.0%CI 0.536, 0.816] at CT23. This

suggests that expression of ΔCLK influences the molecular clock, although it does not

eliminate cycling.

Given the partial inhibition shown by gene expression analysis, we asked whether is any

evidence that expression of ΔCLK affects habenula function. The habenula regulates the

broad release of serotonin and dopamine (Amo et al., 2010; Satar et al., 2020) and contains

cholinergic neurons (Hong et al., 2013). If expression of ΔCLK has any effect on habenula

function, the global levels of secreted serotonin, acetylcholine and dopamine, which vary in a

circadian fashion (Hut and Zee, 2011; Mendoza, 2017), should be affected. To test this,

brains were isolated from adult zebrafish expressing EGFP-ΔCLK and from their non-

expressing siblings, at ZT3 and ZT15. The supernatant from extracts were analysed by liquid

chromatography and mass spectrometry (LC/MS) (Chatterjee and Gerlai, 2009). While non-

expressing fish had a clear difference in the day- and night-time levels of dopamine (Figure

4E; mean difference = 124 pg/mg, $p < 0.0002$ by permutation t-test), there was almost no

difference in fish expressing ΔCLK protein (mean difference = 6.3 pg/mg, $p = 0.30$ by

permutation t-test). The mean difference in between day and night-time levels of serotonin

and acetylcholine levels was also higher in control fish (117 pg/mg and 454 pg/mg

respectively, $p < 0.0002$ by permutation t-test in both cases), compared to ΔCLK-positive fish

(68.1 pg/mg and 137 pg/mg respectively, $p < 0.0002$ by permutation t-test). This change of

neuromodulator levels is consistent with the notion that expression of ΔCLK in the habenula

affects function.

3.4 Habenula expression of ΔCLK reduces anxiety-like behaviour induced by

the alarm substance

We hypothesized that a stress response would be affected by the expression of ΔCLK in the

habenula. One such response for fish is the alarm response, which is triggered by the

pheromone *Schreckstoff*. In adult zebrafish, Schreckstoff causes an acute change in swimming behaviour (Jesuthasan and Mathuru, 2008; Speedie and Gerlai, 2008). Following transient exposure to *Schreckstoff*, fish display abnormal scototaxis (Maximino et al., 2014) and heightened vigilance. To determine whether manipulation of the habenula clock influenced the acute response, adult fish were placed individually in a test tank. After a period of acclimatization, the alarm substance was introduced into the tank (Figure 5A). Adult *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* zebrafish showed an increase in darting (Figure 5C, paired mean difference of 1.33 seconds [95.0% CI 0.5, 2.67], p = 0.0260 Wilcoxon rank sum test), as did their non-expressing siblings (paired mean difference of 1.25 seconds [95.0% CI 0.417, 3.33], p = 0.0256 Wilcoxon rank sum test). There was no difference in the movement to the tank base (Figure 5D; paired mean difference of 16.4 [95.0%CI 1.02, 41.2], p = 0.530 for ΔCLK fish and 12.6 [95.0%CI -0.997, 28.1], p = 0.182 for siblings).

To monitor expectation of danger (or anxiety-like behaviour), individual fish were exposed to the alarm substance in a beaker, then transferred to a novel tank containing fresh water (Figure 5E). ΔCLK fish, in contrast to their non-expressing siblings, did not freeze, but behaved similarly to fish that had been exposed to fish water (Figure 5F, G)). We also monitored the post-exposure response of larval zebrafish, using a light-dark choice assay as an indicator of stress (Steenbergen et al., 2011) (Figure 5H). Larval *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* fish displayed less dark avoidance after transient exposure to Schreckstoff, as indicated by the increased amount of time spent in the dark side of the tank (Figure 5I, J; mean difference of 0.0971 [95.0% CI 0.0475, 0.146], p = 0.001 permutation t-test; Cohen's d = 1.16 [95% CI 0.481, 1.82]), in contrast to their non-expressing siblings (mean difference of 0.0213 [95.0% CI -0.058, 0.0928], p = 0.588; Cohen's d = 0.224 [95% CI -0.667, 10.8]). Together, these observations suggest that disruption of the habenula clock leaves acute responses unaffected but reduces the expectation of danger that is evoked by transient exposure to the alarm substance.

We next asked whether the expectation of danger induced by the alarm substance varies in a circadian fashion in zebrafish; a circadian change has been noted in another species, *Rasbora heteromorpha* (Thines and Vandenbussche, 1966). To do this, we exposed larval fish to the alarm substance either at ZT3 or ZT15. Larval zebrafish respond to the alarm substance by extended immobility (Jesuthasan et al., 2020). To sensitively detect this, a vibration was

presented five minutes before and after delivery of the alarm substance (Figure 6A). Control

zebrafish showed a reduction in startle after delivery of the alarm substance during the day

(ZT3, p = 0.00248, Pearson's chi-squared test in R), but not at night (ZT15, Figure 6B),

consistent with freezing during the day but not at night. *Tg(gng8:GAL4, UAS:EGFP-*

Δclocka) fish did not show a reduction in startle at day or at night (Figure 6C). These data

suggest that the extended freezing evoked by the alarm response is circadian in zebrafish and

is influenced by the habenula clock.

3.5 Expression of ΔCLK in the habenula does not affect pineal-regulated

circadian behaviour

We hypothesized that while the circadian clock in the habenula affects the response to a

stimulus that influences expectation, circadian behaviours that are under the control of the

pineal clock (Livne et al., 2016) should be unaffected. To test this, we analysed the rhythmic

locomotor activity of ΔCLK-positive larvae and their non-expressing siblings. All larvae

were entrained to the standard LD cycle and their locomotion was monitored for a few hours

during the light phase, followed by the onset of darkness at ZT14 for activity tracking during

the night. Subsequently, the fish were monitored for one day under constant darkness (DD),

in order to eliminate the masking effects of light and dark on clock-regulated behaviour

(Livne et al., 2016). As seen in Figure 7A-D, ΔCLK larvae showed a similar pattern of

locomotion as their non-expressing siblings, becoming mobile during the light phase and

during the subjective day, while exhibiting quiescent periods of inactivity during the 1st and

2nd night of the recording.

In addition to locomotion, we also examined arousal threshold, which is normally elevated

during the sleep-like state at night. Arousal threshold was measured by responsiveness to

auditory stimuli (Zhdanova et al., 2001). Nine different intensities were delivered between

ZT17 and ZT23, and a sudden change in locomotion was taken as an indicator of

responsiveness. A larva was scored as a responder if it moved more than 7 pixels after the

stimulus and the average percentage of responders was calculated for each stimulus intensity

per group for each experimental run (N = 5 experimental runs). Overall, no difference was

seen between ΔCLK larvae and their control siblings (Figure 7E, F). Together, these

experiments indicate that disruption of the habenula clock has relatively specific effects.

4. Discussion

Predictive coding has emerged in recent years as a highly effective paradigm for understanding how the brain functions (Clark, 2013; Friston, 2018). It was applied initially to sensory processing, including the visual system (Rao and Ballard, 1999), and has since been extended to include cognition and emotion (Barrett, 2017). This framework can also be used to understand psychiatric disorders such as depression and other stress-related disorders (Kaye and Krystal, 2020; Smith et al., 2021; Wilkinson et al., 2017). Given the broad relevance of predictive coding, a knowledge of the anatomical and physiological mechanisms underlying its implementation is of considerable interest. A key feature of predictive coding is the updating of expectations based on experience, and this involves the release of neuromodulators that indicate the sign and magnitude of error, as well as precision of prediction. The habenula is one regulator of these modulators, and the pattern of evoked activity and effects of manipulation are consistent with a role in predictive coding underlying motivated behaviour and response to stressors. Recent work has identified a number of mechanisms, such as regulation of membrane potential by astroglial cells (Cui et al., 2018), that influence habenula function by altering the level of spontaneous activity. Spontaneous activity determines the output of a network (Arieli et al., 1996) and is an important element of predictive coding (Hartmann et al., 2015). Here, we propose that the molecular clock, which influences spontaneous activity (Harvey et al., 2020), contributes to habenula function in predictive coding.

We have found that habenula clock in zebrafish is transcriptionally complex, consisting of multiple paralogs. Given the potential redundancy provided by this, we used expression of a truncated *clocka* gene (Dekens and Whitmore, 2008) to investigate the function of the habenula clock. This strategy is based on the discovery in mice that a truncated *clock* gene disrupts the circadian clock (Vitaterna et al., 1994), whereas a null mutant does not (DeBruyne et al., 2006). Expression of this antimorphic allele in the zebrafish pineal was previously shown to disruption in the expression of clock-regulated genes and in locomotion and sleep (Livne et al., 2016), indicating that it is an effective technique. Expression in the habenula appears to affect the molecular clock, as indicated by the change in *per3* gene expression. We did not observe a complete flattening of *per3* expression across the circadian cycle, however, indicating that inhibition is partial. While consistent with previous findings (Dekens and Whitmore, 2008), this level of inhibition appears sufficient to affect habenula

function, as indicated by change in day-night variation in neuromodulators that are under the
control of the habenula.

We observed that the freezing behaviour evoked by the alarm pheromone is lower at night
compared to the day, and this difference is reduced by expression of the truncated clock gene
specifically in the habenula. We also observed that dark avoidance induced by the alarm
substance is reduced by manipulation of the habenula clock. Thus, the data here are
consistent with the hypothesis that the habenula clock influences expectation of danger.
Clock gene expression has been reported in the habenula of other vertebrates, including mice
(Salaberry et al., 2019). A broad expression of a functional clock in the mammalian habenula
is consistent with electrical recordings obtained from slices, which show a change in firing
rate and resting membrane potential across the circadian cycle in both medial and lateral
habenula (Sakhi et al., 2014a, 2014b; Zhao and Rusak, 2005). The ability of an intrinsic clock
to influence habenula-dependent expectations may thus be conserved across vertebrates. This
represents a distinct mechanism from the role of clocks in other regions in the regulation of
stress (Koch et al., 2017). One implication of the findings here is that identifying clock-
regulated genes that regulate spontaneous activity in the habenula may be relevant in the
treatment of stress related disorders.

Acknowledgements

We thank Yoav Gothilf for providing the pT2-UAS:EGFP-2A-ΔCLK-5xMYC construct and
Marnie Halpern for providing the *TgBAC(gng8:GAL4)^{c426}* line. We also thank Hugh Piggins,
David Lyons, Caroline Wee and Ajay Mathuru for comments and discussion, and Erik
Meijering for FeatureJ. This work was funded by grants from the Singapore Ministry of
Education under its Academic Research Fund Tier 2 (MOE2017-T2-058) and the National
Research Foundation (NRF2017-NRF-ISF002-2676) to SJ, an ARAP fellowship from A*Star
to AB, and a Start-Up Grant from the Lee Kong Chian School of Medicine to SRL. The
funding agencies played no role in the study design or decision to publish.

Declaration of competing interests

685 The authors declare that no competing interests exist.

Author contributions

687 AB, RKC, JCSM, GTJH and SJ performed experiments. AB, RKC, JCSM, S, GD, SL and SJ
688 analysed data. AB and SJ designed the study and wrote the manuscript.

Figures

Figure 1. Clock genes expression in the zebrafish habenula based on single cell RNA sequencing

A) Dot plot showing mean log-normalized expression per cluster and percentage of cells per cluster expressing each gene in the SmartSeq-2 dataset from Pandey et al. **B)** t-SNE representation of clustering results in the SmartSeq-2 dataset. Each point is a cell. **C)** same as

A) but in the 10X dataset. **D)** same as C but in the 10X dataset. The cluster assignment for the

10X dataset corresponds to those assigned by Pandey et al 2018.

Figure 2. Dynamics of clock gene expression in the zebrafish habenula. (A-C). Single

optical sections of the brain of 6 day old larvae fixed at six different time points, labelled for

arltn1b (A) and *per3* (A) using hybridization chain reaction. **(C)** Merge of (A) and (B).

Arrowheads indicate the habenula. **(D)** Ratio of *arntl1b/per3* in the habenula of fish in a

normal LD cycle. **(E-G)** Expression of *per3* and *arntl1b* in the habenula of fish grown under

DD conditions. (**H-M**) *cry1a* expression in the habenula of fish grown under DD conditions.

Images in E and F are maximum projections of the entire habenula. Images in panels H-M are

maximum projections spanning 5 µm. The arrowheads indicate autofluorescent blood cells.

A, B, C, E , F, H-M are dorsal views with anterior to the top. The dots in panels D, G and N

indicate values for individual fish. The horizontal bars indicate illumination conditions. lHb:

left habenula; rHb: right habenula; Pi: pineal; Pa: pallium; OT: optic tectum. Scale bar = 25

µm.

Figure 3. Circadian variation in cytoplasmic calcium in habenula neurons. (**A-E**) Long

duration multiplane recording of the habenula of *Et(GAL4s1011t),Tg(UAS:GCaMP6s)* larvae

in constant darkness. (A-D) Examples of imaging data from one fish at two different time

points, ZT5 (A, B) and ZT14 (C, D), and two different focal planes, dorsal (A, C) and ventral

(B, D). Scale bar = 25 µm, dorsal view with anterior to the top. (E) Average level of

cytoplasmic calcium in the habenula, as shown by intensity of GCaMP6s. Each coloured

trace represents one fish. The thicker black line is the mean. There is a reduction in

intracellular calcium levels during the subjective night. (F-G) Difference in relative activity

levels of habenula neurons between day (ZT3) and night (ZT15) as measured by both

cytoplasmic and nuclear GCaMP6s. (F) Distribution of the mean of vectors containing z-

scores of neurons with high variance in their activity. (G) K-means cluster distribution for the

same vectors.

Figure 4. The effects of expressing truncated *clocka* in the habenula. (**A**) Specific

expression of EGFP in the habenula of a *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* fish. (**B**) At

higher magnification, EGFP is visible in the cytoplasm of habenula neurons. (**C**) Detection of

Myc, which is fused to the C-terminus of ΔCLK, in the habenula of transgenic fish. **(D)** Level of *per3* in the habenula of fish with mosaic expression of EGFP-ΔCLK, as determined by qHCR imaging. The lower graph shows Cohen's d, indicating the degree of difference between ΔCLK-expressing and non-expressing cells at each time point. Values are given in the main text. **(E)** Effects of ΔCLK expression in the habenula on global neuromodulator levels. Plot showing the amount of secreted serotonin, dopamine and acetylcholine in the whole brain of and *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* and sibling fish (N=3) at ZT3 and ZT15. Fish with ΔCLK expression in the habenula have a reduced day-night change in levels. Scale bar = 100 μm in panel A, 20 μm in panels B and C; *** indicates $p < 0.0002$ by permutation t-test.

Figure 5. The effects of the alarm substance *Schreckstoff* on *Tg(gng8:GAL4,*

***UAS:EGFP-Δclocka)* fish**. (A-D) Acute effects in adult fish (N=12). (A) Schematic of the

experiment. (B) Speed distribution in fish before and after exposure to the alarm substance,

where the substance remained in the tank after delivery. (C-D) Time spent darting (C),

restricted to the tank base (D) in the presence of the alarm substance. As indicated by the

paired mean difference plots (lower plot), fish with and without expression of truncated

clocka behave similarly. (E-G) Post-exposure behaviour in adult fish (N=8). (E) Schematic

diagram of the experiment. (F) Speed distribution in a novel tank containing clean water,

after exposure to alarm substance or clean water. (G) Only sibling fish exhibited freezing

after exposure to alarm substance. (H-J) Post-exposure behaviour in 2 week-old zebrafish.

(H) Schematic diagram of the experiment. (I) Percentage of time in the dark area of a tank,

after 10 minutes of exposure to embryo water. (J) Percentage of time spent in the dark side

after exposure to the alarm substance. *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* fish spend more

time in the dark side, with a Cohen's D of 1.2.

Figure 6. Day-night variation in the response of zebrafish to *Schreckstoff*. (A) Schematic

of the experiment. To assess freezing in larval fish following exposure to the alarm

substance, fish were exposed to vibrations in the day (ZT3) (B) or at night (ZT15) (C).

Control fish, which are non-expressing siblings of *Tg(gng8:GAL4, UAS:Δclocka)* fish, show

a reduction in startle after exposure during the day, but not at night. This response to alarm

substance was not seen in *Tg(gng8:GAL4, UAS:Δclocka)* fish in the day or night.

Figure 7. Sleep is unaltered in *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* larvae

(**A, B**) Distances moved (cm/10min) by individual larvae from (A) Sibling group and (B)

ΔCLK group are plotted against time (hr) over the time-course of the whole recording. CT =

circadian time. (**C, D**) Mean distances moved (cm/10min) of ΔCLK (blue) and Sibling

(orange) groups are plotted against time (hr) (C) during the first 14 hours of recording,

including 4 hours of light-phase and 10 hours in the dark during the night and (D) during the

subsequent 24 hours of recording under DD. 95% CIs are indicated by the shaded areas;

horizontal brackets indicate 5 analyzed timeframes. Black and white bars represent periods of

dark and light, respectively. (**E**) The Cumming estimation plot shows the mean differences in

5 timeframes of the recording: ZT10-14 (light-phase), ZT14-24 (1st night), CT0-7 (early

subjective day), CT7-14 (late subjective day) and CT14-24 (subjective night). The upper axis

shows the raw data, with each dot representing the overall distance moved (cm) by a single

larva; gapped vertical lines to the right of each group indicate mean ± SD. Each mean

difference is plotted on the lower axes as a bootstrap sampling distribution. Mean differences

are depicted as black dots and 95% CIs are indicated by the end of vertical error bars. (**F**)

Analysis of sensory responsiveness of the *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* larvae (7

dpf) during the delivery of auditory stimuli of 9 different intensities. The Cumming

estimation plot shows the mean differences for 9 comparisons, corresponding to 9 stimulus

intensities (0-100%). The raw data is plotted on the upper axis, each dot representing the

percentage of responders per single experimental run.

Figure S1. Examples of *Tg(gng8:GAL4, UAS:EGFP-Δclocka)* larval zebrafish with broad (A,

B) and sparse (C, D) expression of EGFP in the habenula. All images are dorsal view, with

anterior to the top. Scale bar = 10 μm.

References

Agetsuma, M., Aizawa, H., Aoki, T., Nakayama, R., Takahoko, M., Goto, M., Sassa, T.,
Amo, R., Shiraki, T., Kawakami, K., Hosoya, T., Higashijima, S., Okamoto, H., 2010. The
habenula is crucial for experience-dependent modification of fear responses in zebrafish.
Nature neuroscience 13, 1354–1356. https://doi.org/10.1038/nn.2654

Amezquita, R.A., Lun, A.T.L., Becht, E., Carey, V.J., Carpp, L.N., Geistlinger, L., Marini,
F., Rue-Albrecht, K., Risso, D., Soneson, C., Waldron, L., Pagès, H., Smith, M.L., Huber,
W., Morgan, M., Gottardo, R., Hicks, S.C., 2020. Orchestrating single-cell analysis with
Bioconductor. Nat Methods 17, 137–145. https://doi.org/10.1038/s41592-019-0654-x

Amo, R., Aizawa, H., Takahoko, M., Kobayashi, M., Takahashi, R., Aoki, T., Okamoto, H.,
2010. Identification of the zebrafish ventral habenula as a homolog of the mammalian
lateral habenula. The Journal of neuroscience : the official journal of the Society for
Neuroscience 30, 1566–1574. https://doi.org/10.1523/jneurosci.3690-09.2010

Amo, R., Fredes, F., Kinoshita, M., Aoki, R., Aizawa, H., Agetsuma, M., Aoki, T., Shiraki,
T., Kakinuma, H., Matsuda, M., Yamazaki, M., Takahoko, M., Tsuboi, T., Higashijima,
S., Miyasaka, N., Koide, T., Yabuki, Y., Yoshihara, Y., Fukai, T., Okamoto, H., 2014.
The Habenulo-Raphe Serotonergic Circuit Encodes an Aversive Expectation Value
Essential for Adaptive Active Avoidance of Danger. Neuron 84, 1034–1048.
https://doi.org/10.1016/j.neuron.2014.10.035

Andalman, A.S., Burns, V.M., Lovett-Barron, M., Broxton, M., Poole, B., Yang, S.J.,
Grosenick, L., Lerner, T.N., Chen, R., Benster, T., Mourrain, P., Levoy, M., Rajan, K.,
Deisseroth, K., 2019. Neuronal Dynamics Regulating Brain and Behavioral State
Transitions. Cell 177, 970-985.e20. https://doi.org/10.1016/j.cell.2019.02.037

Arieli, A., Sterkin, A., Grinvald, A., Aertsen, A., 1996. Dynamics of Ongoing Activity:
Explanation of the Large Variability in Evoked Cortical Responses. Science 273, 1868–
1871. https://doi.org/10.1126/science.273.5283.1868

Baño-Otálora, B., Piggins, H.D., 2017. Contributions of the lateral habenula to circadian
timekeeping. Pharmacology, biochemistry, and behavior 162, 46–54.
https://doi.org/10.1016/j.pbb.2017.06.007

Barrett, L.F., 2017. The theory of constructed emotion: an active inference account of
interoception and categorization. Soc Cogn Affect Neur 12, 1–23.
https://doi.org/10.1093/scan/nsw154

Berkes, P., Orbán, G., Lengyel, M., Fiser, J., 2011. Spontaneous Cortical Activity Reveals
Hallmarks of an Optimal Internal Model of the Environment. Science (New York, NY)
331, 83–87. https://doi.org/10.1126/science.1195870

Bolte, S., Cordelières, F.P., 2006. A guided tour into subcellular colocalization analysis in
light microscopy. J Microsc-oxford 224, 213–232.

Burgess, H.A., Granato, M., 2007. Modulation of locomotor activity in larval zebrafish
during light adaptation. The Journal of experimental biology 210, 2526–2539.
https://doi.org/10.1242/jeb.003939

Cavallari, N., Frigato, E., Vallone, D., Fröhlich, N., Lopez-Olmeda, J.F., Foà, A., Berti, R.,
Sánchez-Vázquez, F.J., Bertolucci, C., Foulkes, N.S., 2011. A Blind Circadian Clock in
Cavefish Reveals that Opsins Mediate Peripheral Clock Photoreception. Plos Biol 9,
e1001142. https://doi.org/10.1371/journal.pbio.1001142

Chatterjee, D., Gerlai, R., 2009. High precision liquid chromatography analysis of
dopaminergic and serotoninergic responses to acute alcohol exposure in zebrafish. Behav
Brain Res 200, 208–213.

Cheng, R.-K., Krishnan, S., Jesuthasan, S., 2016. Activation and inhibition of tph2
serotonergic neurons operate in tandem to influence larval zebrafish preference for light
over darkness. Sci Rep-uk 6, 20788. https://doi.org/10.1038/srep20788

Chia, J.S.M., Wall, E.S., Wee, C.L., Rowland, T.A.J., Cheng, R.-K., Cheow, K., Guillemin,
K., Jesuthasan, S., 2019. Bacteria evoke alarm behaviour in zebrafish. Nat Commun 10,
3831. https://doi.org/10.1038/s41467-019-11608-9

Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J.,
Cunha, A., Pierce, N.A., 2018. Third-generation in situ hybridization chain reaction:
multiplexed, quantitative, sensitive, versatile, robust. Development 145, dev165753.
https://doi.org/10.1242/dev.165753

Clark, A., 2013. Whatever next? Predictive brains, situated agents, and the future of cognitive
science. Behav Brain Sci 36, 181–204. https://doi.org/10.1017/s0140525x12000477

Csardi, G., Nepusz, T., 2006. The igraph software package for complex network research.
InterJournal Complex Systems, 1695.

Cui, Y., Yang, Y., Ni, Z., Dong, Y., Cai, G., Foncelle, A., Ma, S., Sang, K., Tang, S., Li, Y.,
Shen, Y., Berry, H., Wu, S., Hu, H., 2018. Astroglial Kir4.1 in the lateral habenula drives
neuronal bursts in depression. Nature 554, 323–327. https://doi.org/10.1038/nature25752

DeBruyne, J.P., Noton, E., Lambert, C.M., Maywood, E.S., Weaver, D.R., Reppert, S.M.,
2006. A Clock Shock: Mouse CLOCK Is Not Required for Circadian Oscillator Function.
Neuron 50, 465–477. https://doi.org/10.1016/j.neuron.2006.03.041

Dekens, M.P., Whitmore, D., 2008. Autonomous onset of the circadian clock in the zebrafish
embryo. The EMBO Journal 27, 2757–2765. https://doi.org/10.1038/emboj.2008.183

Delaunay, F., Thisse, C., Thisse, B., Laudet, V., 2003. Differential regulation of Period 2 and
Period 3 expression during development of the zebrafish circadian clock. Gene Expression
Patterns 3, 319–324. https://doi.org/10.1016/s1567-133x(03)00050-4

Friston, K., 2018. Does predictive coding have a future? Nat Neurosci 21, 1019–1021.
https://doi.org/10.1038/s41593-018-0200-7

Guilding, C., Hughes, A.T.L., Piggins, H.D., 2010. Circadian oscillators in the epithalamus.
Neuroscience 169, 1630–1639. https://doi.org/10.1016/j.neuroscience.2010.06.015

Haghani, S., Karia, M., Cheng, R.-K., Mathuru, A.S., 2019. An Automated Assay System to
Study Novel Tank Induced Anxiety. Front Behav Neurosci 13, 180.
https://doi.org/10.3389/fnbeh.2019.00180

Haghverdi, L., Lun, A.T.L., Morgan, M.D., Marioni, J.C., 2018. Batch effects in single-cell
RNA-sequencing data are corrected by matching mutual nearest neighbors. Nat
Biotechnol 36, 421–427. https://doi.org/10.1038/nbt.4091

Hartmann, C., Lazar, A., Nessler, B., Triesch, J., 2015. Where's the Noise? Key Features of
Spontaneous Activity and Neural Variability Arise through Learning in a Deterministic
Network. PLoS Computational Biology 11, e1004640.
https://doi.org/10.1371/journal.pcbi.1004640

Harvey, J.R.M., Plante, A.E., Meredith, A.L., 2020. Ion Channels Controlling Circadian
Rhythms in Suprachiasmatic Nucleus Excitability. Physiol Rev 100, 1415–1454.
https://doi.org/10.1152/physrev.00027.2019

Ho, J., Tumkaya, T., Aryal, S., Choi, H., Claridge-Chang, A., 2019. Moving beyond P
values: data analysis with estimation graphics. Nature methods 16, 565–566.
https://doi.org/10.1038/s41592-019-0470-3

Hong, E., Santhakumar, K., Akitake, C.A., Ahn, S.J., Thisse, C., Thisse, B., Wyart, C.,
Mangin, J.-M., Halpern, M.E., 2013. Cholinergic left-right asymmetry in the habenulo-
interpeduncular pathway. Proc National Acad Sci 110, 21171–21176.
https://doi.org/10.1073/pnas.1319566110

Hsu, Y.-W.A., Wang, S.D., Wang, S., Morton, G., Zariwala, H.A., Iglesia, H.O. de la,
Turner, E.E., 2014. Role of the dorsal medial habenula in the regulation of voluntary
activity, motor function, hedonic state, and primary reinforcement. The Journal of
neuroscience : the official journal of the Society for Neuroscience 34, 11366–11384.
https://doi.org/10.1523/jneurosci.1861-14.2014

Hut, R.A., Zee, E.A.V. der, 2011. The cholinergic system, circadian rhythmicity, and time
memory. Behav Brain Res 221, 466–480. https://doi.org/10.1016/j.bbr.2010.11.039

Jesuthasan, S., Krishnan, S., Cheng, R.-K., Mathuru, A., 2020. Neural correlates of state
transitions elicited by a chemosensory danger cue. Prog Neuro-psychopharmacology
Biological Psychiatry 110110. https://doi.org/10.1016/j.pnpbp.2020.110110

Jesuthasan, S.J., Mathuru, A.S., 2008. The alarm response in zebrafish: innate fear in a
vertebrate genetic model. Journal of Neurogenetics 22, 211–228.
https://doi.org/10.1080/01677060802298475

Jetti, S.K., Vendrell-Llopis, N., Yaksi, E., 2014. Spontaneous activity governs olfactory
representations in spatially organized habenular microcircuits. Current biology : CB 24,
434–439. https://doi.org/10.1016/j.cub.2014.01.015

Johnsson, K., Soneson, C., Fontes, M., 2014. Low Bias Local Intrinsic Dimension Estimation
from Expected Simplex Skewness. Ieee T Pattern Anal 37, 196–202.
https://doi.org/10.1109/tpami.2014.2343220

Kaye, A.P., Krystal, J.H., 2020. Predictive Processing in Mental Illness: Hierarchical
Circuitry for Perception and Trauma. J Abnorm Psychol 129, 629–632.
https://doi.org/10.1037/abn0000628

Kim, U., Chang, S.-Y., 2005. Dendritic morphology, local circuitry, and intrinsic
electrophysiology of neurons in the rat medial and lateral habenular nuclei of the
epithalamus. The Journal of comparative neurology 483, 236–250.
https://doi.org/10.1002/cne.20410

Kobayashi, Y., Ishikawa, T., Hirayama, J., Daiyasu, H., Kanai, S., Toh, H., Fukuda, I.,
Tsujimura, T., Terada, N., Kamei, Y., Yuba, S., Iwai, S., Todo, T., 2000. Molecular
analysis of zebrafish photolyase/cryptochrome family: two types of cryptochromes present
in zebrafish. Genes Cells 5, 725–738. https://doi.org/10.1046/j.1365-2443.2000.00364.x

Koch, C.E., Leinweber, B., Drengberg, B.C., Blaum, C., Oster, H., 2017. Interaction between
circadian rhythms and stress. Neurobiology Stress 6, 57–67.
https://doi.org/10.1016/j.ynstr.2016.09.001

Lee, A., Mathuru, A.S., Teh, C., Kibat, C., Korzh, V., Penney, T.B., Jesuthasan, S., 2010.
The habenula prevents helpless behavior in larval zebrafish. Current biology 20, 2211–
2216. https://doi.org/10.1016/j.cub.2010.11.025

Leung, L.C., Wang, G.X., Madelaine, R., Skariah, G., Kawakami, K., Deisseroth, K., Urban,
A.E., Mourrain, P., 2019. Neural signatures of sleep in zebrafish. Nature 571, 198–204.
https://doi.org/10.1038/s41586-019-1336-7

Livne, Z.B.-M., Alon, S., Vallone, D., Bayleyen, Y., Tovin, A., Shainer, I., Nisembaum,
L.G., Aviram, I., Smadja-Storz, S., Fuentes, M., Falcón, J., Eisenberg, E., Klein, D.C.,
Burgess, H.A., Foulkes, N.S., Gothilf, Y., 2016. Genetically Blocking the Zebrafish Pineal
Clock Affects Circadian Behavior. PLoS genetics 12, e1006445.
https://doi.org/10.1371/journal.pgen.1006445

Lun, A., 2020a. BiocSingular: Singular Value Decomposition for Bioconductor Packages. R
package version 1.6.0.

Lun, A., 2020b. bluster: Clustering Algorithms for Bioconductor.

Lun, A.T.L., McCarthy, D.J., Marioni, J.C., 2016. A step-by-step workflow for low-level
analysis of single-cell RNA-seq data with Bioconductor. F1000research 5, 2122.
https://doi.org/10.12688/f1000research.9501.2

Mathuru, A.S., Kibat, C., Cheong, W.F., Shui, G., Wenk, M.R., Friedrich, R.W., Jesuthasan,
S., 2012. Chondroitin Fragments Are Odorants that Trigger Fear Behavior in Fish. Current
biology : CB 22, 538–544. https://doi.org/10.1016/j.cub.2012.01.061

Matias, S., Lottem, E., Dugué, G.P., Mainen, Z.F., 2017. Activity patterns of serotonin
neurons underlying cognitive flexibility. Elife 6, e20552.
https://doi.org/10.7554/elife.20552

Matsumoto, M., Hikosaka, O., 2009. Representation of negative motivational value in the
primate lateral habenula. Nature neuroscience 12, 77–84. https://doi.org/10.1038/nn.2233

Maximino, C., Lima, M.G., Costa, C.C., Guedes, I.M.L., Herculano, A.M., 2014. Fluoxetine
and WAY 100,635 dissociate increases in scototaxis and analgesia induced by conspecific
alarm substance in zebrafish (Danio rerio Hamilton 1822). Pharmacology, biochemistry,
and behavior 124, 425–433. https://doi.org/10.1016/j.pbb.2014.07.003

McCarthy, D.J., Campbell, K.R., Lun, A.T.L., Wills, Q.F., 2017. Scater: pre-processing,
quality control, normalization and visualization of single-cell RNA-seq data in R.
Bioinformatics btw777. https://doi.org/10.1093/bioinformatics/btw777

Mendoza, J., 2017. Circadian neurons in the lateral habenula: Clocking motivated behaviors.
Pharmacology, biochemistry, and behavior 162, 55–61.
https://doi.org/10.1016/j.pbb.2017.06.013

Moran, R.J., Campo, P., Symmonds, M., Stephan, K.E., Dolan, R.J., Friston, K.J., 2013. Free
Energy, Precision and Learning: The Role of Cholinergic Neuromodulation. J Neurosci
33, 8227–8236. https://doi.org/10.1523/jneurosci.4255-12.2013

Pandey, S., Shekhar, K., Regev, A., Schier, A.F., 2018. Comprehensive Identification and
Spatial Mapping of Habenular Neuronal Types Using Single-Cell RNA-Seq. Current
biology : CB 28, 1052-1065.e7. https://doi.org/10.1016/j.cub.2018.02.040

Pennartz, C.M.A., Jeu, M.T.G. de, Bos, N.P.A., Schaap, J., Geurtsen, A.M.S., 2002. Diurnal
modulation of pacemaker potentials and calcium current in the mammalian circadian
clock. Nature 416, 286–290. https://doi.org/10.1038/nature728

Rao, R.P.N., Ballard, D.H., 1999. Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. Nat Neurosci 2, 79–87.
https://doi.org/10.1038/4580

Ren, J., Qin, C., Hu, F., Tan, J., Qiu, L., Zhao, S., Feng, G., Luo, M., 2011. Habenula
"cholinergic" neurons co-release glutamate and acetylcholine and activate postsynaptic
neurons via distinct transmission modes. Neuron 69, 445–452.
https://doi.org/10.1016/j.neuron.2010.12.038

Sakhi, K., Belle, M.D.C., Gossan, N., Delagrange, P., Piggins, H.D., 2014a. Daily variation
in the electrophysiological activity of mouse medial habenula neurones. The Journal of
Physiology 592, 587–603. https://doi.org/10.1113/jphysiol.2013.263319

Sakhi, K., Wegner, S., Belle, M.D.C., Howarth, M., Delagrange, P., Brown, T.M., Piggins,
H.D., 2014b. Intrinsic and extrinsic cues regulate the daily profile of mouse lateral
habenula neuronal activity. The Journal of physiology 592, 5025–5045.
https://doi.org/10.1113/jphysiol.2014.280065

Salaberry, N.L., Hamm, H., Felder-Schmittbuhl, M.-P., Mendoza, J., 2019. A
suprachiasmatic-independent circadian clock(s) in the habenula is affected by Per gene
mutations and housing light conditions in mice. Brain Struct Funct 224, 19–31.
https://doi.org/10.1007/s00429-018-1756-4

Satar, N.M.A., Ogawa, S., Parhar, I.S., 2020. Kisspeptin-1 regulates forebrain dopaminergic
neurons in the zebrafish. Sci Rep-uk 10, 19361. https://doi.org/10.1038/s41598-020-
75777-0

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source
platform for biological-image analysis. Nat Methods 9, 676–682.
https://doi.org/10.1038/nmeth.2019

Schultz, W., 1998. Predictive reward signal of dopamine neurons. Journal of
neurophysiology 80, 1–27.

Scott, E.K., Baier, H., 2009. The Cellular Architecture of the Larval Zebrafish Tectum, as
Revealed by Gal4 Enhancer Trap Lines. Front Neural Circuit 3, 13.
https://doi.org/10.3389/neuro.04.013.2009

Smith, R., Badcock, P., Friston, K.J., 2021. Recent advances in the application of predictive
coding and active inference models within clinical neuroscience. Psychiat Clin Neuros 75,
3–13. https://doi.org/10.1111/pcn.13138

Speedie, N., Gerlai, R., 2008. Alarm substance induced behavioral responses in zebrafish
(Danio rerio). Behav Brain Res 188, 168–177. https://doi.org/10.1016/j.bbr.2007.10.031

Steenbergen, P.J., Richardson, M.K., Champagne, D.L., 2011. Patterns of avoidance
behaviours in the light/dark preference test in young juvenile zebrafish: a pharmacological
study. Behav Brain Res 222, 15–25. https://doi.org/10.1016/j.bbr.2011.03.025

Stringer, C., Pachitariu, M., 2019. Computational processing of neural recordings from
calcium imaging data. Curr Opin Neurobiol 55, 22–31.
https://doi.org/10.1016/j.conb.2018.11.005

Takahashi, J.S., 2017. Transcriptional architecture of the mammalian circadian clock. Nat
Rev Genet 18, 164–179. https://doi.org/10.1038/nrg.2016.150

Tamai, T.K., Young, L.C., Whitmore, D., 2007. Light signaling to the zebrafish circadian
clock by Cryptochrome 1a. Proc National Acad Sci 104, 14712–14717.
https://doi.org/10.1073/pnas.0704588104

Thines, G., Vandenbussche, E., 1966. The effects of alarm substance on the schooling
behaviour of rasbora heteromorpha Duncker in day and night conditions. Anim Behav 14,
296–302. https://doi.org/10.1016/s0003-3472(66)80086-6

Trivedi, V., Choi, H.M.T., Fraser, S.E., Pierce, N.A., 2018. Multidimensional quantitative
analysis of mRNA expression within intact vertebrate embryos. Development 145,
dev156869. https://doi.org/10.1242/dev.156869

Vitaterna, M., King, D., Chang, A., Kornhauser, J., Lowrey, P., McDonald, J., Dove, W.,
Pinto, L., Turek, F., Takahashi, J., 1994. Mutagenesis and mapping of a mouse gene,
Clock, essential for circadian behavior. Science 264, 719–725.
https://doi.org/10.1126/science.8171325

Wang, H., Yang, Z., Li, X., Huang, D., Yu, S., He, J., Li, Y., Yan, J., 2020. Single-cell in
vivo imaging of cellular circadian oscillators in zebrafish. Plos Biol 18, e3000435.
https://doi.org/10.1371/journal.pbio.3000435

Weger, M., Weger, B.D., Diotel, N., Rastegar, S., Hirota, T., Kay, S.A., Strähle, U.,
Dickmeis, T., 2013. Real-time in vivo monitoring of circadian E-box enhancer activity: A
robust and sensitive zebrafish reporter line for developmental, chemical and neural
biology of the circadian clock. Developmental Biology 380, 259–273.
https://doi.org/10.1016/j.ydbio.2013.04.035

Wilkinson, S., Dodgson, G., Meares, K., 2017. Predictive Processing and the Varieties of
Psychological Trauma. Front Psychol 08, 1840. https://doi.org/10.3389/fpsyg.2017.01840

Woods, I.G., Schoppik, D., Shi, V.J., Zimmerman, S., Coleman, H.A., Greenwood, J., Soucy,
E.R., Schier, A.F., 2014. Neuropeptidergic signaling partitions arousal behaviors in
zebrafish. The Journal of neuroscience : the official journal of the Society for
Neuroscience 34, 3142–3160. https://doi.org/10.1523/jneurosci.3529-13.2014

Yang, Y., Cui, Y., Sang, K., Dong, Y., Ni, Z., Ma, S., Hu, H., 2018. Ketamine blocks
bursting in the lateral habenula to rapidly relieve depression. Nature 554, 317–322.
https://doi.org/10.1038/nature25509

Zappia, L., Oshlack, A., 2018. Clustering trees: a visualisation for evaluating clusterings at
multiple resolutions. Gigascience 7, giy083-. https://doi.org/10.1093/gigascience/giy083

Zhao, H., Rusak, B., 2005. Circadian firing-rate rhythms and light responses of rat habenular
nucleus neurons in vivo and in vitro. 132, 519–528.
https://doi.org/10.1016/j.neuroscience.2005.01.012

Zhdanova, I.V., Wang, S.Y., Leclair, O.U., Danilova, N.P., 2001. Melatonin promotes sleep-
like state in zebrafish. Brain research 903, 263–268.

1051