1

# The habenula clock in zebrafish underlies circadian variation in response to a stressor

Adriana Basnakova[1, 2], Ruey-Kuang Cheng[3], Joanne Chia Shu Ming[3] and *Suresh Jesuthasan[2, 3]

[1]School of Biological Sciences, University of Manchester, UK.

[2]Institute of Molecular and Cell Biology, Singapore.

[3]Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore.

*corresponding author: sureshj@ntu.edu.sg

# Abstract

The lateral habenula, a regulator of neuromodulator release, contains a semi-autonomous molecular clock whose significance is unknown. Here, using zebrafish, we investigate the contribution of this clock to change in internal state that is triggered by an aversive stimulus. We focused on the alarm pheromone, *Schreckstoff*, whose effects vary in a circadian manner. Disruption of habenula clock function, by selective expression of a truncated clock gene (ΔCLK), altered the effects of *Schreckstoff* such that larval responses in the day resembled those at night. Acute responses, including immediate change in swim speed, were unaffected, whereas extended behaviours such as freezing in a novel environment and dark avoidance, were reduced. Circadian behaviours that are under the control of the pineal clock, such as locomotion and sleep, were unchanged. These findings suggest that the habenula clock can act independently to influence the internal state induced by an aversive stimulus.

# Introduction

Sensory stimuli evoke a variety of responses in an animal. A stimulus with high salience, for example, will elicit changes in movement and in internal state. The changes in state, which can include increase in arousal and vigilance, are mediated by the release of neuromodulators such as serotonin (Chaouloff et al., 1999) and dopamine (Abercrombie et al., 1989; Holly and Miczek, 2016). One regulator of these neuromodulators in vertebrates, especially in response to aversive stimuli, is the habenula (Namboodiri et al., 2016). The habenula, which is located in the epithalamus, consists of two domains, medial and lateral, with the latter controlling dopamine and serotonin release from the ventral tegmental area (VTA) and raphe, respectively. Exposure to a stressor increases the level of activity in lateral habenula neurons (Andalman et al., 2019), and abnormal firing in the lateral habenula is associated with maladaptive behaviours (Li et al., 2013; Yang et al., 2018). Thus, a knowledge of factors that influence activity in the lateral habenula is important in understanding what shapes the response to stressors.

One such a factor appears to be an intrinsic molecular clock. Components of the clock machinery (Takahashi, 2017), such as *per2* in mouse (Guilding et al., 2010) and rat (Zhao et al., 2015), *per1* and *bmal*1 in zebrafish (Weger et al., 2013), and *per1b* in goldfish (Sánchez-Bretaño et al., 2015), are expressed in a circadian manner in the lateral habenula. Although the

habenula receives input from the master pacemaker in the suprachiasmatic nucleus (SCN)

(Buijs, 1978; Sakhi et al., 2014), the habenula clock persists even when the SCN is lesioned

(Salaberry et al., 2019). In general, the molecular clock drives daily variations in excitability,

enabling neurons to generate rhythmic electrical output (Belle, 2015; Cao and Nitabach, 2008;

Pennartz et al., 2002). In agreement with this, spontaneous firing in the habenula of rat (Zhao

and Rusak, 2005) and mouse (Guilding et al., 2010; Sakhi et al., 2014) varies across a circadian

cycle. It has been shown that the effect of a stressor on lateral habenula electrical activity is

largest during the active period, when spontaneous activity is lowest (Chastrette et al., 1991).

An implication of this is that the habenula clock influences the response to stimuli that are

processed by the habenula. There is, however, no experimental evidence for this.

We hypothesize that the habenula clock acts in an autonomous manner to influence response

to an aversive stimulus. It is known that the response to stressors varies with time of day (Koch

et al., 2017), but the underlying mechanism is unclear. One strategy that can be used to

investigate the function of the molecular clock in a specific cell is to express a truncated

CLOCK protein (ΔCLK) (Dekens and Whitmore, 2008). The molecular clock consists of a

transcriptional-translational feedback loop (Takahashi, 2017). A heterodimer of CLOCK and

BMAL enters the nucleus, where it drives transcription of *period* (*per*) and *cryptochrome* (*cry*)

genes, which then inhibit the heterodimer. The truncated CLOCK protein binds BMAL, but

the resulting heterodimer is unable to activate transcription of target genes. This strategy has

been successfully used in the pineal of zebrafish, where expression of truncated CLOCK

prevented circadian variation in clock-controlled genes in the pineal and led to a disruption of

circadian rhythmicity in locomotion, sleep and vertical position (Ben-Moshe Livne et al.,

2016). Here, we use the GAL4/UAS system to drive expression of ΔCLK specifically in the

habenula of zebrafish. Our results demonstrate that the intrinsic habenula clock has an essential

role in regulating the behavioural response to an aversive stimulus.

## Results

### The zebrafish habenula displays circadian variation in activity

74  The expression of clock genes in the mouse habenula is correlated with circadian changes in

75  neural activity (Guilding et al., 2010; Sakhi et al., 2014). To assess whether there is a circadian

76  change in neural activity in the zebrafish habenula, we used immunohistochemical detection

of phosphorylated extracellular signal-regulated kinase (pERK), which has the advantage of reporting neural activity in freely swimming animals (Randlett et al., 2015); this method has been used to examine circadian activity in the mouse SCN (Hughes et al., 2004). In 8 day old zebrafish, pERK label was evident in both nuclei and the cytoplasm of scattered neurons. Generally, pERK label was more prominent in the lateral (or ventral) habenula compared to the medial (or dorsal) habenula (compare cells indicated with white arrowheads in Figure 1A and B). Label was also seen in neuropils (Figure 1A). To determine if there is any consistent difference in activity between day and night, registered and scaled image data was first normalized by calculating the pERK/total-ERK (tERK) ratio. The normalized image data was then compared between 'Day' and 'Night' groups, which consisted of > 100 zebrafish larvae fixed at Zeitgeber time ZT3 (3 hours after lights on; Figure 1D, G) and ZT15 (Figure 1C, F) respectively. A pixel-wise 'Night'-over-'Day' comparison, implementing a false discovery rate threshold of 0.1%, (Figure 1E, H) indicates that levels of pERK were higher at night relative to the day, especially in the lateral habenula. The day-night difference was also seen in the lateral habenula of fish held in constant darkness (Figure 2), although to a lesser extent than fish that had undergone a normal light-dark cycle. This suggests that, while light has a role, neural activity in the lateral habenula is influenced by a circadian clock.

## Disruption of the habenula clock with ΔCLK

To selectively express ΔCLK in the habenula, we used the GAL4/UAS system with *TgBAC(gng8:GAL4)* (Hong et al., 2013) as the driver. In addition to ΔCLK, the effector construct also included enhanced green fluorescent protein (EGFP), which was separated from ΔCLK by a 2A peptide linker, thus enabling the expression of two separate proteins. The presence of EGFP in the habenula neurons of *Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* fish (Figure 3A, B) indicates that the construct is expressed appropriately and enables convenient identification of ΔCLK+ individuals (Ben-Moshe Livne et al., 2016). Labelling with an antibody against the Myc tag, which is fused in frame at the C-terminus of ΔCLK, indicates that the truncated CLOCK protein is translated (Fig. 3C). To assess whether ΔCLK expression had an effect on physiological function of the habenula, brains were isolated from adult zebrafish expressing EGFP-ΔCLK, and from their non-expressing siblings, at two different time points, ZT3 and ZT15. The supernatant from extracts were analysed by liquid chromatography and mass spectrometry (LC/MS) (Chatterjee and Gerlai, 2009). While non-expressing fish had a clear difference in the day- and night-time levels of dopamine (DA), there

109     was almost no difference in fish expressing ΔCLK protein (Figure 4). The difference in day-

110     night serotonin (5-HT) levels was also reduced in the ΔCLK+ fish, although to a lesser extent.

111     These results indicate that expression of ΔCLK in the habenula is effective in disrupting the

112     intrinsic molecular clock.

## 113     Habenula expression of ΔCLK affects the alarm response

We hypothesized that the behavioural responses to an aversive stimulus would be affected by

the expression of ΔCLK. One aversive stimulus for fish is the alarm pheromone *Schreckstoff*

(Pfeiffer, 1962). This substance elicits sustained activity in the lateral habenula (Jesuthasan et

al., 2020). The alarm response shows circadian variation in harlequin fish (Thines and

Vandenbussche, 1966) and bleak (Jachner, 1996), with a strong response seen primarily in day,

i.e. the active period. To test whether this is also the case in zebrafish, we exposed larval fish

to the alarm substance either at 11 am or 11 pm. Larval zebrafish respond to the alarm substance

by increased immobility (Jesuthasan et al., 2020). To sensitively detect this, a vibration was

presented five minutes before and after delivery of the alarm substance. Control zebrafish

showed a reduction in startle after delivery of the alarm substance during the day, but not at

night (Figure 5), consistent with freezing during the day but not at night. *Tg(gng8:GAL4,*

*UAS:EGFP-ΔCLK)* fish did not show a reduction in startle at day or at night. These data suggest

that the alarm response is circadian in zebrafish and is influenced by the habenula clock.

The habenula is involved in the regulation of neuromodulators, which would be expected to

alter behaviour that reflect change in state, rather than immediate changes in locomotion in

response to a stimulus. In adult zebrafish, *Schreckstoff* causes an acute change in swimming

speed and tank position (Jesuthasan and Mathuru, 2008; Speedie and Gerlai, 2008).

Additionally, fish transiently exposed to *Schreckstoff* display persistent stress, as indicated by

abnormal scototaxis and elevated levels of serotonin (Maximino et al., 2014), as well as

increased alertness (Pfeiffer and Riegelbauer, 1978). To determine whether manipulation of

the habenula clock influenced the acute response, adult fish were placed individually in a test

tank. After a period of acclimatization, the alarm substance was introduced into the tank. Adult

*Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* zebrafish, like their non-expressing siblings, showed an

increase in darting and moved to the base of the tank, with a number of fish then freezing

(Figure 6A-E). To monitor post-exposure effects, individual fish were exposed to the alarm

substance in a beaker, then transferred to a novel tank. ΔCLK fish, in contrast to their non-

expressing siblings, did not freeze, but behaved similarly to fish that had been exposed to fish water (Figure 6F, G). We also monitored the post-exposure behaviour of larval zebrafish, using a light-dark choice assay (Steenbergen et al., 2011). In comparison to their non-expressing siblings, larval *Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* fish displayed less dark avoidance after transient exposure to *Schreckstoff*, as indicated by the amount of time spent in the dark side of the tank (Figure 6H, I). Together, these observations indicate that disruption of the habenula clock leaves acute responses unaffected but reduces the persistent stress response that is evoked by transient exposure to the alarm substance.

## Expression of ΔCLK in the habenula does not affect pineal-regulated circadian behaviour

We hypothesized that while the circadian clock in the habenula affects the response to an aversive stimulus, circadian behaviours that are under the control of the pineal clock (Ben-Moshe Livne et al., 2016) should be unaffected. To test this, we analysed the rhythmic locomotor activity (Prober et al., 2006) of ΔCLK+ larvae and their control (CTRL) siblings. All larvae were entrained to the standard light-dark (LD) cycle and their locomotion was monitored for a few hours during the light phase, followed by the onset of darkness at ZT14 for activity tracking during the night. Subsequently, the fish were monitored for one day under constant darkness (DD), in order to eliminate the masking effects of light and dark on clock-regulated behaviour (Ben-Moshe Livne et al., 2016). As seen in Figure 7, ΔCLK+ larvae showed a similar pattern of locomotion as their non-expressing siblings, becoming mobile during the light phase and during the subjective day, while exhibiting quiescent periods of inactivity during the 1st and 2nd night of the recording.

In addition to locomotion, we also examined arousal threshold, which is normally elevated during the sleep-like state at night. Arousal threshold was measured by responsiveness to auditory stimuli (Zhdanova et al., 2001). Nine different intensities were delivered between ZT17 and ZT23, and a sudden change in locomotion was taken as an indicator of responsiveness (Figure 7F, G). A larva was scored as a responder if it moved more than 7 pixels after the stimulus and the average percentage of responders was calculated for each stimulus intensity per group for each experimental run (N = 5 experimental runs). Overall, no difference was seen between ΔCLK+ larvae and their control siblings (Figure 7I). Together, these experiments indicate that disruption of the habenula clock has relatively specific effects.

## Discussion

All aspects of life, from arousal state (Scammell et al., 2017) to the ability to handle stress (Daut and Fonken, 2019) and respond to rewards (Antle and Silver, 2016; McClung et al., 2005), vary with time of day. This reflects the tight control exerted by the circadian clock on the physiology and behaviour of animals. Clock genes, which drive the circadian clock, are expressed broadly in the brain (Moore and Whitmore, 2014; Shearman et al., 1997; Shieh, 2003; Sun et al., 1997). In this paper, we investigated the hypothesis that the habenula clock influences the circadian variation in response triggered by a stressor. Such stimuli are known to trigger the release of neuromodulators, which cause an extended change in internal state and thus in behaviour (Anderson and Adolphs, 2014), and the habenula is a part of the circuit mediating stimulus-mediated neuromodulator release. To test this hypothesis, we expressed a dominant negative mutant of the *clock* gene (Dekens and Whitmore, 2008) specifically in the zebrafish habenula. We examined how this affected the alarm response of zebrafish, which we show here to be a circadian behaviour, as has been reported in other fish (Jachner, 1996; Thines and Vandenbussche, 1966).

Expression of ΔCLK protein in the habenula led to a reduced alarm response, such that the day-night difference was no longer present. Disruption of the clock inhibited freezing and dark avoidance induced by the alarm substance. The darting and immediate downward movement caused by alarm substance were not affected, consistent with the hypothesis that the habenula primarily affects the change of internal state that follows exposure to the alarm substance. No change in circadian behaviours that are unlinked to stressors, such as locomotion or arousal, was observed in fish with ΔCLK expression in the habenula. Together, these observations indicate that the behavioural consequence of disruption of the habenula clock is restricted primarily to stress responses, in contrast to the broader phenotype seen with pineal-specific disruption of the clock (Ben-Moshe Livne et al., 2016).

One possible mechanism for the reduced stress response in fish expressing ΔCLK is that the net change in stimulus-evoked activity is lower. As shown in this paper, activity in the zebrafish lateral habenula is higher at night. Expression of ΔCLK blocks transcription of *per* (Dekens and Whitmore, 2008), which is predicted to place habenula neurons in the 'night state'

throughout the circadian cycle. Consistent with this, the day-time levels of dopamine and serotonin in ΔCLK fish were similar to night-time levels. A higher level of activity means that an incoming stimulus will cause a relatively small change in output. In contrast, the same stimulus will have a stronger effect in a fish with low spontaneous activity in the habenula. This interpretation is in line with the finding of Chastrette and colleagues (Chastrette et al., 1991) that the effect of a stressor on the habenula of rats is higher during the active period, when spontaneous activity is low. The observation that the alarm response is reduced at night is consistent with this.

Mechanisms that control spontaneous activity in the lateral habenula are one area of growing interest in mental health, given that increased activity here is linked to abnormal stress responses (Andalman et al., 2019; Li et al., 2013; Yang et al., 2018). The findings here suggest that a potential way forward is to manipulate the clock specifically in the habenula, given the relatively specific effects seen. It may also be useful to screen for clock-regulated genes. This could be done by single-cell RNAseq of habenula neurons at different times of the day, for example, which may lead to the identification of ion channels or other molecules that regulate intrinsic activity and excitability of habenula neurons. This knowledge may contribute to an improvement in the treatment of stress disorders.

# Materials and Methods

## Transgenic lines

TgBAC(gng8:GAL4)[c426] (Hong et al., 2013) was used to drive expression in the habenula. Tg(UAS:EGFP-ΔCLK) was generated by injection of pT2-UAS:EGFP-2A-ΔCLK-5xMYC using Tol2 mediated transgenesis. All experiments were carried out under guidelines approved by the IACUC of Biopolis (181408) and NTU (A19014).

## pERK analysis

### *Immunohistochemistry*

Fish of *nacre* background were randomly split into 'Day' and 'Night' groups. Fish were raised at a density of ~0.125 fish/mL and maintained at 28°C under 14-10 h LD cycle, henceforth referred to as standard conditions. For the comparison under DD, tanks with the larvae were transferred into a dark box at ZT14 the day before fixation.

Larvae at the age of 8 dpf were fixed at Zeitgeber time ZT3 and ZT15 or circadian time CT3 and CT15, based on findings that the peak and trough of *per3* expression occurs at ZT3 and ZT15, respectively (Moore and Whitmore, 2014). Immediate and consistent fixation was achieved by pouring the water containing the larvae through a fine sieve that was immediately placed in ice-cold 4% paraformaldehyde (PFA) in phosphate buffered saline (PBS). Fixations at ZT15, CT3 and CT15 were performed under dim light. Incubation in the fixative was carried out at 4ºC for ~22h. Fixed larvae were then stored in 100% methanol at -20℃ for several days. Larvae were rehydrated by successive 5-minute washes in 70%, 50%, 25% and 0% methanol in PBS, then placed in PBST [1 % Bovine serum albumin (Sigma) + 1 % dimethyl sulfoxide (DMSO) + 0.1% Triton-X 100 in PBS] for 1 h at room temperature. They were then incubated in primary antibodies ((rabbit anti-pERK (Cell Signalling, #4370) and mouse anti-tERK (Cell Signalling, #4696), both diluted at 1:500) at 4°C for 3-4 days. Samples were washed five times in PBST and then incubated in secondary antibodies (Alexa Fluor 647 goat anti-rabbit (ThermoFisher) and Alexa Fluor 488 goat anti-mouse (ThermoFisher), both diluted at 1:500) for 2 h at room temperature. Samples in which the primary antibody was omitted were run in parallel to test the specificity of the secondary antibody binding. Sample size was based on previous experiments (Randlett et al., 2015).

### *Confocal imaging*

Larvae were mounted in 2% low melting agarose (LMA) in PBS and imaged using a Zeiss LSM800 upright confocal microscope equipped with 40X 1.0 NA water dipping objective. The acquisition settings were kept consistent in all experiments. Images were collected at 512x512 resolution. The habenula was imaged with parts of the forebrain (pallium and subpallium) to improve image registration. The number of frames was set to cover the whole depth of habenula and varied across samples, which was later corrected during the registration.

### *Image registration and data analysis*

In order to efficiently compare pERK signals between 'Day' and 'Night' groups, all confocal stacks were first aligned so that each voxel covered an approximately equivalent physical location in each sample. This was achieved by non-rigid registration of the images to a single reference image using the homogenous anti-tERK staining. The registration was carried out with the Computational Morphometry Toolkit (CMTK, http://www.nitrc.org/projects/cmtk/) and the command string (-awr 010203 -T 4 -X 52 -C 8 -G 80 -R 3 -A '--accuracy 0.4' -W '--accuracy 1.6'). Registered stacks were scaled down to 150/150/30 (x/y/z) resolution and the

image noise was reduced with 2D Gaussian filter (sigma = 2) using the
'PrepareStacksForMAPMapping.ijm'              macro              in              FIJI              software
(http://engertlab.fas.harvard.edu/Z-Brain/ (Randlett et al., 2015)). Image data was then
normalized by calculating pERK/tERK ratio and a pairwise comparison between 'Day' and
'Night' groups was carried out at each pixel as follows: first pseudo groups were randomly
generated containing equal number of fish from both groups and Mann-Whitney U statistics
Z-score was calculated over 500 iterations, producing a distribution of Z-scores. Assuming the
normal distribution, a Z-score value corresponding to the false discovery rate of 0.1% was then
used as a threshold so that if the Z-score value of the 'Day' vs 'Night' comparison was higher
than the threshold it was considered significant. Application of the false discovery rate of 0.1%
as a threshold was justified by a control comparison of randomly created mixed groups. The
pixels which were significantly different during the final 'Night'-over-'Day' comparison,
during which 'Day' group served as a baseline, were then assigned a colour where green and
magenta represented significant increase and decrease, respectively, in the 'Night' group.
Those pixels were also allocated intensity values within the range of 0:65535 according to the
size of delta between the median values of the two groups (i.e. the lowest and highest intensities
indicated the smallest and largest deltas within the range of 0:0.5, correspondingly). The pixels
which did not exhibit significant difference were not assigned any colour. All data were
analysed using the 'MakeTheMAPMap.m' script (http://engertlab.fas.harvard.edu/Z-Brain/,
(Randlett et al., 2015)) in Matlab (Mathworks).

## Immunohistochemistry and imaging of EGFP-ΔCLK

The *Tg(gng8:GAL4,UAS:EGFP-ΔCLK)* larvae at the age of 7 dpf were fixed in 4% PFA at 4°C
for ~22 h. For immunostaining of extracted brains, skin covering the brain was manually
removed and exposed brains were washed 3 times in PBST. Antigen retrieval was performed
by incubating the fish in 150 mM Tris-HCl (pH 9.0) for 5 min at room temperature in order to
equilibrate, and by subsequent heating of the samples at 70°C for 15 min. Following several
washes and a blocking step, samples were incubated in the rabbit anti-GFP primary antibody
(Torrey Pines Biolabs Inc., #TP401), diluted 1:500, and the mouse anti-Myc antibody (9E10;
Santa Cruz Biotechnology), diluted 1:100, at 4°C for a few days. The secondary antibodies
used were Alexa Fluor 488 goat anti-rabbit (ThermoFisher and Abcam), diluted 1:1000, and
Alexa Fluor 647 goat anti-mouse (ThermoFisher), diluted 1:500. The immunostained brains

were mounted in 2% LMA in PBS and imaged using a Zeiss LSM800 confocal microscope
equipped with 40X 0.7 NA water immersion objective.

## Screening and selection of *Tg(gng8:GAL4,UAS:EGFP-ΔCLK)* fish

All of the *Tg(gng8:GAL4,UAS:EGFP-ΔCLK)* fish used in the following experiments were first
screened for GFP fluorescence at 3 dpf using a Olympus MVX10 stereo fluorescence
microscope. Since the expression of EGFP-ΔCLK in the habenula was mosaic, only individuals
with expression in the majority of habenula cells were selected and used for further experiments
as ΔCLK-positive (ΔCLK+) fish. Individuals with zero expression as well as larvae exhibiting
a minimal number of randomly labelled cells (less than ~15 cells per habenula) were used as
the control (CTRL) siblings (Figure S1). Following the behavioural experiments, larvae were
immobilized in tricaine methanesulfonate (MS-222; final concentration of 0.01 mg/ml; Sigma),
mounted in 2% LMA in E3 medium and the habenula-specific expression of EGFP was
assessed using a Zeiss LSM800 confocal microscope and 40X water dipping objective.

## Quantification of DA and 5-HT levels

Three-month old female ΔCLK+ fish and their CTRL siblings were immobilized using iced
water and euthanized by decapitation at ZT3 and ZT15. Whole brains, excluding the olfactory
bulb, were extracted in the ice-cold Ringer's solution (116 mM NaCl, 2.9 mM KCl, 1.8 mM
$CaCl_2$, 5.0 mM HEPES, pH 7.2), snap-frozen in liquid nitrogen and stored in -80˚C until further
analysis. Supernatant was collected as described in (Chatterjee and Gerlai, 2009).
Quantification of dopamine (DA) and serotonin (5-HT) was carried out by LC/MS using the
following analytical standards: dopamine hydrochloride (D2960000: Sigma-Aldrich),
dopamine-1,1,2,2-$d_4$hydrochloride (73483: Sigma-Aldrich) and serotonin (14927: Sigma-
Aldrich). Data were analysed and plotted using Estimation Statistics (Ho et al., 2019). Three
biological replicates (i.e. measurements of different samples) were performed.

## Behaviour assays

*Acoustic startle following exposure to Schreckstoff*
Thirteen to sixteen day-old ΔCLK larvae and their non-expressing siblings were placed in two
transparent plastic tanks with dimensions of 80 x 54 x 33 (L x W x H in mm) and their
locomotor activity was recorded from above at 10 fps using a camera (acA2040-90μm USB
3.0; Basler). Each of the two tanks was filled with 20 mL of facility water and held 8 fish at

most. The water level in each tank was 5 mm in height to minimize vertical movement. The

alarm substance was freshly prepared from adult zebrafish (Mathuru et al., 2012) each week,

and was tested for effectiveness on adults before use on larvae. An Arduino Grove Vibration

Motor was positioned in between the two plastic tanks, on the stage holding the tanks, to

provide the stimulus. The vibration was 1 sec in duration and was presented 3 times, at 1 minute

intervals, starting five minutes before and after the delivery of the *Schreckstoff*. A startle

response was defined as a change in body orientation of more than 90 degrees within 5 sec in

response to each vibration tone. The response (yes or no for each fish) was averaged across 3

tones for each group to obtain a response percentage before and after the delivery of the

*Schreckstoff*. Both the vibration tones and the delivery of *Schreckstoff* were controlled by an

Arduino board synched with the video-recording Python codes. Experiments were performed

on 3 separate batches of embryos. A different batch of freshly prepared skin extract was used

for each experiment; the extract was verified for effectiveness by testing on adult fish. Only

extract that evoked freezing was used.

*Locomotor activity*

The *Tg(gng8:GAL4,UAS:EGFP-ΔCLK)* larvae were raised under standard conditions. Each

experimental run involved animals from the same batch, originating from one pair of parents.

The larvae were screened and selected as described in the previous section. At the age of 6 dpf

the larvae were placed individually in a 24-well plate filled with 1.5 ml of tank water per well.

The plate was then transferred into an incubator equipped with an IR camera (acA2040-90µm

USB 3.0: Basler), a white LED box and four IR LED bars (LBS2-00-080-3-IR850-24V,

850 nm: TMS Lite). The camera and the LEDs were connected to a PC via a microcontroller

board (Arduino UNO SMD). At the onset of recording (ZT6 – 9), fish were kept under ambient

light, which was switched off at ZT14 (10 PM) and was kept off during the next day for the

monitoring of activity under DD. A customized script written in Python 2.7, incorporating

functions from the OpenCV library was used to control the Arduino microcontroller and

simultaneously video-track fish locomotor activity. The video was captured using 576 x 864-

pixel resolution at 10 frames per second and the background subtraction method was

implemented to extract the 'x' and 'y' coordinates of each fish online. The position data was

then converted to total distance (mm) travelled by each fish per 1 min time. The data

conversion, further analysis and plotting were carried out using custom-written Excel macros,

Python 2.7 and Estimation Statistics (Ho et al., 2019).

 *Arousal threshold*

Six days-old ΔCLK+ larvae and their CTRL siblings were randomly placed in a 24-well plate

and their locomotor activity was video-tracked as described in the previous section. In order to

assess the arousal during sleep, vibration stimuli were applied through a speaker controlled,

via an Arduino microcontroller board, by a modified version of the activity-tracking code in

Python 2.7. Starting at ZT17, eighteen stimuli of nine different intensities, where 50% of

computer audio output represented the highest intensity, were delivered for 200 ms (starting

from 0%) in 12-16% increments every minute, first in a descending and then ascending order.

The time interval between stimuli was determined based on previous findings that suggest a

30-second inter-stimulus gap is sufficient to avoid behavioural habituation (Burgess and

Granato, 2007). The protocol was repeated at the commencement of every hour until ZT22,

while the onset by ascending or descending order was altered each time. In total, 12 replicates

(i.e. technical replicates) of each stimulus intensity were generated. A fish was scored as

responding if it moved more than 7 pixels (~ 4 mm) following the stimulus delivery and the

average percentage of responding fish was calculated for each of the stimulus intensities

(Woods et al., 2014). The observed values were corrected for variations in the baseline

locomotor activity, by first calculating the background probability of locomotion as a

percentage of larvae which moved more than 7 pixels 5 s before the application of each

stimulus. The average of 108 measures of the background probability of locomotion was then

used to calculate the corrected probability of response at each of the stimulus intensities,

implementing the following equation: corrected probability of response = observed value x

[(observed value - background offset) / (max observed value - background offset)] (Woods et

al., 2014).

*Novel tank assay*

All experiments were conducted between ZT2 and ZT4 on 3-5 month-old adult zebrafish,

following the protocol of (Haghani et al., 2019)). Fish were recorded in pairs in a darkened

room with a black background, where tanks were illuminated from the top with a natural light

LED bar. Novel tanks were made of glass measuring 20 cm (L) by 5 cm (W) by 12 cm (H)

with opaque sides (to prevent fish from viewing each other), placed side by side. Fresh water

from the facility was added to tanks up to the 10 mL mark to make a total volume of 1 L. The

setup was placed within a curtained area to obscure the experimenter.

Prior to exposure to the novel tank, fish were handled carefully to ensure that handling stress
was minimised. Fish were gently netted in pairs from their home tanks in the facility to smaller
crossing tanks, where they were brought into the room with the behavioural setup. Then, fish
were netted into 100 mL beakers with ~30 mL of fish facility water and quickly poured into
the novel tank. The recording was started within 30 s of fish being in the recording tanks. For
exposure to *Schreckstoff,* fish were placed in 100 mL beakers containing 200 μL of heated skin
extract (see (Chia et al., 2019) for extraction method) in 50 mL fish facility water for 2 minutes.
Fish were netted into another beaker containing fresh facility water before being poured into
the novel tank. Videos were acquired at 10 frames per second for 600 s (10 mins) using a Basler
Ace camera (acA1300–200 μm; 1,280 × 1,024). The position (x-y coordinates) of the fish were
tracked, and speed was binned by mm/s into a probability density curve.

For acute exposure to *Schreckstoff*, fish were first habituated in the glass tanks for 5-7 mins.
Then, fish were recorded in the tanks for 5 mins before 1 mL of 1:4 dilution of heated skin
extract was pipetted into the tanks. The recording continued for 5 mins after *Schreckstoff*
delivery for the alarm response to be observed. Two fish were tested at one time, each in a
separate tank, with one tank containing ΔCLK fish and the other a non-expressing sibling. The
tank used for a particular genotype was switched at every trial. Characteristic alarm behaviours
were quantified, such as time spent in the bottom third of the tank, number of darting episodes
(fish speed >5 SD above average swim speed for the first 5 mins), and number of freezing
episodes (movement of <5.5 mm per second). Tracking was automated to prevent observer
bias. Sample size was based on previous experiments (Chia et al., 2019).

*Light/dark choice assay*
All experiments were performed between ZT2 and ZT9 in a behavioural arena within blackout
curtains as described previously (Cheng et al., 2016). Four transparent plastic tanks
(dimensions: 43 mm W × 60 mm L × 30 mm H) filled with 30 mL fresh tank water served as
assay chambers. Opaque cardboard sheets were placed between the 4 tanks. Tanks were placed
on an Apple iPad screen and videos (9 fps) were recorded on a USB3.0 Basler camera (Model#
acA2040-90umNIR 1440 × 1080 pixels) with a long-pass filter (MIDOPT, LP830 830nm)
placed above. Two IR light bars (850 nm peak from TMS-lite) positioned next to the 4 tanks
served as an IR light source. The light/dark compartments were created by white and black
rectangles (with 50% transparency level for the black) on a Microsoft PowerPoint slide

displayed on the iPad Air at highest brightness level. Sample size was based on previous experiments (Cheng et al., 2016).

Larvae were exposed either to *Schreckstoff* or to control water (embryo medium) in a treatment tank of the same dimensions with the same volume of tank water for 10 minutes. Larvae were then gently pipetted from the treatment tank into two wash-out tanks sequentially before being placed into the dark/light test chamber. Real-time tracking of the fish was achieved by custom-written Python codes using the OpenCV library and recorded as coordinates of the centroid of the fish across time. Larval behaviour was recorded for 10 minutes. Offline analysis was performed using custom-written Excel macros to determine the position of the animals, the number of entries into each compartment, the percentage of total time spent in each compartment, etc. A few fish (n=2 in both the Control group and Gng8/ΔCLK group) were excluded from the analysis because they were entirely immobile or because of tracking errors.

# Acknowledgements

We thank Yoav Gothilf for providing the pT2-UAS:EGFP-2A-ΔCLK-5xMYC construct and Marnie Halpern for providing the *TgBAC(gng8:GAL4)$^{c426}$* line. We also thank Hugh Piggins, David Lyons, Caroline Wee and Ajay Mathuru for comments and discussion. This work was funded by grants from the Singapore Ministry of Education under its Academic Research Fund Tier 2 (MOE2017-T2-058) and the National Research Foundation (NRF2017-NRF-ISF002-2676) to SJ, and by an ARAP fellowship from A*Star to AB.

**Author contributions**

AB, RKC and JCSM performed experiments and analysed data. SJ and AB designed the study and wrote the manuscript.

449 Figures

**Figure 1. Day-night difference in ERK phosphorylation in the habenula.** (**A, B**)
Immunohistochemical detection of phosphorylated ERK (pERK; magenta) and total ERK
(tERK, green) at two different focal planes of the habenula, dorsal (A) and ventral (B), of an
8-day old zebrafish. The white arrows indicate labelled cell bodies, while the yellow arrow
indicates the neuropil. (**C-H**) Day-Night comparison of normalized pERK levels in the
habenula. Raw images were registered, downsampled and smoothed and pERK/tERK ratio was
calculated at each pixel. Median pERK/tERK levels in the habenula were calculated based on
N = 102 larvae fixed during the night (C, F) and N = 130 larvae fixed during the day (D, G).

Two different planes, ventral (C, D) and dorsal (F, G) are shown here. The difference in the ventral (E) and dorsal (H) planes, based on 'Night-over-Day' comparison, with green indicating higher levels of pERK at night, while magenta indicates lower levels (see wedge in panel E). (**I-K**) Reconstructed transverse section through the habenula of the stacks of median activity during the night (I), day (J) and difference (K). Strong activity is detected in the lateral region only. All images except I-K are dorsal view. Scale bar = 10 µm. a: anterior; p: posterior; d: dorsal; v: ventral; Pa: pallium; Hb: habenula.

**Figure 2. Circadian variation in habenula activity under constant darkness.** (**A-F**) Median pERK/tERK label at three different focal planes, in larvae fixed in the day (**A-C**; N = 104) or at night (**D-F**; N = 102). A and D are dorsal, B and E are intermediate, whereas C and F are deep. (**G-I**). The difference in pERK levels at Night versus Day, at the 3 different focal planes.

The arrows in I indicate increased pERK in the lateral habenula. Bar = 10 μm. Contrast settings

on panels A-F are identical.

**Figure 3. The *gng8:*GAL*4* driver enables expression of a C-terminus truncated *clock* gene**

**in the habenula of zebrafish.** (**A**) Specific expression of EGFP in the habenula of a

Tg(*gng8:GAL4, UAS:EGFP-ΔCLK*) fish. (**B**) At higher magnification, eGFP is visible in the

cytoplasm of habenula neurons. (**C**) Detection of Myc, which is fused to the C-terminus of

ΔCLK, in the habenula of transgenic fish. Scale bar = 100 μm in panel A, 20 μm in panels B

and C.

**Figure 4. Effects of ΔCLK expression in the habenula on global neuromodulator levels.**

Cumming estimation plots showing the difference in levels of dopamine (A) and serotonin (B) at daytime and night-time in ΔCLK and sibling fish. The raw data is plotted on the upper axis, with each dot representing a value obtained from a single brain. Gapped vertical lines to the right of each group indicate mean ± SD. Mean differences are depicted on the lower axis as black dots, with the vertical bars indicating bootstrap 95% CIs.

**Figure 5. Day-night variation in the response of zebrafish to *Schreckstoff*.** Percentage of larval fish that show a startle response before and after exposure to the alarm substance, during the day (11 am) or at night (11 pm). Control fish (N = 13), which are non-expressing siblings of *Tg(gng8:GAL4, UAS:ΔCLK)* fish (N = 21), show a reduction in startle after exposure during the day, but not at night. Each circle in the top plot represents the percentage of responding fish in one batch. The bottom half shows a Cummings estimation plot of the difference between the startle after and before exposure to the alarm substance. Mean differences are depicted as dots; 95% confidence intervals are indicated by the ends of the vertical error bars.

**Figure 6. The effects of *Schreckstoff* on *Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* fish. (A-E)**

Acute effects in adult zebrafish. (**A, B**) Speed distribution in fish before and during exposure

to the alarm substance Freezing, as indicated by the peak at 0 mm/s, is seen in both wildtype (A) and ΔCLK fish (B). (**C-E**) Time spent darting (C) or at the bottom third of the tank (D), or freezing (E) before and in the presence of *Schreckstoff*. There is no difference in the behaviour of fish with and without expression of truncated clocka. (**F, G**). Post-exposure behaviour in adult fish (N = 8). (**F**) Histogram of speed in a novel tank containing clean water, after exposure to Schreckstoff in a separate container. Fish expressing ΔCLK in the habenula do not freeze (green shading), in contrast to their non-expressing siblings (gray shading). (**G**). Behaviour of non-exposed fish in the novel tank. No freezing was observed. (**H, I**) Post-exposure behaviour in 2 week-old zebrafish. (**H**) Percentage of time in the dark area of a tank, after 10 minutes of exposure to embryo water. (**I**) Percentage of time spent in the dark side after exposure to the alarm substance. *Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* fish spend more time in the dark side, with a Cohen's d of 1.2.

**Figure 7. Circadian behaviour is unaltered in *Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* larvae**

(**A, B**) Distances moved (cm/10min) by individual larvae from (A) CTRL group and (B)

ΔCLK+ group are plotted against time (hr) over the time-course of the whole recording. CT =

circadian time. (**C, D**) Mean distances moved (cm/10min) are plotted against time (hr) (C)

during the first 14 hours of recording, including 4 hours of light-phase and 10 hours in dark

during the night and (D) during the subsequent 24 hours of recording under DD. CTRL (green)

groups Tg(gng8:GAL4;UAS:EGFP-ΔCLK) 95% CIs are indicated by the shaded areas;

interrupted lines indicate 5 analyzed timeframes. Black and white bars represent periods of

dark and light, correspondingly. (**E**) The Cumming estimation plot shows the mean differences

for 5 comparisons, corresponding to 5 timeframes of recording. These are 1 = light-phase; 2 =

1st night; 3 = early subjective day; 4 = late subjective day; 5 = subjective night. The raw data

is plotted on the upper axis, each dot representing the overall distance moved (cm) by a single

larva; gapped vertical lines to the right of each group indicate mean ± SD. Each mean difference

is plotted on the lower axes as a bootstrap sampling distribution. Mean differences are depicted

as black dots and 95% CIs are indicated by the end of vertical error bars. (**F**) Analysis of

sensory responsiveness of the *Tg(gng8:gal4,UAS:EGFP-ΔCLK)* larvae (7 dpf) during the

delivery of auditory stimuli of 9 different intensities. The Cumming estimation plot shows the

mean differences for 9 comparisons, corresponding to 9 stimulus intensities (0-100%). The raw

data is plotted on the upper axis, each dot representing the percentage of responders per single

experimental run. (G)

Figure S1. Examples of *Tg(gng8:GAL4, UAS:EGFP-ΔCLK)* larval zebrafish with broad (A, B)

and sparse (C, D) expression of EGFP in the habenula. All images are dorsal view. Scale bar

= 10 μm.

## References

Abercrombie ED, Keefe KA, DiFrischia DS, Zigmond MJ. 1989. Differential effect of stress

on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex.

*J Neurochem* **52**:1655–1658.

Andalman AS, Burns VM, Lovett-Barron M, Broxton M, Poole B, Yang SJ, Grosenick L,

Lerner TN, Chen R, Benster T, Mourrain P, Levoy M, Rajan K, Deisseroth K. 2019.

Neuronal Dynamics Regulating Brain and Behavioral State Transitions. *Cell* **177**:970–

985.e20.

Anderson DJ, Adolphs R. 2014. A framework for studying emotions across species. *Cell*

**157**:187–200.

Antle MC, Silver R. 2016. Circadian Insights into Motivated Behavior. *Curr Top Behav*

*Neurosci* **27**:137–169.

Belle MDC. 2015. Circadian Tick-Talking Across the Neuroendocrine System and

Suprachiasmatic Nuclei Circuits: The Enigmatic Communication Between the

Molecular and Electrical Membrane Clocks. *J Neuroendocrinol* **27**:567–576.

Ben-Moshe Livne Z, Alon S, Vallone D, Bayleyen Y, Tovin A, Shainer I, Nisembaum LG,

Aviram I, Smadja-Storz S, Fuentes M, Falcón J, Eisenberg E, Klein DC, Burgess HA,

Foulkes NS, Gothilf Y. 2016. Genetically Blocking the Zebrafish Pineal Clock Affects

Circadian Behavior. *PLoS Genet* **12**:e1006445.

Buijs RM. 1978. Intra- and extrahypothalamic vasopressin and oxytocin pathways in the rat.

Pathways to the limbic system, medulla oblongata and spinal cord. *Cell Tissue Res*

**192**:423–435.

Burgess HA, Granato M. 2007. Sensorimotor gating in larval zebrafish. *J Neurosci* **27**:4984–

4994.

Cao G, Nitabach MN. 2008. Circadian control of membrane excitability in Drosophila

melanogaster lateral ventral clock neurons. *J Neurosci* **28**:6493–6501.

Chaouloff F, Berton O, Mormède P. 1999. Serotonin and stress. *Neuropsychopharmacology*

**21**:28S–32S.

Chastrette N, Pfaff DW, Gibbs RB. 1991. Effects of daytime and nighttime stress on Fos-like

immunoreactivity in the paraventricular nucleus of the hypothalamus, the habenula, and

the posterior paraventricular nucleus of the thalamus. *Brain Res* **563**:339–344.

Chatterjee D, Gerlai R. 2009. High precision liquid chromatography analysis of

dopaminergic and serotoninergic responses to acute alcohol exposure in zebrafish.

*Behav Brain Res* **200**:208–213.

Cheng R-K, Krishnan S, Jesuthasan S. 2016. Activation and inhibition of tph2 serotonergic

neurons operate in tandem to influence larval zebrafish preference for light over

darkness. *Sci Rep* **6**:20788.

Chia JSM, Wall ES, Wee CL, Rowland TAJ, Cheng R-K, Cheow K, Guillemin K, Jesuthasan

S. 2019. Bacteria evoke alarm behaviour in zebrafish. *Nat Commun* **10**:3831.

Daut RA, Fonken LK. 2019. Circadian regulation of depression: A role for serotonin. *Front*

*Neuroendocrinol* **54**:100746.

Dekens MPS, Whitmore D. 2008. Autonomous onset of the circadian clock in the zebrafish

embryo. *The EMBO Journal*. doi:10.1038/emboj.2008.183

Guilding C, Hughes ATL, Piggins HD. 2010. Circadian oscillators in the epithalamus.

*Neuroscience* **169**:1630–1639.

Haghani S, Karia M, Cheng R-K, Mathuru AS. 2019. An Automated Assay System to Study

Novel Tank Induced Anxiety. *Front Behav Neurosci* **13**:180.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. 2019. Moving beyond P values: data

analysis with estimation graphics. *Nat Methods* **16**:565–566.

Holly EN, Miczek KA. 2016. Ventral tegmental area dopamine revisited: effects of acute and

repeated stress. *Psychopharmacology* **233**:163–186.

Hong E, Santhakumar K, Akitake CA, Ahn SJ, Thisse C, Thisse B, Wyart C, Mangin J-M,

Halpern ME. 2013. Cholinergic left-right asymmetry in the habenulo-interpeduncular

pathway. *Proc Natl Acad Sci U S A* **110**:21171–21176.

Hughes AT, Fahey B, Cutler DJ, Coogan AN, Piggins HD. 2004. Aberrant gating of photic

input to the suprachiasmatic circadian pacemaker of mice lacking the VPAC2 receptor. *J*

*Neurosci* **24**:3522–3526.

Jachner A. 1996. Alarm reaction in bleak (Alburnus alburnus (L.), Cyprinidae) in response to

chemical stimuli from injured conspecifics. *Hydrobiologia*. doi:10.1007/bf00028275

Jesuthasan SJ, Krishnan S, Cheng R-K, Mathuru A. 2020. Larval zebrafish respond to the

alarm pheromone Schreckstoff by immobility and a change in brain state. *bioRxiv*.

doi:10.1101/2020.04.23.054734

Jesuthasan SJ, Mathuru AS. 2008. The alarm response in zebrafish: innate fear in a vertebrate

genetic model. *J Neurogenet* **22**:211–228.

Koch CE, Leinweber B, Drengberg BC, Blaum C, Oster H. 2017. Interaction between
circadian rhythms and stress. *Neurobiol Stress* **6**:57–67.

Li K, Zhou T, Liao L, Yang Z, Wong C, Henn F, Malinow R, Yates JR 3rd, Hu H. 2013.
βCaMKII in lateral habenula mediates core symptoms of depression. *Science* **341**:1016–
1020.

Mathuru AS, Kibat C, Cheong WF, Shui G, Wenk MR, Friedrich RW, Jesuthasan S. 2012.
Chondroitin fragments are odorants that trigger fear behavior in fish. *Curr Biol* **22**:538–
544.

Maximino C, Lima MG, Costa CC, Guedes IML, Herculano AM. 2014. Fluoxetine and
WAY 100,635 dissociate increases in scototaxis and analgesia induced by conspecific
alarm substance in zebrafish (Danio rerio Hamilton 1822). *Pharmacol Biochem Behav*
**124**:425–433.

McClung CA, Sidiropoulou K, Vitaterna M, Takahashi JS, White FJ, Cooper DC, Nestler EJ.
2005. Regulation of dopaminergic transmission and cocaine reward by the Clock gene.
*Proc Natl Acad Sci U S A* **102**:9377–9381.

Moore HA, Whitmore D. 2014. Circadian rhythmicity and light sensitivity of the zebrafish
brain. *PLoS One* **9**:e86176.

Namboodiri VMK, Rodriguez-Romaguera J, Stuber GD. 2016. The habenula. *Curr Biol*
**26**:R873–R877.

Pennartz CMA, de Jeu MTG, Bos NPA, Schaap J, Geurtsen AMS. 2002. Diurnal modulation
of pacemaker potentials and calcium current in the mammalian circadian clock. *Nature*
**416**:286–290.

Pfeiffer W. 1962. The fright reaction of fish. *Biol Rev Camb Philos Soc* **37**:495–511.

Pfeiffer W, Riegelbauer G. 1978. The effect of the alarm substance on the central nervous
excitation of the black tetraGymnocorymbus ternetzi (Characidae, Ostariophysi, Pisces)
indicated by dorsal light response. *Journal of Comparative Physiology ? A.*
doi:10.1007/bf00656882

Prober DA, Rihel J, Onah AA, Sung R-J, Schier AF. 2006. Hypocretin/orexin overexpression
induces an insomnia-like phenotype in zebrafish. *J Neurosci* **26**:13400–13410.

Randlett O, Wee CL, Naumann EA, Nnaemeka O, Schoppik D, Fitzgerald JE, Portugues R,
Lacoste AMB, Riegler C, Engert F, Schier AF. 2015. Whole-brain activity mapping
onto a zebrafish brain atlas. *Nature Methods*. doi:10.1038/nmeth.3581

Sakhi K, Wegner S, Belle MDC, Howarth M, Delagrange P, Brown TM, Piggins HD. 2014.
Intrinsic and extrinsic cues regulate the daily profile of mouse lateral habenula neuronal

activity. *J Physiol* **592**:5025–5045.

Salaberry NL, Hamm H, Felder-Schmittbuhl M-P, Mendoza J. 2019. A suprachiasmatic-
    independent circadian clock(s) in the habenula is affected by Per gene mutations and
    housing light conditions in mice. *Brain Struct Funct* **224**:19–31.

Sánchez-Bretaño A, Gueguen M-M, Cano-Nicolau J, Kah O, Alonso-Gómez ÁL, Delgado
    MJ, Isorna E. 2015. Anatomical distribution and daily profile of gper1b gene expression
    in brain and peripheral structures of goldfish (Carassius auratus). *Chronobiol Int*
    **32**:889–902.

Scammell TE, Arrigoni E, Lipton JO. 2017. Neural Circuitry of Wakefulness and Sleep.
    *Neuron* **93**:747–765.

Shearman LP, Zylka MJ, Weaver DR, Kolakowski LF Jr, Reppert SM. 1997. Two period
    homologs: circadian expression and photic regulation in the suprachiasmatic nuclei.
    *Neuron* **19**:1261–1269.

Shieh K-R. 2003. Distribution of the rhythm-related genes rPERIOD1, rPERIOD2, and
    rCLOCK, in the rat brain. *Neuroscience* **118**:831–843.

Speedie N, Gerlai R. 2008. Alarm substance induced behavioral responses in zebrafish
    (Danio rerio). *Behav Brain Res* **188**:168–177.

Steenbergen PJ, Richardson MK, Champagne DL. 2011. Patterns of avoidance behaviours in
    the light/dark preference test in young juvenile zebrafish: a pharmacological study.
    *Behav Brain Res* **222**:15–25.

Sun ZS, Albrecht U, Zhuchenko O, Bailey J, Eichele G, Lee CC. 1997. RIGUI, a putative
    mammalian ortholog of the Drosophila period gene. *Cell* **90**:1003–1011.

Takahashi JS. 2017. Transcriptional architecture of the mammalian circadian clock. *Nat Rev
    Genet* **18**:164–179.

Thines G, Vandenbussche E. 1966. The effects of alarm substance on the schooling
    behaviour of Rasbora heteromorpha Duncker in day and night conditions. *Anim Behav*
    **14**:296–302.

Weger M, Weger BD, Diotel N, Rastegar S, Hirota T, Kay SA, Strähle U, Dickmeis T. 2013.
    Real-time in vivo monitoring of circadian E-box enhancer activity: a robust and
    sensitive zebrafish reporter line for developmental, chemical and neural biology of the
    circadian clock. *Dev Biol* **380**:259–273.

Woods IG, Schoppik D, Shi VJ, Zimmerman S, Coleman HA, Greenwood J, Soucy ER,
    Schier AF. 2014. Neuropeptidergic signaling partitions arousal behaviors in zebrafish. *J
    Neurosci* **34**:3142–3160.

Yang Y, Cui Y, Sang K, Dong Y, Ni Z, Ma S, Hu H. 2018. Ketamine blocks bursting in the
lateral habenula to rapidly relieve depression. *Nature* **554**:317–322.

Zhao H, Rusak B. 2005. Circadian firing-rate rhythms and light responses of rat habenular
nucleus neurons in vivo and in vitro. *Neuroscience* **132**:519–528.

Zhao Z, Xu H, Liu Y, Mu L, Xiao J, Zhao H. 2015. Diurnal Expression of the Per2 Gene and
Protein in the Lateral Habenular Nucleus. *Int J Mol Sci* **16**:16740–16749.

Zhdanova IV, Wang SY, Leclair OU, Danilova NP. 2001. Melatonin promotes sleep-like
state in zebrafish. *Brain Res* **903**:263–268.