1
2
3
4

Does hippocampal volume explain performance differences on hippocampal-dependent tasks?

7
8
9
10

Ian A. Clark, Anna M. Monk, Victoria Hotchin, Gloria Pizzamiglio, Alice Liefgreen, Martina F. Callaghan, Eleanor A. Maguire

Wellcome Centre for Human Neuroimaging, UCL Queen Square Institute of Neurology, University College London, London, UK

16
17
18

Correspondence

Eleanor Maguire, Wellcome Centre for Human Neuroimaging, UCL Queen Square Institute of Neurology, University College London, 12 Queen Square, London, WC1N 3AR, UK**;** e.maguire@ucl.ac.uk

24
25
26

Running title

Hippocampal volume and task performance

30
31
32

Highlights

Evidence is mixed about whether hippocampal volume affects cognitive task performance

This is particularly the case concerning individual differences in healthy people

We collected structural MRI data from 217 healthy people

They also had widely-varying performance on cognitive tasks linked to the hippocampus

In-depth analyses showed little evidence hippocampal volume affected task performance

44
45

Abstract

Marked disparities exist across healthy individuals in their ability to imagine scenes, recall autobiographical memories, think about the future and navigate in the world. The importance of the hippocampus in supporting these critical cognitive functions has prompted the question of whether differences in hippocampal grey matter volume could be one source of performance variability. Evidence to date has been somewhat mixed. In this study we sought to mitigate issues that commonly affect these types of studies. Data were collected from a large sample of 217 young, healthy adult participants, including whole brain structural MRI data (0.8mm isotropic voxels) and widely-varying performance on scene imagination, autobiographical memory, future thinking and navigation tasks. We found little evidence that hippocampal grey matter volume was related to task performance in this healthy sample. This was the case using different analysis methods (voxel-based morphometry, partial correlations), when whole brain or hippocampal regions of interest were examined, when comparing different sub-groups (divided by gender, task performance, self-reported ability), and when using latent variables derived from across the cognitive tasks. Hippocampal grey matter volume may not, therefore, significantly influence performance on tasks known to require the hippocampus in healthy people. Perhaps only in extreme situations, as in the case of licensed London taxi drivers, are measurable ability-related hippocampus volume changes consistently exhibited.

KEYWORDS

Hippocampal volume; scene construction; autobiographical memory; future thinking; spatial navigation; individual differences

1. INTRODUCTION

People vary substantially in their ability to perform tasks related to critical aspects of cognition that enable the smooth functioning of our everyday lives. These include imagining scenes (the process of forming and visualising scene imagery in the absence of visual input), autobiographical memory (the recall of past events from one's life), future thinking (imagining future experiences) and spatial navigation (the process of ascertaining one's position in the environment, and planning and following a route). For example, some individuals can recollect decades-old autobiographical memories with great clarity compared to others who struggle to recall what they did last weekend (e.g. Palombo, Sheldon, & Levine, 2018). Spatial navigation can be undertaken with ease by some people while others consistently get lost (e.g. Newcombe, 2018).

Neuroimaging studies of healthy individuals performing tasks related to these cognitive functions reliably show engagement of the hippocampus (Addis, Wong, & Schacter, 2007; Barry, Barnes, Clark, & Maguire, 2018; Brown, Ross, Keller, Hasselmo, & Stern, 2010; Buckner & Carroll, 2007; Dalton, Zeidman, McCormick, & Maguire, 2018; Spiers & Maguire, 2006; Svoboda, McKinnon, & Levine, 2006). Moreover, damage to the hippocampus leads to impaired performance on similar tasks (e.g. Andelman, Hoofien, Goldberg, Aizenstein, & Neufeld, 2010; Hassabis, Kumaran, Vann, & Maguire, 2007; Maguire, Nannery, & Spiers, 2006; Rosenbaum et al., 2005; Rosenbaum et al., 2000; Scoville & Milner, 1957; Winocur & Moscovitch, 2011). The importance of the hippocampus in supporting scene imagination, autobiographical memory, future thinking and navigation, has prompted the question of whether differences in hippocampal structure could be one source of variable task performance among healthy people. While there have been numerous studies in the last two decades that have addressed this issue, typically in relation to hippocampal volume, evidence has been somewhat mixed.

Considering first scene imagination, as far as we are aware, no studies involving healthy

people have yet investigated the association between hippocampal volume and the ability to

construct scene imagery. Recent work documented a relationship between hippocampal grey

matter volume and scene imagination performance in patients with the behavioural variant of

frontotemporal dementia, a link that was not apparent in patients with Alzheimer's disease

(Wilson et al., 2020). However, in that study no direct comparison was made between the two

patient groups, or between the patients and healthy controls, precluding interpretations about

the association between hippocampal volume and scene imagination ability.

A larger number of studies have investigated the relationship between hippocampal

volume and memory ability in healthy individuals, but with mixed results. On the one hand,

"recollective" memory ability (the detailed re-experiencing of individual episodes

characterized by retrieval of items in their context) has been positively associated with posterior

hippocampal grey matter volume (Poppenk & Moscovitch, 2011). Moreover, a single

individual identified with "Highly Superior Autobiographical Memory" was reported to have

greater posterior hippocampal volume compared to matched controls (Mazzoni et al., 2019).

On the other hand, group studies of individuals with superior memory capabilities, including

those with Highly Superior Autobiographical Memory or those who take part in the World

Memory Championships, observed no differences in hippocampal grey matter volume

compared to matched controls (LePort et al., 2012; Maguire, Valentine, Wilding, & Kapur,

2003). Similarly, a meta-analysis of 33 studies found no relationship between hippocampal

grey matter volume and performance on laboratory-based memory tasks (Van Petten, 2004).

Whether a link exists between hippocampal volume and more naturalistic autobiographical

memory ability is, therefore, unclear.

There has been limited work investigating the relationship between future thinking

ability and hippocampal volume. A recent study claimed to have observed a positive

relationship between participant ratings of sensory perceptual qualities in future thinking (e.g. vividness, the amount of visual details) and hippocampal grey matter volume (Yang, Chen, Zhang, Xu, & Feng, 2020). However, the peak voxel coordinates (MNI space; 39, -7.5, -10.5) and cluster were located outside of the hippocampus (Figure 1 of Yang et al., 2020). A more precise investigation into the relationship between hippocampal volume and future thinking ability is, therefore, required.

A definitive link has been identified between hippocampal grey matter volume and extreme spatial navigation ability. Licensed London (UK) taxi drivers must memorise the extensive and complex layout of ~25,000 London streets and thousands of landmarks, known colloquially as acquiring "The Knowledge". In several studies they were found to have greater posterior, and decreased anterior, hippocampal grey matter volume compared to healthy controls (Maguire et al., 2000), London bus drivers, who spend an equivalent amount of time driving but on regular routes instead of requiring a complete knowledge of London's layout (Maguire, Woollett, & Spiers, 2006), and medical doctors, who have high levels of expertise but not primarily in the spatial domain (Woollett, Glensman, & Maguire, 2008). Moreover, longitudinal data collected before and after attempts to acquire The Knowledge identified posterior hippocampal grey matter volume enlargement within subjects, but only in those individuals who went on to qualify as London taxi drivers (Woollett & Maguire, 2011). Greater posterior, but less anterior, hippocampal grey matter volume is, therefore, reliably associated with extreme spatial navigation expertise.

In the general population, however, the relationship between hippocampal grey matter volume and navigation ability is less clear. One study involving navigation in a virtual environment found no association between hippocampal volume and navigation performance in healthy people (Maguire, Spiers, et al., 2003), a finding that has since been replicated in a larger sample of 90 individuals (Weisberg, Newcombe, & Chatterjee, 2019). However, a re-

analysis of the Weisberg et al. (2019) data purports to have identified a relationship between right posterior hippocampal volume and navigation performance, but one that is dependent on self-reported navigation ability or being grouped according to navigation performance ability (He & Brown, 2020). These interactions suggested that in low performing participants a negative relationship existed between right posterior hippocampal volume and task performance. However, interactions were only observed for one of five performance measures and no multiple comparison corrections were made to the statistical threshold used to determine significance. Given the reasonably large sample size and the number of tests performed on the data (by both Weisberg et al., 2019 and He & Brown, 2020) the results of this re-analysis require replication before firm conclusions can be drawn.

Posterior hippocampal grey matter volume has been positively associated with an individual's ability to learn new environments in the real world (Schinazi, Nardi, Newcombe, Shipley, & Epstein, 2013), in virtual reality (Brown, Whiteman, Aselcioglu, & Stern, 2014), and with first person perspective taking ability during navigation (Sherrill, Chrastil, Aselcioglu, Hasselmo, & Stern, 2018). In addition, greater posterior compared to anterior hippocampal grey matter volume has been linked to the use of "map-based" strategies, and consequently with better navigation performance (Brunec et al., 2019). By contrast, greater anterior hippocampal grey matter volume has been related to better topographical memory ability (Hartley & Harlow, 2012), path integration (Chrastil, Sherrill, Aselcioglu, Hasselmo, & Stern, 2017), and performance on a virtual reality eight arm radial maze (Bohbot, Lerch, Thorndycraft, Iaria, & Zijdenbos, 2007). Therefore, the literature on associations between navigational ability and hippocampal grey matter volume is disparate, and lacks consistency.

Overall, across scene imagination, autobiographical memory, future thinking and navigation, the relationship between hippocampal grey matter volume and task performance in healthy individuals is ambiguous. Why might this be the case?

First, the sample sizes used have typically been small, with most studies including 20-

30 participants. Recent empirical work has highlighted the non-replicability of brain structure-

function associations when using small samples (Kharabian Masouleh, Eickhoff, Hoffstaedter,

Genon, & Alzheimer's Disease Neuroimaging Initiative, 2019). Notably, in the navigation

study where a larger sample of 90 participants was utilised, no relationship between

hippocampal volume and task performance was identified (Weisberg et al., 2019). The mixed

results may, therefore, simply be due to using insufficiently powered samples.

Second, there is a large variability across studies in how analyses were performed. For

example, while several analyses included covariates to account for age, gender and total

intracranial volume, others included covariates for only some of these variables, and numerous

analyses involved no covariates at all. Given that these variables are known to be potential

confounds (e.g. Ridgway et al., 2008), their inclusion (or not) as covariates could significantly

influence the observed results.

Third, studies differ in terms of the level of statistical correction that was applied. For

instance, some studies use more lenient statistical thresholds compared to others, and while

correction for multiple comparisons is evident when testing for multiple relationships in several

studies, in others it is not. Differences in statistical thresholds could not only contribute to

contradictory results across the literature, but may also have increased the number of false

positive findings.

Fourth, there are differences in how hippocampal anatomy was examined. Many studies

investigate the hippocampus as a whole, while others divide the hippocampus into anterior and

posterior portions, or focus on the volumetric ratio between the anterior and posterior segments.

Different functions have been ascribed to anterior and posterior portions (Poppenk,

Evensmoen, Moscovitch, & Nadel, 2013; Strange, Witter, Lein, & Moser, 2014; Zeidman &

Maguire, 2016), and this could also have influenced extant results.

Finally, in some studies, differing hippocampal volume-task relationships have been

identified for sub-groups of participants that were not apparent across the whole sample.

However, these participant sub-groups are not always directly compared to each other making

it difficult to have confidence in any reported differences. In addition, sub-group analyses are

often deemed "exploratory" and "requiring replication", but it is rare that such follow-up

occurs.

In the current study we sought to overcome these issues in a comprehensive

investigation of the relationship between hippocampal grey matter volume and scene

imagination, autobiographical memory, future thinking and spatial navigation task

performance in a healthy sample. To ensure an adequate sample size, data were collected from

217 individuals with a wide range of ability. All participants performed tasks relating to each

of the aforementioned cognitive functions and hippocampal grey matter volume was derived

from structural MRI scans with an isotropic voxel resolution of 0.8mm. Analyses were first

performed on each task's main outcome measure. As concerns could be raised that these four

measures are over-general, for completeness, we also examined the sub-measures of each task

(an additional 21 performance variables). This allowed us to investigate potential nuances that

may exist in hippocampal volume-task performance relationships. As reviewed above, the

mixed results in the literature made the formulation of clear hypotheses difficult. Consequently,

our aim in the data analyses was to reflect the range of approaches previously pursued in the

literature, and to perform a broad and inclusive evaluation of the relationship between

hippocampal grey matter volume and task performance.

We have reported the main outcome measures of the four cognitive tasks in two

previous papers that asked different research questions to those under consideration here. Clark

et al. (2019) assessed the relationships between scene imagination, autobiographical memory,

future thinking and navigation in terms of task performance. Using mediation analyses, they

found that scores on the scene construction task explained the relationships between

autobiographical memory, imagining the future and spatial navigation task performance. Clark

& Maguire (2020) investigated how subjective questionnaire data were related to task

performance. This revealed that imagination and navigation questionnaires reflected

performance on their related tasks. Memory questionnaires, on the other hand, were associated

with autobiographical memory vividness. In the current study, we examined whether

performance on the cognitive tasks, including the main outcome measures but also their sub-

measures, were related to hippocampal grey matter volume. The structural MRI data have not

been published previously.

2. MATERIALS AND METHODS

2.1 Participants

Two hundred and seventeen healthy individuals were recruited, including 109 females and 108

males. The age range was restricted to 20-41 years old to limit the possible effects of ageing

(mean age = 29.0 years, *SD* = 5.60). Participants had English as their first language and

reported no history of psychological, psychiatric or neurological conditions. Our aim was to

assess people from the general population who would not be classed as having extreme

expertise in the domains of interest. Consequently, people with vocations such as taxi driving

(or those training to be taxi drivers), ship navigators, aeroplane pilots, or those with regular

hobbies including orienteering, or taking part in memory sports and competitions, were

excluded. Participants were reimbursed £10 per hour for taking part which was paid at study

completion. All participants gave written informed consent and the study was approved by the

University College London Research Ethics Committee (project ID: 6743/001) and conducted

in accordance with the guidelines of the Declaration of Helsinki.

2.2 Procedure

Participants completed the study over multiple visits. Structural MRI scans were acquired

during the first visit. Cognitive testing was conducted during visits two and three. The scene

imagination, autobiographical memory and future thinking tasks were completed during one

visit, while the navigation tasks were performed on a separate visit. All participants completed

all parts of the study.

2.3 Cognitive tasks

All tasks are published and were performed and scored as per their published use. Here, for the

reader's convenience, we describe each task briefly.

2.3.1 Scene imagination: the scene construction task

The scene construction task (Hassabis, Kumaran, Vann, et al., 2007) measures a participant's

ability to mentally construct an atemporal visual scene, meaning that the scene is not grounded

in the past or the future. Participants construct different scenes of commonplace settings. For

each of seven scenes, a short cue is provided (e.g. imagine lying on a beach in a beautiful

tropical bay) and the participant is asked to imagine the scene that is evoked and then describe

it out loud in as much detail as possible. Recordings are transcribed for later scoring.

Participants are explicitly told not to describe a memory, but to create a new atemporal scene

that they have never experienced before.

The main outcome measure is the "experiential index" which is calculated for each

scene and then averaged. In brief, it is composed of four elements: the content, participant

ratings of their sense of presence (how much they felt like they were really there) and perceived

vividness, participant ratings of the spatial coherence of the scene, and an experimenter rating

of the overall quality of the scene.

For the scene construction sub-measures, we separately investigated the four categories

that make up the content score, and also the spatial coherence rating.

To score the content, four categories of statement are identified; spatial references,

entity presence, sensory description and thoughts/emotions/actions. The spatial reference

category encompasses statements regarding the relative position of entities within the

environment or directions relative to the participant's vantage point. The entity category is a

count of how many distinct entities (e.g., objects, people, animals) were mentioned. The

sensory descriptions category consists of any statements describing (in any modality)

properties of an entity or the environment in general. Finally, the thoughts/emotions/actions

category covers any introspective thoughts or emotional feelings as well as the thoughts,

intentions, and actions of other entities in the scene. The final score of each category is the

average across the seven scenes.

The imagined scenes are also examined using the spatial coherence index. After each

scene is mentally constructed, participants are presented with a set of 12 statements, each

providing a possible qualitative description of the imagined scene. Participants are instructed

to indicate which statements they felt accurately described their construction, identifying as

many or as few as they thought appropriate. Eight of the statements indicate that aspects of the

scene were integrated (e.g. "I could see the whole scene in my mind's eye"), whereas four

indicate that aspects of the scene were fragmented (e.g. "It was a collection of separate

images"). One point is awarded for each integrated statement selected and one point taken away

for each fragmented statement. This yields a score between –4 and +8 that is then normalised

around zero. Any construction with a negative score is considered to be incoherent and

fragmented and scored at 0 so as not to over-penalise fragmented descriptions. The final spatial

coherence index was the average score of the seven scenes, ranging between 0 (totally

fragmented) and +6 (completely integrated).

Double scoring was performed on 20% of the data. We took the most stringent approach

to identifying across-experimenter agreement. Inter-class correlation coefficients, with a two-

way random effects model looking for absolute agreement indicated excellent agreement

among the experimenters (minimum score of 0.9; see Supplementary Methods Table S1). For

reference, a score of 0.8 or above is considered excellent agreement beyond chance.

2.3.2 Autobiographical memory: the autobiographical interview

In the autobiographical interview (AI; Levine, Svoboda, Hay, Winocur, & Moscovitch, 2002)

participants are asked to provide autobiographical memories from a specific time and place

over four time periods – early childhood (up to age 11), teenage years (aged from 11-17),

adulthood (from age 18 years to 12 months prior to the interview; two memories are requested)

and the last year (a memory from the last 12 months). Recordings are transcribed for later

scoring.

The AI has two main outcome measures; the number of "internal" and "external" details

included in the description of an event. Importantly, these two scores represent different aspects

of autobiographical memory recall. Internal details are those describing the event in question

(i.e. episodic details), and were of primary interest here. External details describe semantic

information concerning the event, or non-event information. Internal events are, therefore,

thought to be hippocampal-dependent, while external events are not. The two AI scores are

obtained by separately averaging performance for the internal and external details across the

five autobiographical memories.

For the AI sub-measures, we examined the five separate categories that comprise the

internal details outcome measure, as well as considering AI vividness ratings. While external

details can also be split into component categories, too few details were provided by the

participants to assess each of these individually.

Internal details are composed of event, place, time, perceptual, and thoughts/emotions

categories. The event category contains details regarding happenings or the unfolding of the

story, including individuals present, actions and reactions. The time category refers to any

details of the year, season, day or time of day wherein the event occurred. The place category

contains details that localise the event, both at the general level (e.g. a city), and more

specifically (e.g. to parts of a room). The perceptual category involves descriptions (in any

modality) of any aspect of the event. Finally, the thoughts/emotions category includes

descriptions of emotional states, thoughts or implications. The final score of each category is

the average across the five memories.

AI vividness ratings were examined given a recent finding that responses on various

memory questionnaires were associated with AI vividness, suggesting that vividness may be a

key process in autobiographical memory recall (Clark & Maguire, 2020). Vividness ratings are

collected for each memory in response to the question "How clearly can you visualize this

event?" on a 6-point scale from 1 (vague memory, no recollection) to 6 (extremely clear as if

it's happening now). An overall vividness rating was the average of the vividness ratings

provided for each autobiographical memory.

Double scoring was performed on 20% of the data, and there was excellent agreement

across the experimenters (minimum score of 0.81; see Supplementary Methods Table S2).

2.3.3 Future thinking

The future thinking task (Hassabis, Kumaran, Vann, et al., 2007) follows the same procedure

as the scene construction task, but requires participants to imagine three plausible future scenes

involving themselves (an event at the weekend; next Christmas; the next time they will meet a

friend). There are two main differences between the future thinking task and the scene

construction task. First, unlike scenes in the scene construction task, scenes in the future

thinking task involve 'mental time travel' to the future, so they have a clear temporal

dimension. Second, the cues for the scene construction task are somewhat more specific than

those employed in the future thinking task (see Hurley, Maguire, & Vargha-Khadem, 2011 for

more on this issue).

The scoring procedures for both the main outcome measure and the sub-measures are

the same as for the scene construction task. Double scoring identified excellent agreement

across the experimenters (minimum score of 0.88; see Supplementary Methods Table S3).

2.3.4 Navigation

Navigation ability was assessed using the paradigm described by (Woollett & Maguire, 2010).

A participant watches movie clips of two overlapping routes through an unfamiliar real town

(Blackrock, Dublin, Ireland) four times.

The main outcome measure for navigation performance is calculated by combining the

scores from the five tasks used to assess navigational ability. The sub-measures are the

performance scores of each individual task.

The tasks were as follows. First, was a movie clip recognition task; following each

viewing of the route movies, participants are shown four short movie clips – two from the

actual routes, and two distractors. Participants indicate whether they have seen a movie clip

before or not. Second, after all four route viewings are completed, recognition memory for

scenes from the routes is tested. The third task, proximity judgements, involves assessing

knowledge of the spatial relationships between landmarks from the routes. Fourth, the route

knowledge task, requires participants to place scene photographs from the routes in the correct

order as if travelling through the town. Finally, the sketch map task involves participants

drawing a sketch map of the two routes including as many landmarks as they can remember.

Sketch maps are scored in terms of the number of road segments, road junctions, correct

landmarks, landmark positions, the orientation of the routes and an overall map quality score

from the experimenters. Double scoring was performed on 20% of the sketch maps, and there

was excellent agreement among the experimenters (minimum of 0.89; see Supplementary

Methods Table S4).

2.4 Self-report measures

2.4.1 Plymouth Sensory Imagery Questionnaire, Appearance subscale (Andrade, May,

Deeprose, Baugh, & Ganis, 2014). The PSIQ Appearance subscale assesses the vividness of

visual imagery. The subscale requires participants to imagine three scenarios: a bonfire, a

sunset, and a cat climbing a tree. They then rate the visual image they generated on an 11-point

scale from 0 (no image at all) to 10 (vivid as real life). Scores on the three scenarios are summed

to create a total score out of 30. Responses on the PSIQ have previously been associated with

performance on the scene construction, autobiographical memory, and future thinking tasks

(Clark & Maguire, 2020).

2.4.2 Santa Barbara Sense of Direction Scale (Hegarty, Richardson, Montello, Lovelace, &

Subbiah, 2002). The SBSODS assesses spatial and navigational abilities, preferences and

experiences. Fifteen statements are presented, with participants indicating their level of

agreement with each statement. Ratings are made on a 7-point scale from 1 (strongly agree) to

7 (strongly disagree). Seven statements are positively coded ("I am very good at giving

directions") and eight are reverse scored ("I don't have a very good "mental map" of my

environment"). Scores are summed across the 15 statements to create a total score out of 105

(where a low score reflects good navigation ability). Responses on the SBSODS have been

associated with multiple measures of navigation, including the spatial navigation task used in

the current study (Clark & Maguire, 2020; Hegarty et al., 2002).

2.5 Statistical analyses of the behavioural data

Data were summarised using means and standard deviations, calculated in SPSS v22. There were no missing data, and no data needed to be removed from any analysis.

2.6 MRI data acquisition

Three MRI scanners were used to collect the structural neuroimaging data. All scanners were Siemens Magnetom TIM Trio systems with 32 channel head coils and were located at the same imaging centre, running the same software. The sequences were loaded identically onto the individual scanners. Participant set-up and positioning followed the same protocol for each scanner.

Whole brain volumetric images had an isotropic resolution of 800μm × 800μm × 800μm and were obtained using magnetisation transfer (MT) saturation maps derived from a multi-parameter mapping (MPM) quantitative imaging protocol (Callaghan et al., 2015; Weiskopf et al., 2013). This protocol consisted of the acquisition of three multi-echo gradient acquisitions with either proton density (PD), T1 or MT weighting. Each acquisition had a repetition time, TR, of 25 ms. PD weighting was achieved with an excitation flip angle of 6^0, which was increased to 21^0 to achieve T1 weighting. MT weighting was achieved through the application of a Gaussian RF pulse 2 kHz off resonance with 4ms duration and a nominal flip angle of 220^0. This acquisition had an excitation flip angle of 6^0. The field of view was 256mm head-foot, 224mm anterior-posterior (AP), and 179mm right-left (RL). The multiple gradient echoes per contrast were acquired with alternating readout gradient polarity at eight equidistant echo times ranging from 2.34 to 18.44ms in steps of 2.30ms using a readout bandwidth of 488 Hz/pixel. Only six echoes were acquired for the MT weighted volume to facilitate the off-resonance pre-saturation pulse within the TR. To accelerate the data acquisition, partially

parallel imaging using the GRAPPA algorithm was employed in each phase-encoded direction

(AP and RL) with forty reference lines and a speed up factor of two. Calibration data were also

acquired at the outset of each session to correct for inhomogeneities in the RF transmit field

(Lutti, Hutton, Finsterbusch, Helms, & Weiskopf, 2010; Lutti et al., 2012).

2.7 MRI data pre-processing

The data from the MPM protocol were processed for each participant using the hMRI toolbox

(Tabelow et al., 2019) within SPM12 (www.fil.ion.ucl.ac.uk/spm). The default toolbox

configuration settings were used, with the exception that correction for imperfect spoiling was

additionally enabled. The output MT saturation map used for the volumetric analyses

quantified the degree of saturation of the steady state signal having accounted for spatially

varying T_1 times and RF field inhomogeneity (Helms, Dathe, & Dechent, 2008; Weiskopf et

al., 2013).

Each participant's MT saturation map was then segmented into grey matter probability

maps using the unified segmentation approach (Ashburner & Friston, 2005), but with no bias

field correction (since the MT saturation map does not suffer from any bias field modulation).

Inter-subject registration was performed using DARTEL, a nonlinear diffeomorphic algorithm

(Ashburner, 2007), as implemented in SPM12. This algorithm estimates the deformations that

best align the tissue probability maps by iteratively registering them with their average. The

resulting DARTEL template and deformations were used to normalize the tissue probability

maps to the stereotactic space defined by the Montreal Neurological Institute (MNI) template

(at 1 x 1 x 1mm resolution), with an isotropic Gaussian smoothing kernel of 8mm full width at

half maximum (FWHM).

2.8 Primary VBM analyses

Our primary analyses investigating the associations between cognitive task performance and

hippocampal grey matter volume were performed using voxel-based morphometry (VBM)

(Ashburner & Friston, 2000; Mechelli, Price, Friston, & Ashburner, 2005).

First, we investigated the relationship between hippocampal volume and the main

outcome measures for each of the cognitive tasks assessing scene imagination,

autobiographical memory, future thinking and navigation. We then examined the associations

between hippocampal volume and each of the sub-measures from these tasks.

For each performance measure, statistical analyses were carried out using multiple

linear regression models. Six regressors were included in each model, the cognitive task

performance measure of interest and five covariates: one each for age, gender and total

intracranial volume, and two regressors to model the effects of the different scanners (only two

regressors are required because one scanner is taken as the baseline, with the two regressors

then modelling any effects that are different between the other scanners and the baseline

scanner). The dependent variable was the smoothed and normalised grey matter volume.

2.8.1 Whole brain VBM

Analyses were first carried out voxel-wise across whole brain grey matter using an explicitly

defined mask which was generated by averaging the smoothed grey matter probability maps in

MNI space across all subjects. Voxels for which the grey matter probability was below 80%

were excluded from the analysis. Two-tailed t-tests were used to investigate the relationships

between cognitive task performance and grey matter volume, with statistical thresholds applied

at $p < 0.05$ family-wise error (FWE) corrected for the whole brain, and a minimum cluster size

of 5 voxels.

As our main focus was on the relationship between cognitive task performance and

hippocampal grey matter volume, in the main text we report only findings pertaining to the

hippocampus. However, performing analyses at the whole brain level meant that we could

apply the recommended statistical threshold to the analyses (Nichols et al., 2017; Poldrack et

al., 2008). Consequently any significant relationships identified would be supported by the

strongest evidence. In addition, performing analyses at the whole brain level also allowed us

to investigate whether any non-hippocampal brain regions were associated with cognitive task

performance – these results are reported in the Supplementary Results, although there were

very few.

2.8.2 Hippocampal ROI VBM

Following the whole brain analysis, we focused on the hippocampus using anatomical

hippocampal masks. This allowed us to investigate whether any weaker relationships existed

between hippocampal grey matter volume and task performance that did not reach the statistical

threshold required for the whole brain analyses. The masks were manually delineated on the

group-averaged MT saturation map in MNI space (1mm $\times$ 1mm $\times$ 1 mm) using ITK-SNAP

(www.itksnap.org). As in Poppenk & Moscovitch (2011) and Brunec et al. (2019), the anterior

hippocampus was delineated using an anatomical mask that was defined in the coronal plane

and proceeded from the first slice where the hippocampus can be observed in its most anterior

extent until the final slice of the uncus. The posterior hippocampus was defined from the first

slice following the uncus until the final slice of observation in its most posterior extent (see

Dalton, Zeidman, Barry, Williams, & Maguire, 2017 for more details). The whole hippocampal

mask comprised the combination of the anterior and posterior masks. The masks were

examined separately for the left and right hippocampus, and as bilateral masks.

Two-tailed t-tests were used to investigate the relationships between cognitive task

performance and grey matter volume within the hippocampal masks. Voxels were regarded as

significant when falling below an initial whole brain uncorrected voxel threshold of $p < 0.001$,

and then a small volume correction threshold of $p < 0.05$ FWE corrected for each mask, with

a minimum cluster size of 5 voxels. Given that all significant results observed using the anterior

and posterior unilateral masks were also evident using the bilateral masks, we report the results

of the bilateral masks. Where significant results within the masks were identified, the whole

brain analysis was re-examined with the threshold set to $p < 0.001$ uncorrected to assess

whether the effects identified within the hippocampal masks were due to leakage from adjacent

brain regions.

2.9 Auxiliary analyses using extracted hippocampal volumes

Following our main analyses, we performed a series of auxiliary investigations to further

scrutinise potential relationships between hippocampal grey matter volume and scene

imagination, autobiographical memory, future thinking and navigation task performance.

The basis of these auxiliary analyses was the hippocampal grey matter volume for each

participant that was extracted using 'spm_summarise'. The whole, anterior and posterior

anatomical hippocampal masks (described above) were applied to each participant's smoothed

and normalised grey matter volume maps, and the total volume within each mask was

extracted. It has been suggested that the posterior:anterior hippocampal volume ratio shows

stronger relationships to memory and navigation performance than raw anterior and posterior

hippocampal volumes alone (Brunec et al., 2019; Poppenk & Moscovitch, 2011). Therefore,

in line with these studies, we also calculated each participant's posterior:anterior hippocampal

volumetric ratio. We did this by dividing the posterior hippocampal volume by the anterior

hippocampal volume, providing a ratio whereby values greater than 1 were indicative of a

larger posterior over anterior hippocampus and those less than 1 indicated a larger anterior over

posterior hippocampus.

Overall, across all our auxiliary investigations (described in detail below) we examined

the relationships between four different measures of hippocampal grey matter volume (whole,

anterior, posterior, ratio) for each of our 26 task performance measures. Given the extensive

nature of these analyses, we needed to ensure that appropriate correction was made to the

statistical thresholds. Typically, FWE corrections (such as Bonferroni) are employed in this

regard. However, FWE correction acts to ensure that the probability of a *single* false positive

result remains at the critical alpha (typically $p < 0.05$); a highly effective method if the need to

avoid false positives is high. However, this level of control comes at a price, especially when

controlling for a large number of statistical tests, as it means that the effects of any single

variable have to be particularly large to reach the adjusted significance level. Thus, while false

positives are avoided, the risk of incorrectly rejecting a true result is introduced.

An alternative method of statistical correction is to control the false discovery rate

(Benjamini & Hochberg, 1995) which influences the *proportion* of significant results identified

that are false positives. Therefore, a FDR of $p < 0.05$ allows for 5% false positive results across

the tests performed. Controlling for the FDR is, therefore, more lenient than performing FWE

corrections, as it accommodates the potential for a greater number of false positive results but,

by doing so, also reduces the chances of incorrectly rejecting a true result. Applying our

statistical correction in terms of the FDR provided, therefore, a compromise between the need

to adjust our statistical thresholds for multiple comparisons, while not removing all possibility

of identifying true relationships among the data features.

Across the auxiliary analyses, the FDR was set to $p < 0.05$ for each family of tests. A

family was defined as all the tests performed for each set of auxiliary analyses for one cognitive

task. For example, when testing for relationships between hippocampal volume and scene

construction task performance, we had 4 measures of hippocampal volume (whole, anterior,

posterior, ratio) and 6 measures of performance (experiential index, spatial references, entities

present, sensory descriptions, thoughts/emotions/actions, spatial coherence), totalling 24

separate statistical tests. The FDR was thus set to $p < 0.05$ for these 24 tests.

The following auxiliary analyses were performed in SPSS v25 with FDR thresholding

and Benjamini-Hochberg adjusted p values calculated using the resources provided by

McDonald (2014).

2.9.1 Partial correlations using extracted hippocampal volumes

We performed partial correlation analyses between each of the hippocampal volumetric

measures (whole, anterior, posterior, ratio) and each of our measures of task performance. As

with the primary VBM analyses, age, gender, total intracranial volume and MRI scanner were

included as covariates.

2.9.2 Sub-group analyses using extracted hippocampal volumes

2.9.2.1 Effects of gender

To directly investigate the effects of gender, we divided the sample into male (n = 108) and

female (n = 109) participants. We first performed separate partial correlations for the two

participant groups for each of the hippocampal volumetric measures (whole, anterior, posterior,

ratio) and each of our measures of task performance. Age, total intracranial volume and MRI

scanner were included as covariates. If a significant volume-task relationship was identified in

either the male or female group, our intention was to then compare the male and female

relationships directly. However, this was never required (see section 3.3.2).

2.9.2.2 Median split: direct comparison

Participants were first allocated to groups depending on their performance on a task compared to the median score for that task. If their score was less than or equal to the median value they were assigned to the low performing group, scores greater the median resulted in allocation to the high performing group. Due to the distribution of performance scores, the navigation movie clip recognition test could not be divided into groups (as most participants performed at ceiling) and so was not included in these analyses. Our four measures of hippocampal volume were then compared between the low and high performing groups for each task using univariate ANCOVAs with the hippocampal volume measure as the dependent variable, performance group as the main predictor and age, gender, total intracranial volume and MRI scanner as covariates.

2.9.2.3 Median split: partial correlations

We next investigated if different associations between hippocampal volume and cognitive task scores existed in the high and low performing participants. As before, because the navigation movie clip recognition test could not be divided into groups it was not included in these analyses. For each task, partial correlations were performed separately for the two groups between the hippocampal volume measures and the task performance measures, with age, gender, total intracranial volume and MRI scanner included as covariates. If a significant relationship was identified in either the low or high performance group, our intention was to then compare the group correlations directly. However, this was never required (see section 3.3.3).

2.9.2.4 Best versus worst

The best and worst performers for each task were identified. This was approximately the top

and bottom 10% (n ~ 20 in each group). The exact number of participants allocated to each

group varied for each task depending on the distribution of performance scores and to ensure

the number of participants in each group was as similar as possible. As the navigation movie

clip recognition test could not be split into approximately equal groups of best and worst

performers it was not included in these analyses. Our four measures of hippocampal volume

were then compared between the best and worst performing participants for each task using

univariate ANCOVAs with the hippocampal volume measure as the dependent variable,

performance group as the main predictor and age, gender, total intracranial volume and MRI

scanner as covariates.

2.9.2.5 Effects of self-reported ability

Self-reported ability was assessed using the PSIQ (Andrade et al., 2014) for scene imagination,

autobiographical memory and future thinking, and the SBSODS (Hegarty et al., 2002) for

navigation. Our aim here was to investigate whether the interaction between self-reported

ability and hippocampal volume was associated with task performance, following the method

of He & Brown (2020). As such, eight interaction terms were created: responses on the PSIQ

and SBSODS by the four measures of hippocampal grey matter volume. Partial correlation

analyses were then performed between the interaction terms and task performance, with age,

gender, total intracranial volume and MRI scanner as covariates as in the previous analyses,

with the additional covariates of the relevant questionnaire and hippocampal volume

measurement to ensure that any identified relationships were due to the interaction of self-

report and volume and not either individually.

2.10 Multivariate analyses using latent variables

The analyses detailed above are all univariate and task-based. A multivariate approach using

latent variables derived from across the cognitive data may provide additional insights into

associations between task performance and hippocampal grey matter volume. For example,

this method offered increased statistical power and enabled data driven investigations,

eschewing a reliance on conceptually driven cognitive constructs, and so had the potential to

increase the generalisability of findings beyond the specific tasks in question.

To identify latent variables across the cognitive data, two Principal Component

Analyses (PCA) were performed using SPSS v25 with varimax rotation and an eigenvalue cut-

off of 1. The first PCA investigated possible latent variables across the four main outcome

measures of the scene imagination, autobiographical memory, future thinking and navigation

tasks, and the second PCA examined all 21 task sub-measures. Factor scores for the identified

latent variables were extracted using the Anderson-Rubin method. The relationships between

the latent variable factor scores and hippocampal grey matter volume were examined using the

VBM and partial correlation approaches described above.

3. RESULTS

3.1 Cognitive task performance

A summary of the outcome measures for the cognitive tasks is shown in Table 1. A wide range

of scores was obtained for all variables with the exception of navigation movie clip recognition,

where performance was close to ceiling.

  Table 1. Means and standard deviations for the main outcome measures and sub-measures of
  the cognitive tasks and questionnaires.

Variable	Mean	Standard Deviation
Main outcome measures		
Scene construction experiential index (/60)	40.50	6.08
Autobiographical interview internal details (total number)	23.95	7.25
Autobiographical interview external details (total number)	5.35	3.20
Future thinking experiential index (/60)	39.12	7.23
Navigation (/250)	143.46	35.83
Sub-measures		
Scene construction		
Spatial references (total number)	3.47	1.56
Entities present (total number)	9.90	2.92
Sensory descriptions (total number)	12.25	3.46
Thoughts/emotions/actions (total number)	3.36	1.70
Spatial coherence index (/6)	2.88	1.57
Autobiographical interview		
Internal events (total number)	11.02	3.88
Internal time (total number)	1.43	0.69
Internal place (total number)	2.23	0.63
Internal perceptual (total number)	5.91	3.07
Internal thoughts/emotions (total number)	3.37	1.69
Vividness ratings (/6)	4.62	0.74
Future thinking		
Spatial references (total number)	2.53	1.65
Entities present (total number)	10.31	3.28
Sensory descriptions (total number)	8.81	3.59
Thoughts/emotions/actions (total number)	5.18	2.28
Spatial coherence index (/6)	2.63	1.77
Navigation		
Movie clip recognition (/16)	15.44	0.86
Scene recognition (/32)	29.45	1.89
Proximity judgements (/10)	7.48	1.43
Route knowledge (/24)	11.98	4.62
Sketch map	79.11	30.91
Questionnaires		
Plymouth Sensory Imagery Questionnaire; Appearance	21.89	5.63
Santa Barbara Sense of Direction Scale	50.83	15.56

3.2 Primary VBM analyses

3.2.1 Whole brain VBM

As our main focus was on the relationship between cognitive task performance and

hippocampal grey matter volume, here we report only findings pertaining to the hippocampus

– any regions identified outside the hippocampus are reported in the Supplementary Results.

No significant relationships between cognitive task performance and hippocampal grey

matter volume were identified for any of the main outcome measures of the tasks assessing

scene imagination, autobiographical memory, future thinking or navigation. This was also the

case for the sub-measures of these tasks.

3.2.2 Hippocampal ROI VBM

No relationships between cognitive task performance and hippocampal grey matter volume

were identified using any of the hippocampal masks for the main outcome measures from the

tasks examining scene imagination, autobiographical memory, future thinking or navigation.

No associations were observed between performance on the scene construction task

sub-measures and hippocampal grey matter volume using any of the hippocampal masks.

For the AI sub-measures, one potential relationship was identified. When using the

posterior hippocampal mask, a cluster of 10 voxels, located towards the anterior portion of the

posterior mask in the left hippocampus, was found to be negatively associated with the number

of internal place references (Figure 1a; peak coordinates = -34, -23, -12, peak t = 3.28, p $_{FWE}$

$_{posterior\ hippocampus\ ROI\ corrected}$ = 0.046). This relationship, however, was not significant when

correcting for the whole hippocampus mask (p $_{FWE\ whole\ hippocampus\ ROI\ corrected}$ = 0.069). Reducing

the statistical threshold to p < 0.001 uncorrected for the whole brain confirmed the cluster was

confined to the hippocampus and not due to leakage from adjacent brain regions (Figure 1b).

However, visualisation at this lower threshold also demonstrated that the identified relationship

was not unique to the hippocampal voxels, with a number of other brain regions also being

negatively associated with the AI internal place references at this liberal statistical threshold.

For the future thinking sub-measures, no associations were observed between cognitive

task performance and hippocampal grey matter volume.

Considering the navigation sub-measures, one potential relationship was identified.

When using the anterior hippocampal mask, a cluster of 17 voxels, located at the edge of the

mask in the right hippocampus, was found to be positively associated with performance on the

navigation route knowledge sub-measure (Figure 1c; peak coordinates = 27 -8 -29, peak t =

3.31, p $_{\text{FWE anterior hippocampus ROI corrected}}$ = 0.028). This relationship, however, was not significant

when correcting for the whole hippocampus mask (p $_{\text{FWE whole hippocampus ROI corrected}}$ = 0.062).

Reducing the statistical threshold to p < 0.001 uncorrected for the whole brain clearly indicated

that the identified hippocampal voxels were located on the edge of a larger cluster that was in

fact centred on the right anterior fusiform and parahippocampal cortices (Figure 1d, cluster size

= 1557, peak coordinates = 29 -5 -37, peak t = 3.87).

3.3 Auxiliary analyses using extracted hippocampal volumes

We next performed a series of auxiliary analyses to further investigate possible relationships

between cognitive task performance and hippocampal grey matter volume. The latter was

extracted for each participant using the predefined anatomical hippocampal masks. The mean

grey matter volume for the whole hippocampus was 4456.49 mm^3 (SD = 373.74 mm^3), for the

anterior mask was 2303.70 mm^3 (SD = 215.07 mm^3) and for the posterior mask was 2152.79

mm^3 (SD = 182.66 mm^3), with a mean posterior/anterior ratio of 0.94 (SD = 0.057), where a

ratio less than 1 indicates greater anterior relative to posterior hippocampal volume.

3.3.1 Partial correlations using extracted hippocampal volumes

No relationships were identified between hippocampal grey matter volume and performance for any of the main or sub-measures of the tasks assessing scene imagination, autobiographical memory, future thinking or navigation (see Supplementary Results Table S5 for details). These partial correlation findings, therefore, support those of the primary VBM analyses, as well as extending this to include the posterior:anterior hippocampal volume ratio.

3.3.2 Sub-group analyses using extracted hippocampal volumes

3.3.2.1 Effects of gender

We next divided the sample into male and female participants. No relationships between cognitive task performance and hippocampal grey matter volume were observed in either group for any of the main or sub-measures (see Supplementary Results Tables S6 and S7 for details). As no relationships were detected in either group, direct comparison of the relationships was not relevant.

3.3.2.2 Median split: direct comparison

Details of the groups that were created when the sample was divided by median splits of performance are provided in the Supplementary Results (Tables S8). Comparing hippocampal grey matter volume measurements between the two groups determined by their median performance on each main and sub-measure found no differences between the groups for any task (see Supplementary Results Table S9 for details).

3.3.2.3 Median split: partial correlations

No significant relationships between hippocampal grey matter volume and cognitive task main and sub-measures were identified in either the low or high performing groups (see

Supplementary Results Tables S10 and S11 for details). As no relationships were detected in

either group, direct comparison of the relationships was not relevant.

3.3.2.4 Best versus worst

Details of the groups that were created when the sample was divided in terms of best and worst

performers are provided in the Supplementary Results (Tables S12). Taking the best and worst

performing participants for each cognitive task main and sub-measure and comparing the

hippocampal grey matter volume measurements between the two groups also showed no

significant differences between the groups (see Supplementary Results Table S13 details).

3.3.2.5 Effects of self-reported ability

The results of these analysis are provided in the Supplementary Results (Table S14).

No significant associations were observed between any of the scene construction task

main and sub-measures and the interaction between self-reported ability (determined by the

PSIQ) and hippocampal grey matter volume.

For autobiographical memory, a potential relationship was identified for AI vividness,

but no other main or sub-measure. For AI vividness, an association was observed between AI

vividness ratings and the PSIQ by whole hippocampus volume interaction ($r = 0.24$, p_{FDR}

$_{corrected} = 0.016$), and with the PSIQ by posterior hippocampal volume interaction ($r = 0.22$, p

$_{FDR\ corrected} = 0.013$). This suggests that self-reported imagery ability may be influencing the

relationship between hippocampal volume and AI vividness ratings.

To better understand these interactions, simple slopes and region of significance

analyses were performed using the interactions toolbox (Long, 2019) in R 3.6.1 (R Core Team,

2017), with regions of significance calculated using the Johnson-Neyman technique (Johnson

& Fay, 1950).

First, we focused on the relationship between AI vividness and the PSIQ by whole

hippocampus interaction. Repeating the partial correlation analysis as a regression (with AI

vividness as the dependent variable, PSIQ, whole hippocampal volume and the PSIQ by whole

hippocampal volume interaction as predictors, and age, gender, scanner and ICV as covariates),

identified that for the PSIQ by whole hippocampus volume interaction, the unstandardized beta

value was 0.0000710. Performing simple slopes to decompose this relationship identified that

for individuals who were 1 standard deviation below the mean PSIQ score, the unstandardized

beta value was -0.000859 (t = -4.15, $p_{\text{FDR corrected}} < 0.001$). For individuals with a mean PSIQ

score, the unstandardized beta value was -0.000459 (t = -2.78, $p_{\text{FDR corrected}} = 0.009$), while for

individuals who were 1 standard deviation above the mean PSIQ score, the unstandardized beta

value was -0.0000583 (t = -0.28, $p_{\text{FDR corrected}} = 0.92$). As such, in individuals with low PSIQ

scores, there was a negative relationship between whole hippocampal volume and AI vividness

that was not evident in individuals with high PSIQ scores. A region of significance analysis

found that the negative relationship existed in individuals with PSIQ scores less than 23.65.

We then investigated the relationship between AI vividness and the PSIQ by posterior

hippocampus interaction. Repeating the partial correlation analysis as a regression (with AI

vividness as the dependent variable, PSIQ, posterior hippocampal volume and the PSIQ by

posterior hippocampal volume interaction as predictors, and age, gender, scanner and ICV as

covariates), identified that for the PSIQ by posterior hippocampus volume interaction, the

unstandardized beta value was 0.000157. Performing simple slopes identified that for

individuals who were 1 standard deviation below the mean PSIQ score, the unstandardized beta

value was -0.00180 (t = -4.53, $p_{\text{FDR corrected}} < 0.001$). For individuals with a mean PSIQ score,

the unstandardized beta value was -0.000919 (t = -3.24, $p_{\text{FDR corrected}} = 0.003$), while for

individuals who were 1 standard deviation above the mean PSIQ score, the unstandardized beta

value was -0.0000343 (t = -0.098, $p_{\text{FDR corrected}} = 0.92$). As with the whole hippocampal volume

interaction, in individuals with low PSIQ scores, there was a negative relationship between

posterior hippocampal volume and AI vividness that was not apparent in individuals with high

PSIQ scores. A region of significance analysis found that the negative association existed in

individuals with PSIQ scores less than 24.13.

For all of the future thinking and navigation main and sub-measures, no significant

findings were observed when investigating the association between cognitive task performance

and the interaction between self-reported ability and hippocampal grey matter volume.

3.4 Multivariate analyses using latent variables

The detailed outcomes of the multivariate analyses can be found in the Supplementary Results.

In summary, performing a PCA with the four main outcome measures of the scene imagination,

autobiographical memory, future thinking and navigation tasks resulted in a single component

that explained 56.05% of the variance (Supplementary Results Table S15). Conducting a VBM

analysis with this latent variable identified no significant relationship between hippocampal

grey matter volume and the latent variable factor scores, both at the whole brain level and when

using the hippocampal ROIs. Performing partial correlations between the extracted

hippocampal grey matter volumes and the latent variable factor scores also identified no

significant associations (Supplementary Results).

The second PCA using all 21 task sub-measures identified 6 components that explained

65.01% of the variance (Supplementary Results Tables S16 and S17, and Figure S1). One

reassuring finding of this PCA was the broad correspondence of the components to the

cognitive tasks. For example, all of the navigation sub-measures loaded onto Component 2,

and all of the autobiographical memory sub-measures, excluding internal time, were placed in

Component 5. The scene construction and future thinking sub-measures showed greater

interrelatedness, which is not surprising given the close similarity of the task requirements.

Component 1 comprised the content sub-measures of both tasks, and Component 3 contained

their spatial coherence indices, as well as autobiographical memory vividness ratings.

Component 4 encompassed the thoughts and emotions elements of scene construction, future

thinking and autobiographical memory. Finally, Component 6 corresponded predominantly to

the autobiographical memory sub-measure of internal time. We then examined whether there

were any relationships between these latent constructs and hippocampal grey matter volume.

Using VBM (at both the whole brain level and in the anatomical ROIs) or when performing

partial correlations using the extracted hippocampal grey matter volumes (Supplementary

Results Table S18), no significant relationships between task performance and hippocampal

grey matter volume were evident.

4. DISCUSSION

Marked differences exist across healthy individuals in their ability to imagine visual scenes,

recall autobiographical memories, think about the future and navigate in the world. One

possible source of this variance may be an individual's hippocampal grey matter volume. Here,

however, in a large sample of young, healthy adult participants, we found little evidence that

hippocampal grey matter volume was related to task performance. This was the case across

different methodologies (VBM and partial correlations), whole brain or hippocampal ROIs,

different sub-groups of participants (divided by gender, task performance and self-reported

ability) and when examining latent variables derived from across the cognitive tasks.

Hippocampal grey matter volume may not, therefore, significantly influence performance on

tasks known to require the hippocampus in healthy people.

816  Of the 52 main VBM analyses we performed, only two significant associations between

hippocampal grey matter volume and task performance were noted. First, a negative

association between a cluster of ten voxels and the measure of AI internal place was observed.

However, this result was only significant when correcting for the posterior hippocampal mask,

and was not replicated when a partial correlation analysis was performed on the extracted

hippocampal volume data. Indeed, the correlation coefficient between the posterior

hippocampal volume and AI internal place was only -0.053. In the second result, a cluster of

seventeen voxels was positively associated with navigation route knowledge when correcting

for the anterior hippocampal mask only. However, as shown in Figure 1d, these voxels are

clearly on the edge of a larger cluster that was in fact centred on the anterior fusiform and

parahippocampal cortices. Overall, given the number of individual analyses performed and the

nature of the findings, neither of these results is particularly convincing.

Of the 712 auxiliary analyses performed, only two were significant – AI vividness

ratings and the PSIQ by whole hippocampal volume interaction, and AI vividness ratings and

the PSIQ by posterior hippocampal volume interaction. Further examination of these

interactions found that in individuals with low PSIQ scores (less than 23.65 and 24.13

respectively), there was a negative relationship between whole and posterior hippocampal

volume and AI vividness. However, it should be borne in mind that the effect size of the

interactions was small (r values of 0.24 and 0.22) and observing only two significant results

across the large number of analyses performed is well within the number of false positive

results expected, even with statistical correction applied.

The overwhelming number of our analyses produced null results. We are, of course,

mindful that null results can be difficult to interpret and that an absence of evidence is not

evidence of absence. Nevertheless, several factors motivate confidence in interpreting the

current results as showing little evidence of a relationship between hippocampal grey matter

volume and performance on our tasks of interest.

First, we had a sample of 217 participants. Samples of this size are much better powered

to find true and reliable relationships between brain structure and cognitive ability than the

smaller sample sizes (typically between 20 and 30 participants) often tested in previous studies

(Kharabian Masouleh et al., 2019; Kong & Francks, 2019). Indeed, the only other study to

employ a substantial sample was that of Weisberg et al. (2019; n = 90) who also found no

relationship between hippocampal grey matter volume and navigation task performance.

Second, our investigation was wide-ranging and thorough with 26 different outcome

measures, each relating to different aspects of scene imagination, autobiographical memory,

future thinking and navigation. It seems unlikely that if a relationship existed between

hippocampal grey matter volume and these cognitive functions that it would not have been

identified by at least one of these performance measures.

Third, our sample was specifically recruited to provide a wide variance in our measures

of interest. With the exception of the navigation movie clip recognition sub-measure (where

performance was at, or very close to, ceiling) there were large individual differences in ability

on all of the tasks. In addition, hippocampal grey matter volume was heterogeneous across the

sample. We were, therefore, well-placed to identify any potential relationships between

hippocampal volume and task performance, yet few were found.

Fourth, we investigated possible relationships across the whole brain, but also using

anatomically-defined hippocampal ROIs. Despite the reduced level of statistical correction

required when constraining analyses to hippocampal anatomical ROIs, we still found no

significant relationships. It could be argued that the already liberal statistical thresholds should

be reduced even further, but this would simply increase the chances of false positive results,

given our large sample size and the considerable number of analyses that were performed.

Fifth, similar outcomes were observed using two different analysis methods. While our

primary analyses involved VBM, by extracting the whole, anterior and posterior hippocampal

volumes for each participant and additionally calculating the posterior:anterior volume ratio,

we could also perform partial correlations between each of the volume measurements and our

measures of task performance. The results, therefore, do not seem to be consequent upon a

specific analysis technique, but instead there was a consistent pattern of null results.

Sixth, we also divided the sample into multiple sub-groups, investigating male and

female participants, assessing whether there was any effect of cognitive performance (using

both a median split, and by examining only the best and worst performers for each task

measure), and self-reported ability. No significant associations between hippocampal volume

and performance were identified.

Finally, we also conducted multivariate analyses, using PCA to identify latent variables

from across the cognitive tasks. This approach offered increased statistical power and enabled

data driven investigations, eschewing a reliance on conceptually driven cognitive constructs,

and so had the potential to increase the generalisability of findings beyond the specific tasks in

question. To the best of our knowledge, formal PCA analyses using the sub-measures of the

tasks employed here have not been reported previously. It was reassuring to find broad

correspondence between the components identified by the PCA and the cognitive tasks, and to

note that the spatial coherence and the emotional aspects of mental representations cut across

cognitive tasks. Of particular relevance for the questions under consideration here, and as with

the other analyses, no significant associations between hippocampal grey matter volume and

the identified latent variables were observed.

How do our null results correspond to the existing literature? We are not alone in

reporting a lack of association between hippocampal grey matter volume and cognitive ability

in healthy adults. No previous studies have investigated the hippocampal volume-scene

imagination relationship. Considering memory, a meta-analysis of 33 studies identified no

relationship between performance on laboratory-based memory tasks and hippocampal volume

(Van Petten, 2004). The one previous study that investigated future thinking found no evidence

of an association between hippocampal volume and future thinking performance – with voxels

instead being located outside of the hippocampus (Yang et al., 2020). Regarding navigation,

two previous studies have found no relationship between hippocampal volume and

navigational ability in samples drawn from the general population (Maguire, Spiers, et al.,

2003; Weisberg et al., 2019). Given the well-known "file drawer" problem (Franco, Malhotra,

& Simonovits, 2014; Rosenthal, 1979), the number of null results is likely to be higher than

those published in the literature. Our findings, therefore, of no relationship between

hippocampal volume and task performance, are not unique to the current sample.

We appreciate that other studies have reported hippocampal volume-task performance

relationships. For example, a positive association was identified between posterior

hippocampal grey matter volume and recollective memory ability (Poppenk & Moscovitch,

2011), and both posterior and anterior hippocampal grey matter volume have been associated

with various measures of spatial navigation, sometimes within specific subgroups (e.g. Bohbot

et al., 2007; Brown et al., 2014; Brunec et al., 2019; Chrastil et al., 2017; Hartley & Harlow,

2012; He & Brown, 2020; Schinazi et al., 2013; Sherrill et al., 2018). These disparate results,

however, may have been influenced by methodological factors. As alluded to, the sample sizes

used were typically small, with the largest being only a quarter of the current study (see

Kharabian Masouleh et al., 2019 for more on this issue). Moreover, previous studies have not

always been optimally analysed. For example, in some cases relevant covariates were not

included, or appropriate levels of statistical correction were not applied, or results were found

only in specific smaller sub-groups. Any of these issues could have affected the reliability of

the observed results.

The current results differ from those involving licensed London taxi drivers, where

increased posterior, but decreased anterior, hippocampal grey matter volume has been

consistently identified compared to a variety of control groups, including the same individuals

tested before and after they acquired The Knowledge (Maguire et al., 2000; Maguire, Woollett,

et al., 2006; Woollett et al., 2008; Woollett & Maguire, 2011). The training to become a London

taxi driver takes 3-4 years and approximately 50% of trainees fail to qualify. Those who

successfully qualify as London taxi drivers are, therefore, a unique population with extreme

spatial navigation expertise. Consequently, the divergence between our null results and those

in London taxi drivers is not surprising. Indeed, interpreting the current null results (and those

of Maguire, Spiers, et al., 2003, and Weisberg et al. 2019) together with the London taxi driver

findings may help to refine our understanding of the relationship between hippocampal volume

and navigation ability. Together these results suggest that measurable hippocampal volume

changes associated with ability may only be apparent following extreme spatial training rather

than being a marker of ability in the general population.

Why might measureable hippocampal volume changes only be detected in the context

of extreme expertise? As we did not address this question in the current study we can only

speculate. Taking the example of navigation, the vast majority of humans never need to acquire

the huge amount of spatial layout knowledge associated with being a licensed London taxi

driver. Hence, the hippocampus has likely evolved to accommodate representations of

environments that we use day-to-day, with sufficient capacity for a wide range of navigation

experiences. Moreover, proficiency, especially at the extreme end of the scale, may come at a

cost. For instance, in licensed London taxi drivers, grey matter volume is increased in the

posterior hippocampus relative to control groups, but this is accompanied by decreased anterior

hippocampal volume (Maguire et al., 2000; Maguire, Woollett, & Spiers, 2006), and also

poorer performance on some anterograde visuospatial memory testes (Woollett & Maguire,

2009; 2012; Woollett, Spiers, & Maguire, 2009). Consequently, hippocampal volume may

reflect a balance between sufficient functionality for the activities that a typical human

undertakes whilst not compromising other hippocampal operations in the service of extreme

expertise.

While we found that hippocampal volume seemed to be unrelated to task performance
in the general population, it is likely that some neural variations exist that can, at least in part,
explain the marked differences in performance observed in tasks such as those we employed.
One possibility is that while volumetric differences for the whole hippocampus, or even its
anterior or posterior portions, are not related to performance, effects might be observed for the
volume of specific hippocampal subfields. For example, larger CA3 volume has been
associated with reduced subjective confusion when recalling highly similar memories
(Chadwick, Bonnici, & Maguire, 2014). In addition, subiculum volume and combined left
dentate gyrus/CA2,3 volume has been positively associated with the number of internal details
on the autobiographical interview (Palombo et al., 2018). However, while these studies
highlight the potential for relationships between hippocampal subfield volumes and task
performance, they also suffer from the limitation of small sample sizes, and so additional work
is required to address this issue further.

Another possibility is that the volume of brain regions other than the hippocampus are
related to performance on our tasks of interest. Regions including, but not limited to, the
parahippocampal, retrosplenial, posterior cingulate, parietal, and medial prefrontal cortices
have been associated with scene imagination, autobiographical memory, future thinking and
navigation (e.g. Ciaramelli et al., 2017; Hassabis, Kumaran, & Maguire, 2007; Ramanan et al.,
2018; Schacter et al., 2012; Simons, Peers, Mazuz, Berryhill, & Olson, 2010; Stawarczyk &
D'Argembeau, 2015). However, as reported in the Supplementary Results, when using a $p <$
0.05 FWE statistical threshold across the whole brain we observed only one relationship – a
negative association between 22 voxels in the right parahippocampal cortex and AI vividness
ratings. More in-depth analyses using specific ROIs may highlight other relationships (e.g. the
positive association between the parahippocampal/fusiform cortices and navigation route
knowledge sub-measure we observed when using an uncorrected threshold), but these ROIs

would need to be motivated in advance to justify their use, going beyond the remit of the current

study.

Advances in neuroimaging mean it is now possible to investigate brain structure in more

detail than just grey matter volume. Quantitative neuroimaging techniques (Weiskopf,

Mohammadi, Lutti, & Callaghan, 2015) can be used to model different properties of tissue

microstructure, in particular that of myelination and iron. Studies using these techniques have,

for example, found relationships between myelination and iron in ageing (Callaghan et al.,

2014) and, in older participants, increased myelination and decreased iron tissue content has

been associated with verbal memory performance in the ventral striatum (Steiger, Weiskopf,

& Bunzeck, 2016). Tissue microstructure, both at the whole hippocampus level or within

specific subfields, or even in regions outside of the hippocampus, may offer explanations about

the variance observed in scene imagination, autobiographical memory, future thinking and

navigation ability and will be an interesting target for future work.

The connectivity between brain regions may also be an important factor in explaining

individual differences in task performance. Structural measures of connectivity, for example

diffusion-weighted imaging of hippocampal white matter pathways, have been associated with

both autobiographical memory and navigation ability (Hodgetts et al., 2017; Hodgetts et al.,

2020; Iaria, Lanyon, Fox, Giaschi, & Barton, 2008; Metzler-Baddeley, Jones, Belaroussi,

Aggleton, & O'Sullivan, 2011). Moreover, functional connectivity between the medial

temporal lobe and parietal-occipital regions may be related to questionnaire measures of

autobiographical memory (Sheldon, Farb, Palombo, & Levine, 2016), while the functional

connectivity of the posterior hippocampus and retrosplenial cortex has been linked with

responses on navigation questionnaires (Sulpizio, Boccia, Guariglia, & Galati, 2016). Both

structural and functional connectivity measures may, therefore, be associated with task

performance, perhaps to a greater extent than hippocampal grey matter volume, and represent

important avenues to explore in future studies.

As with functional connectivity, patterns of activity during rest or task-based fMRI may

be related to individual differences in performance. For example, higher vividness ratings when

performing mental imagery tasks have been reported to correlate with activity in the fusiform,

posterior cingulate, parahippocampal (Fulford et al., 2018) and visual (Dijkstra, Bosch, & van

Gerven, 2017; Keogh, Bergmann, & Pearson, 2020) cortices. In relation to autobiographical

memory, more vivid recall has been linked with higher levels of hippocampal (Gilboa,

Winocur, Grady, Hevenor, & Moscovitch, 2004) and precuneus (Richter, Cooper, Bays, &

Simons, 2016) activity, while superior memory precision has been related to increased angular

gyrus engagement during task performance (Richter et al., 2016). Better performance during

navigation tasks has also been associated with increased hippocampal activity (Hartley,

Maguire, Spiers, & Burgess, 2003; Ohnishi, Matsuda, Hirakata, & Ugawa, 2006). Individual

differences in ability may, therefore, be detectable in fMRI activity during task performance

although, as with the structural MRI studies outlined in Section 1, there is a need for large-

scale functional neuroimaging studies to examine this further.

1009

5. CONCLUSION

In this study we attempted to circumvent issues known to affect studies of healthy people that

assess relationships between hippocampal grey matter volume and performance on cognitive

tasks that require the hippocampus. Having tested a large sample of healthy participants with a

wide range of ability, no credible associations were identified. Consequently, variability in

hippocampal grey matter volume seems unlikely to explain individual differences in scene

imagination, autobiographical memory, future thinking and spatial navigation performance in

the general population. It may be that in healthy participants it is only in extreme situations, as

in the case of licensed London taxi drivers, that measurable ability-related hippocampus

volume changes are exhibited.

1020

1021

1022 **DECLARTION OF COMPETING INTEREST**
1023 The authors declare no competing interests.
1024
1025
1026 **DATA AVAILABILITY**
1027 Data and materials will be open-access once the construction of a dedicated data-sharing portal
1028 has been finalised. In the meantime, requests for data and materials can be sent to
1029 e.maguire@ucl.ac.uk.
1030
1031
FUNDING INFORMATION
The authors were supported by a Wellcome Principal Research Fellowship to E.A. Maguire
(101759/Z/13/Z) and the Centre by a Strategic Award from Wellcome (203147/Z/16/Z). The
funding sources had had no involvement in study design, the collection, analysis or
interpretation of data, in the writing of the report or in the decision to submit the article for
publication.
1038
1039
CRediT AUTHORSHIP CONTRIBUTION STATEMENT
Ian A. Clark: Conceptualization, Methodology, Investigation, Formal analysis, Writing -
original draft, Writing - review & editing. **Anna M. Monk:** Investigation, Writing - review &
editing. **Victoria Hotchin:** Investigation, Writing - review & editing. **Gloria Pizzamiglio:**
Investigation, Writing - review & editing. **Alice Liefgreen:** Investigation, Writing - review &
editing. **Martina F. Callaghan:** Formal analysis, Writing - review & editing. **Eleanor A.**
Maguire: Conceptualization, Methodology, Formal analysis, Writing - original draft, Writing
- review & editing

1048

1049

1050 **REFERENCES**

1051

Addis, D. R., Wong, A. T., & Schacter, D. L. (2007). Remembering the past and imagining
the future: Common and distinct neural substrates during event construction and
elaboration. *Neuropsychologia, 45*(7), 1363-1377. doi:
http://dx.doi.org/10.1016/j.neuropsychologia.2006.10.016
Andelman, F., Hoofien, D., Goldberg, I., Aizenstein, O., & Neufeld, M. Y. (2010). Bilateral
hippocampal lesion and a selective impairment of the ability for mental time travel.
Neurocase, 16(5), 426-435. doi: http://dx.doi.org/10.1080/13554791003623318
Andrade, J., May, J., Deeprose, C., Baugh, S.-J., & Ganis, G. (2014). Assessing vividness of
mental imagery: The Plymouth Sensory Imagery Questionnaire. *British Journal of
Psychology, 105*(4), 547-563. doi: http://dx.doi.org/10.1111/bjop.12050
Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *Neuroimage, 38*(1),
95-113. doi: http://dx.doi.org/10.1016/j.neuroimage.2007.07.007

Ashburner, J., & Friston, K. J. (2000). Voxel based morphometry - The methods. *Neuroimage, 11*(6), 805-821. doi: http://dx.doi.org/10.1006/nimg.2000.0582

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *Neuroimage, 26*(3), 839-851. doi: http://dx.doi.org/10.1016/j.neuroimage.2005.02.018

Barry, D. N., Barnes, G. R., Clark, I. A., & Maguire, E. A. (2018). The neural dynamics of novel scene imagery. *Journal of Neuroscience, 32*(22), 4375-4386. doi: http://dx.doi.org/10.1523/JNEUROSCI.2497-18.2019

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society B, 57*, 289-300. doi: http://dx.doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bohbot, V. D., Lerch, J., Thorndycraft, B., Iaria, G., & Zijdenbos, A. P. (2007). Gray matter differences correlate with spontaneous strategies in a human virtual navigation task. *Journal of Neuroscience, 27*(38), 10078-10083. doi: http://dx.doi.org/10.1523/jneurosci.1763-07.2007

Brown, T. I., Ross, R. S., Keller, J. B., Hasselmo, M. E., & Stern, C. E. (2010). Which way was I going? Contextual retrieval supports the disambiguation of well learned overlapping navigational routes. *Journal of Neuroscience, 30*(21), 7414. doi: http://dx.doi.org/10.1523/JNEUROSCI.6021-09.2010

Brown, T. I., Whiteman, A. S., Aselcioglu, I., & Stern, C. E. (2014). Structural differences in hippocampal and prefrontal gray matter volume support flexible context-dependent navigation ability. *Journal of Neuroscience, 34*(6), 2314-2320. doi: http://dx.doi.org/10.1523/jneurosci.2202-13.2014

Brunec, I. K., Robin, J., Patai, E. Z., Ozubko, J. D., Javadi, A.-H., Barense, M. D., . . . Moscovitch, M. (2019). Cognitive mapping style relates to posterior–anterior hippocampal volume ratio. *Hippocampus, 29*(8), 748-754. doi: http://dx.doi.org/10.1002/hipo.23072

Buckner, R. L., & Carroll, D. C. (2007). Self-projection and the brain. *Trends in Cognitive Sciences, 11*(2), 49-57. doi: http://dx.doi.org/10.1016/j.tics.2006.11.004

Callaghan, M. F., Freund, P., Draganski, B., Anderson, E., Cappelletti, M., Chowdhury, R., . . . Weiskopf, N. (2014). Widespread age-related differences in the human brain microstructure revealed by quantitative magnetic resonance imaging. *Neurobiology of Aging, 35*(8), 1862-1872. doi: http://dx.doi.org/10.1016/j.neurobiolaging.2014.02.008

Callaghan, M. F., Josephs, O., Herbst, M., Zaitsev, M., Todd, N., & Weiskopf, N. (2015). An evaluation of prospective motion correction (PMC) for high resolution quantitative MRI. *Frontiers in Neuroscience, 9*(97). doi: http://dx.doi.org/10.3389/fnins.2015.00097

Chadwick, M. J., Bonnici, H. M., & Maguire, E. A. (2014). CA3 size predicts the precision of memory recall. *Proceedings of the National Academy of Sciences USA, 111*(29), 10720-10725. doi: http://dx.doi.org/10.1073/pnas.1319641111

Chrastil, E. R., Sherrill, K. R., Aselcioglu, I., Hasselmo, M. E., & Stern, C. E. (2017). Individual differences in human path integration abilities correlate with gray matter volume in retrosplenial cortex, hippocampus, and medial prefrontal cortex. *eneuro, 4*(2). doi: http://dx.doi.org/10.1523/eneuro.0346-16.2017

Ciaramelli, E., Faggi, G., Scarpazza, C., Mattioli, F., Spaniol, J., Ghetti, S., & Moscovitch, M. (2017). Subjective recollection independent from multifeatural context retrieval following damage to the posterior parietal cortex. *Cortex, 91*, 114-125. doi: http://dx.doi.org/10.1016/j.cortex.2017.03.015

Clark, I. A., Hotchin, V., Monk, A., Pizzamiglio, G., Liefgreen, A., & Maguire, E. A. (2019). Identifying the cognitive processes underpinning hippocampal-dependent tasks.

Journal of Experimental Psychology: General, 148(11), 1861-1881. doi: http://dx.doi.org/10.1037/xge0000582

Clark, I. A., & Maguire, E. A. (2020). Do questionnaires reflect their purported cognitive functions? *Cognition, 195*, 104114. doi: http://dx.doi.org/10.1016/j.cognition.2019.104114

Dalton, M. A., Zeidman, P., Barry, D. N., Williams, E., & Maguire, E. A. (2017). Segmenting subregions of the human hippocampus on structural magnetic resonance image scans: An illustrated tutorial. *Brain and Neuroscience Advances, 1*, 1-36. doi: http://dx.doi.org/10.1177/2398212817701448

Dalton, M. A., Zeidman, P., McCormick, C., & Maguire, E. A. (2018). Differentiable processing of objects, associations and scenes within the hippocampus. *Journal of Neuroscience, 38*(38), 8146-8159. doi: http://dx.doi.org/10.1523/JNEUROSCI.0263-18.2018

Dijkstra, N., Bosch, S. E., & van Gerven, M. A. J. (2017). Vividness of visual imagery depends on the neural overlap with perception in visual areas. *Journal of Neuroscience, 37*(5), 1367. doi: http://dx.doi.org/10.1523/JNEUROSCI.3022-16.2016

Franco, A., Malhotra, N., & Simonovits, G. (2014). Publication bias in the social sciences: Unlocking the file drawer. *Science, 345*(6203), 1502-1505. doi: http://dx.doi.org/10.1126/science.1255484

Fulford, J., Milton, F., Salas, D., Smith, A., Simler, A., Winlove, C., & Zeman, A. (2018). The neural correlates of visual imagery vividness – An fMRI study and literature review. *Cortex, 105*, 26-40. doi: http://dx.doi.org/10.1016/j.cortex.2017.09.014

Gilboa, A., Winocur, G., Grady, C. L., Hevenor, S. J., & Moscovitch, M. (2004). Remembering our past: Functional neuroanatomy of recollection of recent and very remote personal events. *Cerebral Cortex, 14*(11), 1214-1225. doi: http://dx.doi.org/10.1093/cercor/bhh082

Hartley, T., & Harlow, R. (2012). An association between human hippocampal volume and topographical memory in healthy young adults. *Frontiers in Human Neuroscience, 6*(338). doi: http://dx.doi.org/10.3389/fnhum.2012.00338

Hartley, T., Maguire, E. A., Spiers, H. J., & Burgess, N. (2003). The well-worn route and the path less traveled: Distinct neural bases of route following and wayfinding in humans. *Neuron, 37*(5), 877-888. doi: http://dx.doi.org/10.1016/S0896-6273(03)00095-3

Hassabis, D., Kumaran, D., & Maguire, E. A. (2007). Using imagination to understand the neural basis of episodic memory. *Journal of Neuroscience, 27*(52), 14365-14374. doi: http://dx.doi.org/10.1523/jneurosci.4549-07.2007

Hassabis, D., Kumaran, D., Vann, S. D., & Maguire, E. A. (2007). Patients with hippocampal amnesia cannot imagine new experiences. *Proceedings of the National Academy of Sciences USA, 104*(5), 1726-1731. doi: http://dx.doi.org/10.1073/pnas.0610561104

He, Q., & Brown, T. I. (2020). Heterogeneous correlations between hippocampus volume and cognitive map accuracy among healthy young adults. *Cortex, 124*, 167-175. doi: http://dx.doi.org/10.1016/j.cortex.2019.11.011

Hegarty, M., Richardson, A. E., Montello, D. R., Lovelace, K., & Subbiah, I. (2002). Development of a self-report measure of environmental spatial ability. *Intelligence, 30*(5), 425-447. doi: http://dx.doi.org/10.1016/S0160-2896(02)00116-2

Helms, G., Dathe, H., & Dechent, P. (2008). Quantitative FLASH MRI at 3T using a rational approximation of the Ernst equation. *Magnetic Resonance in Medicine, 59*(3), 667-672. doi: http://dx.doi.org/10.1002/mrm.21542

Hodgetts, C. J., Postans, M., Warne, N., Varnava, A., Lawrence, A. D., & Graham, K. S. (2017). Distinct contributions of the fornix and inferior longitudinal fasciculus to

episodic and semantic autobiographical memory. *Cortex, 94*, 1-14. doi: http://dx.doi.org/10.1016/j.cortex.2017.05.010

Hodgetts, C. J., Stefani, M., Williams, A. N., Kolarik, B. S., Yonelinas, A. P., Ekstrom, A. D., . . . Graham, K. S. (2020). The role of the fornix in human navigational learning. *Cortex, 124*, 97-110. doi: http://dx.doi.org/10.1016/j.cortex.2019.10.017

Hurley, N. C., Maguire, E. A., & Vargha-Khadem, F. (2011). Patient HC with developmental amnesia can construct future scenarios. *Neuropsychologia, 49*(13), 3620-3628. doi: http://dx.doi.org/10.1016/j.neuropsychologia.2011.09.015

Iaria, G., Lanyon, L. J., Fox, C. J., Giaschi, D., & Barton, J. J. S. (2008). Navigational skills correlate with hippocampal fractional anisotropy in humans. *Hippocampus, 18*(4), 335-339. doi: http://dx.doi.org/10.1002/hipo.20400

Johnson, P. O., & Fay, L. C. (1950). The Johnson-Neyman technique, its theory and application. *Psychometrika, 15*(4), 349-367. doi: http://dx.doi.org/10.1007/BF02288864

Keogh, R., Bergmann, J., & Pearson, J. (2020). Cortical excitability controls the strength of mental imagery. *eLife, 9*, e50232. doi: http://dx.doi.org/10.7554/eLife.50232

Kharabian Masouleh, S., Eickhoff, S. B., Hoffstaedter, F., Genon, S., & Alzheimer's Disease Neuroimaging Initiative. (2019). Empirical examination of the replicability of associations between brain structure and psychological variables. *eLife, 8*, e43464. doi: http://dx.doi.org/10.7554/eLife.43464

Kong, X.-Z., & Francks, C. (2019). An illustration of reproducibility in neuroscience research in the absence of selective reporting. *bioRxiv*, 866301. doi: http://dx.doi.org/10.1101/866301

LePort, A. K. R., Mattfeld, A. T., Dickinson-Anson, H., Fallon, J. H., Stark, C. E. L., Kruggel, F., . . . McGaugh, J. L. (2012). Behavioral and neuroanatomical investigation of Highly Superior Autobiographical Memory (HSAM). *Neurobiology of Learning and Memory, 98*(1), 78-92. doi: http://dx.doi.org/10.1016/j.nlm.2012.05.002

Levine, B., Svoboda, E., Hay, J. F., Winocur, G., & Moscovitch, M. (2002). Aging and autobiographical memory: Dissociating episodic from semantic retrieval. *Psychology and Aging, 17*(4), 677-689. doi: http://dx.doi.org/10.1037/0882-7974.17.4.677

Long J. A. (2019). *Interactions: Comprehensive, user-friendly toolkit for probing interactions*. R package version 1.1.0, http://cran.r-project.org/package=interactions.

Lutti, A., Hutton, C., Finsterbusch, J., Helms, G., & Weiskopf, N. (2010). Optimization and validation of methods for mapping of the radiofrequency transmit field at 3T. *Magnetic Resonance in Medicine, 64*(1), 229-238. doi: http://dx.doi.org/doi:10.1002/mrm.22421

Lutti, A., Stadler, J., Josephs, O., Windischberger, C., Speck, O., Bernarding, J., . . . Weiskopf, N. (2012). Robust and fast whole brain mapping of the RF transmit field B1 at 7T. *PloS One, 7*(3), e32379. doi: http://dx.doi.org/10.1371/journal.pone.0032379

Maguire, E. A., Gadian, D. G., Johnsrude, I. S., Good, C. D., Ashburner, J., Frackowiak, R. S. J., & Frith, C. D. (2000). Navigation-related structural change in the hippocampi of taxi drivers. *Proceedings of the National Academy of Sciences USA, 97*(8), 4398-4403. doi: http://dx.doi.org/10.1073/pnas.070039597

Maguire, E. A., Nannery, R., & Spiers, H. J. (2006). Navigation around London by a taxi driver with bilateral hippocampal lesions. *Brain, 129*(11), 2894-2907. doi: http://dx.doi.org/10.1093/brain/awl286

Maguire, E. A., Spiers, H. J., Good, C. D., Hartley, T., Frackowiak, R. S. J., & Burgess, N. (2003). Navigation expertise and the human hippocampus: A structural brain imaging analysis. *Hippocampus, 13*(2), 250-259. doi: http://dx.doi.org/10.1002/hipo.10087

Maguire, E. A., Valentine, E. R., Wilding, J. M., & Kapur, N. (2003). Routes to remembering: the brains behind superior memory. *Nature Neuroscience, 6*, 90-95. doi: http://dx.doi.org/10.1038/nn988

Maguire, E. A., Woollett, K., & Spiers, H. J. (2006). London taxi drivers and bus drivers: A structural MRI and neuropsychological analysis. *Hippocampus, 16*(12), 1091-1101. doi: http://dx.doi.org/10.1002/hipo.20233

Mazzoni, G., Clark, A., De Bartolo, A., Guerrini, C., Nahouli, Z., Duzzi, D., . . . Venneri, A. (2019). Brain activation in highly superior autobiographical memory: The role of the precuneus in the autobiographical memory retrieval network. *Cortex, 120*, 588-602. doi: http://dx.doi.org/10.1016/j.cortex.2019.02.020

McDonald, J. H. (2014). Handbook of Biological Statistics (3rd ed., pp. 254-260). Baltimore, Maryland: Sparky House Publishing. Retrieved from http://www.biostathandbook.com/multiplecomparisons.html.

Mechelli, A., Price, C. J., Friston, K. J., & Ashburner, J. (2005). Voxel based morphometry of the human brain: Methods and applications. *Current Medical Imaging Reviews, 1*(2), 105-113. doi: http://dx.doi.org/10.2174/1573405054038726

Metzler-Baddeley, C., Jones, D. K., Belaroussi, B., Aggleton, J. P., & O'Sullivan, M. J. (2011). Frontotemporal connections in episodic memory and aging: A diffusion MRI tractography study. *Journal of Neuroscience, 31*(37), 13236-13245. doi: http://dx.doi.org/10.1523/jneurosci.2317-11.2011

Nichols, T. E., Das, S., Eickhoff, S. B., Evans, A. C., Glatard, T., Hanke, M., . . . Yeo, B. T. T. (2017). Best practices in data analysis and sharing in neuroimaging using MRI. *Nature Neuroscience, 20*(3), 299-303. doi: http://dx.doi.org/10.1038/nn.4500

Ohnishi, T., Matsuda, H., Hirakata, M., & Ugawa, Y. (2006). Navigation ability dependent neural activation in the human brain: An fMRI study. *Neuroscience Research, 55*(4), 361-369. doi: http://dx.doi.org/10.1016/j.neures.2006.04.009

Palombo, D. J., Bacopulos, A., Amaral, R. S. C., Olsen, R. K., Todd, R. M., Anderson, A. K., & Levine, B. (2018). Episodic autobiographical memory is associated with variation in the size of hippocampal subregions. *Hippocampus, 28*(2), 69-75. doi: http://dx.doi.org/10.1002/hipo.22818

Poldrack, R. A., Fletcher, P. C., Henson, R. N., Worsley, K. J., Brett, M., & Nichols, T. E. (2008). Guidelines for reporting an fMRI study. *Neuroimage, 40*(2), 409-414. doi: http://dx.doi.org/10.1016/j.neuroimage.2007.11.048

Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis specialization of the human hippocampus. *Trends in Cognitive Sciences, 17*(5), 230-240. doi: http://dx.doi.org/10.1016/j.tics.2013.03.005

Poppenk, J., & Moscovitch, M. (2011). A hippocampal marker of recollection memory ability among healthy young adults: Contributions of posterior and anterior segments. *Neuron, 72*(6), 931-937. doi: http://dx.doi.org/10.1016/j.neuron.2011.10.014

R Core Team. (2017). R: A language and environment for statistical computing. Vienna, Austria R Foundation for Statistical Computing. Retrieved from http://www.R-project.org

Ramanan, S., Alaeddin, S., Goldberg, Z.-l., Strikwerda-Brown, C., Hodges, J. R., & Irish, M. (2018). Exploring the contribution of visual imagery to scene construction – Evidence from Posterior Cortical Atrophy. *Cortex, 106*, 261-274. doi: http://dx.doi.org/10.1016/j.cortex.2018.06.016

Ridgway, G. R., Henley, S. M. D., Rohrer, J. D., Scahill, R. I., Warren, J. D., & Fox, N. C.
(2008). Ten simple rules for reporting voxel-based morphometry studies.
Neuroimage, 40(4), 1429-1435. doi:
http://dx.doi.org/10.1016/j.neuroimage.2008.01.003
Richter, F. R., Cooper, R. A., Bays, P. M., & Simons, J. S. (2016). Distinct neural
mechanisms underlie the success, precision, and vividness of episodic memory. *eLife,*
5, e18260. doi: http://dx.doi.org/10.7554/eLife.18260
Rosenbaum, R. S., Köhler, S., Schacter, D. L., Moscovitch, M., Westmacott, R., Black, S. E.,
. . . Tulving, E. (2005). The case of K.C.: Contributions of a memory-impaired person
to memory theory. *Neuropsychologia, 43*(7), 989-1021. doi:
http://dx.doi.org/10.1016/j.neuropsychologia.2004.10.007
Rosenbaum, R. S., Priselac, S., Köhler, S., Black, S. E., Gao, F., Nadel, L., & Moscovitch,
M. (2000). Remote spatial memory in an amnesic person with extensive bilateral
hippocampal lesions. *Nature Neuroscience, 3*(10), 1044-1048. doi:
http://dx.doi.org/10.1038/79867
Rosenthal, R. (1979). The file drawer problem and tolerance for null results. *Psychological*
Bulletin, 86(3), 638-641. doi: http://dx.doi.org/10.1037/0033-2909.86.3.638
Schacter, D. L., Addis, D. R., Hassabis, D., Martin, V. C., Spreng, R. N., & Szpunar, K. K.
(2012). The future of memory: Remembering, imagining, and the brain. *Neuron,*
76(4), 677-694. doi: http://dx.doi.org/10.1016/j.neuron.2012.11.001
Schinazi, V. R., Nardi, D., Newcombe, N. S., Shipley, T. F., & Epstein, R. A. (2013).
Hippocampal size predicts rapid learning of a cognitive map in humans.
Hippocampus, 23(6), 515-528. doi: http://dx.doi.org/10.1002/hipo.22111
Scoville, W. B., & Milner, B. (1957). Loss of recent memory after bilateral hippocampal
lesions. *Journal of Neurology, Neurosurgery and Psychiatry, 20*, 11-21. doi:
http://dx.doi.org/10.1136/jnnp.20.1.11
Sheldon, S., Farb, N., Palombo, D. J., & Levine, B. (2016). Intrinsic medial temporal lobe
connectivity relates to individual differences in episodic autobiographical
remembering. *Cortex, 74*, 206-216. doi:
http://dx.doi.org/10.1016/j.cortex.2015.11.005
Sherrill, K. R., Chrastil, E. R., Aselcioglu, I., Hasselmo, M. E., & Stern, C. E. (2018).
Structural differences in hippocampal and entorhinal gray matter volume support
individual differences in first person navigational ability. *Neuroscience, 380*, 123-131.
doi: http://dx.doi.org/10.1016/j.neuroscience.2018.04.006
Simons, J. S., Peers, P. V., Mazuz, Y. S., Berryhill, M. E., & Olson, I. R. (2010). Dissociation
between memory accuracy and memory confidence following bilateral parietal
lesions. *Cerebral Cortex, 20*(2), 479-485. doi:
http://dx.doi.org/10.1093/cercor/bhp116
Spiers, H. J., & Maguire, E. A. (2006). Thoughts, behaviour, and brain dynamics during
navigation in the real world. *Neuroimage, 31*(4), 1826-1840. doi:
http://dx.doi.org/10.1016/j.neuroimage.2006.01.037
Stawarczyk, D., & D'Argembeau, A. (2015). Neural correlates of personal goal processing
during episodic future thinking and mind-wandering: An ALE meta-analysis. *Human*
Brain Mapping, 36(8), 2928-2947. doi: http://dx.doi.org/10.1002/hbm.22818
Steiger, T. K., Weiskopf, N., & Bunzeck, N. (2016). Iron level and myelin content in the
ventral striatum predict memory performance in the aging brain. *Journal of*
Neuroscience, 36(12), 3552-3558. doi: http://dx.doi.org/10.1523/jneurosci.3617-
15.2016

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the hippocampal longitudinal axis. *Nature Reviews Neuroscience, 15*(10), 655-669. doi: http://dx.doi.org/10.1038/nrn3785

Sulpizio, V., Boccia, M., Guariglia, C., & Galati, G. (2016). Functional connectivity between posterior hippocampus and retrosplenial complex predicts individual differences in navigational ability. *Hippocampus, 26*(7), 841-847. doi: http://dx.doi.org/10.1002/hipo.22592

Svoboda, E., McKinnon, M. C., & Levine, B. (2006). The functional neuroanatomy of autobiographical memory: A meta-analysis. *Neuropsychologia, 44*(12), 2189-2208. doi: http://doi.org/10.1016/j.neuropsychologia.2006.05.023

Tabelow, K., Balteau, E., Ashburner, J., Callaghan, M. F., Draganski, B., Helms, G., . . . Mohammadi, S. (2019). hMRI – A toolbox for quantitative MRI in neuroscience and clinical research. *Neuroimage, 194*, 191-210. doi: http://dx.doi.org/10.1016/j.neuroimage.2019.01.029

Van Petten, C. (2004). Relationship between hippocampal volume and memory ability in healthy individuals across the lifespan: review and meta-analysis. *Neuropsychologia, 42*(10), 1394-1413. doi: http://dx.doi.org/10.1016/j.neuropsychologia.2004.04.006

Weisberg, S. M., Newcombe, N. S., & Chatterjee, A. (2019). Everyday taxi drivers: Do better navigators have larger hippocampi? *Cortex, 115*, 280-293. doi: http://dx.doi.org/10.1016/j.cortex.2018.12.024

Weiskopf, N., Suckling, J., Williams, G., Correia, M., Inkster, B., Tait, R., . . . Lutti, A. (2013). Quantitative multi-parameter mapping of R1, PD*, MT, and R2* at 3T: a multi-center validation. *Frontiers in Neuroscience, 7*(95). doi: http://dx.doi.org/10.3389/fnins.2013.00095

Wilson, N.-A., Ramanan, S., Roquet, D., Goldberg, Z.-L., Hodges, J. R., Piguet, O., & Irish, M. (2020). Scene construction impairments in frontotemporal dementia: Evidence for a primary hippocampal contribution. *Neuropsychologia, 137*, 107327. doi: http://dx.doi.org/10.1016/j.neuropsychologia.2019.107327

Winocur, G., & Moscovitch, M. (2011). Memory transformation and systems consolidation. *Journal of the International Neuropsychological Society, 17*(5), 766-780. doi: http://dx.doi.org/10.1017/s1355617711000683

Woollett, K., Glensman, J., & Maguire, E. A. (2008). Non-spatial expertise and hippocampal gray matter volume in humans. *Hippocampus, 18*(10), 981-984. doi: http://dx.doi.org/10.1002/hipo.20465

Woollett, K., & Maguire, E.A. (2009). Navigational expertise may compromise anterograde associative memory. *Neuropsychologia, 47*, 1088-1095.

Woollett, K., Spiers, H.J., & Maguire, E.A. (2009). Talent in the taxi: a model system for exploring expertise. *Philosophical Transactions of the Royal Society, London: Series B, 364*, 1407-1416.

Woollett, K., & Maguire, E. A. (2010). The effect of navigational expertise on wayfinding in new environments. *Journal of Environmental Psychology, 30*(4), 565-573. doi: http://dx.doi.org/10.1016/j.jenvp.2010.03.003

Woollett, K., & Maguire, E. A. (2011). Acquiring "the Knowledge" of London's layout drives structural brain changes. *Current Biology, 21*(24), 2109-2114. doi: http://dx.doi.org/10.1016/j.cub.2011.11.018

Woollett, K., & Maguire, E.A. (2012). Exploring anterograde associative memory in London taxi drivers. *Neuroreport, 23*, 885-888.

Yang, Y., Chen, Z., Zhang, R., Xu, T., & Feng, T. (2020). Neural substrates underlying episodic future thinking: A voxel-based morphometry study. *Neuropsychologia, 138*, 107255. doi: http://dx.doi.org/10.1016/j.neuropsychologia.2019.107255

1357   Zeidman, P., & Maguire, E. A. (2016). Anterior hippocampus: the anatomy of perception,
1358   imagination and episodic memory. *Nature Reviews Neuroscience, 17*(3), 173-182.
1359   doi: http://dx.doi.org/10.1038/nrn.2015.24
1360

1361
1362

1363

1364

Fig. 1. Significant results from the VBM analyses. For all panels the results are shown on the averaged brain of the whole sample (n = 217) and the coloured bar indicates the t value associated with each voxel. (a) Negative association between the Autobiographical Interview (AI) internal place sub-measure and grey matter volume limited to the bilateral hippocampus mask - 10 hippocampal voxels were identified, significant (p < 0.05) only when correcting for the posterior hippocampal mask. (b) Negative association between the AI internal place sub-measure and grey matter volume at p < 0.001 uncorrected for the whole brain. (c) Positive association between navigation route knowledge and grey matter volume limited to the bilateral hippocampus mask; 17 hippocampal voxels were identified, significant (p < 0.05) only when correcting for the anterior hippocampal mask. (d) Positive association between navigation route knowledge and grey matter volume at p < 0.001 uncorrected for the whole brain, showing that the hippocampal voxels are in fact located on the edge of a cluster centred on the right anterior fusiform and parahippocampal cortices.

1378

Supplementary Methods

Table S1. Double scoring of the scene construction task.

	Rating				
	Spatial References	Entities Present	Sensory Descriptions	Thoughts/ Emotions/Actions	Quality Ratings
For each individual scene					
n = 308	0.90	0.96	0.94	0.90	0.90
For each individual participant (i.e. score is averaged across the seven scenes)					
n = 44	0.91	0.99	0.97	0.91	0.93

Note. Inter-class correlation coefficients from a two way random effect model looking for absolute agreement for each content score and for the quality ratings. Four experimenters scored the whole data set (n = 217 participants, 1519 individual scenes) with double scoring performed on 20% of the data (n = 44 participants, 308 scenes) proportionally for each original experimenter.

Table S2. Double scoring of the autobiographical interview.

	Rating						
	Internal Event	Internal Place	Internal Time	Internal Perceptual	Internal Emotion	Internal Sum	External Sum
For each individual memory							
n = 215	0.92	0.85	0.94	0.92	0.86	0.94	0.84
For each individual participant (i.e. score is averaged across the five memories)							
n = 43	0.95	0.88	0.96	0.94	0.81	0.97	0.87

Note. Inter-class correlation coefficients from a two way random effects model looking for absolute agreement for each score on the autobiographical interview. Three experimenters scored the whole data set (n = 217 participants, 1085 individual memories) and double scoring was performed 20% of the data (n = 43 participants, 215 individual memories) proportionally for each original experimenter.

Table S3. Double scoring of the future thinking task.

	Rating				
	Spatial References	Entities Present	Sensory Descriptions	Thoughts/ Emotions/Actions	Quality Ratings
For each individual scene					
n = 132	0.90	0.94	0.93	0.88	0.90
For each individual participant (i.e. score is averaged across the three future scenes)					
n = 44	0.94	0.95	0.96	0.88	0.92

Note. Inter-class correlation coefficients from a two way random effects model looking for absolute agreement for each content score and for the quality ratings. Four experimenters scored the whole data set (n = 217 participants, 651 individual future scenes) with double scoring performed on 20% of the data (n = 44 participants, 132 future scenes) proportionally for each original experimenter.

Table S4. Double scoring of the navigation sketch maps.

	Rating					
	Road Segments	Road Junctions	Number of Landmarks	Landmark Placement	Map Orientation	Map Categorisation
n = 42	0.95	0.96	0.97	0.96	0.96	0.89

Note. Inter-class correlation coefficients from a two way random effects model looking for absolute agreement for each score on the navigation sketch maps. Three experimenters scored the whole data set (n = 217) and double scoring was performed on 20% of the data (n = 42 participants) proportionally for each original experimenter.

Supplementary Results

Primary VBM analyses: Results outside of the hippocampus

Only one association was identified between cognitive task performance and grey matter volume outside of the hippocampus (when using a statistical threshold of p < 0.05 FWE whole brain corrected). A negative association was observed between 22 voxels in the right parahippocampal cortex and AI vividness ratings (peak coordinates = 24 -23 -28, peak t = 4.78, p $_{\text{FWE whole brain corrected}}$ = 0.025).

Table S5. Partial correlations between task performance and hippocampal grey matter volume with age, gender, total intracranial volume and MRI scanner as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
Scene construction								
Experiential index	0.054	0.89	0.061	0.89	0.037	0.89	-0.021	0.89
Spatial references	0.082	0.89	0.079	0.89	0.068	0.89	0.00	1.00
Entities present	-0.013	0.89	-0.062	0.89	0.034	0.89	0.10	0.89
Sensory descriptions	0.023	0.89	0.069	0.89	-0.024	0.89	-0.095	0.89
Thoughts/emotions/actions	-0.043	0.89	-0.066	0.89	-0.013	0.89	0.037	0.89
Spatial coherence index	0.029	0.89	0.033	0.89	0.018	0.89	-0.021	0.89
Autobiographical interview								
Internal details	-0.054	0.87	-0.004	0.95	-0.089	0.71	-0.11	0.69
External details	-0.013	0.92	-0.067	0.87	0.039	0.87	0.10	0.69
Internal events	-0.069	0.87	-0.014	0.92	-0.11	0.69	-0.11	0.69
Internal time	-0.027	0.87	-0.025	0.87	-0.024	0.87	-0.013	0.92
Internal place	-0.054	0.87	-0.044	0.87	-0.053	0.87	-0.031	0.87
Internal perceptual	-0.005	0.95	0.036	0.87	-0.042	0.87	-0.091	0.71
Internal thoughts/emotions	-0.033	0.87	-0.027	0.87	-0.032	0.87	-0.027	0.87
Vividness rating	-0.13	0.69	-0.095	0.71	-0.14	0.69	-0.072	0.87
Future thinking								
Experiential index	0.029	0.89	0.064	0.89	-0.010	0.89	-0.075	0.89
Spatial references	0.068	0.89	0.11	0.89	0.020	0.89	-0.071	0.89
Entities present	-0.021	0.89	-0.017	0.89	-0.021	0.89	-0.010	0.89
Sensory descriptions	-0.016	0.89	0.026	0.89	-0.051	0.89	-0.083	0.89
Thoughts/emotions/actions	-0.034	0.89	-0.051	0.89	-0.011	0.89	0.025	0.89
Spatial coherence index	-0.025	0.89	-0.011	0.89	-0.031	0.89	-0.037	0.89
Navigation								
Overall navigation score	0.071	0.55	0.12	0.36	0.015	0.91	-0.10	0.36
Movie clip recognition	0.11	0.36	0.057	0.62	0.14	0.36	0.10	0.36
Scene recognition	-0.18	0.91	-0.016	0.91	-0.016	0.91	-0.002	0.98
Proximity judgements	0.082	0.46	0.060	0.61	0.084	0.46	0.030	0.88
Route knowledge	0.11	0.36	0.16	0.36	0.042	0.78	-0.11	0.36
Sketch map	0.060	0.61	0.11	0.36	0.004	0.98	-0.10	0.36

Note. P values are Benjamini-Hochberg false discovery rate corrected at p < 0.05.

Table S6. Partial correlations between task performance and hippocampal grey matter volume in the <u>male</u> participants with age, total intracranial volume and MRI scanner as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
Scene construction								
Experiential index	-0.054	0.83	-0.073	0.83	-0.023	0.89	0.056	0.83
Spatial references	-0.08	0.83	-0.083	0.83	-0.059	0.83	0.025	0.89
Entities present	-0.11	0.78	-0.17	0.59	-0.032	0.89	0.14	0.72
Sensory descriptions	-0.12	0.78	-0.049	0.83	0.16	0.59	-0.13	0.72
Thoughts/emotions/actions	-0.1	0.83	-0.17	0.59	-0.006	0.95	0.17	0.59
Spatial coherence index	0.034	0.89	0.006	0.95	0.053	0.83	0.058	0.83
Autobiographical interview								
Internal details	-0.086	0.79	-0.039	0.84	-0.11	0.79	-0.10	0.79
External details	-0.037	0.84	-0.080	0.79	0.015	0.91	0.090	0.79
Internal events	-0.038	0.84	0.022	0.90	-0.088	0.79	-0.12	0.79
Internal time	-0.12	0.79	-0.12	0.79	-0.096	0.79	0.010	0.92
Internal place	-0.081	0.79	-0.055	0.82	-0.089	0.79	-0.071	0.82
Internal perceptual	-0.11	0.79	-0.053	0.82	-0.14	0.79	-0.12	0.79
Internal thoughts/emotions	-0.024	0.90	-0.063	0.82	0.02	0.90	0.068	0.82
Vividness rating	-0.079	0.79	-0.10	0.79	-0.036	0.84	0.060	0.82
Future thinking								
Experiential index	-0.12	0.52	-0.056	0.68	-0.15	0.52	-0.11	0.52
Spatial references	-0.12	0.52	-0.013	0.90	-0.19	0.52	-0.20	0.52
Entities present	-0.11	0.52	-0.094	0.55	-0.11	0.52	-0.024	0.87
Sensory descriptions	-0.092	0.55	-0.041	0.78	-0.12	0.52	-0.087	0.55
Thoughts/emotions/actions	-0.12	0.52	-0.14	0.52	-0.066	0.67	0.062	0.67
Spatial coherence index	-0.11	0.52	-0.11	0.52	-0.085	0.55	0.021	0.87
Navigation								
Overall navigation score	0.07	0.77	0.13	0.46	-0.005	0.97	-0.14	0.43
Movie clip recognition	0.24	0.09	0.16	0.38	0.27	0.072	0.15	0.41
Scene recognition	0.12	0.48	0.084	0.67	0.12	0.48	0.061	0.81
Proximity judgements	0.012	0.97	0.017	0.97	0.004	0.97	-0.011	0.97
Route knowledge	0.18	0.35	0.27	0.072	0.044	0.88	-0.24	0.090
Sketch map	0.044	0.88	0.10	0.55	-0.025	0.97	-0.14	0.43

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Table S7. Partial correlations between tasks performance and hippocampal grey matter volume in the <u>female</u> participants with age, total intracranial volume and MRI scanner as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
Scene construction								
Experiential index	0.16	0.38	0.21	0.22	0.083	0.77	-0.1	0.77
Spatial references	0.21	0.22	0.23	0.22	0.16	0.38	-0.027	0.87
Entities present	0.058	0.77	0.028	0.87	0.072	0.77	0.064	0.77
Sensory descriptions	0.16	0.38	0.19	0.28	0.096	0.77	-0.066	0.77
Thoughts/emotions/actions	-0.03	0.87	0.007	0.94	-0.055	0.77	-0.076	0.77
Spatial coherence index	0.014	0.93	0.056	0.77	-0.026	0.87	-0.097	0.77
Autobiographical interview								
Internal details	-0.035	0.98	0.029	0.98	-0.082	0.91	-0.13	0.89
External details	0.015	0.98	-0.052	0.94	0.069	0.94	0.12	0.89
Internal events	-0.098	0.91	-0.049	0.94	-0.12	0.89	-0.10	0.91
Internal time	0.049	0.94	0.062	0.94	0.029	0.98	-0.034	0.98
Internal place	-0.003	0.98	-0.002	0.98	-0.004	0.98	-0.004	0.98
Internal perceptual	0.066	0.94	0.11	0.89	0.013	0.98	-0.086	0.91
Internal thoughts/emotions	-0.074	0.94	-0.006	0.98	-0.12	0.89	-0.13	0.89
Vividness rating	-0.18	0.89	-0.086	0.91	-0.22	0.89	-0.18	0.89
Future thinking								
Experiential index	0.23	0.12	0.24	0.12	0.17	0.34	-0.03	0.88
Spatial references	0.25	0.12	0.23	0.12	0.22	0.12	0.051	0.88
Entities present	0.12	0.64	0.11	0.64	0.10	0.64	0.008	0.95
Sensory descriptions	0.081	0.72	0.12	0.64	0.033	0.88	-0.079	0.72
Thoughts/emotions/actions	0.036	0.88	0.037	0.88	0.028	0.88	-0.006	0.95
Spatial coherence index	0.062	0.85	0.11	0.64	0.012	0.95	-0.098	0.64
Navigation								
Overall navigation score	0.044	0.97	0.079	0.97	0.005	0.97	-0.066	0.97
Movie clip recognition	-0.009	0.97	-0.045	0.97	0.024	0.97	0.058	0.97
Scene recognition	-0.12	0.97	-0.10	0.97	-0.12	0.97	-0.043	0.97
Proximity judgements	0.16	0.97	0.12	0.97	0.16	0.97	0.059	0.97
Route knowledge	0.015	0.97	0.015	0.97	0.012	0.97	0.009	0.97
Sketch map	0.050	0.97	0.091	0.97	0.004	0.97	-0.079	0.97

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Table S8. Details of the groups created for each task when dividing by median performance.

Performance variable	Median performance	N (low performance)	N (high performance)
Scene construction			
Experiential index	41.2	109	108
Spatial references	3.29	112	105
Entities present	9.71	112	105
Sensory descriptions	12.14	112	105
Thoughts/emotions/actions	3.14	114	103
Spatial coherence index	3.0	112	105
Autobiographical interview			
Internal details	23.2	110	107
External details	4.6	111	106
Internal events	10.6	111	106
Internal time	1.4	128	89
Internal place	2.2	123	94
Internal perceptual	5.4	109	108
Internal thoughts/emotions	3.2	111	106
Vividness rating	4.6	114	103
Future thinking			
Experiential index	39.8	111	106
Spatial references	2.3	123	94
Entities present	10.0	114	103
Sensory descriptions	8.67	117	100
Thoughts/emotions/actions	5.0	116	101
Spatial coherence index	2.67	115	102
Navigation			
Overall navigation score	144.0	109	108
Scene recognition	30.0	146	71
Proximity judgements	8.0	159	58
Route knowledge	11.0	114	103
Sketch map	81	111	106

Table S9. Comparison of hippocampal grey matter volumes when dividing the sample into two groups determined by their median performance on each cognitive task, with age, gender, total intracranial volume and MRI scanner included as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	F	p	F	p	F	p	F	p
Scene construction								
Experiential index	0.85	0.70	0.48	0.83	0.88	0.70	0.051	0.99
Spatial references	1.09	0.70	0.95	0.70	0.79	0.70	0.002	0.99
Entities present	0.043	0.99	1.06	0.70	0.35	0.83	2.93	0.70
Sensory descriptions	0.88	0.70	0.013	0.99	2.27	0.70	2.45	0.70
Thoughts/emotions/actions	1.30	0.70	1.26	0.70	0.85	0.70	0.020	0.99
Spatial coherence index	0.031	0.99	0.11	0.99	0.00	0.99	0.39	0.83
Autobiographical interview								
Internal details	0.96	0.81	0.007	0.95	2.59	0.70	4.18	0.62
External details	0.53	0.81	3.65	0.62	0.26	0.81	5.81	0.54
Internal events	0.76	0.81	0.005	0.95	2.06	0.78	2.69	0.70
Internal time	0.27	0.81	0.29	0.81	0.16	0.88	0.004	0.95
Internal place	0.13	0.89	0.009	0.95	0.50	0.81	1.17	0.81
Internal perceptual	0.071	0.94	0.042	0.95	0.42	0.81	1.44	0.81
Internal thoughts/emotions	0.63	0.81	0.27	0.81	0.77	0.81	0.58	0.81
Vividness ratings	1.56	0.81	0.69	0.81	1.90	0.78	0.57	0.81
Future thinking								
Experiential index	0.11	0.96	0.17	0.96	0.033	0.96	0.15	0.96
Spatial references	1.50	0.96	2.71	0.96	0.34	0.96	0.58	0.96
Entities present	0.009	0.96	0.055	0.96	0.003	0.96	0.13	0.96
Sensory descriptions	0.062	0.96	1.58	0.96	0.54	0.96	3.97	0.96
Thoughts/emotions/actions	0.17	0.96	0.11	0.96	1.03	0.96	1.87	0.96
Spatial coherence index	0.005	0.96	0.006	0.96	0.039	0.96	0.020	0.96
Navigation								
Overall navigation score	0.11	0.96	0.49	0.96	0.008	0.96	0.74	0.96
Scene recognition	0.098	0.96	0.23	0.96	0.009	0.96	0.19	0.96
Proximity judgements	0.079	0.96	0.046	0.96	0.081	0.96	0.003	0.96
Route knowledge	0.72	0.96	1.76	0.96	0.052	0.96	1.12	0.96
Sketch map	0.11	0.96	0.29	0.96	0.007	0.96	0.21	0.96

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Table S10. Partial correlations between task performance and hippocampal grey matter volume in the <u>low</u> performing participants only (as determined by a median split for each task) with age, gender, total intracranial volume and MRI scanner as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
Scene construction								
Experiential index	0.002	0.98	0.059	0.98	-0.054	0.98	0.10	0.96
Spatial references	0.019	0.98	-0.003	0.98	0.036	0.98	0.052	0.98
Entities present	0.013	0.98	-0.005	0.98	0.027	0.98	0.042	0.98
Sensory descriptions	0.17	0.88	0.18	0.88	0.14	0.96	-0.008	0.98
Thoughts/emotions/actions	-0.057	0.98	-0.11	0.96	0.008	0.98	0.11	0.96
Spatial coherence index	0.053	0.98	0.097	0.96	-0.007	0.98	-0.10	0.96
Autobiographical interview								
Internal details	-0.032	0.91	-0.022	0.96	-0.033	0.91	-0.004	0.97
External details	-0.082	0.90	-0.047	0.90	-0.10	0.90	-0.065	0.90
Internal events	-0.072	0.90	-0.020	0.96	-0.11	0.90	-0.11	0.90
Internal time	0.047	0.90	0.006	0.97	0.073	0.90	0.064	0.90
Internal place	-0.049	0.90	-0.075	0.90	-0.014	0.97	0.042	0.90
Internal perceptual	-0.054	0.90	-0.099	0.90	0.010	0.97	0.13	0.90
Internal thoughts/emotions	0.047	0.90	-0.034	0.91	0.12	0.90	0.16	0.90
Vividness ratings	-0.11	0.90	-0.077	0.90	-0.11	0.90	-0.067	0.90
Future thinking								
Experiential index	0.075	0.99	0.14	0.99	0.001	0.99	-0.11	0.99
Spatial references	0.032	0.99	0.008	0.99	0.047	0.99	0.063	0.99
Entities present	-0.003	0.99	0.019	0.99	-0.023	0.99	-0.039	0.99
Sensory descriptions	0.056	0.99	0.061	0.99	0.041	0.99	-0.010	0.99
Thoughts/emotions/actions	-0.069	0.99	-0.008	0.99	-0.11	0.99	-0.14	0.99
Spatial coherence index	-0.049	0.99	0.040	0.99	-0.12	0.99	-0.19	0.99
Navigation								
Overall navigation score	0.29	0.06	0.24	0.065	0.27	0.06	0.067	0.60
Scene recognition	-0.077	0.55	-0.094	0.49	-0.044	0.67	0.055	0.60
Proximity judgements	0.082	0.52	0.054	0.60	0.094	0.48	0.057	0.60
Route knowledge	0.18	0.19	0.16	0.24	0.15	0.24	0.032	0.78
Sketch map	0.25	0.065	0.22	0.084	0.21	0.090	0.020	0.84

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Table S11. Partial correlations between performance and hippocampal grey matter volume in the <u>high</u> performing participants only (as determined by a median split for each task) with age, gender, total intracranial volume and MRI scanner as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
Scene construction								
Experiential index	-0.019	0.99	-0.001	0.99	-0.032	0.99	-0.033	0.99
Spatial references	0.057	0.99	0.067	0.99	0.036	0.99	-0.027	0.99
Entities present	-0.017	0.99	-0.021	0.99	-0.009	0.99	0.012	0.99
Sensory descriptions	0.078	0.99	0.066	0.99	0.071	0.99	-0.001	0.99
Thoughts/emotions/actions	0.096	0.99	0.062	0.99	0.11	0.99	0.072	0.99
Spatial coherence index	0.014	0.99	-0.072	0.99	0.080	0.99	0.18	0.99
Autobiographical interview								
Internal details	0.0	1.0	0.004	1.0	-0.003	1.0	-0.003	1.0
External details	0.11	1.0	0.12	1.0	0.076	1.0	-0.023	1.0
Internal events	-0.016	1.0	-0.027	1.0	-0.001	1.0	0.029	1.0
Internal time	-0.067	1.0	0.004	1.0	-0.12	1.0	-0.16	1.0
Internal place	-0.087	1.0	-0.13	1.0	-0.037	1.0	0.10	1.0
Internal perceptual	0.047	1.0	0.12	1.0	-0.018	1.0	-0.13	1.0
Internal thoughts/emotions	-0.040	1.0	0.005	1.0	-0.072	1.0	-0.11	1.0
Vividness ratings	-0.093	1.0	-0.087	1.0	-0.079	1.0	-0.013	1.0
Future thinking								
Experiential index	-0.027	0.91	0.017	0.91	-0.061	0.91	-0.082	0.91
Spatial references	-0.012	0.91	0.035	0.91	-0.054	0.91	-0.11	0.91
Entities present	-0.012	0.91	-0.064	0.91	0.040	0.91	0.10	0.91
Sensory descriptions	-0.11	0.91	-0.16	0.91	-0.047	0.91	0.097	0.91
Thoughts/emotions/actions	-0.099	0.91	-0.089	0.91	-0.088	0.91	-0.022	0.91
Spatial coherence index	-0.066	0.91	-0.073	0.91	-0.046	0.91	0.023	0.91
Navigation								
Overall navigation score	-0.071	0.69	0.039	0.74	-0.15	0.33	-0.22	0.24
Scene recognition	0.11	0.63	0.16	0.40	0.038	0.76	-0.14	0.47
Proximity judgements	0.24	0.33	0.21	0.33	0.20	0.33	0.06	0.73
Route knowledge	0.044	0.73	0.15	0.33	-0.06	0.73	-0.21	0.26
Sketch map	-0.082	0.63	0.048	0.73	-0.18	0.33	-0.27	0.12

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Table S12. Details of the groups created for each task when taking only the best and worst performers.

Performance variable	Worst performers maximum	Best performers minimum	N (worst performers)	N (best performers)
Scene construction				
Experiential index	32.66	48.49	20	20
Spatial references	1.57	5.57	21	22
Entities present	6.29	13.86	20	20
Sensory descriptions	7.71	17.0	21	21
Thoughts/emotions/actions	1.29	6.14	19	18
Spatial coherence index	0.71	5.0	23	25
Autobiographical interview				
Internal details	15.20	34.60	22	21
External details	1.80	10.20	21	21
Internal events	6.20	16.60	21	21
Internal time	2.20	10.0	18	15
Internal place	1.40	3.0	26	32
Internal perceptual	0.40	2.60	20	21
Internal thoughts/emotions	1.20	6.0	19	18
Vividness rating	3.40	5.60	19	20
Future thinking				
Experiential index	30.13	47.33	21	21
Spatial references	0.67	4.33	32	30
Entities present	6.67	14.67	23	22
Sensory descriptions	4.33	13.33	21	21
Thoughts/emotions/actions	2.33	8.33	22	22
Spatial coherence index	0.0	5.0	23	29
Navigation				
Overall navigation score	95.0	191.0	21	21
Scene recognition	27.0	32.0	34	28
Proximity judgements	5.0	10.0	18	13
Route knowledge	6.0	19.0	23	22
Sketch map	39.0	121.0	21	21

Table S13. Comparison of hippocampal grey matter volumes when taking the <u>best and worst</u> performing participants for task, with age, gender, total intracranial volume and MRI scanner included as covariates.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	F	p	F	p	F	p	F	p
Scene construction								
Experiential index	0.092	0.97	0.36	0.97	0.0	1.0	0.39	0.97
Spatial references	0.69	0.97	0.15	0.97	1.24	0.97	1.34	0.97
Entities present	0.0	1.0	0.38	0.97	0.37	0.97	1.94	0.97
Sensory descriptions	0.47	0.97	0.63	0.97	0.27	0.97	0.006	1.0
Thoughts/emotions/actions	0.014	1.0	0.21	0.97	0.49	0.97	1.29	0.97
Spatial coherence index	0.22	0.97	0.34	0.97	0.075	0.97	0.058	0.97
Autobiographical interview								
Internal details	4.09	0.80	1.17	0.88	6.04	0.61	2.44	0.83
External details	0.73	0.88	1.36	0.88	0.089	0.88	0.69	0.88
Internal events	1.56	0.88	0.30	0.88	2.62	0.83	1.67	0.88
Internal time	0.001	0.98	0.053	0.90	0.035	0.91	0.23	0.88
Internal place	0.48	0.88	0.10	0.88	0.85	0.88	0.58	0.88
Internal perceptual	0.18	0.88	0.13	0.88	0.17	0.88	0.092	0.88
Internal thoughts/emotions	0.51	0.88	0.87	0.88	0.17	0.88	0.43	0.88
Vividness ratings	2.89	0.98	1.85	0.88	2.42	0.83	0.097	0.88
Future thinking								
Experiential index	0.17	0.96	0.80	0.96	0.001	0.98	0.85	0.96
Spatial references	0.74	0.96	3.06	0.96	0.001	0.98	1.75	0.96
Entities present	0.20	0.96	0.029	0.98	0.37	0.96	0.44	0.96
Sensory descriptions	0.003	0.98	0.074	0.96	0.12	0.96	0.43	0.96
Thoughts/emotions/actions	0.30	0.96	0.44	0.96	0.12	0.96	0.062	0.96
Spatial coherence index	0.72	0.96	0.24	0.96	1.02	0.96	0.50	0.96
Navigation								
Overall navigation score	0.88	0.47	4.51	0.09	0.071	0.79	8.39	0.06
Scene recognition	0.58	0.56	0.93	0.47	0.21	0.72	0.25	0.72
Proximity judgements	5.38	0.07	2.57	0.22	6.19	0.067	1.84	0.32
Route knowledge	7.34	0.07	6.78	0.065	4.30	0.09	6.15	0.067
Sketch map	0.99	0.47	5.18	0.073	0.098	0.79	10.01	0.06

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Table S14. Partial correlations between task performance and the interaction between hippocampal grey matter volume and self-reported ability (measured by the Plymouth Sensory Imagery Questionnaire for scene construction, autobiographical memory and future thinking and the Santa Barbara Sense of Direction Scale for navigation), with age, gender, total intracranial volume and MRI scanner included as covariates in all analyses, as well as the additional covariates of the relevant questionnaire and hippocampal volume measurement.

Performance variable	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
Scene construction								
Experiential index	0.044	0.94	0.054	0.94	0.029	0.94	-0.023	0.94
Spatial references	-0.029	0.94	-0.01	0.94	-0.047	0.94	-0.022	0.94
Entities present	0.002	0.97	0.009	0.94	-0.017	0.94	-0.038	0.94
Sensory descriptions	-0.076	0.94	-0.057	0.94	-0.078	0.94	-0.008	0.94
Thoughts/emotions/actions	0.055	0.94	0.052	0.94	0.047	0.94	-0.021	0.94
Spatial coherence index	0.015	0.94	0.019	0.94	0.01	0.94	-0.021	0.94
Autobiographical interview								
Internal details	0.014	0.87	0.018	0.87	0.017	0.87	-0.022	0.87
External details	0.084	0.78	0.11	0.50	0.033	0.87	-0.12	0.49
Internal events	0.029	0.87	0.033	0.87	0.031	0.87	-0.032	0.87
Internal time	-0.052	0.87	-0.043	0.87	-0.054	0.87	0.001	0.99
Internal place	0.029	0.87	0.075	0.87	-0.027	0.87	-0.14	0.29
Internal perceptual	-0.034	0.87	-0.015	0.87	-0.042	0.87	-0.028	0.87
Internal thoughts/emotions	0.067	0.87	0.02	0.87	0.11	0.50	0.086	0.78
Vividness ratings	0.22	0.016*	0.18	0.096	0.24	0.013*	0.026	0.87
Future thinking								
Experiential index	-0.018	0.90	0.012	0.91	-0.041	0.82	-0.041	0.82
Spatial references	-0.07	0.82	-0.028	0.82	-0.1	0.82	-0.051	0.82
Entities present	-0.062	0.82	-0.05	0.82	-0.068	0.82	-0.007	0.91
Sensory descriptions	-0.12	0.82	-0.09	0.82	-0.13	0.82	-0.033	0.82
Thoughts/emotions/actions	0.066	0.82	0.076	0.82	0.042	0.82	-0.072	0.82
Spatial coherence index	0.03	0.82	0.045	0.82	0.011	0.91	-0.03	0.82
Navigation								
Overall navigation score	0.049	0.85	0.026	0.85	0.067	0.72	0.035	0.85
Clip recognition	0.084	0.72	0.084	0.72	0.071	0.72	-0.04	0.85
Scene recognition	-0.11	0.72	-0.088	0.72	-0.12	0.72	-0.02	0.88
Proximity judgements	0.1	0.72	0.074	0.72	0.11	0.72	0.047	0.85
Route knowledge	0.002	0.98	-0.012	0.90	0.016	0.89	0.028	0.85
Sketch map	0.056	0.84	0.031	0.85	0.075	0.72	0.037	0.85

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$. * = $p < 0.05$.

Table S15. Details of the principal component analysis on the four main outcome measures.

Cognitive task	Component weighting
Scene construction experiential index	0.90
Future thinking experiential index	0.87
Autobiographical interview internal details	0.66
Navigation	0.49

Note. The factor scores for each participant were extracted using the Anderson-Rubin method. These were included in a VBM analysis with age, gender, scanner and total intracranial volume as covariates, finding no significant relationships with hippocampal volume both at the whole brain level ($p < 0.05$ FWE corrected) and when using the anatomical hippocampal masks ($p < 0.05$ FWE corrected for each mask). Performing partial correlations between the extracted hippocampal grey matter volumes and the factor scores also identified no significant relationships (whole hippocampal volume: $r = 0.032$, $p_{\text{FDR corrected}} = 0.85$; anterior hippocampal volume: $r = 0.073$, $p_{\text{FDR corrected}} = 0.58$; posterior hippocampal volume: $r = -0.012$, $p_{\text{FDR corrected}} = 0.86$; posterior/anterior hippocampal volume ratio: $r = -0.091$, $p_{\text{FDR corrected}} = 0.58$).

Table S16. Proportion of variance explained by each principal component of the sub-measures principal component analysis.

PCA Component	Variance explained
C1. Content scenes and future	19.11
C2. Navigation	11.67
C3. Spatial coherence	9.88
C4. Thoughts/emotions	9.42
C5. Autobiographical memory	9.21
C6. Time	5.73

Table S17. Details of the principal component analysis conducted on all 21 task sub-measures.

Sub-measure	Content scenes and future	Navigation	Spatial coherence	Thoughts/ emotions	Autobiographical memory	Time
Future thinking entities present	0.84					
Future thinking spatial references	0.78					
Future thinking sensory descriptions	0.76					
Scene construction sensory descriptions	0.75					
Scene construction entities present	0.72			0.41		
Scene construction spatial references	0.70					
Navigation route knowledge		0.84				
Navigation sketch map		0.83				
Navigation movie clip recognition		0.64				
Navigation scene recognition		0.59				
Navigation proximity judgements		0.41				
Scene construction spatial coherence index			0.89			
Future thinking spatial coherence index			0.87			
Autobiographical memory vividness			0.61			0.38
Scene construction thoughts/emotions/actions				0.84		
Future thinking thoughts/emotions/actions				0.76		
Autobiographical memory internal thoughts/emotions				0.49	0.57	
Autobiographical memory internal events					0.79	
Autobiographical memory internal perceptual					0.65	
Autobiographical memory internal place					0.57	0.35
Autobiographical memory internal time						0.73

Note. Task order is for display purposes only. Only values over 0.3 are reported for ease of viewing.

Table S18. Partial correlations between the sub-measures principal components and hippocampal grey matter volume with age, gender, total intracranial volume and MRI scanner as covariates.

PCA component	Whole hippocampus		Anterior hippocampus		Posterior hippocampus		Posterior/Anterior hippocampus ratio	
	r	p	r	p	r	p	r	p
C1. Content scenes and future	0.041	0.78	0.048	0.78	0.027	0.88	-0.009	0.97
C2. Navigation	0.12	0.65	0.13	0.65	0.085	0.66	-0.029	0.88
C3. Spatial coherence	-0.044	0.78	-0.021	0.91	-0.056	0.77	-0.055	0.77
C4. Thoughts/emotions	-0.052	0.77	-0.084	0.66	-0.011	0.97	0.060	0.77
C5. Autobiographical memory	-0.052	0.77	0.002	0.98	-0.091	0.66	-0.12	0.65
C6. Time	-0.096	0.66	-0.097	0.66	-0.075	0.75	0.006	0.97

Note. P values are Benjamini-Hochberg false discovery rate corrected at $p < 0.05$.

Figure S1. Scree plot of the principal component analysis performed on the sub-measures.