**Epistatic interactions between *PHOTOPERIOD-1, CONSTANS 1* and *CONSTANS 2***
**modulate the photoperiodic response in wheat**

Lindsay M. Shaw[1, 2]*, Chengxia Li[1, 3]*, Daniel P. Woods[1, 3]*, Maria Alejandra Alvarez[1]*,

Huiqiong Lin[1, 3], Mei Yee Lau[1], Andrew Chen[1, 4], and Jorge Dubcovsky[1, 3#]

*The first four authors contributed equally to this work

[1] Dept. Plant Sciences, University of California, Davis, CA 95616. U.S.A.

[2] Currently at Queensland Alliance for Agriculture and Food Innovation, The University of

Queensland, Brisbane, QLD 4072, Australia

[3] Howard Hughes Medical Institute, Chevy Chase, MD 20815, U.S.A.

[4] Currently at School of Agriculture and Food Science, The University of Queensland, St Lucia,

QLD, 4067, Australia.

Daniel P. Woods: ORCID 0000-0002-1498-5707
Jorge Dubcovsky: ORCID 0000-0002-7571-4345
17
18

20  **Short title:** Regulation of wheat heading time by photoperiod

22  **# Corresponding authors:** Jorge Dubcovsky, Email: jdubcovsky@ucdavis.edu, Phone: 530
23  (752)-5159.

**Abstract**

In Arabidopsis, *CONSTANS* (*CO*) integrates light and circadian clock signals to promote flowering under long days (LD). In the grasses, a duplication generated two paralogs designated as *CONSTANS 1* (*CO1*) and *CONSTANS 2* (*CO2*). Here we show that in tetraploid wheat plants grown under LD, combined loss-of-function mutations in the A and B-genome homeologs of *CO1* and *CO2* (*co1 co2*) result in a small (3 d) but significant ($P<0.0001$) acceleration of heading time both in *PHOTOPERIOD1* (*PPD1*) sensitive (*Ppd-A1b*) and insensitive (*Ppd-A1a*) backgrounds. Under short days (SD), *co1 co2* mutants headed 13 d earlier than the wild type ($P<0.0001$) in the presence of *Ppd-A1a*. However, in the presence of *Ppd-A1b*, spikes from both genotypes failed to emerge by 180 d. These results indicate that *CO1* and *CO2* operate mainly as weak heading time repressors in both LD and SD. By contrast, in the absence of functional *PPD1* alleles (*ppd1*), *Co1* accelerated heading time >60 d relative to *co1* under LD. We detected significant genetic interactions among *CO1*, *CO2* and *PPD1* alleles on heading time, which were reflected in complex interactions among these genes at the transcriptional and protein levels. Loss-of-function mutations in *PPD1* had a stronger effect on heading time than combined *co1 co2* mutations and, more importantly, *PPD1* was able to perceive and respond to differences in photoperiod in the absence of functional *CO1* and *CO2* genes. Similarly, *CO1* was able to accelerate heading time in response to LD in the absence of a functional *PPD1*. Taken together, these results indicate that *PPD1* and *CO1* are able to respond to photoperiod in the absence of each other, and that interactions between these two photoperiod pathways at the transcriptional and protein levels are important to fine-tune the flowering response in wheat.

**Author summary**

An understanding of the mechanisms involved in the regulation of wheat heading time is

required to engineer more productive varieties better adapted to new or changing environments.

A large proportion of wheat's natural variation in heading time is determined by differences in

genes controlling the photoperiodic response. In this study, we show that the wheat

*PHOTOPERIOD1* (*PPD1*) gene has a stronger effect on heading time than *CONSTANS1* (*CO1*)

and *CO2* in the regulation of the photoperiodic response, and that complex genetic interactions

among these genes are important to fine-tune heading time. Using loss-of-function mutants for

both *CO1* and *CO2*, we demonstrate that these genes are not required for *PPD1* to perceive

differences in photoperiod and regulate heading time. Similarly, we show that in the absence of

*PPD1*, *CO1* can accelerate heading time more than 60 days in response to long days. Our results

indicate that each of these two wheat photoperiod pathways can respond to differences in

photoperiod even in the absence of the other one. Differences in the relative importance of these

two pathways and in their epistatic interactions have contributed to the diversity of photoperiodic

responses observed in different grass species.

## Introduction

A precise adjustment of flowering time to optimal environmental conditions is critical to maximize plant reproductive success. Plants have evolved complex regulatory mechanisms to adjust flowering initiation in response to seasonal changes and other environmental cues. Many plants, including the temperate cereals, require exposure to extended periods of cold (vernalization) to gain competence to flower [1]. After their vernalization requirement has been fulfilled, many plants still require inductive photoperiods to accelerate flowering. Based on their light requirements for floral induction, plants can be classified into long day (LD), short day (SD) or day-neutral plants [2].

The regulatory pathway that controls flowering in response to photoperiod is best characterized in the model plant *Arabidopsis thaliana*, a long day plant that flowers earlier when the days are longer than a critical threshold. In this species, the external coincidence model explains the integration of light signals with the circadian clock. The clock determines diurnal oscillations of the gene *CONSTANS* (*CO*), which results in the coincidence of the expression peak with the light phase under LD and with the dark phase under SD. Light stabilizes the CO protein, resulting in the induction of *FLOWERING LOCUS T* (*FT*) and the promotion of flowering under LD. Under SD, the degradation of the CO protein in the dark prevents the activation of *FT* [3-5]. Thus, *co* mutants in Arabidopsis flower later than wild type under LD but at the same time as wild type under SD [6].

CO belongs to a family of transcription factors characterized by two zinc finger B-Box domains at the N-terminus and by a CONSTANS, CONSTANS-like, TIMING OF CAB EXPRESSION 1 (CCT) domain at the C-terminus, a structure conserved in Arabidopsis and the grass homologs [7]. The *Oryza sativa* (rice) genome contains a single *CO* gene designated as *Hd1* [8], whereas

two paralogs designated as *CO1* and *CO2* have been identified in *Brachypodium distachyon*,

barley and wheat [9]. In the last three species, the predicted CO1 protein lacks conserved

cysteine residues at the start of the second B-box, suggesting a modified or non-functional B-box

2 in the temperate grasses [7, 9]. Phylogenetic and comparative genomics analyses suggest that

*CO1* and *CO2* paralogs originated by a segmental duplication at the base of the grass family,

with *CO1* located in the ancestral position colinear with rice *Hd1,* whereas *CO2* was lost in rice

and is likely not functional in sorghum [9]

Overexpression of *CO1* and *CO2* under the maize *UBIQUITIN* promoter in spring barley,

promoted flowering under both LD and SD conditions by upregulating *FT1* [10, 11]. However,

both *Ubi::CO1* and *Ubi::CO2* transgenic lines flowered 42-43 d later in SD than in LD

indicating the existence of additional genes controlling the photoperiodic response [10, 11]. In

barley, the response to photoperiod is determined mainly by the pseudo-response regulator gene

*PHOTOPERIOD1* (*PPD1*) [12], also known as *PRR37* in rice [13] and *Sorghum bicolor*

(sorghum) [14]. This gene is regulated by the circadian clock and the phytochrome-mediated

light signaling pathway and encodes a protein with a pseudo-receiver domain and a CCT domain

[14, 15].

In wheat, natural deletions in the promoter region of *PPD-A1* (*Ppd1-A1a*) and *PPD-D1* (*Ppd-*

*D1a*) are associated with increased transcript levels and accelerated heading under SD relative to

the ancestral *Ppd1b* allele [16, 17]. As a result, plants carrying the *Ppd1a* alleles show reduced

differences in heading time between LD and SD and are designated as photoperiod insensitive

(PI), whereas plants carrying the *Ppd1b* allele are designated as photoperiod sensitive (PS) [16,

17]. However, wheat PI lines still exhibit a significant acceleration of flowering under LD

relative to SD and, therefore, the wheat PI designation is used in this study to indicate a reduced

photoperiodic response rather than a lack of a photoperiodic response.

Under LD conditions, the wheat *PPD1* gene induces the expression of *FT1*, which encodes a

mobile protein that travels to the shoot apical meristem (SAM), where it promotes the expression

of the floral homeotic MADS box transcription factor *VERNALIZATION1* (*VRN1*) and the

synthesis of gibberellic acid (GA) [18]. The simultaneous presence of GA and *VRN1* results in

the up-regulation of *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1-1* (*SOC1*) and

*LEAFY* (*LFY*) genes, which are required for normal and timely spike development, stem

elongation and spike emergence or heading [18]. In photoperiod sensitive wheats grown under

SD, the SAM transitions to the reproductive stage, but stem elongation and spike development

proceed very slowly and heads fail to emerge or emerge extremely late [19]. These results

indicate that in wheat, the photoperiod pathway has a larger impact on the duration of spike

development and stem elongation than on the initial transition between the vegetative and

reproductive stages [18, 19]. The addition of gibberellic acid (GA) to PS spring wheat grown

under SD accelerates spike development, but only in lines that express *VRN1* under SD [18].

In addition to the natural PS and PI alleles, induced loss-of-function mutations in all *PPD1*

homeologs (henceforth, *ppd1*) have been developed for both hexaploid [20] and tetraploid wheat

[19]. These *ppd1* plants show large delays in heading time under LD and usually spikes emerge

very late, fail to emerge or are aborted under SD [19, 20]. The flowering phenotypes of the wheat

*ppd1* mutants are more drastic than those described for the barley photoperiod insensitive allele

*ppd-H1*, which has only amino acid changes and is likely a hypomorphic allele rather than a

complete loss-of-function allele [12]. It is important to differentiate the photoperiod insensitive

allele in barley that refers to delayed heading under LD, from the PI allele in wheat that refers to

accelerated flowering under SD.

*CO1* and *CO2* overexpression results in barley suggest a role of these genes as flowering

promoters [10, 11]. However, overexpression under a constitutive promoter results in elevated

transcript levels and ectopic expression in tissues or developmental phases that are different from

those of the natural alleles, which complicates the interpretation of overexpression results. The

use of loss-of-function mutations can provide a better understanding of the role of these genes

under more natural conditions. In this study, we investigate the role of *CO1* and *CO2* on the

photoperiod response in wheat using loss-of-function mutants for both genes in the tetraploid

spring wheat variety Kronos. In addition, we combine loss-of-function mutations in *co1*, *co2* and

*ppd1* to study the effect of their genetic interactions on heading time and on the transcription

profiles of critical flowering genes. We also investigate the interactions among the proteins

encoded by these genes. Taken together, these analyses show complex genetic interactions

among *CO1*, *CO2* and *PPD1* on heading time that reflect similarly complex interactions at the

transcriptional and protein levels.

**Results**

**Identification of *CO1* and *CO2* loss-of-function mutants in wheat**

We identified a total of 153 mutations in the A and B homeologs of *CO1* and *CO2* in the Kronos

tetraploid TILLING population [21, 22] using primers and PCR conditions listed in Supporting

Information Table S1. Among the 153 mutations, 74 resulted in non-synonymous amino acid

changes, and five introduced early stop codons resulting in truncated non-functional proteins

(Table S2). These five truncation mutations were validated by sequencing and one for each

homeolog was selected to study the function of *CO1* and *CO2* (Figure 1). We backcrossed the

individual mutant lines with wild-type Kronos at least twice to reduce background mutations.

We generated a *co1* loss-of-function mutant by crossing *co-A1* mutant T4-395 (Q242*) with *co-*

*B1* mutant T4-1170 (Exon2 acceptor splice site mutation). To generate the *co2* loss-of-function

mutant, we crossed *co-A2* mutant T4-202 (Exon 2 acceptor splice site mutation) with *co-B2*

mutant T4-391 (Q193*). Finally, we crossed the *co1* and *co2* mutants and generated a *co1 co2*

double mutant. The stop codon mutations in *co-A1* and *co-B2* truncate the carboxyl terminal half

of the CONSTANS protein including the CCT domain (Figure 1), which is critical for protein

function in both Arabidopsis and cereals [23]. Likewise, the splice-site mutations selected in the

*co-B1* and *co-A2* mutants result in intron retention and premature stop codons that also eliminate

the conserved CCT domain. Based on these large truncations we concluded that *co1 co2* has

loss-of-function mutations for all *CO1* and *CO2* homeologs.

**Figure 1: *CO1* and *CO2* gene structure and position of the selected TILLING mutations.**

Both *CO1* and *CO2* genes have two exons, which encode two B-BOX domains at the N-terminus

and a CCT domain at the C-terminus. The selected *co-A1* and *co-B2* mutants have a premature

stop codon, whereas the *co-B1* and *co-A2* mutants have acceptor splice-site mutations that result

in intron retention and premature stop codons that eliminate the CCT domain. The T4 numbers

are the mutant identification. The schematic gene structure is based on *CO-A1* from Kronos

(GenBank MT043302).

**Effect of *PPD1* photoperiod sensitive (PS) and insensitive (PI) alleles on heading time and**

**spike development**

We transferred the four *co1 co2* mutations generated in Kronos-PI into a near-isogenic Kronos-

PS line [18] to study the effect of these mutations in two different backgrounds with large

differences in heading time under SD. To establish a baseline for comparison, we first describe

the effect of the two *PPD1* alleles in lines carrying the wild type *Co1* and *Co2* alleles (Figure

2A, blue bars).

Under LD, heading time was 2.5 d earlier in Kronos-PI ($52.3 \pm 0.4$ d) than in Kronos-PS ($54.8 \pm$

0.3 d). Although the difference was small, it was highly significant ($P < 0.0001$) and consistent

with previously published results [15]. Under SD, the differences between the two genotypes

were much larger (>66 d, $P < 0.0001$), with Kronos-PI heading at $113.2 \pm 0.4$ d and Kronos-PS

spikes failing to emerge before the experiment was terminated at 180 d. The earlier flowering of

Kronos-PI under SD reduced the difference between SD and LD in this genotype (61 d) relative

to Kronos-PS (>125 d, Figure 2A blue bars).

**Figure 2: Heading time of wild type, *co1*, *co2* and *co1 co2* mutants under LD and SD in Kronos-PI (*Ppd-A1a*) and Kronos-PS (*Ppd-A1b*) backgrounds**. (**A**) Averages are based on 15-19 plants in LD-PI (2 experiments), 21-24 plants in LD-PS (3 experiments) and 8 to 14 plants in SD-PI and SD-PS (1 experiment). Red numbers indicate differences in heading time relative to the wild type (mut-WT), with *P* values from a Dunnett test. Vertical arrows above the bars indicate that spikes have not emerged at 180 d (end of the experiment). (**B-C**) Interaction graphs for *CO1* and *CO2* in Kronos-PS (*Ppd-A1b*) and Kronos-PI (*Ppd-A1a*) backgrounds. The interactions was significant in B (*P* = 0.0052) but not in C (Table S3). Error bars are standard errors of the means (henceforth, s.e.m). *= *P* < 0.05, **= *P* < 0.01, ***= *P* < 0.001, ns= not significant.

To determine if the observed differences in heading time were the result of changes in the

duration of the vegetative phase or the spike development and elongation phases, we dissected

SAMs of the main tillers from the different genotypes (from 4- to 14-leaf stage, Figure S1).

Under LD, the SAMs from both Kronos-PI and Kronos-PS reached the early stages of spike

development by leaf four (W3-W3.25, Waddington scale [24]), the terminal spikelet stage

(W3.5) by leaf six, and headed at 52-55 d (Figure S1). Under SD, the SAMs of both genotypes

were at the vegetative stage by leaf four and started to elongate by leaf six. After that point, the

two genotypes diverged, while Kronos-PI transitioned to the terminal spikelet and lemma

elongation stage at the eighth leaf, Kronos-PS reached the same developmental stage at leaf 12

(Figure S1).

**Effect of *CO1* and *CO2* on heading time in the presence of functional *PPD1* alleles**

Loss of *CO1* and *CO2* function resulted in small but significant changes in heading time. The

*co1 co2* double mutant headed 3 d earlier than the wild type in both Kronos-PI and Kronos-PS ($P$

$< 0.0001$) under LD, and 13.5 d earlier than the wild type in Kronos-PI under SD ($P < 0.0001$,

Figure 2A). In the Kronos-PS background under SD, spikes failed to emerge or were aborted in

most of the plants by the end of the experiment at 180 d, limiting our ability to determine

difference on heading time between the *co1* and *co2* mutant and wild type alleles under this

condition.

Even though the differences in heading time between *co1 co2* and wild type alleles were larger

under SD (13.5 d) than under LD (3 d), they were still smaller than the differences between the

*PPD1* photoperiod sensitive and insensitive alleles under SD ($>67$ d), or the differences between

LD and SD in Kronos-PI (61 d) or Kronos-PS ($>125$, Figure 2A blue bars). These results

indicate that, in wheat, the *PPD1* alleles have a stronger effect on the photoperiodic response

than the *CO1* and *CO2* genes.

Except for the *co2* mutant that headed later than wild type in the Kronos-PS background under

LD (+1.5 d, *P* < 0.0001), all the other *co1* and *co2* single and double mutants headed

significantly earlier than their respective wild type controls under both LD and SD (Figure 2A).

These results suggest that under the tested conditions, *CO1* functions as a mild heading time

repressor in all backgrounds, and *CO2* as a mild heading time promoter or repressor depending

on the *PPD1* and *CO1* alleles present and the photoperiod (Figure 2A).

*Long day experiments*

To explore the interactions between *PPD1*, *CO1* and *CO2* under LD, we performed a factorial

ANOVA combining data from three experiments in Kronos-PS and two in Kronos-PI, using

experiments as blocks (nested within genotypes). This analysis showed highly significant effects

on heading time for *PPD1* (*P* < 0.0001), *CO1* (*P* < 0.0001) and *CO2* (*P* = 0.0004) and highly

significant two-way interactions for *PPD1* x *CO1* and *PPD1* x *CO2* (*P* < 0.0001, Table S3A).

The *CO1* x *CO2* interaction was not significant (*P* = 0.3437) but the three-way interaction was

significant (*P* = 0.0046, Table S3A), suggesting the possibility of different *CO1* x *CO2*

interactions in the different *PPD1* backgrounds that cancelled each other in the combined

analysis.

To test this possibility, we performed separate *CO1* x *CO2* factorial ANOVAs for Kronos-PS

and Kronos-PI. In the Kronos-PS background, *CO1* showed a stronger effect (*P* < 0.0001) on

heading time than *CO2* (*P* = 0.1448) and their interaction was significant (*P* =0.0052, Figure 2B,

Table S3B). By contrast, in the Kronos-PI background, *CO2* showed a stronger effect (*P* <

0.0001) than *CO1* (*P* = 0.0157) and the interaction was not significant. (*P* = 0.1934, Figure 2C,

Table S3C). In summary, these results indicate complex genetic interactions between *PPD1*,

*CO1* and *CO2* in the regulation of heading time under LD.

*Short day experiments*

Under SD, the Kronos-PS lines failed to flower before the experiment was terminated (180 d), so

we were only able to study the interactions between *CO1* and *CO2* in the Kronos-PI background.

All four genotypes headed between 100 and 113 d, which is 51 to 61 d later than the same lines

under LD. A factorial ANOVA for heading time showed significant effects for both *CO1* and

*CO2* ($P < 0.0001$) and a significant interaction between the two ($P = 0.0029$, Table S4). This

interaction was associated with the larger acceleration of heading time in the double mutant (13.5

d) than in the single *co1* (2.1 d) and *co2* (4.0 d) mutants (Figure 2A). This result indicates a

stronger effect of each mutation in the presence of the mutant allele of the other gene than in the

presence of the wild type allele (Table S4).

To see if the *co1* and *co2* mutations in the Kronos-PS background were associated with delays in

SAM differentiation or in spike development, we dissected apices for three plants from each

genotype every week from 4 to 11 weeks, and included wild type Kronos-PI as a control. Under

SD, all five genotypes reached the double-ridge stage (W2.5) by week 6 (Figure 3A-B). From

there, the Kronos-PI genotype developed more rapidly than the four Kronos-PS genotypes,

which showed similar developmental rates in the wild type and the *co1* and *co2* mutants (Figure

3B). By week 11, the developing spikes in the Kronos-PI control were 3-13 cm long including

awns, the peduncles were 5 to 14 mm long, and the developing flowers were at the elongated

stigmatic branches stage (>W7.5). By contrast, all four Kronos-PS lines at the same time point

showed no peduncle elongation, produced spikes of less than 0.5 cm, and the flowers still had the

stylar canal open (<W6.0). These results indicate that the *co1* and *co2* mutations have a limited

effect on spike development and elongation in the Kronos-PS background under SD compared

with the strong effect of the *Ppd-A1a* in Kronos-PI.

**Figure 3. Effect of *co1* and *co2* mutations on spike development in Kronos-PS under SD.**

Kronos-PI was included as control. (**A**) Dissecting microscope images of developing apices at

week 6 (double-ridge, W2.5) and week 11 with a detail of their floral organs. (**B**) Developmental

time course of SAM using the Waddington scale from 4- through 11-week-old plants (n = 3).

Error bars are standard errors of the means.

**Effect of *CO1* and *CO2* on heading time in the absence of functional *PPD1* alleles**

To investigate the effect of *CO1* and *CO2* in the absence of *PPD1*, we introgressed the *co1* and

*co2* mutations into a Kronos line with loss-of-function mutations in both *PPD-A1* and *PPD-B1*

(*ppd1*) [19]. As before, we describe first the differences in SAM development and heading time

between Kronos-PS and *ppd1* mutants carrying the wild type *Co1* and *Co2* alleles to establish a

baseline for comparison (Figure 4A, blue bars).

Under LD, the plants with the *ppd1 Co1 Co2* allele combination headed much later (115.5 ± 15.3

d) than the Kronos-PS controls carrying the functional *Ppd1b* allele (52.5 ± 0.7 d, Figure 4A),

similar to results reported before [19]. The *ppd1* mutant plants showed a 79% reduction in

number of grains per spike relative to the Kronos-PS lines (Figure 4B-C). Apex dissections

showed a delayed transition to the double-ridge stage in the *ppd1* mutant (leaf 8 to 10) relative to

Kronos-PS (<leaf 4, Figure S1). Similarly, the terminal spikelet was formed in Kronos-PS by

leaf 6 but was not formed in *ppd1* by leaf 14, when the experiment was terminated (Figure S1).

Under SD, spikes of both the *ppd1* mutant and Kronos-PS failed to emerge by 180 d. Dissection

of the main tillers from these plants showed that spike development in *ppd1* lagged behind that

of Kronos-PS (Figure S1, leaf 12 and 14 and Figure S2). These results indicate that *PPD1* is not

essential for the initial transition of the SAM to the reproductive stage, but in its absence, spike

development and stem elongation are severely compromised.

**Figure 4. Phenotypic effects of *co1* and *co2* mutations in the absence of functional *PPD1***

**genes under LD** (**A**) Average days to heading in *ppd1* mutants carrying mutations in either *co1*

or *co2*. Kronos-PS wild type and *co1 co2* double mutants are included as controls. The arrow on

top of the bar indicates that most plants failed to flower before the experiment was terminated at

180 d. (**B**) Spike phenotypes at the end of the experiment (180 d). (**C**) Number of grains per

spike recovered from the same genotypes as in A. Error bars are standard errors of the means

calculated from 7 to 16 plants per genotype for flowering and from 4 to 8 plants for grain

number.

The introgression of the *co2* mutation in the *ppd1* background had a limited effect on heading

time. Heading time of the double mutant *ppd1 co2* (119.1 ± 12.5 d) was only four days later than

*ppd1*, and the differences were not significant suggesting a limited effect of *CO2* on heading

time in the absence of a functional *PPD1*. The *ppd1 co2* double mutant showed an even more

severe reduction in grain production than the *ppd1* mutant (89% reduction relative to Kronos-

PS), but the differences were not significant.

The most important result was observed when we combined the *ppd1* and *co1* mutations. The

spikes from the *ppd1 co1* plants failed to emerge before the experiment was terminated at 180 d

(Figure 4A), and failed to produce any grains (Figure 4B-C). Even though we were not able to

recover the triple *ppd1 co1 co2* mutant, the delayed heading time of the *ppd1 co1* mutant relative

to the *ppd1* and *ppd1 co2* mutants (>60d) is sufficient to conclude that *CO1* can accelerate

heading time in response to LD even in the absence of functional *PPD1* alleles.

**Effect of mutations in *PPD1*, *CO1* and *CO2* on the transcriptional profiles of flowering**

**genes**

To better understand the regulatory networks underlying the genetic interactions described

above, we characterized the effect of loss-of-function mutations in *PPD1, CO1* and *CO2* in a

Kronos-PS background on the transcription profiles of several flowering genes. We grew the

plants in a growth chamber under LD and sampled the fully expanded fourth leaf every 4 hours

throughout a 24-h period.

*CO1* transcripts were significantly upregulated in both *co2* and *ppd1* mutants 12 hours after the

start of the light phase (Zeitgeber time 12 or ZT12, Figure 5A), suggesting that these two genes

act as transcriptional repressors of *CO1*. The negative effect of *PPD1* on *CO1* transcription was

also observed in a separate experiment using older plants (six-week-old). In leaf samples

collected at ZT4 we observed a significant upregulation of *CO1* in the *ppd1* and *ppd1 co2*

mutants relative to Kronos-PS (Figure S3A). No differences were detected between the *ppd1* and

*ppd1 co2* mutants, which suggests that the loss of *co2* function did not affect *CO1* expression in

the absence of functional *PPD1* genes (Figure S3A). Transcript levels of *CO2* were very low and

showed no significant differences among genotypes (Figure 5B). We further confirmed these

observations in six-week old plants of Kronos-PS, *ppd1* and *ppd1 co1* plants sampled at ZT4

(Figure S3B).

**Figure 5.** Effect of loss-of-function mutations in *CO1*, *CO2* and *PPD1* in Kronos-PS on the

transcriptional profiles of wheat flowering genes under LD. (**A**) *CO1*, (**B**) *CO2*, (**C**) *PPD1*, (**D**)

*FT1*, (**E**) *VRN1*, (**F**) *VRN2* in Kronos-PS, *co1*, *co2*, *co1 co2* and *ppd1* mutants. We grew the

plants in a growth chamber under 16 h light at 22 °C and 8 h darkness at 17 °C (gray areas).

Fully expanded 4th leaves (Zadoks scale =14) were harvested at ZT0, ZT4, ZT8, ZT12, ZR16,

and ZT20. The ZT0 value is the same as at ZT24. Shaded boxes represent night. Values are

averages of four biological replicates and bars are SE of the means. Expression values are

relative to *ACTIN* using the delta Ct method. Primers and primer efficiencies are listed in Table

S5 (all amplify both A and B homologs of the respective genes).

The *PPD1* gene in Kronos-PS showed a main peak in the middle of the light period (ZT8) and a

secondary peak at dusk (ZT16), which is consistent with the *PPD1* profile previously described

in the photoperiod sensitive *T. monococcum* [25]. In the *co1* and *co1 co2* mutants, the main peak

was shifted 4 h earlier (ZT4), whereas in the *co2* mutants the main peak was delayed 4 h (ZT12,

Figure 5C). The ZT16 peak was significantly higher than the wild type in the *co1 co2* double

mutant but not in the single mutants (Figure 5C). The complex effects of the *co1* and *co2*

mutations on *PPD1* transcription profiles reflect the significant two- and three-way genetic

interactions among these three genes on heading time described above.

The *FT1* profile was similar to that of *PPD1* in wild type Kronos-PS, but not in the *co1* and *co2*

mutants (Figure 5D). The two profiles were also similar in the upregulation of the secondary

peak at ZT16 in the *co1 co2* double mutant, which correlates with the earlier heading time of this

mutant under LD. The earlier heading time of *co1 co2* also correlated well with the significantly

higher transcript levels of *VRN1* in this mutant (Figure 5E). We observed a significant reduction

of the transcript levels of *FT1* and *VRN1* in the *ppd1* mutant relative to wild type, which is

consistent with its delayed heading time.

The *VRN2* locus includes two genes, but since *ZCCT1* is not functional in tetraploid wheat [26],

we present only expression data for *ZCCT2*. The transcriptional profiles of *PPD1* and *ZCCT2*

were similar for the wild type Kronos-PS but not for the different mutants (Figure 5). We

observed a small but not significant increase of *ZCCT2* transcript levels in the *ppd1* mutant and a

significant downregulation in the *co1* and *co1 co2* mutants. The interpretation of these results is

complicated by the effects of these mutants on *VRN1* which is a known repressor of *VRN2* [27]

(see Discussion section).

**Effect of *VRN1* on the transcriptional regulation of *CO1* and *CO2***

To explore the potential role of *VRN1* in the transcriptional regulation of *CO1* and *CO2*, we used

Kronos *vrn-A1* and *vrn-B1* single mutants and *vrn-A1 vrn-B1* double mutant (henceforth *vrn1*)

developed previously [27]. The *vrn-B1* mutant has a functional allele of *VRN-A1* that confers a

dominant spring growth habit, whereas the *vrn-A1* mutant has the ancestral allele of *VRN-B1* that

confers a recessive winter growth habit. The *vrn1* double mutant has a delayed heading time and

a residual response to vernalization [27].

*CO1* showed significantly higher transcript levels in the late flowering *vrn-A1* and *vrn1* mutants

than in the rapidly flowering *vrn-B1* mutant, both in vernalized and un-vernalized plants (Figure

6 A-B). In all genotypes, *CO1* was downregulated to very low levels in the flag-leaf at heading

time. In the *vrn1* mutant, we observed a slight decrease of *CO1* transcript levels during

vernalization followed by a rapid upregulation when we moved the plants back to room

temperature (RT). By contrast, in the single *vrn-A1* and *vrn-B1* mutants, which still have one

functional copy of *VRN1*, *CO1* transcripts stayed low after the plants were returned to RT

(Figure 6 A). From previous studies, we know that *VRN1* transcript levels increase slowly during

LD vernalization and then rapidly when plants are moved to RT and start to flower [27]. Taken

together, these results suggest that *VRN1* acts as a transcriptional repressor of *CO1*.

**Figure 6.** Effect of mutations in Kronos *vrn-A1* (winter growth habit), *vrn-B1* (spring growth

habit) and *vrn-A1 vrn-B1* (*vrn1* very late heading) on the transcriptional profiles of *CO1* and

*CO2*. All mutants are homozygous and in the Kronos-PI background. (**A-B**) *CO1* and (**C-D**)

*CO2* transcription profiles. **A & C**) More recently expanded leaves from three weeks old plants

were sampled before vernalization (0w), at 3w and 6 w vernalization (4°C), 2 weeks after return

to room temperature (RT= 22°C day and 18°C) and at heading time (HT), which was different

for each genotype. **B & D**) Same genotypes but grown at RT without vernalization. Points are

averages of eight plants and error bars are standard errors of the means. Means with different

letters are significantly different in Tukey test for that time point ($P < 0.05$). Note that the

upregulation of *CO1* after the return of the plants from vernalization occurred only in the *vrn1*

mutant that has no functional copies of *VRN1*.

*CO2* transcripts were downregulated during vernalization and showed a sharp increase when we

moved the plants to RT. However, in contrast to *CO1*, this increase was observed in all three

genotypes (Figure 6C-D). After this increase, *CO2* transcript levels remained high in the flag

leaves during heading and showed no significant differences between the *vrn1* and the spring

*vrn-B1* mutant (Figure 6C-D). These results indicate that *VRN1* has a smaller effect on the

regulation of *CO2* than on the regulation of *CO1*.

**Effect of photoperiod and phytochromes on the transcriptional regulation of *CO1* and *CO2***

The strong upregulation of *CO1* in the *ppd1* mutant indicated that *PPD1* is a transcriptional

repressor of *CO1* (Figure 4A). Since phytochromes *PHYC* [15] and *PHYB* [28] are both required

for the LD transcriptional activation of *PPD1*, we explored the effect of mutations in these two

phytochromes on the expression profiles of *CO1* and *CO2*. To do this, we used recently

published RNAseq studies for leaf samples (4th leaf, ZT 4h) from loss-of-function mutants *phyB*

and *phyC* in Kronos-PI under both LD [28] and SD conditions [29].

A factorial ANOVA for transcripts per million reads (TPM) using photoperiod (LD and SD) and

genotype (WT, *phyC* and *phyB*) as factors, revealed significant differences between SD and LD

for all four genes (*CO-A1*, *CO-B1*, *CO-A2* and *CO-B2*, Figure S4). We also detected difference

between genotypes and genotype x photoperiod interactions for the *CO1* homeologs but not for

the *CO2* homeologs. The significant interaction was associated with the LD downregulation of

*CO1* homeologs in the wild type and their upregulation in the *phyB* and *phyC* mutants (Figure

S4). Both *CO2* homeologs were upregulated by LD and showed no significant differences

between wild type and *phy* mutants. *CO-B2* showed significantly higher transcript levels than

*CO-A2* for all genotypes (Figure S4), whereas for *CO1* the transcript levels and the differences

were larger in *CO-A1* relative to *CO-B1*.

**Protein-protein interactions among CO1, CO2, PPD1, VRN2, PHYC and PHYB**

To test if the interactions among *CO1*, *CO2* and *PPD1* observed at the genetic and

transcriptional levels were associated with interactions at the protein level, we performed yeast-

two-hybrid (Y2H) and split yellow fluorescent protein (YFP) assays among the three proteins. In

a previous study, we reported positive Y2H interactions between the CCT domains of CO1 and

CO2, but not between these two proteins and a truncated version of PPD1 including the CCT

domain [30]. In this study, we used a complete PPD1 protein and detected strong Y2H

interactions between PPD1 and both CO1 and CO2. The Y2H interaction between PPD1 and

CO1 was stronger than between PPD1 and CO2 (Figure 7A).

We further validated the interactions between CO1 and CO2 (Figure 7B) and between CO1 and

PPD1 (Figure 7C-D) using split-YFP assays in wheat protoplasts. We initially failed to detect the

split-YFP interaction between CO1 and PPD1 in Kronos, so we used protoplasts extracted from

an early flowering loss-of-function mutant for *elf3* in Kronos, which has increased levels of

*PPD1* transcripts [25, 31]. We were able to detect the CO1-CO2 and CO1-PPD1 interactions

(but not the CO2-PPD1) by split-YFP in the *elf3* mutant when protoplasts were exposed to light.

For both CO1 and PPD1, we detected the proteins by Western blots only after we exposed the

protoplasts to light but not when we incubated them in the dark after transfection, suggesting that

light is required to stabilize these proteins.

**Figure 7. Protein interactions**. (**A**) Yeast-two-hybrid (Y2H) interactions between PPD1 and

CO1 or CO2. Left panel: SD medium lacking Leucine and Tryptophan (SD-L-W) to select for

yeast transformants containing both bait and prey. Right panel: interaction on SD-L-W-H-A

medium (no L, W, H and A). Dilution factors = 1, 1:10 and 1:100. (**B-D**) Split-YFP in wheat

protoplast. (**B**) Positive nuclear interactions between CO1 and CO2. (**C-D**) Positive nuclear

interactions between CO1 and PPD1 in two different YFP-N/YFP-C configurations. (**E**)

Summary of Y2H interactions among wheat CO1, CO2, PPD1, VRN2, PHYB, PHYC, and ELF3

from Figure S5. Green lines are based on published results from *Brachypodium* [32] and violet

lines on published results from wheat [24]. Thick lines indicate interactions validated by split-

YFP in wheat protoplasts (blue) or *in vitro* coIP (violet). Curved arrows indicate self-

dimerization**.**

Finally, we tested pairwise interactions among CO1, CO2, PPD1, VRN2, PHYC and PHYB

proteins by Y2H assays (Figures 7E and S5). In addition to interacting with CCT-domain

proteins PPD1 (Figure 7A) and VRN2 [30], CO1 and CO2 showed positive interactions with

both PHYC and PHYB (Figure 7E and S5) in the Y2H assays. The interaction between CO1 and

PHYC was stronger than between CO1 and PHYB, whereas the interaction between CO2 and

PHYB was stronger than between CO2 and PHYC. Both phytochromes showed positive

interactions with VRN2 but not with PPD1 (Figures 7E and S5). In addition, we showed that the

full length PPD1 protein did not interact with VRN2 in Y2H (Figure S5). CO2, VRN2, PHYC

and PHYB were able to form homodimers [15, 30] but not CO1 or PPD1 (Figure S5). The

diagrammatic representation of these interactions reveals an extensive and complex network of

protein-protein interactions, with CO1 and CO2 providing a connection between the LD

flowering promoter PPD1 and the LD flowering repressor VRN2 (Figure 7E).

**Discussion**

**Genetic interactions among *CO1*, *CO2*, *PPD1* and *VRN2* modulate photoperiodic responses in grasses**

*Long day grasses*

Natural variation in wheat heading time in response to photoperiod has been mapped mainly to the three *PPD1* homeologs [16, 17, 33]. Similarly, a genome wide association study of 220 spring barley varieties found that most of the variation in heading time (31%) was associated with the *PPD1* locus and variation in the *CO1* locus had a limited contribution (<5%) [34]. These results agree with the limited effect on heading time detected in this study in the Kronos *co1*, *co2*, and *co1 co2* mutants compared with the large differences detected between *PPD1* alleles (Figure 2) and mutants (Figure 4). Even when both *CO1* and *CO2* homeologs were replaced with truncated mutants (*co1 co2*), plants headed only three days earlier than the wild type under LD and 13 days earlier under SD (Figure 2). By contrast, loss of function mutations in *PPD1* delayed heading time approximately two months in LD (Figure 4), and differences between the *Ppd-A1a* and *Ppd-A1b* alleles were even larger under SD (Figure 2).

The effect of the *co1 co2* mutations on the acceleration of heading time was small, but highly significant ($P < 0.0001$) and consistent across experiments under both LD and SD (Figure 2). This was an unexpected result because transgenic barley plants overexpressing *CO1* [10] or *CO2* [11] flowered earlier than the wild type under both LD and SD. These transgenic results had led to the general assumption that *CO1* and *CO2* operate as flowering promoters in the temperate grasses. However, our results show that when *CO1* and *CO2* are expressed under their native promoters in their expected tissues and developmental stages (and in the presence of functional

*PPD1* and *VRN2* alleles) they operate mainly as mild LD and SD flowering repressors rather

than as flowering promoters (except for *CO2* in Kronos-PS under LD, Figure 2).

This inconsistency may be generated by the elevated and ectopic expression levels of *Ubi::CO1*

and *Ubi::CO2* in the transgenic plants, but also by differences in the genetic backgrounds. The

two transgenic barley lines were developed in the variety Golden Promise [10], which carries the

*ppd-H1* allele with reduced sensitivity to LD and does not have any functional copy of *VRN2*. By

contrast, Kronos has functional *PPD1* and *VRN2* genes. Our results show that in a Kronos line

with no functional copies of *PPD1* (*ppd1*), *CO1* accelerates heading time under LD (Figure 4)

similarly to the overexpression of *CO1* in Golden Promise. In addition, the downregulation of

*VRN2* in the Kronos *co1 co2* mutant may have contributed to the earlier heading time of these

mutants (Figure 2), whereas no interactions with *VRN2* are possible in Golden Promise.

The previous results suggest the existence of complex genetic interactions among the

photoperiodic genes, a hypothesis that is also supported by significant digenic interactions in the

factorial ANOVA including *CO1*, *CO2* and *PPD1* (Table S3). The three-way interaction among

the three genes was also significant, which reflects the significant interaction between *CO1* and

*CO2* in Kronos-PS but not in Kronos-PI (Table S3). The contribution of the *PPD1* x *CO2*

interaction to the total sum of squares (9%) was more than two-fold higher than the main *CO2*

effect (4%), which suggests that a large part of *CO2* effect on heading time was mediated by its

modulation of the *PPD1* function (Table S3). Strong genetic interactions between *CO1* and

*PPD1* on heading time have also been reported in a barley biparental population segregating for

*Ubi::CO1* and the *ppd-H1* mutation [10] and in two genome wide association studies (GWAS) in

spring barley [34, 35]. In both GWAS, significant effects of *CO1* on heading time were detected

among the lines carrying the *ppd-H1* allele but not among the lines carrying the wild type *Ppd-*

*H1* allele [34, 35]. These strong genetic interactions indicate that the effect of a particular

photoperiod gene depends strongly on genetic background, a result that has been also reported in

the SD grasses as described below.

***Short day grasses***

In most sorghum genotypes, *SbCO1* operates as a flowering promoter in SD and as a flowering

repressor in LD [36] and *SbPRR37* as a flowering repressor in LD [14, 37]. By contrast, the

orthologous *PPD1* gene in barley and wheat functions as a LD heading promoter [12, 16, 19]

(Figure 4). In spite of the opposite roles, the genetic interactions between *PRR37/PPD1* and *CO1*

showed some similarities. In a sorghum population segregating for *SbCO1* and *SbPRR37*, *SbCO1*

showed a large effect on heading time among the plants carrying the non-functional *Sbprr37*

allele (48% of the variation), but had no effect among the plants carrying the functional *SbPrr37*

allele [38]. This is similar to the GWAS results in spring barley described above [34, 35] and to

the stronger effect of the *co1* mutation in the *ppd1* mutant than in the Kronos plants with

functional *PPD1* alleles described in this study.

Similar to sorghum, in most rice genotypes, *OsPRR37* (=*Hd2*) works as a LD flowering

repressor [13] and *OsCO1* (=*Hd1*) as both a LD flowering repressor and a SD flowering

promoter. Interestingly, transformation of a rice line deficient in *OsCO1* function with a wheat

*CO2* genomic clone accelerated flowering in SD and delayed flowering in LD, complementing

the *OsCO1* function [39]. Since our results indicate that wheat *CO2* functions as a SD flowering

repressor (Figure 2A), the rice result suggests that the function of the wheat CO2 protein was

altered when it was expressed in rice. The opposite change was observed when a late-flowering

Arabidopsis *prr7* mutant was transformed with rice *OsPRR37* driven by the Arabidopsis *PRR7*

promoter. When expressed in Arabidopsis, *OsPRR37* functioned as a LD promoter of flowering

in spite of its normal role as a LD repressor of flowering in rice. Based on these results, it is

tempting to speculate that the different roles of *PRR37/PPD1* and *CO2* in SD and LD grasses

may be the result of differences in the balance of multiple epistatic interactions among the genes

involved in the photoperiodic response, rather than a difference in the specific function of their

encoded proteins.

This hypothesis is also supported by changes in gene function reported in sorghum and rice

plants carrying loss-of-function mutations in both *PRR37* (*prr37*) and *GRAIN NUMBER, PLANT*

*HEIGHT, AND HEADING DATE7* (*ghd7,* a *VRN2* ortholog [40]). In the *prr37 ghd7* sorghum

and rice mutants, *CO1* function changed from a flowering repressor to a flowering promoter in

LD, while maintaining its usual role as a flowering promoter in SD [38, 41, 42]. In addition, in

most rice genotypes carrying the *ghd7* mutation *OsPRR37* function changes from a SD promoter

to a SD repressor of heading, while maintaining its role as a LD flowering repressor [43].

Genetic interactions between *VRN2* (*GHD7*) and both *CO1* (*Hd1*) [44] and *CO2* [11] have been

reported also in barley, suggesting that similar mechanisms may exist in the temperate cereals.

Taken together, these results indicate that genetic interactions between *PPD1/PRR37*,

*VRN2/GHD7* and *CO1/CO2* play a central role in the regulation of heading time in response to

photoperiod in both LD- and SD-grasses and that, therefore, the effect of particular genes may

vary in different genetic backgrounds.

**Complex interactions at the transcriptional level contribute to the observed genetic**

**interactions on heading time**

The significant upregulation of *CO1* in the *ppd1* mutant (Figure 5A and Figure S3A) suggests

that *PPD1* may function as a transcriptional repressor of *CO1*. This hypothesis is supported by

the upregulation of *CO1* in the *phyB* and *phyC* mutants (Figure S4), where *PPD1* transcription is

strongly downregulated [15, 28], and by the negative correlation between *PPD1* and *CO1*

transcript levels in both *ppd1* triple mutants of hexaploid wheat 'Paragon' under LD [20], and

photoperiod sensitive and insensitive isogenic lines of Paragon under SD [45].

However, since *PPD1* induces the expression of *VRN1*, which represses *CO1* transcription

(Figure 6), we cannot rule out the possibility that the effect of *PPD1* on *CO1* transcription is

mediated by *VRN1*. When Kronos *vrn-A1*, *vrn-B1* and *vrn1* mutants were returned to room

temperature after vernalization, *CO1* was rapidly upregulated in the leaves of the *vrn1* mutant

but not in the other three genotypes with functional *VRN1* copies (Figure 6). An identical

expression profile has been described before for *VRN2* in the same mutants [27], indicating that

*VRN1* functions as a transcriptional repressor of both *CO1* and *VRN2*.

*CO2* transcript levels were not significantly affected by *PPD1*, *CO1* (Figure 5B) or *VRN1*

(Figure 6C-D). Similarly, no changes in *CO2* transcription were observed between Paragon-PI

and Paragon-PS lines grown under SD [45], between Kronos-PI and the *phyB* or *phyC* mutants

(Figure S4), or between Kronos-PS and *ppd1* mutants (Figure S3B). However, mutations in *co2*

affected the diurnal transcriptional profiles of all the flowering genes presented in Figure 5. *CO1*

transcript levels were more than 2-fold higher in the *co2* mutant than in the Kronos-PS control at

the ZT12 peak (Figure 5A), suggesting that CO2 functions as a negative transcriptional regulator

of *CO1*. This is also supported by the downregulation of *CO1* transcripts in a barley line

overexpressing *CO2* [11]. The effect of *CO2* on *CO1* transcription may contribute to the

significant genetic interaction on heading time observed between *CO1* and *CO2* in this study

(Table S3b).

Overexpression of *CO1* and *CO2* in winter barley was associated with the upregulation of *VRN2*, both under LD and SD, where *VRN2* is not normally expressed [11]. A similar upregulation of *VRN2* has been reported in *Brachypodium* plants overexpressing *CO1* under LD [46]. By contrast, knock-down of *CO1* expression in *Brachypodium* [46] or loss-of-function mutants in *co1*, *co2*, or *co1 co2* in Kronos in our study (Figure 5F) were associated with the down regulation of *VRN2* transcripts. Taken together, these results are consistent with *CO1* and *CO2* being promoters of *VRN2* transcription in the temperate grasses, a role that is likely more important during the fall than in the spring, when high expression of *VRN1* results in the downregulation of both *CO1* and *VRN2*.

Photoperiod and vernalization genes converge on the regulation of *FT1*, whose transcript levels are well correlated with heading time [11, 47]. Since *PPD1* is a critical LD transcriptional promoter of *FT1* [12, 32], it was not surprising to see almost undetectable levels of *FT1* in the *ppd1* mutant, which were correlated with low *VRN1* transcript levels and late heading time (Figure 5D-E). The low transcript levels of *VRN1* in the *ppd1* mutant may have contributed to the slight increase in *VRN2* transcript levels observed in this mutant (Figure 5F), because *VRN1* is a known transcriptional repressor of *VRN2* in wheat [27, 48].

The high transcript levels of *VRN2* in the *ppd1* mutant indicates that *PPD1* is not required for the LD expression of *VRN2* (Figure 5F). However, several observations suggest that *PPD1* has a positive effect on *VRN2* transcription. First, *VRN2* transcripts are upregulated under SD in lines carrying the *Ppd-D1a* allele but not in sister lines carrying the *Ppd-D1b* allele [49]. Second, a positive correlation has been reported between *PPD1* and *VRN2* transcript levels in young winter barley plants in spite of a positive correlation between *PPD1* and *VRN1* [11]. Finally, *PPD1* and *VRN2* showed similar transcription profiles in Kronos wild-type (Figure 5C and F). However,

those similarities disappear in the *CO1* and *CO2* mutants, suggesting that the effect of *PPD1* on

*VRN2* is modulated by *CO1* and *CO2*.

To summarize the dynamics of these transcriptional interactions during the seasons, we propose

the following working model for winter wheats. In the fall, *PPD1*, *CO1* and *CO2* promote the

expression of *VRN2* in the leaves, which prevents flowering by repressing the transcriptional

induction of *FT1*. During the winter, the increase in *VRN1* expression results in the gradual

downregulation of both *VRN2* and *CO1*, which facilitates the promotion of *FT1* by *PPD1* and the

initiation of flowering in the spring.

**Protein-protein interactions add an additional level of regulation to wheat heading time**

In addition to the complex transcriptional interactions described above, the proteins encoded by

the photoperiod genes form an intricate network of protein-protein interactions that contribute an

additional layer of complexity to the photoperiodic regulation of heading time. In Arabidopsis,

the central photoperiod gene *CO* is regulated at the transcriptional and posttranscriptional level.

A complex transcriptional regulation is required to establish the presence of *CO* mRNA during

the evening of a long day, when light can stabilize the CO protein, resulting in the activation of

*FT* transcription [23].

Several PRR proteins promote flowering in Arabidopsis by interacting with and stabilizing the

CO protein during the day specifically under LD, and indirectly through their roles in the

circadian clock [50]. Mutations in *PRR* genes delay Arabidopsis flowering under LD but not

under SD (similar to *PPD1* in wheat), with double mutants showing a stronger delay [51]. The

late-flowering Arabidopsis *prr7* mutant can be complemented by rice *OsPRR37* [52], suggesting

some conserved functions in rice and Arabidopsis. In this study, we detected interactions

between wheat PPD1 full-length protein and both CO1 and CO2 in Y2H and validated the CO1-

PPD1 interactions by split-YFP (Figure 7). These results suggest that the interactions between

CO and PRRs likely precede the split between monocot and dicot plants.

In spite of these similarities, there are also important differences between the grass *PPD1/PRR37*

gene and the homologous *PRR* genes from Arabidopsis. The Arabidopsis *PRR* genes are part of

the circadian clock and their mutants affect the expression of circadian clock genes and their

multiple downstream targets. By contrast, the natural mutation in the barley photoperiod

insensitive *ppd-H1* allele did not affect diurnal or circadian cycling of barley clock genes [53].

This result suggests that after the duplication that generated *PRR37* and *PRR73* in the grasses,

which is independent of the Arabidopsis duplication that originated *PRR3* and *PRR7*,

*PRR37/PPD1* acquired a more specialized role in the photoperiodic response with a more limited

role in the circadian clock.

In addition to its interactions with the LD flowering promoter PPD1, CO1 and CO2 can form

heterodimers with the LD flowering repressor VRN2, which is also a member of the CCT

domain protein family [30]. We hypothesize that the interactions with VRN2 might be relevant

to *CO1* and *CO2* function because loss-of-function mutation in *ghd7/vrn2* result in reversals of

functional in *OsPRR37* (*Hd2*) [43] and *OsCO1* (*Hd1*) in rice [41, 42], and *SbCO1* in sorghum

[38]. In rice, GHD7 binds to the transcription-activating domain of OsCO1, probably weakening

or blocking the transcriptional activation of *Ehd1* and *Hd3a* [42]. We have initiated crosses

among Kronos lines with mutations in *vrn2* [26], *co1*, *co2*, and *ppd1* to test these interactions in

wheat.

Recent studies in rice showed that another CCT-domain protein, the NUCLEAR

TRANSCRIPTION FACTOR Y subunit B-11 (NF-YB11 = HEME ACTIVATOR PROTEIN or

HAP-3H = *Ghd8*) has large effects on heading time and significant genetic interactions with

*OsGHD7*, *OsCO1* and *OsPRR37* [43]. We have previously shown that in addition to interacting

with each other, CO2 and VRN2 interact with the same subset of eight NUCLEAR FACTOR-Y

(NF-Y) transcription factors (Figure 7E) [30], suggesting an additional functional connection

between these proteins. This connection through the NF-Y proteins seems to less extensive

between CO1 and VRN2, where only one shared NF-Y interactor has been identified so far

(Figure 7E) [30]. Members of the NF-Y family have been shown to be involved in the regulation

of flowering through interactions with CCT domain proteins CO and COL in other plant species

[54-56]. Using yeast-three-hybrid (Y3H) assays, we previously showed that VRN2 competes

with CO2 for interactions with the NF-Ys, and that mutations in three conserved arginine (R)

residues within the CCT domain of VRN2 reduce the strength of the protein interactions and

competition with CO2 [30]. Since these same mutations abolish or reduce VRN2 function as a

LD flowering repressor, we hypothesize that the protein-protein interactions involving VRN2-

CO2-NF-Y CCT domain could be relevant for their function [30].

Phytochromes PHYB and PHYC are involved in the transcriptional regulation by light of both

*PPD1/PRR37* and *VRN2* in wheat [15, 28], *Brachypodium* [57], rice [58] and sorghum [37]. In

addition, OsPHYA, OsPHYB and OsGI proteins have been shown to interact directly with

OsGHD7, regulating protein stability [59]. We observed similar interactions between VRN2 and

both PHYB and PHYC in wheat. These two phytochromes also showed Y2H interactions with

CO1 and CO2, but not PPD1 (Figure 7E and S5). Additional experiments will be required to test

if these phytochrome interactions can stabilize the VRN2, CO1 and CO2 proteins in wheat as

they stabilize GHD7 in rice [59] or CO in Arabidopsis [60].

In summary, we found that CO1 and CO2 proteins can physically interact with each other and

with both PPD1 (a central LD flowering promoter) and VRN2 (a central LD flowering

repressor), providing a physical link between these central photoperiodic proteins (Figure 7E).

Among the 15 pairwise interactions tested among PHYB, PHYC, PPD1, CO1, CO2 and VRN2

80% showed positive interactions in Y2H assays indicating a complex and dense network of

protein interactions. We hypothesize that this network of protein interactions provides an

additional layer of molecular mechanisms that result in the complex genetic interactions in

heading time detected in this study.

**Two interacting photoperiod pathways in grasses**

A central conclusion of our study is that *PPD1* can perceive the differences in photoperiod and

adjust heading time accordingly even in the absence of any functional copy of *CO1* or *CO2*.

Similarly, *CO1* can accelerate heading time under LD in the absence of a functional copy of

*PPD1*. This is also true in rice, where heading time differences between SD and LD are observed

for *OsCO1* alleles in *Osprr37* mutants, and for *OsPRR37* alleles in rice *Osco1* mutants [43].

These results suggest that grasses have either two photoperiod pathways, each of them capable of

perceiving the length of the day/night and of regulating heading time, or a single point of

perception of the differences in photoperiod upstream of *PPD1* and *CO1* (e.g. *ELF3*), which can

then transmit the signal to both genes.

Another important conclusion from our study and from previous studies in sorghum and rice [37,

41, 43], is that the *CO* and *PRR37/PPD1* photoperiod pathways are extensively interconnected.

Changes in the balance of these interactions seem to play a central role in the diversity of

photoperiod responses in grasses, as suggested by changes in function observed when

photoperiodic genes are swapped between LD and SD species or when the same genes are

expressed in different mutant backgrounds in its own species. We hypothesize that the complex

feedback transcriptional regulatory loops and transcriptional interactions among photoperiod

genes along with the intricate interactions among their encoded proteins contribute to the

complex genetic interactions in heading time observed in this study and to the diversity of

photoperiodic responses in grasses.

**Materials and methods**

**Plant materials**

The materials used in this study included the tetraploid wheat variety Kronos (*Triticum*

*turgidum* ssp. *durum*), which carries a functional and dominant *Vrn-A1* allele for spring growth

habit, a functional but recessive *vrn-B1* allele for winter growth habit, a functional *Vrn-B2* long

day repressor (*ZCCT-B2a* and *ZCCT-B2b* [26]), and the *Ppd-A1a* (photoperiod insensitive or PI)

allele that confers earlier flowering under short days resulting in a reduced photoperiodic

response. We also used in our study near-isogenic lines of Kronos carrying either the

photoperiod sensitive *Ppd-A1b* (PS) allele [18] or no functional alleles in the *PPD-A1* and *PPD-*

*B1* homeologs (*ppd1*) [19].

Using previously published wheat sequences for *CO1* [45] and *CO2* [39], and the Kronos

transcriptome [61] we identified the A and B coding sequences for *CO1* and *CO2* in Kronos and

deposited them in GenBank under accession numbers MT043302 (*CO-A1*), MT043303 (*CO-B1*),

MT043304 (*CO-A2*), MT043305 (*CO-B2*). We used those sequences to identify loss-of-function

mutations in *CO1* and *CO2* in the Kronos TILLING population [21, 22], which are described in

detail in the Results section. We then combined mutations in the A and B homeologs from each

gene to generate loss-of-function *co1* (henceforth *co1*) and *co2* mutants (henceforth *co2*). We

backcrossed the individual mutant lines with wild-type Kronos at least two times to reduce

background mutations and then combined all four homozygous mutations to produce a *co1 co2*

mutant (henceforth *co1 co2*). In addition to the wild type Kronos-PI, the four homozygous *Co1*

*Co2*, *co1 Co2*, *Co1 co2* and *co1 co2* allelic classes were generated in Kronos-PS and the *ppd1*

mutant. We were not able to recover the triple loss-of-function mutant *ppd1 co1 co2* from any of

the tested progenies.

**Phenotyping**

We stratified wheat grains at 4 °C for 2 d in the dark and after germination we planted them in

pots in the soil. We evaluated the mutant and control lines in growth chambers set to 22 °C

during daylight and 17 °C during dark periods. Lights were set to 260-300 µmol m$^{-2}$ s$^{-1}$ and

were on for 16 h in LD experiments and 8 h in short day (SD) experiments. We recorded

flowering time as the total time from planting in soil to full emergences of the spike from the

sheath. Spikes were threshed and total grain number counted per spike.

**Statistical analysis**

We compared heading time in different genotypes using one-way ANOVAs for single locus and

factorial ANOVAs for two or more loci. We tested homogeneity of variances using the Levene's

test and normality of residuals with the Shapiro–Wilks test. We transformed the data when

necessary to satisfy the ANOVA assumptions. We performed all statistical analyses using SAS

9.4 (SAS Institute). Error bars in all graphs are standard errors of the means (s.e.m.).

**Gene expression analysis**

To study the effect of *ppd1*, *co1* and *co2* mutant alleles on the transcription profiles of genes

affecting wheat flowering, we collected leaves from plants grown in a CONVIRON growth

chamber with metal halide bulbs supplemented by florescent bulbs (16 h light / 8 h darkness, 22

°C day temperature and 17 °C night temperature, 300 µmol m$^{-2}$ s$^{-1}$). For the time course

experiment, we collected samples in liquid nitrogen every 4 h throughout the day from the newly

expanded fourth leaf. In this experiment, we included the *ppd1* mutant and all four possible

combinations of the *CO1* and *CO2* wild type and mutant alleles in the Kronos-PS (*Ppd-A1b*)

background.

RNA was isolated from leaves using the Spectrum Plant Total RNA Kit (Sigma-Aldrich)

following the manufacturer's recommendations. We treated the RNA samples with DNase

(Promega) according to the manufacturer's protocol prior to first-strand cDNA synthesis. We

assessed RNA integrity by a combination of A260/A280 and A260/A230 measurements and

running samples on a denaturing formaldehyde-agarose gel as recommended in the Qiagen

RNAeasy Mini Handbook. We used only samples with 260/280 and 260/230 ratios > 1.7 and no

degradation. We synthesized cDNA using 2.0 µg of total RNA per sample and M-MLV Reverse

Transcriptase (Promega) and poly-T primers (Sigma-Aldrich).

We performed the quantitative reverse transcription-PCR (qRT-PCR) assays using SybrGreen

Takara Master Mix in a 7500 Fast Real-Time PCR system (Applied Biosystems). For PCR, we

used one cycle at 95°C for 30 s and 40 cycles of 95°C for 5 s, 60°C for 5 s, and 72°C for 34 s,

followed by a melting curve program. We quantified expression using the delta Ct method ($2^-$

$^{\Delta CT}$) with *ACTIN* as endogenous control [15]. Primer efficiencies were all > 95% and their

sequences are summarized in Table S5. Primer sequences were previously described for *FT1* and

*VRN1* [47], *PPD1* [45], *CO1* and *CO2* [15] and *VRN2* [26].

**Yeast-two-hybrid (Y2H) and split yellow fluorescent protein (YFP) assays**

The full-length coding region of PPD1 was amplified from Kronos and the PHYB truncations

were amplified from *Triticum monococcum* with primers listed in Supplementary Table S5. The

amplified fragments were cloned into yeast vectors pGBKT7 and pGADT7 (Clontech,

http://www.clontech.com/), which were used to transform yeast strain Y2HGold (Clontech)

using the lithium acetate method. Transformants were selected on SD medium lacking leucine

(L) and tryptophan (W) plates and re-plated on SD medium lacking L, W, histidine (H) and

adenine (A) to test the interactions. Yeast vectors for VRN2 (full-length, both bait and prey),

CO1 (prey, including both B-boxes and CCT domain, amino acids 10 to 379) and CO2 (prey,

including both B-boxes and CCT domain, 15 to 356 amino acids) were previously described in

[30], and those for PHYC (N-PHYC, C-PHYC and FL-PHYC) and full-length PHYB (as prey)

in [15]. CO1 and CO2 showed strong auto-activation when used as baits so they were only used

as preys. The full-length PHYB gave strong autoactivation when used as bait, so we generated

two additional truncations N-PHYB and C-PHYB (primers in Table S5). Only N-PHYB can be

used as bait without autoactivation, and all three (N-PHYB, C-PHYB and FL-PHYB) can be

used as preys in Y2H assays. The truncated versions of PHYC include amino acids 1–600 (N-

terminal) and 601-1139 (C-terminal) [15], and those for PHYB amino acids (1-625 (N-terminal)

and 626-1166 (C-terminal).

For the bi-molecular fluorescent complementation (BiFC or split YFP) assays, each set of wheat

proteins was recombined into modified Gateway-compatible pY736 and pY735 which contain

the *UBIQUITIN* promoter to generate YFP-N-terminal fragment and YFP-C-terminal-fragment

fusion proteins. Wheat protoplasts were prepared, transfected and visualized as described in [62]

**ACKNOWLEDEMENTS**

This project was supported by NRI Competitive Grants 2017-67007-25939 and 2016-67013-

24617 from the USDA National Institute of Food and Agriculture (NIFA) and by the Howard

Hughes Medical Institute. Daniel Woods is a Howard Hughes Medical Institute Fellow of the

Life Sciences Research Foundation.

**AUTHOR CONTRIBUTIONS**

Lindsay M. Shaw: Investigation, Writing - Original Draft -Review & Editing. Chengxia Li:

Investigation, Writing - Original Draft -Review & Editing. Daniel Woods: Investigation, Writing

-Review & Editing. Maria A. Alvarez: Investigation, Writing -Review & Editing. Huiqiong:

Investigation, Methodology, Writing -Review & Editing. Mei Yee Lau: Investigation. Andrew

Chen: Investigation, Writing -Review & Editing. Jorge Dubcovsky: Conceptualization, Formal

analysis, Supervision, Project administration and funding acquisition, Writing –extensive Review

& Editing.

**CONFLICT OF INTEREST**

The authors declare that they do not have any conflict of interest.

**REFERENCES**

838    1.    Bouche F, Woods DP, Amasino RM. Winter memory throughout the plant kingdom:

839           different paths to flowering. Plant Physiol. 2017;173(1):27-35. doi: 10.1104/pp.16.01322.

840           PMID: 27756819.

841    2.    Garner WW, Allard HA. Effect of the relative length of day and night and other factors of

842           the environment on growth and reproduction in plants. J Agric Res. 1919;18:0553-606.

843           PMID: WOS:000188347200038.

844    3.    Andres F, Coupland G. The genetic basis of flowering responses to seasonal cues. Nat

845           Rev Genet. 2012;13(9):627-39. doi: 10.1038/nrg3291. PMID: 22898651.

846    4.    Suarez-Lopez P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G.

847           CONSTANS mediates between the circadian clock and the control of flowering in

848           Arabidopsis. Nature. 2001;410(6832):1116-20. doi: 10.1038/35074138. PMID:

849           11323677.

850    5.    Yanovsky MJ, Kay SA. Molecular basis of seasonal time measurement in Arabidopsis.

851           Nature. 2002;419(6904):308-12. doi: 10.1038/nature00996. PMID: 12239570.

852    6.    Putterill J, Robson F, Lee K, Simon R, Coupland G. The CONSTANS gene of

853           Arabidopsis promotes flowering and encodes a protein showing similarities to zinc finger

854           transcription factors. Cell. 1995;80(6):847-57. doi: 10.1016/0092-8674(95)90288-0.

855           PMID: 7697715.

856    7.    Griffiths S, Dunford RP, Coupland G, Laurie DA. The evolution of *CONSTANS*-like

857           gene families in barley, rice, and Arabidopsis. Plant Physiol. 2003;131(4):1855-67. doi:

858           10.1104/pp.102.016188. PMID: 12692345.

8.     Yano M, Katayose Y, Ashikari M, Yamanouchi U, Monna L, Fuse T, et al. Hd1, a major

photoperiod sensitivity quantitative trait locus in rice, is closely related to the Arabidopsis

flowering time gene CONSTANS. Plant Cell. 2000;12(12):2473-84. doi:

10.1105/tpc.12.12.2473. PMID: 11148291.

9.     Higgins JA, Bailey PC, Laurie DA. Comparative genomics of flowering time pathways

using *Brachypodium distachyon* as a model for the temperate grasses. PLoS One.

2010;5(4):e10065. doi: 10.1371/journal.pone.0010065. PMID: 20419097.

10.    Campoli C, Drosse B, Searle I, Coupland G, von Korff M. Functional characterisation of

*HvCO1*, the barley (*Hordeum vulgare*) flowering time ortholog of *CONSTANS*. Plant J.

2012;69(5):868-80. doi: 10.1111/j.1365-313X.2011.04839.x. PMID: 22040323.

11.    Mulki MA, von Korff M. CONSTANS controls floral repression by up-regulating

*VERNALIZATION2* (*VRN-H2*) in Barley. Plant Physiol. 2016;170(1):325-37. doi:

10.1104/pp.15.01350. PMID: 26556793.

12.    Turner A, Beales J, Faure S, Dunford RP, Laurie DA. The pseudo-response regulator

*Ppd-H1* provides adaptation to photoperiod in barley. Science. 2005;310(5750):1031-4.

doi: 10.1126/science.1117619. PMID: 16284181.

13.    Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, et al. Natural variation in

*OsPRR37* regulates heading date and contributes to rice cultivation at a wide range of

latitudes. Mol Plant. 2013;6(6):1877-88. doi: 10.1093/mp/sst088. PMID: 23713079.

14.    Murphy RL, Klein RR, Morishige DT, Brady JA, Rooney WL, Miller FR, et al.

Coincident light and clock regulation of pseudoresponse regulator protein 37 (PRR37)

controls photoperiodic flowering in sorghum. Proc Natl Acad Sci U S A.

2011;108(39):16469-74. doi: 10.1073/pnas.1106212108. PMID: 21930910.

15.   Chen A, Li C, Hu W, Lau MY, Lin H, Rockwell NC, et al. Phytochrome C plays a major

role in the acceleration of wheat flowering under long-day photoperiod. Proc Natl Acad

Sci U S A. 2014;111(28):10037-44. doi: 10.1073/pnas.1409795111. PMID: 24961368.

16.   Beales J, Turner A, Griffiths S, Snape JW, Laurie DA. A pseudo-response regulator is

misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat (Triticum aestivum

L.). Theor Appl Genet. 2007;115(5):721-33. doi: 10.1007/s00122-007-0603-4. PMID:

17634915.

17.   Wilhelm EP, Turner AS, Laurie DA. Photoperiod insensitive *Ppd-A1a* mutations in

tetraploid wheat (*Triticum durum* Desf.). Theor Appl Genet. 2009;118(2):285-94. doi:

10.1007/s00122-008-0898-9. PMID: 18839130.

18.   Pearce S, Vanzetti LS, Dubcovsky J. Exogenous gibberellins induce wheat spike

development under short days only in the presence of *VERNALIZATION1*. Plant Physiol.

2013;163(3):1433-45. doi: 10.1104/pp.113.225854. PMID: 24085801.

19.   Pearce S, Shaw LM, Lin H, Cotter JD, Li C, Dubcovsky J. Night-break experiments shed

light on the *Photoperiod1*-mediated flowering. Plant Physiol. 2017;174(2):1139-50. doi:

10.1104/pp.17.00361. PMID: 28408541.

20.   Shaw LM, Turner AS, Herry L, Griffiths S, Laurie DA. Mutant alleles of *Photoperiod-1*

in wheat (*Triticum aestivum* L.) that confer a late flowering phenotype in long days.

PLoS One. 2013;8(11):e79459. doi: 10.1371/journal.pone.0079459. PMID: 24244507.

21.   Krasileva KV, Vasquez-Gross HA, Howell T, Bailey P, Paraiso F, Clissold L, et al.

Uncovering hidden variation in polyploid wheat. Proc Natl Acad Sci U S A.

2017;114(6):E913-E21. doi: 10.1073/pnas.1619268114. PMID: 28096351.

22.    Uauy C, Paraiso F, Colasuonno P, Tran RK, Tsai H, Berardi S, et al. A modified

905           TILLING approach to detect induced mutations in tetraploid and hexaploid wheat. BMC

906           Plant Biol. 2009;9:115. doi: 10.1186/1471-2229-9-115. PMID: 19712486.

907    23.    Valverde F. CONSTANS and the evolutionary origin of photoperiodic timing of

908           flowering. J Exp Bot. 2011;62(8):2453-63. doi: 10.1093/jxb/erq449. PMID: 21239381.

909    24.    Waddington SR, Cartwright PM, Wall PC. A quantitative scale of spike initial and pistil

910           development in barley and wheat. Ann Bot-London. 1983;51(1):119-30. PMID:

911           WOS:A1983QC01200013.

912    25.    Alvarez MA, Tranquilli G, Lewis S, Kippes N, Dubcovsky J. Genetic and physical

913           mapping of the earliness *per se* locus *Eps-A$^m$1* in *Triticum monococcum* identifies *EARLY*

914           *FLOWERING 3* (*ELF3*) as a candidate gene. Funct Integr Genomic. 2016;16(4):365-82.

915           doi: 10.1007/s10142-016-0490-3. PMID: 27085709.

916    26.    Distelfeld A, Tranquilli G, Li C, Yan L, Dubcovsky J. Genetic and molecular

917           characterization of the *VRN2* loci in tetraploid wheat. Plant Physiol. 2009;149(1):245-57.

918           doi: 10.1104/pp.108.129353. PMID: 19005084.

919    27.    Chen A, Dubcovsky J. Wheat TILLING mutants show that the vernalization gene *VRN1*

920           down-regulates the flowering repressor *VRN2* in leaves but is not essential for flowering.

921           PLoS Genet. 2012;8(12):e1003134. doi: 10.1371/journal.pgen.1003134. PMID:

922           23271982.

923    28.    Pearce S, Kippes N, Chen A, Debernardi JM, Dubcovsky J. RNA-seq studies using wheat

924           *PHYTOCHROME B* and *PHYTOCHROME C* mutants reveal shared and specific

925           functions in the regulation of flowering and shade-avoidance pathways. BMC Plant Biol.

926           2016;16(1):141. doi: 10.1186/s12870-016-0831-3. PMID: 27329140.

29.    Kippes N, van Gessel C, Hamilton J, Akpinar A, Budak H, Dubcovsky J, et al. Effect of

*phyB* and *phyC* loss-of-function mutations on wheat transcriptome under short and long

day photoperiods. bioRxiv. 2020. doi: https://doi.org/10.1101/2020.04.07.030197.

30.    Li C, Distelfeld A, Comis A, Dubcovsky J. Wheat flowering repressor VRN2 and

promoter CO2 compete for interactions with NUCLEAR FACTOR-Y complexes. Plant J.

2011;67(5):763-73. doi: 10.1111/j.1365-313X.2011.04630.x. PMID: 21554456.

31.    Faure S, Turner AS, Gruszka D, Christodoulou V, Davis SJ, von Korff M, et al. Mutation

at the circadian clock gene *EARLY MATURITY 8* adapts domesticated barley (*Hordeum*

*vulgare*) to short growing seasons. Proc Natl Acad Sci U S A. 2012;109(21):8328-33.

doi: 10.1073/pnas.1120496109. PMID: 22566625.

32.    Gao M, Geng F, Klose C, Staudt A-M, Huang H, Nguyen D, et al. Phytochromes

measure photoperiod in *Brachypodium*. bioRxiv. 2019. doi: 10.1101/697169.

33.    Diaz A, Zikhali M, Turner AS, Isaac P, Laurie DA. Copy number variation affecting the

*Photoperiod-B1* and *Vernalization-A1* genes is associated with altered flowering time in

wheat (*Triticum aestivum*). PLoS One. 2012;7(3):e33234. doi:

10.1371/journal.pone.0033234. PMID: 22457747.

34.    Stracke S, Haseneyer G, Veyrieras JB, Geiger HH, Sauer S, Graner A, et al. Association

mapping reveals gene action and interactions in the determination of flowering time in

barley. Theor Appl Genet. 2009;118(2):259-73. doi: 10.1007/s00122-008-0896-y. PMID:

18830577.

35.    Alqudah AM, Sharma R, Pasam RK, Graner A, Kilian B, Schnurbusch T. Genetic

dissection of photoperiod response based on GWAS of pre-anthesis phase duration in

spring barley. PLoS One. 2014;9(11):e113120. doi: 10.1371/journal.pone.0113120.

PMID: 25420105.

36.  Casto AL, Mattison AJ, Olson SN, Thakran M, Rooney WL, Mullet JE. *Maturity2*, a

novel regulator of flowering time in *Sorghum bicolor*, increases expression of *SbPRR37*

and *SbCO* in long days delaying flowering. PLoS One. 2019;14(4):e0212154. doi:

10.1371/journal.pone.0212154. PMID: 30969968.

37.  Yang S, Murphy RL, Morishige DT, Klein PE, Rooney WL, Mullet JE. Sorghum

phytochrome B inhibits flowering in long days by activating expression of *SbPRR37* and

*SbGHD7*, repressors of *SbEHD1*, *SbCN8* and *SbCN12*. PLoS One. 2014;9(8):e105352.

doi: 10.1371/journal.pone.0105352. PMID: 25122453.

38.  Yang S, Weers BD, Morishige DT, Mullet JE. *CONSTANS* is a photoperiod regulated

activator of flowering in sorghum. BMC Plant Biol. 2014;14:148. doi: 10.1186/1471-

2229-14-148. PMID: 24884377.

39.  Nemoto Y, Kisaka M, Fuse T, Yano M, Ogihara Y. Characterization and functional

analysis of three wheat genes with homology to the *CONSTANS* flowering time gene in

transgenic rice. Plant J. 2003;36(1):82-93. doi: 10.1046/j.1365-313x.2003.01859.x.

PMID: 12974813.

40.  Woods DP, McKeown MA, Dong Y, Preston JC, Amasino RM. Evolution of

*VRN2/Ghd7*-like genes in vernalization-mediated repression of grass flowering. Plant

Physiol. 2016;170(4):2124-35. doi: 10.1104/pp.15.01279. PMID: 26848096.

41.  Fujino K, Yamanouchi U, Nonoue Y, Obara M, Yano M. Switching genetic effects of the

flowering time gene *Hd*1 in LD conditions by *Ghd7* and *OsPRR37* in rice. Breeding Sci.

2019;69(1):127-32. doi: 10.1270/jsbbs.18060. PMID: 31086490.

42.    Zhang Z, Hu W, Shen G, Liu H, Hu Y, Zhou X, et al. Alternative functions of Hd1 in

973             repressing or promoting heading are determined by Ghd7 status under long-day

974             conditions. Sci Rep-Uk. 2017;7(1):5388. doi: 10.1038/s41598-017-05873-1. PMID:

975             28710485.

976    43.    Zhang B, Liu H, Qi F, Zhang Z, Li Q, Han Z, et al. Genetic interactions among *Ghd7*,

977             *Ghd8*, *OsPRR37* and *Hd1* contribute to large variation in heading date in rice. Rice (N

978             Y). 2019;12(1):48. doi: 10.1186/s12284-019-0314-x. PMID: 31309345.

979    44.    Maurer A, Draba V, Jiang Y, Schnaithmann F, Sharma R, Schumann E, et al. Modelling

980             the genetic architecture of flowering time control in barley through nested association

981             mapping. BMC Genomics. 2015;16:290. doi: 10.1186/s12864-015-1459-7. PMID:

982             25887319.

983    45.    Shaw LM, Turner AS, Laurie DA. The impact of photoperiod insensitive *Ppd-1a*

984             mutations on the photoperiod pathway across the three genomes of hexaploid wheat

985             (*Triticum aestivum*). Plant J. 2012;71(1):71-84. doi: 10.1111/j.1365-313X.2012.04971.x.

986             PMID: 22372488.

987    46.    Qin Z, Bai Y, Muhammad S, Wu X, Deng P, Wu J, et al. Divergent roles of *FT-like 9* in

988             flowering transition under different day lengths in *Brachypodium distachyon*. Nat

989             Commun. 2019;10(1):812. doi: 10.1038/s41467-019-08785-y. PMID: 30778068.

990    47.    Yan L, Fu D, Li C, Blechl A, Tranquilli G, Bonafede M, et al. The wheat and barley

991             vernalization gene *VRN3* is an orthologue of *FT*. Proc Natl Acad Sci U S A.

992             2006;103(51):19581-6. doi: 10.1073/pnas.0607142103. PMID: 17158798.

993    48.    Deng W, Casao MC, Wang P, Sato K, Hayes PM, Finnegan EJ, et al. Direct links

994           between the vernalization response and other key traits of cereal crops. Nat Commun.

995           2015;6:5882. doi: 10.1038/ncomms6882. PMID: 25562483.

996    49.    Turner AS, Faure S, Zhang Y, Laurie DA. The effect of day-neutral mutations in barley

997           and wheat on the interaction between photoperiod and vernalization. Theor Appl Genet.

998           2013;126(9):2267-77. doi: 10.1007/s00122-013-2133-6. PMID: 23737074.

50.    Hayama R, Sarid-Krebs L, Richter R, Fernandez V, Jang S, Coupland G. PSEUDO

RESPONSE REGULATORs stabilize CONSTANS protein to promote flowering in

response to day length. Embo J. 2017;36(7):904-18. doi: 10.15252/embj.201693907.

PMID: 28270524.

51.    Nakamichi N, Kita M, Ito S, Yamashino T, Mizuno T. PSEUDO-RESPONSE

REGULATORS, PRR9, PRR7 and PRR5, together play essential roles close to the

circadian clock of *Arabidopsis thaliana*. Plant Cell Physiol. 2005;46(5):686-98. doi:

10.1093/pcp/pci086. PMID: 15767265.

52.    Murakami M, Tago Y, Yamashino T, Mizuno T. Characterization of the rice circadian

clock-associated pseudo-response regulators in *Arabidopsis thaliana*. Biosci Biotech

Bioch. 2007;71(4):1107-10. doi: 10.1271/bbb.70048. PMID: 17420570.

53.    Campoli C, Shtaya M, Davis SJ, von Korff M. Expression conservation within the

circadian clock of a monocot: natural variation at barley *Ppd-H1* affects circadian

expression of flowering time genes, but not clock orthologs. BMC Plant Biol.

2012;12:97. doi: 10.1186/1471-2229-12-97. PMID: 22720803.

54.    Ben-Naim O, Eshed R, Parnis A, Teper-Bamnolker P, Shalit A, Coupland G, et al. The

CCAAT binding factor can mediate interactions between CONSTANS-like proteins and

DNA. Plant J. 2006;46(3):462-76. doi: 10.1111/j.1365-313X.2006.02706.x. PMID:

16623906.

55. Kumimoto RW, Adam L, Hymus GJ, Repetti PP, Reuber TL, Marion CM, et al. The

Nuclear Factor Y subunits NF-YB2 and NF-YB3 play additive roles in the promotion of

flowering by inductive long-day photoperiods in Arabidopsis. Planta. 2008;228(5):709-

23. doi: 10.1007/s00425-008-0773-6. PMID: 18600346.

56. Wenkel S, Turck F, Singer K, Gissot L, Le Gourrierec J, Samach A, et al. CONSTANS

and the CCAAT box binding complex share a functionally important domain and interact

to regulate flowering of Arabidopsis. Plant Cell. 2006;18(11):2971-84. doi:

10.1105/tpc.106.043299. PMID: 17138697.

57. Woods DP, Ream TS, Minevich G, Hobert O, Amasino RM. PHYTOCHROME C is an

essential light receptor for photoperiodic flowering in the temperate grass, *Brachypodium*

*distachyon*. Genetics. 2014;198(1):397-408. doi: 10.1534/genetics.114.166785. PMID:

25023399.

58. Osugi A, Itoh H, Ikeda-Kawakatsu K, Takano M, Izawa T. Molecular dissection of the

roles of phytochrome in photoperiodic flowering in rice. Plant Physiol.

2011;157(3):1128-37. doi: 10.1104/pp.111.181792. PMID: 21880933.

59. Zheng T, Sun J, Zhou S, Chen S, Lu J, Cui S, et al. Post-transcriptional regulation of

Ghd7 protein stability by phytochrome and OsGI in photoperiodic control of flowering in

rice. New Phytol. 2019;224(1):306-20. doi: 10.1111/nph.16010. PMID: 31225911.

60. Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G.

Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. Science.

2004;303(5660):1003-6. doi: 10.1126/science.1091761. PMID: 14963328.

61.    Krasileva KV, Buffalo V, Bailey P, Pearce S, Ayling S, Tabbita F, et al. Separating

homeologs by phasing in the tetraploid wheat transcriptome. Genome Biol.

2013;14(6):R66. doi: 10.1186/gb-2013-14-6-r66. PMID: 23800085.

62.    Shan Q, Wang Y, Li J, Gao C. Genome editing in rice and wheat using the CRISPR/Cas

system. Nat Protoc. 2014;9(10):2395-410. doi: 10.1038/nprot.2014.157. PMID:

25232936.

1045

1046

**SUPPORTING INFORMATION CAPTIONS**

**Supplementary Figure S1.** Shoot apical meristem (SAM) and spike development.

**Supplementary Figure S2.** Dissection of developing spikes.

**Supplementary Figure S3.** Effect of *ppd1*, *co1* and *co2* loss-of-function mutations and

photoperiod on *CO1* and *CO2* transcript levels.

**Supplementary Figure S4.** Effect of *phyB*-null and *phyC*-null mutations in Kronos-PI on the

transcriptional profiles of *CO1* and *CO2* under SD and LD.

**Supplementary Figure S5.** Yeast-two-hybrid (Y2H) assays.

**Supplementary Table S1.** Genome-specific primer sequences and PCR conditions for

TILLING.

**Supplementary Table S2.** Number of mutations detected in the targeted regions of wheat *CO1*

and *CO2* homologs in the Kronos TILLING population.

**Table S3**. Analysis of variance for heading time under long days.

**Table S4**. Analysis of variance for heading time under short days.

**Table S5**. Primers used in the qRT-PCR and Y2H experiments.