Epistatic interactions between *PHOTOPERIOD-1, CONSTANS 1* and *CONSTANS 2*
modulate the photoperiodic response in wheat

Lindsay M. Shaw[1, 2]*, Chengxia Li[1, 3]*, Daniel P. Woods[1, 3]*, Maria A. Alvarez[1]*, Huiqiong

Lin[1, 3], Mei Y. Lau[1], Andrew Chen[1, 4], and Jorge Dubcovsky[1, 3#]

*The first four authors contributed equally to this work

[1] Department of Plant Sciences, University of California, Davis, CA 95616. U.S.A.

[2] Currently at Queensland Alliance for Agriculture and Food Innovation, The University of

Queensland, Brisbane, QLD 4072, Australia

[3] Howard Hughes Medical Institute, Chevy Chase, MD 20815, U.S.A.

[4] Currently at School of Agriculture and Food Science, The University of Queensland, St Lucia,

QLD, 4067, Australia.

Lindsay Shaw: ORCID 0000-0001-8078-2409
Chengxia Li: ORCID 0000-0003-4634-8451
Maria Alejandra Alvarez: ORCID 0000-0003-2916-4255
Daniel P. Woods: ORCID 0000-0002-1498-5707
Jorge Dubcovsky: ORCID 0000-0002-7571-4345
20
21

Short title: Regulation of wheat heading time by photoperiod

 # Corresponding authors: Jorge Dubcovsky, Email: jdubcovsky@ucdavis.edu, Phone: 530
(752)-5159.

Abstract

In Arabidopsis, *CONSTANS* (*CO*) integrates light and circadian clock signals to promote

flowering under long days (LD). In the grasses, a duplication generated two paralogs designated

as *CONSTANS 1* (*CO1*) and *CONSTANS 2* (*CO2*). Here we show that in tetraploid wheat plants

grown under LD, combined loss-of-function mutations in the A and B-genome homeologs of

CO1 and *CO2* (*co1 co2*) result in a small (3 d) but significant (*P*<0.0001) acceleration of heading

time both in *PHOTOPERIOD1* (*PPD1*) sensitive (*Ppd-A1b*, functional ancestral allele) and

insensitive (*Ppd-A1a*, functional dominant allele) backgrounds. Under short days (SD), *co1 co2*

mutants headed 13 d earlier than the wild type (*P*<0.0001) in the presence of *Ppd-A1a*. However,

in the presence of *Ppd-A1b*, spikes from both genotypes failed to emerge by 180 d. These results

indicate that *CO1* and *CO2* operate mainly as weak heading time repressors in both LD and SD.

By contrast, in *ppd1* mutants with loss-of-function mutations in both *PPD1* homeologs, the wild

type *Co1* allele accelerated heading time >60 d relative to the *co1* mutant allele under LD. We

detected significant genetic interactions among *CO1*, *CO2* and *PPD1* genes on heading time,

which were reflected in complex interactions at the transcriptional and protein levels. Loss-of-

function mutations in *PPD1* delayed heading more than combined *co1 co2* mutations and, more

importantly, *PPD1* was able to perceive and respond to differences in photoperiod in the absence

of functional *CO1* and *CO2* genes. Similarly, *CO1* was able to accelerate heading time in

response to LD in the absence of a functional *PPD1*. Taken together, these results indicate that

PPD1 and *CO1* are able to respond to photoperiod in the absence of each other, and that

interactions between these two photoperiod pathways at the transcriptional and protein levels are

important to fine-tune the flowering response in wheat.

Author summary

An understanding of the mechanisms involved in the regulation of wheat heading time is required to engineer more productive varieties better adapted to new or changing environments. A large proportion of wheat's natural variation in heading time is determined by differences in genes controlling the photoperiodic response. In this study, we show that the wheat *PHOTOPERIOD1* (*PPD1*) gene has a stronger effect on heading time than *CONSTANS1* (*CO1*) and *CO2* in the regulation of the photoperiodic response, and that complex genetic interactions among these genes are important to fine-tune heading time. Using loss-of-function mutants for both *CO1* and *CO2*, we demonstrate that these genes are not required for *PPD1* to perceive differences in photoperiod and regulate heading time. Similarly, we show that in the absence of *PPD1*, *CO1* can accelerate heading time more than 60 days in response to long days. Our results indicate that each of these two wheat photoperiod pathways can respond to differences in photoperiod even in the absence of the other one. Differences in the relative importance of these two pathways and in their epistatic interactions have contributed to the diversity of photoperiodic responses observed in different grass species.

Introduction

A precise adjustment of flowering time to optimal environmental conditions is critical to maximize plant reproductive success. Plants have evolved complex regulatory mechanisms to adjust flowering initiation in response to seasonal changes and other environmental cues. Many plants, including the temperate cereals, require exposure to extended periods of cold (vernalization) to gain competence to flower [1]. After their vernalization requirement has been fulfilled, many plants still require inductive photoperiods to accelerate flowering. Based on their light requirements for floral induction, plants can be classified into long day (LD), short day (SD) or day-neutral plants [2].

The regulatory pathway that controls flowering in response to photoperiod is best characterized in the model plant *Arabidopsis thaliana*, a long day plant that flowers earlier when the days are longer than a critical threshold. In this species, the external coincidence model explains the integration of light signals with the circadian clock. The clock determines diurnal oscillations of the gene *CONSTANS* (*CO*), which results in the coincidence of the expression peak with the light phase under LD and with the dark phase under SD. Light stabilizes the CO protein, resulting in the induction of *FLOWERING LOCUS T* (*FT*) and the promotion of flowering under LD. Under SD, the degradation of the CO protein in the dark prevents the activation of *FT* [3-5]. Thus, *co* mutants in Arabidopsis flower later than wild type under LD but at the same time as wild type under SD [6].

CO belongs to a family of transcription factors characterized by two zinc finger B-Box domains at the N-terminus and by a CONSTANS, CONSTANS-like, TIMING OF CAB EXPRESSION 1 (CCT) domain at the C-terminus, a structure conserved in Arabidopsis and the grass homologs [7]. The *Oryza sativa* (rice) genome contains a single *CO* gene designated as *Hd1* [8], whereas

two paralogs designated as *CO1* and *CO2* have been identified in *Brachypodium distachyon*,

barley and wheat [9]. In the last three species, the predicted CO1 protein lacks conserved

cysteine residues at the start of the second B-box, suggesting a modified or non-functional B-box

2 in the temperate grasses [7, 9]. Phylogenetic and comparative genomics analyses suggest that

CO1 and *CO2* paralogs originated by a segmental duplication at the base of the grass family,

with *CO1* located in the ancestral position colinear with rice *Hd1,* whereas *CO2* was lost in rice

and is likely not functional in sorghum [9]

Overexpression of *CO1* and *CO2* under the maize *UBIQUITIN* promoter in spring barley,

promoted flowering under both LD and SD conditions by upregulating *FT1* [10, 11]. However,

both *Ubi::CO1* and *Ubi::CO2* transgenic lines flowered 42-43 d later in SD than in LD

indicating the existence of additional genes controlling the photoperiodic response [10, 11]. In

barley, the response to photoperiod is determined mainly by the pseudo-response regulator gene

PHOTOPERIOD1 (*PPD1*) [12], also known as *PRR37* in rice [13] and *Sorghum bicolor*

(sorghum) [14]. This gene is regulated by the circadian clock and the phytochrome-mediated

light signaling pathway and encodes a protein with a pseudo-receiver domain and a CCT domain

[14, 15].

In wheat, natural deletions in the promoter region of *PPD-A1* (*Ppd-A1a* allele) and *PPD-D1*

(*Ppd-D1a* allele) are associated with increased transcript levels and accelerated heading under

SD relative to the ancestral *Ppd1b* allele [16, 17]. As a result, plants carrying the *Ppd1a* alleles

show reduced differences in heading time between LD and SD and are designated as photoperiod

insensitive (PI), whereas plants carrying the *Ppd1b* allele are designated as photoperiod sensitive

(PS) [16, 17]. However, wheat PI lines still exhibit a significant acceleration of flowering under

LD relative to SD and, therefore, the wheat PI designation is used in this study to indicate a

reduced photoperiodic response rather than a lack of a photoperiodic response.

Under LD conditions, the wheat *PPD1* gene induces the expression of *FT1*, which encodes a

mobile protein that travels to the shoot apical meristem (SAM), where it promotes the expression

of the floral homeotic MADS box transcription factor *VERNALIZATION1* (*VRN1*) and the

synthesis of gibberellic acid (GA) [18]. The simultaneous presence of GA and *VRN1* results in

the up-regulation of *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1-1* (*SOC1*) and

LEAFY (*LFY*) genes, which are required for normal and timely spike development, stem

elongation and spike emergence or heading [18]. In photoperiod sensitive wheats grown under

SD, the SAM transitions to the reproductive stage, but stem elongation and spike development

proceed very slowly and heads fail to emerge or emerge extremely late [19]. These results

indicate that in wheat, the photoperiod pathway has a larger impact on the duration of spike

development and stem elongation than on the initial transition between the vegetative and

reproductive stages [18, 19]. The addition of gibberellic acid (GA) to PS spring wheat grown

under SD accelerates spike development, but only in lines that express *VRN1* under SD [18].

The photoperiod and vernalization pathways converge at the regulation of *FT1* in the leaves. The

role of *PPD1* as a LD flowering promoter is antagonized in winter wheats by the role of *VRN2* as

a LD flowering repressor [20, 21], which prevents the induction of *FT1* during the fall. Long

exposures to cold temperatures during the winter (vernalization) result in the induction of *VRN1*

both in the leaves and in the SAM [22]. In the leaves, the presence of *VRN1* prevents the

upregulation of *VRN2* in the spring [23], which results in the upregulation of *FT1* under LD. In

the SAM, the upregulation of *VRN1* initiates the transition of the SAM to the reproductive phase,

but spike development proceeds slowly until the LD induction of *FT1*. The arrival of *FT1* to the

SAM further up-regulates *VRN1* and induces GA accumulation, which results in accelerated

spike development and stem elongation [18].

In addition to the natural *PPD1* alleles (PS and PI), induced loss-of-function mutations in all

PPD1 homeologs (henceforth, *ppd1*) have been developed for both hexaploid [24] and tetraploid

wheat [19]. These *ppd1* plants show large delays in spike development and heading time under

LD, and spikes usually emerge very late, fail to emerge or are aborted under SD [19, 24]. The

flowering phenotypes of the wheat *ppd1* mutants are more drastic than those described for the

barley photoperiod insensitive allele *ppd-H1*, which has only amino acid changes and is likely a

hypomorphic allele rather than a complete loss-of-function allele [12]. It is important to

differentiate the photoperiod insensitive allele in barley that refers to delayed heading under LD,

from the PI allele in wheat that refers to accelerated flowering under SD.

CO1 and *CO2* overexpression results in barley suggest a role of these genes as flowering

promoters [10, 11]. However, overexpression under a constitutive promoter results in elevated

transcript levels and ectopic expression in tissues or developmental phases that are different from

those of the natural alleles, which complicates the interpretation of results. The use of loss-of-

function mutations can provide a better understanding of the role of these genes under more

natural conditions. In this study, we investigate the role of *CO1* and *CO2* on the photoperiod

response in wheat using loss-of-function mutants for both genes in the tetraploid spring wheat

variety Kronos. In addition, we combine loss-of-function mutations in *co1*, *co2* and *ppd1* to

study the effect of their genetic interactions on heading time and on the transcription profiles of

critical flowering genes. We also investigate the interactions among the proteins encoded by

these genes. Taken together, these analyses show complex genetic interactions among *CO1*, *CO2*

and *PPD1* on heading time that reflect similarly complex interactions at the transcriptional and

protein levels.

Results

Identification of *CO1* and *CO2* loss-of-function mutants in wheat

We identified a total of 153 mutations in the A and B homeologs of *CO1* and *CO2* in the Kronos

tetraploid TILLING population [25, 26] using primers and PCR conditions listed in Supporting

Information Table S1. Among the 153 mutations, 74 resulted in non-synonymous amino acid

changes, and five introduced early stop codons resulting in truncated non-functional proteins

(Table S2). These five truncation mutations were validated by sequencing and one for each

homeolog was selected to study the function of *CO1* and *CO2* (Fig 1). We backcrossed the

individual mutant lines with wild-type Kronos at least twice to reduce background mutations.

Fig 1: *CO1* and *CO2* gene structure and position of the selected TILLING mutations. Both *CO1* and

CO2 genes have two exons, which encode two B-BOX domains at the N-terminus and a CCT domain at

the C-terminus. The selected *co-A1* and *co-B2* mutations, located in the first exon after the two B-Box

domains, are predicted to generate premature stop codons, whereas the *co-B1* and *co-A2* mutants have

acceptor splice-site mutations that result in intron retention and premature stop codons that eliminate the

complete exon 2 including the CCT domain. The T4 numbers are the mutant identification. The schematic

gene structure is based on *CO-A1* from Kronos (GenBank MT043302).

We generated a *co1* loss-of-function mutant by crossing *co-A1* mutant T4-395 (Q242*) with *co-*

B1 mutant T4-1170 (Exon2 acceptor splice site mutation). To generate the *co2* loss-of-function

mutant, we crossed *co-A2* mutant T4-202 (Exon 2 acceptor splice site mutation) with *co-B2*

mutant T4-391 (Q193*). Finally, we crossed the *co1* and *co2* mutants and generated a *co1 co2*

double mutant. The stop codon mutations in *co-A1* and *co-B2* truncate the carboxyl terminal half

of the CONSTANS protein including the CCT domain (Fig 1), which is critical for protein

function in both Arabidopsis and cereals [27]. Likewise, the splice-site mutations selected in the

co-B1 and *co-A2* mutants result in intron retention and premature stop codons that also eliminate

the conserved CCT domain. Based on these large truncations we concluded that *co1 co2* has

loss-of-function mutations for all *CO1* and *CO2* homeologs.

Effect of *PPD1* photoperiod sensitive (PS) and insensitive (PI) alleles on heading time and

spike development

We transferred the four *co1 co2* mutations generated in Kronos-PI into a near-isogenic Kronos-

PS line [18] to study the effect of these mutations in two different backgrounds with large

differences in heading time under SD. To establish a baseline for comparison, we first describe

the effect of the two *PPD1* alleles in lines carrying the wild type *Co1* and *Co2* alleles (Fig 2A,

blue bars).

Under LD, heading time was 2.5 d earlier in Kronos-PI (52.3 ± 0.4 d) than in Kronos-PS ($54.8 \pm$

0.3 d). Although the difference was small, it was highly significant ($P < 0.0001$) and consistent

with previously published results [15]. Under SD, the differences between the two genotypes

were much larger (>66 d, $P < 0.0001$), with Kronos-PI heading at 113.2 ± 0.4 d and Kronos-PS

spikes failing to emerge before the experiment was terminated at 180 d. The earlier flowering of

Kronos-PI under SD reduced the difference between SD and LD in this genotype (61 d) relative

to Kronos-PS (>125 d, Fig 2A blue bars).

Fig. 2: Heading time of wild type, *co1, co2* and *co1 co2* mutants under LD and SD in Kronos-PI

(*Ppd-A1a*) and Kronos-PS (*Ppd-A1b*) backgrounds. (**A**) Averages are based on 15-19 plants in LD-PI

(2 experiments), 21-24 plants in LD-PS (3 experiments) and 8 to 14 plants in SD-PI and SD-PS (1

experiment). Red numbers indicate differences in heading time relative to the wild type (mut-WT), with *P*

values from a Dunnett test. Vertical arrows above the bars indicate that spikes have not emerged at 180 d

(end of the experiment). (**B-C**) Interaction graphs for *CO1* and *CO2* in Kronos-PS (*Ppd-A1b*) and Kronos-

PI (*Ppd-A1a*) backgrounds. The interaction was significant in B (*P* = 0.0052) but not in C (Table S3).

Error bars are standard errors of the means (henceforth, s.e.m). *= *P* < 0.05, **= *P* < 0.01, ***= *P* <

0.001, ns= not significant.

To determine if the observed differences in heading time were the result of changes in the

duration of the vegetative phase or the spike development and elongation phases, we dissected

SAMs of the main tillers from the different genotypes (from 4- to 14-leaf stage, S1 Fig). Under

LD, the SAMs from both Kronos-PI and Kronos-PS reached the early stages of spike

development by leaf four (W3-W3.25, Waddington scale [28]), the terminal spikelet stage

(W3.5) by leaf six, and headed at 52-55 d (S1 Fig). Under SD, the SAMs of both genotypes were

at the vegetative stage by leaf four and started to elongate by leaf six. After that point, the two

genotypes diverged, while Kronos-PI transitioned to the terminal spikelet and lemma elongation

stage at the eighth leaf, Kronos-PS reached the same developmental stage at leaf 12 (S1 Fig).

Effect of *CO1* and *CO2* on heading time in the presence of functional *PPD1* alleles

Loss of *CO1* and *CO2* function resulted in small but significant changes in heading time. The

co1 co2 double mutant headed 3 d earlier than the wild type in both Kronos-PI and Kronos-PS (*P*

< 0.0001) under LD, and 13.5 d earlier than the wild type in Kronos-PI under SD (*P* < 0.0001,

Fig 2A). In the Kronos-PS background under SD, spikes failed to emerge or were aborted in

most of the plants by the end of the experiment at 180 d, limiting our ability to determine

difference on heading time between the *co1* and *co2* mutant and wild type alleles under this

condition.

Even though the differences in heading time between *co1 co2* and wild type alleles were larger

under SD (13.5 d) than under LD (3 d), they were still smaller than the differences between the

PPD1 photoperiod sensitive and insensitive alleles under SD (>67 d), or the differences between

LD and SD in Kronos-PI (61 d) or Kronos-PS (>125, Fig 2A blue bars). These results indicate

that, in wheat, the *PPD1* alleles have a stronger effect on the photoperiodic response than the

CO1 and *CO2* genes.

Except for the *co2* mutant that headed later than the wild type in the Kronos-PS background

under LD (+1.5 d, *P* < 0.0001), all the other *co1* and *co2* single and double mutants headed

significantly earlier than their respective wild type controls under both LD and SD (Fig 2A).

These results suggest that under the tested conditions, *CO1* functions as a mild heading time

repressor in all backgrounds, and *CO2* as a mild heading time promoter or repressor depending

on the *PPD1* and *CO1* alleles present and the photoperiod (Fig 2A).

Long day experiments

To explore the interactions between *PPD1*, *CO1* and *CO2* under LD, we performed a factorial

ANOVA combining data from three experiments in Kronos-PS and two in Kronos-PI, using

experiments as blocks (nested within genotypes). This analysis showed highly significant effects

on heading time for *PPD1* (*P* < 0.0001), *CO1* (*P* < 0.0001) and *CO2* (*P* = 0.0004) and highly

significant two-way interactions for *PPD1* x *CO1* and *PPD1* x *CO2* (*P* < 0.0001, Table S3A).

The *CO1* x *CO2* interaction was not significant (*P* = 0.3437) but the three-way interaction was

significant (*P* = 0.0046, Table S3A), suggesting the possibility of different *CO1* x *CO2*

interactions in the different *PPD1* backgrounds that cancelled each other in the combined

analysis.

To test this possibility, we performed separate *CO1* x *CO2* factorial ANOVAs for Kronos-PS

and Kronos-PI. In the Kronos-PS background, *CO1* showed a stronger effect (*P* < 0.0001) on

heading time than *CO2* (*P* = 0.1448) and their interaction was significant (*P* =0.0052, Fig 2B,

Table S3B). By contrast, in the Kronos-PI background, *CO2* showed a stronger effect (*P* <

0.0001) than *CO1* (*P* = 0.0157) and the interaction was not significant. (*P* = 0.1934, Fig 2C,

Table S3C). In summary, these results indicate complex genetic interactions between *PPD1*,

CO1 and *CO2* in the regulation of heading time under LD.

Short day experiments

Under SD, the Kronos-PS lines failed to flower before the experiment was terminated (180 d), so

we were only able to study the interactions between *CO1* and *CO2* in the Kronos-PI background.

All four genotypes headed between 100 and 113 d, which is 51 to 61 d later than the same lines

under LD. A factorial ANOVA for heading time showed significant effects for both *CO1* and

CO2 ($P < 0.0001$) and a significant interaction between the two ($P = 0.0029$, Table S4). This

interaction was associated with the larger acceleration of heading time in the double mutant (13.5

d) than in the single *co1* (2.1 d) and *co2* (4.0 d) mutants (Fig 2A). This result indicates a stronger

effect of each mutation in the presence of the mutant allele of the other gene than in the presence

of the wild type allele (Table S4).

To see if the *co1* and *co2* mutations in the Kronos-PS background were associated with delays in

SAM differentiation or in spike development, we dissected apices for three plants from each

genotype every week from 4 to 11 weeks, and included wild type Kronos-PI as a control. Under

SD, all five genotypes reached the double-ridge stage (W2.5) by week 6 (Fig 3A-B). From there,

the Kronos-PI genotype developed more rapidly than the four Kronos-PS genotypes, which

showed similar developmental rates in the wild type and the *co1* and *co2* mutants (Fig 3B). By

week 11, the developing spikes in the Kronos-PI control were 3-13 cm long including awns, the

peduncles were 5 to 14 mm long, and the developing flowers were at the elongated stigmatic

branches stage (>W7.5). By contrast, all four Kronos-PS lines at the same time point showed no

peduncle elongation, produced spikes of less than 0.5 cm, and the flowers still had the stylar

canal open (<W6.0). These results indicate that the *co1* and *co2* mutations have a limited effect

on spike development and elongation in the Kronos-PS background under SD compared with the

strong effect of the *Ppd-A1a* in Kronos-PI.

Fig 3. Effect of *co1* and *co2* mutations on spike development in Kronos-PS under SD. Kronos-PI was

included as control. (**A**) Dissecting microscope images of developing apices at week 6 (double-ridge,

W2.5) and week 11 with a detail of their floral organs. a = anthers, elongated in Kronos-PI and primordia

in other genotypes, c = carpel and s = styles well developed in Kronos-PI and c&o = carpel extending

around the ovule in other genotypes (**B**) Developmental time course of SAM using the Waddington scale

from 4- through 11-week-old plants (n = 3). W1 = vegetative apex, W2 = double ridge, W3 = glume

primordium, W4 = stamen primordium, W5 = carpel extending round three sides of ovule, W6 = short

style primordia and stylar canal open, W7 = stigmatic branches just differentiating as swollen cells on

styles, W8 = stigmatic branches elongating, W9 = styles and stigmatic branches erect and stigmatic hairs

differentiating.

Effect of *CO1* and *CO2* on heading time in the absence of functional *PPD1* alleles

To investigate the effect of *CO1* and *CO2* in the absence of *PPD1*, we introgressed the *co1* and

co2 mutations into a Kronos line with loss-of-function mutations in both *PPD-A1* and *PPD-B1*

(*ppd1*) [19]. As before, we describe first the differences in SAM development and heading time

between Kronos-PS and *ppd1* mutants carrying the wild type *Co1* and *Co2* alleles to establish a

baseline for comparison (Fig 4A, blue bars).

Under LD, the plants with the *ppd1 Co1 Co2* allele combination headed much later (115.5 ± 15.3

d) than the Kronos-PS controls carrying the functional *Ppd1b* allele (52.5 ± 0.7 d, Fig 4A),

similar to results reported before [19]. The *ppd1* mutant plants showed a 79% reduction in

number of grains per spike relative to the Kronos-PS lines (Fig 4B-C). Apex dissections showed

a delayed transition to the double-ridge stage in the *ppd1* mutant (leaf 8 to 10) relative to

Kronos-PS (<leaf 4, S1 Fig). Similarly, the terminal spikelet was formed in Kronos-PS by leaf 6

but was not formed in *ppd1* by leaf 14, when the experiment was terminated (S1 Fig). Under SD,

spikes of both the *ppd1* mutant and Kronos-PS failed to emerge by 180 d. Dissection of the main

tillers from these plants showed that spike development in *ppd1* lagged behind that of Kronos-PS

(S1 Fig, leaf 12 and 14 and S2 Fig). These results indicate that *PPD1* is not essential for the

initial transition of the SAM to the reproductive stage, but in its absence, spike development and

stem elongation are severely compromised.

Fig 4. Phenotypic effects of *co1* and *co2* mutations in the absence of functional *PPD1* genes under

LD (**A**) Average days to heading in *ppd1* mutants carrying mutations in either *co1* or *co2*. Kronos-PS

wild type and *co1 co2* double mutants are included as controls. Floral transition and spike formation

occurred in all genotypes, but in the *ppd1 co1* double mutant floral development was delayed or arrested

and spikes failed to emerge before the experiment was terminated (arrow on top of the bar indicates

heading time > 180 d) (**C**) Number of grains per spike recovered from the same genotypes as in A. Error

bars are standard errors of the means calculated from 7 to 16 plants per genotype for flowering and from 4

to 8 plants for grain number.

The introgression of the *co2* mutation in the *ppd1* background had a limited effect on heading

time. Heading time of the double mutant *ppd1 co2* (119.1 ± 12.5 d) was only four days later than

ppd1, and the differences were not significant suggesting a limited effect of *CO2* on heading

time in the absence of a functional *PPD1*. The *ppd1 co2* double mutant showed an even more

severe reduction in grain production than the *ppd1* mutant (89% reduction relative to Kronos-

PS), but the differences were not significant.

The most important result was observed when we combined the *ppd1* and *co1* mutations. The

spikes from the *ppd1 co1* plants failed to emerge before the experiment was terminated at 180 d

(Fig 4A), and failed to produce any grains (Fig 4B-C). Even though we were not able to recover

the triple *ppd1 co1 co2* mutant, the delayed heading time of the *ppd1 co1* mutant relative to the

ppd1 and *ppd1 co2* mutants (>60d) is sufficient to conclude that *CO1* can accelerate heading

time in response to LD even in the absence of functional *PPD1* alleles.

Effect of mutations in *PPD1*, *CO1* and *CO2* on the transcriptional profiles of flowering

genes

To better understand the regulatory networks underlying the genetic interactions described

above, we characterized the effect of loss-of-function mutations in *PPD1, CO1* and *CO2* on the

transcription profiles of several flowering genes under LD and SD. Since Kronos-PI and Kronos-

PS have very similar heading times under LD but not under SD, we included both genotypes in

the SD experiment but only Kronos-PS under LD. To perform this experiment with a

manageable number of plants, we included the *co1 co2* double mutants in both experiments but

the individual *co1* and *co2* mutants only in the LD experiment We grew the plants in a growth

chamber under LD or SD and sampled the fully expanded fourth leaf every 4 hours throughout a

24-h period.

Under LD, *CO1* transcripts were significantly upregulated in both *co2* and *ppd1* mutants 12

hours after the start of the light phase (Zeitgeber time 12 or ZT12, Fig 5A), suggesting that these

two genes act as transcriptional repressors of *CO1*. The negative effect of *PPD1* on *CO1*

transcription was also observed in a separate experiment using older plants (six-week-old). In

leaf samples collected at ZT4 we observed a significant upregulation of *CO1* in the *ppd1* and

ppd1 co2 mutants relative to Kronos-PS (S3A Fig). No differences were detected between the

ppd1 and *ppd1 co2* mutants, which suggests that the loss of *co2* function did not affect *CO1*

expression in the absence of functional *PPD1* genes (S3A Fig). Under SD, *CO1* transcripts

showed a very different profile than under LD, with a major peak in the middle of the night

(ZT16) in Kronos-PS and at the end of the day (ZT8) in Kronos-PI, and a different profile in the

ppd1 mutant. These results indicate significant effects of *PPD1* on *CO1* transcription under both

LD and SD (Fig 5A).

Transcript levels of *CO2* were very low compared with *CO1* both under LD and SD, and showed

no significant differences among most genotypes (Fig 5B). We further confirmed these

observations in six-week old plants of Kronos-PS, *ppd1* and *ppd1 co1* plants sampled at ZT4 in

LD (S3B Fig).

Fig 5. Effect of loss-of-function mutations in *CO1*, *CO2* and *PPD1* on the transcriptional profiles of six wheat flowering genes in Kronos-PS and Kronos-PI (SD-only) plants grown under LD and SD. (**A**) *CO1*, (**B**) *CO2*, (**C**) *PPD1*, (**D**) *FT1*, (**E**) *VRN1*, and (**F**) *VRN2*. LD experiments included Kronos-PS, *co1*, *co2*, *co1 co2*, and *ppd1* mutants; whereas SD experiments included Kronos-PS, Kronos-PI, and their respective *co1 co2* and *ppd1* mutants. We grew the plants in a growth chamber under 16 h light at 22 °C and 8 h darkness at 17 °C for LD, and under 8 h light and 16 h darkness for SD (same temperatures). Fully expanded 4[th] leaves (Zadoks scale =14) were harvested at ZT0, ZT4, ZT8, ZT12, ZR16, and ZT20. The ZT0 value is the same as at ZT24. Shaded boxes represent night. Values are averages of four biological replicates and bars are SE of the means. Expression values are relative to *ACTIN* using the delta Ct method. Primers and primer efficiencies are listed in Table S5 (all amplify both A and B homologs of the respective genes).

Under LD, the *PPD1* gene in Kronos-PS showed a main peak in the middle of the light period

(ZT8) and a secondary peak at dusk (ZT16), which is consistent with the *PPD1* profile

previously described in photoperiod sensitive *T. monococcum* [29]. In the *co1* and *co1 co2*

mutants, the main peak was shifted 4 h earlier (ZT4), whereas in the *co2* mutants the main peak

was delayed 4 h (ZT12, Fig 5C). The ZT16 peak was significantly higher than the wild type in

the *co1 co2* double mutant but not in the single mutants (Fig 5C). The complex effects of the *co1*

and *co2* mutations on *PPD1* transcription profiles reflect the significant two- and three-way

genetic interactions among these three genes on heading time described above.

Under SD, the *PPD1* profiles were very different than under LD. Kronos-PS and its respective

co1 co2 mutant showed similar profiles with a peak at ZT4. By contrast, Kronos-PI showed a

very strong peak at ZT8 that was not observed in Kronos-PI with the *co1 co2* mutations (Fig 5C).

In addition, Kronos-PI and Kronos-PI *co1co2* showed significantly higher *PPD1* transcript levels

during the night relative to the other genotypes. The differences were largest at ZT16 (middle of

the night), which has been shown to be the optimum time for a short pulse of light (night break)

to accelerate heading time in Kronos- PS under SD [19]. The upregulation of *PPD1* during the

night under SD is consistent with previously published results in tetraploid wheat lines carrying

the *Ppd-A1a* allele (PI) relative to F_8-near-isogenic sister lines carrying the *Ppd-A1b* allele (PS)

[17].

The *FT1* profiles correlated well with heading time both in LD and SD experiments. The earlier

heading time of Kronos-PS *co1 co2* under LD was paralleled by significantly higher transcript

levels of *FT1* at the end of the day than all other genotypes (Fig 5D). Under SD, *FT1* transcript

levels were upregulated only in Kronos-PI genotypes, which were the only ones that headed

under this condition (Fig 2). Between the two Kronos-PI lines, the *co1 co2* mutant showed

higher *FT1* transcript levels and flowered earlier (13 d, Fig 2) than the wild type (Fig 5D). We

observed a significant reduction of the transcript levels of *FT1* in the *ppd1* mutant relative to

wild type, which was also consistent with its delayed heading time under LD (Fig 4) and its

inability to head under SD.

The transcript levels of *VRN1* correlated well with the transcript levels of *FT1* under LD but not

under SD (Fig 5E). Under SD at dawn, *VRN1* transcript levels were higher in *ppd1* relative to the

other genotypes, and was generally higher in Kronos-PS than in Kronos-PI genotypes. No

significant differences in *VRN1* expression were detected between the *co1 co2* mutants and their

respective Kronos-PS and Kronos-PI isogenic lines at that time point. The Kronos-PI genotypes

showed limited variation during the day compared with the Kronos-PS and *ppd1* mutant (Fig.

5E).

The *VRN2* locus includes two genes, but since *ZCCT1* is not functional in tetraploid wheat [30],

we present only expression data for *ZCCT2*. Under LD, the transcriptional profiles of *PPD1* and

ZCCT2 were similar for the wild type Kronos-PS but not for the different mutants (Fig 5). We

observed a small but not significant increase of *ZCCT2* transcript levels in the *ppd1* mutant and a

significant downregulation in the *co1* and *co1 co2* mutants. The interpretation of these results is

complicated by the effects of these mutations on *VRN1*, which is a known repressor of *VRN2*

under LD [23] (see Discussion section). Under SD, *VRN2* was upregulated only in the Kronos-PI

lines, with higher transcript levels in the presence of the functional *CO1 CO2* genes than in the

co1 co2 mutants.

Effect of *VRN1* on the transcriptional regulation of *CO1* and *CO2* under LD

To explore the potential role of *VRN1* in the transcriptional regulation of *CO1* and *CO2*, we used

Kronos *vrn-A1* and *vrn-B1* single mutants and the *vrn-A1 vrn-B1* double mutant (henceforth

vrn1) developed previously [23]. The *vrn-B1* mutant has a functional allele of *VRN-A1* that

confers a dominant spring growth habit, whereas the *vrn-A1* mutant has the ancestral allele of

VRN-B1 that confers a recessive winter growth habit. The *vrn1* double mutant has a delayed

heading time and a residual response to vernalization [23].

CO1 showed significantly higher transcript levels in the late flowering *vrn-A1* and *vrn1* mutants

than in the rapidly flowering *vrn-B1* mutant, both in vernalized and un-vernalized plants (Fig 6

A-B). In all genotypes, *CO1* was downregulated to very low levels in the flag-leaf at heading

time. In the *vrn1* mutant, we observed a slight decrease of *CO1* transcript levels during

vernalization followed by a rapid upregulation when we moved the plants back to room

temperature (RT). By contrast, in the single *vrn-A1* and *vrn-B1* mutants, which still have one

functional copy of *VRN1*, *CO1* transcripts stayed low after the plants were returned to RT (Fig 6

A). From previous studies, we know that *VRN1* transcript levels increase slowly during LD

vernalization and then rapidly when plants are moved to RT and start to flower [23]. Taken

together, these results suggest that *VRN1* acts as a transcriptional repressor of *CO1*.

Fig 6. Effect of mutations in Kronos *vrn-A1* (winter growth habit), *vrn-B1* (spring growth habit) and *vrn-A1 vrn-B1* (*vrn1*, very late heading) on the transcriptional profiles of *CO1* and *CO2*. All mutants are homozygous and in the Kronos-PI background. (**A-B**) *CO1* and (**C-D**) *CO2* transcription profiles. **A & C**) More recently expanded leaves from three weeks old plants were sampled before vernalization (0w), at 3w and 6 w vernalization (4°C), 2 weeks after return to room temperature (RT= 22°C day and 18°C) and at heading time (HT), which was different for each genotype. **B & D**) Same genotypes but grown at RT without vernalization. Points are averages of eight plants and error bars are standard errors of the means. Means with different letters are significantly different in Tukey test for that time point (*P* < 0.05). Note that the upregulation of *CO1* after the return of the plants from vernalization occurred only in the *vrn1* mutant that has no functional copies of *VRN1*.

CO2 transcripts were downregulated during vernalization and showed a sharp increase when we moved the plants to RT. However, in contrast to *CO1*, this increase was observed in all three genotypes (Fig 6C-D). After this increase, *CO2* transcript levels remained high in the flag leaves during heading and showed no significant differences between the *vrn1* and the spring *vrn-B1*

mutant (Fig 6C-D). These results indicate that *VRN1* has a smaller effect on the regulation of

CO2 than on the regulation of *CO1*.

Effect of photoperiod and phytochromes on the transcriptional regulation of *CO1* and *CO2*

The strong upregulation of *CO1* in the *ppd1* mutant indicated that *PPD1* is a transcriptional

repressor of *CO1* (Fig 4A). Since phytochromes *PHYC* [15] and *PHYB* [31] are both required for

the LD transcriptional activation of *PPD1*, we explored the effect of mutations in these two

phytochromes on the expression profiles of *CO1* and *CO2*. To do this, we used recently

published RNAseq studies for leaf samples (4th leaf, ZT 4h) from loss-of-function mutants *phyB*

and *phyC* in Kronos-PI under both LD [31] and SD conditions [32].

A factorial ANOVA for transcripts per million reads (TPM) using photoperiod (LD and SD) and

genotype (WT, *phyC* and *phyB*) as factors, revealed significant differences between SD and LD

for all four genes (*CO-A1*, *CO-B1*, *CO-A2* and *CO-B2*, S4 Fig). We also detected difference

between genotypes and genotype x photoperiod interactions for the *CO1* homeologs but not for

the *CO2* homeologs. The significant interaction was associated with the LD downregulation of

CO1 homeologs in the wild type and their upregulation in the *phyB* and *phyC* mutants (S4 Fig).

Both *CO2* homeologs were upregulated by LD and showed no significant differences between

wild type and *phy* mutants. *CO-B2* showed significantly higher transcript levels than *CO-A2* for

all genotypes (S4 Fig), whereas for *CO1* the transcript levels and the differences were larger in

CO-A1 relative to *CO-B1*.

Protein-protein interactions among CO1, CO2, PPD1, VRN2, PHYC and PHYB

To test if the interactions among *CO1*, *CO2* and *PPD1* observed at the genetic and

transcriptional levels were associated with interactions at the protein level, we performed yeast-

two-hybrid (Y2H) and split yellow fluorescent protein (YFP) assays among the three proteins. In

a previous study, we reported positive Y2H interactions between the CCT domains of CO1 and

CO2, but not between these two proteins and a truncated version of PPD1 including the CCT

domain [33]. In this study, we used a complete PPD1 protein and detected strong Y2H

interactions between PPD1 and both CO1 and CO2. The Y2H interaction between PPD1 and

CO1 was stronger than between PPD1 and CO2 (Fig 7A).

We further validated the interactions between CO1 and CO2 (Fig 7B) and between CO1 and

PPD1 (Fig 7C-D) using split-YFP assays in wheat protoplasts. We initially failed to detect the

split-YFP interaction between CO1 and PPD1 in Kronos, so we used protoplasts extracted from

an early flowering loss-of-function mutant for *elf3* in Kronos, which has increased levels of

PPD1 transcripts [29, 34]. We were able to detect the CO1-CO2 and CO1-PPD1 interactions

(but not the CO2-PPD1) by split-YFP in the *elf3* mutant when protoplasts were exposed to light.

For both CO1 and PPD1, we detected the proteins by Western blots only after we exposed the

protoplasts to light but not when we incubated them in the dark after transfection, suggesting that

light is required to stabilize these proteins.

Fig 7. Protein interactions. (**A**) Yeast-two-hybrid (Y2H) interactions between PPD1 and CO1 or CO2. Left panel: SD medium lacking Leucine and Tryptophan (SD-L-W) to select for yeast transformants containing both bait and prey. Right panel: interaction on SD-L-W-H-A medium (no L, W, H and A). Dilution factors = 1, 1:10 and 1:100. (**B-D**) Split-YFP in wheat protoplast. (**B**) Positive nuclear interactions between CO1 and CO2. (**C-D**) Positive nuclear interactions between CO1 and PPD1 in two different YFP-N/YFP-C configurations. (**E**) Summary of Y2H interactions among wheat CO1, CO2, PPD1, VRN2, PHYB, PHYC, and ELF3 from S5 Fig. Green lines are based on published results from *Brachypodium* [35] and violet lines on published results from wheat [24]. Thick lines indicate interactions validated by split-YFP in wheat protoplasts (blue) or *in vitro* coIP (violet). Curved arrows indicate self-dimerization**.**

Finally, we tested pairwise interactions among CO1, CO2, PPD1, VRN2, PHYC and PHYB

proteins by Y2H assays (Figs 7E and S5 Fig). In addition to interacting with CCT-domain

proteins PPD1 (Fig 7A) and VRN2 [33], CO1 and CO2 showed positive interactions with both

PHYC and PHYB (Fig 7E and S5 Fig) in the Y2H assays. The interaction between CO1 and

PHYC was stronger than between CO1 and PHYB, whereas the interaction between CO2 and

PHYB was stronger than between CO2 and PHYC. Both phytochromes showed positive

interactions with VRN2 but not with PPD1 (Fig 7E and S5 Fig). In addition, we showed that the

full length PPD1 protein did not interact with VRN2 in Y2H (S5 Fig). CO2, VRN2, PHYC and

PHYB were able to form homodimers [15, 33] but not CO1 or PPD1 (S5 Fig). The diagrammatic

representation of these interactions reveals an extensive and complex network of protein-protein

interactions, with CO1 and CO2 providing a connection between the LD flowering promoter

PPD1 and the LD flowering repressor VRN2 (Fig 7E).

Discussion

Genetic interactions among *CO1*, *CO2*, *PPD1* and *VRN2* modulate photoperiodic responses

in grasses

Long day grasses

Natural variation in wheat heading time in response to photoperiod has been mapped mainly to

the three *PPD1* homeologs [16, 17, 36]. Similarly, a genome wide association study of 220

spring barley varieties found that most of the variation in heading time (31%) was associated

with the *PPD1* locus and variation in the *CO1* locus had a limited contribution (<5%) [37].

These results agree with the limited effect on heading time detected in this study in the Kronos

co1, *co2*, and *co1 co2* mutants compared with the large differences detected between *PPD1*

alleles (Fig 2) and mutants (Fig 4). Even when both *CO1* and *CO2* homeologs were replaced

with truncated mutants (*co1 co2*), plants headed only three days earlier than the wild type under

LD and 13 days earlier under SD (Fig 2). By contrast, loss of function mutations in *PPD1*

delayed heading time approximately two months in LD (Fig 4), and differences between the

Ppd-A1a and *Ppd-A1b* alleles were even larger under SD (Fig 2).

The effect of the *co1 co2* mutations on the acceleration of heading time was small, but highly

significant ($P < 0.0001$) and consistent across experiments under LD in Kronos-PI and Kronos-

PS and under SD in Kronos-PI (Fig 2). This was an unexpected result because transgenic barley

plants overexpressing *CO1* [10] or *CO2* [11] flowered earlier than the wild type under both LD

and SD. These transgenic results had led to the general assumption that *CO1* and *CO2* operate as

flowering promoters in the temperate grasses. However, our results show that when *CO1* and

CO2 are expressed under their native promoters in their expected tissues and developmental

stages (and in the presence of functional *PPD1* and *VRN2* alleles) they operate mainly as mild

LD and SD flowering repressors rather than as flowering promoters (except for *CO2* in Kronos-

PS under LD, Fig 2).

This inconsistency may be generated by the elevated and ectopic expression levels of *Ubi::CO1*

and *Ubi::CO2* in the transgenic plants, but also by differences in the genetic backgrounds. The

two transgenic barley lines were developed in the variety Golden Promise [10], which carries the

ppd-H1 allele with reduced sensitivity to LD and does not have any functional copy of *VRN2*. By

contrast, Kronos has functional *PPD1* and *VRN2* genes. Our results show that in a Kronos line

with no functional copies of *PPD1* (*ppd1*), *CO1* accelerates heading time under LD (Fig 4)

similarly to the overexpression of *CO1* in Golden Promise. In addition, the downregulation of

VRN2 in the Kronos *co1 co2* mutant may have contributed to the earlier heading time of these

mutants (Fig 2), whereas no interactions with *VRN2* are possible in Golden Promise.

The previous results suggest the existence of complex genetic interactions among the

photoperiodic genes, a hypothesis that is also supported by significant digenic interactions in the

factorial ANOVA including *CO1*, *CO2* and *PPD1* (Table S3). The three-way interaction among

the three genes was also significant, which reflects the differences in digenic interactions

between *CO1* and *CO2* in Kronos-PS (significant) and Kronos-PI (not significant, Table S3).

The contribution of the *PPD1* x *CO2* interaction to the total sum of squares (9%) was more than

two-fold higher than the main *CO2* effect (4%), which suggests that a large part of *CO2* effect on

heading time was mediated by its modulation of the *PPD1* function (Table S3). Strong genetic

interactions between *CO1* and *PPD1* on heading time have also been reported in a barley

biparental population segregating for *Ubi::CO1* and the *ppd-H1* mutation [10] and in two

genome wide association studies (GWAS) in spring barley [37, 38]. In both GWAS, significant

effects of *CO1* on heading time were detected among the lines carrying the *ppd-H1* allele but not

among the lines carrying the functional wild type *Ppd-H1* allele [37, 38]. These strong genetic

interactions indicate that the effect of a particular photoperiod gene depends strongly on genetic

background, a result that has also been reported in the SD grasses as described in the next

section.

Although the effect of *CO1* on heading time is limited in the presence of functional *PPD1*

alleles, in the *ppd1* mutant background the presence of *CO1* was required for heading under LD

(Fig 4). It has been previously shown that hexaploid wheat plants with no functional *PPD1* genes

(but with wild type *CO1* alleles) were able to head under LD, and that these plants express *FT1*

although at lower levels [24]. These levels should be sufficient to send enough florigen to the

SAM to up-regulate both *VRN1* and GA biosynthesis and the subsequent induction of *LFY* and

SOC1, which are critical for timely spike development and stem elongation [18]. We hypothesize

that the inability of the spikes to emerge in the *ppd1 co1* mutants grown under LD or in the

Kronos-PS or *ppd1* mutants grown under SD, is associated with insufficient production of *FT1* in

the leaves, which results in limited levels of GA, *LFY* and *SOC1* in the SAM. Addition of

exogenous GA to photoperiod sensitive *T. monococcum* genotypes that express *VRN1* under SD

(due to a deletion in the *VRN1* promoter) has previously been shown to partially rescue the arrest

in spike development and stem elongation under SD [18].

Short day grasses

In most sorghum genotypes, *SbCO1* operates as a flowering promoter in SD and as a flowering

repressor in LD [39] and *SbPRR37* as a flowering repressor in LD [14, 40]. By contrast, the

orthologous *PPD1* gene in barley and wheat functions as a LD heading promoter [12, 16, 19]

(Fig 4). In spite of the opposite roles, the genetic interactions between *PRR37/PPD1* and *CO1*

showed some similarities. In a sorghum population segregating for *SbCO1* and *SbPRR37*, *SbCO1*

showed a large effect on heading time among the plants carrying the non-functional *Sbprr37*

allele (48% of the variation), but had no effect among the plants carrying the functional *SbPrr37*

allele [41]. This is similar to the GWAS results in spring barley described above [37, 38] and

consistent with the stronger effect of the *co1* mutation in the *ppd1* mutant than in the sister lines

carrying functional *PPD1* alleles described in this study.

Similar to sorghum, in most rice genotypes, *OsPRR37* (=*Hd2*) works as a LD flowering

repressor [13] and *OsCO1* (=*Hd1*) as both a LD flowering repressor and a SD flowering

promoter. Interestingly, transformation of a rice line deficient in *OsCO1* function with a wheat

CO2 genomic clone accelerated flowering in SD and delayed flowering in LD, complementing

the *OsCO1* function [42]. Since our results indicate that wheat *CO2* functions as a SD flowering

repressor (Fig 2A), the rice result suggests that the function of the wheat CO2 protein was altered

when it was expressed in rice. The opposite change was observed when a late-flowering

Arabidopsis *prr7* mutant was transformed with rice *OsPRR37* driven by the Arabidopsis *PRR7*

promoter. When expressed in Arabidopsis, *OsPRR37* functioned as a LD promoter of flowering

in spite of its normal role as a LD repressor of flowering in rice. Based on these results, it is

tempting to speculate that the different roles of *PRR37/PPD1* and *CO2* in SD and LD grasses

may be the result of differences in the balance of multiple epistatic interactions among the genes

involved in the photoperiodic response, rather than a difference in the specific function of their

encoded proteins.

This hypothesis is also supported by changes in gene function reported in sorghum and rice

plants carrying loss-of-function mutations in both *PRR37* (*prr37*) and *GRAIN NUMBER, PLANT*

HEIGHT, AND HEADING DATE7 (*ghd7,* a *VRN2* ortholog [43]). In the *prr37 ghd7* sorghum

and rice mutants, *CO1* function changed from a flowering repressor to a flowering promoter in

LD, while maintaining its usual role as a flowering promoter in SD [41, 44, 45]. In addition, in

most rice genotypes carrying the *ghd7* mutation *OsPRR37* function changes from a SD promoter

to a SD repressor of heading, while maintaining its role as a LD flowering repressor [46].

Genetic interactions between *VRN2* (*GHD7*) and both *CO1* (*Hd1*) [47] and *CO2* [11] have been

reported also in barley, suggesting that similar mechanisms may exist in the temperate cereals.

Taken together, these results indicate that genetic interactions between *PPD1/PRR37*,

VRN2/GHD7 and *CO1/CO2* play a central role in the regulation of heading time in response to

photoperiod in both LD- and SD-grasses and that, therefore, the effect of particular genes may

vary in different genetic backgrounds.

Complex interactions at the transcriptional level contribute to the observed genetic

interactions on heading time

The significant upregulation of *CO1* in the *ppd1* mutant under LD (Fig 5A and S3A Fig)

suggests that *PPD1* may function as a transcriptional repressor of *CO1* under this condition. This

hypothesis is supported by the upregulation of *CO1* in the *phyB* and *phyC* mutants (S4 Fig),

where *PPD1* transcription is strongly downregulated [15, 31], and by the negative correlation

between *PPD1* and *CO1* transcript levels in both *ppd1* triple mutants of hexaploid wheat

'Paragon' under LD [24], and photoperiod sensitive and insensitive isogenic lines of Paragon

under SD [48]. Under SD, however, the *CO1* transcription peak in the *ppd1* mutant was slightly

lower than the peaks of the genotypes with functional *PPD1* alleles, which showed an 8 h shift

(Fig 5A). These results indicate that *PPD1* has an effect on *CO1* expression under SD, but not as

a transcriptional repressor as in LD (Fig. 5A). Under SD, both *CO1* and *PPD1* have a

transcription peak at ZT8 (Fig 5A and 5C).

Since *PPD1* induces the expression of *VRN1* under LD, and *VRN1* represses *CO1* transcription

(Fig 6), we cannot rule out the possibility that the effect of *PPD1* on *CO1* transcription is

mediated, at least in part, by *VRN1*. When Kronos *vrn-A1*, *vrn-B1* and *vrn1* mutants were

returned to room temperature after vernalization, *CO1* was rapidly upregulated in the leaves of

the *vrn1* mutant but not in the other three genotypes with functional *VRN1* copies (Fig 6). An

identical expression profile has been described before for *VRN2* in the same mutants [23],

indicating that *VRN1* functions as a LD transcriptional repressor of both *CO1* and *VRN2*.

CO2 transcript levels were not significantly affected by the different *PPD1* alleles under LD or

SD (Fig 5B) or by the loss-of-function mutants of *VRN1* in a LD experiment (Fig 6C-D).

Similarly, no changes in *CO2* transcription were observed between Paragon-PI and Paragon-PS

lines grown under SD [48], between Kronos-PI and the *phyB* or *phyC* mutants (S4 Fig), or

between Kronos-PS and *ppd1* mutants (S3B Fig). However, mutations in *co2* affected the LD

diurnal transcriptional profiles of all the flowering genes presented in Fig 5. Under LD, *CO1*

transcript levels were more than 2-fold higher in the *co2* mutant than in the Kronos-PS control at

the ZT12 peak (Fig 5A), suggesting that CO2 functions as a negative transcriptional regulator of

CO1. This is also supported by the downregulation of *CO1* transcripts in a barley line

overexpressing *CO2* [11]. The effect of *CO2* on *CO1* transcription may contribute to the

significant genetic interaction on heading time observed between *CO1* and *CO2* in this study

(Table S3b).

Overexpression of *CO1* and *CO2* in winter barley was associated with the upregulation of *VRN2*,

both under LD and SD, where *VRN2* is not normally expressed [11]. A similar upregulation of

VRN2 has been reported in *Brachypodium* plants overexpressing *CO1* under LD [49]. By

contrast, knock-down of *CO1* expression in *Brachypodium* [49] or loss-of-function mutants in

co1, *co2*, or *co1 co2* in Kronos in our study (Fig 5F) were associated with the down regulation of

VRN2 transcripts in LD (and SD in Kronos PI). The regulation of *VRN2* by CO1 may be direct,

as it was recently shown in rice that a complex formed by Hd1 (CO1), GHD8 and OsHAP5b can

bind the promoter of *GHD7* (*VRN2*) and activate its transcription [50]. Taken together, these

results are consistent with *CO1* and *CO2* being promoters of *VRN2* transcription in the temperate

grasses, a role that is likely more important during the fall than in the spring, when high

expression of *VRN1* results in the downregulation of both *CO1* and *VRN2*.

Photoperiod and vernalization genes converge on the regulation of *FT1*, whose transcript levels

are well correlated with heading time [11, 51]. Here we also observed a good correlation

between *FT1* transcript levels and heading time both under LD and SD (Fig 2 and 5D). Since

PPD1 is a critical LD transcriptional promoter of *FT1* [12, 35]*,* it was not surprising to see

almost undetectable levels of *FT1* in the *ppd1* mutant, which were correlated with low *VRN1*

transcript levels and late heading time under LD (Fig 5D-E). The low transcript levels of *VRN1*

in the *ppd1* mutant may have contributed to the slight increase in *VRN2* transcript levels

observed in this mutant (Fig 5F), because *VRN1* is a known transcriptional repressor of *VRN2* in

wheat under LD [23, 52].

The transcriptional profiles of *VRN1* under SD were unexpected. The increased levels of *FT1* in

the two Kronos-PI genotypes (Fig 5D), were not reflected in the upregulation of *VRN1* (Fig 5E).

This suggests that either some factors required for the *VRN1* up-regulation by *FT1* [53] are

missing, or the presence of one or more additional SD repressors. If has been previously shown

that deletions in a CArG box located in the *VRN1* promoter in photoperiod sensitive *T.*

monococcum are associated with increased *VRN1* transcript levels under SD, suggesting the

existence of a SD repression mechanism of *VRN1* transcription [20]. The upregulation of *VRN1*

in the *ppd1* mutant under SD (Fig 5E) suggests the possibility that *PPD1* may be involved in this

SD repression mechanism. A strong SD upregulation of *VRN1* has also been reported in *phyC*

and *phyB* mutants of Kronos-PI, which also showed a significant down-regulation of *PPD1* [32]*.*

Under LD, the high transcript levels of *VRN2* in the *ppd1* mutant indicates that *PPD1* is not

required for the LD expression of *VRN2* (Fig 5F). Under SD, however, *VRN2* was expressed only

in Kronos-PI and Kronos-PI *co1 co2* (Fig 5F), which are the only genotypes that show a

significant increase in *PPD1* transcripts during the night (ZT16, Fig 5C). Since a short pulse of

light in the middle of the night induces *PPD1* transcription and is sufficient to greatly accelerate

Kronos-PS flowering in SD [19], we speculate that the higher transcript levels of *PPD1* during

the night in the two Kronos-PI lines (particularly at ZT16) may be important for the acceleration

of heading time under SD. The positive role of *PPD1* on the upregulation of *VRN2* under SD has

also been reported in hexaploid wheat lines carrying the photoperiod insensitive *Ppd-D1a* allele

but not in sister lines carrying the photoperiod sensitive *Ppd-D1b* allele [54], and in young winter

barley plants [11].

The role of *PPD1* on the upregulation of *VRN2* and *FT1* is also reflected in their similar

transcription profiles in Kronos-PS under LD (Fig 5C, D and F, blue lines). However, those

similarities disappear in the different *CO1* and *CO2* mutants, which suggests that the effect of

PPD1 on *FT1* and *VRN2* under LD is modulated by *CO1* and *CO2*. Under SD, the altered

expression of *PPD-A1a* (promoter deletion) in Kronos-PI has two opposing effects, the up-

regulation of *FT1* that promotes flowering (Fig 5D) and the up-regulation of *VRN2* that represses

flowering (Fig 5F). This may contribute to the lower *FT1* transcript levels in Kronos-PI (higher

VRN2) relative to Kronos-PI *co1 co2* (lower *VRN2* levels) under SD (Fig 5D and F).

To summarize the dynamics of these transcriptional interactions during the seasons, we propose

the following working model for the ancestral photoperiod sensitive winter wheats. In the fall,

PPD1, *CO1* and *CO2* promote the expression of *VRN2* in the leaves, which prevents flowering

by repressing the transcriptional induction of *FT1*. During the winter, the increase in *VRN1*

expression in the apex [22] promotes the initial stage of reproductive development whereas its

upregulation in the leaves results in the gradual downregulation of both *VRN2* and *CO1*, which

facilitates the promotion of *FT1* by *PPD1* and the acceleration of spike development and stem

elongation in the spring.

Protein-protein interactions add an additional level of regulation to wheat heading time

In addition to the complex transcriptional interactions described above, the proteins encoded by

the photoperiod genes form an intricate network of protein-protein interactions that contribute an

additional layer of complexity to the photoperiodic regulation of heading time. In Arabidopsis,

the central photoperiod gene *CO* is regulated at the transcriptional and posttranscriptional level.

A complex transcriptional regulation is required to establish the presence of *CO* mRNA during

the evening of a long day, when light can stabilize the CO protein, resulting in the activation of

FT transcription [27].

Several PRR proteins promote flowering in Arabidopsis by interacting with and stabilizing the

CO protein during the day specifically under LD, and indirectly through their roles in the

circadian clock [55]. Mutations in *PRR* genes delay Arabidopsis flowering under LD but not

under SD (similar to *PPD1* in wheat), with double mutants showing a stronger delay [56]. The

late-flowering Arabidopsis *prr7* mutant can be complemented by rice *OsPRR37* [57], suggesting

some conserved functions in rice and Arabidopsis. In this study, we detected interactions

between wheat PPD1 full-length protein and both CO1 and CO2 in Y2H and validated the CO1-

PPD1 interactions by split-YFP (Fig 7). These results suggest that the interactions between CO

and PRRs likely precede the split between monocot and dicot plants.

In spite of these similarities, there are also important differences between the grass *PPD1/PRR37*

gene and the homologous *PRR* genes from Arabidopsis. The Arabidopsis *PRR* genes are part of

the circadian clock and their mutants affect the expression of circadian clock genes and their

multiple downstream targets. By contrast, the natural mutation in the barley photoperiod

insensitive *ppd-H1* allele did not affect diurnal or circadian cycling of barley clock genes [58].

This result suggests that after the duplication that generated *PRR37* and *PRR73* in the grasses,

which is independent of the Arabidopsis duplication that originated *PRR3* and *PRR7*,

PRR37/PPD1 acquired a more specialized role in the photoperiodic response with a more limited

role in the circadian clock.

In addition to its interactions with the LD flowering promoter PPD1, CO1 and CO2 can form

heterodimers with the LD flowering repressor VRN2, which is also a member of the CCT

domain protein family [33]. We hypothesize that the interactions with VRN2 might be relevant

to *CO1* and *CO2* function because loss-of-function mutation in *ghd7/vrn2* result in reversals of

functional in *OsPRR37* (*Hd2*) [46] and *OsCO1* (*Hd1*) in rice [44, 45], and *SbCO1* in sorghum

[41]. In rice, GHD7 binds to the transcription-activating domain of OsCO1, probably weakening

or blocking the transcriptional activation of *Ehd1* and *Hd3a* [45]. We have initiated crosses

among Kronos lines with mutations in *vrn2* [30], *co1*, *co2*, and *ppd1* to test these interactions in

wheat.

Recent studies in rice showed that another CCT-domain protein, the NUCLEAR

TRANSCRIPTION FACTOR Y subunit B-11 (NF-YB11 = HEME ACTIVATOR PROTEIN or

HAP-3H = *Ghd8*) has large effects on heading time and significant genetic interactions with

OsGHD7, *OsCO1* and *OsPRR37* [46]. We have previously shown that in addition to interacting

with each other, CO2 and VRN2 interact with the same subset of eight NUCLEAR FACTOR-Y

(NF-Y) transcription factors (Fig 7E) [33], suggesting an additional functional connection

between these proteins. This connection through the NF-Y proteins seems to less extensive

between CO1 and VRN2, where only one shared NF-Y interactor has been identified so far (Fig

7E) [33]. Members of the NF-Y family have been shown to be involved in the regulation of

flowering through interactions with CCT domain proteins CO and COL in other plant species

[59-61]. Using yeast-three-hybrid (Y3H) assays, we previously showed that VRN2 competes

with CO2 for interactions with the NF-Ys, and that mutations in three conserved arginine (R)

residues within the CCT domain of VRN2 reduce the strength of the protein interactions and

competition with CO2 [33]. Since these same mutations abolish or reduce VRN2 function as a

LD flowering repressor, we hypothesize that the protein-protein interactions involving VRN2-

CO2-NF-Y CCT domain could be relevant for their function [33].

Phytochromes PHYB and PHYC are involved in the transcriptional regulation by light of both

PPD1/PRR37 and *VRN2* in wheat [15, 31], *Brachypodium* [62], rice [63] and sorghum [40]. In

addition, OsPHYA, OsPHYB and OsGI proteins have been shown to interact directly with

OsGHD7, regulating protein stability [64]. We observed similar interactions between VRN2 and

both PHYB and PHYC in wheat. These two phytochromes also showed Y2H interactions with

CO1 and CO2, but not PPD1 (Fig 7E and S5). Additional experiments will be required to test if

these phytochrome interactions can stabilize the VRN2, CO1 and CO2 proteins in wheat as they

stabilize GHD7 in rice [64] or CO in Arabidopsis [65].

In summary, we found that CO1 and CO2 proteins can physically interact with each other and

with both PPD1 (a central LD flowering promoter) and VRN2 (a central LD flowering

repressor), providing a physical link between these central photoperiodic proteins (Fig 7E).

Among the 15 pairwise interactions tested among PHYB, PHYC, PPD1, CO1, CO2 and VRN2

80% showed positive interactions in Y2H assays indicating a complex and dense network of

protein interactions. We hypothesize that this network of protein interactions provides an

additional layer of molecular mechanisms that result in the complex genetic interactions in

heading time detected in this study.

Two interacting photoperiod pathways in grasses

A central conclusion of our study is that *PPD1* can perceive the differences in photoperiod and adjust heading time accordingly even in the absence of any functional copy of *CO1* or *CO2*. Similarly, *CO1* can accelerate heading time under LD in the absence of a functional copy of *PPD1*. This is also true in rice, where heading time differences between SD and LD are observed for *OsCO1* alleles in *Osprr37* mutants, and for *OsPRR37* alleles in rice *Osco1* mutants [46]. These results suggest that grasses have either two photoperiod pathways, each of them capable of perceiving the length of the day/night and of regulating heading time, or a single point of perception of the differences in photoperiod upstream of *PPD1* and *CO1*, which can then transmit the signal to both genes. Recently, it has been suggested that the dark reversion of the Pfr form of PHYC may be involved in measuring night length (molecular hourglass model), by modulating the accumulation of the ELF3 protein, a direct transcriptional repressor of *PPD1* [35]. This mechanism is different from the external coincidence model proposed for Arabidopsis [66]. Although we favour the hypothesis of separate mechanisms for *PPD1* and *CO1* perception of photoperiod, we currently cannot rule out the possibility that, in the temperate grasses, ELF3 mediates both the *PPD1* and *CO1* responses to photoperiod.

Another important conclusion from our study and from previous studies in sorghum and rice [40, 44, 46], is that the *CO* and *PRR37/PPD1* photoperiod pathways are extensively interconnected. Changes in the balance of these interactions seem to play a central role in the diversity of photoperiod responses in grasses, as suggested by changes in function observed when photoperiodic genes are swapped between LD and SD species or when the same genes are expressed in different mutant backgrounds in its own species. The different expression profiles of the photoperiodic genes in LD and SD also show that these interactions are dynamic and

affected by photoperiod. We hypothesize that the complex feedback transcriptional regulatory

loops and transcriptional interactions among photoperiod genes along with the intricate

interactions among their encoded proteins contribute to the complex genetic interactions in

heading time observed in this study and to the diversity of photoperiodic responses in grasses.

Materials and methods

Plant materials

The materials used in this study included the tetraploid wheat variety Kronos (*Triticum*

turgidum ssp. *durum*), which carries a functional and dominant *Vrn-A1* allele for spring growth

habit, a functional but recessive *vrn-B1* allele for winter growth habit, a functional *Vrn-B2* long

day repressor (*ZCCT-B2a* and *ZCCT-B2b* [30]), and the *Ppd-A1a* (photoperiod insensitive or PI)

allele that confers earlier flowering under short days resulting in a reduced photoperiodic

response. We also used in our study near-isogenic lines of Kronos carrying either the

photoperiod sensitive *Ppd-A1b* (PS) allele [18] or no functional alleles in the *PPD-A1* and *PPD-*

B1 homeologs (*ppd1*) [19].

Using previously published wheat sequences for *CO1* [48] and *CO2* [42], and the Kronos

transcriptome [67] we identified the A and B coding sequences for *CO1* and *CO2* in Kronos and

deposited them in GenBank under accession numbers MT043302 (*CO-A1*), MT043303 (*CO-B1*),

MT043304 (*CO-A2*), MT043305 (*CO-B2*). We used those sequences to identify loss-of-function

mutations in *CO1* and *CO2* in the Kronos TILLING population [25, 26], which are described in

detail in the Results section. We then combined mutations in the A and B homeologs from each

gene to generate loss-of-function *co1* (henceforth *co1*) and *co2* mutants (henceforth *co2*). We

backcrossed the individual mutant lines with wild-type Kronos at least two times to reduce

background mutations and then combined all four homozygous mutations to produce a *co1 co2*

mutant (henceforth *co1 co2*). In addition to the wild type Kronos-PI, the four homozygous *Co1*

Co2, *co1 Co2*, *Co1 co2* and *co1 co2* allelic classes were generated in Kronos-PS and the *ppd1*

mutant. We were not able to recover the triple loss-of-function mutant *ppd1 co1 co2* from any of

the tested progenies.

Phenotyping

We stratified wheat grains at 4 °C for 2 d in the dark and after germination we planted them in

pots in the soil. We evaluated the mutant and control lines in CONVIRON growth chambers with

metal halide bulbs supplemented by florescent bulbs set to 22 °C during daylight and 17 °C

during dark periods. Lights were set to 260-300 µmol m^{-2} s^{-1} and were on for 16 h in LD

experiments and 8 h in short day (SD) experiments. We recorded heading time as the total time

from planting in soil to full emergence of the spike from the sheath. Spikes were threshed and

total grain number counted per spike.

Statistical analysis

We compared heading time in different genotypes using one-way ANOVAs for single locus and

factorial ANOVAs for two or more loci. We tested homogeneity of variances using the Levene's

test and normality of residuals with the Shapiro–Wilks test. We transformed the data when

necessary to satisfy the ANOVA assumptions. We performed all statistical analyses using SAS

9.4 (SAS Institute). Error bars in all graphs are standard errors of the means (s.e.m.).

Gene expression analysis

To study the effect of *ppd1*, *co1* and *co2* mutant alleles on the transcription profiles of genes

affecting wheat flowering, we collected leaves from plants grown in a growth chamber under LD

using the same settings as described in the previous phenotypic section. For the time course

experiment, we collected samples in liquid nitrogen every 4 h throughout the day from the newly

expanded fourth leaf. In this experiment, we included the *ppd1* mutant and all four possible

combinations of the *CO1* and *CO2* wild type and mutant alleles in the Kronos-PS (*Ppd-A1b*)

background.

RNA was isolated from leaves using the Spectrum Plant Total RNA Kit (Sigma-Aldrich)

following the manufacturer's recommendations. We treated the RNA samples with DNase

(Promega) according to the manufacturer's protocol prior to first-strand cDNA synthesis. We

assessed RNA integrity by a combination of A260/A280 and A260/A230 measurements and

running samples on a denaturing formaldehyde-agarose gel as recommended in the Qiagen

RNAeasy Mini Handbook. We used only samples with 260/280 and 260/230 ratios > 1.7 and no

degradation. We synthesized cDNA using 2.0 μg of total RNA per sample and M-MLV Reverse

Transcriptase (Promega) and poly-T primers (Sigma-Aldrich).

We performed the quantitative reverse transcription-PCR (qRT-PCR) assays using SybrGreen

Takara Master Mix in a 7500 Fast Real-Time PCR system (Applied Biosystems). For PCR, we

used one cycle at 95°C for 30 s and 40 cycles of 95°C for 5 s, 60°C for 5 s, and 72°C for 34 s,

followed by a melting curve program. We quantified expression using the delta Ct method (2^{-}

$^{\Delta CT}$) with *ACTIN* as endogenous control [15]. Primer efficiencies were all > 95% and their

sequences are summarized in Table S5. Primer sequences were previously described for *FT1* and

VRN1 [51], *PPD1* [48], *CO1* and *CO2* [15] and *VRN2* [30].

Yeast-two-hybrid (Y2H) and split yellow fluorescent protein (YFP) assays

The full-length coding region of PPD1 was amplified from Kronos and the PHYB truncations

were amplified from *Triticum monococcum* with primers listed in Supplementary Table S5. The

amplified fragments were cloned into yeast vectors pGBKT7 and pGADT7 (Clontech,

http://www.clontech.com/), which were used to transform yeast strain Y2HGold (Clontech)

using the lithium acetate method. Transformants were selected on SD medium lacking leucine

(L) and tryptophan (W) plates and re-plated on SD medium lacking L, W, histidine (H) and

adenine (A) to test the interactions. Yeast vectors for VRN2 (full-length, both bait and prey),

CO1 (prey, including both B-boxes and CCT domain, amino acids 10 to 379) and CO2 (prey,

including both B-boxes and CCT domain, 15 to 356 amino acids) were previously described in

[33], and those for PHYC (N-PHYC, C-PHYC and FL-PHYC) and full-length PHYB (as prey)

in [15]. CO1 and CO2 showed strong auto-activation when used as baits so they were only used

as preys. The full-length PHYB gave strong autoactivation when used as bait, so we generated

two additional truncations N-PHYB and C-PHYB (primers in Table S5). Only N-PHYB can be

used as bait without autoactivation, and all three (N-PHYB, C-PHYB and FL-PHYB) can be

used as preys in Y2H assays. The truncated versions of PHYC include amino acids 1–600 (N-

terminal) and 601-1139 (C-terminal) [15], and those for PHYB amino acids (1-625 (N-terminal)

and 626-1166 (C-terminal).

For the bi-molecular fluorescent complementation (BiFC or split YFP) assays, each set of wheat

proteins was recombined into modified Gateway-compatible pY736 and pY735 which contain

the *UBIQUITIN* promoter to generate YFP-N-terminal fragment and YFP-C-terminal-fragment

fusion proteins. Wheat protoplasts were prepared, transfected and visualized as described in [68]

ACKNOWLEDEMENTS

This project was supported by NRI Competitive Grants 2017-67007-25939 and 2016-67013-

24617 from the USDA National Institute of Food and Agriculture (NIFA) and by the Howard

Hughes Medical Institute. Daniel Woods is a Howard Hughes Medical Institute Fellow of the

Life Sciences Research Foundation.

AUTHOR CONTRIBUTIONS

Lindsay M. Shaw: Investigation, Writing - Original Draft -Review & Editing. Chengxia Li:

Investigation, Writing - Original Draft -Review & Editing. Daniel Woods: Investigation, Writing

-Review & Editing. Maria A. Alvarez: Investigation, Writing -Review & Editing. Huiqiong:

Investigation, Methodology, Writing -Review & Editing. Mei Yee Lau: Investigation. Andrew

Chen: Investigation, Writing -Review & Editing. Jorge Dubcovsky: Conceptualization, Formal

analysis, Supervision, Project administration and funding acquisition, Writing –extensive Review

& Editing.

CONFLICT OF INTEREST

915 The authors declare that they do not have any conflict of interest.

REFERENCES

1. Bouche F, Woods DP, Amasino RM. Winter memory throughout the plant kingdom:

different paths to flowering. Plant Physiol. 2017;173(1):27-35. doi: 10.1104/pp.16.01322.

PMID: 27756819.

2.   Garner WW, Allard HA. Effect of the relative length of day and night and other factors of

the environment on growth and reproduction in plants. J Agric Res. 1919;18:0553-606.

PMID: WOS:000188347200038.

3.   Andres F, Coupland G. The genetic basis of flowering responses to seasonal cues. Nat

Rev Genet. 2012;13(9):627-39. doi: 10.1038/nrg3291. PMID: 22898651.

4.   Suarez-Lopez P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G.

CONSTANS mediates between the circadian clock and the control of flowering in

Arabidopsis. Nature. 2001;410(6832):1116-20. doi: 10.1038/35074138. PMID:

11323677.

5.   Yanovsky MJ, Kay SA. Molecular basis of seasonal time measurement in Arabidopsis.

Nature. 2002;419(6904):308-12. doi: 10.1038/nature00996. PMID: 12239570.

6.   Putterill J, Robson F, Lee K, Simon R, Coupland G. The CONSTANS gene of

Arabidopsis promotes flowering and encodes a protein showing similarities to zinc finger

transcription factors. Cell. 1995;80(6):847-57. doi: 10.1016/0092-8674(95)90288-0.

PMID: 7697715.

7.   Griffiths S, Dunford RP, Coupland G, Laurie DA. The evolution of *CONSTANS*-like

gene families in barley, rice, and Arabidopsis. Plant Physiol. 2003;131(4):1855-67. doi:

10.1104/pp.102.016188. PMID: 12692345.

8.   Yano M, Katayose Y, Ashikari M, Yamanouchi U, Monna L, Fuse T, et al. Hd1, a major

photoperiod sensitivity quantitative trait locus in rice, is closely related to the Arabidopsis

flowering time gene CONSTANS. Plant Cell. 2000;12(12):2473-84. doi:

10.1105/tpc.12.12.2473. PMID: 11148291.

9.   Higgins JA, Bailey PC, Laurie DA. Comparative genomics of flowering time pathways

using *Brachypodium distachyon* as a model for the temperate grasses. PLoS One.

2010;5(4):e10065. doi: 10.1371/journal.pone.0010065. PMID: 20419097.

10.   Campoli C, Drosse B, Searle I, Coupland G, von Korff M. Functional characterisation of

HvCO1, the barley (*Hordeum vulgare*) flowering time ortholog of *CONSTANS*. Plant J.

2012;69(5):868-80. doi: 10.1111/j.1365-313X.2011.04839.x. PMID: 22040323.

11.   Mulki MA, von Korff M. CONSTANS controls floral repression by up-regulating

VERNALIZATION2 (*VRN-H2*) in Barley. Plant Physiol. 2016;170(1):325-37. doi:

10.1104/pp.15.01350. PMID: 26556793.

12.   Turner A, Beales J, Faure S, Dunford RP, Laurie DA. The pseudo-response regulator

Ppd-H1 provides adaptation to photoperiod in barley. Science. 2005;310(5750):1031-4.

doi: 10.1126/science.1117619. PMID: 16284181.

13.   Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, et al. Natural variation in

OsPRR37 regulates heading date and contributes to rice cultivation at a wide range of

latitudes. Mol Plant. 2013;6(6):1877-88. doi: 10.1093/mp/sst088. PMID: 23713079.

14.   Murphy RL, Klein RR, Morishige DT, Brady JA, Rooney WL, Miller FR, et al.

Coincident light and clock regulation of pseudoresponse regulator protein 37 (PRR37)

controls photoperiodic flowering in sorghum. Proc Natl Acad Sci U S A.

2011;108(39):16469-74. doi: 10.1073/pnas.1106212108. PMID: 21930910.

15.   Chen A, Li C, Hu W, Lau MY, Lin H, Rockwell NC, et al. Phytochrome C plays a major

role in the acceleration of wheat flowering under long-day photoperiod. Proc Natl Acad

Sci U S A. 2014;111(28):10037-44. doi: 10.1073/pnas.1409795111. PMID: 24961368.

16. Beales J, Turner A, Griffiths S, Snape JW, Laurie DA. A pseudo-response regulator is

misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat (Triticum aestivum

L.). Theor Appl Genet. 2007;115(5):721-33. doi: 10.1007/s00122-007-0603-4. PMID:

17634915.

17. Wilhelm EP, Turner AS, Laurie DA. Photoperiod insensitive *Ppd-A1a* mutations in

tetraploid wheat (*Triticum durum* Desf.). Theor Appl Genet. 2009;118(2):285-94. doi:

10.1007/s00122-008-0898-9. PMID: 18839130.

18. Pearce S, Vanzetti LS, Dubcovsky J. Exogenous gibberellins induce wheat spike

development under short days only in the presence of *VERNALIZATION1*. Plant Physiol.

2013;163(3):1433-45. doi: 10.1104/pp.113.225854. PMID: 24085801.

19. Pearce S, Shaw LM, Lin H, Cotter JD, Li C, Dubcovsky J. Night-break experiments shed

light on the *Photoperiod1*-mediated flowering. Plant Physiol. 2017;174(2):1139-50. doi:

10.1104/pp.17.00361. PMID: 28408541.

20. Dubcovsky J, Loukoianov A, Fu D, Valarik M, Sanchez A, Yan L. Effect of photoperiod

on the regulation of wheat vernalization genes *VRN1* and *VRN2*. Plant Mol Biol.

2006;60:469-80.

21. Yan L, Loukoianov A, Blechl A, Tranquilli G, Ramakrishna W, SanMiguel P, et al. The

wheat *VRN2* gene is a flowering repressor down-regulated by vernalization. Science.

2004;303(5664):1640-4. doi: 10.1126/science.1094305. PMID: 15016992.

22. Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J. Positional

cloning of wheat vernalization gene *VRN1*. Proc Natl Acad Sci USA. 2003;100:6263-8.

23. Chen A, Dubcovsky J. Wheat TILLING mutants show that the vernalization gene *VRN1*

down-regulates the flowering repressor *VRN2* in leaves but is not essential for flowering.

PLoS Genet. 2012;8(12):e1003134. doi: 10.1371/journal.pgen.1003134. PMID:

23271982.

24. Shaw LM, Turner AS, Herry L, Griffiths S, Laurie DA. Mutant alleles of *Photoperiod-1*

in wheat (*Triticum aestivum* L.) that confer a late flowering phenotype in long days.

PLoS One. 2013;8(11):e79459. doi: 10.1371/journal.pone.0079459. PMID: 24244507.

25. Krasileva KV, Vasquez-Gross HA, Howell T, Bailey P, Paraiso F, Clissold L, et al.

Uncovering hidden variation in polyploid wheat. Proc Natl Acad Sci U S A.

2017;114(6):E913-E21. doi: 10.1073/pnas.1619268114. PMID: 28096351.

26. Uauy C, Paraiso F, Colasuonno P, Tran RK, Tsai H, Berardi S, et al. A modified

TILLING approach to detect induced mutations in tetraploid and hexaploid wheat. BMC

Plant Biol. 2009;9:115. doi: 10.1186/1471-2229-9-115. PMID: 19712486.

27. Valverde F. CONSTANS and the evolutionary origin of photoperiodic timing of

flowering. J Exp Bot. 2011;62(8):2453-63. doi: 10.1093/jxb/erq449. PMID: 21239381.

28. Waddington SR, Cartwright PM, Wall PC. A quantitative scale of spike initial and pistil

development in barley and wheat. Ann Bot-London. 1983;51(1):119-30. PMID:

WOS:A1983QC01200013.

29. Alvarez MA, Tranquilli G, Lewis S, Kippes N, Dubcovsky J. Genetic and physical

mapping of the earliness *per se* locus *Eps-A^{m}1* in *Triticum monococcum* identifies *EARLY*

FLOWERING 3 (*ELF3*) as a candidate gene. Funct Integr Genomic. 2016;16(4):365-82.

doi: 10.1007/s10142-016-0490-3. PMID: 27085709.

30. Distelfeld A, Tranquilli G, Li C, Yan L, Dubcovsky J. Genetic and molecular

characterization of the *VRN2* loci in tetraploid wheat. Plant Physiol. 2009;149(1):245-57.

doi: 10.1104/pp.108.129353. PMID: 19005084.

31.   Pearce S, Kippes N, Chen A, Debernardi JM, Dubcovsky J. RNA-seq studies using wheat

PHYTOCHROME B and *PHYTOCHROME C* mutants reveal shared and specific

functions in the regulation of flowering and shade-avoidance pathways. BMC Plant Biol.

2016;16(1):141. doi: 10.1186/s12870-016-0831-3. PMID: 27329140.

32.   Kippes N, van Gessel C, Hamilton J, Akpinar A, Budak H, Dubcovsky J, et al. Effect of

phyB and *phyC* loss-of-function mutations on wheat transcriptome under short and long

day photoperiods. bioRxiv. 2020. doi: https://doi.org/10.1101/2020.04.07.030197.

33.   Li C, Distelfeld A, Comis A, Dubcovsky J. Wheat flowering repressor VRN2 and

promoter CO2 compete for interactions with NUCLEAR FACTOR-Y complexes. Plant J.

2011;67(5):763-73. doi: 10.1111/j.1365-313X.2011.04630.x. PMID: 21554456.

34.   Faure S, Turner AS, Gruszka D, Christodoulou V, Davis SJ, von Korff M, et al. Mutation

at the circadian clock gene *EARLY MATURITY 8* adapts domesticated barley (*Hordeum*

vulgare) to short growing seasons. Proc Natl Acad Sci U S A. 2012;109(21):8328-33.

doi: 10.1073/pnas.1120496109. PMID: 22566625.

35.   Gao M, Geng F, Klose C, Staudt A-M, Huang H, Nguyen D, et al. Phytochromes

measure photoperiod in *Brachypodium*. bioRxiv. 2019. doi: 10.1101/697169.

36.   Diaz A, Zikhali M, Turner AS, Isaac P, Laurie DA. Copy number variation affecting the

Photoperiod-B1 and *Vernalization-A1* genes is associated with altered flowering time in

wheat (*Triticum aestivum*). PLoS One. 2012;7(3):e33234. doi:

10.1371/journal.pone.0033234. PMID: 22457747.

37.   Stracke S, Haseneyer G, Veyrieras JB, Geiger HH, Sauer S, Graner A, et al. Association

mapping reveals gene action and interactions in the determination of flowering time in

barley. Theor Appl Genet. 2009;118(2):259-73. doi: 10.1007/s00122-008-0896-y. PMID:

18830577.

38.   Alqudah AM, Sharma R, Pasam RK, Graner A, Kilian B, Schnurbusch T. Genetic

dissection of photoperiod response based on GWAS of pre-anthesis phase duration in

spring barley. PLoS One. 2014;9(11):e113120. doi: 10.1371/journal.pone.0113120.

PMID: 25420105.

39.   Casto AL, Mattison AJ, Olson SN, Thakran M, Rooney WL, Mullet JE. *Maturity2*, a

novel regulator of flowering time in *Sorghum bicolor*, increases expression of *SbPRR37*

and *SbCO* in long days delaying flowering. PLoS One. 2019;14(4):e0212154. doi:

10.1371/journal.pone.0212154. PMID: 30969968.

40.   Yang S, Murphy RL, Morishige DT, Klein PE, Rooney WL, Mullet JE. Sorghum

phytochrome B inhibits flowering in long days by activating expression of *SbPRR37* and

SbGHD7, repressors of *SbEHD1*, *SbCN8* and *SbCN12*. PLoS One. 2014;9(8):e105352.

doi: 10.1371/journal.pone.0105352. PMID: 25122453.

41.   Yang S, Weers BD, Morishige DT, Mullet JE. *CONSTANS* is a photoperiod regulated

activator of flowering in sorghum. BMC Plant Biol. 2014;14:148. doi: 10.1186/1471-

2229-14-148. PMID: 24884377.

42.   Nemoto Y, Kisaka M, Fuse T, Yano M, Ogihara Y. Characterization and functional

analysis of three wheat genes with homology to the *CONSTANS* flowering time gene in

transgenic rice. Plant J. 2003;36(1):82-93. doi: 10.1046/j.1365-313x.2003.01859.x.

PMID: 12974813.

43.   Woods DP, McKeown MA, Dong Y, Preston JC, Amasino RM. Evolution of

VRN2/Ghd7-like genes in vernalization-mediated repression of grass flowering. Plant

Physiol. 2016;170(4):2124-35. doi: 10.1104/pp.15.01279. PMID: 26848096.

44.   Fujino K, Yamanouchi U, Nonoue Y, Obara M, Yano M. Switching genetic effects of the

flowering time gene *Hd*1 in LD conditions by *Ghd7* and *OsPRR37* in rice. Breeding Sci.

2019;69(1):127-32. doi: 10.1270/jsbbs.18060. PMID: 31086490.

45.   Zhang Z, Hu W, Shen G, Liu H, Hu Y, Zhou X, et al. Alternative functions of Hd1 in

repressing or promoting heading are determined by Ghd7 status under long-day

conditions. Sci Rep-Uk. 2017;7(1):5388. doi: 10.1038/s41598-017-05873-1. PMID:

28710485.

46.   Zhang B, Liu H, Qi F, Zhang Z, Li Q, Han Z, et al. Genetic interactions among *Ghd7*,

Ghd8, *OsPRR37* and *Hd1* contribute to large variation in heading date in rice. Rice (N

Y). 2019;12(1):48. doi: 10.1186/s12284-019-0314-x. PMID: 31309345.

47.   Maurer A, Draba V, Jiang Y, Schnaithmann F, Sharma R, Schumann E, et al. Modelling

the genetic architecture of flowering time control in barley through nested association

mapping. BMC Genomics. 2015;16:290. doi: 10.1186/s12864-015-1459-7. PMID:

25887319.

48.   Shaw LM, Turner AS, Laurie DA. The impact of photoperiod insensitive *Ppd-1a*

mutations on the photoperiod pathway across the three genomes of hexaploid wheat

(*Triticum aestivum*). Plant J. 2012;71(1):71-84. doi: 10.1111/j.1365-313X.2012.04971.x.

PMID: 22372488.

49.　Qin Z, Bai Y, Muhammad S, Wu X, Deng P, Wu J, et al. Divergent roles of *FT-like 9* in

flowering transition under different day lengths in *Brachypodium distachyon*. Nat

Commun. 2019;10(1):812. doi: 10.1038/s41467-019-08785-y. PMID: 30778068.

50.　Wang P, Gong R, Yang Y, Yu S. *Ghd8* controls rice photoperiod sensitivity by forming a

complex that interacts with *Ghd7*. BMC Plant Biol. 2019;19(1):462. doi:

10.1186/s12870-019-2053-y. PMID: 31675987.

51.　Yan L, Fu D, Li C, Blechl A, Tranquilli G, Bonafede M, et al. The wheat and barley

vernalization gene *VRN3* is an orthologue of *FT*. Proc Natl Acad Sci U S A.

2006;103(51):19581-6. doi: 10.1073/pnas.0607142103. PMID: 17158798.

52.　Deng W, Casao MC, Wang P, Sato K, Hayes PM, Finnegan EJ, et al. Direct links

between the vernalization response and other key traits of cereal crops. Nat Commun.

2015;6:5882. doi: 10.1038/ncomms6882. PMID: 25562483.

53.　Li C, Lin H, Dubcovsky J. Factorial combinations of protein interactions generate a

multiplicity of florigen activation complexes in wheat and barley. Plant J. 2015;84:70-82.

54.　Turner AS, Faure S, Zhang Y, Laurie DA. The effect of day-neutral mutations in barley

and wheat on the interaction between photoperiod and vernalization. Theor Appl Genet.

2013;126(9):2267-77. doi: 10.1007/s00122-013-2133-6. PMID: 23737074.

55.　Hayama R, Sarid-Krebs L, Richter R, Fernandez V, Jang S, Coupland G. PSEUDO

RESPONSE REGULATORs stabilize CONSTANS protein to promote flowering in

response to day length. Embo J. 2017;36(7):904-18. doi: 10.15252/embj.201693907.

PMID: 28270524.

56.　Nakamichi N, Kita M, Ito S, Yamashino T, Mizuno T. PSEUDO-RESPONSE

REGULATORS, PRR9, PRR7 and PRR5, together play essential roles close to the

circadian clock of *Arabidopsis thaliana*. Plant Cell Physiol. 2005;46(5):686-98. doi:

10.1093/pcp/pci086. PMID: 15767265.

57. Murakami M, Tago Y, Yamashino T, Mizuno T. Characterization of the rice circadian

clock-associated pseudo-response regulators in *Arabidopsis thaliana*. Biosci Biotech

Bioch. 2007;71(4):1107-10. doi: 10.1271/bbb.70048. PMID: 17420570.

58. Campoli C, Shtaya M, Davis SJ, von Korff M. Expression conservation within the

circadian clock of a monocot: natural variation at barley *Ppd-H1* affects circadian

expression of flowering time genes, but not clock orthologs. BMC Plant Biol.

2012;12:97. doi: 10.1186/1471-2229-12-97. PMID: 22720803.

59. Ben-Naim O, Eshed R, Parnis A, Teper-Bamnolker P, Shalit A, Coupland G, et al. The

CCAAT binding factor can mediate interactions between CONSTANS-like proteins and

DNA. Plant J. 2006;46(3):462-76. doi: 10.1111/j.1365-313X.2006.02706.x. PMID:

16623906.

60. Kumimoto RW, Adam L, Hymus GJ, Repetti PP, Reuber TL, Marion CM, et al. The

Nuclear Factor Y subunits NF-YB2 and NF-YB3 play additive roles in the promotion of

flowering by inductive long-day photoperiods in Arabidopsis. Planta. 2008;228(5):709-

23. doi: 10.1007/s00425-008-0773-6. PMID: 18600346.

61. Wenkel S, Turck F, Singer K, Gissot L, Le Gourrierec J, Samach A, et al. CONSTANS

and the CCAAT box binding complex share a functionally important domain and interact

to regulate flowering of Arabidopsis. Plant Cell. 2006;18(11):2971-84. doi:

10.1105/tpc.106.043299. PMID: 17138697.

62. Woods DP, Ream TS, Minevich G, Hobert O, Amasino RM. PHYTOCHROME C is an

essential light receptor for photoperiodic flowering in the temperate grass, *Brachypodium*

distachyon. Genetics. 2014;198(1):397-408. doi: 10.1534/genetics.114.166785. PMID:

25023399.

63.   Osugi A, Itoh H, Ikeda-Kawakatsu K, Takano M, Izawa T. Molecular dissection of the

roles of phytochrome in photoperiodic flowering in rice. Plant Physiol.

2011;157(3):1128-37. doi: 10.1104/pp.111.181792. PMID: 21880933.

64.   Zheng T, Sun J, Zhou S, Chen S, Lu J, Cui S, et al. Post-transcriptional regulation of

Ghd7 protein stability by phytochrome and OsGI in photoperiodic control of flowering in

rice. New Phytol. 2019;224(1):306-20. doi: 10.1111/nph.16010. PMID: 31225911.

65.   Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G.

Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. Science.

2004;303(5660):1003-6. doi: 10.1126/science.1091761. PMID: 14963328.

66.   Hayama R, Coupland G. The molecular basis of diversity in the photoperiodic flowering

responses of Arabidopsis and rice. Plant Physiol. 2004;135(2):677-84. doi: DOI

10.1104/pp.104.042614. PMID: ISI:000222165300012.

67.   Krasileva KV, Buffalo V, Bailey P, Pearce S, Ayling S, Tabbita F, et al. Separating

homeologs by phasing in the tetraploid wheat transcriptome. Genome Biol.

2013;14(6):R66. doi: 10.1186/gb-2013-14-6-r66. PMID: 23800085.

68.   Shan Q, Wang Y, Li J, Gao C. Genome editing in rice and wheat using the CRISPR/Cas

system. Nat Protoc. 2014;9(10):2395-410. doi: 10.1038/nprot.2014.157. PMID:

25232936.

1141

1142

1143

SUPPORTING INFORMATION CAPTIONS

1144

Supplementary Fig S1. Shoot apical meristem (SAM) and spike development.

Supplementary Fig S2. Dissection of developing spikes.

Supplementary Fig S3. Effect of *ppd1*, *co1* and *co2* loss-of-function mutations and photoperiod

on *CO1* and *CO2* transcript levels.

Supplementary Fig S4. Effect of *phyB*-null and *phyC*-null mutations in Kronos-PI on the

transcriptional profiles of *CO1* and *CO2* under SD and LD.

Supplementary Fig S5. Yeast-two-hybrid (Y2H) assays.

Supplementary Table S1. Genome-specific primer sequences and PCR conditions for

TILLING.

Supplementary Table S2. Number of mutations detected in the targeted regions of wheat *CO1*

and *CO2* homologs in the Kronos TILLING population.

Table S3. Analysis of variance for heading time under long days.

Table S4. Analysis of variance for heading time under short days.

Table S5. Primers used in the qRT-PCR and Y2H experiments.