Suicidal Thoughts, Behaviors, and Event-Related Potentials: A Meta-Analysis

Austin J. Gallyer, M.S.[1]; Sean P. Dougherty, M.S.[1]; Kreshnik Burani, M.S.[1]; Brian J. Albanese,

M.S.[1]; Thomas E. Joiner, Ph.D.[1]; Greg Hajcak, Ph.D.[1, 2]

[1]Florida State University, Department of Psychology

[2]Florida State University, Department of Biomedical Sciences

Author Note

This work was supported by a grant from the National Institute of Mental Health

(T32MH093311). This work was also supported in part by the Military Suicide Research

Consortium (MSRC), an effort supported by the Office of the Assistant Secretary of Defense for

Health Affairs under Award No. (W81XWH-16-2-0004). Opinions, interpretations, conclusions,

and recommendations are those of the author and are not necessarily endorsed by the MSRC or

the Department of Defense.

Correspondence concerning this article should be addressed to Austin J. Gallyer, 1107

West Call St., Tallahassee, Florida, 32306. E-mail: gallyer@psy.fsu.edu.

Abstract

Suicidal thoughts and behaviors (STBs) are thought to be the result, at least in part, of

abnormalities in various neural systems. Event-related potentials (ERPs) are a useful method for

studying neural activity, and can be leveraged to study neural deficits related to STBs, but it is

unknown how effective ERPs are at differentiating various STB groups. The present meta-

analysis examined how well ERPs can differentiate those with and without suicidal ideation (SI),

suicide attempts (SA), and those determined to be at different levels of suicide risk, or

suicidality. Our meta-analysis included 192 effect sizes from 24 studies. We used a random

effects meta-analysis and found a small-to-moderate relationship between ERPs and STBs.

However, we also found evidence that the literature may be suffering from small sample

publication bias. Our results suggest that ERPs may compliment other approaches, such as fMRI,

in the study of the neurobiology of individuals with STBs. However, the clinical utility of ERPs

to predict STBs remains to be tested, and larger samples are needed to confirm the results of the

present meta-analysis.

Keywords: suicide, event-related potentials, meta-analysis, suicidal ideation, suicide

attempt

Suicidal Thoughts, Behaviors, and Event-Related Potentials: A Meta-Analysis

Suicide is a concern around the world. Estimates indicate that well over 800,000 people die by suicide around the world annually (World Health Organization, 2014). In the United States (U.S.) alone, over 47,000 people die by suicide each year, making suicide the 10[th] leading cause of death (Centers for Disease Control [CDC], 2017). In contrast to trends in other western countries, suicide rates in the U.S. have been increasing over the last two decades (Hedegaard et al., 2018). This increase in the suicide rate has contributed to the decreasing trend in U.S. life expectancy every year since 2014 (Woolf & Schoomaker, 2019).

Many scientists have suggested that suicidal thoughts and behaviors (STBs) are partially the result of differences or abnormalities in neurobiological systems (Joiner et al., 2005; Mann, 2003; Van Heeringen & Mann, 2014). This suggestion has led to many studies investigating the brain regions that may be implicated in STBs, with most studies using magnetic resonance imaging (MRI) to examine structural and functional brain differences between those with and without a history of STBs.

A recent qualitative systematic review of the neuroimaging literature of STBs concluded that there is tentative but converging evidence for two brain networks implicated in STBs (Schmaal et al., 2019). First, the authors proposed that the ventral prefrontal cortex and many of its connections are involved in increasing negative and decreasing positive internal states, which may lead to suicidal ideation (SI). Second, they identified other regions, including the dorsomedial prefrontal cortex, the dorsolateral prefrontal cortex, and the inferior frontal gyrus, as a separate network that may be important for suicide attempts (SAs) due to these regions' roles in planning and cognitive control. The authors concluded that further studies investigating brain regions could aid in the development of novel treatments for STBs.

In contrast, a quantitative meta-analysis of neuroimaging studies using whole-brain analyses did not find significant associations between SA and any brain region; the study concluded that there were not enough studies to analyze the relationship between SI and neural structure and function (Huang, Rootes-Murdy, et al., 2020). Though these two reviews reached different conclusions about the state of the STB brain imaging literature, both highlight the need for larger sample sizes in future studies examining the relationship between the brain and STBs.

An alternative approach to examining the neurobiology of STBs is to use event-related potentials (ERPs). ERPs are derived from electroencephalographic (EEG) data and reflect neural responses to specific, repeated events (Luck, 2014). ERP waveforms are calculated by averaging the electrocortical responses to a specific task event (e.g., stimulus presentation, error commission) across multiple trials. Although ERPs generally lack the excellent spatial resolution of MRI (i.e., the ability to accurately localize neural functions to regions in the brain), this methodology has several strengths. First, ERPs are direct measures of brain activity with excellent temporal resolution (i.e., over 1000 samples per second), which allows researchers to examine distinct neural activity that might typically overlap with that assessed in MRI (e.g., ERPs can separately index multiple distinct processes that occur within hundreds of milliseconds of one another). Many ERPs have been shown to have excellent internal consistency with sufficient trial numbers (e.g., Meyer et al., 2013; Moran et al., 2013). Electrocortical data are also less expensive to collect relative to other forms of neuroimaging, thereby making larger samples sizes more feasible. Importantly, the extant literature has demonstrated that many ERPs reflect different neural functions, such as emotion regulation, response inhibition, reward processing, reward anticipation, and error monitoring (Luck & Kappenman, 2011), that might be relevant for understanding STBs.

Because of their potential clinical utility (Hajcak et al., 2019), researchers have begun to employ ERPs in the study of STBs. For example, Gibb and Tsypes (2019) suggested that the inclusion of ERPs in machine learning algorithms may improve the prediction of STBs. Though there have been studies examining differences in ERPs between those with SI or a history of SA and those without SI or SAs (e.g., Albanese et al., 2019), the literature on the relationship between ERPs and STBs is quite mixed. For example, one study found that a reduced late positive potential (LPP; a slow wave ERP that indexes emotional salience of stimuli) to pleasant pictures was related to increased SI (Weinberg et al., 2016). However, a follow-up study did not find a relationship between the late positive potential to rewarding stimuli and SI (Weinberg et al., 2017), and another study found a relationship only when individuals were asked to volitionally increase their reward responses (Albanese, Macatee, Stanley, et al., 2019). Though there are many reasons for these mixed findings, a possibility involves the definitions of STBs used across studies. According to ideation-to-action theories of suicide (e.g., interpersonal theory of suicide, 3-step theory; Joiner, 2005; Klonsky & May, 2015), factors that predict SI do not necessarily predict SAs or death by suicide (Klonsky et al., 2018). This is supported by a meta-analysis showing that most risk factors for suicide (e.g., depression) have stronger associations with SI than SAs (May & Klonsky, 2016). Given these findings, ERPs may have different associations with various STBs.

In addition to these uncertainties, recent meta-analyses have called into question the clinical significance of most risk factors for STBs when examined in isolation (Chang et al., 2016; Franklin et al., 2017; Ribeiro et al., 2018). Given these mixed findings and previous meta-analyses, the current meta-analysis aims to: (1) examine whether ERPs, in general, differentiate between individuals who have and have not engaged in STBs; (2) examine which ERP, if any, is

the most effective at differentiating those who have and have not engaged in STBs; and (3)

investigate whether other factors moderate the relationship between ERPs and STBs. We

examined each of these aims in three separate STBs: between those with and without SI, between

those with and without a SA history, and among those with varying degrees of suicide risk.

Method

Inclusion and Exclusion Criteria

All of our data, analysis scripts, coding spreadsheets, figures, and supplementary material

are available on the Open Science Framework (https://osf.io/k4cpe/). Our meta-analysis was

conducted in adherence with the PRISMA Statement (Moher et al., 2009). The following criteria

were established to select relevant effect sizes.

Language

Only articles written in English were included.

Event-related potential

Only articles that had an effect size that included an event-related potential (ERP) were

included. ERPs were defined as measures using EEG responses to specific, time-locked events of

mental processes. Given these criteria, studies that used cortical recordings or that stimulated the

brain to produce neural activity were excluded.

Suicide-related outcomes

Effect sizes were required to include a measure of suicidal thoughts, behaviors, or suicide

risk/suicidality. Suicidal thoughts were defined as any measure of severity or history of suicidal

ideation. We did not distinguish between death/suicidal ideation, or passive/active suicidal

ideation. Suicidal behaviors were defined as a history of engaging in a self-harm behavior with

nonzero intent to die. Importantly, we did not distinguish among aborted, interrupted, and actual

SAs. Moreover, effect sizes that included SA frequency were not included. Last, suicide risk/suicidality was used when a measure/group with STBs that did not fit in the SI or SA categories was obtained. For example, studies that used the Suicidal Behavior Questionnaire—Revised (Osman et al., 2001) were categorized as "suicide risk." Given these definitions, studies that examined nonsuicidal self-injury or a history of nonsuicidal self-injury were excluded.

Published in print or online by October 2019

Our search results were limited to articles published in print or online by October 25, 2019.

Literature Search

We conducted literature searches across PubMed, PsycInfo, Web of Science, and ProQuest Dissertations & Theses to find relevant literature. Full search terms for each database are available on an online repository stored on the Open Science Framework (https://osf.io/k4cpe/). To summarize, search terms included different permutations related to ERPs and suicidal thoughts and behaviors, including: "event-related potential," "event-related potentials," "electroencephalography," "EEG," "evoked potential," "evoked potentials," "late positive potential," "reward positivity," "P3a," "P3b," "error-related negativity," "suicide," "suicidal," "suicidal ideation," "self-harm," "self-injury," "parasuicide," "suicide ideation," "suicide attempt," "suicide plan," "suicidality," "suicide risk," "attempter," "attempters," and "ideators." We also checked the reference list of every article included in the final meta-analysis for additional studies. In addition to these efforts, we made several attempts to gather unpublished literature. These efforts included: (1) social media announcements on Twitter and (2) contacting researchers who were either a first or last author on at least two papers included in the meta-analysis. Of the nine researchers we emailed asking for unpublished or yet-to-be

published studies, six responded, with three providing data and three indicating that they did not have any other unpublished data. The remaining three did not respond.

Data Extraction and Coding

For each effect size, the following were also coded: mean age of the sample, suicide measure, EEG reference, scoring ERP procedure, specific ERP, type of suicidal group, and the type of comparison group. All studies were initially coded by the first author (AJG) in an Excel spreadsheet. We also had one doctoral student (SPD) recode data included in our meta-analysis. We conducted reliability analyses for our codes, using agreement rate and Cohen's kappa for categorical variables and using intercoder correlation and two-way random effects intraclass correlation (ICC) for continuous variables. For categorical variables, agreement rate ranged from 87.5% to 100%, and Cohen's kappa were excellent ($\kappa = .83 - 1.00$). For continuous variables, intercoder correlations ranged from .99 to 1.0, and ICCs also ranged from .99 to 1.0. Disagreements for categorical variables were resolved through discussion.

Using our search terms across all our included databases produced 270 papers. Figure 1 displays an overview of our search process and number of articles within each search process. After accounting for duplicates, we examined the abstracts of 200 papers and excluded studies that did not meet inclusion criteria. After examining abstracts, 51 studies entered full-text review. In the end, 24 studies had at least one effect size that met inclusion criteria, resulting in 192 total effect sizes. The 27 studies that were not included after full-text review were excluded for the following reasons: (1) no measure that met definition of ERP ($n = 3$), (2) no measure that met one of our definitions of STBs ($n = 6$), (3) no, or not enough, information to calculate effect sizes, including when access to the full text was inaccessible due to the article's being a poster or dissertation ($n = 17$), and (4) paper was a review article ($n = 1$). For articles for which the full

text was available but for which we did not have enough information to calculate effect sizes, we

sought to email the corresponding or last author. We emailed four researchers for more

information to calculate effect sizes from their papers. One responded and indicated that they no

longer had access to the data. No other researchers responded, and we were unable to locate

contact information for authors for two studies.

Effect Size

Information was extracted from each study to calculate Hedges' *g*. For most effect sizes,

means, standard deviations, and sample size were used to calculate effect sizes. However, in

some cases, existing effect sizes had to be converted to Hedges' *g*. Specifically, several studies

reported Pearson's *r* or an *F* value. For one effect size, Pearson's *r* was estimated by using the

computer program WebPlotDigitizer (Version 4.2; Rohatgi, 2019). All effect sizes were

calculated using the "compute.es" package (Version 0.2-4; Del Re, 2013). Because the aim of

this meta-analysis was to examine how well ERPs can differentiate those with and without STBs,

we converted all Hedges' *g* effect sizes to |*g*|. This conversion was done because ERPs may have

differing relationships with STBs, with some studies reporting blunted ERPs in those with STBs

and other studies reporting increased ERPs in those with STBs. Moreover, we made this change

because a blunted or larger ERP could be conceptualized as a risk factor, depending on the neural

processes the ERP represents.

Combining Effect Sizes

All of our analyses were conducted in R (Version 3.5.2; R Core Team, 2018), using the

"metafor" package (Version 2.1; Viechtbauer, 2010). We opted to use a random-effects model

that estimated heterogeneity using restricted maximum-likelihood. We chose this model for two

reasons: First, our literature search produced studies with many effect sizes. This dependence

between effect sizes violates the assumptions of fixed-effect models, which would lead to an incorrectly estimated pooled effect size and would increase the probability of a Type I error (Hedges et al., 2010). Second, given that we combined many different ERPs that measure different neural functions, it is likely that our meta-analysis would result in significant heterogeneity in effect sizes. We also used robust variance estimation to account for the small number of clusters (i.e., studies) in our analyses (Hedges et al., 2010). To address outliers, we first conducted our meta-analysis using all effect sizes. Then, any effect sizes whose confidence interval did not overlap with the pooled effect size was removed. We then reran our analyses with outliers removed.

To examine whether publication bias was present, we used two approaches: (1) we visually inspected the funnel plot for each outcome and (2) we conducted moderation analyses using study sample size as the moderating variable. We also ran moderation analyses examining whether specific ERPs, ERP scoring method, and sample age moderated the pooled effect size. In line with our second aim, we examined which specific ERP best differentiated those with and without each of our three STBs. We planned to examine suicide measurement, location of the reference electrode(s), and method of quantifying ERPs (e.g., mean, area-around-the-peak) as moderators, but, for each of these variables, either there were not enough studies reporting these measures, or there was not enough variation in the methods used. We also conducted unplanned sensitivity analyses by comparing our results to the results if (a) outlier effect sizes were included and if (b) only published studies' effect sizes were included.

Results

Suicidal Ideation

Overall SI Results and Publication Bias

After excluding one outlier effect size, our SI meta-analysis consisted of eight studies and 104 effect sizes (M = 13.00 effect sizes per study). The difference in ERPs between those with SI and those without SI was significant, with an average weighted Hedges' g = 0.27 (95% CI [0.14, 0.39]). We did not find evidence of significant heterogeneity (Q[103] = 79.84, p = 0.956, I^2 = 19.75%). Regarding publication bias, visually inspecting the funnel plot (see Figure 2) revealed small-sample publication bias, with smaller samples more frequently having larger effect sizes. As a test of this observation, we examined study sample size as a moderator of the relation between ERPs and SI. The model was significant (F[1, 6] = 25.63, p = .002), with sample size revealing a negative relationship with the pooled effect size estimate (b = -0.001, p = .002). This result suggests that studies with smaller sample sizes tended to find significantly larger effects than studies with larger sample sizes. Though this unstandardized effect appears small, it results in our smallest sample having an estimated effect size of Hedges' g = 0.35; in comparison, our largest sample had an estimated effect size of Hedges' g = 0.10 (see Figure 3).

Moderation of SI by Specific ERP

We tested specific ERPs as a moderator of the relation between ERPs and SI. All ERPs that had 10 or more effect sizes were given their own code; ERPs that did not meet this cutoff were grouped together. Thus, ERPs were coded as follows: loudness dependence of auditory evoked potential (LDAEP) = 0; late positive potential (LPP) = 1; reward positivity (RewP) = 2; positive 300 (P3) = 3; and negative 100 (N1), contingent negative variation (CNV), negative 200

(N2), positive 200 (P2), post-imperative negative variation (PINV), and stimulus-preceding

negativity (SPN) were all coded collectively as 4. The moderation analysis was significant (F[4,

3] = 52.40, p = .004), suggesting that specific ERP contributed to the heterogeneity in effect

sizes. Individual estimates did not find evidence that any other specific ERP differed from the

LDAEP effect (b = 0.330, p < .001), suggesting that specific ERP contributes to significant

heterogeneity in effect sizes, but that no other ERPs have a significantly different relation with SI

than the LDAEP.

Moderation of SI by Scoring Method

We tested whether the ERP scoring method contributed to the heterogeneity in effect

sizes observed in our meta-analysis. In the end, we only had three different scoring methods: (1)

raw, (2) difference score, and (3) slope procedure used to score the LDAEP. Thus, we coded raw

= 0, and difference score and slope procedure as 1. The moderation analysis was not statistically

significant (F[1, 6] = 1.82, p = .226). Thus, we did not find evidence that ERP scoring method

contributed to the heterogeneity observed in our effect sizes.

Moderation of SI by Sample Age

Our analysis examining age as a moderator did not find evidence of age as a significant

contributor to the heterogeneity in our effect sizes for SI (F[1, 5] = 2.67, p = .163).

Overall SA Results and Publication Bias

After excluding three outlier effect sizes, our SA meta-analysis consisted of 12 studies

with 49 effect sizes (M = 4.08 effect sizes per study). The difference in ERPs between those with

a previous suicide attempt and those without a previous suicide attempt was significant, with an

average weighted Hedges' g = 0.40 (95% CI [0.26, 0.54]). Our model did not find evidence of

significant heterogeneity in effect sizes (Q[45] = 42.93, p = .560, I^2 = 19.91%). For a forest plot

of our suicide attempt results, see Figure 4. Regarding publication bias, visual inspection of the funnel plot (see Figure 5) suggested small-sample publication bias, with smaller samples consistently having larger effect sizes than the pooled effect size. We tested this observation by examining sample size as a moderator. The analysis was not statistically significant ($F[1, 10] = 3.24$, $p = .102$). Therefore, we did not find clear evidence that sample size contributed to differences in effect size estimates.

Moderation of SA by Specific ERP

For this analysis, ERPs were coded similarly as it was in the corresponding analysis with SI. For this analysis, the moderation analysis was significant ($F[4, 7] = 31.89$, $p < .001$), suggesting that specific ERPs contributed to the heterogeneity in effect sizes of SA. Individual estimates suggested that the LPP ($b = -0.268$, $p = .029$) and RewP ($b = -0.321$, $p = .014$) each had significantly smaller effect size estimates than the LDAEP ($b = 0.560$, $p < .001$).

Moderation of SA by Scoring Method

Our analysis did not find evidence that ERP scoring method contributed to the heterogeneity in effect sizes ($F[1, 6] = 4.10$, $p = 0.089$).

Moderation of SA by Sample Age

Our analysis examining whether age moderated the relation between ERPs and SA did not converge. After changing the optimizer to "optim" and the optimizer method to "Nelder-Mead," the analysis converged and was statistically significant ($F[1, 6] = 26.11$, $p = .002$), suggesting that the relation between ERPs and SA is smaller in older samples compared to younger samples ($b = -0.019$, $p = .002$; see Figure 6).

Overall Suicide Risk Results and Publication Bias

We did not find any outliers for our suicide risk meta-analysis, resulting in six studies with 38 effect sizes ($M = 6.33$ effect sizes per study). The difference in ERPs between those with higher suicide risk/suicidality and those with lower suicide risk/suicidality was significant, with an average weighted Hedges' $g = 0.33$ (95% CI [0.18, 0.48]). Our model did not find evidence of significant heterogeneity in effect sizes ($Q[37] = 20.93$, $p = .385$, $I^2 = 12.27\%$). For a forest plot of our suicidality results, see Figure 7. Visual inspection of the funnel plot (see Figure 8) suggested small-sample publication bias, with smaller samples consistently having larger effect sizes than the pooled effect size. Results of the analysis examining study sample size as a moderator was not statistically significant ($F[1, 4] = 0.26$, $p = .636$); therefore, we did not find evidence that sample size contributed to differences in effect size estimate in this meta-analysis.

Moderation of Suicide Risk by Specific ERP

For this analysis, the moderation analysis was not significant ($F[2, 3] = 0.79$, $p = .531$), suggesting that measuring different ERPs did not contribute to the heterogeneity in effect sizes of suicide risk.

Moderation of Suicide Risk by Scoring Method

Our analysis did not find evidence that ERP scoring method contributed to the heterogeneity in effect sizes ($F[1, 3] = 0.07$, $p = 0.808$).

Moderation of Suicide Risk by Sample Age

Our analysis examining whether age moderated the relation between ERPs and suicide risk was not statistically significant ($F[1, 4] = 8.82$, $p = .059$); therefore, we did not find evidence that the relation between ERPs and suicide risk differs as a function of the study sample age.

Sensitivity Analyses

SI Sensitivity Analyses

In our SI meta-analysis without outlier exclusion, the pooled effect size increased slightly to Hedges' $g = 0.28$ (95% CI [0.14, 0.41]). In our SI meta-analysis excluding effect sizes from unpublished studies, the pooled effect size estimate increased slightly to Hedges' $g = 0.30$ (95% CI [0.13, 0.47]).

SA Sensitivity Analyses

In our SA meta-analysis without outlier exclusion, the pooled effect size increased significantly to Hedges' $g = 0.53$ (95% CI [0.32, 0.75]), which is nearly outside the 95% CI of our meta-analysis that excluded these outliers. Excluding these outliers but only using published studies also resulted in a larger pooled effect size of Hedges' $g = 0.45$ (95% CI [0.28, 0.62]).

Suicide Risk Sensitivity Analysis

Because we did not find any outliers for the suicide risk variable, we did not conduct a sensitivity analysis reintroducing outliers. A meta-analysis analysis using only published studies resulted in a much larger pooled effect size of $g = 0.42$ (95% CI [0.21, 0.63]).

Discussion

Scientists have proposed that suicidal thoughts and behaviors (STBs) are at least in part the result of differences in neural architecture and functioning (Mann, 2003; Van Heeringen & Mann, 2014). Evidence from fMRI studies has led to divergent conclusions, with some papers concluding that there are neural differences between those with and without STBs, and others suggesting that there is not enough evidence to conclude that there are differences in neural functioning between individuals with and without STBs (Huang, Rootes-Murdy, et al., 2020; Schmaal et al., 2019). Event-related potentials (ERPs) offer an alternative approach to

quantifying neural functioning in those with STBs that is more cost-effective, provides excellent

temporal resolution, and directly measures neurophysiological processes. However, the

effectiveness of ERPs at differentiating individuals with versus without STBs was unknown.

The current meta-analysis revealed a significant small-to-moderate relationship between

ERPs and STBs. On average, those with STBs exhibit differences in neural functioning that can

be measured using ERPs. Though these effects were significant, the potential clinical utility of

ERPs to predict SI or SA is unknown. In our review of the ERP and STB literature, we found no

longitudinal studies examining the ability of ERPs to predict changes in STBs. Moreover, in light

of recent meta-analyses and the size of the cross-sectional effects in our study, it is unlikely that

ERPs used in isolation can provide enough clinically meaningful information to predict STBs

(Franklin et al., 2017). However, ERPs may provide useful and, importantly, unique information

that can increase the predictive value of machine-learning algorithms (Gibb & Tsypes, 2019).

This concept is supported by work that has shown that adding self-report measures to

administrative data can increase prediction accuracy of perpetrating violence or being a victim of

violence among United States Army personnel (Bernecker et al., 2018). Moreover, ERPs may be

used in conjunction with other measures to improve the predictive ability of linear models

(Hajcak et al., 2019). For example, combining ERPs and self-report measures has been used

successfully to predict the development of depression and anxiety (Meyer et al., 2018; Nelson et

al., 2016).

This meta-analysis also sought to examine whether specific ERPs had stronger effects for

different STBs. For SI and suicide risk, we found no evidence that any specific ERP was

significantly better at differentiating those with versus without SI or differentiating those with or

without suicide risk. However, we found that the late positive potential (LPP) and the reward

positivity (RewP) are both significantly *weaker* than the loudness dependence of auditory evoked

potentials (LDAEP) in differentiating between those with versus without previous SAs. Because

our suicide-risk meta-analysis may have been driven by differences in SI, our results may

suggest that differences in severity of SI can be due to disruptions in neural systems. This

suggestion aligns with recent work suggesting that the number of paths leading to SI is

indeterminate (Huang, Ribeiro, et al., 2020; Ribeiro et al., 2019). Another possibility is that what

has been termed the general factor of psychopathology, or *p* factor, is driving this lack of

differences (Caspi et al., 2014; Lahey et al., 2012). Specifically, in addition to more specific

factors such as internalizing and externalizing factors, the *p* factor may cause disruptions across

neural systems, thereby increasing risk for SI and causing differences across various ERPs. For

now, though, these ideas are speculative and point toward potential future studies.

Regarding our SA results, we see at least two possibilities accounting for the LPP and

RewP having smaller effect sizes. First, the LPP and RewP may be better at detecting other STB

constructs, such as SI, than at detecting SA. This possibility is underscored by work showing that

the LPP and the RewP independently relate to depression (Klawohn et al., 2020), and research

has shown that accounting for depression reduces SI's integrity as a construct (Rogers et al.,

2016). Therefore, the LPP and RewP may better relate to depression/SI. Another possibility is

that smaller effect sizes were driven by larger samples and unpublished work that used the LPP

and RewP. For example, the mean sample size of studies in our SA analyses that used the LPP

was 235, compared to a mean sample size of 38.9 for the LDAEP. Thus, these lower effect sizes

for the LPP and RewP may reflect greater accuracy in detecting the true effect of the relation

between the ERP and SA, whereas other ERP's effect sizes may be inflated due to small-sample

publication bias.

We also found that the mean age of the sample moderated the relation between ERPs and SA history, such that older samples corresponded with smaller effect sizes. One explanation for this result is that large differences in neural functioning early in life are markers of particularly high risk for suicide. Over time, these differences may diminish as those with the largest neural differences die by suicide, restricting the range of ERPs in older samples with STBs[1]. However, given that death by suicide is relatively rare, this effect, if it exists, may be relatively small. Another possibility is that being younger generally serves as a protective factor that can only be overridden by large perturbations in neural functioning, leading to increased risk for SAs. Indeed, while suicide rates see a large increase in late childhood and early adolescence, even very young children can exhibit STBs (CDC, 2017; Rosenthal & Rosenthal, 1984).

Though we found significant differences in ERPs between various STB groups, these results should be interpreted within the context of possible small-sample publication bias across STB outcomes. Specifically, all three funnel plots (i.e., for suicide risk/suicidality, SI, and SA) suggested that smaller studies had inflated effect sizes and that many nonsignificant effect sizes might be missing from the published literature. In our SI meta-analysis, we also found that study sample size moderated the relation between ERPs and SI, such that studies with larger samples had significantly smaller effect size estimates compared to studies with smaller samples (see Figure 3). Though we did not find this significant moderation in our SA and suicidality meta-analyses, we may have been underpowered to detect moderation effects in these meta-analyses.

---

[1] The authors thank Keanan Joyner for making us aware of this possibility.

Future of the Study of STBs Using ERPs

The results of our meta-analysis show that ERPs have promise in aiding our understanding of the neural functioning of those with STBs. ERPs are relatively cheap to collect and can be used in myriad settings, increasing the possibility of measuring ERPs in more ecologically valid settings (Shamay-Tsoory & Mendelsohn, 2019). Capitalizing on the strengths of ERPs can complement the spatial resolution of fMRI to give a fuller picture of the neural functioning of those with STBs. However, as with the fMRI literature, where a meta-analysis found a median sample size of 45 (Huang, Rootes-Murdy, et al., 2020), most studies using ERPs to investigate STBs suffer from small sample sizes. In our review, the mean sample size across studies was 72.95 ($SD = 63.36$), with most studies across the different STB categories having fewer than 60 participants (see Figure 9). Low statistical power increases the frequency of both false negatives and false positives, which limits the replicability of studies of STBs (Button et al., 2013). There are likely many factors contributing to low statistical power in the neuroimaging and ERP literature, including institutional incentives (Smaldino & McElreath, 2016), the high financial cost to conduct neuroimaging studies, and the relatively low base-rate of STBs (Fazel & Runeson, 2020). To address low statistical power, consortia have been developed to construct large neuroimaging datasets (e.g., Thompson et al., 2014). A similar consortium using ERPs, as well as greater collaboration in general, would substantially move the field forward in understanding the neurobiology of STBs.

Conclusions

Our results indicate that event-related potentials (ERPs) have a small-to-moderate relationship with suicidal thoughts and behaviors (STBs), such that ERPs generally reveal differences between those with versus without suicidal thoughts, previous suicide attempts, and

suicide risk/suicidality. These findings suggest that ERPs can complement neuroimaging in studies of the neurobiology of STBs. Our review revealed that all research to date has examined the relation between ERPs and STBs cross-sectionally, with no attempts at prediction (e.g., longitudinal studies were absent from the available literature). Therefore, it remains to be seen whether ERPs can be used in conjunction with other measures to create suicide risk algorithms that can more accurately predict STBs, including suicidal ideation (SI) and suicide attempts (SA). For SI and suicide risk/suicidality, we did not find that any single ERP had a significantly stronger relationship with SI or with suicide risk/suicidality than other ERPs. Therefore, future research may consider using multiple ERPs, or use ERPs with other measures, to predict STBs. Moreover, we found that the late positive potential (an index of the neural response to emotional salience) and the reward positivity (an ERP that measures response to rewards) each had a significantly smaller relationship with suicide attempt status than other ERPs, suggesting an important area for future research. Our results suggest that the extant literature examining ERPs and STBs may be suffering from small-sample publication bias, with smaller samples having much larger effect sizes. We also found that the modal study on ERPs and STBs is likely underpowered to estimate the expected effect size. Thus, future researchers should consider collaborating to obtain larger, more representative samples in order to better detect effects and elucidate the neurobiology of STBs using ERPs.

References marked by * indicate studies included in meta-analysis before outliers removed.

References

*Albanese, B. J. (2020). *Unpublished Data*.

*Albanese, B. J., Macatee, R. J., Gallyer, A. J., Stanley, I. H., Joiner, T. E., & Schmidt, N. B. (2019). Impaired Conflict Detection Differentiates Suicide Attempters From Ideating Nonattempters: Evidence From Event-Related Potentials. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *4*(10), 902–912. https://doi.org/Suicidal behaviors among American Indian/Alaska Native firefighters: Evidence for the role of painful and provocative events

*Albanese, B. J., Macatee, R. J., Stanley, I. H., Bauer, B. W., Capron, D. W., Bernat, E., Joiner, T. E., & Schmidt, N. B. (2019). Differentiating suicide attempts and suicidal ideation using neural markers of emotion regulation. *Journal of Affective Disorders*, *257*, 536–550. https://doi.org/10.1016/j.jad.2019.07.014

*Ashton, C. H., Marshall, E. F., Hassanyeh, F., Marsh, V. R., & Wright-Honari, S. (1994). Biological correlates of deliberate self-harm behaviour: A study of electroencephalographic, biochemical and psychological variables in parasuicide. *Acta Psychiatrica Scandinavica*, *90*(5), 316–323. https://doi.org/10.1111/j.1600-0447.1994.tb01600.x

*Baik, S. Y., Jeong, M., Kim, H. S., & Lee, S.-H. (2018). ERP investigation of attentional disengagement from suicide-relevant information in patients with major depressive disorder. *Journal of Affective Disorders*, *225*, 357–364. https://doi.org/10.1016/j.jad.2017.08.046

Bernecker, S. L., Rosellini, A. J., Nock, M. K., Chiu, W. T., Gutierrez, P. M., Hwang, I., Joiner, T.

    E., Naifeh, J. A., Sampson, N. A., Zaslavsky, A. M., Stein, M. B., Ursano, R. J., &

    Kessler, R. C. (2018). Improving risk prediction accuracy for new soldiers in the U.S.

    Army by adding self-report survey data to administrative data. *BMC Psychiatry*, *18*(87).

    https://doi.org/10.1186/s12888-018-1656-4

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., &

    Munafò, M. R. (2013). Power failure: Why small sample size undermines the reliability

    of neuroscience. *Nature Reviews Neuroscience*, *14*. https://doi.org/10.1038/nrn3475

Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., Meier,

    M. H., Ramrakha, S., Shalev, I., Poulton, R., & Moffitt, T. E. (2014). The p Factor: One

    General Psychopathology Factor in the Structure of Psychiatric Disorders? *Clinical*

    Psychological Science, *2*(2), 119–137. https://doi.org/10.1177/2167702613497473

Centers for Disease Control. (2017). *WISQARS: Web-Based Injury Statistics Query and*

    Reporting System [Government Database]. www.cdc.gov/injury/wisqars

Chang, B. P., Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., & Nock, M. K. (2016).

    Biological risk factors for suicidal behaviors: A meta-analysis. *Translational Psychiatry*,

    6. https://doi.org/10.1038/tp.2016.165

*Chen, T.-J., Yu, Y. W.-Y., Chen, M.-C., Wang, S.-Y., Tsai, S.-J., & Lee, T.-W. (2005). Serotonin

    Dysfunction and Suicide Attempts in Major Depressives: An Auditory Event-Related

    Potential Study. *Neuropsychobiology*, *52*(1), 28–36. https://doi.org/10.1159/000086175

Del Re, A. C. (2013). *compute.es: Compute effect sizes*. http://cran.r-

    project.org/web/packages/compute.es

Fazel, S., & Runeson, B. (2020). Suicide. *New England Journal of Medicine*, *382*(3), 266–274.

    https://doi.org/10.1056/NEJMra1902944

Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., Kleiman, E. M., Huang, X., Musacchio,

    K. M., Jaroszewski, A. C., Chang, B. P., & Nock, M. K. (2017). Risk factors for suicidal

    thoughts and behaviors: A meta-analysis of 50 years of research. *Psychological Bulletin*,

    143(2), 187–232. https://doi.org/10.1037/bul0000084

Gibb, B. E., & Tsypes, A. (2019). Using Event-Related Potentials to Improve Our Prediction of

    Suicide Risk. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *4*(10),

    854–855. https://doi.org/10.1016/j.bpsc.2019.08.003

*Graßnickel, V., Illes, F., Juckel, G., & Uhl, I. (2015). Loudness dependence of auditory evoked

    potentials (LDAEP) in clinical monitoring of suicidal patients with major depression in

    comparison with non-suicidal depressed patients and healthy volunteers: A follow-up-

    study. *Journal of Affective Disorders*, *184*, 299–304.

    https://doi.org/10.1016/j.jad.2015.06.007

Hajcak, G., Klawohn, J., & Meyer, A. (2019). The Utility of Event-Related Potentials in Clinical

    Psychology. *Annual Review of Clinical Psychology*, *15*(1), 71–95.

    https://doi.org/10.1146/annurev-clinpsy-050718-095457

*Hansenne, M., Pitchot, W., Gonzalez Moreno, A., Urcelay Zaldua, I., & Ansseau, M. (1996).

    Suicidal behavior in depressive disorder: An event-related potential study. *Biological

    Psychiatry*, *40*(2), 116–122. https://doi.org/10.1016/0006-3223(95)00372-X

*Hansenne, M., Pitchot, W., Moreno, A. G., Torrecilas, J. G., Mirel, J., & Ansseau, M. (1994).

    Psychophysiological correlates of suicidal behavior in depression. *Neuropsychobiology*,

    30(1), 1–3.

Hedegaard, H., Curtin, S. C., & Warner, M. (2018). *Suicide Mortality in the United States, 1999-2017* [Government Report]. Centers for Disease Control.

Hedges, L. V., Tipton, E., & Johnson, M. C. (2010). Robust variance estimation in meta-regression with dependent effect size estimates. *Research Synthesis Methods*, *1*(1), 39–65. https://doi.org/10.1002/jrsm.5

Huang, X., Ribeiro, J. D., & Franklin, J. C. (2020). The Differences Between Suicide Ideators and Suicide Attempters: Simple, Complicated, or Complex? *Journal of Consulting and Clinical Psychology*, 17.

Huang, X., Rootes-Murdy, K., Bastidas, D. M., Nee, D. E., & Franklin, J. C. (2020). Brain Differences Associated with Self-Injurious Thoughts and Behaviors: A Meta-Analysis of Neuroimaging Studies. *Scientific Reports*, *10*(1), 2404. https://doi.org/10.1038/s41598-020-59490-6

*Jandl, M., Steyer, J., & Kaschka, W. P. (2010). Suicide risk markers in Major Depressive Disorder: A study of Electrodermal Activity and Event-Related Potentials. *Journal of Affective Disorders*, *123*(1–3), 138–149. https://doi.org/10.1016/j.jad.2009.09.011

Joiner, T. E. (2005). *Why people die by suicide*. Harvard University Press.

Joiner, Thomas E, Brown, J. S., & Wingate, L. R. (2005). The psychology and neurobiology of suicidal behavior. *Annual Review of Psychology*, *56*, 287–314. https://doi.org/10.1146/annurev.psych.56.091103.070320

*Juckel, G., & Hegerl, U. (1994). Evoked Potentials, Serotonin, and Suicidality. *Pharmacopsychiatry*, *27*(S 1), 27–29. https://doi.org/10.1055/s-2007-1014323

*Kim, D.-H., & Park, Y.-M. (2013). The association between suicidality and serotonergic

dysfunction in depressed patients. *Journal of Affective Disorders*, *148*(1), 72–76.

https://doi.org/10.1016/j.jad.2012.11.051

Klawohn, J., Burani, K., Bruchnak, A., Santopetro, N., & Hajcak, G. (2020). Reduced neural

response to reward and pleasant pictures independently relate to depression.

Psychological Medicine, 1–9. https://doi.org/10.1017/S0033291719003659

Klonsky, E. D., & May, A. M. (2015). The Three-Step Theory (3ST): A new theory of suicide in

the "ideation-to-action" framework. *International Journal of Cognitive Therapy*, *8*(2),

114–129.

Klonsky, E. D., Saffer, B. Y., & Bryan, C. J. (2018). Ideation-to-action theories of suicide: A

conceptual and empirical update. *Current Opinion in Psychology*, *22*, 38–43.

https://doi.org/10.1016/j.copsyc.2017.07.020

*Kudinova, A. Y., Owens, M., Burkhouse, K. L., Barretto, K. M., Bonanno, G. A., & Gibb, B. E.

(2016). Differences in emotion modulation using cognitive reappraisal in individuals with

and without suicidal ideation: An ERP study. *Cognition and Emotion*, *30*(5), 999–1007.

https://doi.org/10.1080/02699931.2015.1036841

Lahey, B. B., Applegate, B., Hakes, J. K., Zald, D. H., Hariri, A. R., & Rathouz, P. J. (2012). Is

there a general factor of prevalent psychopathology during adulthood? *Journal of

Abnormal Psychology*, *121*(4), 971–977. https://doi.org/10.1037/a0028355

*Lee, S.-H., Park, Y.-C., Yoon, S., Kim, J.-I., & Hahn, S. W. (2014). Clinical implications of

loudness dependence of auditory evoked potentials in patients with atypical depression.

Progress in Neuro-Psychopharmacology and Biological Psychiatry, *54*, 7–12.

https://doi.org/10.1016/j.pnpbp.2014.05.010

Luck, S. J. (2014). *An introduction to the event-related potential technique* (2nd ed.). The MIT

    Press.

Luck, S. J., & Kappenman, E. S. (2011). *The Oxford Handbook of Event-Related Potential*

    Components. Oxford University Press.

Mann, J. J. (2003). Neurobiology of suicidal behaviour. *Nature Reviews: Neuroscience*, *4*, 819–

    828. https://doi.org/10.1038/nrn1220

*Marsic, A. (n.d.). *THE RELATIONSHIP BETWEEN SUICIDE IDEATION AND*

    PARASUICIDE: AN ELECTROPHYSIOLOGICAL INVESTIGATION USING THE

    LOUDNESS DEPENDENCE OF AUDITORY EVOKED POTENTIAL.

*Marsic, A., Berman, M. E., Barry, T. D., & McCloskey, M. S. (2015). The Relationship

    Between Intentional Self-Injurious Behavior and the Loudness Dependence of Auditory

    Evoked Potential in Research Volunteers: Self-Injurious Behavior and the LDAEP.

    Journal of Clinical Psychology, *71*(3), 250–257. https://doi.org/10.1002/jclp.22136

May, A. M., & Klonsky, E. D. (2016). What distinguishes suicide attempters from suicide

    Ideators? A meta-analysis of potential factors. *Clinical Psychology: Science and Practice*,

    23(1), 5–20. https://doi.org/10.1111/cpsp.12136

Meyer, A., Nelson, B., Perlman, G., Klein, D. N., & Kotov, R. (2018). A neural biomarker, the

    error-related negativity, predicts the first onset of generalized anxiety disorder in a large

    sample of adolescent females. *Journal of Child Psychology and Psychiatry*, *59*(11),

    1162–1170. https://doi.org/10.1111/jcpp.12922

Meyer, A., Riesel, A., & Proudfit, G. H. (2013). Reliability of the ERN across multiple tasks as a

    function of increasing errors: Reliability of the ERN across multiple tasks.

    Psychophysiology, *50*(12), 1220–1225. https://doi.org/10.1111/psyp.12132

*Min, J.-A., Lee, S.-H., Lee, S.-Y., Chae, J.-H., Lee, C.-U., Park, Y.-M., & Bae, S.-M. (2012).
Clinical characteristics associated with different strengths of loudness dependence of
auditory evoked potentials (LDAEP) in major depressive disorder. *Psychiatry Research*,
200(2–3), 374–381. https://doi.org/10.1016/j.psychres.2012.06.038

Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred Reporting Items for
Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annals of Internal
Medicine*, 264–269.

Moran, T. P., Jendrusina, A. A., & Moser, J. S. (2013). The psychometric properties of the late
positive potential during emotion processing and regulation. *Brain Research*, *1516*, 66–
75. https://doi.org/10.1016/j.brainres.2013.04.018

Nelson, B. D., Perlman, G., Klein, D. N., Kotov, R., & Hajcak, G. (2016). Blunted Neural
Response to Rewards as a Prospective Predictor of the Development of Depression in
Adolescent Girls. *American Journal of Psychiatry*, *173*(12), 1223–1230.
https://doi.org/10.1176/appi.ajp.2016.15121524

Osman, A., Bagge, C. L., Gutierrez, P. M., Konick, L. C., Kopper, B. A., & Barrios, F. X. (2001).
The Suicidal Behaviors Questionnaire-Revised (SBQ-R): Validation with clinical and
nonclinical samples. *Assessment*, *8*(4), 443–454.
https://doi.org/10.1177/107319110100800409

R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation
for Statistical Computing. https://www.r-project.org/

Ribeiro, J. D., Huang, X., Fox, K. R., & Franklin, J. C. (2018). Depression and hopelessness as
risk factors for suicide ideation, attempts and death: Meta-analysis of longitudinal
studies. *British Journal of Psychiatry*. https://doi.org/10.1192/bjp.2018.27

Ribeiro, J. D., Huang, X., Fox, K. R., Walsh, C. G., & Linthicum, K. P. (2019). Predicting

Imminent Suicidal Thoughts and Nonfatal Attempts: The Role of Complexity. *Clinical

Psychological Science*, 216770261983846. https://doi.org/10.1177/2167702619838464

Rogers, M. L., Stanley, I. H., Hom, M. A., Chiurliza, B., Podlogar, M. C., & Joiner, T. E. (2016).

Conceptual and empirical scrutiny of covarying depression out of suicidal ideation.

Assessment. https://doi.org/10.1177/1073191116645907

Rohatgi, A. (2019). *WebPlotDigitizer* (Version 4.2) [Computer software].

https://automeris.io/WebPlotDigitizer

Rosenthal, P. A., & Rosenthal, S. (1984). Suicidal behavior by preschool children. *American

Journal of Psychiatry*, *April*, 520–525.

Schmaal, L., van Harmelen, A.-L., Chatzi, V., Lippard, E. T. C., Toenders, Y. J., Averill, L. A.,

Mazure, C. M., & Blumberg, H. P. (2019). Imaging suicidal thoughts and behaviors: A

comprehensive review of 2 decades of neuroimaging studies. *Molecular Psychiatry*.

https://doi.org/10.1038/s41380-019-0587-x

Shamay-Tsoory, S. G., & Mendelsohn, A. (2019). Real-Life Neuroscience: An Ecological

Approach to Brain and Behavior Research. *Perspectives on Psychological Science*, *14*(5),

841–859. https://doi.org/10.1177/1745691619856350

Smaldino, P. E., & McElreath, R. (2016). The natural selection of bad science. *Royal Society

Open Science*, *3*(9), 160384. https://doi.org/10.1098/rsos.160384

*Song, W., Li, H., Guo, T., Jiang, S., & Wang, X. (2019). Effect of Affective Reward on

Cognitive Event-related Potentials and its Relationship with Psychological Pain and

Suicide Risk among Patients with Major Depressive Disorder. *Suicide and Life-

Threatening Behavior*, *49*(5), 1290–1306. https://doi.org/10.1111/sltb.12524

*Tavakoli, P., Boafo, A., Dale, A., Robillard, R., Greenham, S. L., & Campbell, K. (2018). Event-Related Potential Measures of Attention Capture in Adolescent Inpatients With Acute Suicidal Behavior. *Frontiers in Psychiatry*, *9*, 85. https://doi.org/10.3389/fpsyt.2018.00085

Thompson, P. M., Stein, J. L., Medland, S. E., Hibar, D. P., Vasquez, A. A., Renteria, M. E., Toro, R., Jahanshad, N., Schumann, G., Franke, B., Wright, M. J., Martin, N. G., Agartz, I., Alda, M., Alhusaini, S., Almasy, L., Almeida, J., Alpert, K., Andreasen, N. C., … Drevets, W. (2014). The ENIGMA Consortium: Large-scale collaborative analyses of neuroimaging and genetic data. *Brain Imaging and Behavior*, *8*(2), 153–182. https://doi.org/10.1007/s11682-013-9269-5

*Tsypes, A., & Gibb, B. E. (2020). *Unpublished Data (1)*.

*Tsypes, A., & Gibb, B. E. (2020). *Unpublished Data (2)*.

*Tsypes, A., Owens, M., & Gibb, B. E. (2019). Blunted Neural Reward Responsiveness in Children With Recent Suicidal Ideation. *Clinical Psychological Science*, 216770261985634. https://doi.org/10.1177/2167702619856341

Van Heeringen, K., & Mann, J. J. (2014). The neurobiology of suicide. *The Lancet Psychiatry*, *1*(1), 63–72. https://doi.org/10.1016/S2215-0366(14)70220-2

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software*, *36*(3). https://doi.org/10.18637/jss.v036.i03

*Weinberg, A., May, A. M., Klonsky, E. D., Kotov, R., & Hajcak Greg. (2017). Decreased neural response to threat differentiates patients who have attempted suicide from nonattempters with current ideation. *Clinical Psychological Science*. https://doi.org/10.1177/2167702617718193

*Weinberg, A., Perlman, G., Kotov, R., & Hajcak, G. (2016). Depression and reduced neural

      response to emotional images: Distinction from anxiety, and importance of symptom

      dimensions and age of onset. *Journal of Abnormal Psychology*, *125*(1), 26–39.

      https://doi.org/10.1037/abn0000118

Woolf, S. H., & Schoomaker, H. (2019). Life Expectancy and Mortality Rates in the United

      States, 1959-2017. *JAMA*, *322*(20), 1996. https://doi.org/10.1001/jama.2019.16932

World Health Organization. (2014). *Preventing suicide: A global imperative*. World Health

      Organization. https://doi.org/10.1002/9780470774120

Figure 1. PRISMA diagram for the selection of studies for the present meta-analysis.

Figure 2. Funnel plot for the suicidal ideation (SI) meta-analysis.

Figure 3. Scatterplot illustrating moderation of average suicidal ideation (SI) effect size by effect

sample size.

Figure 4. Average difference (Hedges' *g*) in event-related potentials (ERPs) between those with

and those without a previous suicide attempt.

Figure 5. Funnel plot for the suicide attempt meta-analysis.

Figure 6. Scatterplot illustrating moderation of average suicide attempt (SA) effect size by effect

sample age.

Figure 7. Average weighted absolute difference Hedges' *g* in event-related potentials (ERPs)

between those at lower risk for suicide and those at higher risk for suicide.

Figure 8. Funnel plot for the suicide risk/suicidality meta-analysis.

Figure 9. Distribution of study sample sizes across the three suicidal groups in the present study.