Invasive freshwater snails form novel microbial symbioses

Bankers L.[1,*,†,‡], Dahan D.[2,†,], Neiman M.[1], Adrian-Tucci C.[1,§], Frost C.[3], Hurst G.D.D[3]., King K.C.[4]

Author Affiliations:

[1] University of Iowa, Department of Biology, Iowa City, IA, USA

[2]Stanford University, School of Medicine, Stanford, CA, USA

[3] University of Liverpool, Institute of Integrative Biology, Liverpool, UK

[4] University of Oxford, Department of Zoology, Oxford, UK

*Corresponding author:

Laura Bankers: laura.bankers@cuanschutz.edu

[†] LB and DD contributed equally to this work and should be considered joint first authors.
[‡] Current address: University of Colorado Anschutz Medical Campus, Aurora, CO, USA
[§] Current address: 1150 4th St. SW Apt. 607 Washington DC, USA

ABSTRACT

Resident microbes (microbiota) can shape host organismal function and adaptation in the face

of environmental change. Invasion of new habitats exposes hosts to novel selection pressures,

but the impacts on microbiota and their relationship with hosts after this transition (e.g., how

rapidly symbioses are formed, whether microbes influence invasion success) are unclear. We

used high-throughput 16S rRNA sequencing of New Zealand (native) and European (invasive)

populations of the freshwater snail *Potamopyrgus antipodarum* to find that while invaders do

carry over some core microbial taxa from New Zealand, most of their microbial community is

distinct. This finding highlights that invasions can result in the formation of novel symbioses.

We further show that the native microbiome is composed of fewer core microbes than invasive

snails, suggesting that the microbiota is streamlined to essential members. Together, our

findings demonstrate that microbiota comparisons across native and invasive populations can

reveal the impact of a long coevolutionary history and specialization of microbes in the native

host range, as well as new associations occurring after invasion.

Keywords: *Potamopyrgus antipodarum*, invasion, microbiome, coevolution, symbiosis,

infection

 INTRODUCTION

All organisms are home to numerous microbes. The relationship between hosts and

their microbial communities (microbiota) can play a role in key host functions (1, 2) from

nutrition (3) to behaviour (4) and immune responses (5). The microbiota has also been shown

to impact host evolution (6, 7). Microbial community composition can be determined by host

traits such as genotype (8), sex (9), infection of the host by parasites or pathogens (10),

environmental conditions (6, 11), and genotype by environment interactions (12). Therefore,

host organisms and the microbes on which they depend for vital processes can both be

impacted by environmental changes, with potential consequences for host adaptation (13).

When species invade novel habitats, they are exposed to new environmental conditions

and selection pressures. Because host and microbe fitness can be mutually dependent, invasion

is predicted to influence both the host and its microbiome (6). The fitness effects of microbes

on their host may affect invasion success and the potential for subsequent adaptation to a new

habitat (14). The performance of introduced species could also be altered by the availability of

microbes outside their native range if there are core microbes that the host needs to thrive. If

host-associated microbes are not all carried over from the native environment, lost upon

invasion, and/or unavailable in the new environment, the diversity of the microbiomes of

invasive lineages could be low. The invader's microbiome might thus only represent a subset of

that residing in the native lineages (15, 16). This pattern is directly analogous to the loss of

genetic diversity that often characterizes invasive lineages (15, 17). Whether such subsampling

occurs at the microbiome level remains an important open question with implications for host

fitness in the new environment.

New invasions may also affect the level of specialization of microbiota relative to hosts

in the native range. This outcome could be a consequence of the comparatively short time

period for coevolution between the invasive host and newly acquired microbes vs. the microbes

from the native environment (15, 16). These effects of colonization could influence microbiome

structure. These consequences could be harmful for the host, such as in a situation where

suitable mutualistic microbes are not in the new habitat. It is also possible that invasive hosts

form new symbioses with the microbes available (18). A more diverse microbiota may result if

there is decreased immune-mediated control of novel microbes (19). By this logic, we might

therefore expect native host individuals to have a less rich microbiota. In particular, established

and stable microbial communities, in which novel competitive interactions between microbes

have long since played out, are likely to be of lower diversity (20). This prediction might be

particularly likely to hold if the host has evolved to cultivate beneficial associations (19).

Addressing whether this prediction is met requires characterization of how invasion impacts

microbial diversity and composition.

While molluscs are one of the most species-rich phyla (21), and represent an

ecologically and economically important class of invaders (22, 23), little is known about the

relationships between molluscs and their symbiotic microbes with the exception of a handful of

recent studies (24, 25). Here, we leveraged a powerful model system for host-parasite

coevolution, the New Zealand freshwater snail *Potamopyrgus antipodarum*, to characterize the

impact of invasion on host microbiota. There is substantial across-lake population variation in

microbiome composition among native *P. antipodarum* (26). *Potamopyrgus antipodarum* in

these native New Zealand populations also feature wide individual-level variation in

reproductive mode (sexual vs. asexual) and ploidy level (sexual snails are diploid while asexual

snails are triploid or tetraploid; (27, 28)), and are coevolving with sterilizing trematode parasites

(29, 30). By contrast, invasive lineages of *P. antipodarum* in Europe and North America are

polyploid asexuals with reduced genetic variation compared to the native range (17, 31) and

have escaped from parasite infection (32, 33).

We conducted 16S rRNA sequencing of field-collected *P. antipodarum* from waterbodies

across New Zealand and Europe to assess changes in microbiota diversity and composition

across and within native and invasive host populations. We predicted that, 1) similar to

patterns often described for genetic diversity, invasive *P. antipodarum* would have a subset of

native microbial diversity because invaders are often subsamples of native diversity and

because key symbionts needed to thrive should be maintained in the new environment; 2)

invaders would possess lower microbial diversity due to subsampling and because they have

had less time in the new environment to acquire new symbioses; 3) invaders will have fewer

conserved microbes than native counterparts as a consequence of relatively recent contact

with symbionts in the new environment, and therefore, relatively low time for the evolution of

specialization. To our knowledge, our study of *P. antipodarum* is the first comparison of

diversification in host-associated microbial communities between native and invasive molluscs.

We also address associations between the fundamental and often-variable organismal traits of

sex, reproductive mode, and infection status on microbiota within those host populations.

Study System

Native New Zealand *P. antipodarum* are primarily found in freshwater lakes and streams

(34). These snails have gained prominence as a model for studying the evolutionary

maintenance of sexual reproduction (27, 28) as well as host-parasite coevolution (27, 30).

Multiple triploid and tetraploid asexual lineages have been separately derived from diploid

sexual conspecifics (27, 28), and New Zealand populations are commonly infected by a

sterilizing trematode parasite, *Atriophallophorus winterbourni* (formerly *Microphallus livelyi*)

(27, 30, 35). A previous study hinted that the microbiome composition of these snails varies

substantially among native populations and between sexuals and asexuals (26). However, these

snails were either laboratory cultured or housed in a laboratory for several months before

harvest (Maurine Neiman, personal communication) meaning that whether these results hold

in snails sampled directly from the field remains unclear.

Only asexual lineages of *P. antipodarum* are highly successful invaders; why remains

unclear. Nevertheless, *P. antipodarum* can survive a range of harsh conditions that may

facilitate travel to and establish in new habitats (22, 36). Invaders are susceptible to few

biological enemies (22), passing alive through the digestive systems of trout (37) and are only

rarely infected by parasites (36). Once they invade, these snails can influence ecosystems due

to rapid population growth (22, 38), high population densities (32, 39), and competitive

exclusion of native invertebrates (22, 38). Invasive *P. antipodarum* can even harm the native

predators that ingest them (37).

116   **MATERIALS AND METHODS**

Sample Collection, Processing, and Sequencing

We collected adult *Potamopyrgus antipodarum* from shallow (lake depth < 1 m) rocks

and vegetation from three New Zealand sites in January 2015 (Table 1; Fig. S1) and ten

European collection sites (Table 1; Fig. S1) in spring 2016. Snails were maintained in 15 L tanks

separated by source population in a 16° C room with a 16:8 hour light:dark cycle for less than

one month before dissection. Snails were fed dried *Spirulina* cyanobacteria *ad libitum* (40).

All snails were sexed prior to DNA extraction and assessed for *A. winterbourni* infection

based on the presence of metacercariae cysts via dissection (41) (Table 1). All metacercariae

were removed from infected snails using a micropipette. After dissection, we used flow

cytometry (28, 42) to determine ploidy status as a proxy for reproductive mode of New Zealand

samples (diploids are sexual and polyploids are asexual). Because it is already established that

the European invasive lineages are virtually all polyploid asexuals (17, 43), we did not perform

flow cytometry on these samples.

We used the Qiagen DNeasy Plant Mini Kit (QIAGEN Inc.) for DNA extractions, following

manufacturer protocol, but eluting DNA in 40 µl 100:1 TE buffer. Samples with a 260/280 ratio

> 1.6 and containing > 20 ng of total DNA, based on Nanodrop® 1000 (Thermo Fisher Scientific),

were analysed on a 1% agarose gel. Samples that reached our quality criteria and produced

clear gel bands were shipped to the W.M. Keck Center for Comparative Functional Genomics

(University of Illinois at Urbana-Champaign) for sequencing. See Electronic Supplementary

Methods for more details about sample collection and processing.

The 16S rRNA V4 region was amplified from the *P. antipodarum* microbiome gDNA using

the 515F Golay-barcoded primers and 806R primers (44, 45). Samples were prepared in

accordance with the standard Earth Microbiome Project 16S rRNA protocol (46). Please see our

Electronic Supplementary Methods for PCR reaction mixtures and thermocycler conditions.

gDNA was quantified using the Qubit 2.0 fluorometer (Thermofisher Scientific,), amplicons were

pooled at equimolar ratios (~240 ng per sample), and amplicon pools were cleaned using the

Qiagen PCR Purification Kit (QIAGEN Inc.). The multiplexed library was quality checked and

sequenced with the MiSeq 2x250 bp PE v2 protocol at the W.M. Keck Center for Comparative

Functional Genomics (University of Illinois).

Computational and Statistical Analyses

We removed PhiX sequences from sequencing libraries using Bowtie2 (47) (PhiX genome

obtained from: support.illumina.com/sequencing/sequencing_software/igenome.html).

Demultiplexed paired-end fastq files were processed using DADA2 in R (3.4.0), with the

suggested filtering and trimming parameters, as previously described (48). We then merged

paired-end reads and constructed an amplicon sequence variant (ASV) table. We used the

native implementation of the DADA2 Ribosomal Database Project (RDP) naïve Bayesian

classifier (49) trained against the GreenGenes 13.8 release reference fasta

(https://zenodo.org/record/158955#.WQsM81Pyu2w) to classify ASVs taxonomically.

We used phyloseq v. 1.16.2 to perform estimate_richness and vegan's pd function to

calculate alpha diversity measurements of observed ASVs, Shannon's index, PD Whole Tree,

Pielou's evenness, and Chao1 (50), and to perform ordinations using PCoA on unweighted and

weighted UniFrac distance scores (51). We used the Songbird reference frames approach to

calculate taxa differentials (52). Data visualization and statistical analyses were performed in R;

see Electronic Supplemental Methods for details.

We corrected for effects of snail sex, infection status, and reproductive mode (Table 1)

when testing for geographic associations with microbiota by only conducting analyses on

female uninfected asexual (polyploid) snails, allowing us to directly compare native and invasive

snails. For alpha diversity analyses, we rarefied samples to 10,000 ASVs per sample and

discarded two samples that had fewer reads than this threshold. To test covariate effects on

microbiota alpha diversity, we used Welch's Two Sample t-tests and adjusted p-values ("adj-p")

with a Bonferroni correction for multiple tests. We conducted beta diversity analyses on all

ASVs after removing singletons. We normalized ASV counts by adding one and then $\log_e$-

transforming ASV counts (48). To evaluate beta diversity we performed PCoA on the distance

matrices built on the unweighted and weighted UniFrac scores of each sample (51).

We used Analysis of Similarity (ANOSIM; 999 permutations, R-statistics (R^2) and exact p-

values reported in Results) to evaluate effects of geographic location (sample site) on

microbiota beta diversity . Because all European snails were female uninfected asexual snails

and many New Zealand snails were sexual, infected, and/or male, we only performed the

ANOSIM comparing Europe and New Zealand on female uninfected asexual snails.

To analyse population specificity, we focused on the core microbiome of snails within

locations, defined as taxa found in at least 90% of samples from a given location. We calculated

the proportion of shared core microbiome taxa among all sampling locations and also

compared all European samples vs. all New Zealand samples. We used a t-test to compare the

proportion of reads that mapped as core microbiome taxa between the combined Europe and

combined New Zealand sample groupings.

We performed a Permutational Multivariate Analysis of Variance Using Distance

Matrices analysis (ADONIS) to test the effects of reproductive mode, sex, and infection status

on microbiota beta diversity. The ADONIS analysis was limited to the Lake Alexandrina (New

Zealand) sample site as it was the only site for which we were able to obtain all conditions

(sexual, asexual, male, female, infected, and uninfected). ADONIS tests were conducted with

999 permutations.

We employed the Analysis of Composition of Microbes (ANCOM) algorithm to conduct a

differential count analysis (53). These analyses were limited to the Lake Alexandrina (New

Zealand) sample site for the reasons noted above. We tested the effect of sex in uninfected

male vs. uninfected female sexual snails, tested the effect of reproductive mode on uninfected

female sexual vs. uninfected female asexual snails, and tested the effect of infection status on

asexual female trematode-infected vs. asexual female uninfected snails. We used machine

learning to model microbiome classification by geography using female uninfected asexual

snails and ASVs agglomerated phylogenetically using default settings (phyloseq tax_glom; h =

0.2). Machine Learning was trained on a random forest model using the caret package (v6.0-81)

with a test split of 80:20 and fit a random forest classifier over the tuning parameter of snail

origin (New Zealand or Europe). Please see the Electronic Supplemental Methods for more

details on bioinformatic and statistical analyses.

202   **RESULTS**

203   Variation in *P. antipodarum* microbiota was in large part driven by whether a snail was

from Europe or New Zealand (Fig. 1; ANOSIM; R = 0.97*; p* < 0.01; Permutations: 999; ADONIS; R^2

= 0.33, *p* < 0.01; weighted UniFrac dissimilarity distances are presented in Fig. S2). We did not

find significant differences among microbiomes among sample sites within European or within

New Zealand samples. We thus attribute the geographic distance-to-microbiome distance

correlation to major differences between the most geographically distant European and New

Zealand microbiomes. As validation, we performed a Mantel test on coordinate and microbiota

distance matrices and again found geographic distance was significantly associated with

microbiome distance (Mantel test; Observation = 0.13; *p* < 0.01).

212   To identify taxa that significantly predicted ecosystem clustering, we used machine

learning classification via a random forest model to classify snails as sampled from Europe or

New Zealand based on microbiome composition. The model classified samples based on

relative microbial abundance into European or New Zealand ecosystems with 99.6% accuracy.

We corroborated these results using a reference frames approach and identified taxa that are

associated with snails from Europe or New Zealand (52). There were eight phyla that were

exclusively associated with snails from Europe and none that were exclusively associated with

snails from New Zealand (Table 2; Fig. 1). However, there were more Firmicutes taxa associated

with snails from New Zealand (9 taxa) than Europe (6 taxa) (Table 2; Table S1).

To test the degree to which microbiota exhibit host population-specific distributions, we

calculated the core microbiome of snails from Europe and New Zealand, defined at the tip

agglomerated level. European snails had a core microbiome (found in >=90% of samples)

consisting of 30 taxa while snails from New Zealand contained a core of one taxon (Fig 2A),

Arenimonas. This taxon, which defined the New Zealand snail core microbiome, was also a part

of the European snail core microbiome. Our results departed from this prediction of higher core

microbiome specificity within New Zealand sites by showing relatively consistent core

microbiome composition, containing fewer taxa, across the New Zealand sample sites

compared to European sample sites. We found that the core microbiome of European snails

constituted, on average, 13.8% of their microbiome. By contrast, the core microbiome of New

Zealand snails constituted 70.1% of their microbiome, a marked and statistically significant

difference between native and invasive snails (Fig 2B; T-test; $p < 0.01$). Proteobacteria were the

main constituent of the core microbiota in both snail populations, making up 100% (1/1) of the

New Zealand snail core and 68.9% of the taxa in the European snail core (20/29).

Because the majority of predictive taxa were in higher relative abundance in European

samples and most taxa from the core microbiome of European samples were not found in New

Zealand, we expected that European snail microbiomes would have higher species richness

than the New Zealand snails. Indeed, we found that European snails (mean: 116 ASVs, s.e.:

2.51) had on average 3.52x more ASVs than New Zealand snails (mean: 32.9 ASVs, s.e.: 2.56)

(Fig. 3; T-test; adj-p < 0.01; t = -23.1). There was significantly higher Chao1 and PD Whole Tree

diversity amongst European snail microbiota, suggesting higher richness of rare species at

relatively low abundance and higher phylogenetic diversity in European snail microbiota

compared to New Zealand snail microbiota (Fig. 3; T-test; adj-p values < 0.01; Chao1 t = -25.2;

PD t = -24.8). Shannon diversity was also significantly higher in European snail microbiota (Fig.

3; T-test; adj-p < 0.01; t = -4.52). Our subsequent prediction that New Zealand snail microbiota

may have higher Pielou's evenness than the European snails was upheld (Fig. 3; T-test; adj-p <

0.01; t = 22.4). There were thus more species, more rare species, and higher phylogenetic

richness in European snail microbiota, as well as higher species evenness in New Zealand snail

microbiota.

Our comparisons of snails from Lake Alexandrina show that sex (R^2 = 0.147; adj-p <

0.05), reproductive mode (R^2 = 0.0467; adj-p < 0.05), and infection status (R^2 = 0.0330; adj-p <

0.05) all are associated with microbiota composition (Table 3; ADONIS; Permutations: 999). We

corrected for these covariate effects and highlight taxa that were significantly different by sex

and reproductive mode (Fig. S3; ANCOM; adj-p < 0.05). No taxa significantly differed by

infection status. We identified ten ASVs (all Xanthomonadaecae) that were significantly higher

in abundance in asexual vs. sexual snails (Fig S3). 50 ASVs (*Niabella* (n = 16), *Bacillus* (n = 15),

and OM60 (n = 19)) were significantly more abundant in male vs. female snails. There was

significantly higher ASV richness in uninfected compared to infected snails (T-test; adj-p < 0.01;

Fig S4). There were no significant differences in observed ASV richness when comparing

reproductive mode or sex (Fig. S4).

262   **DISCUSSION**

263   There is a potential for loss of mutualistic microbes upon invasion and the reduced time

for host-microbiota coevolution in the new range. Thus, we predicted that invasion of an

organism would be reflected in its microbiota. The native New Zealand and invasive European

snails had distinct microbiotas. Whilst invasive snails carried the core taxa from New Zealand,

they harboured more core taxa overall and higher phylogenetic diversity. These results suggest

that invaders seem to readily acquire new symbionts, which may play a role in the success of

this worldwide invader. Our results also suggest that the symbionts associated with invasive

lineages may be more opportunistic and less specialized. This pattern may be a consequence of

the comparatively short time for host-microbe coevolution such that host-control mechanisms

have not yet evolved to keep microbial diversity in check (19).

273   The core microbe from native New Zealand snails was maintained in European invasive

snails. This result suggests that there could be selection to maintain certain symbiont

associations in both ranges, which may be important to host biology. Because we were unable

to obtain environmental samples, and it remains unclear whether some core microbes are

inherited, we cannot say whether core microbes found in European and New Zealand

populations were present in both environments, and thus carried over from the latter.

Nevertheless, European samples had 30 times more core taxa overall, and their most predictive

taxa were absent or relatively infrequent in native snails. In other systems, it has been shown

that invaders have lower microbial diversity and/or a subset of microbes relative to that

present in native populations (15, 16). Invasive *P. antipodarum* populations feature low

genotypic diversity relative to the native range (31, 54, 55) and mitochondrial haplotypes of

invaders represent genetic subsets of native mitochondrial haplotype diversity (17). However,

this genetic subsampling in invasive *P. antipodarum* populations is not reflected in their

microbiota diversity.

Across both ranges, proteobacteria dominate among the core microbes. Proteobacteria

and the majority of the other core microbial taxa observed here are consistent with Takacs-

Vesbach et al. (26). These results suggest a shared coevolutionary history between snails and

microbial symbionts has driven the loss of non-essential partners, and ultimately streamlined

the microbiome down to specialized taxa (19). A similar phenomenon occurs in a number of

insect systems, whereby the host harbours few core microbes that are key players in

organismal function (56, 57). Conversely, other insect taxa are host to relatively diverse core

microbial populations (58).

It is well established that antagonistic coevolutionary interactions such as between

hosts and pathogens can drive rapid evolution and diversification (59, 60). In the case of

mutualisms, theory has predicted relatively low rates of evolution compared to antagonistic

relationships (61, 62), although empirical evidence in support of this conjecture remains mixed

(63). For example, mutualisms can drive diversification, such as that observed in plant-animal

mutualisms (64). Even in the absence of host evolution, beneficial resident microbes can evolve

within days (65). Furthermore, competitive interactions between microbes that overlap in

resource use can also drive diversification (20).

Coevolution between *P. antipodarum* and *A. winterbourni* is well documented (27, 29,

30). Reciprocal antagonistic selection between *P. antipodarum* and *A. winterbourni* is expected

to be strong because infected snails are completely sterilized (66), and parasites die if they

encounter resistant individuals (67). In other systems, the potential for mutualist microbes to

be involved in host defence and even host-parasite coevolution has been demonstrated (68,

69). However, the extent to which coevolution operates in diverse communities that feature

both mutualists and antagonists is unclear. Our results do not suggest that the microbiota is a

dominant player in this host-parasite coevolutionary relationship: some taxa differed only

marginally, but not statistically significantly, in relative abundance between infected and

uninfected snails, although we did observe significantly higher ASV richness in uninfected snails

compared to infected snails.

In other systems, it does seem that microbial diversity could confer some degree of

parasite resistance (70), perhaps analogous to the host genetic diversity conferred by

outcrossing in *P. antipodarum* (71). However, that asexual *P. antipodarum* are more successful

invaders strongly suggests that this diversity cannot outweigh the costs of being sexual. Here,

we found that reproductive mode was associated with microbiome composition and ten ASVs

that were significantly more abundant in asexuals than sexuals. Takacs-Vesbach et al. (26)

observed significant differences in microbiome composition between sexuals and asexuals and

among lake populations of *P. antipodarum*. Because the mechanism(s) leading to transitions

from diploid sexuality to polyploid asexuality in *P. antipodarum* are still unclear, future work

could focus on the possible links between microbes and reproductive mode and/or

polyploidization. For example, *Wolbachia*-mediated transitions to asexuality are well

characterized in arthropods (72). Besides a handful of studies (73, 74), however, very little is

known about potential links between host ploidy level and microbiome composition, especially

in animals.

While it is established that symbiotic microbes have important effects on host fitness,

the relationship between microbes and molluscan invaders has received little attention. Here,

we find that the process of invasion is associated with some carryover of a core microbiota, but

mostly involves the formation of numerous new symbiotic relationships, suggesting that

invasive *P. antipodarum* are able to take advantage of readily available microbes. Despite some

impact of biological trait variation on the microbiota in native snail populations, there is a

consistent and strong pattern suggesting that host-microbiota coevolution (or host evolution

alone (19)) streamlines the microbiota of native snails to essential members. Future research

will need to establish the fitness impacts of those new microbes on invaders, and whether the

invading snails have adapted to accommodate them. Notwithstanding this possibility, we show

that the potential for invasion to alter the microbiome is clear.

ACKNOWLEDGEMENTS

We thank Katelyn Larkin, Dunja Lamatsch, Tanja Schwander, Kaitlin Hatcher, Bennett Brown,

Kyle McElroy, and Curtis Bankers for assistance with snail field collections. We are grateful to

the King Lab for feedback on the results, especially Kieran Bates for manuscript comments.

Sequencing was conducted at the W.M. Keck Center for Comparative Functional Genomics

(University of Illinois). Flow cytometry data were obtained at the Flow Cytometry Facility, a

Carver College of Medicine/Holden Comprehensive Cancer Center core research facility at the

University of Iowa.

349   **FUNDING**

350 This work was supported by the University of Iowa T. Anne Cleary International Dissertation

351 Research Fellowship (LB), the University of Iowa Center for Global and Regional Environmental

352 Research Travel Award (LB), a BBSRC Doctoral Training Grant (BB/M011224/1, DD), and by a

353 British Ecological Society Grant (KCK).

355   **DATA AVAILABILITY**

356 Raw sequencing data will be made publicly available on NCBI Sequence Read Archive and

357 accession numbers will be added upon article acceptance.

359   **AUTHOR CONTRIBUTIONS**

360 KCK conceived the project. KCK, LB, DD, MN, and GH designed project. LB, MN, and KCK

361 conducted field collections. LB, DD, CAT, and CF performed DNA extraction, flow cytometry,

362 and prepared sequencing libraries. DD carried out the bioinformatics analysis. The manuscript

363 was written by LB, DD, and KCK with input and approval from the other authors.

 FIGURE AND TABLE CAPTIONS
365

Fig 1. Snail microbiota ecosystem clustering. Biplot PCoA on unweighted UniFrac dissimilarity
distances between snail microbiota profiles across European and New Zealand sample sites.
Each point represents a single snail and are colour coded based on whether a point is a
microbial taxon (grey filled with outline colour indicating sample site) or snail (colour filled by
sample site) and indicates collection site of the snail. Taxa represented on the plot are the top
20-ranked differentials in predicting whether a snail was from New Zealand or Europe and are
annotated at the deepest available taxonomic level. Density plot below PCo1 shows snail
sample density across PCo1 labelled by sample site. Density plots to the left of PCo2 show snail
sample density across PCo2 labelled by sex (male or female), infection status (infected or
uninfected), and reproductive mode (sexual or asexual).
376

377 **Figure 2. Core microbiota of European and New Zealand snails**. (A) 16S rRNA phylogeny of core
378 microbiota (>90% samples) of European (EU) and New Zealand (NZ) snails. Fractional
379 abundance averaged across samples within countries is represented by dot size. (B) Violin plot
380 summarizing the proportion of amplicon reads mapped to core. Significance is denoted as *** =
381 *p* < 0.001 (T-test).
382

383 **Fig 3. Alpha diversity measurements across sample sites.** Figures are faceted by diversity
384 metric. X-axis shows sampling site. Y-axis shows value of diversity metric. NZ = New Zealand
385 samples and EU = European samples.
386

387 **Table 1. Sample collection sites.** The coordinates and location names of sample sites for both
388 native and invasive snails are list below. Ploidy is a proxy for reproductive mode; diploids are
389 sexual, polyploids are asexual. We also include the sex, infection status, and number of
390 individuals sequenced from each condition and location and condition.
391

392 **Table 2. Europe and New Zealand microbiome phyla predictors.** We used a multinomial
393 regression to predict whether snails were from Europe or New Zealand samples sites.
394 Represented here is a summary of counts of phyla that were associated with snails from EU or
395 NZ.
396

397 **Table 3. ADONIS testing effects of sex, ploidy, and infection status on microbiome UniFrac**
398 **distances.** Results from Analysis of Variance Using Distance Matrices (ADONIS) testing of the
399 effects of sex, ploidy, and infection status on unweighted microbiome UniFrac distances
400 between snails. All tests were performed on snails from Lake Alexandrina with 999
401 permutations.
402

REFERENCES

403
404
405   1.   Archie EA, Theis KR. Animal behaviour meets microbial ecology. Animal Behaviour.
406   2011;82(3):425-36.
407   2.   Brucker RM, Bordenstein SR. Speciation by symbiosis. Trends in ecology & evolution.
408   2012;27(8):443-51.
409   3.   Adams D, Douglas AE. How symbiotic bacteria influence plant utilisation by the
410   polyphagous aphid, Aphis fabae. Oecologia. 1997;110(4):528-32.
411   4.   Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I, Rosenberg E. Commensal
412   bacteria play a role in mating preference of Drosophila melanogaster
413   Proceedings of the National Academy of Sciences. 2010;107(46):20051.
414   5.   Koch H, Schmid-Hempel P. Socially transmitted gut microbiota protect bumble bees
415   against an intestinal parasite. Proceedings of the National Academy of Sciences.
416   2011;108(48):19288.
417   6.   Aires T, Serrão EA, Engelen AH. Host and Environmental Specificity in Bacterial
418   Communities Associated to Two Highly Invasive Marine Species (Genus Asparagopsis).
419   Frontiers in microbiology. 2016;7:559.
420   7.   Brucker RM, Bordenstein SR. The Hologenomic Basis of Speciation: Gut Bacteria Cause
421   Hybrid Lethality in the Genus Nasonia. Science. 2013;341(6146):667.
422   8.   Parker BJ, Hrček J, McLean AHC, Godfray HCJ. Genotype specificity among hosts,
423   pathogens, and beneficial microbes influences the strength of symbiont-mediated
424   protection. Evolution; international journal of organic evolution. 2017;71(5):1222-31.
425   9.   Cong X, Xu W, Janton S, Henderson WA, Matson A, McGrath JM, et al. Gut Microbiome
426   Developmental Patterns in Early Life of Preterm Infants: Impacts of Feeding and Gender.
427   PLOS ONE. 2016;11(4):e0152751.
428   10.   Jani AJ, Briggs CJ. The pathogen Batrachochytrium dendrobatidis disturbs the frog skin
429   microbiome during a natural epidemic and experimental infection. Proceedings of the
430   National Academy of Sciences. 2014;111(47):E5049.
431   11.   Wong S, Rawls JF. Intestinal microbiota composition in fishes is influenced by host ecology
432   and environment. Molecular Ecology. 2012;21(13):3100-2.
433   12.   Macke E, Callens M, De Meester L, Decaestecker E. Host-genotype dependent gut
434   microbiota drives zooplankton tolerance to toxic cyanobacteria. Nature Communications.
435   2017;8(1):1608.
436   13.   Corbin C, Heyworth ER, Ferrari J, Hurst GDD. Heritable symbionts in a world of varying
437   temperature. Heredity (Edinb). 2017;118(1):10-20.
438   14.   Coats VC, Rumpho ME. The rhizosphere microbiota of plant invaders: an overview of
439   recent advances in the microbiomics of invasive plants. Frontiers in microbiology.
440   2014;5:368.
441   15.   Minard G, Tran FH, Van VT, Goubert C, Bellet C, Lambert G, et al. French invasive Asian
442   tiger mosquito populations harbor reduced bacterial microbiota and genetic diversity
443   compared to Vietnamese autochthonous relatives. Front Microbiol. 2015;6.

16.   Zepeda-Paulo F, Ortiz-Martínez S, Silva AX, Lavandero B. Low bacterial community
diversity in two introduced aphid pests revealed with 16S rRNA amplicon sequencing.
PeerJ. 2018;6:e4725.

17.   Städler T, Frye M, Neiman M, Lively CM. Mitochondrial haplotypes and the New Zealand
origin of clonal European Potamopyrgus, an invasive aquatic snail. Molecular Ecology.
2005;14(8):2465-73.

18.   Himler AG, Adachi-Hagimori T, Bergen JE, Kozuch A, Kelly SE, Tabashnik BE, et al. Rapid
spread of a bacterial symbiont in an invasive whitefly is driven by fitness benefits and
female bias. Science. 2011;332(6026):254-6.

19.   Foster KR, Schluter J, Coyte KZ, Rakoff-Nahoum S. The evolution of the host microbiome
as an ecosystem on a leash. Nature. 2017;548(7665):43-51.

20.   Ghoul M, Mitri S. The Ecology and Evolution of Microbial Competition. Trends in
Microbiology. 2016;24(10):833-45.

21.   Smith SA, Wilson NG, Goetz FE, Feehery C, Andrade SCS, Rouse GW, et al. Resolving the
evolutionary relationships of molluscs with phylogenomic tools. Nature.
2011;480(7377):364-7.

22.   Alonso A, Castro-Díez P. What explains the invading success of the aquatic mud snail
Potamopyrgus antipodarum (Hydrobiidae, Mollusca)? Hydrobiologia. 2008;614(1):107-16.

23.   Clusa L, Miralles L, Basanta A, Escot C, García-Vázquez E. eDNA for detection of five highly
invasive molluscs. A case study in urban rivers from the Iberian Peninsula. PloS one.
2017;12(11):e0188126-e.

24.   King GM, Judd C, Kuske CR, Smith C. Analysis of Stomach and Gut Microbiomes of the
Eastern Oyster (Crassostrea virginica) from Coastal Louisiana, USA. PLOS ONE.
2012;7(12):e51475.

25.   Neu AT, Allen EE, Roy K. Diversity and composition of intertidal gastropod microbiomes
across a major marine biogeographic boundary. Environmental Microbiology Reports.
2019;11(3):434-47.

26.   Takacs-Vesbach C, King K, Van Horn D, Larkin K, Neiman M. Distinct Bacterial Microbiomes
in Sexual and Asexual Potamopyrgus antipodarum, a New Zealand Freshwater Snail. PLoS
One. 2016;11(8):e0161050.

27.   Lively CM. Evidence from a New Zealand snail for the maintenance of sex by parasitism.
Nature. 1987;328(6130):519-21.

28.   Neiman M, Paczesniak D, Soper DM, Baldwin AT, Hehman G. Wide variation in ploidy level
and genome size in a New Zealand freshwater snail with coexisting sexual and asexual
lineages. Evolution; international journal of organic evolution. 2011;65(11):3202-16.

29.   Dybdahl MF, Lively CM. The geography of coevolution: comparative population structures
for a snail and its trematode parasite. Evolution; international journal of organic
evolution. 1996;50(6):2264-75.

30.   King KC, Delph LF, Jokela J, Lively CM. The geographic mosaic of sex and the Red Queen.
Current biology : CB. 2009;19(17):1438-41.

31.   Verhaegen G, McElroy KE, Bankers L, Neiman M, Haase M. Adaptive phenotypic plasticity
in a clonal invader. Ecology and evolution. 2018;8(9):4465-83.

32.   Alonso Á, Castro-Díez P. The exotic aquatic mud snail Potamopyrgus antipodarum
487   (Hydrobiidae, Mollusca): state of the art of a worldwide invasion. Aquatic Sciences.
488   2012;74(3):375-83.
489   33.   Żbikowski J, Żbikowska E. Invaders of an invader – Trematodes in Potamopyrgus
490   antipodarum in Poland. Journal of Invertebrate Pathology. 2009;101(1):67-70.
491   34.   Winterbourn MJ. The systematics of the New Zealand species of Potamopyrgus (Mollusca:
492   Hydrobiidae) and studies on the biology of Potamopyrgus antipodum : a thesis presented
493   in partial fulfilment of the requirements for the degree of Doctor of Philosophy in Zoology
494   at Massey University [Doctoral]: Massey University; 1968.
495   35.   Blasco-Costa I, Seppälä K, Feijen F, Zajac N, Klappert K, Jokela J. A new species of
496   Atriophallophorus Deblock & Rosé, 1964 (Trematoda: Microphallidae) described
497   from in vitro-grown adults and metacercariae from Potamopyrgus antipodarum (Gray,
498   1843) (Mollusca: Tateidae). Journal of Helminthology. 2020;94:e108.
499   36.   Gérard C, Blanc A, Costil K. Potamopyrgus antipodarum(Mollusca:Hydrobiidae) in
500   continental aquatic gastropod communities: impact of salinity and trematode parasitism.
501   Hydrobiologia. 2003;493(1):167-72.
502   37.   Vinson MR, Baker MA. Poor Growth of Rainbow Trout Fed New Zealand Mud Snails
503   Potamopyrgus antipodarum. North American Journal of Fisheries Management.
504   2008;28(3):701-9.
505   38.   Quinn GP, Lake PS, Schreiber ESG. Littoral benthos of a Victorian lake and its outlet
506   stream: Spatial and temporal variation. Australian Journal of Ecology. 1996;21(3):292-301.
507   39.   Hall RO, Tank JL, Dybdahl MF. Exotic Snails Dominate Nitrogen and Carbon Cycling in a
508   Highly Productive Stream. Frontiers in Ecology and the Environment. 2003;1(8):407-11.
509   40.   Zachar N, Neiman M. Profound Effects of Population Density on Fitness-Related Traits in
510   an Invasive Freshwater Snail. PLOS ONE. 2013;8(11):e80067.
511   41.   Bankers L, Fields P, McElroy KE, Boore JL, Logsdon JM, Jr., Neiman M. Genomic evidence
512   for population-specific responses to co-evolving parasites in a New Zealand freshwater
513   snail. Molecular Ecology. 2017;26(14):3663-75.
514   42.   Krist AC, Bankers L, Larkin K, Larson MD, Greenwood DJ, Dyck MA, et al. Phosphorus
515   availability in the source population influences response to dietary phosphorus quantity in
516   a New Zealand freshwater snail. Oecologia. 2017;185(4):595-605.
517   43.   Liu HP, Hershler R, Marn J, Worsfold TM. Microsatellite evidence for tetraploidy in
518   invasive populations of the New Zealand mudsnail, Potamopyrgus antipodarum (Gray,
519   1843). Journal of Molluscan Studies. 2012;78(2):227-30.
520   44.   Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene
521   primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial Ecology.
522   2015;75(2):129-37.
523   45.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.
524   Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
525   Proceedings of the National Academy of Sciences. 2011;108(Supplement 1):4516.

46. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-
throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. The
ISME journal. 2012;6:1621.
47. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods.
2012;9(4):357-9.
48. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nature Methods. 2016;13:581.
49. Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, et al. Ribosomal Database Project:
data and tools for high throughput rRNA analysis. Nucleic Acids Res. 2014;42(Database
issue):D633-42.
50. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. PLoS One. 2013;8(4):e61217.
51. Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing microbial
communities. Applied and environmental microbiology. 2005;71(12):8228-35.
52. Morton JT, Marotz C, Washburne A, Silverman J, Zaramela LS, Edlund A, et al. Establishing
microbial composition measurement standards with reference frames. Nature
Communications. 2019;10(1):2719.
53. Mandal S, Van Treuren W, White RA, Eggesbo M, Knight R, Peddada SD. Analysis of
composition of microbiomes: a novel method for studying microbial composition. Microb
Ecol Health Dis. 2015;26:27663.
54. Donne C, Neiman M, Woodell JD, Haase M, Verhaegen G. A layover in Europe:
reconstructing the invasion route of asexual lineages of a New Zealand snail to North
America. Authorea. 2020.
55. Dybdahl MF, Drown DM. The absence of genotypic diversity in a successful
parthenogenetic invader. Biological Invasions. 2011;13(7):1663-72.
56. Chandler JA, Morgan Lang J, Bhatnagar S, Eisen JA, Kopp A. Bacterial Communities of
Diverse Drosophila Species: Ecological Context of a Host–Microbe Model System. PLOS
Genetics. 2011;7(9):e1002272.
57. Raymann K, Moran NA. The role of the gut microbiome in health and disease of adult
honey bee workers. Current Opinion in Insect Science. 2018;26:97-104.
58. Tinker KA, Ottesen EA. The Core Gut Microbiome of the American Cockroach, Periplaneta
americana, Is Stable and Resilient to Dietary Shifts. Applied and Environmental
Microbiology. 2016;82(22):6603.
59. Laine A-L. Role of coevolution in generating biological diversity: spatially divergent
selection trajectories. Journal of Experimental Botany. 2009;60(11):2957-70.
60. Paterson S, Vogwill T, Buckling A, Benmayor R, Spiers AJ, Thomson NR, et al. Antagonistic
coevolution accelerates molecular evolution. Nature. 2010;464(7286):275-8.
61. Bergstrom CT, Lachmann M. The Red King effect: when the slowest runner wins the
coevolutionary race. Proceedings of the National Academy of Sciences of the United
States of America. 2003;100(2):593-8.
62. Veller C, Hayward LK, Hilbe C, Nowak MA. The Red Queen and King in finite populations.
Proceedings of the National Academy of Sciences. 2017;114(27):E5396.

63.   Rubin BE, Moreau CS. Comparative genomics reveals convergent rates of evolution in ant-
569   plant mutualisms. Nat Commun. 2016;7:12679.
570   64.   Weber MG, Agrawal AA. Defense mutualisms enhance plant diversification. Proceedings
571   of the National Academy of Sciences. 2014;111(46):16442.
572   65.   King KC, Brockhurst MA, Vasieva O, Paterson S, Betts A, Ford SA, et al. Rapid evolution of
573   microbe-mediated protection against pathogens in a worm host. The ISME journal.
574   2016;10:1915.
575   66.   Winterbourn MJ. A guide to the freshwater mollusca of New Zealand. Tuatara.
576   1973;20(3):141-59.
577   67.   King KC, Jokela J, Lively CM. Trematode parasites infect or die in snail hosts. Biology
578   letters. 2011;7(2):265-8.
579   68.   Ford SA, King KC. Harnessing the Power of Defensive Microbes: Evolutionary Implications
580   in Nature and Disease Control. PLOS Pathogens. 2016;12(4):e1005465.
581   69.   Vorburger C, Perlman SJ. The role of defensive symbionts in host-parasite coevolution.
582   Biological reviews of the Cambridge Philosophical Society. 2018;93(4):1747-64.
583   70.   Dillon RJ, Vennard CT, Buckling A, Charnley AK. Diversity of locust gut bacteria protects
584   against pathogen invasion. Ecology Letters. 2005;8(12):1291-8.
585   71.   Morran LT, Schmidt OG, Gelarden IA, Parrish RC, Lively CM. Running with the Red Queen:
586   Host-Parasite Coevolution Selects for Biparental Sex. Science. 2011;333(6039):216.
587   72.   Boivin T, Henri H, Vavre F, Gidoin C, Veber P, Candau JN, et al. Epidemiology of asexuality
588   induced by the endosymbiotic Wolbachia across phytophagous wasp species: host plant
589   specialization matters. Molecular Ecology. 2014;23(9):2362-75.
590   73.   Cavé-Radet A, Monard C, El-Amrani A, Salmon A, Ainouche M, Yergeau É. Phenanthrene
591   contamination and ploidy level influence the rhizosphere microbiome of Spartina. bioRxiv.
592   2019:625657.
593   74.   Forrester NJ, Ashman T-L. The direct effects of plant polyploidy on the legume–rhizobia
594   mutualism. Annals of Botany. 2017;121(2):209-20.
595

Fig. 1

Legend:
- female
- male
- infected
- uninfected
- sexual
- asexual

PCo2 [4.6%]

PCo1 [30.4%]

Labels on scatter plot:
Flavobacterium, Sphingopyxis, Legionella, Salmonella, Rhizobiaceae, Undibacterium, Hydrogenophaga, Flavobacterium, Sphingorhabdus, Arcicella, Fluviicola, Verrucomicrobiaceae, Pseudomonas, Bacteria, Caldicellulosiruptor, Thermoanaerobacterium, Bacteria, Bacteria, Pseudoxanthomonas, OLB12

density

Legend:
- Brakel
- Geraardsbergen
- Lake Geneva
- Lake Mondsee
- Melle
- West Holme
- Oxford
- Pulsen
- Bovington
- Wrohe
- Lake Alexandrina
- Lake Kaniere
- Lake Rotoroa

Fig. 2

Fig. 3

Table 1. Sample collection sites.

Country	Location	Geographic Coordinates	Ploidy	Infection	Sex	No. Snails
New Zealand	Lake Alexandrina	43° 56' 12.9" S, 170° 27' 35.4" E	Diploid	Infected	Female	9
			Diploid	Infected	Male	2
			Diploid	Uninfected	Female	22
			Diploid	Uninfected	Male	19
			Polyploid	Unknown	Female	16
			Polyploid	Infected	Female	10
			Polyploid	Uninfected	Female	20
New Zealand	Lake Kaniere	42° 48' 21.8" S, 171° 7' 40.5" E	Diploid	Infected	Female	2
			Diploid	Uninfected	Female	6
New Zealand	Lake Rotoroa	41° 47' 44.5" S, 172° 36' 22.4" E	Diploid	Infected	Female	1
			Diploid	Uninfected	Female	10
			Polyploid	Infected	Female	1
			Polyploid	Uninfected	Female	10
Austria	Lake Mondsee	47° 50' 37.7" N, 13° 20' 30.7" E	Polyploid	Uninfected	Female	17
Belgium	Brakel	50° 48' 12.1" N, 3° 45' 20.9" E	Polyploid	Uninfected	Female	12
Belgium	Geraardsbergen	50° 47' 30.4" N, 3° 54' 15.0" E	Polyploid	Uninfected	Female	13
Belgium	Melle	51° 1' 11.6" N, 3° 48' 59.6" E	Polyploid	Uninfected	Female	12
Germany	Wrohe	54° 16' 7.4" N, 9° 57' 35.9" E	Polyploid	Uninfected	Female	3
Germany	Pulsen	54° 19' 17.3" N, 10° 27' 9.8" E	Polyploid	Uninfected	Female	12
Switzerland	Lake Geneva	46° 30' 45.2" N, 6° 36' 27.4" E	Polyploid	Uninfected	Female	15
United Kingdom	West Holme	50° 40' 20.1" N, 2° 10' 29.03" W	Polyploid	Uninfected	Female	11
United Kingdom	Bovington	50° 41' 26.1" N, 2° 13' 29.02" W	Polyploid	Uninfected	Female	12
United Kingdom	Oxford	51° 45' 6.5" N, 1° 13' 0.35" W	Polyploid	Uninfected	Female	12

Table 2. Europe and New Zealand microbiome phyla predictors.

Phylum	Europe	New Zealand
Acidobacteria	3	0
Actinobacteria	4	0
Armatimonadetes	2	0
Bacteroidetes	31	13
Chlamydiae	0	0
Chloroflexi	0	0
Cyanobacteria	4	3
Deinococcus-Thermus	1	0
Epsilonbacteraeota	1	0
Firmicutes	6	9
Gemmatimonadetes	0	0
Lentisphaerae	0	0
Nitrospirae	1	1
Planctomycetes	2	0
Proteobacteria	65	32
Spirochaetes	1	0
Tenericutes	1	0
Verrucomicrobia	1	2

Table 3. ADONIS testing effects of sex, ploidy, and infection status on microbiome UniFrac distances.

Condition	Df	SS	MS	F Model	R2	P
Sex	1	1.2899	1.28991	14.5946	0.14665	0.001
Reproductive mode	1	0.4105	0.41049	4.6444	0.04667	0.001
Infection	1	0.29	0.29	3.2813	0.03297	0.005
Residuals	77	6.8055	0.08838		0.77371	
Total	80	8.7959	1			