Invasive freshwater snails form novel microbial relationships

Running Title: Native and Invasive snail microbiomes

Bankers L.[1,*,†,‡], Dahan D.[2,†,], Neiman M.[1], Adrian-Tucci C.[1,§], Frost C.[3], Hurst G.D.D[3]., King K.C.[4]

[1]University of Iowa, Department of Biology, Iowa City, IA, USA

[2]Stanford University, School of Medicine, Stanford, CA, USA

[3]University of Liverpool, Institute of Integrative Biology, Liverpool, UK

[4]University of Oxford, Department of Zoology, Oxford, UK

Corresponding author: Laura Bankers: laura.bankers@cuanschutz.edu

[†] LB and DD contributed equally to this work and should be considered joint first authors.
[‡] Current address: University of Colorado Anschutz Medical Campus, Aurora, CO, USA
[§] Current address: 1150 4th St. SW Apt. 607 Washington DC, USA

ABSTRACT

Resident microbes (microbiota) can shape host organismal function and adaptation in the face

of environmental change. Invasion of new habitats exposes hosts to novel selection pressures,

but little is known about the impact of invasion on microbiota and the host-microbiome

relationship after this transition (e.g., how rapidly symbioses are formed, whether microbes

influence invasion success). We used high-throughput 16S rRNA sequencing of New Zealand

(native) and European (invasive) populations of the freshwater snail *Potamopyrgus*

antipodarum and found that while invaders do carry over some core microbial taxa from New

Zealand, most of their microbial community is distinct. This finding highlights that invasions can

result in the formation of novel symbioses. We further show that the native microbiome is

composed of fewer core microbes than the microbiome of invasive snails, suggesting that the

microbiota is streamlined to essential members. Together, our findings demonstrate that

microbiota comparisons across native and invasive populations can reveal the impact of a long

coevolutionary history and specialization of microbes in the native host range, as well as new

associations occurring after invasion. We lay essential groundwork for understanding how

microbial relationships affect invasion success and how microbes may be utilized in the control

of invasive hosts.

Keywords: *Potamopyrgus antipodarum*, invasion, microbiome, coevolution, symbiosis,

infection

INTRODUCTION

All organisms are home to numerous microbes. The relationship between hosts and

their microbial communities (microbiota) can play a role in key host functions (1, 2) from

nutrition (3) to behaviour (4) and immune responses (5). The microbiota has also been shown

to impact host evolution (6, 7). Microbial community composition can be determined by host

traits such as genotype (8), sex (9), infection of the host by parasites or pathogens (10),

environmental conditions (6, 11), and genotype by environment interactions (12). Therefore,

both host organisms and the microbes on which they depend for vital processes can both be

impacted by environmental changes, with potential consequences for host adaptation (13).

When species invade novel habitats, they are exposed to new environmental conditions

and selection pressures. Because host and microbe fitness can be mutually dependent, invasion

is predicted to influence both the host and its microbiome (6). The fitness effects of microbes

on their host may affect invasion success and the potential for subsequent adaptation to a new

habitat (14). The performance of introduced species could also be altered by the availability of

microbes outside their native range if there are core microbes that the host needs to thrive. If

host-associated microbes are not all carried over from the native environment, lost upon

invasion, and/or unavailable in the new environment, the diversity of the microbiomes of

invasive lineages could be low. The invader's microbiome might thus only represent a subset of

that residing in the native lineages (15, 16). This pattern is directly analogous to the loss of

genetic diversity that often characterizes invasive lineages (15, 17). Whether such subsampling

occurs at the microbiome level remains an important open question with implications for host

fitness in the new environment.

New invasions may also affect the level of specialization of microbiota relative to hosts

in the native range. This outcome could be a consequence of the comparatively short time

period for coevolution between the invasive host and newly acquired microbes vs. the microbes

from the native environment (15, 16). These effects of colonization could influence microbiome

structure. These consequences could be harmful for the host, such as in a situation where

suitable mutualistic microbes are not in the new habitat. It is also possible that invasive hosts

form new symbioses with the microbes available (18). A more diverse microbiota may result if

there is decreased immune-mediated control of novel microbes (19). By this logic, we might

therefore expect native host individuals to have a less rich microbiota. In particular, established

and stable microbial communities, in which novel competitive interactions between microbes

have long since played out, are likely to be of lower diversity (20). This prediction might be

particularly likely to hold if the host has evolved to cultivate beneficial associations (19).

Addressing whether this prediction is met requires characterization of how invasion impacts

microbial diversity and composition.

Molluscs are one of the most species-rich phyla (21). They represent an ecologically and

economically important class of invaders (22, 23), however, little is known about the

relationships between molluscs and their symbiotic microbes with the exception of a handful of

recent studies (24, 25). Here, we leveraged a powerful model system for host-parasite

coevolution, the New Zealand freshwater snail *Potamopyrgus antipodarum*, to characterize the

impact of invasion on host microbiota. There is substantial across-lake population variation in

microbiome composition among native *P. antipodarum* (26). *Potamopyrgus antipodarum* in

these native New Zealand populations also feature wide individual-level variation in

reproductive mode (sexual vs. asexual) and ploidy level (sexual snails are diploid while asexual

snails are polyploid (triploid or tetraploid); (27, 28)), and are coevolving with sterilizing

trematode parasites (29, 30). By contrast, invasive lineages of *P. antipodarum* in Europe and

North America are polyploid asexuals with reduced genetic variation compared to the native

range (17, 31) and have escaped from parasite infection (32, 33).

We conducted 16S rRNA sequencing of field-collected *P. antipodarum* from waterbodies

across New Zealand and Europe to assess changes in microbiota diversity and composition

across and within native and invasive host populations. We hypothesized that, 1) invasive *P.*

antipodarum would have a subset of native microbial diversity because invaders are often

subsamples of native diversity, which could also likely lead to lower microbial diversity

(analogous to patterns often described for genetic variation post-invasion). Lower microbial

diversity could arise because they simply have had less time in the new environment to acquire

new symbioses; 2) key symbionts needed to thrive should be maintained in the new

environment; 3) invaders will have fewer conserved microbes than native counterparts as a

consequence of relatively recent contact with symbionts in the new environment, and

therefore, relatively low time for the evolution of specialization. To our knowledge, our study of

P. antipodarum is the first comparison of diversification in host-associated microbial

communities between native and invasive molluscs. We also address associations between the

fundamental and often-variable organismal traits of sex, reproductive mode, and infection

status on microbiota within those host populations. More broadly, the results of this study

provide an important starting point to evaluate whether and how microbes could be utilized in

efforts to control invasive species.

Study System

Native New Zealand *P. antipodarum* are primarily found in freshwater lakes and streams

(34). These snails have gained prominence as a model for studying the evolutionary

maintenance of sexual reproduction (27, 28) as well as host-parasite coevolution (27, 30).

Multiple triploid and tetraploid asexual lineages have been separately derived from diploid

sexual conspecifics (27, 28), and New Zealand populations are commonly infected by a

sterilizing trematode parasite, *Atriophallophorus winterbourni* (formerly *Microphallus livelyi*)

(27, 30, 35). A previous study hinted that the microbiome composition of these snails varies

substantially among native populations and between sexuals and asexuals (26). However, these

snails were either laboratory cultured or housed in a laboratory for several months before

harvest (Maurine Neiman, personal communication) meaning that whether these results hold

in snails sampled directly from the field remains unclear.

Only asexual lineages of *P. antipodarum* are highly successful invaders – why this is

remains unclear. Nevertheless, *P. antipodarum* can survive a range of harsh conditions that may

facilitate travel to and establish in new habitats (22, 36). Invaders are susceptible to few

biological enemies (22), passing alive through the digestive systems of trout (37) and are only

rarely infected by parasites (36). Once they invade, these snails can influence ecosystems due

to rapid population growth (22, 38), high population densities (32, 39), and competitive

exclusion of native invertebrates (22, 38). Invasive *P. antipodarum* can even harm the native

predators that ingest them (37).

123   **MATERIALS AND METHODS**

Sample Collection, Processing, and Sequencing

We collected adult *Potamopyrgus antipodarum* during warm seasons from shallow (lake

depth < 1 m) rocks and vegetation from three New Zealand collection sites (early January 2015)

and ten European collection sites (late May 2016) (Table 1; Fig. S1). Snails were maintained in

identical 15 L tanks separated by source population in a 16° C room with a 16:8 hour light:dark

cycle for less than one month before dissection. Snails were fed dried *Spirulina* cyanobacteria

ad libitum (40).

All snails were sexed and shells were removed prior to DNA extraction and assessed for

A. winterbourni infection based on the presence of metacercariae cysts via dissection (41)

(Table 1). Snails containing non- *A. winterbourni* infections were excluded from the study. All

metacercariae were removed from infected snails using a micropipette. After dissection, we

used flow cytometry (28, 42) to determine ploidy status as a proxy for reproductive mode of

New Zealand samples (diploids are sexual and polyploids are asexual). Because it is already

established that the European invasive lineages are virtually all polyploid asexuals (17, 43), we

did not perform flow cytometry on these samples.

We used the Qiagen DNeasy Plant Mini Kit (QIAGEN Inc.) to extract DNA from whole

snails (shells and metacercariae removed), following manufacturer protocol, but eluting DNA in

40 µl 100:1 TE buffer. The Plant kit was used as it better handles the polysaccharides present in

snail mucus, compared to other DNA extraction kits. Samples with a 260/280 ratio > 1.6 and

containing > 20 ng of total DNA, based on Nanodrop® 1000 (Thermo Fisher Scientific), were

analysed on a 1% agarose gel. Samples that reached our quality criteria and produced clear gel

bands were shipped to the W.M. Keck Center for Comparative Functional Genomics (University

of Illinois at Urbana-Champaign) for sequencing. See Electronic Supplementary Methods for

more details about sample collection and processing.

The 16S rRNA V4 region was amplified from the *P. antipodarum* microbiome gDNA using

the 515F Golay-barcoded primers and 806R primers (44, 45). Samples were prepared in

accordance with the standard Earth Microbiome Project 16S rRNA protocol (46). Please see our

Electronic Supplementary Methods for PCR reaction mixtures and thermocycler conditions.

gDNA was quantified using the Qubit 2.0 fluorometer (Thermofisher Scientific), amplicons were

pooled at equimolar ratios (~240 ng per sample), and amplicon pools were cleaned using the

Qiagen PCR Purification Kit (QIAGEN Inc.). The multiplexed library was quality checked and

sequenced with the MiSeq 2x250 bp PE v2 protocol at the W.M. Keck Center for Comparative

Functional Genomics (University of Illinois).

Computational and Statistical Analyses

We removed PhiX sequences from sequencing libraries using Bowtie2 (47) (PhiX genome

obtained from: support.illumina.com/sequencing/sequencing_software/igenome.html).

Demultiplexed paired-end fastq files were processed using DADA2 in R (3.4.0), with the

suggested filtering and trimming parameters, as previously described (48). We then merged

paired-end reads and constructed an amplicon sequence variant (ASV) table. We used the

native implementation of the DADA2 Ribosomal Database Project (RDP) naïve Bayesian

classifier (49) trained against the GreenGenes 13.8 release reference fasta

(https://zenodo.org/record/158955#.WQsM81Pyu2w) to classify ASVs taxonomically.

We used phyloseq v. 1.16.2 to perform estimate_richness and vegan's pd function to

calculate alpha diversity measurements of observed ASVs, Shannon's index, PD Whole Tree,

Pielou's evenness, and Chao1 (50), and to perform ordinations using PCoA on unweighted and

weighted UniFrac distance scores (51). We used the Songbird reference frames approach to

calculate taxa differentials (52). Data visualization and statistical analyses were performed in R;

see Electronic Supplemental Methods for details.

We controlled for effects of snail sex, reproductive mode, and infection status (Table 1)

when testing for geographic associations with microbiota by only conducting analyses on

female uninfected asexual snails, allowing us to directly compare native and invasive snails

without the confounding factors of sex, reproductive mode, and infection (comparisons of sex,

reproductive mode, and infection status are described below). For alpha diversity analyses, we

rarefied samples to 10,000 ASVs per sample and discarded two samples that had fewer reads

than this threshold. To test covariate effects on microbiota alpha diversity, we used Welch's

180 Two Sample t-tests and adjusted p-values ("adj-p") with a Bonferroni correction for multiple

181 tests. We conducted beta diversity analyses on all ASVs after removing singletons. We

182 normalized ASV counts by adding one and then $\log_e$-transforming ASV counts (48). To evaluate

183 beta diversity we performed PCoA on the distance matrices built on the unweighted and

184 weighted UniFrac scores of each sample (51).

185  We used Analysis of Similarity (ANOSIM; 999 permutations, R-statistics (R^2) and exact p-

186 values reported in Results) to evaluate effects of geographic location (sample site) on

187 microbiota beta diversity. Because all European snails were female uninfected asexual snails

188 and many New Zealand snails were sexual, infected, and/or male, we only performed the

189 ANOSIM comparing Europe and New Zealand on female uninfected asexual snails.

190  To analyse population specificity, we focused on the core microbiome of snails within

191 Europe or New Zealand, defined as taxa found in at least 90% of samples within Europe or New

192 Zealand. We used a t-test to compare the proportion of reads that mapped as core microbiome

193 taxa between the combined Europe and combined New Zealand sample groupings.

194  We performed a Permutational Multivariate Analysis of Variance Using Distance

195 Matrices analysis (ADONIS) to test the effects of reproductive mode, sex, and infection status

196 on microbiota beta diversity. The ADONIS analysis was limited to the Lake Alexandrina (New

197 Zealand) sample site as it was the only site for which we were able to obtain all conditions

198 (sexual, asexual, male, female, infected, and uninfected). ADONIS tests were conducted with

199 999 permutations.

We employed the Analysis of Composition of Microbes (ANCOM) algorithm to conduct a

differential count analysis (53). These analyses were limited to the Lake Alexandrina (New

Zealand) sample site for the reasons noted above. We tested the effect of sex in uninfected

male vs. uninfected female sexual snails, tested the effect of reproductive mode on uninfected

female sexual vs. uninfected female asexual snails, and tested the effect of infection status on

asexual female trematode-infected vs. asexual female uninfected snails. We used machine

learning to model microbiome classification by geography using female uninfected asexual

snails and ASVs agglomerated phylogenetically using default settings (phyloseq tax_glom; h =

0.2). Machine Learning was trained on a random forest model using the caret package (v6.0-81)

with a test split of 80:20 and fit a random forest classifier over the tuning parameter of snail

origin (New Zealand or Europe). Please see the Electronic Supplemental Methods for more

details on bioinformatic and statistical analyses.

213   **RESULTS**

Variation in *P. antipodarum* microbiota was in large part driven by whether a snail was

from Europe or New Zealand (Fig. 1; ANOSIM; R = 0.97*; p* < 0.01; Permutations: 999; ADONIS; R^2

= 0.33, *p* < 0.01; weighted UniFrac dissimilarity distances are presented in Fig. S2). To validate

this analysis, we performed a Mantel test on coordinate and microbiota distance matrices and

again found geographic distance was significantly associated with microbiome distance (Mantel

test; Observation = 0.13; *p* < 0.01). Within the two European and New Zealand regions, we did

not find significant differences in microbial variation among sample sites. We thus attribute the

geographic distance-to-microbiome distance correlation to major differences between

microbiomes from the most geographically distant regions of Europe and New Zealand, rather

than differences among sites within the two larger regions. To confirm that beta diversity

differences between microbiomes of snails from Europe and New Zealand were not due to

uneven sampling (10 European sample sites vs 3 New Zealand samples sites), we permuted

PCoA on the distance matrices built on the unweighted UniFrac on all combinations of three

Europe sites vs. the three New Zealand sites (Fig S3).

To identify taxa that significantly predicted ecosystem clustering, where ecosystems are

defined by the two regions of Europe and New Zealand, we used machine learning classification

via a random forest model to classify snails as sampled from Europe or New Zealand based on

microbiome composition. The model classified samples based on relative microbial abundance

into European or New Zealand ecosystems with 99.6% accuracy. We corroborated these results

using a reference frames approach and identified taxa that are associated with snails from

Europe or New Zealand (52). There were eight phyla that were exclusively associated with snails

from Europe and none that were exclusively associated with snails from New Zealand (Table 2;

Fig. 1). However, there were more Firmicutes taxa associated with snails from New Zealand (9

taxa) than Europe (6 taxa) (Table 2; Table S1).

To test the degree to which microbiota exhibit host population-specific distributions, we

calculated the core microbiome of snails from Europe and New Zealand, defined at the tip

agglomerated level. European snails had a core microbiome (found in >=90% of samples)

consisting of 30 taxa while snails from New Zealand contained a core of one taxon (Fig 2A),

Arenimonas, which was also a part of the European snail sore microbiome. Our results departed

from this prediction of higher core microbiome specificity for sites within the New Zealand

region by showing relatively consistent core microbiome composition, containing fewer taxa,

across the New Zealand sample sites compared to European sample sites. We found that the

core microbiome of European snails constituted, on average, 13.8% of their microbiome. By

contrast, the core microbiome of New Zealand snails constituted 70.1% of their microbiome, a

marked and statistically significant difference between native and invasive snails (Fig 2B; T-test*;*

p < 0.01). Proteobacteria were the main constituent of the core microbiota in both snail

populations, making up 100% (1/1) of the New Zealand snail core and 68.9% of the taxa in the

European snail core (20/29).

After finding that the majority of predictive taxa were in higher relative abundance in

European samples and most taxa from the core microbiome of European samples were not

found in New Zealand, we next evaluated whether there were differences in species richness

between European and New Zealand snails. We found that European snails (mean: 116 ASVs,

s.e.: 2.51) had on average 3.52x more ASVs than New Zealand snails (mean: 32.9 ASVs, s.e.:

2.56) (Fig. 3; T-test; adj-*p* < 0.01; t = -23.1). There was significantly higher Chao1 and PD Whole

Tree diversity amongst European snail microbiota, suggesting higher richness of rare species at

relatively low abundance and higher phylogenetic diversity in European snail microbiota

compared to New Zealand snail microbiota (Fig. 3; T-test; adj-*p* values < 0.01; Chao1 t = -25.2;

PD t = -24.8). Shannon diversity was also significantly higher in European snail microbiota (Fig.

3; T-test; adj-*p* < 0.01; t = -4.52). Our subsequent prediction that New Zealand snail microbiota

may have higher Pielou's evenness than the European snails was upheld (Fig. 3; T-test; adj-p <

0.01; t = 22.4). To confirm that alpha diversity differences between microbiomes of snails from

Europe and New Zealand were not due to uneven sampling (10 European sample sites vs. 3

New Zealand samples sites), we permuted alpha diversity tests on all combinations of three

Europe sites vs. the three New Zealand sites (Fig. S3). Amongst the 600 combination tests,

98.3% remained significant (Table S2; T-test; adj-p < 0.05), where the non-significant findings

were for Shannon diversity comparisons between European sites including West Holme and

Melle compared to New Zealand sites. There were thus more species, more rare species, and

higher phylogenetic richness in European snail microbiota, as well as higher species evenness in

New Zealand snail microbiota.

Our comparisons of snails from Lake Alexandrina show that sex (R^2 = 0.147; adj-p =

0.001), reproductive mode (R^2 = 0.0467; adj-p = 0.001), and infection status (R^2 = 0.0330; adj-p

= 0.005) all are associated with microbiota composition (Table 3; ADONIS; Permutations: 999).

We controlled for these covariate effects and highlight taxa that significantly differed in

abundance by sex and reproductive mode (Fig. S4; ANCOM; adj-p < 0.05). No taxa significantly

differed in abundance based on infection status. We identified ten ASVs (all

Xanthomonadaecae) that were significantly higher in abundance in asexual vs. sexual snails (Fig

S4). We observed 50 ASVs (*Niabella* (n = 16), *Bacillus* (n = 15), and OM60 (n = 19)) that were

significantly more abundant in male vs. female snails. There was significantly higher ASV

richness in uninfected compared to infected snails (T-test; adj-p < 0.01; Fig S5). There were no

significant differences in observed ASV richness when comparing reproductive mode or sex (Fig.

S5).

DISCUSSION

There is a potential for loss of mutualistic microbes upon invasion due to subsampling

and/or the reduced time for host-microbiota coevolution in the new range. Thus, we predicted

that invasion of new habitat by a host organisms would reflected in changes to the composition

and/or diversity of host-associated microbes. The native New Zealand and invasive European

snails had distinct microbiotas. Whilst invasive snails carried the core taxa from New Zealand,

they harboured more core taxa overall and higher phylogenetic diversity. These results suggest

that invaders seem to readily acquire new symbionts, which may play a role in the success of

this worldwide invader. Because numerous new microbes appear to be readily acquired by

invasive snails, our results also suggest that the microbes associated with invasive lineages may

be opportunistic and potentially less specialized. This pattern may be a consequence of the

comparatively short time for host-microbe coevolution, relative to the long coevolutionary time

between microbes and New Zealand snails, such that host-control mechanisms have not yet

evolved to keep microbial diversity in check (19).

The core microbe from native New Zealand snails was found in European invasive snails.

This result suggests that there could be selection to maintain certain symbiont associations in

both ranges, which may be important to host biology. Because we were unable to obtain

environmental samples due to insufficient amounts of DNA in water samples, and it remains

unclear whether some core microbes are inherited, we cannot say whether core microbes

found in European and New Zealand populations were present in both environments, or carried

over from the latter. Nevertheless, European samples had 30 times more core taxa overall, and

their most predictive taxa were absent or relatively infrequent in native snails. In other systems,

it has been shown that invaders have lower microbial diversity and/or a subset of microbes

relative to that present in native populations (15, 16). Invasive *P. antipodarum* populations

feature low genotypic diversity relative to the native range (31, 54, 55) and mitochondrial

haplotypes of invaders represent genetic subsets of native mitochondrial haplotype diversity

(17). However, this genetic subsampling in invasive *P. antipodarum* populations is not reflected

in their microbiota diversity.

Across both ranges, proteobacteria dominate among the core microbes. Proteobacteria

and the majority of the other core microbial taxa observed here are consistent with Takacs-

Vesbach et al. (26). The predominance of proteobacteria among the core microbes suggest a

shared coevolutionary history between snails and microbial symbionts has driven the loss of

non-essential partners, and ultimately streamlined the microbiome down to specialized taxa

(19). A similar phenomenon occurs in a number of insect systems, whereby the host harbours

few core microbes that are key players in organismal function (56, 57). Conversely, other insect

taxa are host to relatively diverse core microbial populations (58).

It is well established that antagonistic coevolutionary interactions such as between

hosts and pathogens can drive rapid evolution and diversification (59, 60). In the case of

mutualisms, theory has predicted relatively low rates of evolution compared to antagonistic

relationships (61, 62), although empirical evidence in support of this conjecture remains mixed

(63). For example, mutualisms can drive diversification, such as that observed in plant-animal

mutualisms (64). Even in the absence of host evolution, beneficial resident microbes can evolve

within days (65). Furthermore, competitive interactions between microbes that overlap in

resource use can also drive diversification (20).

Coevolution between *P. antipodarum* and *A. winterbourni* is well documented (27, 29,

30). Reciprocal antagonistic selection between *P. antipodarum* and *A. winterbourni* is expected

to be strong because infected snails are completely sterilized (66), and parasites die if they

encounter resistant individuals (67). In other systems, the potential for mutualist microbes to

be involved in host defence and even host-parasite coevolution has been demonstrated (68,

69). However, the extent to which coevolution operates in diverse communities that feature

both mutualists and antagonists is unclear. Our results do not suggest that the microbiota is a

dominant player in this host-parasite coevolutionary relationship: some taxa differed only

marginally, but not statistically significantly, in relative abundance between infected and

uninfected snails, although we did observe significantly higher ASV richness in uninfected snails

compared to infected snails.

In other systems, it does seem that microbial diversity could confer some degree of

parasite resistance (70), perhaps analogous to the host genetic diversity conferred by

outcrossing in *P. antipodarum* (71). However, asexual *P. antipodarum* are more successful

invaders strongly, which suggests that this diversity cannot outweigh the costs of being sexual.

Here, we found that reproductive mode was associated with microbiome composition and ten

ASVs that were significantly more abundant in asexuals than sexuals. Takacs-Vesbach et al. (26)

observed significant differences in microbiome composition between sexuals and asexuals and

among lake populations of *P. antipodarum*. Because the mechanism(s) leading to transitions

from diploid sexuality to polyploid asexuality in *P. antipodarum* are still unclear, future work

could focus on the possible links between microbes and reproductive mode and/or

polyploidization. For example, *Wolbachia*-mediated transitions to asexuality are well

characterized in arthropods (72). In regard to potential effects of ploidy level, Takacs-Vesbach

et al. (26) did not observe significant differences in in microbiome composition between

asexual triploids and tetraploids, and we were unable to obtain sufficient sampling of

tetraploids to perform ploidy level comparisons. Furthermore, besides a handful of studies (73,

74), very little is known about potential links between host ploidy level and microbiome

composition, especially in animals

While it is established that symbiotic microbes have important effects on host fitness,

the relationship between microbes and molluscan invaders has received little attention. Here,

we find that the process of invasion is associated with some carryover of a core microbiota, but

mostly involves the formation of numerous new symbiotic relationships, suggesting that

invasive *P. antipodarum* are able to take advantage of readily available microbes. Despite some

impact of biological trait variation on the microbiota in native snail populations, there is a

consistent and strong pattern suggesting that host-microbiota coevolution (or host evolution

alone (19)) streamlines the microbiota of native snails to essential members. Future research

will need to establish the fitness impacts of those new microbes on invaders, and whether the

invading snails have adapted to accommodate them, which could be accomplished by

reciprocal microbial transplant experiment in which germ-free snails are exposed to snail-

associated microbes from different environments. Notwithstanding this possibility, we show

that the potential for invasion to alter the microbiome is clear. Future research seeking to

control invasive species should focus on the new associations being formed with microbes

which might be contributing substantially to invader success.

374   **DATA AVAILABILITY**

375 Raw sequencing data will be made publicly available on NCBI Sequence Read Archive and

376 accession numbers will be added upon article acceptance.

378   **ACKNOWLEDGEMENTS**

We thank Katelyn Larkin, Dunja Lamatsch, Tanja Schwander, Kaitlin Hatcher, Bennett Brown,

Kyle McElroy, and Curtis Bankers for assistance with snail field collections. We are grateful to

the King Lab for feedback on the results, especially Kieran Bates for manuscript comments.

Sequencing was conducted at the W.M. Keck Center for Comparative Functional Genomics

(University of Illinois). Flow cytometry data were obtained at the Flow Cytometry Facility, a

Carver College of Medicine/Holden Comprehensive Cancer Center core research facility at the

University of Iowa. This work was supported by the University of Iowa T. Anne Cleary

International Dissertation Research Fellowship (LB), the University of Iowa Center for Global

and Regional Environmental Research Travel Award (LB), a BBSRC Doctoral Training Grant

(BB/M011224/1, DD), and by a British Ecological Society Grant (KCK).

 FIGURE AND TABLE CAPTIONS

Fig 1. Snail microbiota ecosystem clustering. Biplot PCoA on unweighted UniFrac dissimilarity
distances among snail microbiota profiles across European and New Zealand sample sites. Each
point represents a single snail and points are colour coded based on whether a point is a
microbial taxon (grey filled with outline colour indicating sample site) or snail (colour filled by
sample site) and indicates collection site of the snail. Taxa represented on the plot are the top
20-ranked differentials in predicting whether a snail was from New Zealand or Europe and are
annotated at the deepest available taxonomic level. Density plot below PCo1 shows snail
sample density across PCo1 labelled by sample site. Density plots to the left of PCo2 show snail
sample density across PCo2 labelled by sex (male or female), infection status (infected or
uninfected), and reproductive mode (sexual or asexual).

Figure 2. Core microbiota of European and New Zealand snails. (A) 16S rRNA phylogeny of core
microbiota (>90% samples) of European (EU) and New Zealand (NZ) snails. Fractional
abundance averaged across samples within the EU and NZ regions is represented by dot size.
(B) Violin plot summarizing the proportion of amplicon reads mapped to core. Significance is
denoted as *** = $p < 0.001$ (T-test).

Fig 3. Alpha diversity measurements across sample sites. Figures are faceted by diversity
metric. X-axis shows sampling site. Y-axis shows value of diversity metric. NZ = New Zealand
samples and EU = European samples.

Table 1. Sample collection sites. The coordinates and location names of sample sites for both
native and invasive snails are list below. Ploidy is a proxy for reproductive mode; diploids are
sexual, polyploids are asexual. We also include the sex, infection status, and number of
individuals sequenced from each condition and location and condition.

Table 2. Europe and New Zealand microbiome phyla predictors. We used a multinomial
regression to predict whether snails were from Europe or New Zealand samples sites.
Represented here is a summary of counts of phyla that were associated with snails from EU or
NZ.

Table 3. ADONIS testing effects of sex, ploidy, and infection status on microbiome UniFrac
distances. Results from Analysis of Variance Using Distance Matrices (ADONIS) testing of the
effects of sex, ploidy, and infection status on unweighted microbiome UniFrac distances
between snails. All tests were performed on snails from Lake Alexandrina with 999
permutations.

429
430
431 **Fig. S1. Map of sampling sites**. (A) Locations of our ten European sampling sites. (B) Locations
432 of our three New Zealand sampling sites. Country names and boundaries are in yellow.
433 Sampling sites are represented by red dots and the location names are in white. Zoomed out
434 inlayed maps show the broader regions with the sampled regions highlighted in the red boxes.
435 Maps were made using Google Earth Pro v. 7.3.2.5776.
436
Fig. S2. Snail microbiota ecosystem clustering. Biplot PCoA on weighted UniFrac dissimilarity
distances between snail microbiota profiles across European and New Zealand sampling sites.
Each point represents a single snail, and points are colour-coded by whether a point is a
microbial taxon or snail and indicates snail collection site. Taxa represented on the plot are the
top 20-ranked differentials in predicting whether a snail is from New Zealand or Europe and are
annotated at the deepest available taxonomic level. Density plot below PCo1 shows snail
sample density across PCo1 labelled by sampling site. Density plots to the left of PCo2 show
snail sample density across PCo2 labelled by sex (male or female), infection status (infected or
uninfected), and reproductive mode (sexual or asexual).
446
447 **Fig. S3. Combination PCoAs on unweighted UniFrac dissimilarity distances between snail**
448 **microbiota profiles across European and New Zealand sample sites.** Each point represents a
449 single snail, and points are colour-coded by collection site of snail and shaped by whether a
450 snail is from Europe or New Zealand. Include all combinations of snails from three European
    sites compared to snails from all three New Zealand sites.

Fig. S4. ASVs that significantly differed in abundance in snail microbiota based on
reproductive mode or sex. Points represent significantly differentially abundant taxa and are
plotted by log2fold change in abundance (ANCOM; adj-p < 0.01). Comparisons are
asexual/sexual and male/female. There were no significantly differentially abundant taxa
between infected and uninfected snails. Figure is labelled by phyla and deepest available
taxonomy, where G = genus and F = family. Snails are only from Lake Alexandrina, NZ.
457
458 **Fig. S5. Observed ASVs faceted by infection status, reproductive mode, and sex.** Each point
459 represents a snail sample. Comparing observed ASVs, as a measure of species richness, across
460 snail metadata variables of infection status, reproductive mode, and sex. Data from Lake
461 Alexandrina sails, the only collection site where all metadata variables were tested (T-test; ***
462 = adj-p < 0.01).
463
464 **Table S1.** Complete list of differential taxa.
465
466 **Table S2. Combination alpha diversity comparisons.** Combinations include all combinations of
467 snails from three European sites compared to snails from all three New Zealand sites (T-tests;
468 FDR-corrected p-values).

469   References

1. Archie EA, Theis KR. Animal behaviour meets microbial ecology. Animal Behaviour.
2011;82(3):425-36.
2. Brucker RM, Bordenstein SR. Speciation by symbiosis. Trends in ecology & evolution.
2012;27(8):443-51.
3. Adams D, Douglas AE. How symbiotic bacteria influence plant utilisation by the polyphagous
aphid, Aphis fabae. Oecologia. 1997;110(4):528-32.
4. Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I, Rosenberg E. Commensal bacteria
play a role in mating preference of Drosophila melanogaster. Proceedings of
the National Academy of Sciences. 2010;107(46):20051.
5. Koch H, Schmid-Hempel P. Socially transmitted gut microbiota protect bumble bees against
an intestinal parasite. Proceedings of the National Academy of Sciences.
2011;108(48):19288.
6. Aires T, Serrão EA, Engelen AH. Host and Environmental Specificity in Bacterial Communities
Associated to Two Highly Invasive Marine Species (Genus Asparagopsis). Frontiers in
microbiology. 2016;7:559.
7. Brucker RM, Bordenstein SR. The Hologenomic Basis of Speciation: Gut Bacteria Cause
Hybrid Lethality in the Genus Nasonia. Science. 2013;341(6146):667.
8. Parker BJ, Hrček J, McLean AHC, Godfray HCJ. Genotype specificity among hosts, pathogens,
and beneficial microbes influences the strength of symbiont-mediated protection. Evolution;
international journal of organic evolution. 2017;71(5):1222-31.
9. Cong X, Xu W, Janton S, Henderson WA, Matson A, McGrath JM, et al. Gut Microbiome
Developmental Patterns in Early Life of Preterm Infants: Impacts of Feeding and Gender.
PLOS ONE. 2016;11(4):e0152751.
10.   Jani AJ, Briggs CJ. The pathogen Batrachochytrium dendrobatidis disturbs the frog skin
microbiome during a natural epidemic and experimental infection. Proceedings of the
National Academy of Sciences. 2014;111(47):E5049.
11.   Wong S, Rawls JF. Intestinal microbiota composition in fishes is influenced by host
ecology and environment. Molecular Ecology. 2012;21(13):3100-2.
12.   Macke E, Callens M, De Meester L, Decaestecker E. Host-genotype dependent gut
microbiota drives zooplankton tolerance to toxic cyanobacteria. Nature Communications.
2017;8(1):1608.
13.   Corbin C, Heyworth ER, Ferrari J, Hurst GDD. Heritable symbionts in a world of varying
temperature. Heredity (Edinb). 2017;118(1):10-20.
14.   Coats VC, Rumpho ME. The rhizosphere microbiota of plant invaders: an overview of
recent advances in the microbiomics of invasive plants. Frontiers in microbiology.
2014;5:368.
15.   Minard G, Tran FH, Van VT, Goubert C, Bellet C, Lambert G, et al. French invasive Asian
tiger mosquito populations harbor reduced bacterial microbiota and genetic diversity
compared to Vietnamese autochthonous relatives. Front Microbiol. 2015;6.

16.   Zepeda-Paulo F, Ortiz-Martínez S, Silva AX, Lavandero B. Low bacterial community
diversity in two introduced aphid pests revealed with 16S rRNA amplicon sequencing. PeerJ.
2018;6:e4725.
17.   Städler T, Frye M, Neiman M, Lively CM. Mitochondrial haplotypes and the New Zealand
origin of clonal European Potamopyrgus, an invasive aquatic snail. Molecular Ecology.
2005;14(8):2465-73.
18.   Himler AG, Adachi-Hagimori T, Bergen JE, Kozuch A, Kelly SE, Tabashnik BE, et al. Rapid
spread of a bacterial symbiont in an invasive whitefly is driven by fitness benefits and female
bias. Science. 2011;332(6026):254-6.
19.   Foster KR, Schluter J, Coyte KZ, Rakoff-Nahoum S. The evolution of the host microbiome
as an ecosystem on a leash. Nature. 2017;548(7665):43-51.
20.   Ghoul M, Mitri S. The Ecology and Evolution of Microbial Competition. Trends in
Microbiology. 2016;24(10):833-45.
21.   Smith SA, Wilson NG, Goetz FE, Feehery C, Andrade SCS, Rouse GW, et al. Resolving the
evolutionary relationships of molluscs with phylogenomic tools. Nature.
2011;480(7377):364-7.
22.   Alonso A, Castro-Díez P. What explains the invading success of the aquatic mud snail
Potamopyrgus antipodarum (Hydrobiidae, Mollusca)? Hydrobiologia. 2008;614(1):107-16.
23.   Clusa L, Miralles L, Basanta A, Escot C, García-Vázquez E. eDNA for detection of five
highly invasive molluscs. A case study in urban rivers from the Iberian Peninsula. PloS one.
2017;12(11):e0188126-e.
24.   King GM, Judd C, Kuske CR, Smith C. Analysis of Stomach and Gut Microbiomes of the
Eastern Oyster (Crassostrea virginica) from Coastal Louisiana, USA. PLOS ONE.
2012;7(12):e51475.
25.   Neu AT, Allen EE, Roy K. Diversity and composition of intertidal gastropod microbiomes
across a major marine biogeographic boundary. Environmental Microbiology Reports.
2019;11(3):434-47.
26.   Takacs-Vesbach C, King K, Van Horn D, Larkin K, Neiman M. Distinct Bacterial
Microbiomes in Sexual and Asexual Potamopyrgus antipodarum, a New Zealand Freshwater
Snail. PLoS One. 2016;11(8):e0161050.
27.   Lively CM. Evidence from a New Zealand snail for the maintenance of sex by parasitism.
Nature. 1987;328(6130):519-21.
28.   Neiman M, Paczesniak D, Soper DM, Baldwin AT, Hehman G. Wide variation in ploidy
level and genome size in a New Zealand freshwater snail with coexisting sexual and asexual
lineages. Evolution; international journal of organic evolution. 2011;65(11):3202-16.
29.   Dybdahl MF, Lively CM. The geography of coevolution: comparative population
structures for a snail and its trematode parasite. Evolution; international journal of organic
evolution. 1996;50(6):2264-75.
30.   King KC, Delph LF, Jokela J, Lively CM. The geographic mosaic of sex and the Red Queen.
Current biology : CB. 2009;19(17):1438-41.
31.   Verhaegen G, McElroy KE, Bankers L, Neiman M, Haase M. Adaptive phenotypic
plasticity in a clonal invader. Ecology and evolution. 2018;8(9):4465-83.

32.   Alonso Á, Castro-Díez P. The exotic aquatic mud snail Potamopyrgus antipodarum
(Hydrobiidae, Mollusca): state of the art of a worldwide invasion. Aquatic Sciences.
2012;74(3):375-83.
33.   Żbikowski J, Żbikowska E. Invaders of an invader – Trematodes in Potamopyrgus
antipodarum in Poland. Journal of Invertebrate Pathology. 2009;101(1):67-70.
34.   Winterbourn MJ. The systematics of the New Zealand species of Potamopyrgus
(Mollusca: Hydrobiidae) and studies on the biology of Potamopyrgus antipodum : a thesis
presented in partial fulfilment of the requirements for the degree of Doctor of Philosophy in
Zoology at Massey University [Doctoral]: Massey University; 1968.
35.   Blasco-Costa I, Seppälä K, Feijen F, Zajac N, Klappert K, Jokela J. A new species of
Atriophallophorus Deblock & Rosé, 1964 (Trematoda: Microphallidae) described from in
vitro-grown adults and metacercariae from Potamopyrgus antipodarum (Gray, 1843)
(Mollusca: Tateidae). Journal of Helminthology. 2020;94:e108.
36.   Gérard C, Blanc A, Costil K. Potamopyrgus antipodarum(Mollusca:Hydrobiidae) in
continental aquatic gastropod communities: impact of salinity and trematode parasitism.
Hydrobiologia. 2003;493(1):167-72.
37.   Vinson MR, Baker MA. Poor Growth of Rainbow Trout Fed New Zealand Mud Snails
Potamopyrgus antipodarum. North American Journal of Fisheries Management.
2008;28(3):701-9.
38.   Quinn GP, Lake PS, Schreiber ESG. Littoral benthos of a Victorian lake and its outlet
stream: Spatial and temporal variation. Australian Journal of Ecology. 1996;21(3):292-301.
39.   Hall RO, Tank JL, Dybdahl MF. Exotic Snails Dominate Nitrogen and Carbon Cycling in a
Highly Productive Stream. Frontiers in Ecology and the Environment. 2003;1(8):407-11.
40.   Zachar N, Neiman M. Profound Effects of Population Density on Fitness-Related Traits in
an Invasive Freshwater Snail. PLOS ONE. 2013;8(11):e80067.
41.   Bankers L, Fields P, McElroy KE, Boore JL, Logsdon JM, Jr., Neiman M. Genomic evidence
for population-specific responses to co-evolving parasites in a New Zealand freshwater snail.
Molecular Ecology. 2017;26(14):3663-75.
42.   Krist AC, Bankers L, Larkin K, Larson MD, Greenwood DJ, Dyck MA, et al. Phosphorus
availability in the source population influences response to dietary phosphorus quantity in a
New Zealand freshwater snail. Oecologia. 2017;185(4):595-605.
43.   Liu HP, Hershler R, Marn J, Worsfold TM. Microsatellite evidence for tetraploidy in
invasive populations of the New Zealand mudsnail, Potamopyrgus antipodarum (Gray, 1843).
Journal of Molluscan Studies. 2012;78(2):227-30.
44.   Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R
gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial
Ecology. 2015;75(2):129-37.
45.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.
Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
Proceedings of the National Academy of Sciences. 2011;108(Supplement 1):4516.

46.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-
593   throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. The
594   ISME journal. 2012;6:1621.
595   47.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods.
596   2012;9(4):357-9.
597   48.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
598   resolution sample inference from Illumina amplicon data. Nature Methods. 2016;13:581.
599   49.   Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, et al. Ribosomal Database
600   Project: data and tools for high throughput rRNA analysis. Nucleic Acids Res.
601   2014;42(Database issue):D633-42.
602   50.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis
603   and graphics of microbiome census data. PLoS One. 2013;8(4):e61217.
604   51.   Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing microbial
605   communities. Applied and environmental microbiology. 2005;71(12):8228-35.
606   52.   Morton JT, Marotz C, Washburne A, Silverman J, Zaramela LS, Edlund A, et al.
607   Establishing microbial composition measurement standards with reference frames. Nature
608   Communications. 2019;10(1):2719.
609   53.   Mandal S, Van Treuren W, White RA, Eggesbo M, Knight R, Peddada SD. Analysis of
610   composition of microbiomes: a novel method for studying microbial composition. Microb
611   Ecol Health Dis. 2015;26:27663.
612   54.   Donne C, Neiman M, Woodell JD, Haase M, Verhaegen G. A layover in Europe:
613   reconstructing the invasion route of asexual lineages of a New Zealand snail to North
614   America. Authorea. 2020.
615   55.   Dybdahl MF, Drown DM. The absence of genotypic diversity in a successful
616   parthenogenetic invader. Biological Invasions. 2011;13(7):1663-72.
617   56.   Chandler JA, Morgan Lang J, Bhatnagar S, Eisen JA, Kopp A. Bacterial Communities of
618   Diverse Drosophila Species: Ecological Context of a Host–Microbe Model System. PLOS
619   Genetics. 2011;7(9):e1002272.
620   57.   Raymann K, Moran NA. The role of the gut microbiome in health and disease of adult
621   honey bee workers. Current Opinion in Insect Science. 2018;26:97-104.
622   58.   Tinker KA, Ottesen EA. The Core Gut Microbiome of the American Cockroach,
623   Periplaneta americana, Is Stable and Resilient to Dietary Shifts. Applied and Environmental
624   Microbiology. 2016;82(22):6603.
625   59.   Laine A-L. Role of coevolution in generating biological diversity: spatially divergent
626   selection trajectories. Journal of Experimental Botany. 2009;60(11):2957-70.
627   60.   Paterson S, Vogwill T, Buckling A, Benmayor R, Spiers AJ, Thomson NR, et al.
628   Antagonistic coevolution accelerates molecular evolution. Nature. 2010;464(7286):275-8.
629   61.   Bergstrom CT, Lachmann M. The Red King effect: when the slowest runner wins the
630   coevolutionary race. Proceedings of the National Academy of Sciences of the United States
631   of America. 2003;100(2):593-8.
632   62.   Veller C, Hayward LK, Hilbe C, Nowak MA. The Red Queen and King in finite populations.
633   Proceedings of the National Academy of Sciences. 2017;114(27):E5396.

63. Rubin BE, Moreau CS. Comparative genomics reveals convergent rates of evolution in
ant-plant mutualisms. Nat Commun. 2016;7:12679.
64. Weber MG, Agrawal AA. Defense mutualisms enhance plant diversification. Proceedings
of the National Academy of Sciences. 2014;111(46):16442.
65. King KC, Brockhurst MA, Vasieva O, Paterson S, Betts A, Ford SA, et al. Rapid evolution of
microbe-mediated protection against pathogens in a worm host. The ISME journal.
2016;10:1915.
66. Winterbourn MJ. A guide to the freshwater mollusca of New Zealand. Tuatara.
1973;20(3):141-59.
67. King KC, Jokela J, Lively CM. Trematode parasites infect or die in snail hosts. Biology
letters. 2011;7(2):265-8.
68. Ford SA, King KC. Harnessing the Power of Defensive Microbes: Evolutionary
Implications in Nature and Disease Control. PLOS Pathogens. 2016;12(4):e1005465.
69. Vorburger C, Perlman SJ. The role of defensive symbionts in host-parasite coevolution.
Biological reviews of the Cambridge Philosophical Society. 2018;93(4):1747-64.
70. Dillon RJ, Vennard CT, Buckling A, Charnley AK. Diversity of locust gut bacteria protects
against pathogen invasion. Ecology Letters. 2005;8(12):1291-8.
71. Morran LT, Schmidt OG, Gelarden IA, Parrish RC, Lively CM. Running with the Red
Queen: Host-Parasite Coevolution Selects for Biparental Sex. Science. 2011;333(6039):216.
72. Boivin T, Henri H, Vavre F, Gidoin C, Veber P, Candau JN, et al. Epidemiology of
asexuality induced by the endosymbiotic Wolbachia across phytophagous wasp species: host
plant specialization matters. Molecular Ecology. 2014;23(9):2362-75.
73. Cavé-Radet A, Monard C, El-Amrani A, Salmon A, Ainouche M, Yergeau É. Phenanthrene
contamination and ploidy level influence the rhizosphere microbiome of Spartina. bioRxiv.
2019:625657.
74. Forrester NJ, Ashman T-L. The direct effects of plant polyploidy on the legume–rhizobia
mutualism. Annals of Botany. 2017;121(2):209-20.
661

Fig. 1

Fig. 2

A

Sphingomonadaceae
Rhizorhapis
Sphingopyxis
Sphingorhabdus
Rhizobiaceae
Rhodobacter
Hyphomicrobium
Brevundimonas
Bacteria
Geobacillus
Solitalea
Arcicella
Sphingobacteriales
Chryseobacterium
Bergeyella
Flavobacterium
Flavobacterium
Flavobacterium
Hydrogenophaga
Methylibium
Limnohabitans
Methyloversatilis
Undibacterium
Rheinheimera
Shewanella
Salmonella
Arenimonas
Legionella
Pseudomonas
Pseudomonas

Abundance
- 0.0032
- 0.0080
- 0.0160
- 0.0400
- 0.0800

Location
- EU
- NZ

B

***

Fractional abundance

0.75
0.50
0.25
0.00

Fig. 3

Table 1. Sample collection sites.

Country	Location	Geographic Coordinates	Reproductive Mode	Infection	Sex	No. Snails
New Zealand	Lake Alexandrina	43° 56' 12.9" S, 170° 27' 35.4" E	Sexual	Infected	Female	9
			Sexual	Infected	Male	2
			Sexual	Uninfected	Female	22
			Sexual	Uninfected	Male	19
			Asexual	Unknown	Female	16
			Asexual	Infected	Female	10
			Asexual	Uninfected	Female	20
New Zealand	Lake Kaniere	42° 48' 21.8" S, 171° 7' 40.5" E	Sexual	Infected	Female	2
			Sexual	Uninfected	Female	6
New Zealand	Lake Rotoroa	41° 47' 44.5" S, 172° 36' 22.4" E	Sexual	Infected	Female	1
			Sexual	Uninfected	Female	10
			Asexual	Infected	Female	1
			Asexual	Uninfected	Female	10
Austria	Lake Mondsee	47° 50' 37.7" N, 13° 20' 30.7" E	Asexual	Uninfected	Female	17
Belgium	Brakel	50° 48' 12.1" N, 3° 45' 20.9" E	Asexual	Uninfected	Female	12
Belgium	Geraardsbergen	50° 47' 30.4" N, 3° 54' 15.0" E	Asexual	Uninfected	Female	13
Belgium	Melle	51° 1' 11.6" N, 3° 48' 59.6" E	Asexual	Uninfected	Female	12
Germany	Wrohe	54° 16' 7.4" N, 9° 57' 35.9" E	Asexual	Uninfected	Female	3
Germany	Pulsen	54° 19' 17.3" N, 10° 27' 9.8" E	Asexual	Uninfected	Female	12
Switzerland	Lake Geneva	46° 30' 45.2" N, 6° 36' 27.4" E	Asexual	Uninfected	Female	15
United Kingdom	West Holme	50° 40' 20.1" N, 2° 10' 29.03" W	Asexual	Uninfected	Female	11
United Kingdom	Bovington	50° 41' 26.1" N, 2° 13' 29.02" W	Asexual	Uninfected	Female	12
United Kingdom	Oxford	51° 45' 6.5" N, 1° 13' 0.35" W	Asexual	Uninfected	Female	12

Table 2. Europe and New Zealand microbiome phyla predictors.

Phylum	Europe	New Zealand
Acidobacteria	3	0
Actinobacteria	4	0
Armatimonadetes	2	0
Bacteroidetes	31	13
Chlamydiae	0	0
Chloroflexi	0	0
Cyanobacteria	4	3
Deinococcus-Thermus	1	0
Epsilonbacteraeota	1	0
Firmicutes	6	9
Gemmatimonadetes	0	0
Lentisphaerae	0	0
Nitrospirae	1	1
Planctomycetes	2	0
Proteobacteria	65	32
Spirochaetes	1	0
Tenericutes	1	0
Verrucomicrobia	1	2

Table 3. ADONIS testing effects of sex, ploidy, and infection status on microbiome UniFrac distances.

Condition	Df	SS	MS	F Model	R2	P
Sex	1	1.2899	1.28991	14.5946	0.14665	0.001
Reproductive mode	1	0.4105	0.41049	4.6444	0.04667	0.001
Infection	1	0.29	0.29	3.2813	0.03297	0.005
Residuals	77	6.8055	0.08838	0.77371		
Total	80	8.7959	1			