1 **Title: α-Synuclein facilitates clathrin assembly in synaptic vesicle**
2 **endocytosis**
3
4 Short title: α-Synuclein regulates clathrin assembly
5

6 Karina J Vargas[1*#], P. L. Colosi[1,2*#], Eric Girardi[1*], Sreeganga S Chandra[1,3]
7
8 [1] Departments of Neurology; Neuroscience, Yale University, New Haven, CT
9 06536
10 [2] PREP Program, Yale University, New Haven, CT 06511
11 [3] To whom correspondence should be addressed to:
12 sreeganga.chandra@yale.edu ; 203-785-6172
13
14 *Authors contributed equally to this study
15 # Present Address:
16 KJV, Marine Biological Laboratory, MA
17 PLC, University of Pennsylvania, PA

18
19

20 **Summary** (40 word limit): Synucleins can sense and generate membrane
21 curvature. We previously showed that synuclein null mice exhibit deficits in
22 synaptic vesicle endocytosis. Here, Vargas et al. provide evidence that α-synuclein
23 functions specifically in clathrin assembly during early steps of synaptic vesicle
24 endocytosis.
25
26 **Abstract** (160 word limit):
27
α-Synuclein plays a central role in Parkinson's disease (PD); hence,
elucidating its normal physiological function(s) is important. α-Synuclein and family
members β-, and γ-synuclein, are presynaptically enriched proteins. Synucleins
sense and generate membrane curvature, properties consistent with their
described roles in synaptic vesicle (SV) cycling. We have previously shown SV
endocytosis (SVE) deficits in αβγ-synuclein knockout (KO) neurons. Here, we
investigate which steps of SVE are regulated by α-synuclein. Immuno-electron
microscopy (EM) of synaptosomes reveals that α-synuclein relocalizes from SVs
to the synaptic membrane upon stimulation, allowing α-synuclein to function there
during or after stimulation. Using membrane recruitment assays, we show that α-
synuclein is co-localized with clathrin patches. We also observe that recruitment
of clathrin and its adaptor, AP180, to synaptic membranes is altered in the absence
of synucleins. Visualizing clathrin assembly on membranes in an *in vitro*
reconstitution system reveal that synucleins increase clathrin patch size and
curvature, facilitating clathrin coated pit maturation during the early steps of SVE.

43
44 **Introduction**
45
α-Synuclein became a principal focus of neurodegenerative research when
it was identified as the major constituent of Lewy Bodies, the pathological protein
aggregates found in the brains of PD patients [1]. The importance of α-synuclein
was further underscored by the identification of families with Mendelian forms of
PD arising from causal point mutations and gene multiplications of *SNCA*, the α-
synuclein gene [2-7]. Genome-wide association studies have shown that
sequence variants in *SNCA* are also associated with sporadic PD [8, 9]. Based on
these observations, many current therapeutic strategies for PD are focused on
eliminating or reducing α-synuclein levels in the brain. Therefore, there is a growing
interest in understanding α-synuclein's physiological functions and how loss of α-
synuclein impacts neuronal functions.
57
Since its discovery as a SV-associated protein in the electric organ of
Torpedo [10], several distinct approaches have been used to determine the
physiological function(s) of α-synuclein. Structural studies have revealed that α-
synuclein can adopt several conformations, principally, unfolded in solution, but
also α-helical on phospholipid membranes. In α-helical conformations, the N-
terminus folds into either a single elongated amphipathic helix on flatter
membranes or a 'broken' helix when bound to curved lipid membranes [11, 12]. α-
Synuclein can switch between the two membrane conformations but has a

stronger affinity for highly curved membranes [13]. These biophysical properties
are shared by β-, and γ-synuclein. The conformational plasticity of synucleins
allows these proteins to sense and generate membrane curvature.
69
The initial finding in *Torpedo*, itself suggested an α-synuclein function
related to SV trafficking. The membrane-bending properties of synucleins suggest
that they regulate SV exocytosis and/or endocytosis. Biochemical studies have
indicated a role for α-synuclein in exocytosis [14, 15]. For instance, α-synuclein
has been shown to interact with synaptobrevin2, a v-SNARE protein required for
SV fusion [15-17]. Somewhat surprisingly, electrophysiological studies suggest
that neurotransmission is normal or even improved in αβγ-synuclein KO mice [15,
18, 19], raising questions about the relevance of this interaction for exocytosis.
Other studies have convincingly demonstrated α-synuclein's role in modulating SV
mobilization and recycling, specifically in SVE [20, 21]. Additionally, analysis of
dense core vesicles (which have larger diameter than SVs) in synuclein KOs
suggest a role in fusion pore expansion [22]. Taken together, the preponderance
of functional studies supports synucleins acting post-fusion, in SVE, and in SV
clustering [23].
84
Through pHluorin imaging and cholera toxin labelling, we determined that
αβγ-synuclein KO hippocampal neurons exhibit slower kinetics of SVE [20].
Consistently, our unbiased proteomics of synaptosomes from αβγ-synuclein KO
mice show significant changes in SVE proteins: increases in clathrin and
endophilins and a decrease in levels of the v-SNARE adaptor protein AP180
(neuronal homolog of CALM) [18, 20, 24]. Recent proteomic analyses of α-
synuclein-interacting proteins using proximity labelling in cortical neurons revealed
SVE as one of the most robustly represented pathways [25]. In addition, we have
shown that the interaction between α-synuclein and synaptobrevin2 observed by
other researchers [15, 16, 26] is part of a larger complex of which AP180 is a major
component [20]. AP180 along with AP2 are the main recruiters of clathrin to
synaptic membranes and act to initiate clathrin coated pit (CCP) formation. These
adaptors are critical for both clathrin assembly and SV protein sorting [27-30]. This
suggests that the transient α-synuclein-AP180-synaptobrevin2 complex may play
a role post-fusion in SV protein retrieval.
100
Building on our previous finding of slowed SVE in αβγ-synuclein KOs, we
aimed to identify which step(s) of SVE are regulated by synucleins. We compared
the ultrastructural features of endocytic mutant synapses to those seen in αβγ-
synuclein KOs [18, 20, 21]. AP180 KO mice exhibit a higher density of vesicles
[27] similar to αβγ-synuclein KOs [18]. Dynamin1, endophilins, synaptojanin1, and
auxilin that regulate later stages of endocytosis, such as CCP curvature
generation, scission and uncoating, when deleted show striking synaptic
ultrastructural phenotypes. Dynamin1 KO synapses are unable to scission CCPs
to generate clathrin coated vesicles (CCVs) [31, 32]–consequently, presynaptic
termini become filled with endocytic tubules decorated with CCPs [33].
Contrastingly, endophilin, synaptojanin1, and auxilin KO synapses feature

accumulations of CCVs, due to impaired CCV uncoating. αβγ-synuclein KO
synapses did not display any of the morphological abnormalities
observed–endocytic tubules with CCPs or CCVs–when later stages of SVE are
blocked [20]. This suggests α-synuclein's participation in an earlier stage of SVE
similar to AP180.
117
Here, we performed a wide range of *in vitro* experiments to test whether α-
synuclein modulates clathrin recruitment through its membrane bending properties
and interactions with other proteins. We visualized α-synuclein's subcellular
localization, monitored clathrin polymerization on the membrane, and determined
how clathrin structures are altered by the presence or absence of α-synuclein. In
a minimal reconstitution system, we show that clathrin cage formation is affected
by the addition of recombinant α-synuclein. These experiments collectively
demonstrate that the locus of α-synuclein action in SVE is clathrin assembly.
126
Results and Discussion
The localization of α-synuclein within synapses is dynamic
The majority of α-synuclein is found on SVs at rest [21, 34], consistent with
its preference to bind highly curved membranes [35, 36]. To determine whether
the localization of α-synuclein is changed upon synaptic stimulation, we prepared
fresh synaptosomes from wildtype mouse brains, and tested three physiological
conditions: rest, depolarization with high K^+, and repolarization. The synaptosomes
were then embedded in agarose, fixed, immunolabelled with an α-synuclein or
synaptobrevin2-specific antibody and imaged using EM. At rest, the majority of α-
synuclein is present on SVs (61.1% on SVs, 25.9% on membranes; 5.9% in
cytosol; **Fig 1A**, **B**), similar to the integral SV protein synaptobrevin2 (78.2% on
SV, 12.0% on membranes, 3.1% in cytosol; **Fig 1A**, **C**). These data are in line with
previous publications [21]. Interestingly, upon stimulating synaptosomes with high
K^+, α-synuclein relocates from SVs to the synaptic plasma membrane and cytosol
(30.1% on SVs, 51.1% on membranes, 11.2% in the cytosol; Rest versus K^+
stimulation: 61.1% versus 30.1% on SVs; p<0.001). Our results are congruent with
previous confocal imaging, which showed α-synuclein becoming diffuse upon
neuronal stimulation [37, 38]. As expected, synaptobrevin2 remains mainly on SVs
when stimulated (Rest versus K^+ stimulation: 78.2% versus 68.4% on SVs; p=
0.095). Repolarization leads to relocalization of α-synuclein on SVs (70.1% on
SVs, 18.2% on membranes, 3.8% in cytosol). To validate these results, we
measured the distance from the gold particle to the nearest SV. The distances
measured for α-synuclein at rest, high K^+, and repolarization were 7.73 nm, 11.32
nm and 4.61 nm, while those for synaptobrevin2 were 10.22 nm, 9.47 nm, and 8.8
nm, respectively. Thus, the sub-synaptic location of α-synuclein is dynamic and is
likely to be linked to the SV cycle.
153
Localization of α-synuclein to the synaptic plasma membrane upon
neuronal stimulation strongly supports that α-synuclein functions during SV
endocytosis.   To confirm this, we examined the location of α-synuclein in
dynamin1,3 double KO synapses, which feature few SVs and large CCP-

decorated invaginations of the synaptic plasma membrane. This leads to the
capture of SVE proteins in CCPs, visualized as their apparent clustering in
confocal images. While exocytic and peripheral SV proteins, such as rab3a and
synapsin, appear unchanged or diffuse, respectively [33, 39]. Immunostaining of
dynamin1,3 double KO neurons revealed that α-synuclein behaves as an
endocytic protein, clustering and co-localizing with clathrin and other endocytic
proteins (**Fig. S1**). This is consistent with our previous findings [20] and suggests
that α-synuclein is an accessory endocytic protein that functions prior to dynamin
recruitment.
167
α-Synuclein co-localizes with a fraction of clathrin patches
In order to visualize if α-synuclein co-localizes with clathrin patches at early
stages of clathrin mediated endocytosis, we used a sheared membrane sheet
preparation. Membrane-GFP transfected PTK2 cells were grown on coverslips and
briefly sonicated to unroof the cells, leaving behind adherent membranes. These
coverslips were washed and incubated with cytosol prepared from wildtype mouse
brain and GTPγS (to prevent dynamin function) with an ATP generation system for
increasing amounts of time (t=0-30 minutes). The membranes were then
immunostained for clathrin and α-synuclein to visualize their distribution on the
membrane sheets. As seen in **Fig. 2A-C**, at t=0, both clathrin and α-synuclein are
present in puncta on the membranes. Strikingly, some puncta contain both proteins
in the distinct pattern of a clathrin patch with α-synuclein colocalized at its inner
core (**Fig. 2B**). We measured fluorescence intensity and size of clathrin and α-
synuclein patches for each punctum. We plotted all puncta (n=2071) observed by
their relative clathrin and α-synuclein fluorescence and binned them in 5 size
categories based on the diameter of clathrin patches (**Fig. 2D**). When these data
were plotted as a function of clathrin diameter (**Fig. 2E**), for a given clathrin puncta
size, the average clathrin fluorescence exceeded that of α-synuclein, suggesting
that α-synuclein is found within clathrin patches and not the other way around. As
expected, average clathrin fluorescence increases linearly with puncta size.
However, average α-synuclein fluorescence is largely constant and only increases
in very large patches (diameter > 1.6 μm) (**Fig. 2E**, blue and purple). This indicates
that α-synuclein can colocalize with clathrin, predominantly in large puncta.
191
To classify the puncta into clathrin only, α-synuclein only, and clathrin+α-synuclein
puncta, we set cutoffs at x=0.5 and y=1 (**Fig. 2D**). Using these cutoffs, 21% of
puncta were positive for both proteins, 59% for clathrin only, 20% for α-synuclein
only (**Fig. 2G**; t=0). The average diameters α-synuclein only, clathrin only, and
clathrin+α-synuclein puncta were 0.53 μm, 0.66 μm, and 0.96 μm, respectively
(**Fig. 2F**; n=1174, p=0.001), indicating that clathrin+α-synuclein puncta are
consistently larger than clathrin only puncta.
199
Next, we tested how the distribution and size of the 3 classes of puncta (α-
synuclein only, clathrin only, and clathrin+α-synuclein) change over time (0, 5, 15,
30 minutes) during incubation with brain cytosol (**Fig. 2F, S2A**). Interestingly, we
observe that the fraction of clathrin only puncta decreases (59% at t=0 to 0% at

t=30) with a concomitant increase in clathrin+α-synuclein puncta (20% at t=0 to
50% at t=30), indicating that α-synuclein and clathrin might be co-assembling (**Fig.**
2G, **S2B**). The average size of α-synuclein only and clathrin only puncta remain
constant, while that of clathrin+α-synuclein puncta diameter changes over time
(**Fig. 2F**, **S2A**; 0.96 μm at t=0 and 0.77 μm at t=30), mirroring the dynamics of
clathrin+α-synuclein puncta. It is known that incubation with brain cytosol (t=30
minutes) leads to the growth and curvature of the clathrin lattice to form mature
CCPs that are ready to be scissioned [40]. So, at this time point, clathrin+α-
synuclein puncta may represent these structures.
213
To test if clathrin and α-synuclein co-localize on synaptic membranes, we
purified synaptosomes from the brains of wildtype and αβγ-synuclein KO mice. We
identified proteins in close proximity to α-synuclein, by treating the synaptosomes
with a membrane permeable crosslinker, and the crosslinked proteins were then
immunoprecipitated with an α-synuclein antibody (**Fig. S3**). Unique bands of
immunoprecipitated proteins from wildtype samples were processed for mass
spectroscopy. The αβγ-synuclein KO samples served as a negative control to
identify non-specific interactions. This unbiased analysis revealed that clathrin
heavy chain was present in wildtype samples but absent in the αβγ-synuclein KO
samples (**Fig. S3**), indicating that events observed on PTK2 membranes might
also occur on synaptic membranes.
225
Membrane recruitment of endocytic proteins is altered by the lack of
synucleins
To interrogate α-synuclein involvement in endocytic protein recruitment to
the synaptic membrane, we performed an *in vitro* protein recruitment assay.
Stripped synaptic membranes isolated from wildtype mouse brain were incubated
with cytosol obtained from wildtype or αβγ-synuclein KO mouse brains. αβγ-
synuclein KO mouse brains show compensatory increases in clathrin, endophilin,
and synapsins, decreases in AP180 levels and altered AAK1 splicing (**Fig. 3**; [20]).
The incubations were supplemented with an ATP generation system and GTPγS
similar to the PTK2 membrane experiments (**Fig. 2**). As a positive control, we used
synapsin, whose levels are increased in αβγ-synuclein KO and known to bind
synaptic membranes [41] (**Fig. 3**). As expected, synapsin binds membranes in
proportion to its expression. In the same fashion, we found increased recruitment
of clathrin heavy and light chains when synucleins are absent from the cytosol (1.6
± 0.2 fold, p=0.0061 for clathrin heavy chain; 3.4 ± 0.4 fold, p=0.0016 for clathrin
light chain; **Fig. 3**), largely reflecting the increased clathrin levels in the αβγ-
synuclein KO cytosol [18, 42]. Fascinatingly, we observed increased AP180
recruitment (1.4 ± 0.1 fold, p=0.033), despite decreased AP180 levels in αβγ-
synuclein KO cytosol [20]. We then calculated the clathrin:AP180 ratio on synaptic
membranes and determined it was also increased in αβγ-synuclein KOs (1.14-2.42
fold for heavy and light chain, respectively).
247
Other key endocytic proteins were tested in the membrane recruitment
assay with no significant results. These include AP-2, FBP17, epsin1, Eps-15,

AAK1, endophilin-A1, necap and dynamin1, all proteins that function in SVE. We
also did not observe altered recruitment of Rab3. Based on these findings, we infer
that synucleins may modulate recruitment of clathrin and AP180, proteins involved
in the initiation of SVE.
254
Lipid monolayer assay to investigate clathrin assembly
Our previous SVE kinetic measurements [20] showed slower endocytosis
in αβγ-synuclein KO neurons and now we observe increased membrane
recruitment of clathrin and AP180 *in vitro* (**Fig. 3**). Therefore, we sought to
determine the state of clathrin assembly on the membrane. Also, as the
clathrin:adaptor ratio is one determinant of CCP curvature [43], we wanted to test
if the increased clathrin:AP180 ratio seen in the αβγ-synuclein KO condition
indicates shallower CCPs.
263
In order to visualize clathrin assembly, lipid monolayer assays (**Fig. 4A**)
designed for purified proteins were adapted for brain cytosol [30], and EM was
performed (**Fig. 4B**). The assays were performed with wildtype and αβγ-synuclein
KO cytosol and the time dependence of clathrin assembly into flat lattices and
mature CCPs was quantified. We established that incubation of the lipid monolayer
with wildtype cytosol leads to assembly of flat and curved clathrin lattices, with a
typical hexagonal pattern by 15 minutes (**Fig. 4B**, **C**), in line with previous
publications [30]. On occasion, mature CCPs were also seen (**Fig. 4B**). The size
of individual clathrin patches were quantified. As seen in **Fig. 4**, absence of
synucleins initially leads to significantly smaller clathrin patches (**Fig. 4C**, t=15,
wildtype (WT)= 30,808 ± 2,617 nm^2 and αβγ-synuclein KO (TKO)= 25,544 ± 2,134
nm^2, p<0.001), but by t=23 minutes, the patches are significantly larger (t=23, WT=
36,513 ± 2597 nm^2, and TKO= 64,882 ± 3790 nm^2, p<0.05). Thus, there is delayed
clathrin assembly in the αβγ-synuclein KO, consistent with the functional data in
hippocampal neurons [20]. Since increased clathrin:adaptor ratio is associated
with decreased CCP curvature, we examined the curvature of clathrin patches by
analyzing the fraction of pentagons and heptagons per hexagons in the clathrin
formations (**Fig. 4D**, **E**). The presence of pentagons and/or heptagons indicates
that a curved lattice is being generated. We find that incubation with αβγ-synuclein
KO cytosol leads to fewer non-hexagonal clathrin formations (**Fig. 4E**, **F**),
indicating flatter clathrin lattices.
285
To confirm that clathrin assembly deficits are caused by the absence of
synuclein, we performed rescue experiments, by adding to the αβγ-synuclein KO
cytosol, recombinant mouse α-synuclein matched to the amount in wildtype brain
cytosol. Addition of mouse α-synuclein to the αβγ-synuclein KO cytosol rescued
both the kinetics and the formation of enlarged clathrin patches, at later time points,
to that of wildtype sizes (t=30, WT= 54,434 ± 3,165 nm^2, TKO= 81,753 ± 5,840
nm^2, and TKO + α-synuclein= 58,510 ± 4,127 nm^2; WT versus TKO + α-synuclein,
ns) (**Fig. 4B**, **C**). In addition, there was a greater number of lattices with pentagons
and heptagons, which are associated with curved lattices (**Fig. 4E**, **F**). Together,

these results strongly suggest that synucleins function to assemble clathrin and
curve the lattice.
297
To unequivocally establish α-synuclein's role in clathrin assembly on the
membrane, we performed the lipid monolayer assay (**Fig. 4A**) using purified
proteins in place of cytosol (**Fig. 5A**). Addition of clathrin has no observable effect
on membranes due to its inability to directly bind them (**Fig. 5B**), while AP180 can
bind the lipid monolayer and modestly deform it (**Fig. 5B**). Injection of recombinant
mouse α-synuclein only leads to membrane ruffles and deformations (**Fig. 5C**)
consistent with its ability to bend membranes. The addition of both AP180 and
clathrin leads to the formation of discrete cages (**Fig. 5D**), confirming previous
literature that AP180 is both necessary and sufficient for CCP and clathrin lattice
formation [30]. Further, addition of α-synuclein leads to significantly larger clathrin
assemblies (**Fig. 5D, E**) (Clathrin+AP180 = 8607 ± 413 nm^2, Clathrin+AP180+α-
synuclein= 15,569 ± 2131 nm^2; p<0.01). We were unable to determine the non-
hexagon/hexagon ratio for these images, due to the irregularity of the lattices as
well as the high contrast needed to view these structures. However, we find that
addition of clathrin, AP180 and α-synuclein leads to the formation of juxtaposed
CCPs, congruent with membrane curvature generation.
314

Discussion

316   α-Synuclein has been implicated in several steps of the synaptic cycle. Here
317 we expound on efforts to elucidate precise mechanisms by which synucleins
318 regulate SVE. Our results strongly suggest that the observed slower endocytosis
319 of SVs in αβγ-synuclein KO neurons is due to impairment of clathrin assembly and
320 curvature generation on synaptic membranes.
321

Synucleins are accessory endocytic proteins

Our findings on α-synuclein add to a growing body of work identifying
synucleins as endocytic accessory proteins. First, the biophysical properties of
synucleins are similar to those of SVE proteins. Synucleins bind acidic lipids,
especially PI(4,5)P$_2$, a feature of many endocytic proteins. Synucleins are effective
membrane curvature sensors and generators. The ability to bend membranes is
an important property of many early SVE proteins. Second, α-synuclein is recruited
to the synaptic membrane, the locus of SVE. As shown in **Fig. 1**, α-synuclein is
dynamic and relocalizes to the synaptic membrane upon stimulation, allowing it to
participate in post-fusion events such as SVE. Other SVE proteins such as AP2,
endophilins, and dynamin behave in a similar manner [44-46]. Moreover, in
dynamin1,3 double KO neurons where SVE proteins are trapped in CCPs that do
not resolve, α-synuclein is clustered with other endocytic proteins (**Fig. S2**). α-
Synuclein's behavior in dynamin1,3 double KO differs from those of other
peripheral SV proteins like synapsin and rab3a (**Fig. S2**). Third, the known proteins
that interact with α-synuclein also support a role in SVE. α-Synuclein can
multimerize on membranes, forming patches (**Fig. 2C**). Further, unbiased
proximity labelling [25] and crosslinking suggest that clathrin (**Fig. S3**) and other
SVE proteins are *in situ* binding partners on the membrane [25]. These interactions

are of low affinity and multivalent, similar to other SVE protein interactions. Finally,
the absence of synucleins leads to slowed SVE and enhanced frequency
depression due to decreased SV pool recovery [20]. In this study, we present a
preponderance of evidence that α-synuclein functions in clathrin assembly.
345
One striking finding in our study is that α-synuclein localizes to the center of clathrin
patches (**Fig. 2**). A systematic survey of the sub-cellular localization of endocytic
proteins in clathrin assembled structures has been published on HeLa cell sheets.
[47]. This work was performed using correlative super resolution microscopy
combined with electron microscopy and beautifully showed that most endocytic
proteins are found at the edge of clathrin assemblies (both flat lattices and CCPs
were evaluated) on the membrane. The only two proteins always found in the
middle of the clathrin structures were CALM and VAMP, the non-neuronal
homologs of AP180 and synaptobrevin2 [47]. Together with our finding, the results
of these studies suggest that α-synuclein is localized with AP180 and
synaptobrevin2 at the center of the patch. It is tempting to speculate that this
serves to achieve the high number of synaptobrevin2 molecules present on the SV
(~31/SV). In support for this idea, SVs purified from αβγ-synuclein KO brains show
lower levels of synaptobrevin2 compared to WT brains [20]. Similarly, in AP180
KO mice, activity-dependent defects in synaptobrevin2 recycling resulting in
reduced synaptobrevin2 levels on SVs have been observed [29].
362
What does α-synuclein do in CCP formation? Since clathrin is unable to directly
bind membrane, its recruitment requires adaptor proteins that bind both clathrin
and the membrane. Besides adaptors, accessory and regulatory endocytic
proteins are required for effective clathrin recruitment. α-Synuclein falls into this
category of proteins. Via its multi-valent interactions with PI(4,5)P2, [15, 16, 20,
26] and its ability to self-assemble on the membrane (**Fig. 2C**), and bend it (**Fig.
5C**), α-synuclein can regulate the binding of AP180 to the membrane (**Fig. 3A**, **C**),
thereby promoting clathrin patch initiation. This is supported by our previous finding
that α-synuclein and AP180 are part of a transient protein complex that assembles
post-stimulation [20]. Our present direct observation of α-synuclein within larger
clathrin patches on cell membranes (**Fig. 2B**), and that addition of α-synuclein to
purified AP180 and clathrin in the lipid monolayer assay leads to bigger clathrin
patches (**Fig. 5D-E**) are also in line with this premise. Interpreting the results from
the αβγ-synuclein KO experiments is complicated by the compensatory changes
in the cytosolic levels of clathrin and AP180 (**Fig. 3A**, **C**) and phosphorylation
changes in endocytic proteins [21]. Yet, the time course of clathrin lattice formation
is initially slower mirroring the slow kinetics of SVE (**Fig. 4B-C**) [20]. The larger
patches seen with the αβγ-synuclein KO cytosol at later time points (**Fig. 4C**) likely
occurs due to the upregulated levels of clathrin in the αβγ-synuclein KO cytosol
(**Fig. 3A**, **C**). We speculate that the larger clathrin patches take longer to fully curve
also contributing to the decreased rate of SV endocytosis [20]. Congruently, we do
find CCPs with fewer pentagons and heptagons in the αβγ-synuclein KO condition
in the lipid monolayer assays (**Fig. 4E**). In the rescue condition, the clathrin patch
size is restored (**Fig. 4C**) despite more clathrin in the cytosol, because the resulting

patches are more curved (**Fig. 4E-F**). Thus, α-synuclein appears to coordinate
clathrin assembly with curvature generation.
389
α-Synuclein, SVE and PD
Significantly, both transgenic and acute overexpression models of α-synuclein that
replicate triplication of the α-synuclein PD locus (PARK4) show impairment of SVE
[48-51]. Thus, both deletion and overexpression of α-synuclein lead to a similar
endocytic phenotype. This indicates that SVE is sensitive to levels of α-synuclein.
In the overexpression models, α-synuclein is likely to impact several steps of SVE.
In addition to any possible disruption of clathrin assembly on the membrane, α-
synuclein is also likely to impact the uncaging of CCVs. Indeed, injection of α-
synuclein in the giant Lamprey synapse leads to an accumulation of both CCPs
and CCVs [48]. The uncoating of CCVs is regulated by the heat shock protein
Hsc70 and its co-chaperone auxilin. Hsc70 has been shown to bind both unfolded
monomeric α-synuclein and α-synuclein fibrils, thus excess α-synuclein is likely to
sequester available Hsc70 from the uncoating reaction [52]. Interestingly, loss-of-
function mutations in both auxilin and the PIP phosphatase synaptojanin1 that
function in CCV uncoating cause familial PD [53-56]. Thus, SVE is a key pathway
in the pathogenesis of PD.
406
407
408 **Materials and Methods**
409
Immuno-electron microscopy of mouse brain synaptosomes:
Synaptosome Isolation
Brains from Wildtype mice (n=3) were harvested and washed in homogenizing
buffer (0.32 M Sucrose, 10 mM Tris-HCl pH 7.4, mini cOmplete protease inhibitor
tablet). Cerebella were removed, hemispheres separated, and white matter
removed before adding the remaining brain to 10 mL of homogenizing buffer.
Brains were homogenized in a 55 mL Potter homogenizer at 80% power for 8
strokes. Homogenate was centrifuged at 2,330 x g for 4 min at 4°C in a JA-20
rotor. Pellet (P1) was discarded and supernatant (S1) was transferred to a 50 mL
centrifuge tube. S1 was recentrifuged at 18,850 x g for 12 min at 4°C. Supernatant
(S2) was discarded and pellet (P2) was resuspended in 6 mL of homogenizing
buffer. Gradients of Percoll solutions (from bottom to top: 2.5 mL 23%, 3 mL 10%,
and 2.5 mL 3%) were prepared in 15 mL glass tubes. 2 mL of resuspended P2
was added to the top of each gradient, and tubes were centrifuged at 18,850 x g
for 12 min at 4°C. Synaptosome fractions were collected by Pasteur pipette.
425
Stimulation
Washing buffer Krebs 1 (140 mM NaCl, 10 mM Tris-HCl pH 7.4, 5 mM KCl, 5 mM
$NaHCO_3$, 1.3 mM $MgSO_4$, 1 mM Na-phosphate buffer) was added to the collected
synaptosomes, and sample was centrifuged at 18,850 x g for 12 min. Pellet was
resuspended in 1.5 mL Krebs 1 and divided equally into three 1.5 mL
microcentrifuge tubes labeled for rest, stimulation, and recovery. The tubes were
centrifuged at 16,000 x g for 3 min. Each pellet was resuspended in 500 μL of

washing buffer Krebs 2 (10 mM glucose, 1.2 mM CaCl2 in Krebs 1) and incubated
in a 37°C water bath for 15 min. The rest condition was left in the bath while the
stimulation and recovery conditions were each removed, supplemented to 90 mM
KCl final concentration, rotated by hand, and returned to the bath for 2 min.
Stimulation and recovery tubes were then spun at 16,000 x g for 2 min. The
stimulation pellet was resuspended in 200 μL Krebs 1 and left on ice. The recovery
pellet was resuspended in 500 μL of washing buffer Krebs 3 (10 mM Glucose, 1
mM EGTA, 1 phosSTOP tablet in Krebs 1) and returned to the bath for 10 min.
The recovery and rest tubes were removed from the bath and spun at 16,000 x g
for 2 min. Pellets were resuspended in 200 μL of Krebs 1. All conditions were
transferred into separate 50 mL centrifuge tubes filled with 30 mL of hypotonic
fixative solution (3% paraformaldehyde, 0.25% glutaraldehyde in 5 mM Na-
phosphate buffer) and incubated on ice for 30 min.
446
Agarose Embedding
The samples were centrifuged in their 50 mL tubes at 18,850 x g at 4°C for 12 min.
Each pellet was resuspended in 400 μL of 120 mM Na-phosphate buffer. 180 μL
aliquots of each sample were pipetted into glass test tubes and held on ice until
mixed with agarose. Pasteur pipettes and glass slide frames (see [57]) were
warmed in a 60°C oven. 3% Agarose solution was prepared in 5 mM Na-phosphate
buffer, warmed to 95°C until agarose dissolved, and kept in a 54°C water bath.
180 μL of agarose solution was pipetted over each broken synaptosome sample
while the tube was held in the 54°C water bath and mixed by gentle pipetting with
a Pasteur pipette. The agarose-embedded samples were pipetted into the warmed
glass slide frames and solidified on ice. Once solid, the frames were disassembled,
leaving a solid agarose gel sheet on one glass slide. The gel was cut into 3 mm^3
cubes with a razor blade and washed from the slide with 120 mM Na-phosphate
buffer into petri dishes.
461
Immunolabeling and Electron Microscopy Preparation
Each condition's cubes were divided into 5 wells of glazed porcelain well plates.
Cubes were incubated for 30 min with 5% bovine serum albumin (BSA) in solution
A (0.5 M NaCl, 0.02 M Na-phosphate buffer) at room temperature. BSA solution
was removed and each well was incubated overnight with 200 μL of primary
antibodies in solution A at 4°C. Cubes were washed with 5 changes of solution A
over the course of 1.5 hours. Solution A was removed from the wells and cubes
were incubated with secondary antibodies in solution A and 5% BSA for 6 hours
at room temperature. Cubes were washed with 5 changes of solution A and
incubated at 4°C overnight.
472
473   Agarose samples were prepared for resin embedding, sectioned, and stained with
474   2% uranyl acetate and 1% lead citrate before imaging on a FEI Tecnai Biotwin
475   scope at 48k, 80kV.
476
Reconstitution of endocytosis in PTK2 Cells:

PTK2 cells were transfected to express palmitoylated GFP and cultured for 48
hours in a Mattek dish to confluence. Membrane sheets were prepared by
sonication, using 20% output power, 1 brief pulse on the center of the well. The
sheets were washed gently with cytosolic buffer and used within 20 min. The
membranes were incubated for 0, 5, 15, 30 min with a mix of 2 mg/ml cytosol, 1.5
mM ATP, ATP reconstitution system (16.7mM phosphocreatine,16.7U/ml creatine
phosphokinase), and 150 μM GTPγS. The reaction was stopped by gently wash 2
times with cytosolic buffer and immediately fixed with 4% PFA for 15min in PBS.
Membrane-bound proteins were detected using appropriate primary and
secondary antibodies. The membrane sheets were covered with cytosolic buffer.
Z stack images were captured using spinning disc confocal microscopy and
analyses were performed in ImageJ using Fiji. Z slices were summed to create a
single image. For each circular structure observed on membranes of interest,
linear ROIs were created and analyzed for size and fluorescence intensity.
492
Isolation of synaptic membranes:
All steps were executed at 4°C. Two wildtype mouse brains were washed in Buffer
A (1 mM MgCl$_2$, 1 mM CaCl$_2$, 1 mM NaHCO$_3$, 0.32 M sucrose) for 30 seconds and
transferred to a 55 mL potter homogenizer with 6 mL of Buffer A. The samples
were homogenized at 5,000 rpm for 10 strokes. The homogenate was centrifuged
at 1,400 x g for 10 min. Supernatant was retained. The pellet was resuspended in
6 mL of Buffer A and centrifuged at 700 x g for 10 min. Both supernatants were
combined and centrifuged at 17,500 x g for 15 min. The supernatant was
discarded, and the pellet was resuspended in 3 mL Buffer A. The resuspended
pellet (1.5 mL) was added to the top of freshly prepared 0.65 M, 0.85 M, 1.00 M,
1.20 M sucrose gradients (1.5 mL each concentration). The gradients were
centrifuged at 100,000 x g for 2 hours. Synaptosomes were collected from the
1/1.2 M interface and suspended in an excess of Buffer A. The synaptosomes were
centrifuged at 100,000 x g for 20 min. The pellet was resuspended in 4 mL ice-
cold deionized water, and HEPES pH 7.4 was added to a final concentration of
7.5mM. The suspension was incubated on ice for 30 min and centrifuged at
100,000 x g for 20 min. The pellet was resuspended in 4 mL of 0.1 M Na$_2$CO$_3$ to
strip peripheral proteins, incubated for 15 min at 37°C, and centrifuged at 100,000
x g for 20 min. Pellet was resuspended in 2 mL cytosolic buffer, centrifuged again
at 100,000 x g for 20 min, and resuspended in 2 mL cytosolic buffer. Proteins were
quantified using BCA bioassay. Mini cOmplete protease inhibitor cocktail was
added, and 100 μl aliquots of purified membrane resuspension were flash frozen
in liquid nitrogen and stored at -80°C.
516
Cytosol purification:
Mouse brains were removed and washed in washing buffer (23 mM Tris-HCl, pH
7.4, 320 mM sucrose) for 30 seconds. Two brains were homogenized at 2,500 rpm
in a 5 mL potter homogenizer with 2 mL of homogenization buffer (25mM Tris-HCl,
pH 8.0, 500 mM KCl, 250 mM sucrose, 2 mM EGTA, and 1 mM DTT), using 10
strokes at 5,000 rpm. The homogenate was transferred to a 3.5 mL ultracentrifuge
tube and centrifuged at 160,000 x g for 2 hours at 4°C. A PD-10 column was

equilibrated with 25 mL cytosolic buffer, the supernatant was added to the PD-10
column, and then, eluted with 3.5 mL of cytosolic buffer (25 mM Hepes-NaOH, pH
7.4, 120 mM potassium glutamate, 2.5 mM magnesium acetate, 20 mM KCl, and
5 mM EGTA-NaOH, pH 8.0, filtered and stored at 4°C, with 1 mM DTT added
immediately before use). Protein concentration was quantified by BSA assay. Mini
cOmplete protease inhibitor cocktail was added, and 100 µl aliquots were flash
frozen in liquid nitrogen and stored at -80°C.
531
Synaptic membrane protein recruitment:
Synaptic membranes (400 µg/mL) were mixed with 0.5 mg/mL wildtype or αβγ-
synuclein KO cytosolic proteins in 500 µl cytosolic buffer (see above, with 1 mM
DTT added immediately before use) and supplemented with an ATP regenerating
system (1.5 mM ATP, 16.7 mM phosphocreatine, and 16.7 U/mL creatine
phosphokinase) and 150 µM GTPγS. A control experiment was prepared with 400
µg/mL synaptic membranes in 500µl of cytosolic buffer alone. Mixtures were
pipetted 3 times to mix and incubated at 37°C for 15 min. The samples were
immediately centrifuged at 100,000 x g for 30 min at 4°C. Pellets were thoroughly
resuspended in 500 µl of cytosolic buffer at 4°C. The resuspension was centrifuged
at 100,000 x g for 30 min at 4° resuspended in 45 µl of cytosolic buffer. For each
sample, 20 µl aliquots were mixed with 5 µl of 5X loading buffer. Wildtype and αβγ-
synuclein knockout cytosol reference samples were prepared by diluting 32 µg of
each cytosol to 20 µl with cytosolic buffer and mixing with 5 µl of 5X loading buffer.
Samples were boiled for 10 min at 100°C. Levels of various proteins were
measured for each condition by quantitative immunoblotting.
548
549 **Reconstitution of clathrin recruitment on lipid monolayers:**
550
Purification of proteins
α-Synuclein and AP180 were recombinantly expressed and purified from BL21
(DE3) *E. Coli* as previously described [24, 30, 58]. Clathrin was purified from
porcine or bovine brains by purifying CCVs and disassembling cages as described
[59]. For the described experiments, we used porcine and bovine clathrin
interchangeably with no significant difference.
557
Lipid Monolayer Assay with Purified Proteins
An 8-well Teflon block was arranged in a humid chamber and wells were filled with
40-60 µL of HKM buffer (25 mM HEPES pH 7.4, 125 mM potassium acetate, 5mM
magnesium acetate, 1mM dithiothreitol). Lipid mixture (10% cholesterol, 40% PE,
40% PC, and 10% PI(4,5)P2 to final concentration of 0.1 mg/mL in a 19:1 mixture
of chloroform to methanol) was carefully pipetted onto the buffer in each well. The
blocks were incubated in a humid chamber at room temperature for 1 hour to
evaporate methanol/chloroform. Carbon-coated copper grids were placed carbon-
side down onto each well. Proteins were introduced via side-injection ports beside
each well. Final protein concentrations per well were as follows: 2 µM AP180 and
α-synuclein, 500 nM purified bovine/porcine clathrin. Grids were incubated for 60

min in a humid chamber at room temperature, and then, removed from the block
and immediately negative stained with 1% uranyl acetate. After drying, grids were
imaged on a Phillip CM10 transmission electron microscope at 46k, 80kV). Clathrin
lattices were outlined manually and the areas were quantified in Image J. Non-
hexagonal clathrin lattices could not be quantified due to the high contrast needed
to visualize these structures in these experiments.
575
576
Lipid Monolayer Assay with Mouse Brain Cytosol
5 mg/mL wild type and $\alpha\beta\gamma$-synuclein KO mouse brain cytosol was added to wells
of the Teflon block. Lipid mixture (see above) was pipetted on top of the cytosol.
Blocks were incubated in a humid chamber for 1 hour to evaporate
methanol/chloroform. Carbon-coated copper grids were placed on top of the wells,
and an ATP regenerating system (see above) and 150 μM GTPγS were introduced
through a side injection port. Grids were incubated in a humid chamber at 37°C for
0, 5, 15, or 30 min and then removed from the block and immediately negative
stained as described above. Grids were imaged on a FEI Tecnai Biotwin at scope
42k, 80kV. Clathrin lattices were outlined manually, the area and number of
pentagons, heptagons and hexagons were quantified in Image J.
588
Crosslinking experiments:
Synaptosomes were isolated from wildtype and $\alpha\beta\gamma$-synuclein KO mouse brains
as described. Synaptosomes were incubated with SDAD crosslinker (1 mM;
Thermo) or DMSO as control and incubated on ice for 45 minutes. The reaction
was stopped with 100 μl 1M Tris, pH 8.0 on ice for 15 minutes. The excess
crosslinker was removed by centrifugation. The synaptosomes were then UV
crosslinked for 1 minute with a Strategene UV light (254 nm). Crosslinked synaptic
membranes were solubilized with 1% Triton and used as starting material for α-
synuclein immunoprecipitation experiments. The immunoprecipitants were run on
SDS-PAGE, silver stained with a mass spectrometry compatible silver stain
(Pierce), relevant bands were excised and identified by LC-MS.
600
601 **Statistics:**
602 All experimental analysis was done blinded to condition. Statistical analysis was
603 performed by GraphPad. Data is presented as average ± SEM.
604

Acknowledgments: This work was supported by NIH (R01 NS064963, R01
NS110354, R01 NS083846), Nina Compagnon Hirshfield Parkinson's Disease
Research Fund and DOD (W81XWH-17-1-0564). The proteomic experiments
were supported by the NIDA Neuroproteomic Center (NIH DA018343-11A1). We
would like to thank Dr. Pietro De Camilli and his lab members for Dynamin1,3
double KO pups as well as help in performing the PTK2 cell assays. We thank
Aurelie Nardin for setting up the PTK2 assay, Christopher Westphal and Becket
Greten-Harrison for help with cross-linking experiments. We thank John. E. Lee
for editing the manuscript.

Author contributions: KJV, PC and SSC designed research/experiments; KJV,
PC, EG and SSC performed experiments; KJV, PC, EG and SSC analyzed data;
All authors wrote and edited the paper.
618
619 **Figure Legends**
620
Figure 1. α-Synuclein localization is dynamically regulated by neuronal
activity. (**A**) Electron micrographs of permeabilized synaptosomes at rest,
stimulated with 90 mM KCl, and upon recovery. Immuno-gold labeled α-synuclein
is localized to SVs at rest (arrowhead). During stimulation, α-synuclein
disassociates from SVs and predominantly localizes on the synaptic membrane.
During recovery, α-synuclein returns to the SV membrane. (**B**) Electron
micrographs of permeabilized synaptosomes in rest, stimulation, and recovery
conditions immuno-labeled for synaptobrevin2. Gold-labeled synaptobrevin 2
remains associated with the vesicle in all three conditions. (**C-D**) Quantification of
gold label localization for α-synuclein and synaptobrevin2. Bars represent percent
gold label from n=3 independent experiments for control conditions and n=2 for
remaining conditions. 25-40 micrographs per condition were analyzed. ANOVA
was used to determine significance. * $p < 0.05$; ** $p < 0.01$ and *** $p < 0.0001$. Scale
bar=100 nm.
635
Figure 2. α-Synuclein and clathrin co-localize in patches on the plasma
membrane. (**A**) Membrane sheets from PTK2 cells immunostained for α-synuclein
(red) and clathrin heavy chain (green). α-Synuclein and clathrin are present in
patches on the membrane, with some patches containing both proteins. Scale bar=
1 μm. (**B**) Zoomed-in view of a clathrin+α-synuclein patch. Note that α-synuclein
localizes to the center of the clathrin patch. (**C**) α-Synuclein only patches, which
are on average smaller than clathrin patches. (**D**) Plot of fluorescence of clathrin
versus α-synuclein colored by size of the clathrin patch. The dashed lines (x=0.5
and y=1) indicate the cut-offs used to demarcate patches into clathrin only (x<0.5,
y>1), α-synuclein only (x>0.5, y<1), and clathrin+α-synuclein (x>0.5, y>1) puncta.
Plots show that colocalization of clathrin and α-synuclein is predominant in larger
size patches. (**E**) Intensity of clathrin and α-synuclein in different patch sizes. (**F**)
Average size of clathrin only, α-synuclein only and clathrin+α-synuclein patches.
(**G**) Changes in the three population of patches over the time when incubated with
wildtype purified cytosol. N=3 independent experiments. * $p < 0.05$; ** $p < 0.01$ and
*** $p < 0.0001$. Scale bar=1 μm.
652
Figure 3. Recruitment of endocytic proteins to synaptic membranes from
wildtype and αβγ-synuclein KO brain cytosol. (**A-B**) Western blot of endocytic
proteins recruited from wildtype (WT) and αβγ-synuclein KO (TKO) cytosol to
synaptic membranes (Recruitment). Membrane fraction without addition of cytosol
is used as a control. N-cadherin was used as a membrane loading control.
Abbreviations: Memb, synaptic membrane fraction; CLC, clathrin light chain; CHC,
clathrin heavy chain; AAK1, AP2 Associated Kinase-1. (**C**) Quantification of
membrane recruitment of endocytic proteins. Western blot quantifications (n=3

independent experiments) after normalizing for added membranes (N-Cadherin
signal). * p<0.05, ** p<0.01, Student's t-test.
663
Figure 4. Lipid monolayer assay to visualize clathrin recruitment from brain
cytosol. (**A**) Schematic of Teflon block and experimental setup used in monolayer
reconstitutions. Wells (W) and Side Injection Ports (SIP) labeled. (**B**) Electron
micrographs showing the clathrin assembly into flat lattices and CCPs on
membrane monolayers over time (0,15, 23, and 30 min). These monolayers were
incubated with cytosol from wild type (WT), αβγ-synuclein KO (TKO), and mouse
α-synuclein added to TKO cytosol (rescue). Clathrin patches are highlighted with
red outlines. Arrow indicates CCP. N=3 independent experiments with 10 images
per experiment. Scale bar=100 nm. (**C**) Comparison of clathrin patch areas in the
three conditions mentioned above. (**D**) Examples of pentagon, hexagon and
heptagon clathrin lattices. (**E**) Ratio of non-hexagons (pentagons and heptagons)
to hexagon in clathrin lattices for each of the three conditions, t=23 min. (**F**)
Histogram of non-hexagon to hexagon ratio for the three conditions. The sum of
all weights was used to normalize the data. For all graphs: * p<0.05; ** p<0.01 and
*** p<0.0001. Welsh's t-test.
679
Figure 5. Reconstitution of clathrin assembly using recombinant proteins.
(**A**) Coomassie stain showing proteins used in reconstitution conditions. (**B**)
Electron micrographs of negative controls. Left: Lipid monolayer with no added
protein. Center: recombinant AP180 added to lipid monolayer. Right: Brain purified
clathrin added to lipid monolayer. Lower: All proteins without a lipid monolayer
Under all four conditions no clathrin cages form. (**C-E**) Electron micrographs of
constituent proteins added to lipid monolayers. (**C**) Left: Recombinant wildtype
mouse α-synuclein. Only membrane ruffles are observed. Center Recombinant
mouse AP180 and α-synuclein. Left: α-Synuclein and isolated clathrin added to
lipid monolayer. (**D**) Electron micrographs of clathrin and AP180 added to lipid
monolayer. Characteristic fullerene-shaped clathrin cages form. Inset: Zoomed
image outlined in green (**E**) Electron micrographs of α-synuclein, clathrin, and
AP180 added to lipid monolayer. Inset: Zoomed image outlined in blue. Scale bar
= 200 nm. (**F**) Quantification of area of clathrin patches obtained in conditions **D**
and **E**. Addition of α-synuclein increases the area of clathrin patches. N=3
independent experiments with at least 10 micrographs per experiment. * p<0.05;
** p<0.01 and *** p<0.0001. Scale bar=100 nm.
697
698

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

SUPPLEMENTARY FIGURES

Title: α-Synuclein facilitates clathrin assembly in synaptic vesicle endocytosis

Short title: α-Synuclein regulates clathrin assembly

Karina J Vargas[1]*#, P. L. Colosi[1,2]*#, Eric Girardi[1]*, Sreeganga S Chandra[1,3]

[1] Departments of Neurology; Neuroscience, Yale University, New Haven, CT 06536
[2] PREP Program, Yale University, New Haven, CT 06511
[3] To whom correspondence should be addressed to:
sreeganga.chandra@yale.edu ; 203-785-6172

*Authors contributed equally to this study
Present Address:
KJV, Marine Biological Laboratory, MA
PLC, University of Pennsylvania, PA

Supplementary Figure 1. α-Synuclein is localized to endocytic clusters in Dynamin1,3 double KO neurons. Control and Dynamin1,3 double KO neurons were immunostained with antibodies to an endocytic protein (clathrin-CLC or Endophilin A1-Endo1) and α-synuclein. Clustering of endocytic proteins is observed in the Dynamin1,3 double KO but not in the Control. α-Synuclein colocalizes with other trapped endocytic proteins. Scale bar =10 um.

Supplementary Figure 2. Temporal changes of size and colocalization of α-synuclein and clathrin patches. PTK2 cells were sheared and processed to detect α-synuclein and clathrin on the resulting membrane sheets. **(A)** Size comparisons of clathrin only, α-synuclein only, and clathrin+α-synuclein patches at the denoted time points. Welsh's t-test was used to determine significance. * $p<0.05$; ** $p<0.01$ and *** $p<0.0001$. **(B)** Fraction of the three type of patches as a function of time.

Supplementary Figure 3. Immunoprecipitation of crosslinked α-synuclein from wild type and αβγ-synuclein KO synaptosomes. Wild type (WT) and αβγ-synuclein KO (TKO) synaptosomes were crosslinked, solubilized and immunoprecipitated with an antibody against α-synuclein. The immunoprecipitants were load in a polyacrylamide gel and stained with silver. Bands marked with boxes were cut and processed for mass spectroscopy.

References

1. Spillantini, M.G., et al., *Alpha-synuclein in Lewy bodies.* Nature, 1997. **388**(6645): p. 839-40.

2. Ferese, R., et al., *Four Copies of SNCA Responsible for Autosomal Dominant Parkinson's Disease in Two Italian Siblings.* Parkinsons Dis, 2015. **2015**: p. 546462.

3. Ibanez, P., et al., *Causal relation between alpha-synuclein gene duplication and familial Parkinson's disease.* Lancet, 2004. **364**(9440): p. 1169-71.

4. Kruger, R., et al., *Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease.* Nat Genet, 1998. **18**(2): p. 106-8.

5. Polymeropoulos, M.H., et al., *Mutation in the alpha-synuclein gene identified in families with Parkinson's disease.* Science, 1997. **276**(5321): p. 2045-7.

6. Singleton, A.B., et al., *alpha-Synuclein locus triplication causes Parkinson's disease.* Science, 2003. **302**(5646): p. 841.

7. Zarranz, J.J., et al., *The new mutation, E46K, of alpha-synuclein causes Parkinson and Lewy body dementia.* Ann Neurol, 2004. **55**(2): p. 164-73.

8. Satake, W., et al., *Genome-wide association study identifies common variants at four loci as genetic risk factors for Parkinson's disease.* Nat Genet, 2009. **41**(12): p. 1303-7.

9. Simon-Sanchez, J., et al., *Genome-wide association study reveals genetic risk underlying Parkinson's disease.* Nat Genet, 2009. **41**(12): p. 1308-12.

10. Maroteaux, L., J.T. Campanelli, and R.H. Scheller, *Synuclein: a neuron-specific protein localized to the nucleus and presynaptic nerve terminal.* J Neurosci, 1988. **8**(8): p. 2804-15.

11. Lokappa, S.B. and T.S. Ulmer, *Alpha-synuclein populates both elongated and broken helix states on small unilamellar vesicles.* J Biol Chem, 2011. **286**(24): p. 21450-7.

12. Ferreon, A.C., et al., *Interplay of alpha-synuclein binding and conformational switching probed by single-molecule fluorescence.* Proc Natl Acad Sci U S A, 2009. **106**(14): p. 5645-50.

13. Middleton, E.R. and E. Rhoades, *Effects of curvature and composition on alpha-synuclein binding to lipid vesicles.* Biophys J, 2010. **99**(7): p. 2279-88.

14. Burre, J., M. Sharma, and T.C. Sudhof, *alpha-Synuclein assembles into higher-order multimers upon membrane binding to promote SNARE complex formation.* Proc Natl Acad Sci U S A, 2014. **111**(40): p. E4274-83.

15. Burre, J., et al., *Alpha-synuclein promotes SNARE-complex assembly in vivo and in vitro.* Science, 2010. **329**(5999): p. 1663-7.

16. Sun, J., et al., *Functional cooperation of alpha-synuclein and VAMP2 in synaptic vesicle recycling.* Proc Natl Acad Sci U S A, 2019. **116**(23): p. 11113-11115.

17. Diao, J., et al., *Native alpha-synuclein induces clustering of synaptic-vesicle mimics via binding to phospholipids and synaptobrevin-2/VAMP2.* Elife, 2013. **2**: p. e00592.

18. Greten-Harrison, B., et al., *alphabetagamma-Synuclein triple knockout mice reveal age-dependent neuronal dysfunction.* Proc Natl Acad Sci U S A, 2010. **107**(45): p. 19573-8.

19.　Anwar, S., et al., *Functional alterations to the nigrostriatal system in mice*
lacking all three members of the synuclein family. J Neurosci, 2011. **31**(20): p.
7264-74.
20.　Vargas, K.J., et al., *Synucleins regulate the kinetics of synaptic vesicle*
endocytosis. J Neurosci, 2014. **34**(28): p. 9364-76.
21.　Vargas, K.J., et al., *Synucleins Have Multiple Effects on Presynaptic Architecture.*
Cell Rep, 2017. **18**(1): p. 161-173.
22.　Logan, T., et al., *alpha-Synuclein promotes dilation of the exocytotic fusion*
pore. Nat Neurosci, 2017. **20**(5): p. 681-689.
23.　Atias, M., et al., *Synapsins regulate alpha-synuclein functions.* Proc Natl Acad
Sci U S A, 2019. **116**(23): p. 11116-11118.
24.　Westphal, C.H. and S.S. Chandra, *Monomeric synucleins generate membrane*
curvature. J Biol Chem, 2013. **288**(3): p. 1829-40.
25.　Chung, C.Y., et al., *In Situ Peroxidase Labeling and Mass-Spectrometry Connects*
Alpha-Synuclein Directly to Endocytic Trafficking and mRNA Metabolism in
Neurons. Cell Syst, 2017. **4**(2): p. 242-250 e4.
26.　Burre, J., M. Sharma, and T.C. Sudhof, *Systematic mutagenesis of alpha-*
synuclein reveals distinct sequence requirements for physiological and
pathological activities. J Neurosci, 2012. **32**(43): p. 15227-42.
27.　Koo, S.J., D. Puchkov, and V. Haucke, *AP180 and CALM: Dedicated endocytic*
adaptors for the retrieval of synaptobrevin 2 at synapses. Cell Logist, 2011.
1(4): p. 168-172.
28.　Moshkanbaryans, L., L.S. Chan, and M.E. Graham, *The Biochemical Properties*
and Functions of CALM and AP180 in Clathrin Mediated Endocytosis.
Membranes (Basel), 2014. **4**(3): p. 388-413.
29.　Koo, S.J., et al., *Vesicular Synaptobrevin/VAMP2 Levels Guarded by AP180*
Control Efficient Neurotransmission. Neuron, 2015. **88**(2): p. 330-44.
30.　Ford, M.G., et al., *Simultaneous binding of PtdIns(4,5)P2 and clathrin by AP180*
in the nucleation of clathrin lattices on membranes. Science, 2001. **291**(5506):
p. 1051-5.
31.　Miller, S.E., et al., *CALM regulates clathrin-coated vesicle size and maturation*
by directly sensing and driving membrane curvature. Dev Cell, 2015. **33**(2): p.
163-75.
32.　Qualmann, B., D. Koch, and M.M. Kessels, *Let's go bananas: revisiting the*
endocytic BAR code. EMBO J, 2011. **30**(17): p. 3501-15.
33.　Ferguson, S.M., et al., *A selective activity-dependent requirement for dynamin 1*
in synaptic vesicle endocytosis. Science, 2007. **316**(5824): p. 570-4.
34.　Clayton, D.F. and J.M. George, *Synucleins in synaptic plasticity and*
neurodegenerative disorders. J Neurosci Res, 1999. **58**(1): p. 120-9.
35.　Antonny, B., *Mechanisms of membrane curvature sensing.* Annu Rev Biochem,
2011. **80**: p. 101-23.
36.　Rhoades, E., et al., *Quantification of alpha-synuclein binding to lipid vesicles*
using fluorescence correlation spectroscopy. Biophys J, 2006. **90**(12): p. 4692-
700.
37.　Fortin, D.L., et al., *Neural activity controls the synaptic accumulation of alpha-*
synuclein. J Neurosci, 2005. **25**(47): p. 10913-21.

38. Wang, L., et al., *alpha-synuclein multimers cluster synaptic vesicles and*
attenuate recycling. Curr Biol, 2014. **24**(19): p. 2319-26.
39. Raimondi, A., et al., *Overlapping role of dynamin isoforms in synaptic vesicle*
endocytosis. Neuron, 2011. **70**(6): p. 1100-14.
40. Wu, M., et al., *Coupling between clathrin-dependent endocytic budding and F-*
BAR-dependent tubulation in a cell-free system. Nat Cell Biol, 2010. **12**(9): p.
902-8.
41. Hosaka, M., R.E. Hammer, and T.C. Sudhof, *A phospho-switch controls the*
dynamic association of synapsins with synaptic vesicles. Neuron, 1999. **24**(2):
p. 377-87.
42. Chandra, S., et al., *Alpha-synuclein cooperates with CSPalpha in preventing*
neurodegeneration. Cell, 2005. **123**(3): p. 383-96.
43. Bucher, D., et al., *Clathrin-adaptor ratio and membrane tension regulate the*
flat-to-curved transition of the clathrin coat during endocytosis. Nat Commun,
2018. **9**(1): p. 1109.
44. Evergren, E., et al., *A pre-embedding immunogold approach for detection of*
synaptic endocytic proteins in situ. J Neurosci Methods, 2004. **135**(1-2): p.
169-74.
45. Pechstein, A., et al., *Regulation of synaptic vesicle recycling by complex*
formation between intersectin 1 and the clathrin adaptor complex AP2. Proc
Natl Acad Sci U S A, 2010. **107**(9): p. 4206-11.
46. Ringstad, N., et al., *Endophilin/SH3p4 is required for the transition from early*
to late stages in clathrin-mediated synaptic vesicle endocytosis. Neuron, 1999.
24(1): p. 143-54.
47. Sochacki, K.A., et al., *Endocytic proteins are partitioned at the edge of the*
clathrin lattice in mammalian cells. Nature Cell Biology, 2017. **19**(4): p. 352-+.
48. Busch, D.J., et al., *Acute increase of alpha-synuclein inhibits synaptic vesicle*
recycling evoked during intense stimulation. Mol Biol Cell, 2014. **25**(24): p.
3926-41.
49. Xu, J., et al., *alpha-Synuclein Mutation Inhibits Endocytosis at Mammalian*
Central Nerve Terminals. J Neurosci, 2016. **36**(16): p. 4408-14.
50. Nemani, V.M., et al., *Increased expression of alpha-synuclein reduces*
neurotransmitter release by inhibiting synaptic vesicle reclustering after
endocytosis. Neuron, 2010. **65**(1): p. 66-79.
51. Eguchi, K., et al., *Wild-Type Monomeric alpha-Synuclein Can Impair Vesicle*
Endocytosis and Synaptic Fidelity via Tubulin Polymerization at the Calyx of
Held. J Neurosci, 2017. **37**(25): p. 6043-6052.
52. Susan M. L. Banks, A.T.M., Molly McQuillan, David J. Busch, Ana Sofia
Ibarraran-Viniegra, Subhojit Roy, Rui Sousa, Eileen M. Lafer, Jennifer R.
Morgan, *Hsc70 Ameliorates the Vesicle Recycling Defects Caused by Excess α-*
Synuclein at Synapses. Neuroscience Online, 2020.
53. Cao, M., et al., *Parkinson Sac Domain Mutation in Synaptojanin 1 Impairs*
Clathrin Uncoating at Synapses and Triggers Dystrophic Changes in
Dopaminergic Axons. Neuron, 2017. **93**(4): p. 882-896 e5.
54. Cremona, O., et al., *Essential role of phosphoinositide metabolism in synaptic*
vesicle recycling. Cell, 1999. **99**(2): p. 179-88.

55.   Yim, Y.I., et al., *Endocytosis and clathrin-uncoating defects at synapses of*
auxilin knockout mice. Proc Natl Acad Sci U S A, 2010. **107**(9): p. 4412-7.
56.   Olgiati, S., et al., *DNAJC6 Mutations Associated With Early-Onset Parkinson's*
Disease. Ann Neurol, 2016. **79**(2): p. 244-56.
57.   De Camilli, P., et al., *Synapsin I (Protein I), a nerve terminal-specific*
phosphoprotein. II. Its specific association with synaptic vesicles demonstrated
by immunocytochemistry in agarose-embedded synaptosomes. J Cell Biol, 1983.
96(5): p. 1355-73.
58.   Taguchi, Y.V., et al., *Glucosylsphingosine Promotes alpha-Synuclein Pathology*
in Mutant GBA-Associated Parkinson's Disease. J Neurosci, 2017. **37**(40): p.
9617-9631.
59.   Chen, Y., et al., *Dynamic instability of clathrin assembly provides proofreading*
control for endocytosis. J Cell Biol, 2019. **218**(10): p. 3200-3211.
850