1   Article type: Research article

2   Germination ecology of wild and domesticated *Ensete*
3   *ventricosum*: Evidence for maintenance of sexual
4   reproductive capacity in a vegetatively propagated
5   perennial crop

Solomon Tamrat[1*], James S. Borrell[2*†], Eleni Shiferaw[3], Simon Kallow[4,5], Rachael M. Davies[4], John

B. Dickie[4], Gizachew W. Nuraga[1], Oliver White[2], Feleke Woldeyes[3], Sebsebe Demissew[1], Paul

Wilkin[2]

*Denotes joint first authors

[†]Corresponding author

[1]Department of Plant Biology and Biodiversity Management, Addis Ababa University, Addis Ababa,

P.O. Box 3293,5 Ethiopia.

[2]Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK.

[3]Ethiopian Biodiversity Institute, Addis Ababa, P.O.Box: 30726, Ethiopia.

[4]Royal Botanic Gardens Kew, Millennium Seed Bank, Wakehurst, Ardingly, Sussex, RH17 6TN, UK.

[5]Katholieke Universiteit Leuven, Department of Biosystems, Willem de Croylaan 42, 3001, Leuven,

Belgium.

Running head: Germination ecology of wild and domestic enset

ABSTRACT

PREMISE: Loss of sexual reproductive capacity has been proposed as a syndrome of domestication in vegetatively propagated crops, but there are relatively few examples from agricultural systems. Compared to sexually propagated crops banked as seeds, vegetative crop diversity is typically conserved in living collections that are more costly and insecure. This vulnerability may disproportionately impact tropical crop diversity where vegetative reproduction predominates. In this study we assess sexual reproductive capacity in wild (sexual) and domesticated (vegetative) populations of enset (*Ensete ventricosum* (Welw.) Cheesman)*, a giant tropical banana relative and Ethiopian food security crop.

METHODS: We harvested seeds from 20 wild and domesticated enset to survey variation in seed weight, viability, external and internal morphology. We germinated seeds across a range of constant and alternating temperature regimes to characterise optimum germination requirements, and evaluate differences in performance.

KEY RESULTS: We found no significant difference between wild and domesticated enset seed viability, days to germination and internal morphological traits. However, we report a significant and unexpected shift in seed weight and germination behaviour, with seed from domesticated plants responding to cooler temperatures with greater diurnal range. Shifts in germination behaviour appear concordant with a climatic envelope shift in the domesticated distribution.

CONCLUSIONS: Our findings suggest sexual reproductive capacity has been maintained despite long-term near-exclusive vegetative propagation in domesticated enset. Furthermore, certain traits such as germination behaviour, may be under continued selection through rare sexually reproductive events. Improved understanding of sexual propagation in vegetative crops may have applications in germplasm conservation and plant breeding.

Keywords: clonal reproduction; Crop Wild Relative; domestication; Enset; Ethiopia; germination biology; Perennial crops; Seed banking; vegetative propagation

INTRODUCTION

The erosion of plant genetic resources poses a substantial threat to food security and the diverse benefits we derive from useful plants (Borrell et al., 2019; Powell et al., 2018). To address this challenge a significant effort has been made to conserve >2 million unique accessions, representing >16,500 plant species in 1750 genebanks worldwide (Commission for genetic resources on food and agriculture, 2010; Fu, 2017). However an emphasis on conventional seed crops may overlook the majority of perennial fruit crops for which vegetative propagation is the predominant means of replication and seed generation may be rare or absent (Miller and Gross, 2011; Castañeda-álvarez et al., 2016; Migicovsky and Myles, 2017a). Vegetatively propagated crops are especially important in the tropics (Denham et al., 2020), where *ex situ* or *in vitro* germplasm collections are currently the only viable approach for conserving genetic resources (Thormann and Dulloo, 2006). Maintaining such collections is often logistically challenging and prohibitively expensive, particularly for developing countries (Dulloo et al., 2013). Nevertheless, the majority of vegetatively propagated crops have sexually reproducing wild progenitors (Miller and Gross, 2011). Improving our understanding of reproductive biology in vegetative propagated perennial crops has potential to address these challenges by enabling a broader range of useful plants to be banked as seed and used in breeding programmes, enhancing conservation and use of genetic diversity (even if this does not conserve specific vegetative genotypes) (McKey et al., 2010; Borrell, Dodsworth, et al., 2019; Pironon et al., 2019; Denham et al., 2020).

Clonally propagated food crops encompass at least 34 families and a wide variety of morphological diversity (McKey et al., 2010). This diversity has hindered attempts to define a domestication syndrome in clonally reproduced crops, but a number of commonalities have been proposed (McKey et al., 2010; Miller and Gross, 2011; Denham et al., 2020). Foremost is the suggestion that prolonged vegetative reproduction can lead to the loss of sexual reproductive capacity (Eckert, 2002; McKey et al., 2010; Barrett, 2015; Denham et al., 2020). For example, cultivated bananas (*Musa*) differ from their wild relatives by being seedless and parthenocarpic (Heslop-Harrison and Schwarzacher, 2007)

whilst in *Citrus*, fertile varieties co-exist with and asexually propagated cultivars that reproduce via

nucellar embryony (Wu et al., 2018). Domestication of Pineapple (*Annas*) was also associated with a

combination of sexual and asexual selection, together with reduced seed production through lower

fertility and self-incompatibility (Chen et al., 2019). Loss of sexual reproductive capacity, if

widespread across domesticated perennial crops, could significantly hinder future perennial crop

breeding programmes and integration of diversity from crop wild relatives (Dempewolf et al., 2014;

Migicovsky and Myles, 2017a). Importantly, sexual reproductive capacity does not have to be

completely lost – reduced fertility, viability, altered germination behaviour or other changes as a

result of deleterious mutations and genetic drift in reproductive traits could still significantly hinder

programmes that seek to recombine or conserve diversity as seeds (e.g. McClure, Sawler, Gardner,

Money, & Myles, 2014; Migicovsky & Myles, 2017). However, surprisingly few studies have

attempted to survey the sexual reproductive capacity of vegetatively propagated crops in

agricultural systems (Elias et al., 2001; Scarcelli et al., 2006).

Here, we investigate the impact of domestication on seed and germination biology of the perennial

food security crop enset (*Ensete ventricosum* (Welw) Cheesman) (Figure 1). Enset is a giant

monocarpic herb, in the sister genus to the more widely known bananas (*Musa),* which provides a

staple starch source for 20 million people of South and South-west Ethiopia (Janssens et al., 2016;

Borrell, Biswas, et al., 2019). Wild enset reproduces sexually, generating inedible fruits that contain

hundreds to thousands of seeds (Ssali and Sheil, 2019), but unlike bananas, enset does not produce

vegetative suckers. However, during domestication (a process which also improved palatability and

diversified uses (Tamrat et al., 2020)), sophisticated agronomic practices were developed to enable

exclusively vegetative propagation (Borrell et al., 2020) (Figure 1B, 1C). Farmers achieve this by

removing the apical meristem and meristematic tissue of a two to three year old corm, resulting in

the generation of adventitious buds (Karlsson, Dalbato, Tamado, & Mikias, 2014). Currently, up to

1500 named clonal landraces have been recorded and are maintained by millions of rural farmers

(Borrell, Biswas, et al., 2019). Domesticated enset is restricted to Ethiopia and has been increasingly

overlooked in favour of introduced species (Borrell, Biswas, et al., 2019), whilst permanent

international transfer of enset plant material out of Ethiopia is not permitted. As a result,

domesticated enset is not banked as seed in international (or national) collections (Guzzon and

Müller, 2016), and only a handful of local institutes maintain field germplasm collections

(specifically; Areka, Yerefezy, Angacha and Hawassa, situated in southern Ethiopia). With increasing

pressures due to climate change (Conway and Schipper, 2011) and emerging pests and pathogens

(Blomme et al., 2017), this represents a significant risk for the future sustainability of enset

agriculture.

Conserving domesticated enset diversity as seeds has been considered challenging for several

reasons. Firstly, enset is monocarpic and is harvested before flowering (unlike banana), to avoid

reallocation of resources from edible storage organs to the inedible inflorescence (Borrell et al.,

2020). This means that developed fruits are rarely encountered in cultivation. Secondly, in clonally

propagated crops generally, reduction of sexual fertility has been identified as a characteristic of the

domestication syndrome (Denham et al., 2020). Indeed a recent genomic analysis concluded that

accumulation of mutations in genes associated with sexual reproduction may have contributed to

enset's domestication (Tesfamicael et al., 2020). However, there has been no systematic study of

whether the majority of domesticated enset landraces retain the ability to produce viable seeds.

Thirdly, enset is thought to be an obligate out-crosser, yet the pollinator(s) of enset in Ethiopia are

unknown (though see recent reports of the Nectar bat *Megaloglossus woermanni* visiting flowers

from Ssali & Sheil (2019) in Uganda), and significant conversion of wild habitat to agriculture has

taken place, meaning the pollinator may not be present in an agricultural setting. Finally, unlike

other tuberous perennials, such as Yam (*Dioscorea*) (Mengesha et al., 2013) or Cassava (*Manhiot*)

(Rival and Mckey, 2008), wild or sexually reproduced enset seedlings are not (knowingly)

incorporated into cultivated populations (Borrell et al., 2020), except for the localised harvest of

leaves from wild plants (see Discussion). Therefore, little to no indigenous knowledge pertains to

enset seed germination.

Several previous studies have found enset germination to be highly variable (0-90% success)

(Tesfaye, 1992; Messele, 1994; Diro et al., 2003). The most detailed study to date by Karlsson et al

(2013) found 5-55% germination depending on the accession. They found that exposure to sulphuric

acid, sodium hydroxide, ammonium nitrate, sodium hypochlorite or hot water had no significant

positive effect on germination; and indeed, scarification and 70% ethanol had significant negative

effects. Whilst seed germination ecology is also poorly understood in *Musa,* hindering conservation

and use in breeding (Laliberté, 2016), it has long been know that an alternating temperature regime

is a key requirement for germination in *M. balbisiana* (Stotzky et al., 1962). Separately, Simmonds

(1962) reported that immediately after clearing of tropical forest, banana seedlings will readily

emerge. This is thought to be due to exposure of the seeds on the soil surface to alternating day and

night temperatures. Using alternating temperatures as a germination cue may be a strategy

conserved in genus *Ensete*, and is therefore a useful starting point for investigating enset

germination behaviour.

Here, we present the largest survey to date of wild and domesticated enset seed morphology and

germination behaviour, aimed at evaluating evidence for reduced sexual reproductive potential in

this domesticated vegetative crop. We attempt to distinguish between multiple plausible scenarios.

Specifically, if sexual reproductive capacity has been conserved through domestication, we

hypothesise that we should observe comparable seed morphology, viability and germination

behaviour between wild and cultivated enset plants (see for example maintenance of silicon

herbivore-defence through domestication in grasses, Simpson et al., 2017), or potentially, continued

selection for the domesticated environmental niche (Meyer et al., 2012). An alternative hypothesis

would be that, after being released from selection pressure, enset reproductive traits would be

subject to deleterious somatic mutations and genetic drift. Therefore, traits may be expected to

display higher variance in domesticated individuals than wild plants. A third potential scenario

entails a bottleneck (though these are likely weaker in perennial vegetative crops (Gaut et al., 2015))

in which domesticated enset represents a subset of wild diversity, and therefore a subset of the

morphological or behavioural diversity in reproductive traits.

To address these scenarios, we frame our analysis with three questions; i) Is there evidence for

changes in seed morphology or viability in domesticated compared to wild enset? ii) Does seed

germination behaviour differ between wild and domesticated enset, and how do these germination

requirements relate to the climate of wild and domesticated enset distributions? And iii) do

domesticated enset seed traits exhibit higher variance than under neutral expectations. Finally, we

conclude by outlining the most favourable enset seed germination protocol to support germplasm

conservation of a regionally important food security crop.

MATERIALS AND METHODS

Observations, collection, processing and storage

Enset in Ethiopia is readily distinguished in the field as the only member of its genus and may be

subdivided into i) domesticated clonal landraces under farmers' vernacular names, and ii) wild

occurring, sexually reproducing populations. In this study we collected a total of 20 seed accessions

from seven wild and 13 domesticated individuals, in Spring 2018 (Table 1). Domesticated accessions

were collected with permission from farmers' fields, and wild collections were made in river valleys

at least 1 km from settlements cultivating enset, to mitigate the risk of feral or recently introgressed

individuals. Where possible domesticated and wild accessions were collected in the same localities

to mitigate differences arising from the climate the parent plants were exposed to. Through our

extensive fieldwork for this, and related studies, we maintained a database of a) vernacular names

of all observed and documented landraces, and b) those observed flowering, to help resolve the

question of whether all landraces produce flowers. We ranked both lists by the frequency at which

landraces were observed and compared ranks using a Wilcoxon rank sum test implemented in R

software (R Core Team, 2017) to ascertain whether landraces observed flowering are a random

sample of all observed landraces.

After collection, seeds were extracted from ripe fruits by hand and the pulp containing seeds was

washed thoroughly until all the flesh was removed. After extraction and cleaning, seeds were air

dried at room temperature (~20°C), packaged and transferred to Kew where they were stored at a

constant 60% relative humidity for 1-2 weeks (approximate humidity of collection region) in

preparation for germination trials. In addition to field collected enset seed, we also used non-

Ethiopian enset seeds legally sourced from the online horticultural distributor Rare Palm Seeds

(https://www.rarepalmseeds.com/) which are available in large quantities (hereafter RPS),

permitting us to screen a wider range of experimental conditions and act, to a certain extent, as a

control across all experiments.

Morphological characterisation and viability

For each accession, 20-50 seeds were weighed individually, using a balance accurate to 0.001g. Seed

size (diameter on an x, y, and z axis) was recorded for 10-20 seeds per accession using a digital

caliper accurate to 0.01mm. Due to the highly non-uniform shape of enset seeds, with no consistent

long or short axis, seed volume (estimated as the cube of the three measured axes) was used for

subsequent analyses. We used a Faxitron Ultrafocus x-ray (Faxitron Bioptics, LLC), to measure total

seed area (cross section), endosperm area and testa thickness (outer integument) in five seeds per

accession (Figure 2). Seeds were positioned with the proximal end facing upwards, and results

averaged by accession. We tested each dataset for homogeneity of variances and normality before

applying the appropriate unpaired t-test to evaluate differences in population means between

domesticated and wild seeds. All analyses were conducted in R software (R Core Team, 2017).

Tetrazolium tests were used to detect living tissue and viable seeds. Briefly, seeds were imbibed for

24 hours, chipped to expose the endosperm and placed in 1% buffered 2,3,5-triphenyl tetrazolium

chloride for two days in the dark at 30°C. Subsequently, seeds were carefully dissected, and the

embryo staining pattern recorded. Germination proportions were corrected for viability during

subsequent analysis.

Germination trials

We performed four germination experiments (Table 2), with temperature ranges selected based on WorldClimV2 values for the region (Fick and Hijmans, 2017), previous reports from *Musa* (Stotzky et al., 1962; Messele, 1994) and our knowledge of enset ecology in Ethiopia.

i) First, using RPS seed we screened a range of 15 constant and alternating temperature regimes to guide our experimental design for subsequent germination tests (Exp 1). Constant conditions comprised six regimes: 10°C, 15°C, 20°C, 25°C, 30°C and 40°C. Alternating conditions included 10-20°C of diurnal variation to simulate larger temperature shifts, comprising an additional nine regimes: 20/10°C, 25/10°C, 25/15°C, 30/10°C, 30/15°C, 30/20°C, 35/20°C, 40/20°C, 40/25°C.

ii) Based on results from the initial screening, seeds from seven wild accessions and seven domesticated accessions (those with sufficient seeds available for fully replicated trials) were exposed to a refined range of 11 temperature regimes (Exp 2).

iii) To evaluate the relative importance of an absolute shift in ambient temperature (e.g. warming due to disturbance) and regular diurnal temperature shifts (i.e. our alternating temperature regimes) we exposed a subset of domesticated seed to five constant temperatures for three months, and then moved them to the 'optimum' alternating temperature identified in earlier tests (Exp 3).

iv) To evaluate the extent and influence of dormancy, a subset of domesticated accessions was stratified at 10°C for three months, and then transferred to five other temperature regimes (Exp 4).

In all germination tests seeds were placed on moist sand (300g sand, 42ml de-ionised water) and sealed in clear plastic boxes (120 x 180 mm). Boxes were then sealed in plastic bags to further minimise moisture loss and contamination. Each box was considered a replicate and contained 60 seeds (except where specified), with a total of 235 replicates evaluated in this study. Seed boxes

were placed in the corresponding incubators with either constant temperature or 12 hour

alternating temperature cycles. All treatments included a 12-hour photoperiod, although Musaceae

seeds are unlikely to be photosensitive (data not shown; pers. comm. S. Kallow 2019). Germination

was defined by radicle emergence ≥2 mm, and tests were scored every 3-18 days depending on

activity for 120 days. We calculated the mean days to germination retrospectively as the number of

days required for 50% of the final germination count in each experimental replicate. We aggregated

these data by individual, excluding any replicate with zero germination.

To test for a significant difference between wild and domesticated germination behaviour (EXP2),

we fitted polynomial regression models for the logit transformed germination proportion against a)

the daily temperature change the replicate was exposed to and b) the mean temperature the

replicate was exposed to. For each variable we fitted two models, the first with all accessions and

the second with an additional variable grouping the data by type (wild v domesticated). We then

used ANOVA to test whether grouping produced a significantly better model fit. To evaluate the role

of stratification and dormancy (EXP3), we plotted the absolute temperature change between the

first temperature and the mean of the second temperature of the treatments, against germination

proportion and applied linear regression. The effect of a 10°C stratification treatment was compared

to non-stratification using an unpaired t-test (EXP4).

Climate data for Ethiopia were sourced from WorldClim v2 (Fick and Hijmans, 2017) at 2.5 arc

minute resolution (~10 km). In the first instance, we extracted climate values for our study

accessions, and tested for significant differences in Annual Mean Temperaure and Mean Diurnal

Range using unpaired t-tests. We then collated 472 enset localities from GBIF (GBIF.org, 2018),

publications (Borrell et al., 2019; Pironon et al., 2019) and personal observations, and subsampled

these to a 10 km grid consistent with the environmental data layers, retaining 94 unique

domesticated records and 19 unique wild records. We extracted climate data for these cells and

aggregated them for domesticated and wild enset separately.

P$_{ST}$ — F$_{ST}$ comparison

To test whether divergence of quantitative seed and germination traits are putatively as a result of

natural selection or genetic drift we compared variance in trait values with neutral genetic

differentiation. Genetic samples were collected for ongoing study (White et al. *in prep*) and are not

identical individuals to those used for seed and germination traits due to difficulties extracting DNA

from fully mature and senescing plants. Instead we selected a larger dataset of 21 domesticated

individuals matched to the same named landraces surveyed for seed and germination traits, and 14

wild individuals for comparison. DNA was extracted, prepared as libraries and sequenced following

the protocol of Ott et al., (2017). SNPs were called following the method of (Borrell et al., 2018) (see

Data Accessibility for scripts). We calculated neutral F$_{ST}$ between wild and domesticated enset

populations based on 11,412 putatively neutral SNPs, using the R packages Adegenet and Genepop

(Jombart, 2008; Rousset, 2012). Population differentiation for quantitative traits was estimated as

P$_{ST}$ (an approximation of Q$_{ST}$ where the relative contribution of environmental variation is difficult to

ascertain, see Silva and Silva, 2018) in the package Pstat (Silva and Silva, 2018). Due to limited

knowledge, we generated estimates for a range of heritability values (0.2, 0.4 and 0.6) based on

Visscher et al., (2008).

RESULTS

Observations, collection, processing and storage

We observed domesticated enset flowering in all enset cultivating zones, and across a wide range of

environmental conditions (albeit at a very low frequency), but only a very small number were un-

harvested and therefore able to mature and produce ripe fruits (Table S1). Location and altitude

data for the 20 collected enset seed accessions is provided in Table 1. In our collected accessions,

the number of harvested seeds per plant ranged from 19 to 3200 (mean 600). It was not possible to

ascertain reproductive effort (through, for example, total seed production) due to the longevity of

the infloresence and continued predation by frugivores. We note, however, that on several

occasions throughout our fieldwork in Ethiopia apparently ripe fruits were observed on

domesticated enset plants that had developed infructescences, but that seeds were absent,

undeveloped or very few in number (observed in landraces Nechwe, Kiticho and Badadet). We

believe this is most likely attributed to lack of pollinators. During field surveys, we documented 1864

observations of 453 named landraces from across the enset growing region. In addition, we

recorded 39 flowering individuals of 26 landraces. After ranking by frequency of observation we

found no significant difference in rank order (W = 6251.5, P = 0.60).

Morphological characterisation and viability

The basic structure of enset seeds is similar to that of *Musa* (McGahan, 1961), though enset seeds

are larger. At the proximal end is a micropyle forming a plug, behind which is a capitate embryo, the

base of which is embedded within the endosperm (Figure 2). At the opposite end, in a separate

chamber is the chalazal mass which forms a cavity. We observed that enset seeds appear highly

variable in overall surface texture and the degree of striations.

Seed volume ranged from 1.26 - 6.29 cm^3 and seed weight from 0.20 g - 2.85 g in domesticated

accessions, and 1.15 - 5.05 cm^3 and 0.78 g - 2.66 g respectively in wild accessions (Figure 3). Overall,

wild enset accessions had significantly higher mean weight than domesticated enset (t = -2.52, df =

12.63, P = 0.04), but we found no significant difference in mean seed volume between domesticated

and wild (t = 0.07, df = 19, p = 0.95). Similarly, we found non-significant differences for mean testa

thickness (t = 0.29, df = 14, p = 0.78), mean total seed area (t = -1.57, df = 14, p = 0.14) and mean

endosperm area (t= -1.85, df = 14, p = 0.08) (Figure 3). These patterns of (non)significance were

consistent even if poorly germinating accessions are removed (*i.e.* Deri'ea, Suitiya landraces). Full

morphological data are available in Appendix S1 and S2. Tetrazolium tests showed high variation in

viability across accessions, with both wild and domesticated accessions ranging from 0-100%

viability. Mean viability was 55% and 49.5% for wild and domesticated respectively with no

significant difference (t = 0.12, df = 12.9, p = 0.84) (Table 1).

Germination trials

Mean time to germination was 36 days (sd = 15.7) for domesticated enset and 35 days (sd = 8.6) for

wild enset, with no significant difference detected (t = 0.51, df = 11.89, p = 0.62). In Exp 1.

alternating temperature regimes outperformed constant temperatures, with the exception of

constant 25°C (Figure 4). Based on these data we reduced our suite of temperature regimes in

subsequent experiments (Table 2.). ANOVA analysis of polynomial regression models for

germination behaviour in domesticated and wild enset were significantly different (i.e. grouping by

accession type resulted in significantly better model fit) for both the alternating temperature range

($F_{136,132}$ = 4.32, p = 0.003) and the mean experimental temperature ($F_{136,132}$ = 2.71, p = 0.033)

(Figure 5A). Specifically, we found that domesticated accessions had an improved germination

response in cooler mean temperatures with higher alternating temperature amplitude. Comparison

of germination requirements to regional climatic conditions for wild and domesticated enset found

that domesticated enset is found in local climates that have significantly cooler Annual Mean

Temperatures (AMT) (t = -5.52, df = 31, p = <0.001), with significantly greater Mean Diurnal Range

(MDR) (t = 3.42, df = 19.2, p = 0.003), than wild enset (Figure 5B). Importantly, there was no

significant difference in the AMT (t = -1.77, df = 12.56, p = 0.10) or MDR (t = -1.63, df = 13.96, p =

0.13) of our collected accessions, therefore this is unlikely to be solely a maternal effect.

Analysis of Exp 3. found a significant positive relationship between germination proportion and the

absolute temperature change from a constant to alternating temperature regime ($F_{1,38}$ = 6.37, p =

0.0159) (Figure 6A). Analysis of Exp 4. found that a three-month period of cold stratification at 10°C

prior to an experimental treatment also significantly improved germination compared to seeds

immediately exposed to the experimental treatment (t = 5.22, df = 26.4, p = <0.001) (Figure 6B). Full

germination data are available in Appendix S3 and S4.

P_{ST} — F_{ST} comparison

Estimated P_{ST} values varied substantially, with values for seed volume and testa thickness being

consistently close to zero, and seed weight, endosperm area and total seed area relatively high. Full

results across a range of assumed heritability values are reported in Table 3. Pairwise F_{ST} between

domesticated and wild enset populations was estimated at 0.188.

Optimum enset germination conditions

Based on a broad range of experimental conditions, we found an alternating temperature regime of

25/15°C, ideally with a period of moist cool stratification at 10°C for up to 3 months prior to be most

effective for domesticated enset. Most germination can be expected over 20-50 days, with a

germination proportion of ~30-50%. Considering that this germination success is still low, there may

be further opportunities to improve this protocol.

DISCUSSION

It has frequently been suggested that prolonged vegetative reproduction during domestication can

lead to the loss of sexual reproductive capacity (Eckert, 2002; Barrett, 2015; Denham et al., 2020;

Tesfamicael et al., 2020). In this study we show evidence that the indigenous Ethiopian vegetative

crop enset has retained viable sexual reproductive potential comparable to that of wild plants.

Currently, the duration over which enset has been domesticated, and the temporal advent of

vegetative reproduction is unclear (Borrell, Biswas, et al., 2019). However, if we consider the

extensive accumulation of indigenous knowledge associated with enset cultivation, particularly

clonal propagation (Garedew et al., 2017), and its origins in the Ethiopian Highlands, an important

and ancient center of crop domestication (Harlan, 1971), it is reasonable to conclude that enset has

a modest to long domestication history, and that vegetative propagation has been practiced for a

significant portion. This suggests that the potential for sexual reproduction has been maintained

despite prolonged vegetative propagation.

In our surveys, we found no evidence that certain landraces have lost the propensity to flower,

which appears consistent with farmer perceptions (pers. obs. J. Borrell). We also found no evidence

of differing seed viability rates between wild and domesticated enset. Though similar to other

authors (Karlsson et al., 2013), we did find high variability in seed viability across accessions. In a

comparison of wild and domesticated enset seed morphology, we found a significant difference only

for seed weight, with domesticated seeds generally being lighter. There was also a similar, but non-

significant trend for domesticated seeds to have a smaller endosperm area, whilst seed volume

showed no such pattern (Figure 3). Less dense seeds could be explained by reduced resource

provisioning, however this seems unlikely to be due to reduced maternal resource acquisition in an

agricultural context where enset receives substantial fertiliser (manure) input and cultural practices

aim to delay maturation to maximise energy accumulation (Borrell et al., 2020). An alternative

explanation is that highly fertile agricultural systems have relaxed selection on energy investment in

seeds, meaning that seeds with reduced resources may still be sufficiently fit for establishment,

though we are cautious against over interpreting these data.

Our germination trials indicate that not only has enset maintained the capacity for viable seed

production, but surprisingly, that the optimum germination requirements significantly differ

between domesticated and wild enset (Figure 5). Domesticated enset has an increased germination

response in cooler mean temperatures (~22°C) and with an increased amplitude of alternating

temperature (Figure 5A) compared to wild enset. Naturally, wild enset occupies consistently warm

moist tropical forest in Western Ethiopia, whereas the predominant region of contemporary enset

cultivation is a region with lower Annual Mean Temperature and higher Mean Diurnal Range (Figure

5B). This suggests, that wild and domesticated enset are potentially adapted to their respective

environments. Importantly, there was no significant difference between the Annual Mean

Temperature and Mean Diurnal Range of the wild and domesticated seed collection sites surveyed

here, suggesting that this observation is unlikely to be solely a maternal effect.

When we consider enset seed morphology and germination behaviour together, there are several

evolutionary explanations for these observations. First, a scenario where selection pressure has

been relaxed as a result of domestication, would culminate in domesticated enset reproductive

traits that have been subject to deleterious somatic mutations and genetic drift. In this situation we

may expect to see reduced seed viability, or at least consistently higher variance in domesticated

enset traits and $P_{ST} \approx F_{ST}$, which is not the case. If pervasive genetic drift is indeed in progress, then

potentially insufficient time or generations have passed for it to become apparent. Alternatively, in a

scenario where 'perfect' vegetative propagation has been maintained, seed morphology and

germination behaviour are expected to remain consistent between wild and domesticated enset, as

clonal lineages are essentially in stasis. Our data do not provide strong evidence for this, as we

detect significant differences in seed weight and germination behaviour. Instead, this could be

explained by a domestication bottleneck, whereby the progenitors of domesticated enset landraces

represent a subset of the total trait variation (i.e. domesticated enset arose from a population of

wild enset preferring cooler, more variable climes with lighter seeds). However this also seems

unlikely, given that i) in perennial crops we expect to observe a weak domestication bottleneck

(Gaut et al., 2015), ii) wild and domesticated appear adapted to their respective environmental

conditions and iii) based on climax vegetation maps of Ethiopia it is unlikely that wild populations

originated from further East in Ethiopia, in the cooler, more variable climates which enset cultivation

currently occupies (Friis et al., 2010).

Seeking a more parsimonious scenario, we suggest that despite virtually exclusive clonal propagation

in cultivation, it is possible that a small number of escaped or neglected domesticated plants are

continuing to reproduce sexually. Where this has occurred in a novel agricultural environment,

certain seed and germination traits have been subjected to directional selection, whilst others have

been maintained under balancing selection. This explanation is supported by both significantly

different germination behaviour concordant with local environment, evidence for local adaptation in

seed weight and endosperm area ($P_{ST} > F_{ST}$), and reduced variance in domestic versus wild indicating

putatively stabilising selection ($P_{ST} < F_{ST}$) in traits such as seed volume and testa thickness (or weak,

undetectable genetic drift). Previous work has shown that a comparatively low rate of sexual

reproduction would be sufficient to maintain this balancing selection (Rice and Chippindale, 2001;

Cutter, 2019). We conclude that this best describes the patterns we have observed, not only based

on the data report here, but also based on observation in other clonal crops that farmers actively

manage infrequent sexual reproduction. Examples include the practice of enoblement in Yams

(*Dioscorea*) (Cornet et al., 2010; Mengesha et al., 2013), or tolerance of volunteer manioc seedlings

in Cassava (*Manhiot*) (Rival and Mckey, 2008).

To further support this inference, we provide two qualitative observations of enset agronomic

practices that could contribute to the persistence of sexual reproduction. First, in Western regions

(e.g. Gesha and Bonga), small numbers of enset seedlings are collected from the wild and integrated

into the farm for the purpose of harvesting leaves (leaves are harvested from inedible wild plants,

where available, to avoid damaging and slowing the growth of edible domesticated plants). Similarly,

in a small number of cases, comparatively wealthy farmers occasionally allow a single domesticated

enset, often in a highly visible location near the front of their compound, to flower as a sign of

prosperity (pers. obs. J Borrell). Both these practices could increase the frequency of sexual

recombination, and the subsequent survival of seedlings, with associated impacts on the evolution

and domestication of enset.

More broadly, we note that enset germination requirements appear consistent with reports from

Musa (Stotzky et al., 1962, Kallow et al., in review). Specifically, in enset, alternating temperatures

elicit a stronger germination response than constant temperatures, though in *Musa* this is virtually

an absolute requirement. Using alternating temperatures as an environmental cue is hypothesised

to be a strategy for detecting disturbance and canopy gaps whereby solar radiation warms the seeds

in the day followed by a cooler ambient temperature at night. Enset is also reported to colonise

disturbed areas (Stotzky et al., 1962), and thus this trait appears to be conserved across the two

major branches of the Musaceae. Surprisingly, the magnitude of the transition from constant to

alternating temperature was also significantly associated with germination (Figure 6A). Kallow et al

(in review) hypothesise that in *Musa* seeds, alternating temperatures reduce the ratio of Abscisic

acid to Gibberellic acid (GA), reducing water potential and initiating elongation and cell growth. This

suggests that future approaches involving application of GA to seeds or the growing medium may

provide another mechanism for initiating germination. In addition, increased germination is also

observed where seeds were stratified at a constant temperature prior to germination. Climate data

indicates that the domesticated distribution of enset may reach a minimum of 8.2°C in the coolest

month, which coincides with higher rainfall. Whilst no seasonality has been reported in enset

flowering, this may be an additional, putatively conserved, mechanism for optimising germination

timing.

Despite the importance of understanding the germination biology of enset, this study was limited by

the fact that we were not able to determine total seed production of enset individuals, as

inflorescences ripen over a considerable time, and may be damaged by animals in the process,

though this would provide an important, comparative metric for sexual reproductive potential.

Similarly, we were also limited by the available sample size, in a rarely flowering crop, compounded

by significant variation associated with the accession. We anticipate that numerous factors such as

fruit maturity, length and type of storage, epigenetic and other factors may influence variability,

though we do not have sufficient power to resolve these in this study. To further evaluate

maintenance of enset sexual reproductive potential, we suggest that comparative analysis of the

floral structures of enset, to detect loss of function arising from genetic drift, may corroborate our

findings.

CONCLUSIONS

In the future, development of enset seed germination protocols, underpinned by diverse seed

collections, is a compelling strategy to safeguard the genetic diversity of a food security crop, with a

lower risk of provenance information loss than living plants in a germplasm collection (Thormann

and Dulloo, 2006). Moreover, we show that this is achievable, by demonstrating the maintenance of

sexual reproductive potential in enset, despite a significant period of near-exclusive vegetative

propagation. Furthermore, we provide initial evidence that very low levels of sexual reproduction

may be facilitating the adaptation of enset germination biology to domestication, and that enset

cultural practices have unrecognised parallels with the agronomy of other clonal crops. An improved

understanding of enset germination biology is a useful prerequisite for future crop development

through sexual recombination of existing landraces, developing mapping populations and the

breeding of novel genotypes, with significant reciprocal potential in bananas (*Musa*) a closely related

and globally important group of clonally produced crops. In conclusion, we advocate for a broader

effort in developing germination protocols for vegetatively propagated crops to provide an

important alternative germplasm conservation strategy that is likely to disproportionately benefit

tropical species and developing country agriculture in the global south.

ACKNOWLEDGEMENTS

The authors thank the Ethiopian Biodiversity Institute for facilitating the export of enset seeds for

germination studies. We thank John Adams and Pablo Barreiro at the Millennium Seed Bank for lab

support and Dr Yann Dussert, Prof Richard Nichols, Prof Pat J.S. Heslop-Harrison and Dr Manosh

Biswas for fruitful discussions on the evolutionary ecology of vegetative reproduction. We also thank

Robert Kessler for permitting publication of enset photographs. Finally, we thank Mr Tobias Spanner

from Rare Palm Seeds for helpful information on seed provenance and morphological variation. This

work was supported by the GCRF Foundation Awards for Global Agricultural and Food Systems

Research, entitled, 'Modelling and genomics resources to enhance exploitation of the sustainable

and diverse Ethiopian starch crop enset and support livelihoods' [Grant No. BB/P02307X/1]. In

addition, part of this work was funded as a subgrant from the University of Queensland from the Bill

& Melinda Gates Foundation project 'BBTV mitigation: Community management in Nigeria, and

screening wild banana progenitors for resistance' [OPP1130226].

AUTHOR CONTRIBUTIONS

484 PW and SD conceived the project and acquired funding. JB, SK and RD designed the study. ST, JB, ES,

485 GW and FW collected material. ST, JB, ES, SK and RD performed experiments. JB analysed data and

486 wrote the first draft of the manuscript. All authors contributed to and approved the final version of

487 the manuscript.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information section at the

end of the article

Appendix S1 – Seed_morphology_(size).txt

Appendix S2 – Seed_morphology_(weight).txt

Appendix S3 – Germination_Exp_1-2.txt

Appendix S4 – Germination_Exp_3-4.txt

Genetic processing and SNP calling scripts are available here: https://github.com/o-william-

white/Enset_tGBS.

LITERATURE CITED

Anon. 2018. GBIF.org. *Page, GBIF Home*.

Barrett, S. C. H. 2015. Influences of clonality on plant sexual reproduction. *Proceedings of the*

National Academy of Sciences 112: 8859–8866.

Blomme, G., M. Dita, K. S. Jacobsen, L. Pérez Vicente, A. Molina, W. Ocimati, S. Poussier, and P. Prior.

2017. Bacterial Diseases of Bananas and Enset: Current State of Knowledge and Integrated

Approaches Toward Sustainable Management. *Frontiers in Plant Science* 8: 1–25.

Borrell, J. S., M. K. Biswas, M. Goodwin, G. Blomme, T. Schwarzacher, P. J. S. Heslop-Harrison, A. M.

Wendawek, et al. 2019. Enset in Ethiopia: a poorly characterised but resilient starch staple.

Annals of Botany 123 (5): 747–766.

Borrell, J. S., S. Dodsworth, F. Forest, O. A. Perez-Escobar, M. A. Lee, E. Mattana, P. C. Stevenson, et

al. 2019. The climatic challenge: Which plants will people use in the next century?

Environmental and Experimental Botany 170.

Borrell, J. S., M. Goodwin, G. Blomme, K. Jacobsen, A. M. Wendawek, D. Gashu, E. Lulekal, et al.

2020. Enset based agri-systems in Ethiopia: A systematic review of production trends,

agronomy, processing and the wider food security applications of a neglected banana relative.

Plants, People, Planet 00: 1–17.

Borrell, J. S., N. Wang, R. A. Nichols, and R. J. A. Buggs. 2018. Genetic diversity maintained among

fragmented populations of a tree undergoing range contraction. *Heredity* 121: 304–318.

Castañeda-álvarez, N. P., C. K. Khoury, H. A. Achicanoy, V. Bernau, H. Dempewolf, R. J. Eastwood, L.

Guarino, et al. 2016. Global conservation priorities for cropwild relatives. *Nature Plants* 2: 1–6.

Chen, L., R. Vanburen, M. Paris, H. Zhou, X. Zhang, C. M. Wai, H. Yan, et al. 2019. The bracteatus

pineapple genome and domestication of clonally propagated crops. *Nature Genetics* 51.

Commission for genetic resources on food and agriculture. 2010. The Second Report on the state of

the world's plant genetic resources for food and agriculture. Rome.

Conway, D., and E. L. F. Schipper. 2011. Adaptation to climate change in Africa: Challenges and

opportunities identified from Ethiopia. *Global Environmental Change* 21: 227–237.

Cornet, D., M. Deu, M. N. Baco, A. Agbangla, M. F. Duval, and J. L. Noyer. 2010. Impact of farmer

selection on yam genetic diversity. *Conservation Genetics* 11: 2255–2265.

Cutter, A. D. 2019. Reproductive transitions in plants and animals : selfing syndrome , sexual

selection and speciation. *New Phytologist* 224: 1080–1094.

Dempewolf, H., R. J. Eastwood, L. Guarino, C. K. Khoury, J. V. Müller, and J. Toll. 2014. Adapting

Agriculture to Climate Change: A Global Initiative to Collect, Conserve, and Use Crop Wild

Relatives. *Agroecology and Sustainable Food Systems* 38: 369–377.

Denham, T., H. Barton, C. Castillo, A. Crowther, E. Dotte-sarout, S. A. Florin, J. Pritchard, et al. 2020.

The domestication syndrome in vegetatively propagated field crops. *Annals of Botany* XX: 1–17.

Diro, M., J. van Staden, and C. H. Bornman. 2003. In vitro regeneration of Ensete ventricosum from

zygotic embryos of stored seeds. *South African Journal of Botany* 69: 364–369.

Dulloo, M. E., I. Thormann, E. Fiorino, S. De Felice, V. R. Rao, and L. Snook. 2013. Trends in Research

using Plant Genetic Resources from Germplasm Collections: From 1996 to 2006. *Crop Science*

53: 1217–1227.

Eckert, C. G. 2002. The loss of sex in clonal plants. *In* J. F. Stuefer, B. Erschbamer, H. Huber, and J.-I.

Suzuki [eds.], Ecology and Evolutionary Biology of Clonal Plants, 279–298. Springer

Netherlands, Dordrecht.

Elias, M., L. Penet, P. Vindry, D. McKey, O. Panaud, and T. Robert. 2001. Unmanaged sexual

reproduction and the dynamics of genetic diversity of a vegetatively propagated crop plant,

cassava (Manihot esculenta Crantz), in a traditional farming system. *Molecular Ecology* 10:

1895–1907.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. *International Journal of Climatology*.

Friis, I., S. Demissew, and P. Breugel. 2010. Atlas of the Potential Vegetation of Ethiopia. *Biologiske*

Skrifter (Biol.Skr.Dan.Vid.Selsk.) 58: 307.

Fu, Y. 2017. The Vulnerability of Plant Genetic Resources Conserved Ex Situ. *Crop Science* 57.

Garedew, B., A. Ayiza, B. Haile, and H. Kasaye. 2017. Indigenous Knowledge of Enset (Ensete

ventricosum (Welw .) Cheesman) Cultivation and Management Practice by Shekicho People ,

Southwest Ethiopia. *Journal of Plant Sciences* 5: 6–18.

Gaut, B. S., C. M. Díez, and P. L. Morrell. 2015. Genomics and the Contrasting Dynamics of Annual

and Perennial Domestication. 31: 709–719.

Guzzon, F., and J. V. Müller. 2016. Current availability of seed material of enset (Ensete ventricosum,

Musaceae) and its Sub-Saharan wild relatives. *Genetic Resources and Crop Evolution* 63: 185–

191.

Harlan, J. 1971. Agricultural Origins: Centers and Noncenters. *Science* 239: 17–18.

Heslop-Harrison, J. S., and T. Schwarzacher. 2007. Domestication, genomics and the future for

banana. *Annals of Botany* 100: 1073–1084.

Janssens, S. B., F. Vandelook, E. De Langhe, B. Verstraete, E. Smets, I. Vandenhouwe, and R.

Swennen. 2016. Evolutionary dynamics and biogeography of Musaceae reveal a correlation

between the diversification of the banana family and the geological and climatic history of

Southeast Asia. *New Phytologist* 210: 1453–1465.

Jombart, T. 2008. Adegenet: A R package for the multivariate analysis of genetic markers.

Bioinformatics 24: 1403–1405.

Karlsson, L. M., A. L. Dalbato, T. Tamado, and Y. Mikias. 2014. Effect of cultivar, traditional corm pre-

treatment and watering on sprouting and early growth of enset (Ensete ventricosum) suckers.

Experimental Agriculture 51: 232–243.

Karlsson, L. M., T. Tamado, A. L. Dalbato, and Y. Mikias. 2013. Seed morphology and germination of

Ensete ventricosum (Musaceae). *Seed Science and Technology* 41: 357–370.

Laliberté, B. 2016. Global strategy for the conservation and use of Musa (banana) genetic resources:

a consultative document prepared by the Global Musa Genetic Resources Network (MusaNet).

Bioversity International, Rome.

McClure, K. A., J. Sawler, K. M. Gardner, D. Money, and S. Myles. 2014. Genomics: A potential

panacea for the perennial problem. *American Journal of Botany* 101: 1780–1790.

McGahan, M. W. 1961. Studies on the Seed of Banana. I. Anatomy of the Seed and Embryo of Musa

balbisiana. *American Journal of Botany* 48: 230–238.

McKey, D., M. Elias, M. E. Pujol, and A. Duputié. 2010. The evolutionary ecology of clonally

propagated domesticated plants. *New Phytologist* 186: 318–332.

Mengesha, W. A., S. Demissew, M. F. Fay, R. J. Smith, I. Nordal, and P. Wilkin. 2013. Genetic diversity

and population structure of Guinea yams and their wild relatives in South and South West

Ethiopia as revealed by microsatellite markers. *Genetic Resources and Crop Evolution* 60: 529–

541.

Messele, T. 1994. Enset fruit and seed processing, germination and seedling development.

Meyer, R. S., A. E. Duval, and H. R. Jensen. 2012. Patterns and processes in crop domestication: an

historical review and quantitative analysis of 203 global food crops. *New Phytologist* 196: 29–

48.

Migicovsky, Z., and S. Myles. 2017a. Exploiting Wild Relatives for Genomics-assisted Breeding of

Perennial Crops. 8: 1–16.

Migicovsky, Z., and S. Myles. 2017b. Exploiting Wild Relatives for Genomics-assisted Breeding of

Perennial Crops. *Frontiers in Plant Science* 8: 1–16.

Miller, A. J., and B. L. Gross. 2011. From forest to field: Perennial fruit crop domestication. *American*

Journal of Botany 98: 1389–1414.

Ott, A., S. Liu, J. C. Schnable, C. T. Yeh, K. S. Wang, and P. S. Schnable. 2017. tGBS®genotyping-by-

sequencing enables reliable genotyping of heterozygous loci. *Nucleic Acids Research* 45.

Pironon, S., T. R. Etherington, J. S. Borrell, N. Kuhn, M. Macias-Fauria, I. Ondo, C. Tovar, et al. 2019.

Potential adaptive strategies for 29 Sub-Saharan crops under future climate change. *Nature*

Climate Change 9: 758–763.

Powell, W., M. Halewood, T. Chiurugwi, R. S. Hamilton, B. Kurtz, E. Marden, E. Welch, et al. 2018.

Tansley review Plant genetic resources for food and agriculture : opportunities and challenges

emerging from the science and information technology revolution. *New Phytologist* 217: 1407–

1419.

R Core Team. 2017. R: A language and environment for statistical computing. v3.4.1.

Rice, W. R., and A. K. Chippindale. 2001. Sexual Recombination and the Power of Natural Selection.

Science 294: 555–559.

Rival, L., and D. Mckey. 2008. Domestication and Diversity in Manioc (Manihot esculenta Crantz ssp.

esculenta, Euphorbiaceae). *Current Anthropology* 49: 1119–1128.

Rousset, F. 2012. Genepop 4.2.

Scarcelli, N., S. Tostain, Y. Vigouroux, C. Agbangla, O. Dainou, and J. -L. Pham. 2006. Farmers' use of

wild relative and sexual reproduction in a vegetatively propagated crop . The case of yam in

Benin. *Molecular Ecology* 15: 2421–2431.

Silva, S. B. Da, and A. Da Silva. 2018. Pstat : An R Package to Assess Population Differentiation in

Phenotypic Traits. *The R Journal* 10: 447–454.

Simmonds, N. W. 1962. The evolution of the bananas. Longmans, Green, London.

Simpson, K. J., R. N. Wade, S. E. Hartley, M. Rees, C. P. Osborne, and S. E. Hartley. 2017. Still armed

after domestication? Impacts of domestication and agronomic selection on silicon defences in

cereals. *Functional Ecology* 31: 2108–2117.

Ssali, F., and D. Sheil. 2019. Some Initial Observations Concerning the African Wild Banana Ensete

ventricosum as a Resource for Vertebrates. *Tropical Conservation Science* 12: 1–7.

Stotzky, G., E. A. Cox, and R. D. Goos. 1962. Seed Germination Studies in Musa. I. Scarification and

Aseptic Germination of Musa balbisiana. *American Journal of Botany* 49: 515–520.

Tamrat, S., J. S. Borrell, M. K. Biswas, D. Gashu, T. Wondimu, and A. Carlos. 2020. Micronutrient

composition and microbial community analysis across diverse landraces of the Ethiopian

orphan crop enset. *BioRxiv*.

Tesfamicael, K. G., E. Gebre, T. J. March, B. Sznajder, D. E. Mather, and C. M. R. Lopez. 2020.

Accumulation of mutations in genes associated with sexual reproduction contributed to the

domestication of a vegetatively propagated staple crop, enset. *bioRxiv*: 1–31.

Tesfaye, M. 1992. The effect of soaking, temperature and other pretreatments on the germination

of" ensete" seed. Proceedings of the International Seed Testing Association,.

Thormann, I., and M. E. Dulloo. 2006. Techniques for ex Situ for Situ Plant Conservation. *In* H. RJ

[ed.], Plant Conservation Genetics, Haworth Press.

Visscher, P. M., W. G. Hill, and N. R. Wray. 2008. Heritability in the genomics era — concepts and

misconceptions. *Nature Reviews Genetics* 9: 255–266.

Wu, G. A., J. Terol, V. Ibanez, A. López-garcía, E. Pérez-román, C. Borredá, C. Domingo, et al. 2018.

Genomics of the origin and evolution of Citrus. *Nature* 554: 311–317.

    TABLES

658   **Table 1.** Wild and domesticated enset accessions used for germination studies

Origin	Landrace name	Elevation (m a.s.l.)	Latitude (°N)	Longitude (°E)	Tetrazolium viability (%)	Annual Mean Temp. (°C)	Mean Diurnal Range (°C)
Domesticated	Addis Ababa	2430	9.03	38.76	84	16.02	13.25
Domesticated	Maze	2120	6.87	37.79	64	18.78	14.44
Domesticated	Hala	2120	6.87	37.79	66	18.78	14.44
Domesticated	Deri'ea	2700	7.93	37.90	0	15.37	12.97
Domesticated	Wanadiya	2125	6.86	37.79	48	18.78	14.44
Domesticated	Gefetano	2125	6.86	37.79	46	18.78	14.44
Domesticated	Gefetano	1930	6.83	37.75	75	20.24	14.72
Domesticated	Ganticho	1840	6.47	38.35	30	18.41	14.25
Domesticated	Suitiya	2120	6.87	37.79	0	18.78	14.44
Domesticated	Addis Ababa	2420	9.02	38.78	100	16.02	13.25
Domesticated	Midasho	2786	6.47	38.54	14	12.89	13.67
Domesticated	Kiticho	2787	6.48	38.54	50	12.89	13.67
Domesticated	Lemat	2053	8.45	38.03	67	17.97	13.49
Wild	W1	1880	7.17	36.22	0	18.78	14.35
Wild	W2	1850	7.16	36.20	38	18.47	14.30
Wild	W3	1936	7.16	36.20	54	18.47	14.30
Wild	W4	1936	7.16	36.20	68	18.47	14.30
Wild	W5	1927	7.16	36.20	78	18.47	14.30
Wild	W6	1936	7.19	36.20	52	18.11	14.14
Wild	W7	1930	7.29	36.14	100	17.97	13.99
Wild (non-Ethiopian)	'Rare Palm Seeds'	-	-	-	94	-	-

667   **Table 2.** Overview of the four main germination experiments performed

Experiment	No. of conditions	Replicates	Temperature regimes
EXP1: Full range (RPS seeds) to test preferred temperature	6 constant, 9 alternating	15 replicates (900 seeds)	10°C, 15°C, 20°C, 25°C, 30°C, 40°C, 20/10°C, 25/10°C, 25/15°C, 30/10°C, 30/15°C, 30/20°C, 35/20°C, 40/20°C, 40/25°C
EXP2: Wild and domestic	5 constant, 6 alternating	77 replicates (3213) wild; 88 replicates (3844 seeds) domestic	10°C, 15°C, 20°C, 25°C, 30°C, 20/10°C, 25/10°C, 25/15°C, 30/10°C, 30/15°C, 30/20°C
EXP3: Domestic and RPS - testing shift from constant to alternating temperature	5 constant, moved to 1 alternating	40 replicates (1719 seeds)	10°C, 15°C, 20°C, 25°C, 30°C moved to 25/10°C
EXP4: Domestic and RPS - subset tested after cold stratification at 10°C	1 constant, 4 alternating	15 replicated (850 seeds)	10°C moved to 25°C, 25/10°C, 25/15°C, 30/20°C, 30/15°C

680 **Table 3.** Phenotypic trait differentiation and trait variance in domesticated and wild enset.

Morphological trait	P_{ST} heritability estimates			Trait variance	
	0.2	0.4	0.6	Domesticated	Wild
Seed volume (mm^3)	0.00	0.01	0.01	578,127	324,302
Total seed area (mm^2)	0.20	0.33	0.42	1,678	679
Endosperm area (mm^2)	0.25	0.41	0.51	162	210
Seed weight (g)	0.36	0.53	0.62	0.13	0.22
Testa thickness (mm)	0.01	0.02	0.02	0.45	0.18
Mean F_{ST}	0.19	0.19	0.19	-	-

697 **FIGURES**

700 **Figure 1.** Enset cultivation in Ethiopia. A) Mature enset flowering in a neglected field near Checha. B)

701 A farmer removing the meristematic tissue of a two-year old enset, as part of processing for

702 vegetative propagation. C) Numerous adventitious buds sprouting from prepared corm, near Bonga.

714 **Figure 2.** X-ray images of *Ensete ventricosum* seeds. Scale bar denotes 5mm. A) Domesticated

715 accession 'Wanadiya'. A poorly filled seed can be observed in the lower center of the image. B) Wild

716 accession six 'W6'. C) Major enset seed morphological traits referred to in this study: 1. Hilum. 2.

717 Micropylar plug. 3. Embryo. 4. Endosperm surrounding embryo. 5. Inner integument. 6. Outer

718 integument (testa) 7. Chalazal mass.

731   **Figure3.** Boxplots comparing seed traits in wild and domesticated enset. Asterisks denote traits in

732   which we observed a significant difference.

Figure 4. Percentage germination across a range of environmental conditions for wild and domesticated enset (Exp 2), after 120 days. Data are corrected for variation in seed viability.

Figure 5. Germination behaviour and regional climate variables for wild and domesticated enset in

Ethiopia. A) Polynomial regression of logit transformed germination proportion in wild and domestic

accessions under varying mean temperature and alternating temperature regimes. Each point

denotes a germination experimental replicate comprising 60 seeds of a single accession, corrected to

account for variation in seed viability. B) Boxplots of regional climate for domestic and wild enset

records in Ethiopia.

769 **Figure 6.** Analysis of the influence of temperature shifts and dormancy on domesticated enset

770 germination (EXPs 3 and 4). A) The influence of absolute temperature change from constant to

771 alternating temperature, showing a positive relationship with increased germination response for

772 larger temperature shifts. B) Comparison of germination response for stratified seeds (3 months at

773 10°C), versus no stratification (immediate exposure to experimental conditions).