Genome-wide mutagenesis identifies factors involved in *Enterococcus faecalis* vaginal adherence

and persistence.

Norhan Alhajjar[1], Anushila Chatterjee[1], Brady L. Spencer[1], Lindsey R. Burcham[1], Julia L. E. Willett[2],

Garry M. Dunny[2], Breck A. Duerkop[1*], and Kelly S. Doran[1*].

[1]Department of Immunology and Microbiology, University of Colorado School of Medicine, Anschutz

Medical Campus, Aurora, CO, USA.

[2]Department of Microbiology and Immunology, University of Minnesota Medical School, Minneapolis,

MN, USA.

*Corresponding authors: Breck A. Duerkop and Kelly S. Doran, Department of Immunology and

Microbiology, University of Colorado School of Medicine, 12800 East 19th Avenue Aurora, CO, USA.

Phone: (303) 724-3539, Fax: (303) 724-4226, E-mail: breck.duerkop@cuanschutz.edu,

kelly.doran@cuanschutz.edu

RUNNING TITLE: *Enterococcus faecalis* **Vaginal Colonization**

ABSTRACT

Enterococcus faecalis is a Gram-positive commensal bacterium native to the gastrointestinal tract and an opportunistic pathogen of increasing clinical concern. *E. faecalis* also colonizes the female reproductive tract and reports suggest vaginal colonization increases following antibiotic treatment or in patients with aerobic vaginitis. Currently, little is known about specific factors that promote *E. faecalis* vaginal colonization and subsequent infection. We developed a mouse model of *E. faecalis* vaginal carriage and demonstrate that both vancomycin resistant and sensitive strains colonize the murine vaginal tract. Following vaginal colonization, we observed *E. faecalis* in vaginal, cervical and uterine tissue. A mutant lacking endocarditis- and biofilm-associated pili (Ebp) exhibited a decreased ability to associate with human vaginal and cervical cells *in vitro* as well as ascend to the cervix and uterus *in vivo*. To further identify novel factors that promote vaginal colonization, we screened a low-complexity transposon (Tn) mutant library to identify genes important for *E. faecalis* persistence in the vaginal tract. This screen revealed 383 mutants that were underrepresented during vaginal colonization at all time points compared to growth in culture medium. We confirmed that mutants deficient in ethanolamine catabolism and the type VII secretion system were attenuated in persisting during vaginal colonization. These results reveal the complex nature of vaginal colonization and suggest that multiple factors contribute to *E. faecalis* persistence in the reproductive tract.

IMPORTANCE

Despite increasing prevalence and association of *E. faecalis* with aerobic vaginitis, essentially nothing is known about the bacterial factors that influence *E. faecalis* vaginal colonization. We have established an animal model of vaginal colonization that supports colonization of multiple *E. faecalis* strains. Additionally, we determined that pili, ethanolamine utilization, and type VII secretion system genes

contribute to vaginal persistence and ascension to reproductive tract tissues. Identification of factors important for vaginal colonization and persistence provides potential targets for the development of therapeutics. This study is the first to identify key determinants that promote vaginal colonization by *E. faecalis*, which may represent an important reservoir for antibiotic resistant enterococci.

INTRODUCTION

Enterococcus faecalis is an opportunistic pathogen that resides in the human gastrointestinal and urogenital tracts [1, 2]. While *E. faecalis* colonization is normally asymptomatic, certain populations are at risk for severe disease including urinary tract infections (UTIs), wound infections, pelvic inflammatory disease (PID), infective endocarditis, and adverse birth effects during pregnancy [3, 4]. Enterococcal infections are often associated with the production of biofilms, assemblages of microbes enclosed in an extracellular polymeric matrix that exhibit cell-to-cell interactions. These biofilms have been observed on catheters, diabetic ulcers, and wounds resulting in severe infection. Treatment of enterococcal infections is becoming increasingly problematic due to their augmented resistance to antibiotics, including "last-line-of-defense" antibiotics such as vancomycin [5, 6]. Recently, there has been an increase in the emergence of vancomycin resistant enterococci [7], putting immunocompromised individuals at risk for developing severe chronic enterococcal infections. The emergence of vancomycin resistant enterococci (VRE) and its prevalence in both community and nosocomial settings is concerning and necessitates the development of alternative therapeutics to treat enterococcal infections.

E. faecalis encodes a multitude of virulence factors that allow the bacterium to colonize and persist in different sites of the human body. Surface proteins such as the adhesin to collagen (Ace), enterococcal fibronectin binding protein A (EfbA), aggregation substance (AS), and the endocarditis-and biofilm-associated pilin (Ebp) have been previously shown to play important roles in infective endocarditis and

UTIs [8]. Secreted factors such as gelatinase (GelE), autolysin A (AtlA), and serine protease (SprE) are

biofilm-associated factors that are involved in the degradation of host substrates, including collagen, fibrin

and certain complement components [9]. Many of these virulence factors are regulated via quorum

sensing, which may be responsible for the switch from a commensal to pathogenic state [10-12].

Certain risk factors are associated with the transition of *E. faecalis* from commensalism to pathogenicity

such as immune   status,   prolonged   hospital   stay, and   the   use   of   antibiotics   [13]. *E.*

faecalis colonization and infection is often polymicrobial and these interactions have been observed in the

intestine, bloodstream, and wounds [14]. Furthermore, *E. faecalis* is frequently found in the vaginal tract

of healthy women [15, 16] and its prevalence is increased in women diagnosed with aerobic vaginitis

(AV),   an   inflammatory   response   accompanied   by   depletion   of   commensal *Lactobacillus*

sp. and   increased   presence   of   opportunistic   pathogens   such   as *E.   faecalis*,   Group

B *Streptococcus* (GBS), *Staphylococcus aureus*, and *Escherichia coli* [17, 18]. Symptoms of AV include

malodor and discomfort, but AV can transition to more serious complications such as PID, severe UTIs,

and complications during pregnancy. While it is evident that *E. faecalis* colonizes the human vaginal tract,

the molecular determinants that allow enterococci to colonize and persist in the vaginal tract remain to be

identified.

In this study, we developed a murine model of *E. faecalis* vaginal colonization and persistence. We

determined that *E. faecalis* OG1RF (a rifampicin and fusidic acid derivative of strain OG1) and

vancomycin resistant *E. faecalis* V583 can colonize and persist in the vaginal tract of CD1 and C57BL/6

mouse strains. We detected fluorescent *E. faecalis* in the vaginal lumen as well as the cervical and uterine

tissues of colonized mice. Further, we demonstrate that an *E. faecalis* strain lacking Ebp pili is attenuated

in both *in vitro* and *in vivo* models of vaginal colonization. Lastly, we screened an *E. faecalis* OG1RF

transposon (Tn) mutant library for mutants that are underrepresented in the vaginal tract compared the

culture input, revealing multiple factors for *E. faecalis* persistence within the vagina. These factors include

the Ebp pili, ethanolamine utilization genes, and genes involved in type VII secretion system (T7SS)

machinery. We confirmed that mutant strains lacking these genes are significantly attenuated in their

ability to colonize and ascend in the female reproductive tract. This work is an important first step in

identifying factors required for enterococcal vaginal colonization and will provide insight into potential

therapeutic targets aimed at mitigating *E. faecalis* vaginal colonization in at-risk individuals.

RESULTS

***E. faecalis* colonization of the female reproductive tract.**

To characterize the ability of *E. faecalis* to interact with the epithelial cells of the lower female

reproductive tract, we performed *in vitro* quantitative adherence assays using *E. faecalis* strains V583

[19] and OG1RF [20]. An inoculum of 10^5 CFU/well (multiplicity of infection [MOI] = 1) was added to

a confluent monolayer of immortalized human vaginal and endocervical epithelial cells. Following 30

minutes of incubation, the cells were washed to remove non-adherent bacteria, the epithelial cells were

detached from the plates, and adherent bacteria were plated on agar. Data are expressed as percent

recovered cell-associated *E. faecalis* relative to the initial inoculum. Both strains exhibited substantial

adherence to both cell lines (Fig. 1A). Next, we assessed the ability of *E. faecalis* to establish colonization

of the murine vaginal tract. C57BL/6 mice were treated with β-estradiol 1 day prior to inoculation with

10^7 CFU of *E. faecalis* V583 or OG1RF. After 1 day post-inoculation, the vaginal lumen was swabbed

and bacteria were plated to enumerate *E. faecalis* V583 and OG1RF vaginal colonization levels (Fig. 1B).

Swabs were plated on selective agar to ensure quantification of only the enterococcal strains of interest,

restricting growth of native enterococcus. To determine whether *E. faecalis* ascends into reproductive

tissues during colonization, murine vaginal, cervical, and uterine tissues were collected and homogenized

to enumerate *E. faecalis* V583 abundance. *E. faecalis* was recovered from all mice 1 day post-inoculation

in all tissues tested (Fig. 1C) and the CFU recovered from the vaginal swabs were similar to the total CFU

counts from the vaginal tissue homogenates (Fig. 1B, C). This level and range of recovered *E. faecalis*

CFU is similar to what we have observed using this mouse model for GBS and *S. aureus* vaginal

colonization [21, 22]. To visualize *E. faecalis* within the murine reproductive tract, mice were inoculated

with *E. faecalis* V583 expressing green fluorescent protein (GFP) [23]. We harvested the female

reproductive tract 1 day post-inoculation, made serial sections of these tissues, and performed fluorescent

microscopy to visualize *E. faecalis* (Fig. 1D, E, F). We observed numerous fluorescent bacteria in the

vaginal and uterine lumen (Fig. 1D, F) and embedded in the cervical lamina propria (Fig. 1E).

***E. faecalis* persists in the vaginal tract.**

To assess vaginal persistence, C57BL/6 mice were colonized with *E. faecalis* V583 or OG1RF and

swabbed to determine bacterial load over time. We observed that both strains exhibited similar persistence

within the mouse vagina (Fig. 2A, B). The mean CFU recovered remained constant for the first week and

then declined as mice eventually cleared both strains by 11-13 days. To determine if enterococcal vaginal

persistence differs across mouse strains, C57BL/6 and CD1 mice were inoculated with V583 and swabbed

over time. Both mouse strains were initially colonized with V583, but bacteria in C57BL/6 mice persisted

longer (Fig. 2C, D). By day six only 20% of CD1 mice remained colonized compared to 85% of C57BL/6

mice. Differences in vaginal persistence may be due to differences in the native vaginal microbiota or

immune status between mouse strains. These results show that mouse strain background influences *E.*

faecalis vaginal colonization and that C57BL/6 mice are a sufficient model to assess prolonged *E. faecalis*

vaginal colonization and persistence.

Enterococcal pili contribute to interaction with reproductive tract tissues.

The <u>e</u>ndocarditis- and <u>b</u>iofilm-associated <u>p</u>ilin (Ebp) of *E. faecalis* mediates infective endocarditis and UTIs, thus we hypothesized that Ebp may similarly contribute to vaginal colonization. To determine whether Ebp is important for facilitating interaction with the vaginal epithelium, we used a deletion mutant of *E. faecalis* OG1RF lacking all pilin structural components (Δ*ebpABC*) [24]. We observed that the pilus mutant exhibited significantly reduced adherence to human vaginal and endocervical cells *in vitro* (Fig. 3A, B). To determine if Ebp is important for *in vivo* vaginal colonization and persistence, mice were inoculated with either WT OG1SSp (an isogenic derivative of OG1 that is resistant to streptomycin and spectinomycin) or OG1RF Δ*ebpABC* and colonization was quantified over the course of 13 days. We note that both WT strains, OG1RF and OG1SSp, persist in the vaginal lumen and colonize reproductive tract tissues at similar levels (Fig. S1A, B). We observed no differences in the CFU recovered from the vaginal lumen between WT OG1SSp and OG1RF Δ*ebpABC* strains when inoculated together in competition (Fig. 3C). However, in a subsequent experiment, we recovered significantly less of the Δ*ebpABC* mutant in vaginal and cervical tissues and slightly less Δ*ebpABC* in uterine tissue compared to the WT strain 4 days post-inoculation (Fig. 3D, E). Taken together these data suggest that Ebp contributes to *E. faecalis* attachment to reproductive tract tissues, but additional factors are likely required for persistence in the vaginal lumen.

Identification of additional vaginal colonization factors by transposon mutagenesis analysis.

Although the Ebp pilus mutant showed reduced association with reproductive tract tissues, we observed no significant difference in the ability of this mutant to persist in the mouse vagina (Fig. 3C). Thus, we hypothesized that additional bacterial factors are involved in promoting *E. faecalis* vaginal colonization. To identify genetic determinants that confer vaginal persistence, we used <u>sequence-</u>

defined *mariner* technology transposon sequencing (SMarT TnSeq) to screen an *E. faecalis* OG1RF Tn

library consisting of 6,829 unique mutants [25]. The library was grown to mid-log phase in triplicate and

10^7 CFU of each replicate was vaginally inoculated into a group of 5 C57BL/6 mice (Fig. 4A). Vaginal

swabs were plated on selective media daily for 8 days and CFU were quantified to assess colonization of

the OG1RF Tn library compared to WT OG1RF (Fig. 4A and B). Genomic DNA was isolated from pooled

Tn libraries recovered on days 1, 5, and 8 post-inoculation and Tn insertions in *E. faecalis* genomic DNA

were sequenced as described by Dale *et al.* [25]. Sequenced reads were mapped to the *E. faecalis* OG1RF

genome to identify genes that are necessary for *E. faecalis* vaginal colonization.

We observed that the *in vivo* vaginal environment altered the abundance of select mutants from the Tn

library pool compared to the original culture input (Fig. 4C, D, E) (Suppl. Tables 1 - 6). At day 1, a total

of 667 depleted mutants were identified, along with 544 and 507 at days 5 and 8 respectively; 383 of these

mutants were identified at all three time points (Fig. 5A). Classification by clusters of orthologous groups

of proteins (COGs) could be identified in 196 mutants from all 3 time points, the majority of which are

involved in carbohydrate, amino acid, lipid and nucleotide transport/metabolism, as well as those involved

in transcription and defense mechanisms (Fig. 5B). Interestingly the *ebpC* transposon mutant

(*OG1RF_10871*, which encodes the shaft component of Ebp) was underrepresented at all time points (Fig.

4C, D, E, Suppl Table 1). Additional mutants of interest included those involved in ethanolamine

catabolism as well as T7SS components.

Potential gain-of-function mutations have also been discovered during genome-wide library screens of

fitness determinants in other bacteria [26-30]. In addition to mutations that adversely impact vaginal

colonization, our data shows that Tn insertions in 11 protein coding genes and 11 intergenic regions

potentially enhance bacterial fitness *in vivo*. Nineteen of the 22 enriched mutants were common between

days 1 and 5 whereas the other 3 were unique to day 8 (Suppl Table 5 and 6). Since the majority of the

mutants with increased fitness encode hypothetical proteins, the relationship between these genes and

vaginal colonization is currently unclear and requires further investigation.

Ethanolamine utilization and T7SS genes contribute to *E. faecalis* persistence in the reproductive

tract.

Our TnSeq analysis revealed many potential mutants that exhibited decreased colonization in the murine

vaginal tract. We sought to confirm these results by analyzing mutants from systems with multiple

affected genes. One significantly affected operon was ethanolamine (EA) utilization (*eut*) which consists

of 19 genes in *E. faecalis* [31]. Mutations in 4 *eut* genes were significantly underrepresented *in vivo*

compared to the culture input at all time points. These included transposon mutants of the genes encoding

both subunits for ethanolamine ammonia lyase, *eutB (OG1RF_11344)* and *eutC (OG1RF_11343)* [32], a

carboxysome structural protein, *eutL (OG1RF_11342)*[33], and the response regulator, *eutV*

(OG1RF_11347), of the two-component system involved in the regulation of EA utilization [34] (Suppl

Table 1). To assess the importance of EA utilization on *E. faecalis* vaginal colonization, we co-colonized

mice with *E. fae*calis OG1SSp and an OG1RF Δ*eutBC* mutant [35]. Compared to the WT OG1SSp strain,

the Δ*eutBC* mutant was cleared significantly faster from the mouse vagina as seen by the mean CFU

recovered and competitive index (CI) over time (Fig. 6A, B, C). These results suggest that the utilization

of ethanolamine is important for enterococcal persistence in the vaginal tract.

We also observed that Tn insertion mutants in the T7SS locus were significantly underrepresented at all

time points *in vivo* compared to the culture input. The T7SS has been shown to play an important role in

virulence in multiple bacterial species such as *Staphylococcus*, *Listeria* and *Bacillus* [36-38]. In *E.*

faecalis, genes in the T7SS locus have been shown to be induced during phage infection [39]. We observed

that transposon mutants for *esaB* (*OG1RF_11103)*, a putative cytoplasmic accessory protein;

OG1RF_11109 and *OG1RF_11111*, putative toxin effector proteins; *OG1RF_11113*, a putative toxin,

OG1RF_11114, a transmembrane protein; and *OG1RF_11122,* a potential antitoxin protein were all

significantly underrepresented (Suppl Table 1). To confirm the role of the T7SS in vaginal colonization,

we utilized an *esaB* (*esaB*-Tn) mutant, in which *esaB* is disrupted by a transposon and thus Tn-mediated

polar effects may exist for this strain. Following co-colonization with *E. faecalis* OG1RF and the OG1RF

esaB-Tn mutant strain, we observed that while there were no differences in initial colonization, fewer

esaB-Tn mutant bacteria were recovered from the vaginal lumen at later time points (Fig. 6D). We

repeated this experiment and harvested tissues at an earlier time point, before there were any colonization

differences observed between the two strains (day 11). Here we observed that WT OG1RF outcompeted

the *esaB*-Tn mutant strain and was better able to access reproductive tract tissues (Fig. 6E, F), indicating

that the T7SS may be involved in vaginal persistence and ascension in the female reproductive tract.

DISCUSSION

E. faecalis is associated with a wide spectrum of infections, particularly under immunocompromised states

and during compositional shifts in the host microbiota [40, 41]. Although an increasing body of evidence

links enterococci with bacterial vaginosis (BV) and aerobic vaginitis (AV) [17, 18, 42-44], the molecular

determinants that facilitate *E. faecalis* colonization and persistence in the vaginal tract are largely

unknown. Here, we established *in vitro* and *in vivo* systems to acquire genome-scale interactions that

confer *E. faecalis* fitness within the female reproductive tract. We show that both vancomycin sensitive

enterococci (VSE) and VRE adhere to cell types of vaginal and cervical origin, a signature of bacterial

colonization that precedes tissue invasion and systemic spread. Further, genetic features involved in

biofilm formation, ethanolamine utilization and polymicrobial interactions influence *E. faecalis* vaginal

carriage.

Previous studies have demonstrated the importance of Ebp pili in enterococcal virulence and biofilm

formation [14]. We found that deletion of *ebpABC* attenuated binding to human vaginal and endocervical

cells but did not influence bacterial burden in the vaginal lumen, similar to the observed function of Ebp

pili in the intestine[45]. Enhanced *E. faecalis* adherence in tissue culture compared to *in vivo* colonization

may reflect the lack of liquid and mucus flow that bacteria encounter within the vaginal tract, emphasizing

the significance of our animal model for to investigating *E. faecalis*-vaginal interactions. Consistent with

this observation, an *in vivo* Tn library screen revealed only two underrepresented biofilm-associated

mutants, *ebpC-Tn* and *OG1RF_10506-Tn*, encoding a putative polysaccharide deacetylase homolog

implicated in low biofilm formation *in vitro* [46, 47]. Together, these results show that individual

mutations in *ebp* or other well characterized biofilm genes are not sufficient to impair vaginal niche

establishment and/or persistence of enterococci, which likely depends on the concerted effort of multiple

factors.

Transition from nutrient rich laboratory media to the vaginal tract likely imparts dramatic alterations in *E.*

faecalis metabolism. In support of this hypothesis, our high-throughput Tn mutant screen showed that

mutations in carbohydrate, amino acid and nucleotide metabolic pathways were indispensable in the

vaginal tract. Specifically, we showed that WT bacteria outcompete a *eut* locus mutant during vaginal

colonization. In contrast, Kaval and colleagues demonstrated that mutations in *eut* genes leads to a slight

increase in fitness within the intestine [48]. This observation likely reflects the metabolic requirements of

different host environments. While a number of reports exist on the contributions of EA catabolism in

host-bacteria interactions within the intestine [49], studies are lacking for the relevance of this EA

metabolism in other host-associated environments. Our results raise important questions regarding EA

utilization in the female reproductive tract. Although commensal microbes and the epithelium are rich

sources of EA, the composition and source of EA in the vaginal tract remains unknown. A recent report

showed that *E. faecalis* EA utilization attenuates intestinal colitis in IL10 knockout mice in the presence

of a defined microbiota [50]. Whether EA utilization promotes virulence or commensalism for enterococci

in the context of vaginal tissue remains to be determined. Considering that the by-product of EA

metabolism, acetate, is anti-inflammatory and promotes IgA production in the intestine [51, 52], it is

intriguing to consider that enterococcal EA catabolism might modulate immune responses within the

female reproductive tract.

T7SSs have been implicated in the maintenance of bacterial membrane integrity, virulence, and

interbacterial antagonism [36, 53-59]. In *S. aureus*, T7SS encoded proteins confer protection from

membrane damage caused by host fatty acids [58, 59]. Although *E. faecalis* T7SS genes were shown to

be induced in response to phage driven membrane damage, direct contributions of these genes in cell

envelop barrier function and/or virulence in the context of animal models are poorly defined. Our TnSeq

analysis revealed that insertional mutations in six T7SS genes diminished early and late vaginal

colonization by *E. faecalis* OG1RF. In vaginal co-colonization competition experiments, an *esaB-Tn* strain

reached WT colonization levels early and showed a colonization defect in long term persistence. The

incongruence in the colonization kinetics of T7SS mutant strains compared to T7SS-Tn library mutants

which were observed early after inoculation, presumably stem from the inherent differences in the vaginal

milieu in these two experiments. The TnSeq library employed in this screen is a complex population of

approximately 7,000 unique mutants, but it is still very likely that direct or indirect interactions between

mutants influences fitness. LXG–domain toxins, which are part of the T7SS, have been shown to

antagonize neighboring non-kin bacteria [56]. The fact that two mutants with LXG–domain encoding gene

mutations, *OG1RF_11109* and *OG1RF_11111*, were underrepresented across all time points during

vaginal colonization suggests that these putative antibacterial proteins may influence enterococcal

interactions with the resident microbes of the vaginal tract.

Here, we report the development of a novel mouse model for investigating host-enterococcal interactions

in the vaginal tract. This will be a useful model for analyzing the bacterial and host factors that govern

enterococcal vaginal colonization, as well as characterizing the polymicrobial interactions that may

contribute to *E. faecalis* niche establishment and persistence. Transposon library screening of *E.*

faecalis recovered from the mouse vagina has revealed new insights into our understanding of

enterococcal vaginal carriage. Our results emphasize the importance of pili, ethanolamine utilization, and

T7SS components for successful *E. faecalis* colonization of the female reproductive tract, highlighting the

complex nature of this niche. Continued studies will further elucidate the molecular mechanisms of

vaginal persistence, and may represent an important reservoir for antibiotic resistant enterococci.

METHODS

Bacterial strains and culture conditions.

A detailed list of bacterial strains can be found in Suppl Table 7. *E. faecalis* strains V583 [60] and OG1-

(RF and SSp) [20, 61] were used for these experiments. *E. faecalis* was grown in brain heart infusion (BHI

[62]) broth at 37℃ with aeration and growth was monitored by measuring the optical density at 600nm

(OD_{600}). For selection of *E. faecalis* V583, BHI agar was supplemented with gentamicin (100 µg/ml). For

selection of *E. faecalis* OG1RF, OG1RF Δ*ebpABC* [24] and OG1RF Δ*eutBC* [35], BHI agar was

supplemented with rifampicin (50 μg/ml) and fusidic acid (25 μg/ml). For selection of *E. faecalis*

OG1SSp, BHI agar was supplemented with streptomycin (150 μg/ml) and spectinomycin (100 μg/ml). *E.*

faecalis OG1RF *esaB*-Tn [25] was grown on BHI agar supplemented with rifampicin (50 μg/ml), fusidic

acid (25 μg/L), and chloramphenicol (15 μg/ml).

***In vitro* adherence assays**

Immortalized VK2 human vaginal epithelial cells and End1 human endocervical epithelial cells were

obtained from the American Type Culture Collection (VK2.E6E7, ATCC CRL-2616 and End1/E6E7,

ATCC CRL-2615) and were maintained in keratinocyte serum-free medium (KSFM; Gibco) with 0.1

ng/mL human recombinant epidermal growth factor (EGF; Gibco) and 0.05 mg/ml bovine pituitary extract

(Gibco) at 37°C with 5% CO2. Assays to determine cell surface-adherent *E. faecalis* were performed as

described previously when quantifying GBS adherence [63]. Briefly, bacteria were grown to mid-log

phase (OD_{600} = 0.4 - 0.6) and added to cell monolayers (multiplicity of infection [MOI] = 1). After a 30-

min incubation, cells were washed with PBS three times following detachment with 0.1 mL of 0.25%

trypsin-EDTA solution and lysed with addition of 0.4 mL of 0.025% TritonX-100 in PBS by vigorous

pipetting. The lysates were then serially diluted and plated on Todd Hewitt agar (THA) to enumerate

the bacterial CFU. Experiments were performed at least three times with each condition in triplicate, and

results from a representative experiment are shown.

Murine vaginal colonization model.

Animal experiments were approved by the Institutional Animal Care and Use Committee at University of

Colorado-Anschutz Medical Campus protocols #00316 and #00253 and performed using accepted

veterinary standards. A mouse vaginal colonization model for GBS was adapted for our studies [21]. 8-

week old female CD1 (Charles River) or C57BL/6 (Jackson) mice were injected intraperitoneally with 0.5

mg 17β-estradiol (Sigma) 1 day prior to colonization with *E. faecalis*. Mice were vaginally inoculated

with 10^7 CFU of *E. faecalis* in 10μL PBS and on subsequent days the vaginal lumen was swabbed with a

sterile ultrafine swab (Puritan). For co-colonization, mice were inoculated with *E. faecalis* OG1SSp,

OG1RF or deletion mutants in the OG1RF background. To assess tissue CFU, mice were euthanized

according to approved veterinary protocols and the female reproductive tract tissues were dissected and

placed into 500μL PBS and bead beat for 2 min to homogenize the tissues. The resulting homogenate was

serially diluted and *E. faecalis* CFU enumerated on BHI agar supplemented with antibiotics to select for

the strain of interest.

Histology.

Mice were inoculated with *E. faecalis* V583 containing a plasmid that expresses *gfp* (pMV158GFP) and

contains resistance to tetracycline (15 μg/mL) [23]. After 1 day post-inoculation, the murine female

reproductive tract was harvested, embedded into Optimal Cutting Temperature (OCT) compound

(Sakura), and sectioned at 7μm with a CM1950 freezing cryostat (Leica). For fluorescence microscopy,

coverslips were mounted with VECTASHIELD mounting medium containing 4',6-diamidino-2-

phenylindole (DAPI, Vector Labs). Images were taken with a BZ-X710 microscope (Keyence) [22].

Transposon mutant library growth and vaginal colonization.

The *E. faecalis* OG1RF transposon mutant library was generated previously [25]. The E. *faecalis* OG1RF

pooled transposon library was inoculated into 5 ml of BHI at a total of 10^8 CFU and grown with aeration

to an OD_{600} of 0.5. The library was inoculated into the vaginal tracts of C57BL/6 mice at 10^7 CFU. The

library was plated on BHI agar to confirm the inoculum for all groups of mice. Mice were swabbed daily

and swabs were plated on BHI supplemented with rifampicin (50 μg/ml), fusidic acid (25 μg/ml) and

chloramphenicol (20 µg/ml) to quantify CFU. On days 1, 5, and 8, undiluted swabs were plated on BHI

agar with antibiotics and grown to a bacterial lawn. Bacteria were scraped and re-suspended in PBS and

pelleted. DNA from days 1, 5, 8, and the input culture was extracted from pellets using a

ZymoBIOMICS™ DNA MiniPrep kit (Zymo Research).

Transposon library sequencing and data analysis.

Transposon-junction DNA library preparation and Illumina NovaSeq 6000 DNA sequencing (150 base

paired-end mode) was performed by the Microarray and Genomics Core at the University of Colorado

Anschutz Medical Campus as previously described [39]. For downstream analysis of transposon-

junctions, we used only the R1 reads generated by paired end sequencing. Illumina adapter trimmed raw

reads were mapped to the *E. faecalis* OG1RF reference sequence (NC_017316.1) and differentially

abundant transposon mutants were identified using statistical analysis scripts established by Dale *et*

al.[25]. The abundance of Tn mutants in culture was compared to input library used for culture inoculation

and mutants that are not significantly different ($p > 0.05$) between these two samples were considered for

the next steps of the analysis. For comparisons between *in vivo* and *in vitro* samples, mutants were

considered significantly different if the adjusted P value was < 0.05 and a $\log_2$ (fold change) > 1. The

DNA sequencing reads associated with this study have been deposited at the European Nucleotide Archive

under accession number PRJEB37929.

Data analysis.

GraphPad Prism version 7.0 was used for statistical analysis and statistical significance was accepted at P

values of < 0.05 ($* P < 0.05$; $** P < 0.005$; $*** P < 0.0005$; $**** P < 0.00005$). Specific tests are indicated

in figure legends.

ACKNOWLEGMENTS

We thank Kimberley Kline for providing *E. faecalis* OG1RF Δ*ebpABC*, Danielle Garsin for providing *E. faecalis* OG1RF Δ*eutBC*, Katrina Diener and Monica Ransom for custom TnSeq library preparation and the Microarray and Genomics Core at the University of Colorado Anschutz Medical Campus for DNA sequencing. This study was supported by the NIH 5T32AI007405-28 to B.L.S., the NIH/NIAID R21 AI130857 to K.S.D and the NIH/NIAID R01 AI141479 to B.A.D, and University of Colorado School of Medicine IMPA funding to K.S.D and B.A.D.

FIGURE LEGENDS

Figure 1: *E. faecalis* colonizes the murine female reproductive tract. (A) Adherence of *E. faecalis* V583 and OG1RF to human vaginal and endocervical cells. Experiments were performed in triplicate and error bars represent standard deviations (SDs); the results of a representative experiment are shown. (B) CFU counts of V583 and OG1RF recovered from vaginal swabs 1 day post-inoculation. (C) CFU counts of V583 from vaginal, cervical and uterine tissue 1 day post-inoculation. (D, E, F) Mice were inoculated with V583 expressing *gfp* and 7μm sections of vaginal (D), cervical (E) and uterine (F) tissue were stained with DAPI for fluorescence microscopy. White arrows indicate green fluorescent bacteria present in tissue sections. Images were all taken at 40x magnification. LP = lamina propria, EP = epithelial layer, Lu = lumen.

Figure 2: *E. faecalis* persists in the murine vaginal tract. (A and B) *E. faecalis* V583 and OG1RF in the murine vaginal tract. C57BL/6 mice were inoculated with 10^7 V583 or OG1RF CFU and the vaginal lumen of each mouse was swabbed daily and swabs were serially diluted and plated on selective media to

quantify CFU. Data are presented as recovered CFU per swab (A) and mean recovered CFU (B). (C and D) CD1 (n = 10) and C57BL/6 (n = 7) mice were inoculated with V583 and the vaginal lumen of each mouse was swabbed daily, serially diluted, and plated on selective media to quantify CFU. Data are presented as recovered CFU per swab (C) and mean recovered CFU (D)

Figure 3: Enterococcal pili contribute to colonization of the female reproductive tract. (A and B) *E. faecalis* OG1RF WT and OG1RF Δ*ebpABC* adherence to human vaginal (A) and endocervical (B) cells. Experiments were performed with four technical replicates and error bars represent SDs; the results of a representative experiment are shown and analyzed using an unpaired t-test; ∗∗P < 0.0001; ∗∗∗∗P < 0.00001. (C) C57BL/6 mice were co-inoculated with OG1SSp WT and OG1RF Δ*ebpABC* and the vaginal lumen was swabbed daily. (D and E) C57BL/6 mice were co-inoculated with OG1SSp WT and OG1RF Δ*ebpABC* and reproductive tissue was collected and CFU/gram was quantified. Data are presented as recovered Log_{10}CFU/gram (D) and competitive index (CI) between OG1SSp WT and OG1RF Δ*ebpABC* (E). CI is enumerated by calculating the ratio of WT to mutant *E. faecalis* recovered from the mouse reproductive tract. A CI >1 indicates an advantage to WT *E. faecalis*. Values below the limit of detection were enumerated as one-half the limit of detection. Data were analyzed by a paired t-test; ∗P < 0.05.

Figure 4: Identification of additional factors required for vaginal colonization and persistence by transposon mutant library screen. (A) Schematic representing experimental approach. C57BL/6 mice were treated with 17β-estradiol prior to inoculation with the OG1RF Tn library in triplicate groups or WT OG1RF. The vaginal lumen was swabbed and CFU was enumerated daily. DNA from recovered OG1RF Tn mutants was sequenced on days 1, 5 and 8. (B) CFU recovered from vaginal swabs of triplicate groups

of mice colonized with OG1RF Tn mutagenesis library (T1, T2, T3) and OG1RF WT (n=5 mice per

group). (C, D and E) Volcano plots depicting underrepresented and overrepresented mutants *in vivo*

compared to culture on day 1 (C), day 5 (D) and day 8 (E) post-inoculation. Colored dots represent mutants

of interest. Pink = *ebpC* Tn mutant, Orange = *eut* Tn mutant, Green = T7SS gene Tn mutants. (F)

Figure 5: Classification of transposon insertion mutants by Cluster of Orthologous Groups (COGs).

(A) Euler plot representing number of underrepresented mutants at all time points. (B) Cluster of

Orthologous Groups (COGs) from underrepresented mutants common to all three time points categorized

into functional categories. Total = 196 represents all common mutants that were assigned a COG domain.

Figure 6: Ethanolamine utilization and type VII secretion system genes contribute to enterococcal

persistence in the vaginal tract. (A, B and C) C57BL/6 mice were co-inoculated with OG1SSp WT and

OG1RF Δ*eutBC* and vaginal lumen was swabbed to quantify CFU. Data are presented as recovered CFU

per swab (A), mean recovered CFU (B) and CI between WT and mutant strains (C). Data was analyzed

used a Two-way ANOVA with multiple comparisons; *$P < 0.05$, **$P < 0.005$, ***$P < 0.00005$. (D)

C57BL/6 mice were co-inoculated with OG1RF WT and OG1RF *esaB*-Tn and vaginal lumen was

swabbed to quantify CFU. Data was analyzed using a Two-way ANOVA with multiple comparisons; **P

< 0.005, ***$P < 0.0005$ (E and F) C57BL/6 mice were co-inoculated with OG1RF WT and OG1RF *esaB*-

Tn and reproductive tissue was collected at 11 days post-inoculation. Data are presented as recovered

Log_{10} CFU/gram (E) and CI between WT and Tn-mutant strain (F). CI is enumerated as described in

Figure 4. Data was analyzed using a paired t-test; *$P < 0.05$, **$P < 0.005$.

Supplemental Figure 1: Colonization and persistence of *E. faecalis* OG1- (RF and SSp). (A and B) C57BL/6 mice were co-inoculated with OG1RF WT and OG1SSp WT and the vaginal lumen was swabbed to quantify CFU. Reproductive tissue was collected at 8 days post-inoculation. Data are presented as recovered CFU per swab (A) and Log_{10} CFU/gram (B). Data was analyzed using a Two-way ANOVA with multiple comparisons; $P > 0.05$ (A) and a paired t-test; $P > 0.05$ (B).

Supplemental Table 1: Common underrepresented *E. faecalis* OG1RF transposon mutants during vaginal colonization day 1, 5 and 8 post-inoculation. All significantly underrepresented Tn mutants common to all time points are listed along with their functional categorization from COGs and log_2 adjusted P values comparing *in vitro* to *in vivo* samples.

Supplemental Table 2: Underrepresented *E. faecalis* OG1RF transposon mutants during vaginal colonization day 1 post-inoculation. All significantly underrepresented Tn mutants from *in vivo* samples 1 day post-inoculation are listed along with their functional categorization from COGs and log_2 adjusted P values comparing *in vitro* to *in vivo* samples.

Supplemental Table 3: Underrepresented *E. faecalis* OG1RF transposon mutants during vaginal colonization day 5 post-inoculation. All significantly underrepresented Tn mutants from *in vivo* samples 5 days post-inoculation are listed along with their functional categorization from COGs and log_2 adjusted P values comparing *in vitro* to *in vivo* samples.

Supplemental Table 4: Underrepresented *E. faecalis* OG1RF transposon mutants during vaginal

colonization day 8 post-inoculation. All significantly underrepresented Tn mutants from *in vivo* samples

8 days post-inoculation are listed along with their functional categorization from COGs and $\log_2$ adjusted

P values comparing *in vitro* to *in vivo* samples.

Supplemental Table 5: Overrepresented *E. faecalis* OG1RF transposon mutants during vaginal

colonization day 1 and 5 post-inoculation. All significantly overrepresented Tn mutants from *in vivo*

samples 1 and 5 days post-inoculation are listed along with their functional categorization from COGs and

$\log_2$ adjusted P values comparing *in vitro* to *in vivo* samples.

Supplemental Table 6: Overrepresented *E. faecalis* OG1RF transposon mutants during vaginal

colonization day 8 post-inoculation. All significantly overrepresented Tn mutants from *in vivo* samples

8 days post-inoculation are listed along with their functional categorization from COGs and $\log_2$ adjusted

P values comparing *in vitro* to *in vivo* samples.

Supplemental Table 7: List of bacterial strains and method of selection used in this study.

REFERENCES:

1.   Lebreton, F., R.J.L. Willems, and M.S. Gilmore, *Enterococcus diversity, origins in nature, and*

gut colonization, in *Enterococci: from commensals to leading causes of drug resistant infection*,

M.S. Gilmore, et al., Editors. 2014: Boston.

2.   Coolen, M.J., et al., *Characterization of microbial communities found in the human vagina by*

analysis of terminal restriction fragment length polymorphisms of 16S rRNA genes. Appl

Environ Microbiol, 2005. **71**(12): p. 8729-37.

3.   Huycke, M.M., D.F. Sahm, and M.S. Gilmore, *Multiple-drug resistant enterococci: the nature of*

the problem and an agenda for the future. Emerg Infect Dis, 1998. **4**(2): p. 239-49.

4.   Schindlbeck, C., D. Dziura, and I. Mylonas, *Diagnosis of pelvic inflammatory disease (PID):*

intra-operative findings and comparison of vaginal and intra-abdominal cultures. Arch Gynecol

Obstet, 2014. **289**(6): p. 1263-9.

5.   Arias, C.A. and B.E. Murray, *Emergence and management of drug-resistant enterococcal*

infections. Expert Rev Anti Infect Ther, 2008. **6**(5): p. 637-55.

6.   Uttley, A.H., et al., *Vancomycin-resistant enterococci.* Lancet, 1988. **1**(8575-6): p. 57-8.

7.   Martin, H.L., et al., *Vaginal lactobacilli, microbial flora, and risk of human immunodeficiency*

virus type 1 and sexually transmitted disease acquisition. J Infect Dis, 1999. **180**(6): p. 1863-8.

8.   Madsen, K.T., et al., *Virulence Factors Associated with Enterococcus Faecalis Infective*

Endocarditis: A Mini Review. Open Microbiol J, 2017. **11**: p. 1-11.

9.   Thurlow, L.R., et al., *Gelatinase contributes to the pathogenesis of endocarditis caused by*

Enterococcus faecalis. Infect Immun, 2010. **78**(11): p. 4936-43.

10.   Ali, L., et al., *Molecular Mechanism of Quorum-Sensing in Enterococcus faecalis: Its Role in*

Virulence and Therapeutic Approaches. Int J Mol Sci, 2017. **18**(5).

11.   Waters, C.M., et al., *Role of the Enterococcus faecalis GelE protease in determination of cellular*

chain length, supernatant pheromone levels, and degradation of fibrin and misfolded surface

proteins. J Bacteriol, 2003. **185**(12): p. 3613-23.

12.   Qin, X., et al., *Effect of disruption of a gene encoding an autolysin of Enterococcus faecalis*
503   *OG1RF.* Antimicrob Agents Chemother, 1998. **42**(11): p. 2883-8.

504   13.   Tacconelli, E. and M.A. Cataldo, *Vancomycin-resistant enterococci (VRE): transmission and*
505   *control.* Int J Antimicrob Agents, 2008. **31**(2): p. 99-106.

506   14.   Ch'ng, J.H., et al., *Biofilm-associated infection by enterococci.* Nat Rev Microbiol, 2019. **17**(2):
507   p. 82-94.

508   15.   Leyva-Gomez, G., et al., *Modifications in Vaginal Microbiota and Their Influence on Drug*
509   *Release: Challenges and Opportunities.* Pharmaceutics, 2019. **11**(5).

510   16.   Ravel, J., et al., *Vaginal microbiome of reproductive-age women.* Proc Natl Acad Sci U S A,
511   2011. **108 Suppl 1**: p. 4680-7.

512   17.   Donders, G.G.G., et al., *Aerobic vaginitis: no longer a stranger.* Res Microbiol, 2017. **168**(9-10):
513   p. 845-858.

514   18.   Kaambo, E., et al., *Vaginal Microbiomes Associated With Aerobic Vaginitis and Bacterial*
515   *Vaginosis.* Front Public Health, 2018. **6**: p. 78.

516   19.   Sahm, D.F., et al., *In vitro susceptibility studies of vancomycin-resistant Enterococcus faecalis.*
517   Antimicrob Agents Chemother, 1989. **33**(9): p. 1588-91.

518   20.   Gold, O.G., H.V. Jordan, and J. van Houte, *The prevalence of enterococci in the human mouth*
519   *and their pathogenicity in animal models.* Arch Oral Biol, 1975. **20**(7): p. 473-7.

520   21.   Patras, K.A. and K.S. Doran, *A Murine Model of Group B Streptococcus Vaginal Colonization.* J
521   Vis Exp, 2016(117).

522   22.   Deng, L., et al., *Identification of Key Determinants of Staphylococcus aureus Vaginal*
523   *Colonization.* mBio, 2019. **10**(6).

23. Nieto, C. and M. Espinosa, *Construction of the mobilizable plasmid pMV158GFP, a derivative*

of pMV158 that carries the gene encoding the green fluorescent protein. Plasmid, 2003. **49**(3): p.

281-5.

24. Afonina, I., et al., *Planktonic Interference and Biofilm Alliance between Aggregation Substance*

and Endocarditis- and Biofilm-Associated Pili in Enterococcus faecalis. J Bacteriol, 2018.

200(24).

25. Dale, J.L., et al., *Comprehensive functional analysis of the Enterococcus faecalis core genome*

using an ordered, sequence-defined collection of insertional mutations in strain OG1RF.

mSystems, 2018. **3**(5).

26. Bachman, M.A., et al., *Genome-Wide Identification of Klebsiella pneumoniae Fitness Genes*

during Lung Infection. mBio, 2015. **6**(3): p. e00775.

27. Jung, H.J., et al., *Genome-Wide Screening for Enteric Colonization Factors in Carbapenem-*

Resistant ST258 Klebsiella pneumoniae. mBio, 2019. **10**(2).

28. Shames, S.R., et al., *Multiple Legionella pneumophila effector virulence phenotypes revealed*

through high-throughput analysis of targeted mutant libraries. Proc Natl Acad Sci U S A, 2017.

114(48): p. E10446-E10454.

29. Skurnik, D., et al., *A comprehensive analysis of in vitro and in vivo genetic fitness of*

Pseudomonas aeruginosa using high-throughput sequencing of transposon libraries. PLoS

Pathog, 2013. **9**(9): p. e1003582.

30. Wiles, T.J., et al., *Combining quantitative genetic footprinting and trait enrichment analysis to*

identify fitness determinants of a bacterial pathogen. PLoS Genet, 2013. **9**(8): p. e1003716.

31. Del Papa, M.F. and M. Perego, *Ethanolamine activates a sensor histidine kinase regulating its*

utilization in Enterococcus faecalis. J Bacteriol, 2008. **190**(21): p. 7147-56.

32.   Roof, D.M. and J.R. Roth, *Ethanolamine utilization in Salmonella typhimurium.* J Bacteriol,

548   1988. **170**(9): p. 3855-63.

549   33.   Thompson, M.C., et al., *Structure of a bacterial microcompartment shell protein bound to a*

550   *cobalamin cofactor.* Acta Crystallogr F Struct Biol Commun, 2014. **70**(Pt 12): p. 1584-90.

551   34.   Fox, K.A., et al., *Multiple posttranscriptional regulatory mechanisms partner to control*

552   *ethanolamine utilization in Enterococcus faecalis.* Proc Natl Acad Sci U S A, 2009. **106**(11): p.

553   4435-40.

554   35.   Kaval, K.G., et al., *Ethanolamine Utilization and Bacterial Microcompartment Formation Are*

555   *Subject to Carbon Catabolite Repression.* J Bacteriol, 2019. **201**(10).

556   36.   Cao, Z., et al., *The type VII secretion system of Staphylococcus aureus secretes a nuclease toxin*

557   *that targets competitor bacteria.* Nat Microbiol, 2016. **2**: p. 16183.

558   37.   Huppert, L.A., et al., *The ESX system in Bacillus subtilis mediates protein secretion.* PLoS One,

559   2014. **9**(5): p. e96267.

560   38.   Pinheiro, J., et al., *Listeria monocytogenes encodes a functional ESX-1 secretion system whose*

561   *expression is detrimental to in vivo infection.* Virulence, 2017. **8**(6): p. 993-1004.

562   39.   Chatterjee, A., et al., *Parallel genomics uncover novel enterococcal-bacteriophage interactions.*

563   mBio, 2020. **11**(2).

564   40.   Brandl, K., et al., *Vancomycin-resistant enterococci exploit antibiotic-induced innate immune*

565   *deficits.* Nature, 2008. **455**(7214): p. 804-7.

566   41.   Ubeda, C., et al., *Vancomycin-resistant Enterococcus domination of intestinal microbiota is*

567   *enabled by antibiotic treatment in mice and precedes bloodstream invasion in humans.* J Clin

568   Invest, 2010. **120**(12): p. 4332-41.

42.   Dezwaan, D.C., et al., *Purification and characterization of enterocin 62-6, a two-peptide*

bacteriocin produced by a vaginal strain of Enterococcus faecium: Potential significance in

bacterial vaginosis. Microb Ecol Health Dis, 2007. **19**(4): p. 241-250.

43.   Kelly, M.C., M.J. Mequio, and V. Pybus, *Inhibition of vaginal lactobacilli by a bacteriocin-like*

inhibitor produced by Enterococcus faecium 62-6: potential significance for bacterial vaginosis.

Infect Dis Obstet Gynecol, 2003. **11**(3): p. 147-56.

44.   Tempera, G., et al., *Topical kanamycin: an effective therapeutic option in aerobic vaginitis.* J

Chemother, 2006. **18**(4): p. 409-14.

45.   Banla, L.I., et al., *Sortase-Dependent Proteins Promote Gastrointestinal Colonization by*

Enterococci. Infect Immun, 2019. **87**(5).

46.   Willett, J.L.E., M.M. Ji, and G.M. Dunny, *Exploiting biofilm phenotypes for functional*

characterization of hypothetical genes in Enterococcus faecalis. NPJ Biofilms Microbiomes,

2019. **5**: p. 23.

47.   Balomenou, S., et al., *Distinct functions of polysaccharide deacetylases in cell shape, neutral*

polysaccharide synthesis and virulence of Bacillus anthracis. Mol Microbiol, 2013. **87**(4): p.

867-83.

48.   Kaval, K.G., et al., *Loss of Ethanolamine Utilization in Enterococcus faecalis Increases*

Gastrointestinal Tract Colonization. mBio, 2018. **9**(3).

49.   Kaval, K.G. and D.A. Garsin, *Ethanolamine Utilization in Bacteria.* mBio, 2018. **9**(1).

50.   Lengfelder, I., et al., *Complex Bacterial Consortia Reprogram the Colitogenic Activity of*

Enterococcus faecalis in a Gnotobiotic Mouse Model of Chronic, Immune-Mediated Colitis.

Front Immunol, 2019. **10**: p. 1420.

51.   Maslowski, K.M., et al., *Regulation of inflammatory responses by gut microbiota and*

chemoattractant receptor GPR43. Nature, 2009. **461**(7268): p. 1282-6.

52.   Wu, W., et al., *Microbiota metabolite short-chain fatty acid acetate promotes intestinal IgA*

response to microbiota which is mediated by GPR43. Mucosal Immunol, 2017. **10**(4): p. 946-

956.

53.   Burts, M.L., et al., *EsxA and EsxB are secreted by an ESAT-6-like system that is required for the*

pathogenesis of Staphylococcus aureus infections. Proc Natl Acad Sci U S A, 2005. **102**(4): p.

1169-74.

54.   Kneuper, H., et al., *Heterogeneity in ess transcriptional organization and variable contribution*

of the Ess/Type VII protein secretion system to virulence across closely related Staphyloccus

aureus strains. Mol Microbiol, 2014. **93**(5): p. 928-43.

55.   Ohr, R.J., et al., *EssD, a nuclease effector of the Staphylococcus aureus ESS Pathway.* J

Bacteriol, 2017. **199**(1).

56.   Whitney, J.C., et al., *A broadly distributed toxin family mediates contact-dependent antagonism*

between Gram-positive bacteria. Elife, 2017. **6**.

57.   Ishii, K., et al., *Induction of virulence gene expression in Staphylococcus aureus by pulmonary*

surfactant. Infect Immun, 2014. **82**(4): p. 1500-10.

58.   Lopez, M.S., et al., *Host-derived fatty acids activate type VII secretion in Staphylococcus aureus.*

Proc Natl Acad Sci U S A, 2017. **114**(42): p. 11223-11228.

59.   Tchoupa, A.K., et al., *The type VII secretion system protects Staphylococcus aureus against*

antimicrobial host fatty acids. bioRxiv, 2020: p. 572172.

60.   Paulsen, I.T., et al., *Role of mobile DNA in the evolution of vancomycin-resistant Enterococcus*

faecalis. Science, 2003. **299**(5615): p. 2071-4.

61.   Bourgogne, A., et al., *Large scale variation in Enterococcus faecalis illustrated by the genome*

analysis of strain OG1RF. Genome Biol, 2008. **9**(7): p. R110.

62.   Morgan, D.J., et al., *Comparison of Gram-stained smears prepared from blind vaginal swabs*

with those obtained at speculum examination for the assessment of vaginal flora. Br J Obstet

Gynaecol, 1996. **103**(11): p. 1105-8.

63.   Deng, L., et al., *Characterization of a Two-Component System Transcriptional Regulator, LtdR,*

That Impacts Group B Streptococcal Colonization and Disease. Infect Immun, 2018. **86**(7).

A. *E. faecalis* Adherence *in vitro*

% Adherence (y-axis: 0, 10, 20, 30, 40)

Legend: V583 (filled), OG1RF (open)

x-axis: Vaginal, Endocervical

B. *E. faecalis* Vaginal Colonization

Recovered CFU (Log$_{10}$) (y-axis: 0, 2, 4, 6, 8, 10)

x-axis: V583, OG1RF

C. V583 Colonization

Recovered CFU/g (Log$_{10}$) (y-axis: 0, 2, 4, 6, 8, 10)

x-axis: Vagina, Cervix, Uterus

D. LP, Lu — vagina (40x)

E. EP, Lu, LP — cervix (40x)

F. LP, Lu, EP — uterus (40x)

A. Vaginal Persistence (Recovered CFU/Mouse)

Recovered CFU (Log$_{10}$) vs Days Post-Inoculation

● V583   ○ OG1RF

B. Vaginal Persistence (Mean CFU)

Mean recovered CFU (log$_{10}$) vs Days Post-Inoculation

● V583   ○ OG1RF

C. V583 Vaginal Persistence (Recovered CFU/Mouse)

Recovered CFU (Log$_{10}$) vs Days Post-Inoculation

○ CD1   ● C57BL/6

D. V583 Vaginal Persistence (Mean CFU)

Mean recovered CFU (log$_{10}$) vs Days Post-Inoculation

○ CD1   ● C57BL/6

A.

D8
(507)

D1
(667)

D5
(544)

🟩 Day 1	🟪 Day 1 and 5
🟦 Day 5	🟨 Day 1 and 8
🟥 Day 8	🔵 Day 5 and 8
	🫒 Day 1, 5 and 8

B.

Function unknown

General function prediction only

Secondary metabolites biosynthesis, transport and catabolism

Inorganic ion transport and metabolism

Lipid transport and metabolism

Coenzyme transport and metabolism

Nucleotide transport and metabolism

Amino acid transport and metabolism

Carbohydrate transport and metabolism

Energy production and conversion

Cell motility

Cell wall/membrane/envelope biogenesis

Signal transduction mechanisms

Defense mechanisms

Cell cycle control, cell division, chromosome partitioning

Replication, recombination and repair

Transcription

Translation, ribosomal structure and biogenesis

0   10   20   30

Number of Tn mutants

A. Vaginal Persistence (Recovered CFU/Mouse)

B. Vaginal Persistence (Mean CFU)

C. Competitive Index

● OG1SSp WT   ○ OG1RF Δ*eutBC*

D. Vaginal Persistence (Recovered CFU/Mouse)

E. Competition (Recovered CFU/Mouse)

F. Competitive Index

● OG1RF WT   ○ OG1RF *esaB*-Tn

A. Vaginal Persistence

B. Colonization

OG1RF   OG1SSp