Genome-wide mutagenesis identifies factors involved in *Enterococcus faecalis* vaginal adherence

and persistence.

Norhan Alhajjar[1, #], Anushila Chatterjee[1, #], Brady L. Spencer[1], Lindsey R. Burcham[1], Julia L. E.

Willett[2], Gary M. Dunny[2], Breck A. Duerkop[1*], and Kelly S. Doran[1*].

[1]Department of Immunology and Microbiology, University of Colorado School of Medicine, Anschutz

Medical Campus, Aurora, CO, USA.

[2]Department of Microbiology and Immunology, University of Minnesota Medical School, Minneapolis,

MN, USA.

[#]Designates equal contribution - author order chosen alphabetically

*Corresponding authors: Breck A. Duerkop and Kelly S. Doran, Department of Immunology and

Microbiology, University of Colorado School of Medicine, 12800 East 19th Avenue Aurora, CO, USA.

Phone: (303) 724-3539, Fax: (303) 724-4226, E-mail: breck.duerkop@cuanschutz.edu,

kelly.doran@cuanschutz.edu

RUNNING TITLE: *Enterococcus faecalis* **Vaginal Colonization**

ABSTRACT

Enterococcus faecalis is a Gram-positive commensal bacterium native to the gastrointestinal tract and an opportunistic pathogen of increasing clinical concern. *E. faecalis* also colonizes the female reproductive tract and reports suggest vaginal colonization increases following antibiotic treatment or in patients with aerobic vaginitis. Currently, little is known about specific factors that promote *E. faecalis* vaginal colonization and subsequent infection. We modified an established mouse vaginal colonization model to explore *E. faecalis* vaginal carriage and demonstrate that both vancomycin resistant and sensitive strains colonize the murine vaginal tract. Following vaginal colonization, we observed *E. faecalis* in vaginal, cervical and uterine tissue. A mutant lacking endocarditis- and biofilm-associated pili (Ebp) exhibited a decreased ability to associate with human vaginal and cervical cells *in vitro*, but did not contribute to colonization *in vivo*. Thus, we screened a low-complexity transposon (Tn) mutant library to identify novel genes important for *E. faecalis* colonization and persistence in the vaginal tract. This screen revealed 383 mutants that were underrepresented during vaginal colonization at 1, 5 and 8 days post-inoculation compared to growth in culture medium. We confirmed that mutants deficient in ethanolamine catabolism or in the type VII secretion system were attenuated in persisting during vaginal colonization. These results reveal the complex nature of vaginal colonization and suggest that multiple factors contribute to *E. faecalis* persistence in the reproductive tract.

IMPORTANCE

Despite increasing prevalence and association of *E. faecalis* with aerobic vaginitis, essentially nothing is known about the bacterial factors that influence *E. faecalis* vaginal colonization. We have adapted an animal model of vaginal colonization that supports colonization of multiple *E. faecalis* strains. Additionally, we determined that ethanolamine utilization and type VII secretion system genes

contribute to vaginal colonization and persistence. Identification of factors important for vaginal

colonization and persistence provides potential targets for the development of therapeutics. This study is

the first to identify key determinants that promote vaginal colonization by *E. faecalis*, which may

represent an important reservoir for antibiotic resistant enterococci.

INTRODUCTION

Enterococcus faecalis is an opportunistic pathogen that resides in the human gastrointestinal and

urogenital tracts (1, 2). While *E. faecalis* colonization is normally asymptomatic, certain populations are

at risk for severe disease including urinary tract infections (3), wound infections, pelvic inflammatory

disease (PID), infective endocarditis, and adverse birth effects during pregnancy (reviewed in 4,

5). Enterococcal infections are often associated with the production of biofilms, assemblages of

microbes enclosed in an extracellular polymeric matrix that exhibit cell-to-cell interactions (reviewed in

6). These biofilms have been observed on catheters, diabetic ulcers, and wounds resulting in severe

infection. Treatment of enterococcal infections is becoming increasingly problematic due to their

augmented ability to acquire mobile genetic elements, resulting in increased resistance to antibiotics,

including "last-line-of-defense" antibiotics such as vancomycin (reviewed in 7, 8). Recently, there has

been an increase in the emergence of vancomycin resistant enterococci (9), putting

immunocompromised individuals at risk for developing severe chronic enterococcal infections. The

emergence of vancomycin resistant enterococci (10) and its prevalence in both community and

nosocomial settings is concerning and necessitates the development of alternative therapeutics to treat

enterococcal infections.

E. faecalis encodes a multitude of virulence factors that allow the bacterium to colonize and persist in

different sites of the human body. Surface proteins such as the adhesin to collagen (Ace), enterococcal

fibronectin binding protein A (EfbA), aggregation substance (AS), and the endocarditis-and biofilm-

associated pilin (Ebp) have been previously shown to play important roles in infective endocarditis and

UTIs (reviewed in 11). Secreted factors such as gelatinase (12), autolysin A (13), and serine protease

(SprE) are biofilm-associated factors that are involved in the degradation of host substrates, including

collagen, fibrin and certain complement components (14). Many of these virulence factors are regulated

via quorum sensing, which may be responsible for the switch from a commensal to pathogenic state (15-

17).

Certain risk factors are associated with the transition of *E. faecalis* from commensalism to pathogenicity

such as immune   status,   prolonged   hospital   stay, and   the   use   of   antibiotics   (18). *E.*

faecalis colonization and infection is often polymicrobial and these interactions have been observed in

the intestine, bloodstream, and wounds (reviewed in 19). Furthermore, *E. faecalis* is frequently found in

the vaginal tract of healthy women (20, 21) and its prevalence is increased in women diagnosed

with aerobic   vaginitis   (AV),   an   inflammatory   response   accompanied   by   depletion   of

commensal *Lactobacillus sp.* and increased presence of opportunistic pathogens such as *E. faecalis*,

Group B *Streptococcus* (GBS), *Staphylococcus aureus*, and *Escherichia coli* (22, 23). Symptoms of

AV include malodor and discomfort, but AV can transition to more serious complications such as PID,

severe UTIs, and complications during pregnancy. While it is evident that *E. faecalis* colonizes the

human vaginal tract, the molecular determinants that allow enterococci to colonize and persist in the

vaginal tract remain to be identified.

In this study, we modified our previously established GBS vaginal colonization model to analyze *E.*

faecalis vaginal colonization and persistence. We determined that *E. faecalis* OG1RF (a rifampicin and

fusidic acid derivative of strain OG1) and vancomycin resistant *E. faecalis* V583 can colonize and

persist in the vaginal tract of CD1 and C57BL/6 mouse strains. We detected fluorescent *E. faecalis* in

the vaginal lumen as well as the cervical and uterine tissues of colonized mice. Further, we demonstrated

that an *E. faecalis* strain lacking Ebp pili is less adherent to vaginal cervical epithelium *in vitro,* but not

attenuated *in vivo*. Thus, we screened an *E. faecalis* OG1RF transposon (Tn) mutant library for mutants

that are underrepresented in the vaginal tract compared to the culture input, revealing multiple factors

for *E. faecalis* persistence within the vagina. These factors include sortase-dependent proteins (SDPs),

ethanolamine utilization genes, and genes involved in type VII secretion system (T7SS) machinery. We

confirmed that a mutant strain in ethanolamine catabolism was significantly attenuated in the ability to

colonize the vaginal epithelium, and T7SS was required to ascend in the female reproductive tract. This

work is an important first step in identifying factors required for enterococcal vaginal colonization and

will provide insight into potential therapeutic targets aimed at mitigating *E. faecalis* vaginal colonization

in at-risk individuals.

RESULTS

E. faecalis **colonization of the female reproductive tract.**

To characterize the ability of *E. faecalis* to interact with the epithelial cells of the lower female

reproductive tract, we performed *in vitro* quantitative adherence assays using *E. faecalis* strains V583

(24) and OG1RF (25). An inoculum of 10^5 CFU/well (multiplicity of infection [MOI] = 1) was added to

a confluent monolayer of immortalized human vaginal and endocervical epithelial cells. Following 30

minutes of incubation, the cells were washed to remove non-adherent bacteria, the epithelial cells were

detached from the plates, and adherent bacteria were plated on agar. Both strains exhibited substantial

adherence to both cell lines (Fig. 1A). Next, we assessed the ability of *E. faecalis* to establish

colonization of the murine vaginal tract. The vaginal lumen of C57BL/6 were swabbed and swabs were

plated on CHROM^{TM} agar to determine the presence of native enterococci. While native enterococci are detected on CHROM^{TM} agar, no mice were colonized with strains that resemble *E. faecalis* V583 or OG1RF, as no colonies appeared on agar supplemented with antibiotics that select for V583 and OG1RF. Next, C57BL/6 mice were treated with β-estradiol 1 day prior to inoculation with 10^7 CFU of *E. faecalis* V583 or OG1RF. After 1 day post-inoculation, the vaginal lumen was swabbed and bacteria were plated to enumerate *E. faecalis* V583 and OG1RF vaginal colonization levels (Fig. 1B). Swabs were plated on selective agar to ensure quantification of only the enterococcal strains of interest, restricting growth of native enterococcus. To determine whether *E. faecalis* ascends into reproductive tissues during colonization, murine vaginal, cervical, and uterine tissues were collected and homogenized to enumerate *E. faecalis* V583 abundance. *E. faecalis* was recovered from all mice 1 day post-inoculation in all tissues tested (Fig. 1C) and the CFU recovered from the vaginal swabs were similar to the total CFU counts from the vaginal tissue homogenates (Fig. 1B, C). This level and range of recovered *E. faecalis* CFU is similar to what we have observed using this mouse model for GBS and *S. aureus* vaginal colonization (26, 27). To visualize *E. faecalis* within the murine reproductive tract, mice were inoculated with either WT *E. faecalis* V583 or a V583 strain expressing green fluorescent protein (GFP) (28). These strains both colonize the vaginal tract (Fig. S1A). We harvested the female reproductive tract 1 day post-inoculation to avoid the loss of the GFP plasmid and made serial sections of these tissues, and performed fluorescent microscopy to visualize *E. faecalis*. We observed numerous fluorescent bacteria in the vaginal and uterine lumen (Fig. 1D, F) and embedded in the cervical lamina propria (Fig. 1E). We did not observe background green fluorescence in naïve mice (Fig. S1B, C, D), which coincides with previous experiments performed with GBS and *S. aureus* (26, 27). The presence of fluorescent *E. faecalis* in the cervix and uterus shows that *E. faecalis* can move from the vaginal lumen to the superior organs of the female reproductive tract.

E. faecalis persists in the vaginal tract.

To assess vaginal persistence, C57BL/6 mice were colonized with *E. faecalis* V583 or OG1RF and

swabbed to determine bacterial load over time. Mice were swabbed daily until no colonies appeared on

agar selective for V583 or OG1RF, indicating bacterial clearance from the vaginal lumen. While V583

persisted longer in the mouse vaginal tract, the mean CFU recovered for both V583 and OG1RF

remained constant for the first week and then declined as mice eventually cleared both strains by 11-13

days (Fig. 2A, B). To determine if enterococcal vaginal persistence differs across mouse strains,

C57BL/6 and CD1 mice were inoculated with V583 and swabbed over time. Both mouse strains were

initially colonized with V583, but bacteria in C57BL/6 mice persisted longer (Fig. 2C, D). By day six

only 20% of CD1 mice remained colonized compared to 85% of C57BL/6 mice. Differences in vaginal

persistence may be due to differences in the native vaginal microbiota or immune status between mouse

strains. It is also possible that bacteria occupy different niches within the reproductive tract of different

mouse strains, which warrants further investigation. Overall, these results show that mouse strain

background influences *E. faecalis* vaginal colonization and that C57BL/6 mice are a sufficient model to

assess prolonged *E. faecalis* vaginal colonization and persistence.

Enterococcal pili contribute to interaction with reproductive tract tissues.

The underlineendocarditis- and underlinebiofilm-associated underlinepilin (Ebp) of *E. faecalis* mediates infective endocarditis and

UTIs(29-32), thus we hypothesized that Ebp may similarly contribute to vaginal colonization. To

determine whether Ebp is important for facilitating interaction with the vaginal epithelium, we used a

deletion mutant of *E. faecalis* OG1RF lacking all pilin structural components (Δ*ebpABC*) (33). We

observed that the pilus mutant exhibited significantly reduced adherence to human vaginal and

endocervical cells *in vitro* (Fig. 3A, B). To determine if Ebp is important for *in vivo* vaginal colonization

and persistence, mice were inoculated with either WT OG1RF or OG1RF Δ*ebpABC* and colonization

was quantified over the course of 12 days. We observed no differences in the CFU recovered from the

vaginal lumen between WT OG1RF and OG1RF Δ*ebpABC* strains (Fig. 3C). Taken together these data

suggest that Ebp contributes to *E. faecalis* attachment to reproductive tract tissues, but additional factors

are likely required for persistence in the vaginal lumen *in vivo*.

Identification of additional vaginal colonization factors by transposon mutagenesis analysis.

To identify genetic determinants that confer enterococccal vaginal persistence, we used sequence-

defined <u>mariner</u> technology <u>t</u>ransposon <u>seq</u>uencing (SMarT TnSeq) to screen an *E. faecalis* OG1RF Tn

library consisting of 6,829 unique mutants (34). The library was grown to mid-log phase in triplicate and

10^7 CFU of each replicate was vaginally inoculated into a group of 5 C57BL/6 mice (Fig. 4A). Vaginal

swabs were plated on selective media daily for 8 days and CFU were quantified to assess colonization of

the OG1RF Tn library compared to WT OG1RF (Fig. 4A and B). Genomic DNA was isolated from

pooled Tn libraries recovered on days 1, 5, and 8 post-inoculation and Tn insertion junctions in *E.*

faecalis genomic DNA were sequenced as described by Dale *et al.* (34). Sequenced reads were mapped

to the *E. faecalis* OG1RF genome to identify genes that are necessary for *E. faecalis* vaginal

colonization.

We observed that the *in vivo* vaginal environment altered the abundance of select mutants from the Tn

library pool compared to the original culture input (Fig. 4C, D, E) (Tables S1 - S5). At day 1, a total of

667 depleted mutants were identified (Table S1), along with 544 (Table S2) and 507 (Table S3) at days

5 and 8 respectively; 383 of these mutants were identified at all three time points (Fig. 5A, Table S4A).

Classification by clusters of orthologous groups of proteins (COGs) could be identified in 196 mutants

from all 3 time points, the majority of which are involved in carbohydrate, amino acid, lipid and

nucleotide transport/metabolism, as well as those involved in transcription and defense mechanisms

(Fig. 5B). Of the remaining 187 mutants, 85 had insertions in intergenic regions and 102 were not

assigned a COG domain. Interestingly the *ebpC* transposon mutant (*OG1RF_10871*, which encodes the

shaft component of Ebp) was underrepresented at all time points (Fig. 4C, D, E, Table S4A). Additional

mutants of interest included those involved in ethanolamine catabolism and T7SS components, in which

various components of these systems were underrepresented at all three time points (Tables 1 and S4A).

Furthermore, mutants for multiple sortase-dependent proteins (SDPs), including *ebpC*, were

underrepresented at all time points (Tables 1 and S4A), suggesting that these factors may play important

roles vaginal colonization and persistence.

Potential gain-of-function mutations have also been discovered during genome-wide library screens of

fitness determinants in other bacteria (35-39). In addition to mutations that adversely impact vaginal

colonization, our data shows that Tn insertions in 11 protein coding genes and 11 intergenic regions

potentially enhance bacterial fitness *in vivo*. Nineteen of the 22 enriched mutants were common between

days 1 and 5 whereas the other 3 were unique to day 8 (Table S5A and S5B). Since the majority of the

mutants with increased fitness encode hypothetical proteins, the relationship between these genes and

vaginal colonization is currently unclear and requires further investigation.

Ethanolamine utilization and T7SS genes contribute to *E. faecalis* persistence in the reproductive

tract.

Our TnSeq analysis revealed many potential mutants that exhibited decreased colonization in the murine

vaginal tract. We sought to confirm these results by analyzing mutants from systems with multiple

affected genes. One significantly affected operon was ethanolamine (EA) utilization (*eut*) which consists

of 19 genes in *E. faecalis* (40). Mutants in 4 *eut* genes were significantly underrepresented *in vivo*

compared to the culture input at all time points. These included transposon mutants of the genes

encoding   both   subunits   for   ethanolamine   ammonia   lyase,   *eutB   (OG1RF_11344)* and *eutC*

(OG1RF_11343) (41), a carboxysome structural protein, *eutL (OG1RF_11342)*(42), and the response

regulator, *eutV (OG1RF_11347)*, of the two-component system involved in the regulation of EA

utilization (43) (Table S4A). To assess the importance of EA utilization on *E. faecalis* vaginal

colonization, we co-colonized mice with *E. fae*calis OG1SSp (a derivative of OG1 that is resistant to

streptomycin and spectinomycin) and an OG1RF Δ*eutBC* mutant (44). We note that both WT strains,

OG1RF and OG1SSp, were able to colonize the vaginal tract at similar levels (Fig. S2).   Further

chromosomal DNA sequence comparison of OG1RF and OG1SSp revealed multiple nucleotide

polymorphisms (SNPs) in OG1SSp, but no SNPs in genes in the *eut* locus (Table S6). Compared to the

WT OG1SSp strain, the Δ*eutBC* mutant was cleared significantly faster from the mouse vagina as seen

by CFU from individual mouse swabs, the mean CFU recovered and competitive index (CI) over time

(Fig. 6A, B, C). These results suggest that the utilization of ethanolamine is important for enterococcal

persistence in the vaginal tract.

We also observed that Tn insertion mutants in the T7SS locus were significantly underrepresented at all

time points *in vivo* compared to the culture input. The T7SS has been shown to play an important role in

virulence in multiple bacterial species such as *Staphylococcus*, *Listeria* and *Bacillus* (45). In *E. faecalis*,

genes in the T7SS locus have been shown to be induced during phage infection (46). We observed that

transposon mutants for *esaB* (*OG1RF_11103)*, a putative cytoplasmic accessory protein; *OG1RF_11109*

and *OG1RF_11111*, putative toxin effector proteins; *OG1RF_11113*, a putative toxin, *OG1RF_11114,* a

transmembrane protein; and *OG1RF_11122,* a potential antitoxin protein were all significantly

underrepresented (Table S4A). To confirm the role of the T7SS in vaginal colonization, we utilized an

esaB (*esaB::tn*) mutant, in which *esaB* is disrupted by a transposon and thus Tn-mediated polar effects

may exist for this strain. Following co-colonization with *E. faecalis* OG1RF and the OG1RF *esaB::tn*

mutant strain, we observed that while there were no differences in initial colonization, fewer *esaB::tn*

mutant bacteria were recovered from the vaginal lumen at later time points (Fig. 6D). Since we only

observed differences in colonization between WT OG1RF and OG1RF *esaB::tn* at later time points, we

performed subsequent experiments to determine whether there were differences in ascension between

the two strains. Mice were co-colonized with the two strains and we harvested tissues at day 11, before

there were any colonization differences observed between the two strains. Here, we observed that WT

OG1RF outcompeted the *esaB::tn* mutant strain and was better able to access reproductive tract tissues

(Fig. 6E, F), indicating that the T7SS may be involved in vaginal persistence and ascension in the female

reproductive tract.

DISCUSSION

E. faecalis is associated with a wide spectrum of infections, particularly under immunocompromised

states and during compositional shifts in the host microbiota (47, 48). Although an increasing body of

evidence links enterococci with bacterial vaginosis (BV) and aerobic vaginitis (AV) (22, 23, 49-51), the

molecular determinants that facilitate *E. faecalis* colonization and persistence in the vaginal tract are

largely unknown. Here, we employed *in vitro* and *in vivo* systems to acquire genome-scale interactions

that confer *E. faecalis* fitness within the female reproductive tract. We show that both vancomycin

sensitive enterococci (VSE) and VRE adhere to cell types of vaginal and cervical origin, a signature of

bacterial colonization that precedes tissue invasion and systemic spread. Further, genetic features

involved in biofilm formation, ethanolamine utilization and polymicrobial interactions influence *E.*

faecalis vaginal carriage.

Previous studies have demonstrated the importance of Ebp pili in enterococcal virulence and biofilm

formation (19). We found that deletion of *ebpABC* attenuated binding to human vaginal and

endocervical cells but did not influence bacterial burden in the vaginal lumen, similar to the observed

function of Ebp pili in the intestine (52). Enhanced *E. faecalis* adherence in tissue culture compared to *in*

vivo colonization may reflect the lack of liquid and mucus flow that bacteria encounter within the

vaginal tract, emphasizing the significance of our animal model for investigating *E. faecalis*-vaginal

interactions. Consistent with this observation, an *in vivo* Tn library screen revealed only two

underrepresented biofilm-associated mutants, *ebpC-Tn* and *OG1RF_10506-Tn*, encoding a putative

polysaccharide deacetylase homolog implicated in low biofilm formation *in vitro* (53, 54). Together,

these results show that individual mutations in *ebp* or other well characterized biofilm genes are not

sufficient to impair vaginal niche establishment and/or persistence of enterococci, which likely depends

on the concerted effort of multiple factors. Furthermore, similar to *ebpC-Tn*, we observed that genes for

multiple sortase-dependent proteins (SDPs) were underrepresented at all time points during vaginal

colonization. The genome of OG1RF contains 21 sortase-dependent proteins, including Ebp (52). Other

than *ebpC* (*OG1RF_10871*), we observed that 4 other SDPs are underrepresented during vaginal

colonization, including *OG1RF_10811, OG1RF_11531, OG1RF_11764* and *OG1RF_12054* (Table 1).

Previous reports indicate the importance of SDPs during gastrointestinal colonization by enterococci

(52), implicating the possibility that multiple SDPs also play a role during vaginal colonization.

Transition from nutrient rich laboratory media to the vaginal tract likely imparts dramatic alterations in

E. faecalis metabolism. In support of this hypothesis, our high-throughput Tn mutant screen showed that

mutations in carbohydrate, amino acid and nucleotide metabolic pathways were indispensable in the

vaginal tract. Specifically, we showed that WT bacteria outcompete a *eut* locus mutant during vaginal

colonization. In contrast, Kaval and colleagues demonstrated that mutations in *eut* genes leads to a slight

increase in fitness within the intestine (55). This observation likely reflects varying metabolic

requirements of enterococci in different host environments. While a number of reports exist on the

contributions of EA catabolism in host-bacteria interactions within the intestine (56), studies are lacking

for the relevance of this EA metabolism in other host-associated environments. Our results raise

important questions regarding EA utilization in the female reproductive tract. Although commensal

microbes and the epithelium are rich sources of EA, the composition and source of EA in the vaginal

tract remains unknown. A recent report showed that *E. faecalis* EA utilization attenuates intestinal colitis

in IL10 knockout mice in the presence of a defined microbiota (57). Whether EA utilization promotes

virulence or commensalism for enterococci in the context of vaginal tissue remains to be determined.

Considering that the by-product of EA metabolism, acetate, is anti-inflammatory and promotes IgA

production in the intestine (58, 59), it is intriguing to consider that enterococcal EA catabolism might

modulate immune responses within the female reproductive tract.

T7SSs have been implicated in the maintenance of bacterial membrane integrity, virulence, and inter-

bacterial antagonism (45, 60-66). In *S. aureus*, T7SS encoded proteins confer protection from membrane

damage caused by host fatty acids (65, 66). Although *E. faecalis* T7SS genes were shown to be induced

in response to phage driven membrane damage, direct contributions of these genes in cell envelop

barrier function and/or virulence in the context of animal models are poorly defined. Our TnSeq analysis

revealed that insertional mutations in six T7SS genes diminished early and late vaginal colonization by

E. faecalis OG1RF. In vaginal co-colonization competition experiments, an *esaB::tn* strain reached WT

colonization levels early and showed a defect in long term persistence. The incongruence in the

colonization kinetics of T7SS mutant strains compared to T7SS-Tn library mutants which were observed

early after inoculation, presumably stem from the inherent differences in the vaginal milieu in these two

experiments. The TnSeq library employed in this screen is a complex population of approximately 7,000

unique mutants, and it is very likely that direct or indirect interactions between mutants influences

fitness. LXG–domain toxins, which are part of the T7SS, have been shown to antagonize neighboring

non-kin bacteria (63). The fact that two mutants with LXG–domain encoding gene mutations,

OG1RF_11109 and *OG1RF_11111*, were underrepresented across all time points during vaginal

colonization suggests that these putative antibacterial proteins may influence enterococcal interactions

with the resident microbes of the vaginal tract.

In addition to genes encoding SDPs, ethanolamine utilization and T7SS, other Tn mutants that were

underrepresented in all time points are worth discussing. For example, *OG1RF_12241*, a homolog of the

oxidative stress regulator *hypR*, was underrepresented at all time points (Table S4A). We have recently

shown that this gene is involved in phage VPE25 infection of *E. faecalis* OG1RF (46). Furthermore,

enterococcal mutants in the CRISPR/*cas9* locus (*OG1RF_10404* and *OG1RF_10407*) were

underrepresented at all three time points (Table S4A). While the role of CRISPR-Cas systems in

providing prokaryotic immunity to mobile genetic elements has been extensively investigated, there is

also evidence suggesting that this system may be involved in other prokaryotic processes besides

adaptive immunity. Cas9 has been shown to have various functions in regulation of virulence in a

number of bacteria including *Francisella novicida*, *Campylobacter jejuni*, and *Streptococcus agalactiae*

(67-69). In *E. faecalis*, CRISPR-Cas-harboring strains are associated with increased capacity to form

biofilms and increased mortality in a mouse urinary tract model (70). Our Tn-seq analysis further reveals

the potential importance for Cas9 during vaginal colonization, which warrants follow up studies.

While a majority of underrepresented mutants were common to all time points, we identified certain

mutants unique to either early or late colonization. For example, the ethanolamine utilization protein

EutQ (*OG1RF_11333*), a classified acetate kinase in *Salmonella enterica* (71), was significantly

underrepresented at day 1, but not the later time points. We also observed that Tn mutants for the

transmembrane signaling protein kinase IreK (*OG1RF_12384*) were underrepresented only at day 1. In

E. faecalis, IreK is involved in regulation of cell wall homeostasis (72), long-term persistence in the gut

and has also been shown to be essential for enterococcal T7SS expression and subsequent activity (73).

These proteins may therefore be important contributors to enterococcal vaginal colonization, though

further investigation is required. Our Tn-seq analysis also identified Tn mutants that were unique to day

8 post-inoculation, indicating that these factors may be important for later stage colonization and

persistence. One observed mutant was the *fsrB* gene (*OG1RF_11528*) of the Fsr quorum-sensing system,

which directly regulates virulence factors such as serine protease and gelatinase, while also indirectly

regulating other virulence factors involved in surface adhesion and biofilm development (74-76).

Although it is not well understood whether biofilms are being formed during vaginal colonization,

certain hits in our Tn-seq (i.e. *ebpC* and *fsrB*) analysis suggests that biofilm-associated factors play a

role in enterococcal persistence in the vaginal tract. Bacterial mutants for the response regulator *croR*

(*OG1RF_*12535) were also underrepresented only at the later time point. CroR has shown to be involved

in virulence regulation, cell wall homeostasis and stress response, and antibiotic resistance (77-79).

Finally, underrepresentation of the sortase-associated gene (*OG1RF_10872*) was also unique to late

colonization. The underrepresentation of enterococcal mutants late in vaginal colonization suggests

these factors may be important for long-term persistence of *E. faecalis* in the vaginal tract. While the

majority of underrepresented mutants were common to all time points, mutants unique to certain time

points indicate that some factors may be important for either initial colonization or enterococcal survival

in the vaginal tract.

Here, we report the utilization of a mouse model for investigating host-enterococcal interactions in the

vaginal tract. This will be a useful model for analyzing the bacterial and host factors that govern

enterococcal vaginal colonization, as well as characterizing the polymicrobial interactions that may

contribute to *E. faecalis* niche establishment and persistence. Transposon library screening of *E.*

faecalis recovered from the mouse vagina has revealed new insights into our understanding of

enterococcal vaginal carriage. Our results emphasize the importance of ethanolamine utilization and

T7SS components for successful *E. faecalis* colonization of the female reproductive tract, highlighting

the complex nature of this niche.

METHODS

Bacterial strains and culture conditions.

A detailed list of bacterial strains can be found in Table S7. *E. faecalis* strains V583 (80) and OG1- (RF

and SSp) (70, 81) were used for these experiments. *E. faecalis* was grown in brain heart infusion (BHI

(82)) broth at 37℃ with aeration and growth was monitored by measuring the optical density at 600nm

(OD_{600}). For selection of *E. faecalis* V583, BHI agar was supplemented with gentamicin (100 μg/ml).

For selection of *E. faecalis* OG1RF, OG1RF Δ*ebpABC* (33) and OG1RF Δ*eutBC* (44), BHI agar was

supplemented with rifampicin (50 μg/ml) and fusidic acid (25 μg/ml). For selection of *E. faecalis*

OG1SSp, BHI agar was supplemented with streptomycin (150 µg/ml) and spectinomycin (100 µg/ml).

E. faecalis OG1RF *esaB*::*tn* (34) was grown on BHI agar supplemented with rifampicin (50 µg/ml),

fusidic acid (25 µg/L), and chloramphenicol (15 µg/ml).

In vitro **adherence assays**

Immortalized VK2 human vaginal epithelial cells and End1 human endocervical epithelial cells were

obtained from the American Type Culture Collection (VK2.E6E7, ATCC CRL-2616 and End1/E6E7,

ATCC CRL-2615) and were maintained in keratinocyte serum-free medium (KSFM; Gibco) with 0.1

ng/mL human recombinant epidermal growth factor (EGF; Gibco) and 0.05 mg/ml bovine pituitary

extract (Gibco) at 37°C with 5% CO_2. Assays to determine cell surface-adherent *E. faecalis* were

performed as described previously when quantifying GBS adherence (83). Briefly, bacteria were grown

to mid-log phase (OD_{600} = 0.4 - 0.6) and added to cell monolayers (multiplicity of infection [MOI] = 1).

After a 30 minute incubation, cells were washed with phosphate-buffered saline (PBS) three times

following detachment with 0.1 mL of 0.25% trypsin-EDTA solution and lysed with addition of 0.4 mL

of 0.025% TritonX-100 in PBS by vigorous pipetting. The lysates were then serially diluted and plated

on Todd Hewitt agar (THA) to enumerate the bacterial CFU. Experiments were performed at least three

times with each condition in triplicate, and results from a representative experiment are shown.

Murine vaginal colonization model.

Animal experiments were approved by the Institutional Animal Care and Use Committee at University

of Colorado-Anschutz Medical Campus protocols #00316 and #00253 and performed using accepted

veterinary standards. A mouse vaginal colonization model for GBS was adapted for our studies (26).

Eight-week old female CD1 (Charles River) or C57BL/6 (Jackson) mice were injected intraperitoneally

with 0.5 mg 17β-estradiol (Sigma) 1 day prior to colonization with *E. faecalis*. Mice were vaginally

inoculated by gently pipetting 10^7 CFU of *E. faecalis* in 10μL PBS into the vaginal tract, avoiding

contact with the cervix. On subsequent days the vaginal lumen was swabbed with a sterile ultrafine swab

(Puritan). For co-colonization, mice were inoculated with two of the following *E. faecalis* strains:

OG1SSp, OG1RF or deletion mutants in the OG1RF background. To assess tissue CFU, mice were

euthanized according to approved veterinary protocols and the female reproductive tract tissues were

dissected and placed into 500μL PBS and bead beat for 2 min to homogenize the tissues. The resulting

homogenate was serially diluted and *E. faecalis* CFU enumerated on BHI agar supplemented with

antibiotics to select for the strain of interest.

Histology.

Mice were inoculated with *E. faecalis* V583 containing a plasmid that expresses *gfp* (pMV158GFP) and

contains resistance to tetracycline (15 μg/mL) [23]. After 1 day post-inoculation, the murine female

reproductive tract was harvested, embedded into Optimal Cutting Temperature (OCT) compound

(Sakura), and sectioned at 7μm with a CM1950 freezing cryostat (Leica). For fluorescence microscopy,

coverslips were mounted with VECTASHIELD mounting medium containing 4',6-diamidino-2-

phenylindole (DAPI, Vector Labs). Images were taken with a BZ-X710 microscope (Keyence) [22].

Transposon mutant library growth and vaginal colonization.

The *E. faecalis* OG1RF transposon mutant library was generated previously (34). The E. *faecalis*

OG1RF pooled transposon library was inoculated into 5 ml of BHI at a total of 10^8 CFU and grown with

aeration to an OD_{600} of 0.5. The library was inoculated into the vaginal tracts of C57BL/6 mice at

10^7 CFU. The library was plated on BHI agar to confirm the inoculum for all groups of mice. Mice were

swabbed daily and swabs were plated on BHI supplemented with rifampicin (50 μg/ml), fusidic acid

(25 μg/ml) and chloramphenicol (20 μg/ml) to quantify CFU. On days 1, 5, and 8, undiluted swabs were

plated on BHI agar with antibiotics and grown to a bacterial lawn. Bacteria were scraped and re-

suspended in PBS and pelleted. DNA from days 1, 5, 8, and the input culture was extracted from pellets

using a ZymoBIOMICS™ DNA MiniPrep kit (Zymo Research).

Transposon library sequencing and data analysis.

Transposon-junction DNA library preparation and Illumina NovaSeq 6000 DNA sequencing (150 base

paired end mode) was performed by the Microarray and Genomics Core at the University of Colorado

Anschutz Medical Campus as previously described (46)**.** For downstream analysis of transposon-

junctions, we used only the R1 reads generated by paired end sequencing. Illumina adapter trimmed raw

reads were mapped to the *E. faecalis* OG1RF reference sequence (NC_017316.1) and differentially

abundant transposon mutants were identified using statistical analysis scripts established by Dale *et*

al.(34). The abundance of Tn mutants in culture was compared to input library used for culture

inoculation and mutants that are not significantly different ($p > 0.05$) between these two samples were

considered for the next steps of the analysis. For comparisons between *in vivo* and *in vitro* samples,

mutants were considered significantly different if the adjusted P value was < 0.05 and a $\log_2$ (fold

change) > 1.

Genomic DNA sequencing and comparative analysis

E. faecalis genomic DNA was purified using a ZymoBIOMICS™ DNA Miniprep Kit (Zymo Research)

and 150 bp paired end sequencing was performed on Illumina NextSeq 550 by the Microbial Genome

Sequencing Center, University of Pittsburgh. *E. faecalis* OG1SSp genome DNA was purified using a

Qiagen DNeasy kit and was sequenced on the MiSeq platform (2 X 75 bp) at the University of

Minnesota Genomics Center. All reads were mapped to *E. faecalis* OG1RF reference sequence

(NC_017316.1) using CLC Workbench (Qiagen). The basic variant caller tool in CLC Genomics

Workbench was used to identify single nucleotide polymorphisms using default settings (similarity

fraction = 0.5 and length fraction = 0.8).

Data availability.

The Tn-Seq and genomic DNA reads have been deposited at the European Nucleotide Archive under

accession numbers PRJEB37929 and PRJEB39171, respectively.

Statistical analysis.

GraphPad Prism version 7.0 was used for statistical analysis and statistical significance was accepted at

P values of < 0.05 (* P < 0.05; ** P < 0.005; *** P < 0.0005; **** P < 0.00005). Specific tests are

indicated in figure legends.

ACKNOWLEGMENTS

We thank Kimberly Kline for providing *E. faecalis* OG1RF Δ*ebpABC*, Danielle Garsin for providing *E.*

faecalis OG1RF Δ*eutBC*, Katrina Diener and Monica Ransom for custom TnSeq library preparation and

the Microarray and Genomics Core at the University of Colorado Anschutz Medical Campus for DNA

sequencing. This study was supported by the NIH 5T32AI007405-28 to B.L.S., the NIH/NIAID R21

AI130857 to K.S.D, the NIH/NIAID R01 AI141479 to B.A.D, and a University of Colorado School of

Medicine IMPA to K.S.D and B.A.D.

FIGURE LEGENDS

Figure 1: *E. faecalis* colonizes the murine female reproductive tract. (A) Adherence of *E. faecalis*

V583 and OG1RF to human vaginal and endocervical cells. Data are expressed as percent recovered

cell-associated *E. faecalis* relative to the initial inoculum. Experiments were performed in triplicate and

error bars represent standard deviations (SDs); the results of a representative experiment are shown. (B)

CFU counts of V583 and OG1RF recovered from vaginal swabs 1 day post-inoculation. (C) CFU counts

of V583 from vaginal, cervical and uterine tissue 1 day post-inoculation. (D, E, F) Mice were inoculated

with V583 expressing *gfp* and 7μm sections of vaginal (D), cervical (E) and uterine (F) tissue were

collected 1 day post-inoculation and stained with DAPI for fluorescence microscopy. White arrows

indicate green fluorescent bacteria present in tissue sections. Images were all taken at 40x magnification.

LP = lamina propria, EP = epithelial layer, Lu = lumen.

Figure 2: *E. faecalis* persists in the murine vaginal tract. (A and B) *E. faecalis* V583 and OG1RF in

the murine vaginal tract. C57BL/6 mice (n = 10) were inoculated with 10^7 V583 or OG1RF CFU and the

vaginal lumen of each mouse was swabbed daily and swabs were serially diluted and plated on selective

media to quantify CFU. Data are presented as recovered CFU per mouse (A) and mean recovered CFU

(B). Data was analyzed using a Two-way ANOVA; ∗∗ $P < 0.001$. (C and D) CD1 (n = 10) and C57BL/6

(n = 7) mice were inoculated with V583 and the vaginal lumen of each mouse was swabbed daily,

serially diluted, and plated on selective media to quantify CFU. Data are presented as recovered CFU

per mouse (C) and mean recovered CFU (D). Black lines indicate the median of CFU values.

Figure 3: The role of enterococcal pili during vaginal colonization. (A and B) *E. faecalis* OG1RF

WT and OG1RF Δ*ebpABC* adherence to human vaginal (A) and endocervical (B) cells. Data are

expressed as percent recovered cell-associated *E. faecalis* relative to the initial inoculum. Experiments

were performed with four technical replicates and error bars represent SDs; the results of three

combined biological replicates are shown and analyzed using an unpaired t-test; *** P < 0.0001*; **** P

< 0.00001. (C) C57BL/6 mice were inoculated with either OG1RF WT or OG1RF Δ*ebpABC* and the

vaginal lumen was swabbed daily. Data was analyzed using a Two-way ANOVA with Sidak's multiple

comparisons (P>0.05). Black lines indicate the median of CFU values.

Figure 4: Identification of additional factors required for vaginal colonization and persistence by

transposon mutant library screen. (A) Schematic representing experimental approach. C57BL/6 mice

were treated with 17β-estradiol prior to inoculation with the OG1RF Tn library in triplicate groups or

WT OG1RF. The vaginal lumen was swabbed and CFU was enumerated daily. DNA from recovered

OG1RF Tn mutants was sequenced on days 1, 5 and 8. (B) CFU recovered from vaginal swabs of

triplicate groups of mice colonized with OG1RF Tn mutagenesis library (T1, T2, T3) and OG1RF WT

(n=5 mice per group). (C, D and E) Volcano plots depicting underrepresented and overrepresented

mutants *in vivo* compared to culture on day 1 (C), day 5 (D) and day 8 (E) post-inoculation. (C) 667

underrepresented genes and 21 overrepresented genes *in vivo* compared to culture. (D) 404

underrepresented genes and 6 overrepresented genes *in vivo* compared to culture. (E) 507

underrepresented genes and 3 overrepresented genes *in vivo* compared to culture. Colored dots

represent mutants of interest. Pink = *ebpC* Tn mutant, Orange = *eut* Tn mutants, Green = T7SS gene Tn

mutants. Black solid line represents cut-off for statistical significance.

Figure 5: Classification of transposon insertion mutants by Cluster of Orthologous Groups

(COGs). (A) Euler plot representing number of underrepresented mutants at all time points. (B) Cluster

of Orthologous Groups (COGs) from underrepresented mutants common to all three time points

categorized into functional categories. Total = 196 represents all common mutants that were assigned a

COG domain.

Figure 6: Ethanolamine utilization and type VII secretion system genes contribute to enterococcal

persistence in the vaginal tract. (A, B and C) C57BL/6 mice were co-inoculated with OG1SSp WT

and OG1RF Δ*eutBC* and vaginal lumen was swabbed to quantify CFU. Data are presented as recovered

CFU per swab (A), mean recovered CFU (B) and CI between WT and mutant strains (C). Data was

analyzed using a Two-way ANOVA with Sidak's multiple comparisons; $*P < 0.05$, $**P < 0.005$, $****$

$P < 0.00005$. (D) C57BL/6 mice were co-inoculated with OG1RF WT and OG1RF *esaB::tn* and vaginal

lumen was swabbed to quantify CFU. Data was analyzed using a Two-way ANOVA with Sidak's

multiple comparisons; $**P < 0.005$, $***P < 0.0005$ (E and F) C57BL/6 mice were co-inoculated with

OG1RF WT and OG1RF *esaB*::*tn* and reproductive tissue was collected at 11 days post-inoculation.

Data are presented as recovered Log_{10} CFU/gram (E) and CI between WT and Tn-mutant strain (F). CI

is enumerated by calculating the ratio of WT to mutant *E. faecalis* recovered from the mouse

reproductive tract. A CI >1 indicates an advantage to WT *E. faecalis*. Values below the limit of

detection were enumerated as one-half the limit of detection. Data was analyzed using a paired t-test; $*P$

< 0.05, $**P < 0.005$. Black lines indicate the median of CFU values.

Table 1: Selected list of differentially abundant transposon mutants during vaginal colonization

compared to *in vitro* cultures.

REFERENCES:

1.   Lebreton F, Willems RJL, Gilmore MS. 2014. Enterococcus Diversity, Origins in Nature, and
Gut Colonization. *In* Gilmore MS, Clewell DB, Ike Y, Shankar N (ed), Enterococci: From
Commensals to Leading Causes of Drug Resistant Infection, Boston.
2.   Coolen MJ, Post E, Davis CC, Forney LJ. 2005. Characterization of microbial communities
found in the human vagina by analysis of terminal restriction fragment length polymorphisms of
16S rRNA genes. Appl Environ Microbiol 71:8729-37.
3.   Chaturvedi SK, Manche Gowda S, Ahmed HU, Alosaimi FD, Andreone N, Bobrov A, Bulgari
V, Carra G, Castelnuovo G, de Girolamo G, Gondek T, Jovanovic N, Kamala T, Kiejna A, Lalic
N, Lecic-Tosevski D, Minhas F, Mutiso V, Ndetei D, Rabbani G, Somruk S, Srikanta S, Taj R,
Valentini U, Vukovic O, Wolwer W, Cimino L, Nouwen A, Lloyd C, Sartorius N. 2019. More
anxious than depressed: prevalence and correlates in a 15-nation study of anxiety disorders in
people with type 2 diabetes mellitus. Gen Psychiatr 32:e100076.
4.   Huycke MM, Sahm DF, Gilmore MS. 1998. Multiple-drug resistant enterococci: the nature of
the problem and an agenda for the future. Emerg Infect Dis 4:239-49.
5.   Schindlbeck C, Dziura D, Mylonas I. 2014. Diagnosis of pelvic inflammatory disease (PID):
intra-operative findings and comparison of vaginal and intra-abdominal cultures. Arch Gynecol
Obstet 289:1263-9.
6.   Mohamed JA, Huang DB. 2007. Biofilm formation by enterococci. J Med Microbiol 56:1581-
1588.
7.   Arias CA, Murray BE. 2008. Emergence and management of drug-resistant enterococcal
infections. Expert Rev Anti Infect Ther 6:637-55.
8.   Uttley AH, Collins CH, Naidoo J, George RC. 1988. Vancomycin-resistant enterococci. Lancet
1:57-8.
9.   Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C,
Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outterson K, Patel J, Cavaleri M, Cox EM,
Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N, Group
WHOPPLW. 2018. Discovery, research, and development of new antibiotics: the WHO priority
list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis 18:318-327.
10.   Martin HL, Richardson BA, Nyange PM, Lavreys L, Hillier SL, Chohan B, Mandaliya K,
Ndinya-Achola JO, Bwayo J, Kreiss J. 1999. Vaginal lactobacilli, microbial flora, and risk of
human immunodeficiency virus type 1 and sexually transmitted disease acquisition. J Infect Dis
180:1863-8.
11.   Madsen KT, Skov MN, Gill S, Kemp M. 2017. Virulence Factors Associated with Enterococcus
Faecalis Infective Endocarditis: A Mini Review. Open Microbiol J 11:1-11.
12.   Minhas A, Dewey M, Vavere AL, Tanami Y, Ostovaneh MR, Laule M, Rochitte CE, Niinuma
H, Kofoed KF, Geleijns J, Hoe J, Chen MY, Kitagawa K, Nomura C, Clouse ME, Rybicki FJ,
Tan SY, Paul N, Matheson M, Cox C, Rief M, Maier P, Feger S, Plotkin M, Schonenberger E.
2019. Patient Preferences for Coronary CT Angiography with Stress Perfusion, SPECT, or
Invasive Coronary Angiography. Radiology 291:340-348.
13.   Kumar N, Vishnubhatla S, Wadhawan AN, Minhas S, Gupta P. 2020. A randomized, double
blind, sham-controlled trial of repetitive transcranial magnetic stimulation (rTMS) in the
treatment of negative symptoms in schizophrenia. Brain Stimul 13:840-849.
14.   Thurlow LR, Thomas VC, Narayanan S, Olson S, Fleming SD, Hancock LE. 2010. Gelatinase
contributes to the pathogenesis of endocarditis caused by Enterococcus faecalis. Infect Immun
78:4936-43.

15.   Ali L, Goraya MU, Arafat Y, Ajmal M, Chen JL, Yu D. 2017. Molecular Mechanism of
Quorum-Sensing in Enterococcus faecalis: Its Role in Virulence and Therapeutic Approaches.
Int J Mol Sci 18.
16.   Waters CM, Antiporta MH, Murray BE, Dunny GM. 2003. Role of the Enterococcus faecalis
GelE protease in determination of cellular chain length, supernatant pheromone levels, and
degradation of fibrin and misfolded surface proteins. J Bacteriol 185:3613-23.
17.   Qin X, Singh KV, Xu Y, Weinstock GM, Murray BE. 1998. Effect of disruption of a gene
encoding an autolysin of Enterococcus faecalis OG1RF. Antimicrob Agents Chemother 42:2883-
8.
18.   Tacconelli E, Cataldo MA. 2008. Vancomycin-resistant enterococci (VRE): transmission and
control. Int J Antimicrob Agents 31:99-106.
19.   Ch'ng JH, Chong KKL, Lam LN, Wong JJ, Kline KA. 2019. Biofilm-associated infection by
enterococci. Nat Rev Microbiol 17:82-94.
20.   Leyva-Gomez G, Prado-Audelo MLD, Ortega-Pena S, Mendoza-Munoz N, Urban-Morlan Z,
Gonzalez-Torres M, Gonzalez-Del Carmen M, Figueroa-Gonzalez G, Reyes-Hernandez OD,
Cortes H. 2019. Modifications in Vaginal Microbiota and Their Influence on Drug Release:
Challenges and Opportunities. Pharmaceutics 11.
21.   Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R,
Russell J, Tacket CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal
microbiome of reproductive-age women. Proc Natl Acad Sci U S A 108 Suppl 1:4680-7.
22.   Donders GGG, Bellen G, Grinceviciene S, Ruban K, Vieira-Baptista P. 2017. Aerobic vaginitis:
no longer a stranger. Res Microbiol 168:845-858.
23.   Kaambo E, Africa C, Chambuso R, Passmore JS. 2018. Vaginal Microbiomes Associated With
Aerobic Vaginitis and Bacterial Vaginosis. Front Public Health 6:78.
24.   Sahm DF, Kissinger J, Gilmore MS, Murray PR, Mulder R, Solliday J, Clarke B. 1989. In vitro
susceptibility studies of vancomycin-resistant *Enterococcus faecalis*. Antimicrob Agents
Chemother 33:1588-91.
25.   Gold OG, Jordan HV, van Houte J. 1975. The prevalence of enterococci in the human mouth and
their pathogenicity in animal models. Arch Oral Biol 20:473-7.
26.   Patras KA, Doran KS. 2016. A Murine Model of Group B Streptococcus Vaginal Colonization. J
Vis Exp doi:10.3791/54708.
27.   Deng L, Schilcher K, Burcham LR, Kwiecinski JM, Johnson PM, Head SR, Heinrichs DE,
Horswill AR, Doran KS. 2019. Identification of Key Determinants of Staphylococcus aureus
Vaginal Colonization. mBio 10.
28.   Nieto C, Espinosa M. 2003. Construction of the mobilizable plasmid pMV158GFP, a derivative
of pMV158 that carries the gene encoding the green fluorescent protein. Plasmid 49:281-5.
29.   Nallapareddy SR, Singh KV, Sillanpaa J, Garsin DA, Hook M, Erlandsen SL, Murray BE. 2006.
Endocarditis and biofilm-associated pili of Enterococcus faecalis. J Clin Invest 116:2799-807.
30.   Nallapareddy SR, Sillanpaa J, Mitchell J, Singh KV, Chowdhury SA, Weinstock GM, Sullam
PM, Murray BE. 2011. Conservation of Ebp-type pilus genes among Enterococci and
demonstration of their role in adherence of Enterococcus faecalis to human platelets. Infect
Immun 79:2911-20.
31.   Nielsen HV, Guiton PS, Kline KA, Port GC, Pinkner JS, Neiers F, Normark S, Henriques-
Normark B, Caparon MG, Hultgren SJ. 2012. The metal ion-dependent adhesion site motif of the
Enterococcus faecalis EbpA pilin mediates pilus function in catheter-associated urinary tract
infection. mBio 3:e00177-12.

32.   Nallapareddy SR, Singh KV, Sillanpaa J, Zhao M, Murray BE. 2011. Relative contributions of
Ebp Pili and the collagen adhesin ace to host extracellular matrix protein adherence and
experimental urinary tract infection by Enterococcus faecalis OG1RF. Infect Immun 79:2901-10.
33.   Afonina I, Lim XN, Tan R, Kline KA. 2018. Planktonic Interference and Biofilm Alliance
between Aggregation Substance and Endocarditis- and Biofilm-Associated Pili in Enterococcus
faecalis. J Bacteriol 200.
34.   Dale JL, Beckman KB, Willett JLE, Nilson JL, Palani NP, Baller JA, Hauge A, Gohl DM,
Erickson R, Manias DA, Sadowsky MJ, Dunny GM. 2018. Comprehensive functional analysis of
the *Enterococcus faecalis* core genome using an ordered, sequence-defined collection of
insertional mutations in strain OG1RF. mSystems 3.
35.   Bachman MA, Breen P, Deornellas V, Mu Q, Zhao L, Wu W, Cavalcoli JD, Mobley HL. 2015.
Genome-Wide Identification of Klebsiella pneumoniae Fitness Genes during Lung Infection.
mBio 6:e00775.
36.   Jung HJ, Littmann ER, Seok R, Leiner IM, Taur Y, Peled J, van den Brink M, Ling L, Chen L,
Kreiswirth BN, Goodman AL, Pamer EG. 2019. Genome-Wide Screening for Enteric
Colonization Factors in Carbapenem-Resistant ST258 Klebsiella pneumoniae. mBio 10.
37.   Shames SR, Liu L, Havey JC, Schofield WB, Goodman AL, Roy CR. 2017. Multiple Legionella
pneumophila effector virulence phenotypes revealed through high-throughput analysis of
targeted mutant libraries. Proc Natl Acad Sci U S A 114:E10446-E10454.
38.   Skurnik D, Roux D, Aschard H, Cattoir V, Yoder-Himes D, Lory S, Pier GB. 2013. A
comprehensive analysis of in vitro and in vivo genetic fitness of Pseudomonas aeruginosa using
high-throughput sequencing of transposon libraries. PLoS Pathog 9:e1003582.
39.   Wiles TJ, Norton JP, Russell CW, Dalley BK, Fischer KF, Mulvey MA. 2013. Combining
quantitative genetic footprinting and trait enrichment analysis to identify fitness determinants of
a bacterial pathogen. PLoS Genet 9:e1003716.
40.   Del Papa MF, Perego M. 2008. Ethanolamine activates a sensor histidine kinase regulating its
utilization in Enterococcus faecalis. J Bacteriol 190:7147-56.
41.   Roof DM, Roth JR. 1988. Ethanolamine utilization in Salmonella typhimurium. J Bacteriol
170:3855-63.
42.   Thompson MC, Crowley CS, Kopstein J, Bobik TA, Yeates TO. 2014. Structure of a bacterial
microcompartment shell protein bound to a cobalamin cofactor. Acta Crystallogr F Struct Biol
Commun 70:1584-90.
43.   Fox KA, Ramesh A, Stearns JE, Bourgogne A, Reyes-Jara A, Winkler WC, Garsin DA. 2009.
Multiple posttranscriptional regulatory mechanisms partner to control ethanolamine utilization in
Enterococcus faecalis. Proc Natl Acad Sci U S A 106:4435-40.
44.   Kaval KG, Gebbie M, Goodson JR, Cruz MR, Winkler WC, Garsin DA. 2019. Ethanolamine
Utilization and Bacterial Microcompartment Formation Are Subject to Carbon Catabolite
Repression. J Bacteriol 201.
45.   Cao Z, Casabona MG, Kneuper H, Chalmers JD, Palmer T. 2016. The type VII secretion system
of *Staphylococcus aureus* secretes a nuclease toxin that targets competitor bacteria. Nat
Microbiol 2:16183.
46.   Chatterjee A, Willett JLE, Nguyen UT, Monogue B, Palmer KL, Dunny GM, Duerkop BA.
2020. Parallel genomics uncover novel enterococcal-bacteriophage interactions. mBio 11.
47.   Brandl K, Plitas G, Mihu CN, Ubeda C, Jia T, Fleisher M, Schnabl B, DeMatteo RP, Pamer EG.
2008. Vancomycin-resistant enterococci exploit antibiotic-induced innate immune deficits.
Nature 455:804-7.

48.   Ubeda C, Taur Y, Jenq RR, Equinda MJ, Son T, Samstein M, Viale A, Socci ND, van den Brink
669   MR, Kamboj M, Pamer EG. 2010. Vancomycin-resistant Enterococcus domination of intestinal
670   microbiota is enabled by antibiotic treatment in mice and precedes bloodstream invasion in
671   humans. J Clin Invest 120:4332-41.
672   49.   Dezwaan DC, Mequio MJ, Littell JS, Allen JP, Rossbach S, Pybus V. 2007. Purification and
673   characterization of enterocin 62-6, a two-peptide bacteriocin produced by a vaginal strain of
674   Enterococcus faecium: Potential significance in bacterial vaginosis. Microb Ecol Health Dis
675   19:241-250.
676   50.   Kelly MC, Mequio MJ, Pybus V. 2003. Inhibition of vaginal lactobacilli by a bacteriocin-like
677   inhibitor produced by Enterococcus faecium 62-6: potential significance for bacterial vaginosis.
678   Infect Dis Obstet Gynecol 11:147-56.
679   51.   Tempera G, Abbadessa G, Bonfiglio G, Cammarata E, Cianci A, Corsello S, Raimondi A, Ettore
680   G, Nicolosi D, Furneri PM. 2006. Topical kanamycin: an effective therapeutic option in aerobic
681   vaginitis. J Chemother 18:409-14.
682   52.   Banla LI, Pickrum AM, Hayward M, Kristich CJ, Salzman NH. 2019. Sortase-Dependent
683   Proteins Promote Gastrointestinal Colonization by Enterococci. Infect Immun 87.
684   53.   Willett JLE, Ji MM, Dunny GM. 2019. Exploiting biofilm phenotypes for functional
685   characterization of hypothetical genes in Enterococcus faecalis. NPJ Biofilms Microbiomes 5:23.
686   54.   Balomenou S, Fouet A, Tzanodaskalaki M, Couture-Tosi E, Bouriotis V, Boneca IG. 2013.
687   Distinct functions of polysaccharide deacetylases in cell shape, neutral polysaccharide synthesis
688   and virulence of Bacillus anthracis. Mol Microbiol 87:867-83.
689   55.   Kaval KG, Singh KV, Cruz MR, DebRoy S, Winkler WC, Murray BE, Garsin DA. 2018. Loss of
690   Ethanolamine Utilization in Enterococcus faecalis Increases Gastrointestinal Tract Colonization.
691   mBio 9.
692   56.   Kaval KG, Garsin DA. 2018. Ethanolamine Utilization in Bacteria. mBio 9.
693   57.   Lengfelder I, Sava IG, Hansen JJ, Kleigrewe K, Herzog J, Neuhaus K, Hofmann T, Sartor RB,
694   Haller D. 2019. Complex Bacterial Consortia Reprogram the Colitogenic Activity of
695   Enterococcus faecalis in a Gnotobiotic Mouse Model of Chronic, Immune-Mediated Colitis.
696   Front Immunol 10:1420.
697   58.   Maslowski KM, Vieira AT, Ng A, Kranich J, Sierro F, Yu D, Schilter HC, Rolph MS, Mackay F,
698   Artis D, Xavier RJ, Teixeira MM, Mackay CR. 2009. Regulation of inflammatory responses by
699   gut microbiota and chemoattractant receptor GPR43. Nature 461:1282-6.
700   59.   Wu W, Sun M, Chen F, Cao AT, Liu H, Zhao Y, Huang X, Xiao Y, Yao S, Zhao Q, Liu Z, Cong
701   Y. 2017. Microbiota metabolite short-chain fatty acid acetate promotes intestinal IgA response to
702   microbiota which is mediated by GPR43. Mucosal Immunol 10:946-956.
703   60.   Burts ML, Williams WA, DeBord K, Missiakas DM. 2005. EsxA and EsxB are secreted by an
704   ESAT-6-like system that is required for the pathogenesis of *Staphylococcus aureus* infections.
705   Proc Natl Acad Sci U S A 102:1169-74.
706   61.   Kneuper H, Cao ZP, Twomey KB, Zoltner M, Jager F, Cargill JS, Chalmers J, van der Kooi-Pol
707   MM, van Dijl JM, Ryan RP, Hunter WN, Palmer T. 2014. Heterogeneity in ess transcriptional
708   organization and variable contribution of the Ess/Type VII protein secretion system to virulence
709   across closely related *Staphyloccus aureus* strains. Mol Microbiol 93:928-43.
710   62.   Ohr RJ, Anderson M, Shi M, Schneewind O, Missiakas D. 2017. EssD, a nuclease effector of the
711   *Staphylococcus aureus* ESS Pathway. J Bacteriol 199.
712   63.   Whitney JC, Peterson SB, Kim J, Pazos M, Verster AJ, Radey MC, Kulasekara HD, Ching MQ,
713   Bullen NP, Bryant D, Goo YA, Surette MG, Borenstein E, Vollmer W, Mougous JD. 2017. A

broadly distributed toxin family mediates contact-dependent antagonism between Gram-positive
bacteria. Elife 6.

64.   Ishii K, Adachi T, Yasukawa J, Suzuki Y, Hamamoto H, Sekimizu K. 2014. Induction of
virulence gene expression in *Staphylococcus aureus* by pulmonary surfactant. Infect Immun
82:1500-10.

65.   Lopez MS, Tan IS, Yan D, Kang J, McCreary M, Modrusan Z, Austin CD, Xu M, Brown EJ.
2017. Host-derived fatty acids activate type VII secretion in *Staphylococcus aureus*. Proc Natl
Acad Sci U S A 114:11223-11228.

66.   Tchoupa AK, Watkins KE, Jones RA, Kuroki A, Alam MT, Perrier S, Chen Y, Unnikrishnan M.
2020. The type VII secretion system protects *Staphylococcus aureus* against antimicrobial host
fatty acids. bioRxiv doi:10.1101/572172:572172.

67.   Sampson TR, Saroj SD, Llewellyn AC, Tzeng YL, Weiss DS. 2013. A CRISPR/Cas system
mediates bacterial innate immune evasion and virulence. Nature 497:254-7.

68.   Shabbir MAB, Tang Y, Xu Z, Lin M, Cheng G, Dai M, Wang X, Liu Z, Yuan Z, Hao H. 2018.
The Involvement of the Cas9 Gene in Virulence of Campylobacter jejuni. Front Cell Infect
Microbiol 8:285.

69.   Spencer BL, Deng L, Patras KA, Burcham ZM, Sanches GF, Nagao PE, Doran KS. 2019. Cas9
Contributes to Group B Streptococcal Colonization and Disease. Front Microbiol 10:1930.

70.   Bourgogne A, Garsin DA, Qin X, Singh KV, Sillanpaa J, Yerrapragada S, Ding Y, Dugan-Rocha
S, Buhay C, Shen H, Chen G, Williams G, Muzny D, Maadani A, Fox KA, Gioia J, Chen L,
Shang Y, Arias CA, Nallapareddy SR, Zhao M, Prakash VP, Chowdhury S, Jiang H, Gibbs RA,
Murray BE, Highlander SK, Weinstock GM. 2008. Large scale variation in Enterococcus
faecalis illustrated by the genome analysis of strain OG1RF. Genome Biol 9:R110.

71.   Moore TC, Escalante-Semerena JC. 2016. The EutQ and EutP proteins are novel acetate kinases
involved in ethanolamine catabolism: physiological implications for the function of the
ethanolamine metabolosome in Salmonella enterica. Mol Microbiol 99:497-511.

72.   Banla IL, Kommineni S, Hayward M, Rodrigues M, Palmer KL, Salzman NH, Kristich CJ. 2018.
Modulators of Enterococcus faecalis Cell Envelope Integrity and Antimicrobial Resistance
Influence Stable Colonization of the Mammalian Gastrointestinal Tract. Infect Immun 86.

73.   Chatterjee A, Willett JLE, Dunny GM, Duerkop BA. 2020. Phage infection mediates inhibition
of bystander bacteria. bioRxiv doi: 10.1101/2020.05.11.077669

74.   Qin X, Singh KV, Weinstock GM, Murray BE. 2000. Effects of Enterococcus faecalis fsr genes
on production of gelatinase and a serine protease and virulence. Infect Immun 68:2579-86.

75.   Bourgogne A, Hilsenbeck SG, Dunny GM, Murray BE. 2006. Comparison of OG1RF and an
isogenic fsrB deletion mutant by transcriptional analysis: the Fsr system of Enterococcus faecalis
is more than the activator of gelatinase and serine protease. J Bacteriol 188:2875-84.

76.   Teixeira N, Varahan S, Gorman MJ, Palmer KL, Zaidman-Remy A, Yokohata R, Nakayama J,
Hancock LE, Jacinto A, Gilmore MS, de Fatima Silva Lopes M. 2013. Drosophila host model
reveals new enterococcus faecalis quorum-sensing associated virulence factors. PLoS One
8:e64740.

77.   Kellogg SL, Kristich CJ. 2016. Functional Dissection of the CroRS Two-Component System
Required for Resistance to Cell Wall Stressors in Enterococcus faecalis. J Bacteriol 198:1326-36.

78.   Muller C, Le Breton Y, Morin T, Benachour A, Auffray Y, Rince A. 2006. The response
regulator CroR modulates expression of the secreted stress-induced SalB protein in Enterococcus
faecalis. J Bacteriol 188:2636-45.

79. Comenge Y, Quintiliani R, Jr., Li L, Dubost L, Brouard JP, Hugonnet JE, Arthur M. 2003. The
CroRS two-component regulatory system is required for intrinsic beta-lactam resistance in
Enterococcus faecalis. J Bacteriol 185:7184-92.
80. Paulsen IT, Banerjei L, Myers GS, Nelson KE, Seshadri R, Read TD, Fouts DE, Eisen JA, Gill
SR, Heidelberg JF, Tettelin H, Dodson RJ, Umayam L, Brinkac L, Beanan M, Daugherty S,
DeBoy RT, Durkin S, Kolonay J, Madupu R, Nelson W, Vamathevan J, Tran B, Upton J, Hansen
T, Shetty J, Khouri H, Utterback T, Radune D, Ketchum KA, Dougherty BA, Fraser CM. 2003.
Role of mobile DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. Science
299:2071-4.
81. Dunny GM, Brown BL, Clewell DB. 1978. Induced cell aggregation and mating in
Streptococcus faecalis: evidence for a bacterial sex pheromone. Proc Natl Acad Sci U S A
75:3479-83.
82. Morgan DJ, Aboud CJ, McCaffrey IM, Bhide SA, Lamont RF, Taylor-Robinson D. 1996.
Comparison of Gram-stained smears prepared from blind vaginal swabs with those obtained at
speculum examination for the assessment of vaginal flora. Br J Obstet Gynaecol 103:1105-8.
83. Deng L, Mu R, Weston TA, Spencer BL, Liles RP, Doran KS. 2018. Characterization of a Two-
Component System Transcriptional Regulator, LtdR, That Impacts Group B Streptococcal
Colonization and Disease. Infect Immun 86.
777
778
779

A. *E. faecalis* Adherence *in vitro*

■ V583 □ OG1RF

% Adherence

Vaginal   Endocervical

B. *E. faecalis* Vaginal Colonization

Recovered CFU (Log$_{10}$)

V583   OG1RF

C. V583 Colonization

Recovered CFU/g (Log$_{10}$)

Vagina   Cervix   Uterus

D.

LP

Lu

vagina (40x)

E.

EP   Lu

LP

cervix (40x)

F.

LP

Lu

EP

uterus (40x)

A.

Adherence to Vaginal Cells

B.

Adherence to Endocervical Cells

C.

Recovered CFU/Mouse

A

B

Table 1: Selected list of differentially abundant transposon mutants during vaginal colonization compared to *in vitro* cultures

old_locus_tag	NCBI_description	Difference (D1 - Cul)	Difference (D5 - Cul)	Difference (D8 - Cul)
Sortase-dependent proteins (SDPs)				
OG1RF_10811	collagen adhesion protein	-4.33	-4.14	-5.72
OG1RF_10871	cell wall surface anchor family protein, *ebpC*	-2.54	-2.37	-4.44
OG1RF_11531	glycosyl hydrolase	-3.04	-3.15	-4.25
OG1RF_11764	cell wall surface anchor family protein	-2.80	-3.48	-5.24
OG1RF_12054	cell wall surface anchor family protein	-2.62	-1.50	-6.01
Ethanolamine utilization				
OG1RF_11342	ethanolamine utilization protein EutL	-3.92	-3.25	-6.10
OG1RF_11343	ethanolamine ammonia-lyase small subunit	-4.09	-4.60	-5.85
OG1RF_11344	ethanolamine ammonia-lyase large subunit	-4.27	-4.28	-6.94
Type VII secretion system				
OG1RF_11103	YukD superfamily protein, *esaB*	-3.19	-5.02	-6.95
OG1RF_11109	putative LXG-containing toxin	-3.93	-3.47	-4.87
OG1RF_11111	putative T7SS toxin	-3.76	-4.15	-4.80
OG1RF_11113	putative T7SS toxin	-3.19	-2.87	-3.70
OG1RF_11114	putative immunity protein	-3.96	-3.88	-5.62
OG1RF_11122	immunity protein	-1.31	-1.43	-2.14