Transposable Elements activity and role in *Meloidogyne incognita* genome dynamics and adaptability

Djampa KL KOZLOWSKI, Rahim HASSANALY-GOULAMHOUSSEN, Martine DA-ROCHA, Marc BAILLY-BECHET*, Etienne GJ DANCHIN*.

* co-last authors
Affiliation : Université Côte d'Azur, INRAE, CNRS, ISA, Sophia Antipolis, France

Abstract

Despite reproducing without sexual recombination, the root-knot nematode *Meloidogyne incognita* is adaptive and versatile. Indeed, this species displays a global distribution, is able to parasitize a large range of plants and can overcome plant resistance in a few generations. The mechanisms underlying this adaptability without sex remain poorly known and only low variation at the single nucleotide polymorphism level have been observed so far across different geographical isolates with distinct ranges of compatible hosts. Hence, other mechanisms than the accumulation of point mutations are probably involved in the genomic dynamics and plasticity necessary for adaptability. Transposable elements (TEs), by their repetitive nature and mobility, can passively and actively impact the genome dynamics. This is particularly expected in polyploid hybrid genomes such as the one of *M. incognita*. Here, we have annotated the TE content of *M. incognita*, analyzed the statistical properties of this TE content, and used population genomics approach to estimate the mobility of these TEs across 12 geographical isolates, presenting phenotypic variations. The TE content is more abundant in DNA transposons and the distribution of sequence identity of TE occurences to their consensuses suggest they have been at least recently active. We have identified loci in the genome where the frequencies of presence of a TE showed variations across the different isolates. Compared to the *M. incognita* reference genome, we detected the insertion of some TEs either within coding regions or in the upstream regulatory regions. These predicted TEs insertions might thus have a functional impact. We validated by PCR the insertion of some of these TEs, confirming TE movements probably play a role in the genome plasticity with possible functional impacts.

Introduction

Agricultural pests cause substantial yield loss to the worldwide life-sustaining production (Savary et al. 2019) and threaten the survival of different communities in developing countries. With a constantly growing human population, it becomes more and more crucial to reduce the loss caused by these pests while limiting the impact on the environment. In this context, understanding how pests evolve and adapt both to the control methods deployed against them and to a changing environment is essential. In metazoa, nematodes and insects are the most destructive agricultural pests. Nematodes alone are responsible for

crop yield losses of ca. 11% which represents up to 100 billion € economic loss annually
(Agrios 2005; McCarter 2009). The most problematic nematodes to worldwide agriculture
belong to the genus Meloidogyne (Jones et al. 2013) and are commonly named root-knot
nematodes (RKN) owing to the gall symptoms their infection leaves on the roots. Curiously,
the RKN species showing the wider geographical distribution and infecting the broadest
diversity of plants reproduce asexually via mitotic parthenogenesis (Trudgill and Blok 2001;
Castagnone-Sereno and Danchin 2014). In the absence of sexual recombination, the
genomes are supposed to irreversibly accumulate deleterious mutations, the efficiency of
selection is reduced due to linkage between conflicting alleles while the combination of
beneficial alleles from different individuals is impossible (Muller 1964; Hill and Robertson
1966; Kondrashov 1988; Glémin and Galtier 2012). For these reasons asexual reproduction
is considered an evolutionary dead end and is actually quite rare in animals (Rice 2002). In
this perspective, the parasitic success of the parthenogenetic RKN might represent an
evolutionary paradox.
Previous comparative genomics analyses have shown the genomes of the most devastating
RKN are polyploid as a result of hybridization events (Blanc-Mathieu et al. 2017; Szitenberg
et al. 2017). In the model RKN *M. incognita*, the resulting gene copies not only diverge at the
nucleotide level but also in their expression patterns, suggesting this peculiar genome
structure could support a diversity of functions and might be involved in the parasitic success
despite the absence of sexual reproduction (Blanc-Mathieu et al. 2017). This hypothesis
seems consistent with the 'general purpose genotype' concept, which proposes successful
parthenogens have a generalist genotype with good fitness in a variety of environments
(Vrijenhoek and Parker 2009). An alternative non mutually exclusive hypothesis is the
'frozen niche variation' concept which proposes parthenogens are successful in stable
environments because they have a frozen genotype adapted to this specific environment
(Vrijenhoek and Parker 2009). Interestingly, the frequency of parthenogenetic invertebrates
is higher in agricultural pests, probably because the anthropized environments in which they
live are more stable and uniform (Hoffmann et al. 2008).
However, although a general purpose genotype brought by hybridization might partially
explain the wide host range and geographical distribution of these parthenogenetic RKNs,
this alone, cannot explain how these species evolve and adapt to new hosts or environments
without sex. For instance, initially, avirulent populations of some of these RKN, controlled by
a resistance gene in a tomato, are able to overcome the plant resistance in a few
generations, leading to virulent sub-populations, in controlled laboratory experiments
(Castagnone-Sereno et al. 1994; Castagnone-Sereno 2006). Emergence of virulent
populations, not controlled anymore by resistance genes have also been reported in the field
(Barbary et al. 2015).
The mechanisms underlying the adaptability of parthenogenetic RKN without sex remain
elusive. Recent population genomics analyses showed that only a few single nucleotide
variations (SNV) could be identified by comparing different Brazilian *M. incognita* isolates
showing distinct ranges of host compatibility (ie host races) (Koutsovoulos et al. 2020).
Addition of further isolates from different geographical locations across the world did not
substantially expand the number of variable positions in the genome. Furthermore, the few
identified SNV did not show correlation with either the geographical location, the host range
or the current crop species. However, these SNV could be used as markers to confirm the
absence of sexual meiotic recombination in *M. incognita*. Thus, the low nucleotide variability
that was observed between isolates is probably not the main player in the genomic plasticity
underlying the adaptability of *M. incognita*.

Consistent with these views, convergent gene copy number variations were observed
following resistance breaking down by two originally avirulent populations of *M. incognita*
from distinct geographic origins (Castagnone☐Sereno et al. 2019). The mechanisms
supporting these gene copy numbers and other genomic variations possibly involved in the
adaptive evolution of *M. incognita* remain to be described.
Transposable elements (TEs), by their repetitive and mobile nature, can both passively and
actively impact genome plasticity and stability. Being repetitive, they can be involved in
illegitimate genomic rearrangements leading to loss of genomic portions or expansion of
gene copy numbers. Being mobile, they can insert in coding or regulatory regions and have
a functional impact on the gene expression or gene structure itself. In some fungal
phytopathogens, TEs are a major player of adaptive genome evolution by both passively and
actively impacting the genome structure and sequence (Faino et al. 2016). In parallel,
although TE movements can provide beneficial 'novelty' / plasticity, their uncontrolled activity
can also be highly detrimental and put the organism at risk. Whether TEs also play an
important role in the adaptive evolution of animal genomes and particularly in parasites,
engaged in a continuous 'arms race' with their hosts, remains poorly known. According to
the Red Queen hypothesis, host-parasites arms race is a major justification for the
prevalence of otherwise costly sexual reproduction (Lively 2010) and, in the absence of sex,
other mechanisms should provide the necessary plasticity to sustain this arms race.
From an evolutionary point of view, the parthenogenetic root-knot nematode *M. incognita*
represents an interesting model to study the activity of TEs and their impact. Indeed, being a
plant parasite, *M. incognita* is engaged in an arms race with the plant defence systems and
point mutations alone are not expected to be a major mechanism supporting adaptation in
this species (Koutsovoulos et al. 2020).
In this study, we have tested whether the TE activity could represent a mechanism
supporting genome plasticity and eventually adaptive evolution in *M. incognita*. We have re-
annotated the 185Mb triploid genome of *M. incognita* (Blanc-Mathieu et al. 2017) for TEs,
using stringent filters to identify canonical TEs, possibly active in the genome. We analyzed
the statistical properties of the TE content and the distribution of TE sequence identity levels
to their consensuses was skewed towards high values, suggesting they might have
undergone recent multiplications in the genome. We have then tested whether the
frequencies of presence/absence of these TEs across the genome varied between different
isolates. To test for variations in frequencies, we have used population genomics data from
eleven *M. incognita* isolates collected on different crops and locations and differing in their
ranges of compatible hosts (Koutsovoulos et al. 2020). From the set of TE loci that
presented the most contrasted patterns of presence/absence across the isolates, we
investigated whether some could represent neo-insertions. To estimate the potential impact
of TE insertions, we checked whether some were inserted within coding or possible
regulatory regions. Finally, we validated some of the neo-insertions, predicted by population
genomics data by PCR assays. Overall, our study represents the first estimation of TE
activity as a mechanism possibly involved in the genome plasticity and the associated
functional impact in the most devastating nematode to worldwide agriculture. Because this
study focuses on an allopolyploid and parthenogenetic animal species, it also opens new
evolutionary perspectives on the fate and potential adaptive impact of TEs in these singular
organisms.
132

Results

The *M. incognita* TE landscape is diversified but mostly composed of DNA transposons.

We used the REPET pipeline (Quesneville et al. 2005; Flutre et al. 2011) to predict and annotate the *M. incognita* repeatome (cf. methods). Here, we define the repeatome as all the repeated sequences in the genome, excluding Simple Sequence Repeats (SSR or microsatellites). The repeatome spans 26.38 % of the *M. incognita* genome length. As we wanted to assess whether TEs actively contributed to genomic plasticity, we applied a series of stringent filters on the whole repeatome to retain only repetitive elements presenting canonical signatures of TEs (cf. methods). We identified 480 different TE-consensus sequences that allowed annotation of 9,702 canonical TE, spanning 4.73% of the genome. Both retro (Class I) and DNA (Class II) transposons (Wicker et al. 2007) compose the *M. incognita* TE landscape with 5/7 and 4/5 of the known TE orders represented respectively, showing a great diversity of elements (Fig 1). Retro-transposons and DNA-transposons respectively cover 0.94 and 3.78 % of the genome. TIR (Terminal Inverted Repeats) and MITEs (Miniature Inverted repeat Transposable Elements) DNA-transposons alone represent almost two-thirds of the *M. incognita* TE content (63.64 %). Hence, the *M. incognita* TE landscape is diversified but mostly composed of DNA-transposons.

As a technical validation of our annotation method, we used the same protocol to predict the *C. elegans* TE genomic content (sup. Fig 1 & sup. Table 1), using the PRJNA13758 assembly (The C. elegans Genome Sequencing Consortium 1998), and compared our results to the reference report of the TE landscape in this model nematode (Bessereau 2006). We estimated that the *C. elegans* repeatome spans 11.81% of its genome, which is close to the 12 % described in (Bessereau 2006). The same resource also reported that MITEs and LTR respectively compose ~2% and 0.4% of the *C. elegans* genomes while we predicted 1.8% and 0.2%. Predictions obtained using our protocol are thus in the range of previous predictions for *C. elegans*. In the same study, it was mentioned that most of *C. elegans* TE sequences "are fossil remnants that are no longer mobile", and that active TEs are DNA transposons. This suggests a stringent filtering process is necessary to isolate TEs that are the most likely to be active (e.g. the 'canonical' ones). Using the same post-processing protocol as for *M. incognita*, we estimated that canonical TEs span 3.99% of the *C. elegans* genome and DNA-transposon alone span 3.13% of this genome.

Overall, the similarity of our results with the previous reports in *C. elegans* suggests our filtered annotation of the TE content of *M. incognita* represents an accurate picture of the potentially active TE landscape.

169
170
171 **Fig 1: Canonical TE annotations distribution in *M. incognita* genome**
172 CLASS_1_LIKE and CLASS_2_LIKE elements present sufficient evidence to class them as
173 Retro or DNA transposons respectively, but not enough to further assign them to an order.
174 Genome percentage is based on a M. incognita genome size of 183,531,997 bp
175
176
177 **Table 1: Per-order summary of *M. incognita* TE annotations.**

	order	nb. of features	total length (bp)	genome percentage (%)	median length (bp)	median of median identity with consensus (%)
Retro-transposon	CLASS_1_LIKE	21	81,609	0.044	3,871	97.6
	LARD	45	63,42	0.035	1,433	97.05
	LINE	145	313,224	0.171	1,971	96.6
	LTR	373	1,164,836	0.635	2,415	97
	SINE	9	4,522	0.002	528	99.7
	TRIM	174	104,018	0.057	525	97.7
	Total	**767**	**1,731,629**	**0.944**		

	CLASS_2_LIKE	48	23,132	0.013	508.5	98.9
	HELITRON	18	86,666	0.047	5,080	94.4
DNA-transposon	MAVERICK	189	1,307,068	0.712	6,224	95.3
	MITE	5085	2,755,381	1.501	525	96.2
	TIR	3595	2,777,270	1.513	737	97.3
	Total	**8935**	**6,949,517**	**3.787**		

Canonical TE annotations are highly identical to their consensus sequences and some present evidence for transposition machinery.

Canonical TE annotations have a median nucleotide identity of 97.12% with their respective consensus sequences, but the distribution of identity values varies between TE orders (Fig 2, Table 1). Most of the TEs within an order share a high identity level with their consensuses, except HELITRON and MAVERICK. Even considering our inclusion threshold at minimum 85% identity (cf methods), the overall distribution of average % identities peaks at high values. DNA transposons show a wider dispersion in identity to their consensuses than retrotransposons, as indicated by the elongated boxes in Fig 2B and this distribution peaks at higher values. As a consequence, it is among those elements that annotations sharing a higher similarity to their consensus sequences are found. In particular among TIR, one fourth (Fig 2; sup. Table 2) of the annotations share above 99% identity with their consensus. SINE and CLASS_2_LIKE have similar profiles but are present in very low numbers.

Fig2: **per-copy identity rate with consensus**
Top frequency plots show the distribution of TE copies count per order in function of the identity % they share with their consensus sequence. To facilitate inter-orders comparison, bottom violin plots display the same information as a density curve, but also encompass boxplots. Each colour is specific to a TE order.

Higher identity of TE annotations to their consensus can be considered a proxy of their
recent activity (Bast et al. 2015; Lerat et al. 2019). To further investigate whether some TEs
might be (or have been recently) active, we searched for the presence of genes involved in
the transposition machinery within *M. incognita* canonical TEs (cf methods). Among the
canonical TE annotations, 6.26% (607/9,702) contain at least one predicted protein-coding
gene, with a total of 907 genes. Of these 907 genes, 328 code for proteins with at least one
conserved domain known to be related to transposition machinery. We found that 35.37%
(116/328) of the transposition machinery genes had substantial expression support from
RNA-seq data. In total, 111 canonical TE-annotations contain at least one substantially
expressed transposition machinery gene (see supplementary material 1). These 111 TE
annotations correspond to 41 different TE-consensuses, and as expected, only consensuses
from the autonomous TE orders, e.g. LTRs, LINEs, TIRs, HELITRON, and MAVERICKs
present TE-copies with substantially expressed genes coding for transposition machinery.
Also as expected, the non-autonomous TEs do not contain any transposition machinery
gene at all. This suggests that some of the detected TEs might have functional transposition
machinery, which in turn could be hijacked by the non-autonomous elements.

Overall, the presence of a substantial proportion of TE annotations highly similar to their
consensuses combined with the presence of genes coding for the transposition machinery
and supported by expression data suggest some TE might be active in the genome of *M.*
incognita.

Thousands of loci show variations in TE presence frequencies across *M. incognita* isolates.

We used the PopoolationTE2 (Kofler et al. 2016) pipeline (joint algorithm) on the *M.*
incognita reference genome (Blanc-Mathieu et al. 2017) and the canonical TE annotation to
detect variations in TE frequencies across the genome between 12 geographical isolates (cf.
methods). One isolate comes from Morelos in Mexico, which is the isolate that was used to
produce the *M. incognita* reference genome. The 11 other isolates come from different
locations across Brazil, and present different ranges of compatible hosts (referred to as R1,
R2, R3, R4, see sup. Fig 2) currently infected crop species (Koutsovoulos et al. 2020). Pool-
seq paired-end Illumina data has been generated for all these isolates. For each locus, each
isolate has an associated frequency value representing the proportion of individuals in the
pool having the TE detected at this location.

We identified 3,524 loci where the frequency variation between at least two isolates was
higher than our estimated PopoolationTE2 error rate (0.00972 i.e less than 1%, see
methods).

Overall, the distribution of frequencies is bimodal (Fig 3-A), and this pattern is common to all
the isolates, including the reference Morelos isolate (Fig 3-B). On average per isolate, 21.1%
of the loci have frequencies < 25%, 60.7% have frequencies > 75%, and only 18.2% show
intermediate frequencies Hence, in every isolate, most of the TE frequency values pack
around extreme values e.g. <25% or >75%.

Nevertheless, these statistics provide no information about the frequency variability between
isolates for a given locus. To address this question, for each locus, we computed the
absolute maximum frequency difference between isolates (Fig 3-C). We found that the
maximum frequency variation across the isolates is smaller than 20% in 75% of the loci
(2,643/3,524). Hence, most of the loci show little to moderate variations in frequencies
between isolates. Combined to the previous result, this implies that for most loci, the TEs are
present either at a high or a low frequency among all isolates. However, some TE loci show
more contrasted variations and will be the focus of further studies in our pipeline.

Fig. 3: TE frequency distribution.

The histogram (A) and violin plot (B) represent the TE frequency distribution per isolate. The
colour chart is identical between the two figures. Both representations reveal that in all the
isolates, only a few TE are found with intermediate frequencies. Right boxplot (C) represents
the frequency absolute maximum difference per locus. For a given locus, it illustrates the
frequency variability between isolates. The higher is the value; the more important is the
frequency difference between at least two isolates. A value of 1 implies that the TE is absent
in at least one isolate while it is present in 100% of the individuals of at least another isolate.

Variations of TE frequencies across isolates recapitulate their divergence at the sequence level

We performed a Neighbour-joining phylogenetic analysis of *M. incognita* isolates based on a
distance matrix constructed from TE frequencies (cf methods). We then compared the

obtained phylogenetic tree to the phylogenetic tree based on SNPs in coding regions
performed as part of previous analysis (Koutsovoulos et al. 2020).
The TE-based and SNP-based tree topologies are almost identical. In particular, the two
trees allowed defining three highly supported clades, with maximum support value (Fig.4).
Clades (2) and (3) were identical, including branching orders. Because clade (1) is a
polytomy in the SNP tree (A), relative branching of the three isolates cannot be compared
although clustering remains unchanged. In both trees, isolates R2-1 and R2-6 are outside of
the clusters but their relative positions differ. Similarly, although R1-6 is more closely related
to clusters (1) and (2) than the rest of the isolates in the two trees, its position also slightly
differs between the SNP-based (A) and TE-based (B) trees. Differences in relative positions
of R2-1 and R2-6 can be explained by the low distance observed between these two isolates
in the SNP-based analysis (version with branch length in sup. Fig 3).

Overall, the similarity between the SNP-based and TE frequency-based trees indicates that
most of the phylogenetic signal coming from variations in TE-frequencies between isolates
recapitulates the SNP-based genomic divergence between isolates.
281

282
283
Fig 4: Phylogenetic tree for *M. incognita* isolates.
A- Phylogenetic tree based on SNP present in coding sequences. Maximum Likelihood (ML) tree
reconstruction. Branch length not displayed.
B- Phylogenetic tree based on TE-frequencies euclidean distances between isolates. Neighbor-
Joining (NJ) tree reconstruction. Branch length not displayed. In both trees, bootstrap support
values are indicated on the branches. Isolates enclosed in the dashed area form a super-
cluster composed of the clusters (1) and (2), and the isolate R1-6. Clades with bootstrap
support values <= 20 were collapsed and represented as a polytomy.

Some loci with TE frequency variations across isolates correspond to neo-insertions.

As explained below (and cf methods), we categorized all the loci with TE frequency variations across the isolates by (i) comparing their position to the TE annotation in the reference genome, (ii) analysing TE frequency in the reference isolate Morelos, (iii) comparing TE-frequencies detected for each isolate to the reference isolate Morelos. This allowed defining, on the one hand, non-polymorphic and hence stable reference annotation, and on the other hand, 3 categories of polymorphic (variable) loci (see Fig 5).

Fig 5: Categories of polymorphic TE loci

Orange boxes illustrate the presence of a TE at this position in the reference genome annotation. Purple boxes illustrate the percentage of individuals in the isolates for which the TE is present at this position (i.e. frequency). Frequencies values are reported as colour gradients. A - non-polymorphic TE locus: a TE is predicted in the reference annotation (orange box), is detected in all the isolates (purple box), but the presence frequency does

not vary substantially between the isolates (frequency variation < 1% among all the isolates).
B - polymorphic reference locus: a TE is predicted in the reference annotation, is detected in
the reference isolate Morelos with a frequency > 75%, and the presence frequency varies
between the isolates (frequency variation >= 1% among all the isolates). This category
encompasses 2,096 loci. C - extra-detection: no TE is predicted at this locus in the reference
annotation but one is detected at a frequency >25% in the reference isolate Morelos, and
optionally in other isolates. This category counts 210 loci. D - neo-insertion: no TE is
predicted at this locus in the reference genome annotation and none is detected in the
reference isolate (dashed box, frequency < 1%), but a TE is detected in at least one other
isolate with a frequency >= 25%. This category counts 287 loci.

Only 931 loci could not be assigned to the categories in Fig 5 and were discarded.
Uncategorised loci are cases where all the isolates (Morelos included) show frequencies <
25% or ambiguous cases with a low Morelos frequency (between 1% and 25%) and at least
one isolate showing a frequency > 25%.

Overall, 73.6% (2,593/3,524) of the loci with TE frequency variations could be assigned to
one of the 3 categories of TE-polymorphisms (B, C, D in Fig 5) and the decomposition per
TE order is given in Fig 6 and sup. Table 3.

The vast majority of the polymorphic loci (80.83 %; 2,096/2,593) corresponds to an already
existing TE-annotation in the reference genome and the corresponding TE is fixed
(frequency > 75 %) at least in the reference isolate Morelos but varies in at least another
isolate. These polymorphic loci cover ~21.6% (2,096/9,702) of the canonical TE annotations,
in total. These loci will be referred to as 'polymorphic reference loci' from now on (Fig 5B)
and they encompass both DNA- and Retro-transposons.

Then, we considered as 'neo-insertion' TEs present at a frequency >25% in at least one
isolate at a locus where no TE was annotated in the reference genome and the frequency of
TE presence was < error rate (~1%) in the reference Morelos isolate (Fig 5D). In total, 11.07
% (287/2,593) of the detected TE polymorphisms correspond to such neo-insertions. It
should be noted here that we consider neo-insertions as regard to the reference Morelos
isolate only and some of these so-called neo-insertions might represent TE loss in Morelos.
Comparison with the phylogenetic pattern of presence / absence will allow distinguishing
further the most parsimonious of these two possibilities.

Finally, we classified as 'extra-detection' (Fig 5C) (8.10%; 210/2,593) the loci where no TE
was initially annotated by REPET in the reference genome, but a TE was detected at a
frequency >25% at least in the ref isolate Morelos by PopoolationTE2. It should be noted
that 57.62% (121/210) of these loci correspond to draft annotations that have been
discarded during the filtering process to only select the canonical annotations. These draft
annotations might represent truncated or diverged versions of TE that exist in a more
canonical version in another locus in the genome. Half of the remaining 'extra-detections'
(45/89) are detected with low to moderate frequency (<43.5%) in the reference isolate
Morelos. We hypothesise that because they represent the minority form, these regions were
not taken into account during the assembly of the genome. This would explain why these
TEs could not be detected in the genome assembly by REPET (assembly-based approach)
but were identified with a read mapping approach on the genome plus repearome by

PopoolationTE2. The remaining 'extra-detections' might correspond to REPET false
negatives, PopoolationTE false positives, or a combination of the two. Nonetheless, we can
notice these cases only represent 1.69% (44/2,593) of the detected polymorphic TEs. Loci of
variable frequency that could not be assigned to the 3 above-mentioned B, C, D categories
(26.4% of them) were discarded from the rest of the analysis.

357 TIR and MITEs elements are overrepresented among TE-
358 polymorphisms.

359 By themselves, MITEs and TIRs elements encompass 94.25% (2,444/2,593) of the
360 categorized TE-polymorphisms (Fig 6).
361 We showed that the polymorphism distribution varies significantly between the four
362 categories presented in Fig. 5 (Chi-square test, p-value < 2.2e-16), indicating that some TE
363 orders are characterised by specific polymorphisms types.
364

365
366
367 **Fig 6: TE polymorphisms count per orders and types.**
368 Top left barplot shows TE polymorphisms distribution per type and per order. Bottom-left
369 barplot summarizes TE polymorphisms distribution per type. In both barplots, the values in
370 black represent the count per polymorphism type. Top-right barplot illustrates the total
371 number of polymorphisms per order. Orders are sorted identically in both plots.

372
373

The analysis of the chi-square residuals (sup. Fig 4) shows MITEs and TIRs are the only two
orders presenting a relative lack of non-polymorphic TEs. Hence, in addition to being the
most abundant in the genome, these two TE orders are significantly enriched among
polymorphic loci. MITEs are over-represented in both TE polymorphisms types (polymorphic
ref. loci and neo-insertions, Fig5 B and D), suggesting a variety of activities within this order.
On the other hand, TIRs are found in excess in ref-polymorphisms but lack in neo-insertions.
This lack of neo-insertions in TIRs may indicate a lower activity in this order, or a more
efficient negative selection.
Finally, we observed a strong excess of Maverick among the extra-detection as almost 70%
of Maverick polymorphisms (16/23) (Fig 6) fell into this category. Consistent with the
observation that, globally, >50% of the extra detections were actually draft annotations
eliminated afterwards during filtering steps, ¾ (12/16) of these Maverick elements were also
actually present in the draft annotations but eliminated during filtering and thus only appear
here due to the stringency of our filtering.

Overall, in proportion, MITEs and TIRs elements are significantly over-represented in TE-
polymorphisms. This observation suggests TEs from MITEs and TIRs orders, in addition to
being the most numerous canonical TEs, might have been more active in the genome of *M.*
incognita than elements from other TE-orders.

393 Some polymorphic loci showing contrasted frequency variations
394 between isolates represent neo-insertions.

We investigated the variability of TE presence frequency per locus over the 12 isolates for all
the categorized polymorphic loci in the genome.
In ~ 3/4 (1,919/2,593) of the categorized polymorphic TE loci, the TE presence frequency is
homogeneous between isolates (cf methods). Said differently, it means that although we
observe variations in frequencies above the estimated error rate (<1%) between isolates,
these variations remain at low amplitude (maximum frequency variation between isolates
<=25% for a given locus). The vast majority (97.97%; 1,880/1,919) concerns loci where the
TE is present at a high frequency in all isolates (> 75%). These loci might be considered as
fixed in all the isolates. In the remaining 2.03% (39/1,919), the TE frequency is either
between 25 and 50% or between 50 and 75% in all isolates. As expected given our
methodology, all the high-frequency loci correspond to ref-polymorphisms while all the
intermediate frequency loci belong to extra-detections. Also, we did not detect loci where the
TE was present with low frequency (<25%) in all isolates as they did not meet the
categorisation criteria (1% < freq <= 25 in Morelos isolate).
In the 674 remaining polymorphic TE loci, TE frequency is heterogeneous, meaning the
frequency difference between at least two isolates is > 25%. Among the most extreme cases
of frequency variation per locus, we identified 33 loci for which the presence of the TE is
found with high frequencies (> 75%) for some isolate(s) while it is absent or rare (frequency
<25 %) in the other(s). These loci will be from now on referred to as HCPTEs standing for
"Highly Contrasted Polymorphic TE" loci. Because they are highly contrasted, these loci
might represent differential fixation/loss across isolates and will be the focus of the following
analyses.

HCPTEs encompass 19 MITEs elements, 12 TIRs and 2 LINEs. We can also notice that
some consensuses are more involved in HCPTEs as 4 TE consensuses are responsible for
72.72% (24/33) of these polymorphisms.
Interestingly, all the HCPTEs loci correspond to neo-insertions regarding the reference
genome, meaning that no TE was annotated in the reference genome at this location and
the TE presence frequency is < 1% in the Morelos reference isolate. As described in Fig. 7,
most of these fixed neo-insertions (20/33) are specific to an isolate. However, we also found
neo-insertions shared by 2 (10/33), 3 (2/33) or even 6 isolates (1/33).
Interestingly, all the shared neo-insertions were between isolates present in a same cluster
in the phylogenetic trees (TE-based and SNP-based in Fig. 4), suggesting they might have
been fixed in a common ancestor and then inherited. For example, two neo-insertions are
shared by isolates R4-4, R1-2 and R3-2 which belong to the same cluster 1 and one neo-
insertion is shared by isolates R4-3 and R1-3 which belong to the same cluster 2. Even the
neo-insertion shared by 6 isolates follows this pattern as all the concerned isolates belong to
the same super-cluster composed of the cluster 2 and 3 (dashed line in Fig 4).
Hence, the phylogenetic distribution reinforces the idea that these cases are more likely to
represent branch-specific neo-insertions than multiple independent losses, including in the
reference isolate Morelos.
R1-2, R3-2, and R4-4 show the highest number of neo-insertions among isolates. However,
their profiles are quite different. In 10/12 HCPTEs involving R1-2, the TE is present only in
this isolate while most of the HCPTEs involving R3-2 and R4-4 are neo-insertions shared
with close isolates. This may be related to the relative phylogenetic divergence of those
isolates (sup Fig 3), which shows that R1-2 is the most divergent isolate, while R3-2 is quite
close to its neighbour, and especially to R4-4.

Fig 7: HCPTEs Neo-insertions specificity among the isolates.
The central plot shows how many and which isolate(s) share common HCPTEs neo-insertion(s), every line representing an isolate. Columns with several dots linked by a line indicate shared HCPTEs neo-insertion(s) between isolates. Each dot represents which isolate is involved. Columns with a single dot design isolate-specific HCPTEs neo-insertion(s). The top bar plot indicates how many HCPTEs neo-insertions the corresponding group of isolate shares. The left side barplot specifies how many HCPTEs neo-insertion(s) occurred in a given isolate.

Functional impact of TE neo-insertion and validation of in silico predictions

Interestingly, the vast majority (22/33) of the fixed HCPTEs are inserted inside a gene or in a possible regulatory region (i.e. 1 kb region upstream of a gene). These fixed neo-insertions might have a functional impact in *M. incognita*. Overall, 27 different protein-coding genes are possibly impacted by the 22 neo-insertions, some genes being in the opposite direction at a neo-insertion point (overlapping this insertion point or being at max 1kb downstream). More than 80% of these genes (22/27) show a substantial expression level during at least one life stage of the nematode life cycle, suggesting the impacted genes are functional in the *M. incognita* genome (cf. methods). Some of the impacted genes (37.04%, 10/27) are specific to the *Meloidogyne* genus. Despite being all conserved in multiple Meloidogyne species, reinforcing their importance in the genus, they have no predicted orthologs in other nematodes. Among the remaining genes, one is specific to *M. incognita* (even other Meloidogyne species have no ortholog). Another one is present in multiple *Meloidogyne* species and otherwise only found in other Plant Parasitic Nematodes species (PPN) (*Ditylenchus destructor, Globodera rostochiensis*) (see sup. Table 4). Conservation of these genes across multiple PPN but exclusion from the rest of the nematodes suggest these genes might be involved in important functions relative to these organisms' lifestyle, including plant parasitism itself.

To experimentally validate in-silico predictions of TE neo-insertions with potential functional impact, we performed PCR experiments on 5 of the 24 HCPTEs loci falling in coding or possible regulatory regions. To perform these PCR validations, we used the DNA remaining from previous extractions performed on the *M. incognita* isolates for population genomics analysis (Koutsovoulos et al. 2020). Basically, the principle was to validate whether the highly contrasted frequencies (>75% / <25%) obtained by PopoolationTE2 actually corresponded to absence/presence of a TE at the locus under consideration (cf methods). One isolate (R3-1) presented no amplification in any of the tested loci nor in the positive control. After testing the DNA concentration in the sample, we concluded that the DNA quantity was limiting in this isolate and decided to discard it from the analysis.

For four of the five tested HCPTEs loci, we could validate by PCR the differential presence/absence of a sequence at this position, predicted by PopoolationTE2 across the different isolates (Fig 8; supplementary material 4).

In one of the five tested loci, named locus 1, we could i) validate by PCR the presence of a sequence at this position for the isolates presenting a PopoolationTE2 frequency >75% and absence for those having a frequency <25%; ii) also validate by sequencing that the sequence itself corresponded to the TE (a MITE) under consideration. This case is further explained in detail below and in Fig. 8.

According to PopoolationTE2 frequencies, the locus 1 MITE is inserted and fixed in 3
isolates (R1-2, R3-2, R4-4) as the estimated frequencies are higher than 75% in these
isolates. We assumed the TE is absent from the rest of the isolates as all of them display
frequencies <5%. To validate this differential presence across the isolates, we designed
specific primers from each side of the estimated insertion point so that the amplicon should
measure 973 bp with the TE insertion and 180 bp without.
The PCR results are coherent with the frequency predictions as only R1-2, R3-2, and R4-4
display a ~1 kb amplicon while all the other isolates show a ~0.2 kb amplicon (Fig 8). Hence,
as expected, only the 3 isolates with a predicted TE frequency >75% at this locus exhibit a
longer region, compatible with the MITE insertion.
To make sure the amplified regions corresponded to the expected MITE, we sequenced the
amplicons for the 3 predicted insertions and aligned the sequences to the TE consensus and
the genomic region surrounding the estimated insertion point (see supplementary material
4). R-1_2, R-3_2, and R-4_4 amplicon sequences all covered a significant part of the TE
consensus sequence length (> 78%) with high % identity (> 87%) and only a few gaps
(<5%). These results confirm that the inserted sequence corresponds to the predicted TE
consensus. Moreover, all the 3 amplicons aligned on the genomic region downstream of the
insertion point with high % identity (>= 99%), which helped us to further determine the real
position of the insertion point. The real insertion point is 26 bp upstream of the one predicted
by PopoolationTE2 and falls in the forward primer sequence. This explains why the amplicon
sequences do not align on the region upstream the insertion point.
We also noticed that the inserted TE sequences slightly diverged between the isolates while
the genomic region surrounding the insertion point remains identical. Interestingly, the level
of divergence in the TE sequence does not follow the phylogeny as R-4_4 is closer to R-1_2
than to R-3_2 (see sup. Table 5).
Finally, in the Morelos isolate, as well as R-2_1, and R-2_6 isolates, the sequencing of the
amplicon validated the absence of insertions as the sequences aligned on the genomic
region surrounding the insertion point with high % identity (99, 97, 87 % respectively) but not
with the MITE consensus.
Hence, we fully validated experimentally the presence/absence profile across isolates
predicted in silico at this locus.
In the *M. incognita* genome, this neo-insertion is predicted to occur in the 3' UTR region of a
gene (Minc3s00026g01668). This gene has no obvious predicted function, as no conserved
protein domain is detected and no homology to another protein with an annotated function
could be found. However, orthologs were found in the genomes of several other
Meloidogyne species (*M. arenaria*, *M. javanica*, *M. floridensis*, *M. enterolobii*, and *M.*
graminicola), ruling out the possibility that this gene results from a prediction error from gene
calling software. The broad conservation of this gene in the *Meloidogyne* genus suggests
this gene might be important for *Meloidogyne* biology and survival.
In the Morelos isolate, for which no TE was inserted at this position, this gene is substantially
supported by transcriptomic RNA-seq data during the whole life cycle of the nematode (see
supplementary material 1), suggesting this gene is probably functionally important in *M.*
incognita and other root-knot nematodes. Consequently, the insertion of the TE in R-1_2, R-
3_2, and R-4_4 genome at this locus could have functional impacts.
531
532
533

A

B

535
536
Fig 8: Experimental validation of a predicted neo-insertion.
A- Diagram of the TE neo-insertion. The neo-insertion of the MITE element occurs in the
3'UTR region of the gene (Minc3s00026g01668). Blue boxes illustrate the 3' and 5' UTR
regions of the gene while the yellow boxes picture the exons. Green arrows represent the
primers used to amplify the region. Gene subparts and TE representations are not at scale.
Predicted size of the amplicon: 973 bp with the TE insertion, 180 bp without.
543
544 B- PCR validation of the TE neo-insertion. Estimated freq. values correspond to the
545 proportion of individuals per isolate predicted to have the TE at this position
546 (PopoolationTE2). Isolates in red were predicted to have the TE inserted at this locus. Only
547 these isolates show an amplicon with a size suggesting an insertion. See supplementary
548 material 4 for sequences.

Discussion

The role of TE in genome plasticity and adaptive evolution of root-knot nematodes

M. incognita is a parthenogenetic mitotic nematode with a major agronomic impact. How this pest adapts to its environment in the absence of sexual recombination remains unresolved. In this study, we investigated whether TE movements could constitute a mechanism of genome plasticity compatible with adaptive evolution.

In *M. javanica*, a closely related root-knot nematode, comparison between an avirulent line unable to infest tomato plants carrying a nematode resistance gene and another virulent line that overcame this resistance, led to the identification of a gene present in the avirulent nematodes but absent from the virulent ones. Interestingly, the gene under consideration is present in a TIR-like DNA transposon and its absence in the virulent line suggests this is due to excision of the transposon and thus that TE activity plays a role in *M. javanica* adaptive evolution (Gross and Williamson 2011). However, this is so far the sole report and no large-scale analysis of the global role of TE in the adaptive evolution of root-knot nematodes has been performed.

In *M. incognita*, convergent gene losses at the whole genome level between two virulent populations compared to their avirulent populations of origin were recently reported (Castagnone☐Sereno et al. 2019). However, although TE activity might be involved in these convergent gene losses, this has never been shown.

Hence, the importance of TE activity both in the genome plasticity and adaptive evolution of the root-knot nematodes has never been assessed so far.

More broadly, except for *C. elegans* (Bessereau 2006; Laricchia et al. 2017), little to nothing is known yet about the TE dynamics and its impacts on nematode genomes and possibly in their adaptive evolution.

Here, we analysed the *M. incognita* TE dynamics using TE presence frequencies variations between isolates as a reporter of TE activity. We also assessed whether TE activity could have a functional impact, a necessary prerequisite for natural selection and adaptive evolution.

Our results established that TEs were likely recently active in *M. incognita* and that thousands of loci in the genome show substantial variations in the frequency of reads supporting a TE at this position, across different geographical isolates. It should be noted here that the total impact of TE on the genome dynamics is probably underestimated, in part because of our strategy to eliminate false positives as much as possible by applying a series of stringent filters, in another part because of the intrinsic limitations of the tools, such as the incapacity of PopoolationTE2 to detect nested TEs (Kofler et al. 2016). Despite this, we identified highly contrasted TE loci across the populations and some of these represented neo-insertions of TE, specifically in some isolates or some branches ancestral to several isolates. Because for certain TE, the insertion took place within a coding sequence or in a possible regulatory region, these insertions have a potential impact either by disrupting the encoded protein or modifying the gene expression pattern.

We confirmed insertions of the TEs for some of these functionally-important regions by PCR and thus a possible functional impact. However, this functional impact itself would need to be evaluated in the future either by generating transcriptomics data for the different isolates

enabling study of difference in gene expression patterns or transcript length or by proteomic
studies to directly search for differences at the encoded protein level.
Regardless of the future experimental validation of the functional impact, one important
question concerns the current preliminary evidence for a possible role in the nematode
adaptive evolution. Because some of the impacted genes are specific to plant-parasitic
species and yet conserved in several of these phytoparasites, a role in plant parasitism is
possible. However, in the absence of known protein domains or functional characterization
of these genes, the exact biochemical activity or biological processes in which they might be
involved is unknown. Furthermore, for the functional impact, we have exclusively considered
neo-insertions as the most evident cases. Some TE showing less contrasted changes in
frequencies in coding or possible regulatory regions might be associated to genes with
characterized functions but as these cases are less clear cut, they were not considered.
It should be noted that *M. incognita* is a relatively recent model with a first version of the
genome available only since 2008 (Abad et al. 2008) and only up to a dozen of genes
functionally characterized. In comparison, the model nematode *C. elegans* was the first
animal genome to be sequenced in 1998 (The C. elegans Genome Sequencing Consortium
1998) and as early as 2003, more than 85% of the genes had already been inactivated one
by one via RNAi to monitor the effect on the worm phenotype (Kamath et al. 2003).
Therefore, although no evident role in adaptive evolution for the *M. incognita* genes
impacted by TE insertions could be reported so far, future functional characterization might
bring more evidence.
Taking into account all the TE loci and regardless of the potential functional impact on
genes, we found that the pattern of variations of TE frequencies across the loci between the
different populations recapitulated almost exactly the phylogeny of the isolates built on SNP
in coding regions. Thus, it seems that most of the divergence in terms of TE pattern follows
the divergence at the nucleotide level and thus the phylogeny of the isolates. Almost the
same conclusion was drawn by comparing SNV data to TE variation data across different *C.
elegans* populations (Laricchia et al. 2017). In *M. incognita*, the phylogeny of isolates does
not significantly correlate with the biological traits that had been surveyed, namely
geographical distribution, range of compatible host plants and nature of the crop currently
infected (Koutsovoulos et al. 2020). Interestingly, no correlation was also observed between
variations in TE frequencies and geographical distribution for European Drosophila
populations (Lerat et al. 2019). As for the other traits considered here (range of compatible
host plants and nature of the plant infected), the lack of evident correlation between those
traits and phylogenetic signals regardless whether it is TE-based or SNV-based suggest that
most of the variations follow the drift between isolates and are not necessarily adaptive,
which is not surprising.
It should be noted that, in terms of functional impact, we have so far only considered the
active role of TE and not analyzed yet their possible passive roles. Indeed, being repetitive,
TE can be involved in illegitimate recombination events, and also generate loops in the DNA
which can eventually be excised and lead to gene loss. In parallel, genes can be hitchicked
by TE and multiplied in the genome. These different properties can clearly impact gene copy
number variations (CNV). CNV are known to be involved in genomic plasticity and in
adaptive evolution (Katju and Bergthorsson 2013). As stated above, convergent gene CNV
have been associated with plant resistance breaking down by *M. incognita*, although neither
direct functional link between these CNV and mechanisms of resistance breaking down nor
evidence for involvement of TEs have been shown so far (Castagnone⬜Sereno et al. 2019).
Actually, these analyses have been done in a previous version of the *M. incognita* genome

assembly which was partially incomplete (Abad et al. 2008) in comparison to the most recent
available genome assembly (Blanc-Mathieu et al. 2017). Reinvestigating CNV and the
possible involvement of TEs in association to an adaptive process such as resistance
breaking down on this more complete and more recent assembly would be interesting.
However, the current genome is still fragmentary and limits structural studies such as the
identification of TE-rich vs. TE-poor genome regions in possible association with CNV loci.
Future more contiguous versions of the genomes of root-knot nematodes will undoubtedly
enable such interesting perspectives to be undertaken.
649

TE-load and composition in a clonal allopolyploid species

M. incognita is an asexual (mitotic parthenogenetic), polyploid, and hybrid species. These
three features are expected to impact TE load in the genome with various intensities and
possibly conflicting effects.
Contradictory theories exist concerning the activity/proliferation of TEs as a function of the
reproductive mode. The higher efficacy of selection under sexual reproduction can be
viewed as an efficient system to purge TEs and control their proliferation. Supporting these
views, in parasitoid wasps, TE load was shown to be higher in asexual lineages induced by
the endosymbiotic Wolbachia bacteria than in sexual lineages (Kraaijeveld et al. 2012).
However, whether this higher load is a consequence of the shift in reproductive mode or of
Wolbachia infection remains to be clarified.
In an opposite theory, sexual reproduction can also be considered as a way for TEs to
spread across individuals within the population whereas in clonal reproduction the
transposons are trapped exclusively in the offspring of the holding individual. Under this
view, asexual reproduction is predicted to reduce TE load as TE are unable to spread in
other individuals, and are thus are removed by genetic drift and/or purifying selection in the
long term (Wright and Finnegan 2001). Consistent with this theory, comparison of sexual
and asexual *Saccharomyces cerevisiae* populations showed that the TE loads decrease
rapidly under asexual reproduction (Bast et al. 2019).
Hence, whether the TE-load is expected to be higher or lower in species with clonal vs.
sexual reproduction remains unclear and other conflicting factors such as TE excision rate
and the effective size of the population probably blur the signal (Glémin et al. 2019).
Interestingly, at a broader scale, a comparative analysis of different lineages of sexual and
asexual arthropods revealed no evidence for differences in TE load according to the
reproductive modes (Bast et al. 2015). Similar conclusions were drawn for nematodes
(Szitenberg et al. 2016), although only one asexually-reproducing species was present in the
comparative analysis.
Polyploidy, in contrast, is commonly accepted as a major event initially favouring the
multiplication and activity of TEs. This is clearly described with numerous examples in plants
(Vicient and Casacuberta 2017) and some examples are also emerging in animals
(Rodriguez and Arkhipova 2018). When hybridization and polyploidy are combined, this can
lead to TE bursts in the genome. As proposed by Barbara McClintock, allopolyploidization
produces a "genomic shock", a genome instability associated with the relaxation of the TE
silencing mechanisms and the reactivation of ancient TEs (McClintock 1984; Mhiri et al.
2019).

Hybridization, polyploidy and asexual reproduction are combined in *M. incognita* with relative
effects on the TE load extremely challenging, if not impossible, to disentangle. Initial
comparisons of the TE loads in three allopolyploid clonal *Meloidogyne* against a diploid
facultative sexual relative suggested a higher TE load in the clonal species (Blanc-Mathieu
et al. 2017). However, to differentiate the relative contribution of each of these three features
to the *M. incognita* TE load, it would be necessary to conduct comparative analysis with a
same method on diploid asexuals, on polyploid sexuals as well as on diploid asexuals in the
genus *Meloidogyne,* and ideally with and without hybrid origin. So far, genomic sequences
are only available for other polyploid clonal species, which are all suspected to have a hybrid
origin (Blanc-Mathieu et al. 2017; Szitenberg et al. 2017; Koutsovoulos et al. 2019; Susič et
al. 2020), and, apart from that, only two diploid facultative sexual species (Opperman et al.
2008; Somvanshi et al. 2018). Hence, further sampling of *Meloidogyne* species with diverse
ploidy levels and reproductive modes will be necessary to answer these questions.
698
699 Regardless of the TE load, we found that DNA transposons were majoritary in *M. incognita*.
700 Using the same annotation method, we found a similar result in *C. elegans*. Interestingly,
701 even if the methodology used was different, a similar observation was made at the whole
702 nematoda level (Szitenberg et al. 2016), suggesting a higher abundance of DNA
703 transposons might be a general feature of nematode genomes.
704

705 TE show signs of recent activity in *M. incognita* and they might
706 still be active

In the current analysis, we used variations in TE frequencies between geographical isolates
across loci in the *M. incognita* genome as a reporter of their activity. We have shown 75% of
the polymorphic TE loci display moderate frequency variations between isolates (<25%), a
majority being found with high frequencies (> 75%) in all the isolates simultaneously. Hence,
a substantial part of the TE can be considered as stable and fixed among the isolates.
Nevertheless, the remaining quarter of polymorphic TE loci present frequency variations
across the isolates higher than 25%. This observation concerns both the TE already present
in the reference genome, but also the neo-insertions. We even detected loci where the TE
frequencies were so contrasted between the isolates (HCPTEs) that we could predict the TE
presence/absence pattern among the isolates. Such frequency variations between isolates,
and the fact that part of the HCPTEs are isolate-specific neo-insertions, constitute strong
evidence for TE activity in the *M. incognita* genome.
We then evaluated how recent this activity could be, using % identity of the TE copies with
their respective consensuses as a proxy for their age. We showed that a substantial
proportion of the canonical TE annotations were highly similar to their consensus, indicating
most of these TE copies were recent in the genome. This result suggests a TE burst in the
M. incognita genome, which would be consistent with its likely recent hybrid origin (Blanc-
Mathieu et al. 2017). Indeed, as evoked previously, it is well established that hybridization
events can lead to a relaxation of the TE silencing mechanisms and consequently to a TE
expansion (Belyayev 2014; Guerreiro 2014; Rodriguez and Arkhipova 2018).
However, as suggested in (Bourgeois and Boissinot 2019), the extent of this phenomenon
might differ depending on the TE order. In *M. incognita,* MITEs and TIRs alone account for
~2/3 of the canonical TE annotations, but their fate in the genome seems to have followed

different paths. Indeed, MITE copy numbers almost linearly increase as a function of the
identity rate with their consensus, which suggests they might have progressively invaded the
genome being uncontrolled or poorly controlled as suggested for the rice genome (Lu et al.
2017). On the opposite, almost all the TIR copies share high percentage identity with their
consensuses which could be reminiscent of a rapid and recent burst. Nevertheless, this
burst could have quickly been under control as we observed that the TIR neo-insertions are
less numerous than expected owing to their abundance in the genome.
Because no molecular clock is available for *M. incognita*, it is impossible to evaluate more
precisely when this burst would have happened and how fast each TE from each order
would have spread in the genome. However, while an absolute dating of TE activities is
currently not possible, a relative timing of the events regarding speciation and diversification
can still be deduced from distribution of TE loci frequencies across the isolates. Indeed, we
have shown that some neo-insertion were shared between isolates and that in each case,
the concerned isolates were related according to the phylogeny. This observation indicates
that these neo-insertions occurred after *M. incognita* speciation, but before the diversification
of the phylogenetically-related isolates, in a common ancestor. Other TE neo-insertions, in
contrast, were so far isolate-specific, suggesting some TE movements were more recent and
that TE mobility might be a continuous phenomenon.
As there is no correlation between life history traits and geography among those isolates
(Koutsovoulos 2019), we can make the hypothesis that the Brasilian geographical isolates
we compared were recently spread by human intervention across different cultivated fields
during the modern era of extensive agriculture. Hence we could conclude that these neo-
insertions happened in the last centuries.
Overall, these observations, the distribution of percent identities of some TE copies to their
consensuses shifted towards high value, as well as support for transcriptional activity of
some of the genes involved in the transposition machinery, suggest TE have recently been
active in *M. incognita* and are possibly still active.
757

758 Concluding remarks

In this study we used population genomics technique and statistical analyses of the results
to assess whether TE might contribute to the genome dynamics of *M. incognita* and possibly
to its adaptive evolution. Overall, we provided a body of evidence suggesting TE have been
at least recently active and might still be active. With thousands of loci showing variations in
TE presence frequencies across geographical isolates, there is a clear impact on the *M.*
incognita genome plasticity. Some TE being neo-inserted in coding or regulatory regions
might have a functional impact. Although no clear connection with a role in adaptive
evolution could be made so far, based on the few impacted coding loci we experimentally
checked in this study, this is not to be excluded given the current lack of large-scale
functional information for this species. This pioneering study constitutes a valuable resource
and opens new perspectives for future targeted investigation of the potential effect of TE
dynamics on the evolution, fitness and adaptability of *M. incognita*.

Materials and Methods

Material

The genome of *M. incognita*

We used the genome assembly published in (Blanc-Mathieu et al. 2017) as a reference for TE prediction and annotation as well as for read-mapping of the different geographical isolates (Koutsovoulos et al. 2020), used for prediction of TE presence frequencies.

Briefly, the triploid *M. incognita* genome is 185Mb long with ~12,000 scaffolds and a N50 length of ~38 kb. Although the genome is triploid, because of the high nucleotide divergence between the genome copies (8% on average), most of these genome copies have been correctly separated during genome assembly, which can be considered effectively haploid (Blanc-Mathieu et al. 2017; Koutsovoulos et al. 2020). This reference genome originally came from a *M. incognita* population from the Morelos region of Mexico and was reared on tomato plants from the offspring of one single female in our laboratory.

The genome of *C. elegans*

We used the *C. elegans* genome (The C. elegans Genome Sequencing Consortium 1998) assembly (PRJNA13758) to perform its repeatome prediction and annotation and compare our results to the literature as a methodological validation.

Genome reads for 12 *M. incognita* geographical isolates

To predict the presence frequencies at TE loci across different *M. incognita* isolates, we used whole-genome sequencing data from pools of individuals from 12 different geographical regions (sup. Fig 2 & sup. Table 6). One pool corresponds to the Morelos isolates used to produce the *M. incognita* reference genome itself, as described above. The 11 other pools correspond to different geographical isolates across Brazil as described in (Koutsovoulos et al. 2020).

All the samples were reared from the offspring of one single female and multiplied on tomato plants. Then, approximately 1 million individuals were pooled and sequenced by Illumina paired-end reads (2*150bp). Libraries sizes vary between 74 and 76 million reads (Koutsovoulos et al. 2020).

We used cutadapt-1.15 (Martin 2011) to trim adapters, discard small reads, and trim low-quality bases in reads boundaries (–max-n=5 -q 20,20 -m 51 -j 32 -a AGATCGGAAGAGCACACGTCTGAACTCCAGTCA -A AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT). Then, for each library, we performed a fastqc v-0.11.8 (Andrew S., 2010: http://www.bioinformatics.babraham.ac.uk/projects/fastqc) analysis to evaluate the quality of the reads. FastQC results analyses showed that no additional filtering or cleaning step was needed and no further read was discarded.

Methods

We performed the statistical analysis and the graphical representation using R' v-3.6.3 and the following libraries: ggplot2, reshape2, dplyr, ggpubr, phangorn, plyr, and UpSetr. All codes and analysis workflows are available in the supplementary materials 1 to 3. For experimental validations, see supplementary material 4.

M. incognita and *C. elegans* repeatome predictions and annotations.

We predicted and annotated the *M. incognita* and *C. elegans* repeatomes following the same protocol as thoroughly explained in (Koutsovoulos et al. 2019). We define the repeatome as all the repeated sequences in the genome, excluding Simple Sequence Repeats (SSR) and microsatellites. Then, following the above-mentioned protocol, we further analysed each repeatome to isolate annotations with canonical signatures of Transposable Elements (TEs).

Below, we briefly explain each step and describe protocol adjustments.

Genome pre-processing.

Unknown nucleotides 'Ns' encompass 1.81% of *the M. incognita* reference genome and need to be trimmed before repeatome predictions. We created a modified version of the genome by splitting it at N stretches of length 11 or more and then trimming all N, using dbchunk.py from the REPET package (Quesneville et al. 2005; Flutre et al. 2011). As this increases genome fragmentation and may, in turn, lead to false positives in TE detection, we only kept chunks of length above the L90 chunk length threshold, which is 4,891 bp. This modified version of the genome was only used to perform the *de novo* prediction of the TE consensus library (below). The TE annotation was performed on the whole reference genome.

The *C. elegans* reference genome was entirely resolved (no N), at the chromosome-scale. Hence, we used the whole assembly as is to perform the *de novo* prediction analysis.

De novo prediction: constituting draft TE-consensus libraries.

For each species, we used the TEdenovo pipeline from the REPET package to generate a draft TE-consensus library..

Briefly, TEdenovo pipeline i) realises a self-alignment of the input genome to detect repetitions, ii) clusters the repetitions, iii) performs multiple alignments from the clustered repetitions to create consensus sequences, and eventually, iv) classify the consensus sequence following the Wicker's classification (Wicker et al. 2007) using structural and homology based information. One of the most critical steps of this process concerns the clustering of the repetitions as it requires prior knowledge about assembly ploidy and phasing quality.

We ran the analysis considering the modified *M. incognita* reference assembly previously described as triploid and set the 'minNbSeqPerGroup' parameter to 7 (*i.e* 2n+1). As the *C. elegans* assembly was haploid, we set the same parameter to 3.

All the remaining parameters values set in these analyses can be found in the TEdenovo configuration files (see supplementary material 2).

Automated curation of the TE-consensus libraries.

To limit the redundancy in the previously created TE consensus libraries and the false
positives, we performed an automated curation step. Briefly, for each species, i) we
performed a minimal annotation (steps 1, 2, 3, 7 of TEannot) of their genome with their
respective draft TE-consensus libraries, and ii) only retained consensus sequences with at
least one Full-Length Copy (FLC) annotated in the genome. All parameters values are
described in the configuration files produced in the supplementary material 2.
860

Repeatome annotation
For each species, we performed a full annotation (steps 1, 2, 3, 4, 5, 7, and 8) of their
genome with their respective cleaned TE-consensus libraries using TEannot from the
REPET package. The obtained repeatome annotations (excluding SSR and microsatellites)
were exported for further analyses. All parameters values are described in the configuration
files produced in the supplementary material 2.
867

Repeatome post-processing: identifying annotations with canonical signatures of
TEs.
Using in house scripts (see supplementary material 2), we analysed REPET outputs to retain
annotations with canonical signatures of Transposable Elements (TEs) from the rest of the
repeatomes. The same parameters were set for *M. incognita* and *C. elegans*. Briefly, for
each species, we only conserved TE annotations i) classified as retro-transposons or DNA-
transposons, ii) longer than 250 bp, iii) sharing more than 85% identity with their consensus
sequence, iv) covering more than 33% of their consensus sequence length, v) first aligning
with their consensus sequence in a BLAST analysis against the TE-consensus library, and
vi) not overlapping with other annotations. TE annotations respecting all the described
criterion were referred to as canonical TE annotations.

Putative transposition machinery identification (*M. incognita* only)
We analysed the *M. incognita* predicted proteome and transcriptome (Blanc-Mathieu et al.
2017) and crossed the obtained information with the canonical TE-annotation to identify TE
containing genes putatively involved in the transposition machinery and evaluate TE-related
gene expression levels in comparison to the rest of the genes in the genome.
885

Finding genes coding for proteins with TE-related HMM profiles
We performed an exhaustive HMMprofile search analysis on the whole *M. incognita*
predicted proteome and then looked for proteins with TE-related domains. First, we
concatenated two HMMprofile libraries into one: Pfram32 (Finn et al. 2016) library and
Gypsy DB 2.0 (Llorens et al. 2011), a curated library of HMMprofiles linked to viruses, mobile
genetic elements, and genomic repeats. Then, using this concatenated HMM profile library,
we performed an exhaustive but stringent HMM profile search on the *M. incognita* proteome
using hmmscan (-E 0.00001 --domE 0.001 --noali).
Eventually, using in house script (see supplementary material 1), we selected the best non-
overlapping HMM profiles for each protein and then tagged corresponding genes with TE-
related HMM profiles thanks to a knowledge-based function from the REPET tool

'profileDB4Repet.py'. We kept as genes with TE-related profiles all the genes with at least
one TE-related HMM-profile identified.
899
900
901

Genes expression level
To determine the *M. incognita* protein-coding genes expression patterns, we used data from
a previously published life-stage specific RNA-seq analysis of *M. incognita* transcriptome
during tomato plant infection (Blanc-Mathieu et al. 2017). This analysis encompassed four
different life stages: (i) eggs, (ii) pre-parasitic second stage juveniles (J2), (iii) a mix of late
parasitic J2, third stage (J3) and fourth stage (J4) juveniles and (iv) adult females, all
sequenced in triplicates.
The cleaned RNA-seq reads were retrieved from the previous analysis and re-mapped to the
M. incognita annotated genome assembly (Blanc-Mathieu et al. 2017) using a more recent
version of STAR (2.6.1) (Dobin et al. 2013) and the more stringent end-to-end option (i.e. no
soft clipping) in 2-passes. Expected read counts were calculated on the predicted genes
from the *M. incognita* GFF annotation as FPKM values using RSEM (Li and Dewey 2011) to
take into account the multi-mapped reads via expectation maximization. To reduce
amplitude of variations, raw FPKM values were transformed to Log10(FPKM+1) and the
median value over the 3 replicates was kept as a representative value in each life stage. The
expression   data   are   available   at
https://data.inra.fr/dataset.xhtml?persistentId=doi:10.15454/YM2DHE .
Then, for each life stage independently, i) we ranked the gene expression values, and ii)
defined gene expression level corresponding to the gene position in the ranking. We
considered as substantially expressed all the genes that presented an expression level >=
1st quartile in at least one life stage.
923
TE annotations with potential transposition machinery
To identify TE-annotations including predicted genes involved in transposition machinery
(inclusion >= 95% of the gene length), we performed the intersection of the canonical TE
annotation and the genes annotation BED files (see supplementary material 1) using the
intersect tool (-wo -s -F 0.95) from the bedtools v-2.27.1 suite (Quinlan and Hall 2010).
We then cross-referenced the obtained file with the list of the substantially expressed genes
and the list of the TE-related genes previously elaborated to identify the TEs containing
potential transposition machinery genes and their expression levels.
932

Evaluation of TE presence frequencies across the different *M. incognita*
isolates
We used the popoolationTE2 v-1.10.04 pipeline (Kofler et al. 2016) to compute isolate-
related support frequencies of both annotated, and *de novo* TE-loci across the 12 *M.*
incognita geographical isolates previously described. To that end, we performed a 'joint'
analysis as recommended by the popoolationTE2 manual. Briefly, popoolationTE2 uses both
quantitative and qualitative information extracted from paired-end (PE) reads mapping on the
TE-annotated reference genome and a set of reference TE sequences to detect signatures
of TE polymorphisms and estimate their frequencies in every analysed isolate. Frequency

values correspond to the proportion of individuals in an isolate for which a copy of the TE is
present at a given locus.
944
945
946
947
Preparatory work: creating the TE-hierarchy and the TE-merged-reference files.
We used the canonical TE-annotation set created above (see supplementary material 3) and
the *M. incognita* reference genome to produce the TE-merged reference file and the TE-
hierarchy file necessary to perform the popoolationTE analysis.
We used getfasta and maskfasta commands (default parameters) from the bedtools suite to
respectively extract and mask the sequences corresponding to canonical TE-annotations in
the reference genome. Then we concatenated both resulting sequences in a 'TE-merged
reference' multi fasta file. The 'TE-hierarchy' file was created from the TE-annotation file from
which it retrieves and stores the TE sequence name, the family, and the TE-order for every
entry.
958
959 Reads mapping
960 For each *M. incognita* isolate library, we mapped forward and reverse reads separately on
961 the "TE-merged-references" genome-TE file using the local alignment algorithm bwa bwasw
962 v-0.7.17-r1188 (Li and Durbin 2009) with the default parameters. The obtained sam
963 alignment files were then converted to bam files using samtools view v-1.2 (Li et al. 2009).
964
Restoring paired-end information and generating the ppileup file.
We restored paired-end information from the previous separate mapping using the sep2pe (-
-sort) tool from popoolationTE2. Then, we created the ppileup file using the 'ppileup' tool
from popolationTE2 with a map quality threshold of 15 (--map-qual 15).
For every base of the genome, this file summarises the number of PE reads inserts
spanning the position (physical coverage) but also the structural status inferred from paired-
end read covering this site.
972
Estimating target coverage and subsampling the ppileup to a uniform coverage
As noticed by R. Kofler, heterogeneity in physical coverage between populations may lead to
discrepancies in TE frequency estimation. Hence, we flattened the physical coverage across
the *M. incognita* isolates by a subsampling and a rescaling approach.
We first estimated the optimal target coverage to balance information loss and homogeneity
using the 'stats-coverage' tool from PopoolationTE2 (default parameter) and set this value to
15X. We then used the 'subsamplePpileup' tool (--target-coverage 15) to discard positions
with a physical coverage below 15X and rescale the coverage of the remaining position to
that value.
982
Identify signatures of TE polymorphisms
We identified signatures of TE polymorphisms from the previously subsampled file using the
'identifySignature' tool following the joint algorithm (--mode joint; --min-count 2; --signature-
window minimumSampleMedian; --min-valley minimumSampleMedian).
Then, for each identified site, we estimated TE frequencies in each isolate using the
'frequency' tool (default parameters). Eventually, we paired up the signatures of TE

polymorphisms using 'pairupSignatures' tool (--min-distance -200; --max-distance -- 300 as
recommended by R. Kofler), yielding a final list of potential TE-polymorphisms positions in
the reference genome with their associated frequencies for each one of the isolates.

Evaluation of PopoolationTE2 systematic error rate in the TE-frequency estimation.

To estimate PopoolationTE2 systematic error rate in the TE-frequency estimation, we ran
the same analysis (from the PE information restoration step) but comparing each isolate
against itself (12 distinct analyses).
We then analysed each output individually, measuring the frequency difference between the
two 'replicates' in all the detected loci with FR signatures (see below for more explanations).
We tested the homogeneity of the frequency-difference across the 12 analyses with an
ANOVA and concluded that the mean values of the frequencies differences between the
analysis were not significantly heterogeneous (p. value = 0.102 > 0.05). Hence, we
concatenated the 12 analysis frequency-difference and set the systematic error rate in the
TE-frequency estimation to 2 times the standard deviation of the frequency differences, a
value of 0.97 % .
1004

TE polymorphism analysis

Isolating TE loci with frequency variation across *M. incognita* isolates.
We parsed PopoolationTE2 analysis output to identify TE loci with enough evidence to
characterise them as polymorphic in frequency across the isolates.
PopoolationTE2 output informs for each detected locus i) its position on the reference
genome, ii) its frequency value for every sample of the analysis (e.g each isolate), and iii)
qualitative information about the reads mapping signatures supporting a TE insertion.
In opposition to separate Forward ('F') or Reverse ('R') signatures, 'FR' signatures mean the
locus both boundaries are supported by significant physical coverage. Entries with such type
of signature are more accurate in terms of frequency and position estimation. Hence, we
only retained candidate loci with 'FR' signatures. Then, for each locus, we computed the
maximal frequency variation between all the isolates and discarded the loci with a frequency
difference smaller than the PopoolationTE2 systematic error rate in the TE-frequency
estimation we computed (0.97 %; see above). We also discarded loci where different TEs
were predicted to be inserted. We considered the remaining loci as polymorphic in frequency
across the isolates.
1021

Isolates phylogeny
We reconstructed *M. incognita* isolates phylogeny according to their patterns of
polymorphism in TE frequencies.
We first computed a euclidean distance matrix from the isolates TE frequencies of all the
detected polymorphic loci. We then used the distance matrix to construct the phylogenetic
tree using the Neighbor Joining (NJ) method (R' phangorn package v-2.5.5). We computed
nodes support values with a bootstrap approach (n=500 replicates). We compared the
resulting tree with the topology described in (Koutsovoulos et al. 2020) using Itol v-4.0
viewer (Letunic and Bork 2019).

 Polymorphisms characterisation.

We exported the polymorphic TE positions as an annotation file, and we used bedtools
intersect (-wao) to perform their intersection with the reference canonical TE annotation. We
then cross-referenced the results with the filtered popoolationTE2 output and defined a
decision tree to characterise the TE-polymorphism detected by popoolationTE2 as
'reference-TE polymorphism' (ref-polymorphism), 'extra-detection', or 'neo-insertion'.
We considered a reference TE-annotation as polymorphic (e.g. ref-polymorphism locus) if:
i) The position of the polymorphism predicted by PoPoolationTE2 falls between the
boundaries of the reference TE-annotation
ii) Both the reference TE-annotation and the predicted polymorphism belong to the same TE-
consensus sequence.
iii) The TE has a predicted frequency > 75% in the reference isolate Morelos.
Canonical TE-annotations that did not intersect with polymorphic loci predicted by
PopoolationTE2 were considered as non-polymorphic.
We classified as 'neo-insertions' all the polymorphic loci for which no canonical TE was
predicted in the reference annotation (polymorphism position is not included in a reference
TE-annotation), but which were detected with a frequency > 25% in at least one isolate
different from the reference isolate Morelos, in which the TE frequency should be inferior to
1% and thus considered truly absent in the reference genome.
Finally, we classified as 'extra-detection' all the polymorphic loci which did not correspond to
a reference annotation but which were detected with a frequency > 25% in the reference
isolate Morelos (at least). Polymorphic loci having a frequency between 1% and 25% in
Morelos isolate were considered ambiguous and were discarded.
Then, for each TE polymorphism, we investigated the homogeneity of the TE frequency
between the isolates.We considered TE frequency was homogeneous between isolates
when the maximum frequency variation between isolate was <= to 25%. Above this value,
we considered the TE presence frequency was heterogeneous between isolates.
1058
1059
1060

Highly Contrasted Polymorphic TE loci (HCPTEs): isolation,
characterisation and experimental validation.
HCPTEs isolation
We considered as highly contrasted all the polymorphic loci for which i) all the isolates had
frequency values either < 25% or > 75%, ii) at least one isolate showed a frequency < 25 %
while another presented a frequency > 75%. Polymorphic loci fitting with these requirements
were exported as an annotation file in the bed format.
1068
HCPTEs possible functional impact
We first identified the genes potentially impacted by the HCPTEs by cross-referencing the
HCPTEs annotation file with the gene annotation file, using the bedtools suite. We used the
'closest' program (-D b -fu -io; b being the gene annotation file) to identify the closest (but not
intersecting) gene downstream each HCPTE. We only retained the entries with a maximum
distance of 1 kb between the HCPTE and gene boundaries. We identified the insertions in
the gene using the 'intersect' tool (-wo).
1076

Then, we performed a manual bioinformatic functional analysis for each gene potentially impacted by HCPTEs. Protein sequences were extracted from the *M. incognita* predicted proteome (Blanc-Mathieu et al. 2017) and blasted (blastp; default parameters) against the Non-Redundant protein sequences database (NR) from the NCBI (https://blast.ncbi.nlm.nih.gov/). The same sequences were also used on the InterProScan website (https://www.ebi.ac.uk/interpro/) to perform an extensive search on all the available libraries of conserved protein domains and motifs.

Then, for each gene potentially impacted by HCPTEs, we performed an orthology search on the Wormbase Parasite website (https://parasite.wormbase.org/) using genes accession numbers and the pre-computed ENSEMBL Compara orthology prediction (Herrero et al. 2016).

Finally, we analysed the expression levels of the genes potentially impacted by HCPTEs extracting the information from the RNA-seq analysis of four *M. incognita* life-stages performed previously (see Putative transposition machinery identification section).

Experimental validation of Highly Contrasted Polymorphic TE loci

To experimentally validate in-silico predictions of TE neo-insertions with potential functional impact, we selected 5 candidates among the HCPTEs loci and performed a PCR experiment. To run this experiment, we used DNA remaining from extractions performed on the *M. incognita* isolates for a previous population genomics analysis (Koutsovoulos et al. 2020).

Primer design and PCR amplification.
We designed primers for the PCR analysis using the Primer3Plus web interface (Untergasser et al. 2007). The set of 10 primers with the corresponding sequence and expected amplicon sizes with, or without TE insertion, is shown in (sup. Table 7 & supplementary material 4). We used primers amplifying the whole actin-encoding gene (Minc3s00960g19311) as positive control.

PCR experiments were performed on *M. incognita* Morelos isolate and 11 Brazilian isolates: R1-2, R1-3, R1-6, R2-1, R2-6, R3-1, R3-2, R3-4, R4-1, R4-3 and R4-4.

R3-1 presented no amplification in any of the tested loci nor the positive control (actin) and was thus discarded from this analysis.

PCR mixture contained 0.5µmol of each primer, 1x MyTaq™ reaction buffer and 1.0 U of MyTaq™ DNA polymerase (Bioline Meridian Bioscience) adjusted to a total volume of 20µL. PCR amplification was performed with a TurboCycler2 (Blue-Ray Biotech Corp.). PCR conditions were as follows: initial denaturation at 95°C for 5 min, followed by 35 cycles of 95°C for 30 s, 56°C for 30 s of annealing, and 72°C for 3 min of extension, the program ending with a final extension at 72°C for 10 min. Aliquots of 5µL were migrated by electrophoresis on a 1% agarose gel (Sigma Chemical Co.) for 70 min at 100 V. The size marker used is 1kb Plus DNA Ladder (New England Biolabs Inc.), containing the following size fragments in bp: 100, 200, 300, 400, 500, 600, 700, 900, 1000, 1200, 1500, 2000, 3000, 4000, 5000, 6000, 8000 and 10000.

Purification and sequencing of PCR amplicons.
Amplicon bands were revealed using ethidium bromide and exposure to ultraviolet radiation. PCR products bands were excised from the agarose gel with a scalpel and purified using MinElute Gel Extraction Kit (Qiagen) before sequencing, following the manufacturer's protocol. PCR products were sequenced by Sanger Sequencing (Eurofins Genomics).

Forward (F) and Reverse (R) sequences were blasted individually (https://blast.ncbi.nlm.nih.gov/ ; Optimised for 'Somewhat similar sequences', default

parameters) to the expected TE-consensus sequence and to the genomic region surrounding the predicted insertion point (2 kb region: 1kb upstream the predicted insertion point and 1kb downstream). When no significant hit was found, the sequence was blasted against the *Meloidogyne* reference genomes available (https://meloidogyne.inrae.fr/), the whole TE-consensus library, and the NR database on the NCBI blast website.

Acknowledgements

DKLK would like to thank Joffrey Mejias for all the inspiring discussion, and all the experimental pieces of advice he gave. DKLK would also like to thank the BIG bioinformatics platform from the PlantBios infrastructure as well as the URGI team for providing facilities and technical support.

References

Abad P, Gouzy J, Aury J-M, Castagnone-Sereno P, Danchin EGJ, Deleury E, Perfus-Barbeoch L, Anthouard V, Artiguenave F, Blok VC, et al. 2008. Genome sequence of the metazoan plant-parasitic nematode Meloidogyne incognita. Nat. Biotechnol. 26:909–915.

Agrios GN. 2005. Plant Pathology, 5th Edition. Burlington, USA: Elsevier Academic Press

Barbary A, Djian□Caporalino C, Palloix A, Castagnone□Sereno P. 2015. Host genetic resistance to root-knot nematodes, Meloidogyne spp., in Solanaceae: from genes to the field. Pest Manag. Sci. 71:1591–1598.

Bast J, Jaron KS, Schuseil D, Roze D, Schwander T. 2019. Asexual reproduction reduces transposable element load in experimental yeast populations.Coop G, Tautz D, Coop G, Charlesworth B, editors. eLife 8:e48548.

Bast J, Schaefer I, Schwander T, Maraun M, Scheu S, Kraaijeveld K. 2015. No Accumulation of Transposable Elements in Asexual Arthropods. Mol. Biol. Evol.:msv261.

Belyayev A. 2014. Bursts of transposable elements as an evolutionary driving force. J. Evol. Biol. 27:2573–2584.

Bessereau J-L. 2006. Transposons in C. elegans. WormBook [Internet]. Available from: http://www.wormbook.org/chapters/www_transposons/transposons.html

Blanc-Mathieu R, Perfus-Barbeoch L, Aury J-M, Rocha MD, Gouzy J, Sallet E, Martin-Jimenez C, Bailly-Bechet M, Castagnone-Sereno P, Flot J-F, et al. 2017. Hybridization and polyploidy enable genomic plasticity without sex in the most devastating plant-parasitic nematodes. PLOS Genet. 13:e1006777.

Bourgeois Y, Boissinot S. 2019. On the Population Dynamics of Junk: A Review on the Population Genomics of Transposable Elements. Genes 10:419.

Castagnone-Sereno P. 2006. Genetic variability and adaptive evolution in parthenogenetic root-knot nematodes. Heredity 96:282–289.

Castagnone-Sereno P, Danchin EGJ. 2014. Parasitic success without sex – the nematode experience. J. Evol. Biol. 27:1323–1333.

Castagnone□Sereno P, Mulet K, Danchin EGJ, Koutsovoulos GD, Karaulic M, Rocha MD, Bailly□Bechet M, Pratx L, Perfus□Barbeoch L, Abad P. 2019. Gene copy number variations as signatures of adaptive evolution in the parthenogenetic, plant-parasitic

nematode Meloidogyne incognita. Mol. Ecol. 28:2559–2572.

Castagnone-Sereno P, Wajnberg E, Bongiovanni M, Leroy F, Dalmasso A. 1994. Genetic
variation inMeloidogyne incognita virulence against the tomatoMi resistance gene:
evidence from isofemale line selection studies. Theor. Appl. Genet. 88:749–753.

Faino L, Seidl MF, Shi-Kunne X, Pauper M, Berg GCM van den, Wittenberg AHJ, Thomma
BPHJ. 2016. Transposons passively and actively contribute to evolution of the two-
speed genome of a fungal pathogen. Genome Res. [Internet]. Available from:
http://genome.cshlp.org/content/early/2016/07/12/gr.204974.116

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M,
Qureshi M, Sangrador-Vegas A, et al. 2016. The Pfam protein families database:
towards a more sustainable future. Nucleic Acids Res. 44:D279–D285.

Flutre T, Duprat E, Feuillet C, Quesneville H. 2011. Considering Transposable Element
Diversification in De Novo Annotation Approaches. PLoS ONE 6:e16526.

Glémin S, François CM, Galtier N. 2019. Genome Evolution in Outcrossing vs. Selfing vs.
Asexual Species. In: Anisimova M, editor. Evolutionary Genomics: Statistical and
Computational Methods. Methods in Molecular Biology. New York, NY: Springer. p.
331–369. Available from: https://doi.org/10.1007/978-1-4939-9074-0_11

Glémin S, Galtier N. 2012. Genome Evolution in Outcrossing Versus Selfing Versus Asexual
Species. In: Anisimova M, editor. Evolutionary Genomics. Vol. 855. Totowa, NJ:
Humana Press. p. 311–335. Available from: http://link.springer.com/10.1007/978-1-
61779-582-4_11

Gross SM, Williamson VM. 2011. Tm1: A Mutator/Foldback Transposable Element Family in
Root-Knot Nematodes. PLoS ONE 6:e24534.

Guerreiro MPG. 2014. Interspecific hybridization as a genomic stressor inducing mobilization
of transposable elements in *Drosophila*. Mob. Genet. Elem. 4:e34394.

Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, Vilella AJ, Searle SMJ,
Amode R, Brent S, et al. 2016. Ensembl comparative genomics resources. Database
[Internet] 2016. Available from:
https://academic.oup.com/database/article/doi/10.1093/database/bav096/2630091

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. Genet. Res.
8:269–294.

Hoffmann AA, Reynolds KT, Nash MA, Weeks AR. 2008. A high incidence of
parthenogenesis in agricultural pests. Proc. R. Soc. Lond. B Biol. Sci. 275:2473–
2481.

Jones JT, Haegeman A, Danchin EGJ, Gaur HS, Helder J, Jones MGK, Kikuchi T,
Manzanilla-López R, Palomares-Rius JE, Wesemael WML, et al. 2013. Top 10 plant-
parasitic nematodes in molecular plant pathology. Mol. Plant Pathol. 14:946–961.

Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot N, Moreno
S, Sohrmann M, et al. 2003. Systematic functional analysis of the Caenorhabditis
elegans genome using RNAi. Nature 421:231–237.

Katju V, Bergthorsson U. 2013. Copy-number changes in evolution: rates, fitness effects and
adaptive significance. Front. Genet. [Internet] 4. Available from:
http://www.frontiersin.org/Evolutionary_and_Population_Genetics/10.3389/fgene.201
3.00273/abstract

Kofler R, Gómez-Sánchez D, Schlötterer C. 2016. PoPoolationTE2: Comparative Population
Genomics of Transposable Elements Using Pool-Seq. Mol. Biol. Evol. 33:2759–2764.

Kondrashov AS. 1988. Deleterious mutations and the evolution of sexual reproduction.
Nature 336:435–440.

Koutsovoulos GD, Marques E, Arguel M-J, Duret L, Machado ACZ, Carneiro RMDG,
Kozlowski DK, Bailly☐Bechet M, Castagnone☐Sereno P, Albuquerque EVS, et al.
2020. Population genomics supports clonal reproduction and multiple independent
gains and losses of parasitic abilities in the most devastating nematode pest. Evol.
Appl. 13:442–457.

Koutsovoulos GD, Poullet M, Ashry AE, Kozlowski DK, Sallet E, Rocha MD, Martin-Jimenez
C, Perfus-Barbeoch L, Frey J-E, Ahrens C, et al. 2019. The polyploid genome of the

1224   mitotic parthenogenetic root knot nematode Meloidogyne enterolobii.
1225   bioRxiv:586818.
1226 Kraaijeveld K, Zwanenburg B, Hubert B, Vieira C, De Pater S, Van Alphen JJM, Den Dunnen
1227   JT, De Knijff P. 2012. Transposon proliferation in an asexual parasitoid. Mol. Ecol.
1228   21:3898–3906.
1229 Laricchia KM, Zdraljevic S, Cook DE, Andersen EC. 2017. Natural Variation in the
1230   Distribution and Abundance of Transposable Elements Across the Caenorhabditis
1231   elegans Species. Mol. Biol. Evol. 34:2187–2202.
1232 Lerat E, Goubert C, Guirao☐Rico S, Merenciano M, Dufour A-B, Vieira C, González J. 2019.
1233   Population-specific dynamics and selection patterns of transposable element
1234   insertions in European natural populations. Mol. Ecol. 28:1506–1522.
1235 Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
1236   developments. Nucleic Acids Res. 47:W256–W259.
1237 Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler
1238   transform. Bioinformatics 25:1754–1760.
1239 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
1240   R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics
1241   25:2078–2079.
1242 Lively CM. 2010. A Review of Red Queen Models for the Persistence of Obligate Sexual
1243   Reproduction. J. Hered. 101:S13–S20.
1244 Llorens C, Futami R, Covelli L, Domínguez-Escribá L, Viu JM, Tamarit D, Aguilar-Rodríguez
1245   J, Vicente-Ripolles M, Fuster G, Bernet GP, et al. 2011. The Gypsy Database
1246   (GyDB) of mobile genetic elements: release 2.0. Nucleic Acids Res. 39:D70–D74.
1247 Lu L, Chen J, Robb SMC, Okumoto Y, Stajich JE, Wessler SR. 2017. Tracking the genome-
1248   wide outcomes of a transposable element burst over decades of amplification. Proc.
1249   Natl. Acad. Sci. 114:E10550–E10559.
1250 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
1251   reads. EMBnet.journal 17:10–12.
1252 McCarter JP. 2009. Molecular Approaches Toward Resistance to Plant-Parasitic
1253   Nematodes. In: Berg RH, Taylor CG, editors. Cell Biology of Plant Nematode
1254   Parasitism. Vol. 15. Plant Cell Monographs. Berlin, Heidelberg: Springer Berlin
1255   Heidelberg. p. 239–267. Available from:
1256   http://www.springerlink.com/index/10.1007/978-3-540-85215-5_9
1257 McClintock B. 1984. The significance of responses of the genome to challenge. Science
1258   226:792–801.
1259 Mhiri C, Parisod C, Daniel J, Petit M, Lim KY, Borne FD de, Kovarik A, Leitch AR,
1260   Grandbastien M-A. 2019. Parental transposable element loads influence their
1261   dynamics in young Nicotiana hybrids and allotetraploids. New Phytol. 221:1619–
1262   1633.
1263 Muller HJ. 1964. The Relation of Recombination to Mutational Advance. Mutat Res 106:2–9.
1264 Opperman CH, Bird DM, Williamson VM, Rokhsar DS, Burke M, Cohn J, Cromer J, Diener
1265   S, Gajan J, Graham S, et al. 2008. Sequence and genetic map of Meloidogyne
1266   hapla: A compact nematode genome for plant parasitism. Proc Natl Acad Sci U A
1267   105:14802–14807.
1268 Quesneville H, Bergman CM, Andrieu O, Autard D, Nouaud D, Ashburner M, Anxolabehere
1269   D. 2005. Combined evidence annotation of transposable elements in genome
1270   sequences. PLoS Comput Biol 1:166–175.
1271 Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
1272   features. Bioinformatics 26:841–842.
1273 Rice WR. 2002. Experimental tests of the adaptive significance of sexual recombination.
1274   Nat. Rev. Genet. 3:241–251.
1275 Rodriguez F, Arkhipova IR. 2018. Transposable elements and polyploid evolution in animals.
1276   Curr. Opin. Genet. Dev. 49:115–123.
1277 Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. 2019. The global
1278   burden of pathogens and pests on major food crops. Nat. Ecol. Evol. 3:430–439.

Somvanshi VS, Tathode M, Shukla RN, Rao U. 2018. Nematode Genome Announcement: A
Draft Genome for Rice Root-Knot Nematode, Meloidogyne graminicola. J. Nematol.
50:111–116.
Susič N, Koutsovoulos GD, Riccio C, Danchin EGJ, Blaxter ML, Lunt DH, Strajnar P, Širca
S, Urek G, Stare BG. 2020. Genome sequence of the root-knot nematode
Meloidogyne luci. J. Nematol. 52:1–5.
Szitenberg A, Cha S, Opperman CH, Bird DM, Blaxter ML, Lunt DH. 2016. Genetic drift, not
life history or RNAi, determine long term evolution of transposable elements.
Genome Biol. Evol.:evw208.
Szitenberg A, Salazar-Jaramillo L, Blok VC, Laetsch DR, Joseph S, Williamson VM, Blaxter
ML, Lunt DH. 2017. Comparative Genomics of Apomictic Root-Knot Nematodes:
Hybridization, Ploidy, and Dynamic Genome Change. Genome Biol. Evol. 9:2844–
2861.
The C. elegans Genome Sequencing Consortium. 1998. Genome sequence of the
nematode C. elegans: a platform for investigating biology. Science 282:2012–2018.
Trudgill DL, Blok VC. 2001. Apomictic, polyphagous root-knot nematodes: exceptionally
successful and damaging biotrophic root pathogens. Annu Rev Phytopathol 39:53–
77.
Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JAM. 2007.
Primer3Plus, an enhanced web interface to Primer3. Nucleic Acids Res. 35:W71–
W74.
Vicient CM, Casacuberta JM. 2017. Impact of transposable elements on polyploid plant
genomes. Ann. Bot. 120:195–207.
Vrijenhoek RC, Parker ED. 2009. Geographical Parthenogenesis: General Purpose
Genotypes and Frozen Niche Variation. In: Schön I, Martens K, Dijk P, editors. Lost
Sex. Dordrecht: Springer Netherlands. p. 99–131. Available from:
http://www.springerlink.com/index/10.1007/978-90-481-2770-2_6
Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P,
Morgante M, Panaud O, et al. 2007. A unified classification system for eukaryotic
transposable elements. Nat. Rev. Genet. 8:973–982.
Wright S, Finnegan D. 2001. Genome evolution: Sex and the transposable element. Curr.
Biol. 11:R296–R299.