Queuine is a nutritional regulator of *Entamoeba histolytica* response to oxidative stress and a virulence attenuator

Shruti Nagaraja[1], Maggi W. Cai[2,3], Lotem Sarid[1], Yana Shaulov[1], Meirav Trebicz-Geffen[1], Mohit Mazumdar[5], Hugo Varet[6,7], Rachel Legendre[6,7], Jean-Yves Coppée[6,7], Peter C. Dedon[2,3], Samudrala Gourinath[5], Nancy Guillen[8,9], Yumiko Saito-Nakano[10], Chikako Shimokawa[10], Hajime Hisaeda[10], Serge Ankri[1*].

Affiliation

[1]Department of Molecular Microbiology, Ruth and Bruce Rappaport Faculty of Medicine, Technion, P.O.B. 9649, 31096 Haifa Israel.

[2]Department of Microbiology and Immunology, National University of Singapore, Singapore 117597

[3]Singapore-MIT Alliance for Research and Technology, Singapore 138602

[4]Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139 USA.

[5]Jawaharlal Nehru University School of Life Sciences, New Delhi 110067, India

[6]Plate-forme Technologique Biomics – Centre de Ressources et Recherches Technologiques (C2RT), Institut Pasteur, Paris, France

[7]Hub de Bioinformatique et Biostatistique – Département Biologie Computationnelle, Institut Pasteur, USR 3756 CNRS, Paris, France

[8]Centre National de la Recherche Scientifique, CNRS-ERL9195

[9]Institut Pasteur, Paris, France

[10]Department of Parasitology, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo 162-8640, Japan

*Corresponding author

Email:sankri@tx.technion.ac.il

Abstract

Queuosine is a naturally occurring modified ribonucleoside found in the first position of the anticodon of the transfer RNAs for Asp, Asn, His and Tyr. Eukaryotes lack pathways to synthesize queuine, the nucleobase precursor to queuosine, and must obtain it from diet or gut microbiota. Here we describe the effects of queuine on the physiology of the eukaryotic parasite, *Entamoeba histolytica*, the causative agent of amebic dysentery. Queuine is efficiently incorporated into *E. histolytica* tRNAs by a tRNA-guanine transglycosylase (EhTGT) and this incorporation stimulates the methylation of C_{38} in tRNA$^{Asp}_{GTC}$. Queuine protects the parasite against oxidative stress (OS) and antagonizes the negative effect that oxidation has on translation by inducing the expression of genes involved in OS response, such as heat shock protein 70 (Hsp 70), antioxidant enzymes, and enzymes involved in DNA repair. On the other hand, queuine impairs *E. histolytica* virulence by downregulating the expression of genes previously associated with virulence, including cysteine proteases, cytoskeletal proteins, and small GTPases. Silencing of EhTGT prevents incorporation of queuine into tRNAs and strongly impairs methylation of C_{38} in tRNA$^{Asp}_{GTC}$, parasite growth, resistance to OS, and cytopathic activity. Overall, our data reveal that queuine plays a dual role in promoting OS resistance and reducing parasite virulence.

Introduction

Amebiasis is an enormous global medical problem due to poor sanitary conditions and unsafe hygiene practices in many parts of the world. According to the World Health Organization, 50 million people in India, Southeast Asia, Africa, and Latin America suffer from amebic dysentery, with amebiasis causing the death of at least 100,000 people each year. *Entamoeba histolytica*, the etiologic agent of amebiasis, proliferate in the intestinal lumen and phagocytose resident gut flora. Over the last few decades, it has become evident that *E. histolytica*'s pathogenicity is directly linked to the parasite's interaction with the gut microbiota [1]. This interaction is very selective because only those bacteria with the appropriate recognition molecules are ingested by the parasite [2]. It has been reported that the *E. histolytica*'s association with specific intestinal bacteria changes the parasite's cell surface architecture [3,4]. It has also been reported that phagocytosis of pathogenic bacteria boosts *E. histolytica*'s cytopathogenicity, increases the expression of Gal/GalNAc lectin on the cell surface, and boosts cysteine proteinase activity when trophozoites are co-cultured with the enteropathogenic *Escherichia coli* (EPEC) O55 or *Shigella dysenteriae* [5]. It has also been reported that bacteria-induced augmentation of *E. histolytica*'s virulence is achieved only when the trophozoites phagocytose intact live cells [6]. The composition of the gut flora in patients suffering from amebiasis shows a significant decrease in the population size of *Bacteroides*, *Clostridium coccoides*, *Clostridium leptum,* *Lactobacillus,* and *Campylobacter* and an increase in *Bifidobacterium,* while there was no change in *Ruminococcus* compared to healthy patients [7]. These findings suggest that the pathogenesis of amebiasis might be driven by a dysregulated microbiome or crosstalk between enteropathic bacteria, the parasite, and the intestinal immune system. This crosstalk may be modulated by chemical signaling molecules, such as short-chain fatty acids (SCFAs) released by the bacteria [8], or by bacterial oxaloacetate that regulates parasite virulence and resistance to oxidative stress (OS) [9,10]. The modified ribonucleoside, queuosine, 7-(((4.5-cis-dihydroxy-2-cyclopenten-1-yl)-amino)-methyl)-7-deazaguanosine) may also participate in the crosstalk between the parasite and the microbiota. Eubacteria can synthesize the queuine nucleobase precursor of queuosine *de novo*. However, a recent study showed that some Eubacteria can also salvage precursors of queuosine [11]. Since mammals cannot synthesize queuine and salvage it from diet and gut microbes, these observations suggest a dynamic supply and demand for queuine by both gut microbiota and the host [12-14]. The diverse roles of queuosine in bacterial physiology are only now emerging. tRNA-guanine transglycosylase (TGT), which catalyzes the exchange of queuine for guanine at the wobble position of certain tRNAs, has been

recognized as one of the key enzymes that regulate virulence in *Shigella flexneri* [15,16]. Recently, it has been found that a modification of the wobble position of tRNA with a GUN anticodon by 7-deaza-guanosine derivative queuosine (Q34) stimulates *Schizosaccharomyces pombe* Dnmt2/Pmt1-dependent C38 methylation (m^{5}C$_{38}$) in the tRNA$_{Asp}$ anticodon loop [17], with the latter correlated with resistance to OS in many organisms including *E. histolytica* [18,19]. The fact that both methylation of C$_{38}$-tRNA^{Asp} and queuinosylation of tRNAs occur at the same anticodon loop suggests coordination among these modifications in regulating OS resistance in the parasite. It has been previously reported that queuine promotes the resistance of cancer cells to OS by increasing the activity of antioxidant enzymes [20]. In contrast, queuosine deficiency in tRNAs leads to the accumulation of misfolded proteins that induce a cellular stress response [17]. It has also been found that queuine enhances translational speed and fidelity in eukaryotes [21].

Here we report that queuine: (i) induces C$_{38}$ hypermethylation (m^{5}C$_{38}$) in the tRNA^{Asp} anticodon loop, (ii) promotes resistance of the parasite to OS by triggering the expression of genes associated with stress response, (iii) restores protein synthesis in parasite exposed to OS, and (iv) reduces parasite virulence by reducing expression of genes associated with virulence.

Results

Queuine is incorporated into *E. histolytica* tRNA and induces C38 hypermethylation (m^{5}C$_{38}$) in the tRNA^{Asp} anticodon loop.

Given the established role of queuine as a precursor to Q in tRNA, we first set out to characterize the incorporation of queuine into amoebic tRNA. Here we quantified Q in tRNA in the parasite using LC-MS/MS. tRNA samples were prepared from untreated trophozoites and from trophozoites grown with queuine at a concentration chosen based on studies with the social amoeba *Dictostelium discoideum* [22]. We observed that supplementation of queuine in the Trypticase Yeast Extract Iron Serum (TYI-S-33) growth medium caused a nearly 10-fold increase in the level of Q in tRNA in the trophozoites (Fig 1A). While we do not know if the number of copies of Q-containing tRNAs increased, these data indicate that queuine is indeed incorporated into *E. histolytica* tRNAs, which suggests the presence of an active TGT enzyme in *E. histolytica.* The results also raise concerns about the amount of queuine available in *E. histolytica* growth media and what represents an appropriate amount of queuine to add to the medium as a supplement for consistent growth conditions.

Recently, it has been found that a modification of the wobble position of tRNA with a GUN anticodon

by 7-deaza-guanosine derivative queuosine (Q34) stimulates *S. pombe* Dnmt2/Pmt1-dependent C_{38} methylation (m^5C_{38}) in the $tRNA^{Asp}$ anticodon loop [17]. The presence of an active Dnmt2 in *E. histolytica* [23-25] incited us to check if queuine also enhances the level of m^5C_{38} in the parasite's $tRNA^{Asp}$ anticodon loop (Fig 1B). We used PCR-based bisulfite sequencing to detect cytosine methylation in the tRNA by identifying the location of the methylated cytosine by sequencing. Amplicons were generated from trophozoites grown with and without queuine and sequences of several independent PCR products were sequenced and analyzed. We observed that the percentage of C_{38} $tRNA^{Asp}$ methylation in trophozoites grown with queuine ($55 \pm 7.1\%$) was significantly higher than the percentage of C_{38} $tRNA^{Asp}$ methylation in control trophozoites ($20 \pm 2\%$). In addition, we were unable to detect methylation in C_{32}, C_{33}, C_{37}, C_{48}, and C_{56}, whereas 100% methylation was observed in $C_{49.}$ These results which are in accordance to our previously published results [23] indicate that the bisulfite reaction was successful. These results also indicate that the incorporation of queuine into amoebic tRNAs leads to hypermethylation of $tRNA^{Asp}$.

Characterization of EhTGT

Given the queuine-induced increase in Q in tRNA, we next set out to characterize the *E. histolytica* TGT (EhTGT) and its role in Q incorporation and cell phenotype. We used a homology-based approach with the human QTRT1 gene to identify the subunits of *E. histolytica* TGT. The eukaryotic TGT enzymes, such as human TGT, consist of a queuine tRNA-ribosyltransferase 1 (QTRT1, eubacterial TGT homolog) and a queuine tRNA-ribosyltransferase domain-containing 1 (QTRTD1). To get more information about EhTGT, we performed a protein sequence alignment of *Homo sapiens* QTRT1 subunit (hQTRT1), EhQTRT1, and EhQTRTD1 (Supplementary Figure S1A). Based on annotation of the *E. histolytica* genome, a homolog of hQTRT1 and hQTRTD1 exists in *E. histolytica*, namely EhQTRT1 (XP_656142.1) and EhQTRTD1 (XP_652881.1). The EhQTRT1gene shares a high degree of homology with the human QTRT1 (52% sequence identity), whereas the EhQTRTD1 shares a homology at the C-terminal domain (28.3% sequence identity). Moreover, these two proteins also share a high degree of homology with the *Zymomonas mobilis TGT* (41.6% and 35.6%, respectively). The *Z. mobilis* TGT enzyme is a homodimeric zinc-binding protein, whereas the eukaryotic TGT is a heterodimeric zinc-binding protein. We identified four essential residues that are important for Zn^{2+} binding shown highlighted purple in (Cys 305, Cys307, Cys310, and His342; *E. histolytica* numbering) to be conserved in both EhQTRT1 and EhQTRTD1 (Supplementary Figure S1B)

In order to get information about the EhTGT structure, we built an *in silico* model of EhTGT subunits based on the *Thermotoga maritima* QTRT1 structure and murine QTRT2 structures (Fig 2A). The prediction also suggested that EhQTRT1 and EhQTRTD1 are homodimers (Supplementary Figure S2), which interact together to form a heterodimer like the eukaryotic TGT enzyme (Fig 2A). To get more insights about the EhTGT enzyme, a polyhistidine-tagged EhQTRT1 and untagged EhQTRTD1 was cloned and co-expressed in *E. coli*. During protein expression, the addition of 100μM $ZnS0_4$ and a low-temperature induction (19 ºC) was found to be essential for obtaining an active complex. The EhQTRTD1 was co-purified with the polyhistidine-tagged EhQTRT1 by Ni^{2+} affinity chromatography. The enzyme was further purified by size-exclusion chromatography (Fig 2B) and the fractions were analyzed by SDS-PAGE. The molecular weight of both EhQTRT1 and EhQTRTD1 on SDS-PAGE revealed their molecular mass to be around 44.2kDa and 44.1kDa respectively, which matches the predicted molecular mass (Fig 2C). To confirm the identity of both bands, mass spectrometry was performed on peptides obtained from tryptic digests of the bands excised from the SDS gel. After mapping these peptide fragments against *E. coli* and *E. histolytica* databases, our samples were confirmed to be EhQTRT1 and EhQTRD1 (Supplementary Figure S3).

To verify that the EhTGT is active, the activity of the complex was tested by determining its ability to displace guanine in the $tRNA^{Asp}$ anticodon with 8-$[^{14}C]$-guanine. We observed that EhTGT is an active enzyme and that EhQTRT1-EhQTRTD1 enzyme complex efficiently incorporates 8-$[^{14}C]$-guanine with a specific activity of 2.28 pmol/min/mg of protein, whereas the His-tagged EhQTRT1 showed no activity (Table 1).

The *in silico* model also predicted that the stretch of S102-L125 is essential for binding EhQTRTD1, and that the D267 residue in *Eh*QTRT1 is a substrate binding residue (Fig2A). A punctual mutation (D267 to A267) in the predicted active site of EhQTRT1 (indicated in black circle in Fig 2A) made the enzyme inactive without affecting the formation of the EhQTRT1-EhQTRTD1 complex (Table 1). Moreover, the deletion of the binding site S102-L125 (ΔEhQTRT1-EhQTRTD1) also resulted in an inactive enzyme (Table 1).

Together, these results confirm that the EhQTRT1 requires the presence of its interacting partner (EhQTRTD1) and validate the prediction of the *in silico* model (Fig 2A).

To better understand the role of EhTGT in *E.histolytica*, we silenced the expression of EhQTRT1 by using an epigenetic method based on antisense small RNAs [26]. In this method, a gene-coding region to which large numbers of antisense small RNAs map is used as a 'trigger' to silence the gene fused to

it. The silencing is maintained after that the trigger plasmid has been cured from the silenced strain. Here, we used the siEhQTRT1 plasmid to silence the expression of EhQTRT1. Following transfection of trophozoites with the siEhQTRT1 plasmid and the selection of a population resistant to G418 (6 μg/ml), the siEhQTRT1 plasmid was cured by removal of G418 from the culture for a month. Silencing of EhQTRT1 in the cured EhQTRT1 silenced trophozoites was confirmed by immunoblotting using an antibody against EhTGT (Fig 3A) and by northern blot by using a radiolabeled probe against EhQTRT1 (Fig 4). Using densitometry analysis, we observed that the EhTGT complex in siEhQTRT1 trophozoites was inhibited up to 80% at the protein level (Fig 3 B) and up to 90% at the mRNA level (Fig 4B). The cellular localization of EhTGT in control trophozoites and siEhQTRT1 trophozoites has been performed by confocal immunofluorescence microscopy. In control trophozoites, EhTGT has a cytoplasmic localization which is barely detectable in siEhQTRT1 trophozoites (Fig 5A &B).

We have previously demonstrated that EhTGT can incorporate queuine into *E.histolytica* tRNAs. Consequently, we assume that siEhQTRT1 trophozoites will have a reduced ability to incorporate queuine in their tRNAs. To test this assumption, we determine the level of queuine incorporated into $tRNA_{His}$ by using N-acrylolyl-3-aminophenylboronic acid (APB) polyacrylamide gel to separate queuosine-modified- tRNAs from unmodified tRNAs. We observed a significant reduction of the level of queuosine-modified- $tRNA_{His}$ in siEhQTRT1 trophozoites grown with queuine when compared to control trophozoites grown with queuine (Fig 5 C and D). The amount of m^5C_{38} methylation in $tRNA^{Asp}$ in siEhQTRT1 trophozoites grown with queuine was also significantly reduced (25±7.07 %) compared to the level in control trophozoites grown with queuine (55±7.07%) (Fig 1B).

We also observed that siEhQTRT1 trophozoites have their growth impaired by 50% when compared to the growth rate of the control trophozoites (Fig 6A). Queuine has no significant effect on the growth of control or siEhQTRT1 trophozoites (Fig 6A).

Queuine induces the resistance of *E.histolytica* to OS.

Having established that queuine supplementation leads to increased Q levels in tRNA mediated by EhTGT, we next set out to define the effect of queuine on the parasite phenotype, possibly related to Q-modified tRNA. In order to determine if queuine regulates the resistance of the parasite to OS, we grew *E. histolytica* trophozoites with and without 0.1 μM queuine and then determined H_2O_2 cytotoxicity. We observed that the LD_{50} of H_2O_2 of trophozoites grown with queuine (3.3±0.05 mM) is significantly higher than in the absence of queuine (2.4± 0.3 mM) (Fig 7). These results indicate that queuine protects the parasite against OS. Next we asked whether the beneficial effect that queuine has

on the resistance of *E.histolytica* to OS (Fig 7) depends on EhTGT. For this purpose, we determined the resistance of siEhQTRT1 trophozoites grown with queuine and then exposed to OS (Fig 7). We observed that queuine does not protect siEhQTRT1 trophozoites against OS.

Queuine regulates protein synthesis during OS

The protective effect that queuine has on the parasite exposed to OS suggests that protein synthesis is preserved in the parasite exposed to queuine and OS. In order to test this hypothesis, we used the surface sensing of translation assay (SUnSET) [27] to determine the amount of puromycin that was incorporated into nascent peptide chains (Fig 8A&B). As previously described [28], we found that protein synthesis is strongly inhibited in trophozoites exposed to OS (Fig 8B). In contrast, we found that protein synthesis of trophozoites grown with queuine and then exposed to OS is comparable to protein synthesis of control trophozoites (Fig 8B). Surprisingly, we observed that trophozoites grown with queuine but not exposed to OS (Fig 8B) have reduced protein synthesis. These results indicate that queuine not only regulates total protein synthesis but it also inhibits protein synthesis in response to OS. Next we asked whether the regulatory effect that queuine has on protein synthesis depends on EhTGT. For this purpose, SunSET assay was performed on siEhQTRT1 trophozoites grown with and without queuine and then exposed or not to OS (Fig 9A&B). We observed that queuine does not regulate protein synthesis in siEhQTRT1 trophozoites.

Queuine impairs cytopathic activity and the survival of the parasite within the mouse caecum.

It has been previously shown that TGT and presumably queuine regulates the virulence of *Shigella* by controlling the translation of VirF, a central transcriptional regulator of virulence factors involved in cellular invasion and spreading of this bacteria [29]. In order to test whether queuine has an effect on the virulence of the parasite, we determined the ability of *E. histolytica* trophozoites grown with and without queuine to destroy a monolayer of mammalian cells (cytopathic activity) (Fig 10A). We found that the cytopathic activity of trophozoites grown in presence of queuine was impaired compared to that of trophozoites grown in absence of queuine.

Next, we asked whether the detrimental effect that queuine has on the cytopathic activity of *E.histolytica* (Fig 10A) depends on EhTGT. We found that the cytopathic activity of siEhQTRT1 trophozoites is significantly impaired (Fig 10A) compared to that of control trophozoites (Fig 10A). Moreover, the growth of siEhQTRT1 trophozoites in presence of queuine did not have any significant

effect on their cytopathic activity (Fig 10A). In order to determine if the siEhQTRT1 trophozoites require more time to destroy the HeLa monolayer than control trophozoites, we incubated siEhQTRT1 trophozoites grown with and without queuine on the HeLa cell monolayer for 90 minutes instead of 40 minutes. We found that more than 90 % of the HeLa monolayer is destroyed by siEhQTRT1 trophozoites after 90 minutes of incubation (Fig 10A). The same level of monolayer destruction was observed for siEhQTRT1 trophozoites grown with queuine (Fig 10B). These results indicate that queuine impairs the strong cytopathic activity of control trophozoites (Fig 10A) but does not impair the weak cytopathic activity of siEhQTRT1 trophozoites.

To validate the alteration of amoebic pathogenicity *in vivo*, trophozoites grown in presence of queuine were subjected to the mouse intestinal amoebiasis (Fig 10B). In mouse intestinal amebiasis model, control trophozoites or trophozoites grown in presence of queuine were inoculated to susceptible CBA/J mice cecum. After 7 days of post inoculation, the number of trophozoites present in the stool were determined. The average number of detectable trophozoites from mice inoculated with control trophozoites (14 ± 23 trophozoites/mg stool) was higher than that of trophozoites grown in presence of queuine (0.74 ± 0.71 trophozoites/mg stool) (Fig 10B). These results indicate that queuine affects the survival of the parasite in the mouse caecum.

Queuine has a profound impact on the *E. histolytica* transcriptome.

We used RNA sequencing (RNA-seq) to investigate the molecular basis of the effect of queuine on the resistance of *E.histolytica* to OS (Fig 11A and B). RNA-Seq experiments were performed as described previously [10], and the data were analyzed using the generalized linear model implemented in the DESeq2 package in R. This allowed us to perform pairwise comparisons of gene expression by the *E. histolytica* HM1:IMSS strain under the various tested conditions and to probe the interaction between queuine and OS (Fig. 11B).

We first used a principal component analysis to explore the data's structure (Fig. 12A); this showed that the untreated trophozoites group (T) and the trophozoites grown with queuine (Q) form two distinct clusters that can be separated on the first principal component. Stress effect is mainly supported by the second component and OS-treated and -untreated samples are also forming clusters both in T and in Q conditions. Our comparisons revealed that queuine has a profound impact on the *E. histolytica* transcriptome (C1) with 664 downregulated and 548 upregulated genes (Table 2). Details regarding these genes can found in Supplementary Table S1.

Gene categories affected by queuine supplementation

The differentially regulated genes in Q vs T (comparison C1) were classified according to the protein class which they encode (Fig 13A&B) using the Protein ANalysis THrough Evolutionary Relationship (PANTHER) sequence classification tool (version 14.1)[30]. Of the upregulated genes in the presence of queuine (Fig 13A), genes that are associated with nucleic acid binding (GO:0003676) and nucleosome binding (GO:0031491) such as chromatin-specific transcription elongation factor (EHI_109860) and protein HIRA (EHI_131950), heterocyclic compound binding (GO:1901363) such as 60S ribosomal protein L4 (EHI_064710) and ATP-dependent DNA helicase (EHI_119290), ATPase activity (GO:0016887) such as heat shock protein 70 family (EHI_001950) and RecQ family helicase (EHI_028890), oxidoreductase activity (GO:0016491) such as peroxiredoxin (EHI_145840) and aldehyde-alcohol dehydrogenase 2 (EHI_024240), catalytic activity, acting on RNA (GO:0140098) such as regulator of nonsense transcripts (EHI_035550) and RNA lariat debranching enzyme (EHI_062730) , actin binding (GO:0003779) such as coronin (EHI_122800) and villidin (EHI_150430), cytoskeletal protein binding (GO:0008092) and tubulin binding (GO:0015631) such as dynamin-1-like protein (EHI_013180) and kinesin-like protein (EHI_140230), protein-containing complex binding (GO:0044877) such as elongation factor 2 (EHI_189490) and chromatin-specific transcription elongation factor (EHI_109860), catalytic activity, acting on DNA (GO:0140097) such as DNA replication licensing factor (EHI_076880) and RecQ family helicase EHI_028890 are significantly enriched according to the PANTHER statistical overrepresentation test (Fig 13C).

The categories for functional classification of genes downregulated in trophozoites grown with queuine (using the subset of protein class) is shown (Fig 14A) (Supplementary Table S2). The most abundant class of proteins are nucleic acid binding protein (PC00171) (22.8%), protein-binding activity modulators (PC00095) (22%), Protein modifying enzymes (PC00260) (16.5%), metabolite interconversion enzyme (PC00262) (11%), membrane traffic protein (PC00150) (8.7%), transporter (PC00227) (7.9%), translational protein (PC00263) (3.9%), scaffold/adaptor protein (PC00226) (3.1%).

Of the downregulated genes in Q, no enrichment of a specific biological process was detected according to the PANTHER statistical overrepresentation test (Fig 13B).

Gene categories affected by queuine supplementation and OS

As previously described [10], OS strongly modulates the transcriptome of the parasite with 1646 genes that are differentially expressed compared to the T condition (comparison C2). Interestingly, in presence of queuine, OS affects the expression of only 340 genes, suggesting that queuine influences the response of the parasite to OS. We next addressed the influence of queuine on *E.histolytica*'s response to OS (comparison C5). We identified 932 genes (Table 2), and then performed a hierarchical clustering analysis (Fig. 12B). The average profile of each of the four identified clusters was then established (Fig. 12B) (Supplementary Table S3). Clusters 1&4 represent genes that are mostly upregulated in trophozoites exposed to OS but that are either mostly not differentially expressed (cluster 4) or down-regulated (cluster 1) in trophozoites grown with queuine and exposed to OS. Cluster 2&3 represent genes that are mostly downregulated in control trophozoites exposed to OS but that are mostly not differentially expressed in trophozoites grown with queuine exposed to OS. Detailed analysis of these clusters is described in Supplementary File 2.

Comparative analysis of genes upregulated in virulent *E.histolytica* trophozoites vs genes downregulated in trophozoites grown with queuine.

We compared our RNA Seq data containing genes downregulated in trophozoites grown with queuine with the previously published RNA Seq data of genes upregulated in a highly virulent *E.histolytica* HM1:IMSS strain [31]. Our analysis showed that out of 816 upregulated genes present in the virulent strain vs 1238 genes downregulated in trophozoites grown with queuine, a total of 120 genes were common between both sets (Supplementary Table S4.) These genes include putative cysteine protease (EHI_151440), putative thioredoxin (EHI_152600), (EHI_124400), fatty acid elongase (EHI_009370), putative proteasome regulatory subunit (EHI_200220), ubiquitin putative (EHI_156660), putative eukaryotic elongation factors (EHI_006170), and majority of the genes correspond to hypothetical proteins that are shared between the two data sets.

Functional analysis of the proteins in this set showed that the most abundant class of proteins are protein modifying enzyme (PC00260) (19.6%), nucleic acid binding protein (PC00171) (17.6%), protein-binding activity modulator (PC00095) (13.7%), cytoskeletal protein (PC00085) (13.7%), membrane traffic protein (PC00150) (9.8%), translational protein (PC00263) (7.8%), metabolite interconversion enzyme (PC00262) (7.8%), chaperone (PC00072) and chaperone (PC00072) (3.9%), and scaffold/adaptor protein (PC00226) (2%) (Fig 14B).

Taken together, this transcriptome analysis indicates that queuine induces an upregulation of genes involved in OS stress response and the downregulation of genes previously associated with virulence.

Discussion

We have recently demonstrated that *E.histolytica* uses oxaloacetate produced by *E.coli* to protect itself against OS [9]. In this study, we present another facet of this protective effect through the lens of the human microbiome. Queuine released by the gut bacteria has a protective effect on the parasite during OS. Interestingly, this protective effect against OS conferred by queuine is also found cancer cells where the activity of antioxidant enzymes is improved by the supplementation of queuine in the culture [20], suggesting that this effect is universal.

In this work, we have shown for the first time that *E.histolytica* has an active TGT enzyme. This activity is essential for the incorporation of queuine in its tRNA and for the resistance to OS. EhTGT shares structural similarity with its eukaryotic counterpart which is also a heterodimer. It has been proposed that one subunit (QTRT1) is responsible for the recognition of the anticodon loop of the tRNA and for transglycosylase activity, whereas the second subunit (QTRTD1) is responsible to maintain the orientation of the tRNA [32,33]. It has been suggested that the human QTRTD1 salvage queuine from queuosine-5'-monophosphate that is generated after tRNA turnover [34]. Since the amount of queuine which circulates in the range of 1-10 nM [35], it might be possible that in *E.histolytica,* there is a requirement for a second subunit to salvage queuine from microbial tRNAs.

Different mechanisms can explain why queuine protects *E.histolytica* against OS. The first mechanism described in *S.pombe* and in mammals involves the stimulation of Dnmt2 activity by prior queuosine incorporation at G34 of $tRNA^{Asp}_{GAC}$ [36,37]. Dnmt2 mediated C38 methylation of $tRNA^{Asp}_{GAC}$ during OS protects it against cleavage by the RNA endonuclease angiogenin [38,39]. In *E.histolytica*, we have also correlated the hypermethylation of $tRNA^{Asp}_{GAC}$ catalyzed by the *E.histolytica* Dnmt2 homolog Ehmeth with the resistance of the parasite to OS [18]. Then, this specific $tRNA^{Asp}_{GAC}$ is preferentially used by the cells for the translation of stress proteins [23]. In our current study, we have shown that exogenous supplementation of *E.histolytica* trophozoites with queuine leads to hypermethylation of C38 in $tRNA^{Asp}_{GAC}$ and that this hypermethylation depends on the active EhTGT enzyme. Thus, extrapolating this information it is possible that queuine exerts a protective effect on *E.histolytica* during OS via the hypermethylation of C38 in $tRNA^{Asp}_{GAC}$.

 A second mechanism of protection against OS mediated by queuine relies on its ability to prone the

parasite to resist OS. We have reported here that several genes involved in stress response are upregulated in the presence of queuine including heat shock protein 70 (Hsp 70), antioxidant enzymes, and enzymes involved in DNA repair. Hsp70 is known for its role in refolding of denatured and misfolded proteins, and translocation of secretory proteins [40]. In *E.histolytica,* Hsp70 are produced by cells in response to OS, heat shock response, and their levels are also increased during the formation of liver abscess [41-43]. Antioxidant enzymes such as alcohol dehydrogenases (ADH2) are well known for their role in energy metabolism in *E.histolytica* during OS [44] and help the parasite to resist OS [45,46]. Transketolase is another antioxidant enzyme which is involved in the pentose phosphate pathway. This enzyme is essential for the production of the antioxidant NADPH which ultimately protects cancer cells against OS stress [47,48]. The upregulation of the expression of this enzyme in the parasite exposed to queuine (this work) and the accumulation of sedoheptulose-7-phosphate, the product of transketolase, in the parasite exposed to OS [49] suggest that this enzyme is also involved in the resistance of the parasite to OS. RecQ helicases are conserved across all forms of life ranging from prokaryotes to eukaryotes [50]. They have many metabolic functions in the cell but are widely popular for their role in DNA repair and genome stability [51-53]. Mutation in these enzymes lead to chromosomal instability which in turn are associated with many diseases such as cancer [54]. Human RecQ helicase is involved in the repair of oxidative DNA damage [54,55], and it interacts with Poly[ADP-ribose]polymerase1 (PARP-1) both *in vivo* and *in vitro* aiding in the repair [56]. This interaction with PARP-1 is interesting as PARP-1 itself is known for its involvement in DNA repair and also in regulation of transcription in the cell [57]. Based on these observations, we hypothesize that either queuine stimulates the production of DNA repairing enzymes in *E.histolytica* under normal physiological conditions or it prepares the parasite to resist DNA damage in response to stress. These results suggest that queuine has a role in DNA repair damage in the parasite, however, this assumption needs thorough investigation.

In this work, we have shown that queuine impairs the cytopathic activity of the parasite and its survival in the large intestine of mice with experimentally induced amebiasis. The transcriptome of the parasite exposed to queuine provides some insights into the mechanism that leads to this low virulence phenotype. Many protein modifying enzymes such as cysteine proteases, metalloproteases, phosphatases, and ubiquitin ligases have their expression downregulated in presence of queuine. Protein phosphatases have an essential role in cellular signaling pathways and it may also play a role in the pathogenicity of the parasite [58]. GTPases are an essential part of cell signaling events and we

found many genes belonging to putative Ras GTPases, Rho GTPase, and Rab GTPase proteins. Many of these genes have been linked to vesicular trafficking. For example, Rab-GTPases may play an essential role in phagocytosis [59]. Moreover, a RabGTPase in *E.histolytica* (EhRab7A) is involved in the transport of cysteine protease to phagosomes [60]. Further investigation regarding these enzymes will help us define their role in the virulence of the parasite. The most surprising observation in genes downregulated in response to queuine is the presence of 760 genes encoding hypothetical proteins. These proteins whose functions are unknown may probably play important roles in regulating the virulence of the parasite.

Another explanation as to how or why queuine may affect the virulence of the parasite stems from our comparative analysis of genes upregulated in the virulent strain [31] that are downregulated in the presence of queuine. For example, actin which is a major cytoskeletal proteins involved in motility of the parasite and phagocytosis [61] has its expression upregulated in the virulent strain of *E.histolytica* and downregulated in the parasite exposed to queuine. The same trend is observed for cysteine proteases, which are essential virulence factors in the parasite that promote intestinal invasion by degrading the extracellular matrix [62]. Among the 120 genes that are common, 84 genes correspond to hypothetical proteins. Characterization of these genes presents an opportunity to discover their novel functions in the parasite. Moreover, new hypothetical proteins may serve as good markers and act as possible pharmacological targets against this parasite.

Our results may provide some clues to understand why most of the infections with *E.histolytica* are asymptomatic. Factors leading to such outcome are the level of virulence of the parasite, the immune status of the host and most probably the gut microbiota. Different mechanisms may influence the development of amoebiasis by the gut microbiota including a competition with the parasite to invade the colonic mucosa or the boosting of a host immune responses against the parasite [63,64]. Here, we are adding a new mechanism to this list which is based on the downregulation of virulence genes expression by queuine. The fact that queuine confers resistance to OS seems counterintuitive because this resistance often correlates with a stronger virulence of the parasite [65,66]. However, the resistance to OS conferred by queuine is mild if we compare it to the strong resistance conferred by oxaloacetate, an antioxidant metabolite scavenged by the parasite from the gut microbiota [67]. This mild resistance to OS may be enough to help the parasite survive fluctuation of the oxygen content that occurs in the large intestine due to changes in the composition of the microbiota. For example, the depletion of butyrate-producing species following an antibiotic treatment increased epithelial oxygenation and

allows the expansion of aerobic luminal microbes [68]. The price paid by the parasite to acquire a mild resistance to OS is a lower virulence. The concept of fitness cost is well known and it has been described for many microorganisms [69] that have acquired drug resistance including in *E.histolytica* [70].

Queuine regulates the transcription of more than 1200 genes. Different hypothesis can explain the effect of Q on the transcriptome of *E.histolytica*: (i) The presence of queuosine in the anticodon loop position 34 of tRNA[Asp] and maybe other tRNAs has an influence on the transcriptome of *E.histolytica*. This hypothesis is supported by a recent report showing that the deletion of a gene encoding for KEOPS (an enzyme involved in the catalysis of t6A modification in tRNA) in *S.cerevisiae* leads to the upregulation of the transcription of genes related to mitochondrial function and carbohydrate metabolism [71]. The effect of tRNA modifications on transcription can also be indirect when it impairs the translation of a transcription factor like Gcn4 [71]. It is possible that like for Gcn4, a yet to be discovered transcription factor whose translation depends on the presence of Q-modified tRNAs is involved in the regulation of gene expression in Q-trophozoites. It is also possible that Q is regulating transcription by using a mechanism that does not involved the modification of tRNA. An alternative explanation is that queuine regulates the expression of a transcription factor that control the expression of these genes. The best studied example is in the enteropathogenic bacteria, *Shigella flexneri*. VirF has been recognized as the main transcriptional regulator involved in the virulence of this pathogen. Lack of this VirF mRNA (in the bacteria lacking TGT) reduces the virulence of the bacteria. It has been found that VirF mRNA is a substrate of the *E.coli* TGT, since it has extremely high sequence similarity with *shigella* TGT, and that this recognition causes a site-specific modification of a base in the VirF mRNA and modulates its translation[29]. Based on this information, we cannot rule out that EhTGT and queuine work via a mechanism unrelated to tRNA modification.

To conclude, we have investigated for the first time the role of queuine, a compound produced by the gut microbiota, on the physiology of *E.histolytica*. We showed that queuine is incorporated in the tRNA of the parasite and that it prepares the amoeba to resist oxidative stress *in vitro*. We have also shown that the presence of queuine leads to hypermethylation of tRNA[Asp] in the organism and this agrees with other studies indicating that the role of queuine in methylation is conserved across all species and that its presence may influence Dnmt2 activity. Our study also revealed the role of queuine in shaping the transcriptome of the parasite. Our data also support a role of queuine in the resistance of the parasite to OS but this comes at a cost in term of virulence. Anti-virulence therapeutic strategies have been

proposed to fight against drug resistant pathogens [72]. In this regard, queuine provided directly to the patient or by selected probiotics may represent a new therapeutic approach for the treatment of amoebiasis.

Material and methods

E. histolytica culture.

E. histolytica trophozoites HM-1:IMSS strain were grown under axenic condition at 37 °C in Trypticase Yeast Extract Iron Serum (TYI-S-33) medium prepared according to a previously reported protocol [73]. The trophozoites were harvested during the logarithmic phase of growth by chilling the culture tubes at 4 °C and pelleted by centrifugation at 500 g for 5 min. The pellet was washed twice with ice-cold phosphate-buffered saline. Trophozoites were grown with 0.1 µM queuine (a kind gift from Dr. Klaus Reuter, University of Marburg) for 3 d.

Bacterial strains.

The bacterial strains used in this study are *E. coli* BL21(DE3)pLysS and Dh5 alpha. *E. coli* was grown at 37°C in Luria-Bertani (LB) medium [74].

Cultivation of HeLa cells.

HeLa cells (a kind gift from Dr. Kleinberger, Faculty of Medicine, Technion) were maintained in continuous culture using a previously described protocol [75].

Transfection of *E. histolytica* trophozoites.

The transfection of *E. histolytica* trophozoites was done using a previously described protocol [76] .

Effect of queuine on hydrogen peroxide (H_2O_2) resistance.

E. histolytica trophozoites were grown with and without queuine (0.1 µM) for 3 days at 37 ˚C. Trophozoites were harvested and (1×10^6) exposed in TYI-S-33 medium (without serum) to various concentrations of H_2O_2 (0-5 mM) for 30 min at 37 ˚C. The viability of the trophozoites was determined with eosin dye exclusion method [67], with this value used to calculate the median lethal dose (LD_{50}) of H_2O_2.

Measurement of the cytopathic activity of *E. histolytica*.

The cytopathic activity of *E. histolytica* trophozoites was determined using a previously described protocol [77]. Briefly, *E. histolytica* trophozoites (2.5×10^5 cells/ per well) grown with and without queuine, were exposed or not to 2.5mM H_2O_2 for 30 min in TYI-S-33 medium (without serum). After the incubation, the cells were resuspended in fresh TYI-S-33 medium (without serum) prior to their incubation with HeLa cell monolayers in 24-well tissue culture plates at 37°C for 40 min or for 90 min. The incubation was stopped by placing the plates on ice and unattached trophozoites were removed by washing the plates with cold Phosphate Buffer Saline (PBS) supplemented with 1% Galactose. The cells were fixed with 4% formaldehyde for 10 min, and finally washed twice with cold PBS. HeLa cells remaining attached to the plates were stained with methylene blue (0.1% in 0.1 M borate buffer, pH 8.7). The dye was extracted from the stained cells by 0.1 M HCl, and its color intensity was measured spectrophotometrically at OD_{660}. The amount of dye extracted was proportional to the number of intact cells that remained attached to the tissue culture well. The amount of dye extracted from monolayers of tissue-cultured cells that had not interacted with trophozoites served as control (100% of monolayer integrity). The intensity of color was measured in a spectrophotometer (at 660 nm) after appropriate dilution with HCI (0.1 M). Experiments were carried out in duplicate, repeated at least twice, and standard error determined. To decide the exact time at which to terminate an experiment, the progress in tissue culture damage during the incubation with trophozoites was also monitored by periodic observation under an inverted microscope.

Primers used in this study.

The list of primers used in this study are listed in Supplementary File 1.

Ethics statement.

Animal procedures dealing with the mouse model of intestinal amebiasis were approved by the Institutional Animal Care and Use Committee (No. 211075) and conducted at the AAALAC-accredited National Institute of Infectious Diseases, Japan. Animal procedures dealing with the production of EhTGT antibody were approved by the Technion Animal Care and Use Committee (IL-115-10-16) and conducted at the accredited Faculty of Medicine-Technion, Haifa, Israel.

Detection and quantification of *E. histolytica* in infected mice stool.

The virulent trophozoites were passaged *in vivo* through the CBA/J mouse cecum (purchased from Jackson Laboratory, Japan)[78]. For all intracecal injections, 1×10^6 axenic trophozoites with or without 0.1 μM queuine treatment for three days were inoculated into the proximal and apical region of the cecum [79]. To quantify the number of *E. histolytica* trophozoites in stool, real-time quantitative PCR was performed by using Fast SYBR Green Master Mix (ThermoFisher Scientific, USA) in the StepOne Plus Real-Time PCR System (Applied Biosystems, USA). Primer sets were specific to *E. histolytica* 18S rRNA (EntaF and EhR) [67,80]. A standard curve was generated using the DNA extracted from trophozoites serially diluted from 10^5 to 10^0. Using the standard curve and the stool weight, the number of trophozoites/mg stool was calculated.

Construction of His-tagged EhQTRT1.

For construction of the pET28b EhQTRT1 vector for expression in *E. coli*, the full length EhQTRT1 (EHI_035660) was amplified from *E. histolytica* genomic DNA by PCR using the primers, NcoI EhQTRT1 and XhoI EhQTRT1. The PCR product was cloned in a pGEM-T easy vector, digested with NcoI and XhoI, and then subcloned into the pET28b vector that was previously linearized with NcoI and XhoI. Primers.

Construction of His-tagged EhQTRT1-EhQTRTD1.

For construction of the pETDuet-1 His-tagged EhQTRT1-EhQTRTD1 vector, EhQTRT1 was amplified from *E. histolytica* genomic DNA with the primers BamHI EhQTRT1 and NotI EhQTRT1. The PCR product was cloned in a pGEM-T easy vector, digested with BamHI and NotI, and then subcloned in a pDuet expression vector that was previously linearized with BamHI and NotI to generate the pETDuet-1 His-tagged EhQTRT1 vector. EhQTRTD1 (EHI_079900) was then amplified from *E. histolytica* cDNA by using primers BglII EhQTRTD1 and KpnI EhQTRTD1 primers, and then cloned in pGEM-T easy vector, digested with BglII and KpnI, and then subcloned in the pETDuet-1 His-tagged EhQTRT1 vector that was previously linearized with BglII and KpnI.

Construction of His-tagged ΔEhQTRT1-EhQTRTD1.

For the construction of the pETDuet-1 his-tagged ΔEhQTRT1-EhQTRTD1 lacking the binding region S102- L125 in EhQTRT1, the PCR amplification was done in two parts. First, the EhQTRT1 was

amplified from *E.histolytica* genomic DNA using primers EhQTRT1 5' and 1-101-EhQTRT1 XhoI 3'. The second part of EhQTRT1 was amplified using genomic DNA using primers 126-EhQTRT-XhoI and Standard Reverse Primer EhQTRT1. Both PCR products were purified using QIA PCR purification kit (Qiagen), and digested with XhoI for 1 h at 37°C and purified. The purified products were ligated at 16°C overnight using T4 ligase. Following ligation, the product was amplified by PCR using primers BamHI EhQTRT1 and NotI EhQTRT1. The PCR product was cloned in pGEM-T easy vector, digested with BamHI and NotI, and further subcloned into pETDuet-1 vector that was previously linearized with BamHI and NotI. Following this, the EhQTRTD1 gene was cloned into pETDuet-1 vector as described above. The plasmids were sequenced to ensure the deletion of the sequence and to ensure the presence of unwanted mutations.

Construction of His-tagged mEhQTRT1-EhQTRTD1.

For the construction of the pETDuet-1 his-tagged mEhQTRT1-EhQTRTD1 having a point mutation of D267 to A267 in EhQTRT1, EhQTRT1 was amplified from *E.histolytica* genomic DNA using primers Standard EhQTRT1 and 1-265-EhQTRT1 XhoI. The second part of EhQTRT1 was amplified using genomic DNA using primers 266-EhQTRT-XhoI (where the Aspartate was replaced with Alanine in the primer) and Reverse Primer EhQTRT1. Both PCR products were purified using QIA PCR purification kit (Qiagen), and digested with XhoI for 1 hour at 37°C and purified. The purified products were ligated at 16°C overnight using T4 ligase. Following ligation, the product was amplified by PCR using primers BamHI EhQTRT1 and NotI EhQTRT1. The PCR product was cloned in pGEM-T easy vector, digested with BamHI and NotI, and further subcloned into pETDuet-1 vector that was previously linearized with BamHI and NotI. Following this, the EhQTRTD1 gene was cloned into pETDuet-1 vector as described above. The resulting plasmid was sequenced to ensure the presence of the desired mutation.

Construction of silenced EhQTRT1.

For the construction of siEhQTRT1, the entire protein coding region of EhQTRT1 was amplified from *E.histolytica* genomic DNA with 5' EhQTRT1 BglII and 3' EhQTRT1 XhoI primers. The resulting PCR product was cloned in pGEM-T easy vector system (Promega), digested with BglII and XhoI, and then subcloned in pEhEx-04-trigger vector containing a 142bp trigger region (EHI_048660), to generate siEhQTRT1.

Plasmids were sequenced to ensure the presence of unwanted mutations.

Growth Rate Assays.

Trophozoites (3×10^3/ml) were grown in a 13 ml tube in TYI-S-33 medium at 37 $^{\textbf{o}}$C and the number of viable trophozoites were counted at 24, 48, and 72 h, respectively by using the eosin dye exclusion method.

Preparation of recombinant EhTGT complex.

Recombinant EhTGT complex was expressed as His-tagged protein in *E. coli* BL21(DE3)pLysS competent cells, which were transfected with the pETDuet-1 vector derived plasmids. The overnight culture was supplemented with Ampicillin (100 μg/ml) and grown at 37 °C. The culture was grown with the addition of 100 μM of $ZnSO_4$ until the OD_{600} reached 0.6. The synthesis of the His-tagged protein complex was initiated by adding isopropyl β -D-1-thiogalactopyranoside (IPTG) to the culture at a final concentration of 1 mM. After an overnight incubation in the presence of IPTG at 22 °C, the bacteria were harvested and lysed as described elsewhere [81]. The proteins were then eluted with elution buffer (50 mM NaH_2PO_4, pH 8.0, 300 mM NaCl, and 250 mM imidazole).

Preparation and purification of tRNA^{Asp}.

A commercially synthesized DNA oligomer containing the tRNA^{Asp} sequence of *E. histolytica* and the T7 promoter sequence were amplified by using PCR using primers- Universal primer 5' tRNA^{Asp} and tRNA^{Asp} 3' under the following conditions: under the following conditions: primers (20 pmol each), tRNA^{Asp} template (500 ng), Mg^{2+} (2 mM), dNTPs (0.5 mM each), ThermolPol Taq buffer (1×), Taq DNA polymerase (2 U), in a final volume of 50 μL. The samples were treated at 94 °C (5 min), followed by 29 cycles in the following order- 94 °C (45 s), 60 °C (45 s), 72 °C (45 s). A final extension reaction was allowed to occur at 72 °C for 10 min. The PCR product was purified by using the QIAquick PCR purification kit (Quiagen). The tRNA^{Asp} was subsequently generated by in vitro transcription. The conditions for a 50μl reaction are as follows: T7 RNA Pol Buffer (10X)- 5μl, rNTPs (25mM each)- 4μl, Template- 300- 400ng, T7 RNA Polymerase- 4μl, and purified water up to the final volume. The reaction was incubated overnight at 37°C, after which white precipitates (magnesium pyrophosphate) were removed by centrifugation (13,000 rpm, 5 min). DNAse I was added to remove any unwanted DNA templates in the reaction and incubated for another 30 minutes. The tRNA transcript was ethanol-

precipitated at −20°C and then pelleted by centrifugation (20,000x*g*, 30 min, 4 °C). The resulting pellet was resuspended in DEPC water and the concentration was measured using a nanodrop (Thermo Scientific).

Enzymatic assay with [^{14}C] guanine.

The enzyme activity of the EhTGT complex was performed as described elsewhere using 8-[^{14}C]-guanine to quantify activity by scintillation counting [34].

Production of EhTGT antibody.

Female BALB/c mice were injected intraperitoneally with 700 μg recombinant His-tagged TGT protein that was emulsified in complete Freund's adjuvant (Sigma-Aldrich). Two weeks later, the mice were injected with 700 μg of His-tagged TGT complex protein in incomplete Freund's adjuvant (Sigma-Aldrich). One week after the 6-week injection, the mice were anesthetized and their blood was collected as described elsewhere [81]. Serum that was obtained from mice that were not injected with His-tagged EhTGT complex was used as the control.

Western blotting.

Total protein extracts of *E. histolytica* trophozoites were prepared according a published method [82]. Proteins (40 μg) in the total extract were resolved on a 10% SDS-PAGE in SDS-PAGE running buffer (25 mM Tris, 192 mM glycine, 0.1% SDS). The resultant protein bands were visualized after staining with Ponceau red stain. Next, proteins were electrotransferred in protein transfer buffer (25 mM Tris, 192 mM glycine, 20% methanol, pH 8.3) to nitrocellulose membranes (Whatman, Protran BA83). The blots were first blocked using 3% skim milk, and then probed with 1:1000 mouse polyclonal EhTGT (complex) antibody or 1:1000 mouse monoclonal Actin antibody, clone c-4 (MP biotechnologies, CA, USA) for 16 h at 4 °C. After incubation with one of the previously described primary antibodies, the blots were incubated with 1:5000 secondary antibody for one hour at RT (Jackson ImmunoResearch), and then developed using enhanced chemiluminescence (Bio RAD).

SUnSET assay to measure protein synthesis.

Trophozoites (2×10^6/ml) were grown with and without queuine ((0.1μM) for 3 days at 37˚C) were exposed or not to 2.5mM H_2O_2 for 20 min at 37°C. After treatment with H_2O_2, the trophozoites were

incubated with 10 μg/ml puromycin (Sigma) for 20 min. For pretreatment of the trophozoites with cycloheximide (Sigma), the trophozoites were incubated with 100 μg/ml cycloheximide for 5 min before the addition of puromycin. The trophozoites were lysed with 1% Igepal (Sigma) in phosphate-buffered saline (PBS). Puromycin was detected by immunoblotting with a 12D10 clone monoclonal puromycin antibody (Millipore).

RNA extraction.

Total RNA was extracted from control trophozoites, trophozoites treated with queuine (0.1μM for three days at 37°C), and then exposed (or not) to H_2O_2 (in three biological replicates) using the TRI reagent kit. according to the manufacturer's instructions (Sigma-Aldrich). Libraries were built using a Truseq mRNA-Seq Library Preparation Kit (Illumina), according to the manufacturer's recommendations. Quality control was performed on an Agilent Bioanalyzer. Sequencing was performed on a HiSeq 2500 system (Illumina) and produced 65-base single-end reads.

RNA-Seq data analysis.

Bioinformatics analysis was performed using the RNA-seq pipeline from Sequana [83]. Reads were cleaned of adapter sequences and low-quality sequences using cutadapt version 1.11 [84]. Only sequences at least 25 nt in length were considered for further analysis. Bowtie version 1.2.2 [85], with default parameters, was used for alignment on the reference genome (*Entamoeba histolytica*, from amoebaDB). Genes were counted using featureCounts version 1.4.6-p3 [86] from the Subreads package (parameters: -t gene -g ID -s 1).

Count data were then analyzed using R version 3.5.3[12] and the Bioconductor package DESeq2 version 1.22.2[29]. Normalization and dispersion estimation were performed with DESeq2 (using the default parameters) and statistical tests for differential expression were performed applying the independent filtering algorithm. A generalized linear model including (i) the effect of stress (+OS vs -OS), (ii) the presence of queuine (Q vs. T) and (iii) the interaction term was set up in order to test for inter-condition differences in expression and to test the interaction between stress and queuine. For each pairwise comparison, raw p-values were adjusted for multiple testing using the Benjamini and Hochberg procedure [87]. Genes with an adjusted p-value < 0.05 were considered to be differentially expressed. RNA-Seq data have been deposited at the Gene Expression Omnibus (http://www.ncbi.nlm.nih.gov/geo) under the accession number GSE142211.

Principal component analysis and hierarchical clustering were performed using variance-stabilizing transformed counts. Hierarchical clustering of the genes was based on the correlation distance and the Ward aggregation criterion. The average profile was established by computing the mean $\log_2$-normalized count over the genes in each of the four identified clusters.

The functional classification, GO term analysis and protein class analysis were performed with PANTHER tools (http://pantherdb.org) [88].

Immunofluorescence microscopy.

E. histolytica trophozoites were transferred to microscope slides as described elsewhere [77]. The attached trophozoites were washed in warm (37°C) PBS, fixed with methanol and then permeabilized with FACS buffer (2% bovine serum albumin in PBS that was supplemented with 0.1% tween) as described by [77]. The slides were then blocked with 5% Goat serum in FACS buffer for 15 min at ambient temperature. The samples were then probed with 1:1000 polyclonal mouse EhTGT (complex) antibody. At the end of the incubation, the slides were washed 3 times in FACS buffer, and then incubated with a 1:250 Alexaflour (Jackson ImmunoResearch) for 3 h at 4 °C. At the end of the incubation, the nuclei of the trophozoites were stained with 1:1000 4′, 6-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich). The samples were then washed with FACS buffer and mounted onto microscope slides with mounting medium (Dako). The specimens were then examined under a confocal immunofluorescence microscope (ZEISS- LSM700 Meta Laser Scanning System confocal imaging system) with a 63X oil immersion objective. For densitometry analysis, the intensity of 20 trophozoites per condition was quantified using image J and signal intensities were normalized to the wild type trophozoites.

Northern blot analysis.

Total RNA (10 μg) were suspended in RNA sample buffer comprising of 17.5μl of RNA mix (10 ml RNA Mix: 600μl of 10X MOPS, 2.1 ml Formaldehyde, 6 ml Formamide, 120 μg/ml ethidium bromide) and 2 μl of RNA loading buffer. Samples were gently mixed and incubated at 70°C for 5 min, then cooled immediately on ice and loaded into 1% agarose gel (6.66% formaldehyde in 1X MOPS buffer). The RNA was blotted on to nylon membrane (Amershan Hybond-N) overnight, the membrane was washed briefly in 1xMOPS and cross-linked by UV using 120 milli joules (Stratagene UV linker). Membranes were hybridized overnight with [32]-P labelled EhQTRT1 or enolase probes. The membrane

was washed with two washing buffers (Washing buffer 1:40mM NaP, 1mM EDTA, 5%SDS Washing buffer 2: 40mM NaP, 1mM EDTA, 1% SDS). Images were developed using a phosphor imager and quantification of signal density from two independent experiments was performed using ImageJ.

Detection of queuosine in tRNA by chromatography-coupled tandem quadrupole mass spectrometry (LC-MS/MS).

E.histolytica trophozoites were grown with and without queuine (0.1μM for three days) and total RNA was isolated using the TRI reagent kit according to the manufacturer's instructions (Sigma-Aldrich). Three biological replicates from each treatment condition were processed. Samples were processed as blind samples. tRNA was then purified by size-exclusion HPLC performed using a Bio SEC-3 300Å, 7.8 X 300 (internal diameter X length, mm) with 100 mM ammonium acetate pH 7.4 as the mobile phase. Purity of isolated tRNA species was assessed on a Bioanalyzer 2100 system. The average yield of purified tRNA collected ranged from 0.6 to 0.8 mg per sample. Purified tRNA (5μg) from each sample was hydrolyzed in a digestion cocktail containing 10 U benzonase, 5 U bacterial alkaline phosphatase, 0.05 U phosphodiesterase I, 50 μM desferroxamine, 0.5 μg pentostatin, 50 μM butylated hydroxytoluene, 0.25 μg tetrahydrouridine, and 2.5 nmol of internal standard $[^{15}N]_5$dA in 10 mM Tris buffer, pH 7.9. The digestion mixture was incubated at 37 °C for 3 h. Digestion mixtures were subsequently removed by passing the mixture through 10 kDa MWCO spin filter. Samples were lyophilized and reconstituted at final concentration of 100 ng/μl and analyzed by LC-MS/MS. The ribonucleosides were resolved on a Thermo Scientific Hypersil aQ GOLD column (2.1 mm x 100 mm; 1.9 μm particle size) using a mobile phase gradient consisting of 0.1% formic acid as solvent A and acetonitrile with 0.1% formic acid as solvent B at flow rate of 0.3 mL/min and 25 °C using the following elution gradient for solvent B: 0-12 min, 0%; 12-15.3 min, 0-1%; 15.3-18.7 min, 1-6%; 18.7-20 min, 6%; 20-24 min, 6-100%; 24-27.3 74 min, 100%; 27.3-36 min 100-0%; 36-41 min, 0%. The coupled 6490 Triple Quadrupole LC/MS mass spectrometer was operated with the following optimized source parameters: gas temperature, 50 °C; gas flow, 11 L/min; nebulizer pressure, 20 psi; sheath gas temperature, 300 °C; capillary voltage, 1800 V; nozzle voltage, 2000 V; iFunnel RF voltage, 150/50 V and fragmentor voltage, 380 V. Based on a synthetic standard, Q was detected at a retention time of 11.5 min and identified based on two transitions: precursor *m/z* 410, products *m/z* 163 and 295. Quantitation of Q was achieved by normalizing the area under curve (AUC) for the signal intensity of Q against the total AUC for canonical ribonucleosides U, C, A and G for each injected sample. Analysis

of each sample condition was performed with 12 biological replicates and data presented as mean and standard error (SEM).

Bisulfite sequencing of tRNA^Asp.

Total RNA isolation and bisulfite conversions were done using a previously described protocol (Schaefer et al. 2009). Bisulfite-treated tRNAs were reverse transcribed by using a tRNA 3′-specific stem-loop primer and amplified with primers that bind only to the deaminated sequences at the 5′ end (Universal primer and a specific tRNA^Asp stem-loop primer). Amplicons (20) under each condition were subcloned in pGEM-T Easy (Promega) and sequenced (Multi-Disciplinary Laboratories Unit, Bruce Rappaport Faculty of Medicine, Technion).

APB northern blot.

Acryloyl aminophenylboronic acid gels were prepared and run with a few modifications according to Igloi and Kossel [89]. Briefly, 15 μg of RNA was deacetylated in 100 mM Tris–HCl pH 9 for 30 min at 37 °C. RNA was ethanol-precipitated and resuspended in 10μl DEPC water and 1× RNA-loading dye (Fermentas). Samples were then denatured for 10 min at 70 °C and run at 4 °C on Tris Acetate EDTA (TAE), 8 M urea, 15% acrylamide, and 5 mg/ml aminophenylboronic acid (Sigma). Gels were blotted, hybridized, and quantified as a conventional northern blot.

Modeling of EhQTRT1 and EhQTRTD1.

Swiss Model & Phyre servers were used for template assessment & modelling of the homo dimeric interface of the protein structures. The structure of EhQTRT1 was modelled obtained using combination of homology and ab intio model building approaches. The structure was modelled with high confidence using the high-resolution crystal structure of queuine trna-ribosyltransferase (ec 2.4.2.29)2 (trna-guanine (tm1561) from *thermotoga maritima* (PDB 2ASH). The structure of EhQTRTD1 was modelled using the crystal structure of QTRT2, the non-catalytic subunit of murine tRNA-Guanine transglycosylase (PDB 6FV5).

Modeling of the EhTGT complex.

The homodimers models of both the proteins EhQTRT1 and EhQTRTD1 were energy minimized using AMBER 14 molecular dynamics software (https://ambermd.org/index.php). Using the lowest energy

structures, we performed the protein – Protein docking experiment using the cluspro server (https://cluspro.org/login.php). We performed docking checking for the all the possible conformations and defining the active site. The cluspro algorithm considers greedy clustering of ligand positions after obtaining the best poses by rotating the ligand using a grid. We have used the best poses from the top two clusters. The images were created using the pymol software (https://pymol.org/2/).

Data Availability

RNA-Seq data have been deposited at the Gene Expression Omnibus under the accession number GSE142211.

Acknowledgement

We would like to thank Dr. Klaus Reuter, Marburg for providing us queuine to conduct this study. We also thank Yuko Umeki for technical assistance, and the staff of the Microscopy Imaging Laboratory in the Faculty of Medicine, Technion for help with their confocal microscopy resources, and the excellent support in image recording and analysis.

Additional Information

Supplementary data are available.

Funding

The work was supported by the Israel Ministry of Health within the framework European Research Area NETwork Infect-ERA (031L0004; AMOEBAC project), the Israel Science Foundation (260/16), the ISF-NRF program (3208/19), the National Research Foundation of Singapore through the Singapore-MIT Alliance for Research and Technology Antimicrobial Resistance IRG, the Rappaport Institute, the US–Israel Binational Science Foundation (2015211), and the Niedersachsen program.

Conflict of Interest

None

References

1. Mirelman D (1987) Ameba-bacterium relationship in amebiasis. Microbiol Rev 51: 272-284.
2. Bracha R, Kobiler D, Mirelman D (1982) Attachment and ingestion of bacteria by trophozoites of Entamoeba histolytica. Infect Immun 36: 396-406.
3. Bhattacharya A, Ghildyal R, Prasad J, Bhattacharya S, Diamond LS (1992) Modulation of a surface antigen of Entamoeba histolytica in response to bacteria. Infect Immun 60: 1711-1713.
4. Padilla-Vaca F, Ankri S, Bracha R, Koole LA, Mirelman D (1999) Down regulation of Entamoeba histolytica virulence by monoxenic cultivation with Escherichia coli O55 is related to a decrease in expression of the light (35-kilodalton) subunit of the Gal/GalNAc lectin. Infect Immun 67: 2096-2102.
5. Galvan-Moroyoqui JM, Del Carmen Dominguez-Robles M, Franco E, Meza I (2008) The interplay between Entamoeba and enteropathogenic bacteria modulates epithelial cell damage. PLoS Negl Trop Dis 2: e266.
6. Bracha R, Mirelman D (1984) Virulence of Entamoeba histolytica trophozoites. Effects of bacteria, microaerobic conditions, and metronidazole. Journal of Experimental Medicine 160: 353-368.
7. Verma AK, Verma R, Ahuja V, Paul J (2012) Real-time analysis of gut flora in Entamoeba histolytica infected patients of Northern India. BMC Microbiol 12: 183.
8. Byers J, Faigle W, Eichinger D (2005) Colonic short-chain fatty acids inhibit encystation of Entamoeba invadens. Cell Microbiol 7: 269-279.
9. Shaulov Y, Shimokawa C, Trebicz-Geffen M, Nagaraja S, Methling K, et al. (2018) Escherichia coli mediated resistance of Entamoeba histolytica to oxidative stress is triggered by oxaloacetate. PLOS Pathogens 14: e1007295.
10. Varet H, Shaulov Y, Sismeiro O, Trebicz-Geffen M, Legendre R, et al. (2018) Enteric bacteria boost defences against oxidative stress in Entamoeba histolytica. Sci Rep 8: 9042.
11. Yuan Y, Zallot R, Grove TL, Payan DJ, Martin-Verstraete I, et al. (2019) Discovery of novel bacterial queuine salvage enzymes and pathways in human pathogens. Proc Natl Acad Sci U S A.
12. Katze JR, Basile B, McCloskey JA (1982) Queuine, a modified base incorporated posttranscriptionally into eukaryotic transfer RNA: wide distribution in nature. Science 216: 55-56.
13. Ott G, Kersten H, Nishimura S (1982) Dictyostelium discoideum: a useful model system to evaluate the function of queuine and of the Q-family of tRNAs. FEBS Lett 146: 311-314.
14. Farkas WR (1980) Effect of diet on the queuosine family of tRNAs of germ-free mice. J Biol Chem 255: 6832-6835.
15. Klebe G (2000) Recent developments in structure-based drug design. J Mol Med (Berl) 78: 269-281.
16. Hortner SR, Ritschel T, Stengl B, Kramer C, Schweizer WB, et al. (2007) Potent inhibitors of tRNA-guanine transglycosylase, an enzyme linked to the pathogenicity of the Shigella bacterium: charge-assisted hydrogen bonding. Angew Chem Int Ed Engl 46: 8266-8269.
17. Tuorto F, Legrand C, Cirzi C, Federico G, Liebers R, et al. (2018) Queuosine-modified tRNAs confer nutritional control of protein translation. EMBO J.
18. Fisher O, Siman-Tov R, Ankri S (2006) Pleiotropic phenotype in Entamoeba histolytica overexpressing DNA methyltransferase (Ehmeth). Mol Biochem Parasitol 147: 48-54.
19. Schaefer M, Pollex T, Hanna K, Tuorto F, Meusburger M, et al. (2010) RNA methylation by Dnmt2 protects transfer RNAs against stress-induced cleavage. Genes Dev 24: 1590-1595.
20. Pathak C, Jaiswal YK, Vinayak M (2008) Queuine promotes antioxidant defence system by activating cellular antioxidant enzyme activities in cancer. Biosci Rep 28: 73-81.
21. Muller M, Legrand C, Tuorto F, Kelly VP, Atlasi Y, et al. (2019) Queuine links translational control in eukaryotes to a micronutrient from bacteria. Nucleic Acids Res.

22. Muller M, Hartmann M, Schuster I, Bender S, Thuring KL, et al. (2015) Dynamic modulation of Dnmt2-dependent tRNA methylation by the micronutrient queuine. Nucleic Acids Res.

23. Hertz R, Tovy A, Kirschenbaum M, Geffen M, Nozaki T, et al. (2014) The Entamoeba histolytica Dnmt2 homolog (Ehmeth) confers resistance to nitrosative stress. Eukaryot Cell 13: 494-503.

24. Schulz EC, Roth HM, Ankri S, Ficner R (2012) Structure analysis of Entamoeba histolytica DNMT2 (EhMeth). PLoS One 7: e38728.

25. Tovy A, Hofmann B, Helm M, Ankri S (2010) In vitro tRNA methylation assay with the Entamoeba histolytica DNA and tRNA methyltransferase Dnmt2 (Ehmeth) enzyme. J Vis Exp.

26. Morf L, Pearson RJ, Wang AS, Singh U (2013) Robust gene silencing mediated by antisense small RNAs in the pathogenic protist Entamoeba histolytica. Nucleic Acids Res 41: 9424-9437.

27. Schmidt EK, Clavarino G, Ceppi M, Pierre P (2009) SUnSET, a nonradioactive method to monitor protein synthesis. Nat Methods 6: 275-277.

28. Shahi P, Trebicz-Geffen M, Nagaraja S, Alterzon-Baumel S, Hertz R, et al. (2016) Proteomic Identification of Oxidized Proteins in Entamoeba histolytica by Resin-Assisted Capture: Insights into the Role of Arginase in Resistance to Oxidative Stress. PLoS Negl Trop Dis 10: e0004340.

29. Hurt JK, Olgen S, Garcia GA (2007) Site-specific modification of Shigella flexneri virF mRNA by tRNA-guanine transglycosylase in vitro. Nucleic Acids Res 35: 4905-4913.

30. Mi H, Huang X, Muruganujan A, Tang H, Mills C, et al. (2017) PANTHER version 11: expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool enhancements. Nucleic Acids Res 45: D183-D189.

31. Weber C, Koutero M, Dillies MA, Varet H, Lopez-Camarillo C, et al. (2016) Extensive transcriptome analysis correlates the plasticity of Entamoeba histolytica pathogenesis to rapid phenotype changes depending on the environment. Sci Rep 6: 35852.

32. Xie W, Liu X, Huang RH (2003) Chemical trapping and crystal structure of a catalytic tRNA guanine transglycosylase covalent intermediate. Nat Struct Biol 10: 781-788.

33. Ritschel T, Atmanene C, Reuter K, Van Dorsselaer A, Sanglier-Cianferani S, et al. (2009) An integrative approach combining noncovalent mass spectrometry, enzyme kinetics and X-ray crystallography to decipher Tgt protein-protein and protein-RNA interaction. J Mol Biol 393: 833-847.

34. Chen YC, Kelly VP, Stachura SV, Garcia GA (2010) Characterization of the human tRNA-guanine transglycosylase: confirmation of the heterodimeric subunit structure. RNA 16: 958-968.

35. Fergus C, Barnes D, Alqasem MA, Kelly VP (2015) The queuine micronutrient: charting a course from microbe to man. Nutrients 7: 2897-2929.

36. Muller M, Hartmann M, Schuster I, Bender S, Thuring KL, et al. (2015) Dynamic modulation of Dnmt2-dependent tRNA methylation by the micronutrient queuine. Nucleic Acids Res 43: 10952-10962.

37. Ehrenhofer-Murray AE (2017) Cross-Talk between Dnmt2-Dependent tRNA Methylation and Queuosine Modification. Biomolecules 7.

38. Nawrot B, Sochacka E, Duchler M (2011) tRNA structural and functional changes induced by oxidative stress. Cell Mol Life Sci 68: 4023-4032.

39. Wang X, Matuszek Z, Huang Y, Parisien M, Dai Q, et al. (2018) Queuosine modification protects cognate tRNAs against ribonuclease cleavage. RNA 24: 1305-1313.

40. Radons J (2016) The human HSP70 family of chaperones: where do we stand? Cell Stress Chaperones 21: 379-404.

41. Akbar MA, Chatterjee NS, Sen P, Debnath A, Pal A, et al. (2004) Genes induced by a high-oxygen environment in Entamoeba histolytica. Mol Biochem Parasitol 133: 187-196.

42. Weber C, Guigon G, Bouchier C, Frangeul L, Moreira S, et al. (2006) Stress by heat shock induces massive down regulation of genes and allows differential allelic expression of the Gal/GalNAc lectin in Entamoeba histolytica. Eukaryot Cell 5: 871-875.

43. Santos F, Nequiz M, Hernandez-Cuevas NA, Hernandez K, Pineda E, et al. (2015) Maintenance of intracellular hypoxia and adequate heat shock response are essential requirements for pathogenicity and virulence of Entamoeba histolytica. Cell Microbiol 17: 1037-1051.

44. Pineda E, Encalada R, Rodriguez-Zavala JS, Olivos-Garcia A, Moreno-Sanchez R, et al. (2010) Pyruvate:ferredoxin oxidoreductase and bifunctional aldehyde-alcohol dehydrogenase are essential for energy metabolism under oxidative stress in Entamoeba histolytica. FEBS J 277: 3382-3395.

45. Espinosa A, Clark D, Stanley SL, Jr. (2004) Entamoeba histolytica alcohol dehydrogenase 2 (EhADH2) as a target for anti-amoebic agents. J Antimicrob Chemother 54: 56-59.

46. Espinosa A, Yan L, Zhang Z, Foster L, Clark D, et al. (2001) The bifunctional Entamoeba histolytica alcohol dehydrogenase 2 (EhADH2) protein is necessary for amebic growth and survival and requires an intact C-terminal domain for both alcohol dahydrogenase and acetaldehyde dehydrogenase activity. J Biol Chem 276: 20136-20143.

47. Diaz-Moralli S, Tarrado-Castellarnau M, Alenda C, Castells A, Cascante M (2011) Transketolase-like 1 expression is modulated during colorectal cancer progression and metastasis formation. PLoS One 6: e25323.

48. Xu IM, Lai RK, Lin SH, Tse AP, Chiu DK, et al. (2016) Transketolase counteracts oxidative stress to drive cancer development. Proc Natl Acad Sci U S A 113: E725-734.

49. Husain A, Sato D, Jeelani G, Soga T, Nozaki T (2012) Dramatic increase in glycerol biosynthesis upon oxidative stress in the anaerobic protozoan parasite Entamoeba histolytica. PLoS Negl Trop Dis 6: e1831.

50. Cobb JA, Bjergbaek L (2006) RecQ helicases: lessons from model organisms. Nucleic Acids Res 34: 4106-4114.

51. Sharma S, Doherty KM, Brosh RM, Jr. (2006) Mechanisms of RecQ helicases in pathways of DNA metabolism and maintenance of genomic stability. Biochem J 398: 319-337.

52. Sharma S, Brosh RM, Jr. (2008) Unique and important consequences of RECQ1 deficiency in mammalian cells. Cell Cycle 7: 989-1000.

53. Kawabe T, Tsuyama N, Kitao S, Nishikawa K, Shimamoto A, et al. (2000) Differential regulation of human RecQ family helicases in cell transformation and cell cycle. Oncogene 19: 4764-4772.

54. Brosh RM, Jr., Bohr VA (2007) Human premature aging, DNA repair and RecQ helicases. Nucleic Acids Res 35: 7527-7544.

55. Sharma S, Brosh RM, Jr. (2007) Human RECQ1 is a DNA damage responsive protein required for genotoxic stress resistance and suppression of sister chromatid exchanges. PLoS One 2: e1297.

56. Sharma S, Phatak P, Stortchevoi A, Jasin M, Larocque JR (2012) RECQ1 plays a distinct role in cellular response to oxidative DNA damage. DNA Repair (Amst) 11: 537-549.

57. Krishnakumar R, Kraus WL (2010) The PARP side of the nucleus: molecular actions, physiological outcomes, and clinical targets. Mol Cell 39: 8-24.

58. Anwar T, Gourinath S (2013) Analysis of the Protein phosphotome of Entamoeba histolytica reveals an intricate phosphorylation network. PLoS One 8: e78714.

59. Bosch DE, Siderovski DP (2013) G protein signaling in the parasite Entamoeba histolytica. Exp Mol Med 45: e15.

60. Nakada-Tsukui K, Saito-Nakano Y, Ali V, Nozaki T (2005) A retromerlike complex is a novel Rab7 effector that is involved in the transport of the virulence factor cysteine protease in the enteric protozoan parasite Entamoeba histolytica. Mol Biol Cell 16: 5294-5303.

61. Aguilar-Rojas A, Olivo-Marin JC, Guillen N (2016) The motility of Entamoeba histolytica: finding ways to understand intestinal amoebiasis. Curr Opin Microbiol 34: 24-30.

62. Que X, Reed SL (2000) Cysteine proteinases and the pathogenesis of amebiasis. Clin Microbiol Rev 13: 196-206.

63. Travers MA, Florent I, Kohl L, Grellier P (2011) Probiotics for the control of parasites: an overview. J Parasitol Res 2011: 610769.
64. Nagaraja S, Ankri S (2019) Target identification and intervention strategies against amebiasis. Drug Resist Updat 44: 1-14.
65. Rastew E, Vicente JB, Singh U (2012) Oxidative stress resistance genes contribute to the pathogenic potential of the anaerobic protozoan parasite, Entamoeba histolytica. Int J Parasitol 42: 1007-1015.
66. Pineda E, Perdomo D (2017) Entamoeba histolytica under Oxidative Stress: What Countermeasure Mechanisms Are in Place? Cells 6.
67. Shaulov Y, Shimokawa C, Trebicz-Geffen M, Nagaraja S, Methling K, et al. (2018) Escherichia coli mediated resistance of Entamoeba histolytica to oxidative stress is triggered by oxaloacetate. PLoS Pathog 14: e1007295.
68. Rivera-Chavez F, Zhang LF, Faber F, Lopez CA, Byndloss MX, et al. (2016) Depletion of Butyrate-Producing Clostridia from the Gut Microbiota Drives an Aerobic Luminal Expansion of Salmonella. Cell Host Microbe 19: 443-454.
69. Bleuven C, Landry CR (2016) Molecular and cellular bases of adaptation to a changing environment in microorganisms. Proc Biol Sci 283.
70. Shaulov Y, Nagaraja S, Sarid L, Trebicz-Geffen M, Ankri S (2020) Formation of oxidised (OX) proteins in Entamoeba histolytica exposed to auranofin and consequences on the parasite virulence. Cell Microbiol: e13174.
71. Chou HJ, Donnard E, Gustafsson HT, Garber M, Rando OJ (2017) Transcriptome-wide Analysis of Roles for tRNA Modifications in Translational Regulation. Mol Cell 68: 978-992 e974.
72. Fleitas Martinez O, Cardoso MH, Ribeiro SM, Franco OL (2019) Recent Advances in Anti-virulence Therapeutic Strategies With a Focus on Dismantling Bacterial Membrane Microdomains, Toxin Neutralization, Quorum-Sensing Interference and Biofilm Inhibition. Front Cell Infect Microbiol 9: 74.
73. Diamond LS, Harlow DR, Cunnick CC (1978) A new medium for the axenic cultivation of Entamoeba histolytica and other Entamoeba. Trans R Soc Trop Med Hyg 72: 431-432.
74. Bertani G (1951) Studies on lysogenesis. I. The mode of phage liberation by lysogenic Escherichia coli. J Bacteriol 62: 293-300.
75. Shahi P, Trebicz-Geffen M, Nagaraja S, Hertz R, Alterzon-Baumel S, et al. (2016) N-acetyl ornithine deacetylase is a moonlighting protein and is involved in the adaptation of Entamoeba histolytica to nitrosative stress. Sci Rep 6: 36323.
76. Olvera A, Olvera F, Vines RR, Recillas-Targa F, Lizardi PM, et al. (1997) Stable transfection of Entamoeba histolytica trophozoites by lipofection. Arch Med Res 28 Spec No: 49-51.
77. Trebicz-Geffen M, Shahi P, Nagaraja S, Vanunu S, Manor S, et al. (2017) Identification of S-Nitrosylated (SNO) Proteins in Entamoeba histolytica Adapted to Nitrosative Stress: Insights into the Role of SNO Actin and In vitro Virulence. Front Cell Infect Microbiol 7: 192.
78. Hamano S, Asgharpour A, Stroup SE, Wynn TA, Leiter EH, et al. (2006) Resistance of C57BL/6 mice to amoebiasis is mediated by nonhemopoietic cells but requires hemopoietic IL-10 production. J Immunol 177: 1208-1213.
79. Houpt ER, Glembocki DJ, Obrig TG, Moskaluk CA, Lockhart LA, et al. (2002) The mouse model of amebic colitis reveals mouse strain susceptibility to infection and exacerbation of disease by CD4+ T cells. J Immunol 169: 4496-4503.
80. Hamzah Z, Petmitr S, Mungthin M, Leelayoova S, Chavalitshewinkoon-Petmitr P (2006) Differential detection of Entamoeba histolytica, Entamoeba dispar, and Entamoeba moshkovskii by a single-round PCR assay. J Clin Microbiol 44: 3196-3200.

81. Shahi P, Trebicz-Geffen M, Nagaraja S, Hertz R, Baumel-Alterzon S, et al. (2016) N-acetyl ornithine deacetylase is a moonlighting protein and is involved in the adaptation of Entamoeba histolytica to nitrosative stress. Sci Rep 6: 36323.
82. Lavi T, Isakov E, Harony H, Fisher O, Siman-Tov R, et al. (2006) Sensing DNA methylation in the protozoan parasite Entamoeba histolytica. Mol Microbiol 62: 1373-1386.
83. Cokelaer T, Desvillechabrol, D., LegendreR., Cardon, M. (2017) 'Sequana': a Set of Snakemake NGS pipeline. Journal of Open Source Software 2: 352.
84. Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. 2011 17: 3.
85. Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol 10: R25.
86. Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30: 923-930.
87. Benjamini YH, Y. (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. . Journal of the Royal Statistical Society, series B 57: 289–300.
88. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD (2019) PANTHER version 14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res 47: D419-D426.
89. Igloi GL, Kossel H (1985) Affinity electrophoresis for monitoring terminal phosphorylation and the presence of queuosine in RNA. Application of polyacrylamide containing a covalently bound boronic acid. Nucleic Acids Res 13: 6881-6898.

Tables

Table 1

	Source of the enzyme	Specific Activity of the enzyme (= 1pmol/ min/ mg of protein)
a)	EhTGT	2.28
b)	ΔEhQTRT1-EhQTRTD1	0.63 ± 0.02
c)	mEhQTRT1-EhQTRTD1	0.1 ± 0.02
d)	HIS-tagged EhQTRT1	0

Table 1: Validation of the *in silico* model of EhTGT. Different mutants of the enzyme were tested for their activity by determining their ability to displace guanine in the tRNA$^{Asp}_{GTC}$ anticodon with [8-^{14}C] guanine. Our results indicate that EhTGT is an active enzyme when it forms a functional complex (a), the deletion of the predicted binding region of EhQTRT1 and EhQTRTD1 (S102-L125) makes the enzyme inactive (b), a punctual mutation (D267 to A267) in the predicted active site of EhQTRT1 makes the enzyme inactive without affecting the formation of the EhQTRT1-EhQTRTD1 complex (c). EhQTRT1 subunit expressed alone is inactive (d). The specific activity of EhTGT was significantly different from mEhQTRT1-EhQTRTD1 and ΔEhQTRT1-EhQTRTD1 according to an unpaired student's t test (p≤0.05). The data are expressed as mean ± standard deviation and are representative of three independent experiments performed in duplicates.

Table 2

	Comparison	Modulated Genes		
		# Downregulated	# Upregulated	Total
C1	Q-OS vs T-OS	664	548	1212
C2	T+OS vs T-OS	985	661	1646
C3	Q+OS vs Q-OS	248	92	340
C4	Q+OS vs T+OS	176	265	441
C5	C3 vs C2	485	447	932

Table 2: Number of genes modulated in *E.histolytica* trophozoites in control trophozoites (T) and in trophozoites grown with queuine (q) in response to OS.

Figure Legends

Figure 1: Queuine is efficiently incorporated into *E. histolytica* tRNA and leads to C38 hypermethylation (m⁵C₃₈) in the anticodon loop of tRNA^Asp.

Queuine triggers resistance to OS and it is efficiently incorporated into *E.histolytica* tRNAs.

(A) Quantification of Q in tRNA of *E. histolytica* trophozoites using LC-MS/MS. RNA samples were prepared from control trophozoites (T) and trophozoites grown with queuine (q) supplemented in the culture medium (0.1 µM for 3 days) (T+q). tRNA was purified and analyzed by LC-MS/MS to quantify Q using two transitions: m/z 410 → 163 (left) and m/z 410 → 295 (right). Data represent mean ± SEM for N=12 (Unpaired Student's t-test, p≤0.0001).

B) Bisulfite sequencing of tRNA^Asp $_{GTC}$ has been performed $_{on}$ control trophozoites (T) and trophozoites grown with 0.1µM queuine for three days (T+q). The level of cytosine-38 tRNA^Asp$_{GTC}$ methylation in T+q (55±7.07%;) was significantly different from that of T (20 ± 2 %) according to the analysis with an unpaired Student's t test (p<0.05). Indeed, the level of cytosine-38 tRNA^Asp$_{GTC}$ methylation in siEhQTRT1 trophozoites (siEhQTRT1) and in siEhQTRT1 trophozoites grown with queuine (siEhQTRT1+q) was not significantly different according to the analysis with an Unpaired Student's t test (p<0.05). The data are expressed are mean ± SD obtained from 20 clones each.

Figure 2 : In silico modeling of EhTGT and expression in *E.coli*

(A) *In silico* model analysis showing the interaction of EhQTRT1 and EhQTRTD1 and expression of EhTGT in *E.coli.* The EhQTRT1-EhQRTD1 complex was successfully modelled. EhQTRT1 and EhQTRTD1 were co-expressed in *E.coli* using a pETDuet-1 vector. (B) A size-exclusion chromatogram representing purified EhTGT. (C) An SDS-gel representing fractions (1-4) from the elution corresponding to the peak obtained from size-exclusion chromatography.

Figure 3: **Silencing of EhTGT**

(A): Western blotting was performed on total protein extracts that were prepared from wild type *E. histolytica* trophozoites and EhQTRT1 silenced trophozoites. The proteins were separated on 10% SDS-PAGE gels and analyzed by western blotting using a homemade EhTGT antibody (1:1000) or actin antibody (1:1000). (B): Densitometry analysis showing the percentage of inhibition of the EhTGT complex in EhQTRT1 silenced trophozoites. The data represent mean ± SD for three independent

experiments (Unpaired Student's t test, p = 0.0003).

Figure 4: Determination of the level of EhQTRT1 expression by Northern blot in EhQTRT1 silenced trophozoites. (A) Northern blot analysis of EhQTRT1. Enolase has been used as housekeeping gene because its expression does not change in response to queuine (this work). (B) Quantitative analysis of relative density of mRNA. The signal intensities of the samples were divided by enolase mRNA and were then normalized to the wild type mRNA. Data are represented as mean± standard deviation about the mean for two independent experiments.

Figure 5: Characterization of the EhTGT silenced trophozoites (A) EhTGT was detected in control trophozoites and siEhQTRT1 trophozoites grown with and without queuine using a homemade polyclonal mice EhTGT antibody (1:250) and then incubated with a goat anti mouse secondary antibody Alexa fluor (Jackson Immuno Research) at a concentration of 1:250. Nucleus are stained with 4`,6-diamidino-2-phenylingole (DAPI) (Sigma-Aldrich) at concentration of 1:1000. The samples were examined under a confocal immunofluorescence microscopy (Zeiss-LSM700 Meta laser scanning system confocal imaging system) with a 63X oil immersion objective. Images are scaled to 10μm.
(B). Densitometry analysis showing the percentage of the EhTGT complex in 20 trophozoites in each condition (Normalized to control trophozoites). Data representative of three independent experiments (Unpaired Student's t test, p ≤ 0.0001).
(C) APB Northern Blot analysis of *E.histolytica* trophozoites. (A). Lane 1- Control Trophozoites (T), Lane 2- Control trophozoites grown with queuine (T+q), Lane 3- siEhQTRT1 trophozoites (siEhQTRT1), and Lane 4- siEhQTRT1 trophozoites grown with queuine (siEhQTRT1+q). Total RNA samples was run on a 15% acrylamide gel supplemented with APB, transferred on a nylon membrane, dried and incubated with a probe against tRNA[His].
(D) Polyacrylamide gel stained with ethidium bromide as loading control.

Figure 6: Effect of silencing EhQTRT1 on the growth of *E.histolytica*
Control trophozoites (T) and Trophozoites silenced for EhQTRT1 (siEhQTRT1) ($30X10^3$ /ml of complete TYI media) were grown for 24h, 42, and 72 hours with queuine (+q) or without queuine. The number of viable trophozoites at each time point was determined by using the vital stain eosin. siEhQTRT1 trophozoites have their growth impaired compared to that of control trophozoites. Queuine

has no significant effect on the growth of T or SiEhQTRT1 trophozoites. The data are representative of three independent experiments performed in duplicates. Data are expressed as mean ± SD. (Unpaired Student's t test, T vs siEhQTRT1 24h (p= 0.0006), 48h (p= 0.0082), 72h (p= 0.005).

Figure 7: Queuine triggers resistance to OS and this effect is lost in siEhQTRT1 trophozoites.
Determination of the amount of H_2O_2 required to kill 50% of the population (LD50) of control trophozoites, trophozoites grown with queuine (0.1µM for 3 days), siEhQTRT1 trophozoites and siEhQTRT1 trophozoites cultivated with queuine. The data represent mean ± SD of three independent experiments performed in duplicates. The LD_{50} of trophozoites grown with queuine and exposed to OS (T+q+OS) was significantly higher than the LD_{50} of untreated trophozoites exposed to OS (T) (Unpaired Student's t test, p≤0.005). The LD_{50} of siEhQTRT1 trophozoites grown with queuine and exposed to OS (siEhQTRT1+q+OS) was significantly lower than the LD_{50} of siEhQTRT1 trophozoites exposed to OS (siEhQTRT1+OS) (Unpaired Student's t test, p≤0.005)

Figure 8: Measurement of protein synthesis in trophozoites grown with queuine using puromycin-labeled proteins.
(A) Ponceau stained membrane (loading control). (B) Lane 1- control trophozoites, Lane 2- trophozoites grown with queuine (0.1µM for three days), Lane 3- trophozoites grown with queuine labeled with 10ug/ml puromycin, Lane 4- control trophozoites exposed to 2.5mM H_2O_2 for 20 minutes and then labeled with puromycin for 20 mins. Lane 5- trophozoites grown with queuine prior to their exposure to 2.5mM H_2O_2 and then labeled with puromycin. Lane 6- control trophozoites labeled with 10µg/ml puromycin. Lane 7- trophozoites grown with queuine were treated with cyclohexamide (CHX;100ug/ml) before labeling with puromycin (Control+P+CHX). Total protein extracts were separated on a SDS-PAGE gel and analyzed by Western blotting with a mouse-monoclonal puromycin antibody (1:5000). These results are representative of three independent experiments.

Figure 9: Measurement of protein synthesis in trophozoites silenced for EhQTRT1 grown with queuine using puromycin-labeled proteins.
 A) Ponceau stained membrane (loading control). (B) Lane 1- siEhQTRT1 trophozoites, Lane 2-

siEhQTRT1 trophozoites grown with queuine (0.1µM for three days), Lane 3- siEhQTRT trophozoites labeled with 10ug/ml puromycin, Lane 4- siEhQTRT1 trophozoites grown with queuine and then labeled with puromycin for 20 mins. Lane 5- siEhQTRT1 trophozoites exposed to 2.5mM H_2O_2 and then labeled with puromycin, Lane 6- siEhQTRT1 trophozoites grown with queuine prior to their exposure to 2.5mM H_2O_2 followed by their labeling with puromycin. Lane 7- siEhQTRT1 trophozoites were treated with cyclohexamide (CHX;100ug/ml) before labeling with puromycin (Control+P+CHX). Total protein extracts were separated on a SDS-PAGE gel and analyzed by Western blotting with a mouse-monoclonal puromycin antibody (1:5000). These results are representative of three independent experiments.

Figure 10: Effect of queuine on the virulence of *E.histolytica*

(A). Cytopathic activity of *E. histolytica* trophozoites and siEhQTRT1 trophozoites grown with and without queuine was measured by their ability to destroy a monolayer of HeLa cells. *E. histolytica* trophozoites were grown with and without queuine (0.1µM for 3 days). Data are displayed as the mean ± standard deviation of three independent experiments that were repeated twice. (Unpaired Student's t test, T vs T+q, p= 0.0026). siEhQTRT1 trophozoites grown with and without queuine were incubated with HeLa cell monolayer for 40 or 90 minutes. The cytopathic activity of siEhQTRT1 trophozoites grown with queuine is not different from the cytopathic activity of siEhQTRT1 trophozoites grown without queuine at both 40 and 90 minutes of incubation between the trophozoites and the HeLa cell monolayer. The data are representative of three independent experiments (normalized to wild type) (Unpaired Student's t test, p= 0.01 (T vs T+q)).

 (B) Effects of queuine treatment in intestinal amoebiasis. Number of trophozoites determined in CBA/J mice stool infected with trophozoites grown with queuine (q) or non-treated trophozoites (T). Trophozoites number was quantified by the quantitative real-time PCR method using *E. hisolytica* 18S rRNA specific primers. Data represent the mean ± SD for 10 mice. p < 0.05 by an Unpaired Student's t-test.

Figure 11: Experimental design of the RNA-seq.

(A)Trophozoites have been grown with and without 0.1µM queuine for 3 days (T and q), and then were exposed (or not) to OS (+OS and –OS). The experiments were repeated three times to obtain 12 samples that were purified and sequenced further. (B). Biological conditions compared during differential

RNA-seq analysis. 5 pair-wise comparisons have been performed.

Figure12: Principal component analysis and average profiles

(A) Principal Component Analysis of the RNA-Seq data performed on the variance-stabilizing transformed count matrix.

(B) Average profiles for the four gene clusters defined by the hierarchical clustering. The average transcriptome profile was established by computing the mean $\log_2$-normalized count for the genes in each of the four identified clusters.

Figure 13: Functional categories and classification of genes enriched in trophozoites grown with queuine vs control trophozoites.

Functional categories: **(a)** up regulated genes in trophozoites grown with queuine. **(b)** downregulated genes in trophozoites grown with queuine. The number of genes related to each category was derived by processing the data (Supplementary Table S1) with PANTHER (http://pantherdb.org). **(c)** Enrichment test in queuine treated trophozoites.

Figure 14: Functional classification of genes

(A) Categories for functional classification of genes downregulated in the presence of queuine (using a subset of protein class). A complete list of the genes downregulated in the presence of queuine are given in Supplemental Table S2. (B) Categories for functional classification of genes commonly upregulated in virulent HM1:IMSS strain and genes downregulated in the presence of queuine. The names of the genes in each category are given in Supplementary Table S4.

Figure 1

(A)

(B)

Figure 2

Figure 3

(A)

(B)

Figure 4

(A)

(B)

Figure 5

(A)

(B)

(C)

(D)

Figure 6

Figure 7

Figure 8

(A)
(B)

Figure 9

(A)   (B))

Figure 10

40 minutes

90 minutes

(A)

(B)

Figure 11

(A)

(B)

Figure 12

(A)   **(B)**

Figure 13

Functional categories of differentially regulated genes in Q vs T (C1)

(A) Upregulated genes

(B) Downregulated genes

Figure 13

Enrichment (PANTHER GO-Slim Molecular Function) in Q vs T (C1)

Figure 14

(A)

(B)