Characterizing the body morphology of the first metacarpal in the Homininae

using 3D geometric morphometrics

Jonathan Morley[1], Ana Bucchi[2,3], Carlos Lorenzo[4,3], Thomas A. Püschel[5,1*]

[1]Institute of Cognitive and Evolutionary Anthropology, School of Anthropology,

University of Oxford, 64 Banbury Road, OX2 6PN, Oxford, United Kingdom.

[2]Centro de Investigación en Odontología Legal y Forense (CIO), Universidad de La

Frontera, Uruguay 1836, Temuco, Chile.

[3]Institut Català de Paleoecologia Humana i Evolució Social (IPHES), Zona

Educacional 4, Campus Sescelades URV (Edifici W3), 43007 Tarragona, Spain.

[4] Àrea de Prehistòria, Universitat Rovira i Virgili (URV), Avinguda de Catalunya 35,

43002 Tarragona, Spain.

[5] Ecology and Evolutionary Biology Division, School of Biological Sciences, University

of Reading, Philip Lyle building, RG6 6BX, Reading, United Kingdom.

*Corresponding author:

Thomas A. Püschel

E-mail: thomas.puschelrouliez@anthro.ox.ac.uk

Phone: -

Running title: Characterizing MC1's morphology in the Homininae

Abstract

Objectives: The morphological characteristics of the thumb are of particular interest

due to its fundamental role in enhanced manipulation. Despite its possible importance

regarding this issue, the body of the first metacarcapal (MC1) has not been fully

characterized using morphometrics. This could provide further insights into its

anatomy, as well as its relationship with manipulative capabilities. Hence, this study

quantifies the shape of the MC1's body in the extant Homininae and some fossil

hominins to provide a better characterization of its morphology.

Materials and methods: The sample includes MC1s of modern humans (n=42), gorillas

(n=27) and chimpanzees (n=30), as well as *Homo neanderthalensis*, *Homo naledi* and

Australopithecus sediba. 3D geometric morphometrics were used to quantify the

shape of MC1's body.

Results: The results show a clear distinction among the three extant genera. *H.*

neanderthalensis mostly falls within the modern human range of variation. *H. naledi*

varies slightly from modern humans, although also showing some unique trait

combination, whereas *A. sediba* varies to an even greater extent. When classified

using a discriminant analysis, the three fossils are categorized within the *Homo* group.

Conclusion: The modern human MC1 is characterized by a distinct suite of traits, not

present to the same extent in the great apes, that are consistent with an ability to use

forceful precision grip. This morphology was also found to align very closely with that

of *H. neanderthalensis*. *H. naledi* shows a number of human-like adaptations, whilst

A. sediba presents a mix of both derived and more primitive traits.

Keywords: MC1; 3D geometric morphometrics; Semi-landmarks; Hominins; African

apes

1 Introduction

Thumbs in modern humans are different from those of African apes (e.g., Almécija, Smaers, & Jungers, 2015; Dunmore, Bardo, Skinner, & Kivell, 2020; Galletta, Stephens, Bardo, Kivell, & Marchi, 2019; Green & Gordon, 2008; Stephens et al., 2016; Susman, 1994). Modern humans have a relatively broader shaft for the first metacarpal (MC1) and a higher thumb-to-digit ratio than the African apes, especially chimpanzees (Almécija et al., 2015; Feix, Kivell, Pouydebat, & Dollar, 2015; Green & Gordon, 2008; Rolian & Gordon, 2013). Additionally, compared to our closest living relatives we possess thenar musculature that is relatively more developed than the other hand muscles (Tuttle, 1969), a condition that has been inferred from the hominin fossil record by observing how strong or flawed their bony attachments are (Bush, Lovejoy, Johanson, & Coppens, 1982; Karakostis, Hotz, Tourloukis, & Harvati, 2018; Kivell, 2015; Kivell, Kibii, Churchill, Schmid, & Berger, 2011; Maki & Trinkaus, 2011; Richmond et al., 2020). Modern humans also differ from the extant African apes in the relative size of the epicondyles and degree of curvature of the proximal (Marchi, Proctor, Huston, Nicholas, & Fischer, 2017; Marzke et al., 2010) and distal joint surfaces of the MC1 (Galletta et al., 2019).

These anatomical traits that set apart humans from the African apes have presumably evolved to cope with the different functional demands experienced by these taxa (i.e., manipulation vs. locomotion) (Almécija, Moyà-Solà, & Alba, 2010; Matarazzo, 2015; Püschel, Marcé-Nogué, Chamberlain, Yoxall, & Sellers, 2020; Richmond & Strait, 2000; Tsegai et al., 2013). The more robust human thumb and greater degree of curvature of the joint surfaces allow our species to produce greater force and to better withstand the stresses of tool-related behaviors (Galletta et al., 2019; Key & Dunmore, 2018; Key & Dunmore, 2015; Rolian, Lieberman, & Zermeno, 2011). On the other

hand, the thumb of chimpanzees is slender and shorter relative to the other fingers,

presumably a suspensory-related adaptation (Almécija et al., 2015; Feix et al., 2015),

and although the thumb length and breadth in gorillas differs less from humans than

chimpanzees (Almécija et al., 2015; Feix et al., 2015; Green & Gordon, 2008), it does

show a reduced thenar musculature, which is the primitive condition in the hominidae

(Diogo, Richmond, & Wood, 2012; Tocheri, Orr, Jacofsky, & Marzke, 2008; Tuttle,

1969).

Even though using skeletal proxies of the MC1 to infer the degree of dexterity is

common practice (e.g., Dunmore et al., 2020; Feix et al., 2015; Maki & Trinkaus,

2011), the continuous nature of these traits makes it difficult to quantify how different

hominines are with respect to each other, which consequently complicates the

correlation of these proxies with different functional capabilities. Building upon this

problem, recent research on the MC1 has been conducted using three-dimensional

geometric morphometric (3DGM) techniques, focusing on the joint surfaces in apes

and fossil hominins (Galletta et al., 2019; Marchi et al., 2017). Marchi et al. (2017)

propose that hominins (modern humans, *Paranthropus robustus*/early *Homo* SK84

and *Au. africanus*) are significantly different from non-human hominids in that they

possess a radioulnar and dorsovolar flatter proximal joint, a less projecting volar beak

and a radially extended surface. This would allow our species to better abduct and to

accommodate larger axial loads when pinching objects (Marchi et al., 2017; Marzke

et al., 2010). Humans also vary from apes in having a larger and flatter distal articular

surface in a radioulnar direction and a radial palmar condyle that is larger and more

palmarly projecting than the ulnar one, which would contribute to the stabilization of

the joint during forceful precision grip (Galletta et al., 2019). Neanderthals and *H.*

naledi are located within the modern human range of variation for these traits, whereas

the other analyzed hominins (*Au. africanus, Paranthropus robustus*/early *Homo* SK84

and *Au. sediba*) occupy a position between modern humans and the great apes.

In spite of its possible importance, the body of the MC1 has not been fully analyzed

using 3DGM to assess its possible relevance when correlating its anatomy with

different manipulative capabilities. In addition, fossils are often fragmentary and

epiphyses in the fossil record are often damaged (see for e.g., *H. naledi*'s U.W. 101-

401 MC1) or abraded (see for e.g., *H. naledi*'s U.W. 101-1641 MC1). Therefore, a

method focused only on the MC1 shaft might be particularly useful. Consequently, this

study focuses on the body morphology of the MC1 using 3DGM. The objective was to

provide further information that could contribute towards a better characterization of

the MC1's anatomy, as well as to provide further insights towards the identification of

structures in extant species that may be associated with human-like manipulative

capabilities and to assess if similar morphologies are present in fossil hominins.

Even though the great apes use their hand for manipulatory activities, their

morphology is likely more related to their locomotion (i.e., knuckle-walking and

arborealism) (Almécija, Moyà-Solà, & Alba, 2010; Matarazzo, 2015; Püschel et al.,

2020; Richmond & Strait, 2000; Tsegai et al., 2013). It is therefore expected that the

selective pressures associated with locomotor behavior in chimpanzees and gorillas

will result in an MC1 morphology that varies significantly from that of modern humans.

We also expect gorillas to be closer to humans rather than chimpanzees, as previous

research has indicated that their metacarpals are broader and the thumb-fingers ratio

less different from humans compared with those of chimpanzees (Almécija et al.,

2015; Green & Gordon, 2008; Rolian & Gordon, 2013). Additionally, we also expect

the MC1's shaft morphology of *Au. sediba* to show an intermediate morphology

located between the range of variation of modern humans and that of the great apes,

as previous studies indicate that the hand of this species displays a mosaic anatomy

of primitive and derived traits (Kivell et al., 2011). *Au. sediba* MC1 has gracile

attachments for the opponens pollicis and first dorsal interosseus muscles, but it also

possesses a long thumb relative to the fingers, which is close to the modern human

configuration (Kivell et al., 2011). On the other hand, we expect that the Neanderthal

and *H. naledi* specimens will show a morphology similar to that of modern humans as

previous analyses have suggested that they exhibit similar attachment sites of the

thenar musculature, as well as a relatively similar thumb length (Feix et al., 2015;

Karakostis, Hotz, Tourloukis, & Harvati, 2018; Kivell, 2015; Maki & Trinkaus, 2011).

Consequently, we tested the following hypothesis:

Hypothesis 1: MC1's morphology significantly differs between modern humans and

the extant African ape species. The modern human MC1's shaft is expected to be

more similar to that of gorillas rather than chimpanzees due to its broader shaft, as

well as relative length and breadth.

Hypothesis 2: *H. naledi* and *H. neanderthalensis* specimens exhibit an MC1

morphology more similar to modern humans than other great apes, while *Au. sediba*

shows an intermediate morphology between the African apes and modern humans.

2 Material and methods

2.1 Sample

The extant sample used in this study includes MC1s of modern humans (*Homo*

sapiens; n=42), chimpanzees (*Pan troglodytes*; n=30), and gorillas (*Gorilla gorilla* and

Gorilla beringei; n=27) (Table S1). The human MC1s came from a medieval cemetery

in Burgos, Spain (Casillas Garcia & Alvarez, 2005) and the surface models were

obtained using a Breuckmann SmartSCAN structured light scanner. The non-human

sample came from museum collections and are of different origins (i.e., wild-shot,

captivity and unknown origin). There were no significant shape differences between

wild vs. captive specimens, nor between the two gorilla species included in this study,

hence we felt confident to pool the extant specimens at the genus level (see S2 for

further details). The MC1 surface models were collected using photogrammetry as

described in Bucchi et al. (2020a). The 3D models from the surface scanner were

obtained using a resolution of 0.125 mm, while most of the photogrammetric models

ranged from 400,000 to 600,000 triangles of uniform size. A previous study that

applied the same surface scan and photogrammetry protocols to digitize hand bones

found that both types of 3D models are of comparable quality (Bucchi, Luengo, Bove,

& Lorenzo, 2020b). Additionally, we carried out a comparison a sub-sample of 30

specimens that were digitized using both technologies (i.e., photogrammetry and

structured-light scanning) and we found that differences between models obtained

using the different digitalization technologies are extremely small (less than ~0.17 mm

on average). Hence, we are confident that it is possible to combine these kinds of 3D

models in our analyses. Further details about these 3D model comparisons can be

found in S3. Only adult individuals that show no evident pathologies were included in

the study and right MC1s were preferred (although left MC1s were reflected when their

antimere was not present as indicated in S1).

The fossil sample includes the right metacarpal from a *Homo neanderthalensis*, the

right metacarpal from a *Homo naledi* and the left metacarpal from an *Australopithecus*

sediba. The *H. neanderthalensis* sample (La Ferrassie 1) was found in La Ferrassie

archaeological site in Savignac-de-Miremont, France. This skeleton was discovered

in 1909 and is estimated to be 70–50,000 years old (Guérin et al., 2015). The *Homo*

naledi sample (U.W. 101-1321) was recovered in 2013 from the Rising Star cave

system in South Africa and has been dated to around 236-335 ka years ago (Dirks et

al., 2017). The *A. sediba* sample (MH2) was taken from the near complete wrist and

hand of an adult female (Malapa Hominin 2) discovered in Malapa, South Africa, which

has been dated around 1.98 million years (Berger et al., 2010; Pickering et al., 2011).

The latter fossils were downloaded from Morphosource

https://www.morphosource.org/, whereas the Neanderthal was obtained from a cast

housed at the Catalan Institute of Human Paleoecology and Social Evolution (IPHES).

2.2. 3DGM

3D landmarks were collected using the software Landmark Editor 3.6 (Wiley et al.,

2005) to quantify the MC1's morphology, including relevant functional proxies as the

epicondyles, the shaft curvature and the attachments sites for the opponens pollicis,

abductor pollicis longus and first dorsal interosseus muscles. These attachments sites

are in the MC1 at the lateral margin, body at the ulnar side of the bone and the base

at the radial side, respectively, and are the same for the three genera under study

(Diogo et al., 2011; 2013). Eight curves comprising 20 equidistant landmarks each

were placed at pre-defined points on the MC1 (Figure 1; S4). These landmarks were

chosen to provide a good representation of the shaft of the bone.

We assessed whether sufficient number of landmarks have been sampled to

characterize MC1's shape variation by using the lasec() function of the 'LaMBDA'

0.1.0.9000 R package (Watanabe 2017) (further details about this procedure can be

found in S5). The first and last landmarks from each one of the eight curves were

treated as fixed (i.e., 16 fixed landmarks), whereas all the rest of them (i.e., 144

landmarks) were considered as semi-landmarks. A generalized Procrustes

superimposition was performed on the landmark data to remove differences due to

scale, translation, and rotation in order to obtain shape variables (Bookstein, 1991).

This procedure was done using the gpagen() function available as part of the

'geomorph' R package 3.3.1 (Adams, Collyer, & Kaliontzopoulou, 2020). The semi-

landmarks were slid on the MC1's surface by minimizing bending energy (Bookstein,

1997; Gunz, Mitteroecker, & Bookstein, 2005). This is an iterative process that works

by allowing the semi-landmarks to slide along the surface to remove the effects of

arbitrary spacing by optimizing the location of the semi-landmarks with respect to the

consensus shape configuration (Gunz & Mitteroecker, 2013). There are two main

criteria to slide semi-landmarks (i.e., bending energy and Procrustes distance) which

have been shown to provide relatively similar results when carrying out inter-specific

comparisons (Perez, Bernal, & Gonzalez, 2006). We preferred to use bending energy

as this sliding criterion allows all semi-landmarks to slide together and being influenced

by the other available landmarks and semi-landmarks, whereas when Procrustes

distance is used, each semi-landmark slides individually and, apart from the common

Procrustes superimposition (Gunz & Mitteroecker, 2013). All the Procrustes residuals

analyzed in this work are available in S6.

These obtained shape variables were then used in a principal component analysis

(PCA) to summarize shape variation. The PCA was carried out using the gm.prcomp()

function of the 'geomorph' R package 3.3.1 (Adams, Collyer, & Kaliontzopoulou,

2020). To visualize shape differences warped models representing the shape changes

along the first three principal components (PCs) were generated. The models closest

to the mean shape (i.e., lowest Procrustes distance to the multivariate consensus) was

warped to match the multivariate mean using the thin plate spline method. Then, the

obtained average model was warped to display the variation along the three plotted

PC axes (mag = 1).

The dataset of extant hominoids was then grouped by genus and the Procrustes

variance of observations in each group (i.e., the mean squared Procrustes distance of

each specimen from the mean shape of the respective group) was computed as a

simple measure to assess morphological disparity within each one (Drake &

Klingenberg, 2010; Zelditch, Swiderski, & Sheets, 2012a). Procrustes variance was

applied here as way to evaluate intra-genus variation, and absolute differences in

Procrustes variances were computed to test differences in morphological disparity

among groups (these differences statistically evaluated through permutation [999

rounds]). This procedure was carried out using the morphol.disparity() function

available as part of the 'geomorph' R package 3.3.1 (Adams, Collyer, &

Kaliontzopoulou, 2020).

A multi-group linear discriminant analysis (LDA) (also known as canonical variate

analysis [CVA]) was run to test if it was possible to distinguish among the three genera.

This procedure maximizes the separation between groups. Since our number of

variables (i.e., landmarks and semi-landmarks) exceeded the number of analyzed

specimens, we carried out this analysis using the principal components (PCs) that

accounted for 90% of the sample variance to reduce the dimensionality of the dataset.

The LDA was carried out using the lda() function of the 'MASS' 7.3-51.6 R package

(Venables & Ripley, 2002). Performance was calculated using the confusion matrix

from which the overall classification accuracy was computed, as well as the Cohen's

Kappa statistic (Püschel, Marcé-Nogué, Gladman, et al., 2020; Püschel, Marcé-

Nogué, Gladman, Bobe, & Sellers, 2018). The complete dataset was resampled using

a 'leave-one-subject-out' cross-validation, as a way to asses classification

performance (Kuhn & Johnson, 2013). In addition, by using the obtained discriminant

function we classified the fossil sample into the three extant genera as way to assess

morphological affinities. Pairwise PERMANOVA tests with Bonferroni corrections for

multiple comparisons were performed to assess shape differences among the three

extant genera using the PCs that accounted for 90% of the sample variance. Euclidean

distances computed using the PCs that accounted for ~ 90% of the total variance of

the sample were selected as dissimilarity index. This procedure was performed using

the adonis() function of the 'vegan' 2.5-7 R package (Oksanen et al., 2020).

Additionally, we also decided to compute a curvature metric to better assess how

curve the MC1's shaft is along both its dorsal and palmar aspects (i.e., semi-landmark

curves C1 and C2 in Fig. 1), as well as to facilitate the morphological description of

the morphometric results. Hence, Menger (1930) curvatures were calculated for each

one of the semi-landmark points of the two curves (i.e., C1 for the dorsal side and C2

for the palmar aspect) using a custom-written script in R. The Menger curvature of

three points in n-dimensional Euclidean space $\mathbb{R}^n$ is defined as the reciprocal of the

radius of the circle that passes through the three points (Menger, 1930). Menger

curvature was calculated locally for each semi-landmark point along the curve,

excepting the first and last fixed landmarks, as it is not possible to compute a curvature

value at the starting and ending points of each one of the curves. This resulted in 18

curvature values for each one of the semi-landmark curves (i.e., C1 and C2). The

curvature values of each one of the curves were summed to obtain a measurement of

the overall curvature of C1 and C2 (higher values would correspond to more

pronounced curvatures). This procedure was performed on the six 3D models that

were warped to represent the variation along the first three PC axes.

In addition, as measurement error (ME) has a critical importance when performing

morphometric analyses, a sub-sample of 33 randomly selected MC1s were digitized

twice and compared via a Procrustes ANOVA to assess ME (Klingenberg & McIntyre,

1998). We also carried out a regression of shape variables on centroid size using the

whole sample to assess allometric influence. Both procedures were carried out using

the procD.lm() function available as part of the 'geomorph' R package 3.3.1 (Adams,

Collyer, & Kaliontzopoulou, 2020). All the mentioned morphometric and statistical

analyses were carried out in R 4.0.2 (R Core Team, 2020).

3 Results

3.1 Measurement error and allometric influence

The Procustes ANOVA used to measure intra-observer error in the sub-sample

showed that the mean square for individual variation far exceeded ME, so this type of

error was negligible (see S7 for further details). ME was also quantified as shape

repeatability using a ratio of the among-individual to the sum of the among-individual

and measurement error components as explained in Zelditch, Swiderski, & Sheets

(2012b). Shape repeatability was 0.95, which indicates a minimal ~5% error.

Regarding allometric influence, we found that centroid size only accounted for ~2.7%

of MC1 shape variation. This means that, for the goals of the present study, we can

exclude size as a particularly significant factor contributing to potential inter-generic

variation in shape. Hence, we decided that it was not necessary to 'correct' for

allometric effects as ~97.3% of the shape variation is not explained by size (further

details about this regression are available in S8).

3.2 Principal component analysis

The PCA performed using the shape variables returned 102 PCs. The first 22 PCs

accounted for ~ 90% of the total variance of the sample, hence offering a reasonable

estimate of the total amount of MC1's shape variation, which were then used in the

LDA and pairwise PERMANOVA tests. The first three PCs in the PCA account for ~

58% of the total variance and display a relatively clear separation among the extant

African ape genera (Fig. 2a) (PCA biplots for PC1 vs. PC2, PC1 vs. PC3 and PC2 vs.

PC3 are also available in S9). PC1 explains 41.44%, PC2 11.18% and PC3 5.82% of

total variance, respectively (Fig. 2). To visualize shape differences, the warped 3D

models corresponding to the highest and lowest values at each extreme of the first

three PCs were plotted alongside the violin plots. A violin plot is a combination of a

boxplot and a kernel density plot that is rotated and placed on each side to show the

distribution shape of the data (Adler & Kelly, 2020). In addition, six movies showing

the shape changes along the three first PCs axes are also provided in S10. These

warped models are also displayed in Figure 3 to facilitate the morphological

interpretation of our results. Anatomical descriptions associated with each one of the

positive and negative extremes of the first three PCs are also provided in the same

figure.

Violin plots of PC1 (Fig. 2b) show a notable difference between gorillas and humans

vs. chimpanzees. PC1 separates the mediolaterally narrower MC1 shafts of *Pan* from

the broad MC1 shafts of *H. sapiens* and *Gorilla.* These two genera exhibit the highest

PC1 scores, which correspond to a more developed muscular attachments, straighter

dorsal aspect of the body, and overall robust shaft with broader distal-most and

proximal-most portions of the shaft (i.e., MC1's body and not the articular surfaces

which were not morphometrically characterized in this study) (Fig. 3). Chimpanzees

show the lowest PC1 scores, representing a more gracile shaft, a more pronounced

antero-posterior curvature of the shaft, less marked muscle attachments with narrower

distal-most and proximal-most portions of the body and smaller radial and ulnar

epicondyles (Fig. 3). *H. neanderthalensis* falls within the human and gorilla

distributions and is distinct completely from the chimpanzees. *H naledi* falls within the

gorilla distribution, whilst *A. sediba* is characterized by a lower PC1 score and aligns

closer to the *Pan* distribution. None of the analyzed fossils fall within any of the

interquartile ranges (IQR) (i.e., black bars in Fig. 2b-d) of any of the extant genera.

Violin plots of PC2 (Fig. 2c) shows distinct variation among the extant genera, with a

morphological continuum ranging from *Gorilla* (lower PC2 values), *Pan* (central PC2

values) and extant *Homo* (higher PC2 scores). PC2 seems to summarize the relative

breadth of the middle and distal shaft with respect to the relative size of the proximal

shaft and base. The *Gorilla* sample has the lowest PC2 scores, a more radioulnar and

dorsovolar rounded ends of the shaft, and a medial epicondyle which is more distal

relative to the lateral epicondyle (Fig.3). The modern human distribution shows the

highest PC2 scores, representing flatter distal-most and proximal-most portions of the

body, as well as larger area at the place of origin of the radial palmar condyles at the

distal end of the analyzed area (i.e., the shaft). The chimpanzee sample lies in

between the gorilla and modern human samples displaying an intermediate

morphology. In a similar fashion as chimpanzees, the three fossils are located at

intermediate positions in PC2 distribution. *H. neanderthalensis* and *H. naledi* display

PC2 scores that are within the *Pan* IQR, whilst *A. sediba* has higher values.

Violin plots of PC3 (Fig. 2d) show a similar distribution of PC scores for the three extant

genera. From a morphological perspective, lower values correspond to more gracile

MC1s while higher scores are associated with more robust morphologies displaying

more surface for muscular attachments (for the opponens pollicis, first dorsal

interosseous and abductor pollicis longus muscles). *H. naledi* and *A. sediba* show

values which are within the extant genera distributions, but outside their IQR and at

opposite extremes of the axis. *H. neanderthalensis* lies outside the distribution of any

of the extant genera, probably due to its particularly robust morphology and associated

marked muscular insertion areas, in particular a marked lateral flange for the

opponens pollicis.

3.3 Morphological disparity

The obtained results show that three extant genera show a similar magnitude of

disparity. Nevertheless, gorillas exhibit a higher Procrustes variance as compared to

modern humans and chimpanzees (Table 1a). Gorillas are significantly different to

modern humans, and chimpanzees when comparing absolute variance differences,

whilst modern human do not significantly differ from chimpanzees (Table 1b).

3.4 Linear discriminant analysis

The LDA model using the first 22 PCs clearly distinguishes among the three extant

genera, displaying an outstanding performance with excellent classification results

after cross-validation (Accuracy: 0.97; Cohen's Kappa: 0.95; Fig. 4; Table 2). When

using the obtained discriminant function to classify the fossils into the extant categories

(as a way of assessing morphological affinities) (Table 3), the three of them were

classified into the *Homo* category, even though only *H. neanderthalensis* was located

within the 95% confidence interval of the modern humans (Fig. 4). The posterior

probabilities were extremely close to 1 for *H. naledi and H. neanderthalensis*, hence

indicating that, in spite of their differences, their morphology is closer to that of modern

humans. *A. sediba* was also classified within the *Homo* category (posterior probability:

63%) but this specimen also showed non-trivial posterior probabilities classifying it

within the *Gorilla* category (posterior probability: 29%) or as a member of the *Pan*

group (posterior probability: 7%). There were significant differences among all extant

genera when analyzing the 22 PCs from the PCA carried out using the shape variables

(Table 4).

3.5 Curvature

Table 5 provides the summed Menger curvature values for the six 3D models that

were warped to represent the variation along the first three PC axes. These values

were computed for both the dorsal (C1) and palmar (C2) sides of the MC1's shaft and

represent overall curvature. As expected, the palmar side of the shaft (C2) is more

curved than its dorsal counterpart (C1). This means that the palmar curvature (C2)

values are always higher as compared to the dorsal ones (C1) for all analyzed PCs.

Overall, the shapes associated with the maximum values for each one of the three

PCs corresponded to straighter shafts along the palmar side. In addition, the palmar

curvature value (C2) for the minimum scores along PC1 correspond to the highest

curvature (i.e., C2 curvature value for PC1 min; Table 5). The region of the

morphospace that corresponds to this shape is occupied by the chimpanzees (Fig.

2b). In summary, the shapes associated with modern humans and gorillas are straighter, whereas the shapes that describe chimpanzees exhibit a more pronounced curvature along the palmar side.

4 Discussion

The first hypothesis was that the shape of the human MC1 would differ significantly from that of *Pan* and *Gorilla*, due to the variation in their manipulative capabilities and locomotive behaviors, and that *Gorilla* would show more morphological affinity with humans than chimpanzees. Overall, these analyses provide support for this hypothesis, confirming that there is indeed significant morphological variation between the extant great apes. We also found clear differences between chimpanzees and gorillas, with gorillas closer (i.e., more similar) to humans than to chimpanzees in PC1. This is due to their broader and more robust MC1s of gorillas and humans (i.e., broader shafts and expanded ulnar and radial epicondyles), as compared to the slender and more curved MC1s of chimpanzees (Table 5). The second hypothesis was that fossil hominin species *H. naledi* and *H. neanderthalesis* would exhibit an MC1 morphology more similar to humans than other great apes and *Au. sediba* an intermediate morphology between the African apes and humans. The results also support this hypothesis, as observed in the PCA and LDA plots (Figs. 2a and 3). However, it is important to note that even though the three fossils are more similar to the modern humans, they also display some distinct features, different from those which would typically be expected in extant *Homo*.

4.1 Hominin MC1 shape

The 3DGM data indicate that there is a distinctive suite of morphological traits that

distinguish humans from chimpanzees and gorillas (Fig 3.). The main distinguishing

traits are a straighter and more robust shaft (Fig. 2b and 3; Table 5) accompanied by

larger radial and ulnar epicondyles and flatter distal and proximal ends of the body

(Fig. 2c and 3). It is important to note that African apes are not a homogeneous group.

For instance, gorillas show morphological affinities with humans relative to the shaft

robusticity, although with a more proximo-distally curved shaft along its palmar side

(Table 5), and more rounded distal-most and proximal-most portions of the MC1's

body. The PCA shows that chimpanzees are characterized by a slender and more

gracile MC1, and this trait makes chimpanzees the most distinctive genus amongst

extant taxa (Fig. 2b and 3). Chimpanzees are also characterized by an intermediate

curvature of the radioulnar and dorsovolar ends of the shaft (Fig. 2c and 3).

Interestingly, we found that chimpanzees display a more proximo-distally curved MC1

shaft compared with gorillas and humans (Fig. 3; Table 5). To our knowledge, this

property has been only studied in phalanges 2-5, with those of chimpanzees being

more curved than those of gorillas and humans (Stern, Jungers, & Susman, 1995;

Susman, 1979). This curvature degree at the shaft has been usually interpreted as an

adaptation for suspension and overall arboreal locomotion in digits 2-5 (Rein, 2011).

As for the thumb, there is preliminary data that seems to indicate that it is routinely

recruited during suspension in orangutans (MCclure Phillips, Vogel & Tocheri, 2012),

yet its role has not been fully studied in chimpanzees. Consequently, it is not possible

associate the MC1's curvature observed in chimpanzees with the suspensory

behaviors of this species.

As for the fossils studied here, results indicate that they all show a unique repertoire

of morphological traits, different from those of extant genera. The general scientific

consensus in recent years is that *H. neanderthalensis* had a hand morphology that

was very similar to that of humans (Karakostis, Hotz, Scherf, Wahl, & Harvati, 2017;

Karakostis et al., 2018; Niewoehner, 2001, 2006; Tocheri et al., 2008; Trinkaus &

Villemeur, 1991). Our obtained results align well with this consensus, with the *H.*

neanderthalensis specimen showing several similarities with the modern humans. The

described morphology is one of a flatter (PC2) and broader (PC1) distal-most portion

of the shaft, bigger epicondyles at the distal head (PC1) and a flatter proximal-most

area of the shaft (PC2). However, *H. neanderthalensis* also differs in exhibiting a

particularly robust MC1 with strongly marked muscular insertions, which distinguishes

it from the rest of the sample, particularly along PC3. Neanderthals are known for the

large flanges on their MC1s for the insertion of the opponens pollicis muscle that

results in a waisted appearance of the MC1 in an anterior or posterior view (Maki &

Trinkaus 2011; Trinkaus 1983). This trait also appears, to varying degrees, among

some modern human populations, but rarely to the extent observed in *H.*

neanderthalensis (Trinkaus et al. 2014). In our sample, this trait clearly distinguishes

H. *neanderthalensis* from the rest of the analyzed specimens along PC3.

Previous reports indicate that the general morphology of the *H. naledi* MC1 aligns

more closely with humans than apes, whilst still possessing a number of more primitive

characteristics than the human MC1 (Kivell et al., 2015; Galletta et al. 2019). In our

results we found that *H. naledi* aligns closer to humans in terms of shaft robusticity

(Fig. 2b) and well-developed crest for the insertion of the opponens pollicis muscle

(PC3) as it was reported by Kivell et al., (2015) and Galletta et al. (2019). However, it

is also close to the range of morphological variability of chimpanzees in both PC1 and

PC2, which indicates that the robusticity and curvature of the radioulnar and

dorsovolar ends of the shaft is not similar to what is observed in modern humans. Even

though the LDA robustly classifies *H. naledi* within the *Homo* category, it is worth

mentioning that this specimen occupies a particularly unique position when projected

to the LDA space. This is also observed in its position in the PCA, which seem to

indicate an unusual morphology that can be described as displaying a narrower

proximal end of the body, a relatively broader distal portion of the shaft, as well as

marked attachment sites for the opponens pollicis and dorsal interossei. All these

anatomical attributes contribute to generate the 'pinched' appearance of the palmar

surface of *H. naledi's* MC1 shaft (Kivell et al. 2015b).

Previous analysis of *A. sediba*'s hand morphology has found that it possessed several

advanced *Homo*-like features, such as a longer thumb relative to shorter fingers, that

potentially indicate advanced manipulative capabilities, whilst retaining primitive traits,

such as a gracile MC1, similar to those of other australopiths (Kivell et al., 2011;

Galletta et al., 2019). Our analysis showed that, unlike the Neanderthal and *H. naledi*

specimens, *A. sediba* presented a general morphology that is more similar to

chimpanzees than modern humans. *A. sediba* exhibits smaller epicondyles and a

gracile shaft (Fig. 2b), with relatively flatter muscle attachments at the MC1 (Fig. 2d)

than those observed in *Pan.*

4.2 Functional implications

Even though our study rigorously addresses the anatomical differences among the

MC1s of extant hominines, any functional interpretations that we can advance are

certainly inferred and not directly derived from our results. Hence, caution is required

when interpreting these functional implications because shape differences could result

from several different factors and not only be the result of different manipulative

capabilities. Overall, our 3DGM results are consistent with previous assessments for

the shaft morphology of the extant African apes and fossils hominins, and thus

provides a morphometric support for the functional interpretations made based upon

those features. The large epicondyles and robust shaft presented by the Neanderthal

MC1 sample may suggest that they performed tool use in a very similar fashion to

modern humans (Karakostis, et al., 2018; Niewoehner, 2001, 2006; Tocheri et al.,

2008; Trinkaus & Villemeur, 1991). Nevertheless, the analyzed *H neanderthalensis*

specimen also shows a classic Neanderthal feature (i.e., the opponens pollicis flange),

which clearly distinguish it from the rest of the sample, particularly along PC3. It has

been mentioned that it difficult to evaluate to what extent this trait may reflect muscle

hypertrophy since the actual insertion area is mostly along the radiopalmar margin

rather than across the palmar flange (Trinkaus, 2016). However, it is worth noticing

that the radial extension of the opponens pollicis flange has been interpreted as

increasing the opponens pollicis rotational moment arm, which suggests a greater

mechanical effectiveness of this muscle in this species (Maki & Trinkaus 2011). *H.*

naledi MC1's anatomy suggests that this species was probably able to perform a

certain degree of advanced manipulation, which might imply that this taxon was also

a tool-user due to its robust shaft with marked muscular attachments but small

epicondyles (Berger et al., 2015; Kivell et al., 2015a,b, Galletta et al., 2019). However,

it is also worth considering that *H. naledi* shows an unusual MC1 morphology that

when interpreted in combination with what is known from this species finger anatomy,

may indicate a distinctive behavioral repertoire that could have included tool use as

well as significant amounts of climbing (Kivell et al. 2015b). Finally, *A. sediba*'s

anatomical characteristics suggests incipient tool using capabilities due to its slender

thumb, smaller radial and ulnar epicondyles and curved joint surfaces (Kivell et al.,

2011; Skinner et al., 2015; Galleta et al., 2019). Nevertheless, it is important to keep

in mind that the above interpretations are exclusively based on a morphometric

assessment of the MC1's body anatomy (i.e., we did not directly assess any functional

capabilities). Future studies should try not only to imply functional aspects based on

morphological similarities but rather explicitly include them as part of the study (see

e.g., Bucchi et al., 2020c).

From a functional perspective, the more robust MC1 shaft of humans (Fig. 2b) has

been associated in previous studies with the ability of withstanding higher stresses

placed upon the thumb by sustained power and precision grasping (e.g., Key &

Dunmore, 2015; Marzke, Wullstein, & Viegas, 1992; Rolian, Lieberman, & Zermeno,

2011). These robust thumbs have also been related to a greater development of the

thenar musculature attached to the shaft that is highly active during hard hammer

percussion and favors thumb opposition (Marzke, 2013; Marzke, Toth, Schick, &

Reece, 1998). The pronounced radial and ulnar epicondyles found at the distal head

of the human MC1 (as described by PC1) may help to reduce the range of motion and

stabilize the MCPJ (Imaeda, An, & Cooney, 1992). These epicondyles act as anchor

points for collateral ligaments, which insert at the base of the proximal phalanx. Larger

epicondyles are therefore thought to act as stronger anchors by providing a greater

area for the collateral ligaments to attach to, helping to stabilize the MCPJ during the

high forces that are experienced by the thumb during manipulation (Galletta et al.,

2019). The flatter and larger distal articular surface in humans serves a similar purpose

and has been interpreted as an adaptation that limits dorso-palmar motion whilst

preventing radioulnar motion (Barmakian, 1992), thereby stabilizing the MC1 and

facilitating forceful power and precision grasping.

4.3 Conclusion

The aim of this study was to quantify the morphology of the MC1 shaft in extant African

hominoids, in order to better characterize its morphology. This characterization is not

only relevant to better understand hominine anatomical differences and similarities,

but also to provide further insights about its possible relationship with manipulative

capabilities. This can facilitate more informed functional interpretation of fossil hominin

morphology and contribute towards future studies linking morphology and function in

hominin thumbs. Our study found that each taxon presented a unique repertoire of

morphological traits, not present to the same extent in the others. Overall, the results

obtained both aligned with and added to past functional interpretations of hominin

morphology, thereby reinforcing the validity of 3DGM as a method of quantifying MC1

morphology and providing a deeper insight into the anatomy of the thumb in both

extant hominids and fossil hominins. In addition, fossil MC1s are frequently

fragmentary, and their epiphyses are often damaged. Hence, our applied approach

which exclusively focused on the MC1 shaft might be particularly helpful in

paleoanthropological contexts.

Acknowledgments

We are grateful to the following curators and institutions: Emmanuel Gilissen

(AfricaMuseum), Anneke H. van Heteren and Michael Hiermeier (Zoologische

Staatssammlung München), Javier Quesada (Museu de Ciències Naturals de

Barcelona), José Miguel Carretero (Universidad de Burgos). AB was partially funded

by a Becas Chile scholarship, whilst TP was funded by the Leverhulme Trust Early

Career Fellowship, ECF-2018-264. This study was funded by the research projects

AGAUR 2017 SGR 1040 and MICINN-FEDER PGC2018-093925-B-C32. We are algo

grateful to the two anonymous reviewers, associate editor and editorial board member

who helped us to improve our work.

Author contributions

Jonathan Morley: Data analysis; investigation; methodology; writing-original draft;

writing-review and editing. Ana Bucchi: Conceptualization; methodology; resources;

data curation; writing-review and editing. Carlos Lorenzo: Resources; data curation.

Thomas A. Püschel: Conceptualization; data curation; data analysis; investigation;

visualization; writing-review and editing; project administration.

Data availability statement

The data supporting the findings of this study are available in the supplementary

material (S6) of this article.

ORCID

Ana Bucchi https://orcid.org/0000-0002-1247-230X

Carlos Lorenzo https://orcid.org/0000-0001-5706-293X

Thomas A. Püschel https://orcid.org/0000-0002-2231-2297

References

Adams, D. C., Collyer, M. L., & Kaliontzopoulou, A. (2020). Geomorph: Software for

geometric morphometric analysis. R package version *4.0.2*.

Adler, D., & Kelly, S. T. (2020). vioplot: violin plot. R package version 0.3.5.

603
Almécija, S., Moyà-Solà, S., & Alba, D. M. (2010). Early origin for human-like

precision grasping: A comparative study of pollical distal phalanges in fossil

hominins. *PLOS ONE, 5*(7), e11727.

https://doi.org/10.1371/journal.pone.0011727

Almécija, S., Smaers, J. B., & Jungers, W. L. (2015). The evolution of human and

ape hand proportions. *Nature Communications, 6*, 1–11.

https://doi.org/10.1038/ncomms8717

Barmakian, J. T. (1992). Anatomy of the joints of the thumb. *Hand Clinics, 8*(4), 683–

691.

Berger, L. R., Ruiter, D. J. de, Churchill, S. E., Schmid, P., Carlson, K. J., Dirks, P.

H. G. M., & Kibii, J. M. (2010). Australopithecus sediba: A new species of

Homo-like Australopith from South Africa. *Science, 328*(5975), 195–204.

https://doi.org/10.1126/science.1184944

Bookstein, F. L. (1991). *Morphometric Tools for Landmark Data: Geometry and*

Biology. Cambridge: Cambridge University Press.

Bookstein, F. L. (1997). *Morphometric Tools for Landmark Data: Geometry and*

Biology (2nd edition). Cambridge: Cambridge University Press.

Bucchi, A., Luengo, J., Fuentes, R., Arellano-Villalón, M., Lorenzo, C. (2020a).

Recommendations for improving photo quality in close range

photogrammetry, exemplified in hand bones of chimpanzees and gorillas.

International Journal of Morphology, *38*(2), 348–355.

https://doi.org/10.4067/S0717-95022020000200348

626
627   Bucchi, A., Luengo, J., Bove, A. D., & Lorenzo, C. (2020b). Insertion sites in manual

628   proximal phalanges of African apes and modern humans. *American Journal of*

629   *Physical Anthropology*, 1–12. https://doi.org/10.1002/ajpa.24127

630
Bucchi, A., Püschel, T. A., Lorenzo, C., & Marcé-Nogué, J. (2020c). Finite element

analysis of the proximal phalanx of the thumb in Hominoidea during simulated

stone tool use. Comptes Rendus Palevol, 19(2), 26–39.

Bush, M. E., Lovejoy, C. O., Johanson, D. C., & Coppens, Y. (1982). Hominid carpal,

metacarpal, and phalangeal bones recovered from the Hadar Formation:

1974–1977 collections. *American Journal of Physical Anthropology*, *57*(4),

651–677.

Casillas Garcia, J. A., & Alvarez, G. A. (2005). La actuacion arqueologica en el

"Solar de Caballeria" y el convento de San Pablo de Burgos. *Ayuntamiento de*

Burgos.

Diogo, R., Pastor, J. F., Ferrero, E., Barbosa, M., Potau, J., de Paz, F., … Wood, B.

(2011). Photographic and Descriptive Musculoskeletal Atlas of Gorilla. New

Hampshire: Science Publishers.

Diogo, R., Richmond, B. G., & Wood, B. (2012). Evolution and homologies of primate

and modern human hand and forearm muscles, with notes on thumb

movements and tool use. *Journal of Human Evolution*, *63*(1), 64–78.

https://doi.org/10.1016/j.jhevol.2012.04.001

Diogo, R., Potau, J. M., Pastor, J. F., Paz, F. J. D., Ferrero, E. M., Bello, G., …

Wood, B. (2013). Photographic and Descriptive Musculoskeltal Atlas of

Chimpanzees. New Hampshire: Science Publishers.

Dirks, P. H., Roberts, E. M., Hilbert-Wolf, H., Kramers, J. D., Hawks, J., Dosseto, A.,

… Berger, L. R. (2017). The age of homo naledi and associated sediments in

the rising star cave, South Africa. *ELife*, *6*. https://doi.org/10.7554/eLife.24231

Drake, A. G., & Klingenberg, C. P. (2010). Large-Scale diversification of skull shape

in domestic dogs: Disparity and modularity. *The American Naturalist*, *175*(3),

289–301. https://doi.org/10.1086/650372

Dunmore, C. J., Bardo, A., Skinner, M. M., & Kivell, T. L. (2020). Trabecular variation

in the first metacarpal and manipulation in hominids. *American Journal of*

Physical Anthropology, *171*(2), 219–241.

Feix, T., Kivell, T. L., Pouydebat, E., & Dollar, A. M. (2015). Estimating thumb-index

finger precision grip and manipulation potential in extant and fossil primates.

Journal of the Royal Society, Interface / the Royal Society, *12*(106),

20150176–. https://doi.org/10.1098/rsif.2015.0176

Galletta, L., Stephens, N. B., Bardo, A., Kivell, T. L., & Marchi, D. (2019). Three-

dimensional geometric morphometric analysis of the first metacarpal distal

articular surface in humans, great apes and fossil hominins. *Journal of Human*

Evolution, *132*, 119–136. https://doi.org/10.1016/j.jhevol.2019.04.008

Green, D. J., & Gordon, A. D. (2008). Metacarpal proportions in Australopithecus

africanus. *Journal of Human Evolution*, *54*(5), 705–719.

https://doi.org/10.1016/j.jhevol.2007.10.007

Guérin, G., Frouin, M., Talamo, S., Aldeias, V., Bruxelles, L., Chiotti, L., … Jain, M.

(2015). A multi-method luminescence dating of the Palaeolithic sequence of

La Ferrassie based on new excavations adjacent to the La Ferrassie 1 and 2

skeletons. *Journal of Archaeological Science*, *58*, 147–166.

Gunz, P., & Mitteroecker, P. (2013). Semilandmarks: A method for quantifying

curves and surfaces. *Hystrix, the Italian Journal of Mammalogy*, *24*(1), 103–

109.

Gunz, P., Mitteroecker, P., & Bookstein, F. L. (2005). Semilandmarks in three

dimensions. In D. E. Slice (Ed.), *Modern Morphometrics in Physical*

Anthropology (pp. 73–98). Boston, MA: Springer US.

https://doi.org/10.1007/0-387-27614-9_3

Imaeda, T., An, K. N., & Cooney 3rd, W. P. (1992). Functional anatomy and

biomechanics of the thumb. *Hand Clinics*, *8*(1), 9–15.

Karakostis, F. A., Hotz, G., Scherf, H., Wahl, J., & Harvati, K. (2017). Occupational

manual activity is reflected on the patterns among hand entheses. *American*

Journal of Physical Anthropology, *164*(1), 30–40.

https://doi.org/10.1002/ajpa.23253

Karakostis, F. A., Hotz, G., Tourloukis, V., & Harvati, K. (2018). Evidence for

precision grasping in Neandertal daily activities. *Science Advances*, *4*(9),

eaat2369. https://doi.org/10.1126/sciadv.aat2369

Key, A. J., & Dunmore, C. J. (2018). Manual restrictions on Palaeolithic technological

behaviours. *PeerJ*, *6*, e5399.

Key, A. J. M., & Dunmore, C. J. (2015). The evolution of the hominin thumb and the

influence exerted by the non-dominant hand during stone tool production.

Journal of Human Evolution, *78*, 60–69.

https://doi.org/10.1016/j.jhevol.2014.08.006

Kivell, T. L. (2015a). Evidence in Hand: recent discoveries and the early evolution of

human manual manipulation. *Philosophical Transactions of the Royal Society*,

Kivell, T. L., Deane, A. S., Tocheri, M. W., Orr, C. M., Schmid, P., Hawks, J., …

Churchill, S. E. (2015b). The hand of Homo naledi. *Nature Communications*,

6, 8431.

Kivell, T. L., Kibii, J. M., Churchill, S. E., Schmid, P., & Berger, L. R. (2011).

Australopithecus sediba hand demonstrates mosaic evolution of locomotor

and manipulative abilities. *Science*, *333*(6048), 1411–1417.

https://doi.org/10.1126/science.1202625

Klingenberg, C. P., & McIntyre, G. S. (1998). Geometric morphometrics of

developmental instability: Analyzing patterns of fluctuating asymmetry with

Procrustes methods. *Evolution*, *52*(5), 1363–1375.

https://doi.org/10.2307/2411306

Kuhn, M., & Johnson, K. (2013). Measuring Performance in Classification Models. In

Applied Predictive Modeling (pp. 247–273). Springer, New York, NY.

https://doi.org/10.1007/978-1-4614-6849-3_11

Maki, J., & Trinkaus, E. (2011). Opponens pollicis mechanical effectiveness in

Neandertals and early modern humans. *PaleoAnthropology*, 62−71.

https://doi.org/doi:10.4207/PA.2011.ART43

Marchi, D., Proctor, D. J., Huston, E., Nicholas, C. L., & Fischer, F. (2017).

Morphological correlates of the first metacarpal proximal articular surface with

manipulative capabilities in apes, humans and South African early hominins.

Comptes Rendus Palevol, *16*(5), 645–654.

https://doi.org/10.1016/j.crpv.2016.09.002

Marzke, M. W. (2013). Tool making, hand morphology and fossil hominins.

Philosophical Transactions of the Royal Society B: Biological Sciences,

368(1630), 20120414. https://doi.org/10.1098/rstb.2012.0414

Marzke, M. W., Tocheri, M. W., Steinberg, B., Femiani, J. D., Reece, S. P.,

Linscheid, R. L., … Marzke, R. F. (2010). Comparative 3D quantitative

analyses of trapeziometacarpal joint surface curvatures among living

catarrhines and fossil hominins. *American Journal of Physical Anthropology*,

141(1), 38–51. https://doi.org/10.1002/ajpa.21112

Marzke, Mary W, Toth, N., Schick, K., & Reece, S. (1998). EMG study of hand

muscle recruitment during hard hammer percussion manufacture of Oldowan

tools. *American Journal of Physical Anthropology 105*(3), 315–332.

Matarazzo, S. A. (2015). Trabecular architecture of the manual elements reflects

locomotor patterns in primates. *PLoS ONE*, *10*(3), e0120436.

https://doi.org/10.1371/journal.pone.0120436

McClure, N., Phillips, A., Vogel ER, Tocheri, M. (2012). Unexpected pollex and

hallux use in wild Pongo pygmaeus wurmbii. *American Journal of Physical*

Anthropology, 147(S54): S208.

Menger, K. (1930). Untersuchungen über allgemeine Metrik. Vierte Untersuchung.

Zur Metrik der Kurven. Mathematische Annalen, 103, 466–501.

Niewoehner, W. A. (2001). Behavioral inferences from the Skhul/Qafzeh early

modern human hand remains. *Proceedings of the National Academy of*

Sciences, *98*(6), 2979–2984. https://doi.org/10.1073/pnas.041588898

Niewoehner, W. A. (2006). Neanderthal hands in their proper perspective. In

Neanderthals revisited: New approaches and perspectives (pp. 157–190).

Springer.

746 Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn,

747   D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P.,Henry, M.,

748   Stevens, H., Szoecs, E., & Wagner, H. (2020). vegan:Community Ecology

749   Package. R package version 2.5-7.

750
Perez, S. I., Bernal, V., & Gonzalez, P. N. (2006). Differences between sliding semi-

landmark methods in geometric morphometrics, with an application to human

craniofacial and dental variation. *Journal of Anatomy*, *208*(6), 769–784.

https://doi.org/10.1111/j.1469-7580.2006.00576.x

Pickering, R., Dirks, P. H. G. M., Jinnah, Z., Ruiter, D. J. de, Churchill, S. E., Herries,

A. I. R., … Berger, L. R. (2011). Australopithecus sediba at 1.977 Ma and

Implications for the Origins of the Genus Homo. *Science*, 333(6048), 1421–

1423. https://doi.org/10.1126/science.1203697

Püschel, T. A., Marcé-Nogué, J., Chamberlain, A. T., Yoxall, A., & Sellers, W. I.

(2020). The biomechanical importance of the scaphoid-centrale fusion during

simulated knuckle-walking and its implications for human locomotor evolution.

Scientific Reports, *10*(1), 1–7. https://doi.org/10.1038/s41598-020-60590-6

Püschel, T. A., Marcé-Nogué, J., Gladman, J., Patel, B. A., Almécija, S., & Sellers,

W. I. (2020). Getting its feet on the ground: Elucidating Paralouatta's semi-

terrestriality using the virtual morpho-functional toolbox. *Frontiers in Earth*

Science, *8*. https://doi.org/10.3389/feart.2020.00079

Püschel, T. A., Marcé-Nogué, J., Gladman, J. T., Bobe, R., & Sellers, W. I. (2018).

Inferring locomotor behaviours in Miocene New World monkeys using finite

element analysis, geometric morphometrics and machine-learning

classification techniques applied to talar morphology. *Journal of The Royal*

Society Interface, *15*(146), 20180520. https://doi.org/10.1098/rsif.2018.0520

Rein, T. R. (2011). The correspondence between proximal phalanx morphology and

locomotion: Implications for inferring the locomotor behavior of fossil

catarrhines. *American Journal of Physical Anthropology*, *146*(3), 435–445.

https://doi.org/10.1002/ajpa.21599

Richmond, B. G., Green, D. J., Lague, M. R., Chirchir, H., Behrensmeyer, A. K.,

Bobe, R., … Braun, D. R. (2020). The upper limb of Paranthropus boisei from

Ileret, Kenya. *Journal of Human Evolution*, *141*, 102727.

https://doi.org/10.1016/j.jhevol.2019.102727

Richmond, Brian G., & Strait, D. S. (2000). Evidence that humans evolved from a

knuckle-walking ancestor. *Nature*, *404*(6776), 382.

Rolian, C., & Gordon, A. D. (2013). Reassessing manual proportions in

Australopithecus afarensis. *American Journal of Physical Anthropology*,

152(3), 393–406. https://doi.org/10.1002/ajpa.22365

Rolian, C., Lieberman, D. E., & Zermeno, J. P. (2011). Hand biomechanics during

simulated stone tool use. *Journal of Human Evolution*, *61*(1), 26–41.

https://doi.org/10.1016/j.jhevol.2011.01.008

Stephens, N. B., Kivell, T. L., Gross, T., Pahr, D. H., Lazenby, R. A., Hublin, J. J., …

Skinner, M. M. (2016). Trabecular architecture in the thumb of Pan and Homo:

implications for investigating hand use, loading, and hand preference in the

fossil record. *American Journal of Physical Anthropology*, *161*(4), 603–619.

https://doi.org/10.1002/ajpa.23061

Stern, J. T., Jungers, W. L., & Susman, R. L. (1995). Quantifying phalangeal

curvature: an empirical comparison of alternative methods. *American Journal

of Physical Anthropology*, *97*(1), 1–10.

Susman, R. (1994). Fossil evidence for early hominid tool use. *Science*, *265*(5178),

1570–1573. https://doi.org/10.1126/science.8079169

Susman, R. L. (1979). Comparative and functional morphology of hominoid fingers.

American Journal of Physical Anthropology, *50*(2), 215–236.

Tocheri, M. W., Orr, C. M., Jacofsky, M. C., & Marzke, M. W. (2008). The

evolutionary history of the hominin hand since the last common ancestor of

Pan and Homo. *Journal of Anatomy*, *212*(4), 544–562.

https://doi.org/10.1111/j.1469-7580.2008.00865.x

Trinkaus E. (1983). *The Shanidar Neandertal*s. Academic Press, New York

Trinkaus, E. (2016). The Evolution of the Hand in Pleistocene Homo. In T. L. Kivell,

P. Lemelin, B. G. Richmond, & D. Schmitt (Eds.), The Evolution of the Primate

Hand (pp. 545–571). New York, NY: Springer New York.

https://doi.org/10.1007/978-1-4939-3646-5_19

Trinkaus, E., & Villemeur, I. (1991). Mechanical advantages of the Neandertal thumb

in flexion: A test of an hypothesis. *American Journal of Physical Anthropology*,

84(3), 249–260. https://doi.org/10.1002/ajpa.1330840303

Trinkaus E, Haduch E, Valde-Nowak PW, Wojtal P (2014) The Obłazowa 1 early

modern human pollical phalanx and Late Pleistocene distal thumb

proportions. *Homo* 65, 1–12.

Tsegai, Z. J., Kivell, T. L., Gross, T., Nguyen, N. H., Pahr, D. H., Smaers, J. B., &

Skinner, M. M. (2013). Trabecular bone structure correlates with hand posture

and use in hominoids. *PLoS One*, *8*(11), e78781.

Tuttle, R. H. (1969). Quantitative and functional studies on the hands of the

Anthropoidea. I. The Hominoidea. *Journal of Morphology*, *128*(3), 309–363.

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.).

New York: Springer-Verlag.

Watanabe, A. (2017). LaMBDA: LandMark-Based Data Assessment. R package

version 0.1.0.9000.

Wiley, D. F., Amenta, N., Alcantara, D. A., Ghost, D., Kil, Y. J., Delson, E., …

Hamann, B. (2005). Evolutionary morphing. *EEE Visualization, 2005. VIS 05.*

Zelditch, M. L., Swiderski, D. L., & Sheets, H. D. (2012a). *Geometric Morphometrics*

for Biologists: A Primer (2 edition). Amsterdam: Academic Press.

Zelditch, M. L., Swiderski, D. L., & Sheets, H. D. (2012b). Chapter 9 - General Linear

Models. In *Geometric Morphometrics for Biologists* (2 edition, pp. 209–228).

San Diego: Academic Press.

831
832

Figure legends

Figure 1. Illustration of the 16 landmarks (yellow spheres) and 144 semi-landmarks

(red spheres) used in this study. The numbers of the 16 fixed landmarks and eight

semi-landmark curves (C1-C8) are also plotted.

Figure 2. Principal component analysis of the shape data: the a) three main axes of

morphological variation are displayed (ellipses represent 95% confidence intervals,

red spheres: fossils, orange spheres*: H. sapiens,* green spheres: *P. troglodytes*;

golden spheres: *G. gorilla*, golden cubes: *G. beringei*); Violin plots of the PCs scores

of the analyzed sample are shown for b) PC1, c) PC2 and d) PC3 (fossil values are

displayed as red triangles). The white dot in the middle is the median value, whilst the

thick black bar in the center represents the interquartile range. The thin black line

extended from it corresponds to the upper (maximum) and lower (minimum) adjacent

values in the data. The distribution shape of the data for each one of the three PCs is

represented by a kernel density plots that were rotated and placed on each side of

each one of the boxplots. To visualize shape differences warped models representing

the shape changes along the first three principal components (PCs) were plotted

alongside the violin plots (dorsal views). The models closest to the mean shape was

to match the multivariate mean using the thin plate spline method. Then, the obtained

average model was warped to display the variation along the three plotted PC axes

(mag = 1).

Figure 3. Warped models representing the shape changes along the first three

principal components (PCs). The models closest to the mean shape was to match the

multivariate mean using the thin plate spline method. Then, the obtained average

model was warped to display the variation along the three plotted PC axes (mag = 1).

Corresponding anatomical descriptions are provided alongside each one the warped

models. Please notice that articular surfaces were not morphometrically characterized

and as such, none of the anatomical descriptions refer to them. References to

curvature on this figure are based on the results provided in Table 5.

Figure 4. Multi-group linear discriminant analysis (LDA) of MC1's shape using extant

genera categories. One of the models closest to the mean shape was warped to match

the multivariate mean using the thin plate spline method, then the obtained average

model was warped to represent the variation along the two plotted CV axes.

PC1 (−)
- Slender and gracile shaft
- Marked antero-posterior curvature of the shaft on the palmar side
- Narrower distal-most and proximal-most portions of the shaft
- Reduced radial and ulnar epicondyles

PC1 (+)
- Robust and broader shaft
- Straight dorsal aspect of the body
- Broader distal-most and proximal-most portions of the shaft
- Expanded ulnar and radial epicondyles

41.44%

PC2 (−)
- Extended proximal palmar lip
- More curved shaft on both palmar and dorsal sides with radioulnar and dorsovolar rounded ends
- Medial epicondyle more distal relative to the lateral epicondyle
- Wider proximal-most portion of the shaft

PC2 (+)
- Flatter distal-most and proximal-most portions of the shaft
- Expanded area at the place of origin of the radio-palmar condyles at the distal end
- Proximal-most and distal-most portions of the shaft show similar relative widths
- Medial epicondyle more proximal relative to the proximal epicondyle

11.18%

PC3 (−)
- More curved palmar appearance of the shaft
- Proximally displaced epicondyles
- More gracile overall morphology

PC3 (+)
- Lateral flange for the opponens pollicis
- Waisted palmar and dorsal appearance
- Distally displaced epicondyles
- Broader distal-most portion of the shaft

5.82%