Characterizing the body morphology of the first metacarpal in the Homininae

using 3D geometric morphometrics

Jonathan Morley[1], Ana Bucchi[2,3], Carlos Lorenzo[2,3], Thomas A. Püschel[4*]

[1]Department of Zoology, University of Oxford, South Parks Road, OX1 3PS, Oxford,

United Kingdom.

[2]Àrea de Prehistòria, Universitat Rovira i Virgili (URV), Avinguda de Catalunya 35,

43002 Tarragona, Spain.

[3]Institut Català de Paleoecologia Humana i Evolució Social (IPHES), Zona

Educacional 4, Campus Sescelades URV (Edifici W3), 43007 Tarragona, Spain.

[4] Primate Models for Behavioural Evolution Lab, Institute of Cognitive and Evolutionary

Anthropology, School of Anthropology, University of Oxford, 64 Banbury Road, OX2

6PN, Oxford, United Kingdom.

*Corresponding author:

Thomas A. Püschel

E-mail: thomas.puschelrouliez@anthro.ox.ac.uk

Phone: +44 (0) 7476608464

Running title: Characterizing MC1's morphology in the Homininae

Abstract

Objectives: Extinct hominins can provide key insights into the development of tool use, with the morphological characteristics of the thumb of particular interest due to its fundamental role in enhanced manipulation. This study quantifies the shape of the first metacarpal's body in the extant Homininae and some fossil hominins to provide insights about the possible anatomical correlates of manipulative capabilities.

Materials and methods: The extant sample includes MC1s of modern humans (n=42), gorillas (n=27) and chimpanzees (n=30), whilst the fossil sample included *Homo neanderthalensis*, *Homo naledi* and *Australopithecus sediba*. 3D geometric morphometrics were used to characterize the overall shape of MC1's body.

Results: Humans differ significantly from extant great apes when comparing overall shape. *H. neanderthalensis* mostly falls within the modern human range of variation although also showing a more robust morphology. *H. naledi* varies from modern human slightly, whereas *A. sediba* varies from humans to an even greater extent. When classified using a linear discriminant analysis, the three fossils are categorized within the *Homo* group.

Discussion: The results are in general agreement with previous studies on the morphology of the MC1. This study found that the modern human MC1 is characterized by a distinct suite of traits, not present to the same extent in the great apes, that are consistent with an ability to use forceful precision grip. This morphology was also found to align very closely with that of *H. neanderthalensis*. *H. naledi* shows a number of human-like adaptations consistent with an ability to employ enhanced manipulation, whilst *A. sediba* apparently presents a mix of both derived and more primitive traits.

Keywords: MC1; 3D geometric morphometrics; Semi-landmarks; Hominins; African apes

1 Introduction

There is no doubt that the extremely dexterous human hand is unmatched among animals. As a result, the human hand has been the subject of considerable paleoanthropological research over the years in order to provide a better understanding of the processes that led to its evolution (Almécija, Smaers, & Jungers, 2015a; Key, 2016; Lewis, 1977; Marzke & Marzke, 2000; Susman, 1998). Much of this has been directed at how the hand morphology of higher primates correlates to their manipulative capabilities, and how fossil morphology can be used to infer tool use in extinct hominin species (Almécija & Alba, 2014; Kivell, Kibii, Churchill, Schmid, & Berger, 2011; Marchi, Proctor, Huston, Nicholas, & Fischer, 2017; Napier, 1955). The thumb and its components, most notably the first metacarpal (MC1), plays a fundamental role in object manipulation and the study of its anatomy has therefore been at the center of research in this field (Galletta, Stephens, Bardo, Kivell, & Marchi, 2019; Marchi et al., 2017). However, the constant discovery of new fossils (e.g. *Homo Naledi;* Berger et al., 2015) and the development of new morpho-functional analysis tools mean that there is still much about the tool use behaviors and manipulative capabilities of extinct hominins that is yet to be uncovered.

Amongst the extant great apes, humans possess the superior manipulative capabilities, with the ability not only to adeptly utilize the objects in their environment, but also to manufacture complex tools in ways that require high levels of dexterity. These advanced skills are facilitated in part by a unique thumb morphology: the human thumb is long relative to the length of the fingers compared to other less dexterous

apes, with powerful thenar musculature and robust thumb bones (Almécija, Moyà-
Solà, & Alba, 2010; Almécija et al., 2015; Feix, Kivell, Pouydebat, & Dollar, 2015;
Tuttle, 1969).

In comparison to human hands, non-human great apes, especially chimpanzees and
orangutans, have longer, robust fingers relative to their shorter, more gracile thumbs,
which is probably the result of selective pressures associated with locomotor
behaviors such as suspension and knuckle-walking (Almécija et al., 2015; Püschel,
Marcé-Nogué, Chamberlain, Yoxall, & Sellers, 2020; Richmond & Strait, 2000). Both
gorillas and chimpanzees have been observed using tools in the wild, to varying
degrees; however, their manipulative capabilities are limited due to the constraints
imposed by their hand morphology. Chimpanzees are prolific tool users, known to use
tools both in nature and captivity (Boesch & Boesch, 1990). Examples of chimp tool
use include termite fishing with specially crafted sticks (Sanz, Call, & Morgan, 2009),
hunting bush babies with sharp spears and nut-cracking with stones (Sanz & Morgan,
2007). Gorillas are less reliant on tool use due the fact that they exploit food resources
differently from chimpanzees (e.g., crack nuts with their teeth), but they have been
observed using sticks to test the depth of water and to support themselves when
crossing deep water (Breuer, Ndoundou-Hockemba, & Fishlock, 2005). However,
whilst non-human great apes do regularly use their thumbs to manipulate objects, they
are not as efficient as humans in using forceful precision grips (Marzke, Marchant,
McGrew, & Reece, 2015; Marzke & Wullstein, 1996).

Traditionally it was believed that extinct hominin species also fell into this category,
lacking the manual dexterity of modern humans (Lewis, 1977; Niewoehner, 2001,
2006; Rightmire, 1972). However, there is a growing body of evidence that many

hominins as early as *O. tugenensis* (Gommery & Senut, 2006) show the capacity to

efficiently use tools, with the ability to use forceful precision grips as dexterously as

humans (Alba, Moyà-Solà, & Köhler, 2003; Feix et al., 2015; Karakostis, Hotz, Scherf,

Wahl, & Harvati, 2017; Kivell et al., 2011; Tocheri, Orr, Jacofsky, & Marzke, 2008).

Human-like features would have been present along with traits suitable for arboreal

locomotion, which lead to the suggestion that the hominin hand evolved in a mosaic

fashion showing a manual morphology adapted to these two functional demands (i.e.,

manipulation and locomotion) (Kivell et al., 2011; Kivell, 2015). This mixed morphology

is apparent in the hand of *A. sediba* and *H. naledi* (Kivell et al., 2011; 2015) whereas

the hand of Neanderthals would be fully derived (Tocheri et al., 2008).

Several studies have focused on different anatomical features of the hand in an effort

to understand the extent to which the hand of early hominins is adapted to

manipulative abilities (e.g., Almécija et al., 2010; Galletta et al., 2019; Green & Gordon,

2008; Skinner et al., 2015), in particular regarding the joint areas of the MC1, as this

bone plays a crucial role in complex manipulative behaviors. However, whilst there is

now a greater understanding of the manipulative capabilities of hominins, much of the

research on the MC1 dates back to the past century and often contained only

qualitative assessments (Aubriot & Tubiana, 1981; Barmakian, 1992; Imaeda, An, &

Cooney, 1992; Napier, 1956, 1960; Tuttle, 1969). Even the most recent quantitative

research that has been conducted using three-dimensional geometric morphometric

(3DGM) techniques have focused only on those certain areas of the MC1 deemed to

be most important in controlling manipulation, such as the trapeziometacarpal joint

connecting the thumb to the wrist (i.e., proximal articular surface; Marchi et al., 2017)

and the first metacarpal distal articular surface (Galletta et al., 2019). As a result, most

of MC1's morphology (i.e., its body) has yet to be fully quantitatively analyzed to

assess its possible importance when assessing possible correlates with manipulative

abilities.

Consequently, in this study body morphology of the MC1 was quantified using 3DGM

in order help in the identification of structures in extant species that may be correlated

with human-like manipulative capabilities and determining if similar morphologies are

present in fossil hominins. The sample investigated in this study included three extant

African ape genera (*Homo, Gorilla, Pan*) and three fossil hominins (*Homo*

neanderthalensis, *Homo naledi* and *Australopithecus sediba*). Based on previous

literature about thumb morphology and function, the following hypotheses were tested:

Hypothesis 1: MC1 morphology significantly differs between humans and extant

great ape species

Though great apes use their hand for manipulative activities, their specialisation is

more a consequence of their locomotion (i.e., knuckle-walking and arborealism)

(Almécija, Moyà-Solà, & Alba, 2010). It is therefore expected that the selective

pressures associated with locomotor behaviour in chimpanzees and gorillas will result

in an MC1 morphology that varies significantly from that of bipedal humans.

Furthermore, different use of the human thumb during manipulation and human

adaptations to precise and forceful tool use are expected to lead to an MC1

morphology that differs from other extant apes.

Hypothesis 2: All fossil hominin specimens exhibit an MC1 morphology more

similar to humans than other great apes

H. neanderthalensis, A. sediba and *H. naledi* have overall hand morphologies that

appear to align with human hands to a greater extent than those of non-human great

apes. They possess adaptations, such a long thumb and short fingers, that are associated with advanced manipulative capabilities in modern humans (Holliday et al., 2018; Kivell et al., 2011). Given these morphological characteristics and the inferred tool using abilities of *H. naledi, A. sediba* and *H. neanderthalensis* in previous studies, they would be expected to have an MC1 morphology more closely aligned with humans than gorillas or chimpanzees.

2 Material and methods

2.1 Sample

The extant sample used in this study includes MC1s of modern humans (*Homo sapiens*; n=42), chimpanzees (*Pan troglodytes*; n=30), and gorillas (*Gorilla gorilla* and *Gorilla beringei*; n=27) (Table S1). The human MC1s came from a medieval cemetery in Burgos, Spain (Casillas Garcia & Alvarez, 2005) and the surface models were obtained using a Breuckmann SmartSCAN structured light scanner. The non-human sample came from museum collections and they came from different origins (i.e., wild-shot, captivity and unknown origin). Their surface models were collected using photogrammetry as described in Bucchi et al., (2020). Both scanned and photogrammetry models are high resolution, therefore providing a good representation of the original anatomy. The resolution of the models generated using surface scanner and photogrammetry have been previously tested and found to be comparable (Giacomini et al., 2019), thus allowing us to combine these data types in our analyses. Only adult individuals were included in the study and right MC1s were preferred (although left MC1s were reflected when their antimere was not present).

The fossil sample includes the right metacarpal from a *Homo neanderthalensis*, the right metacarpal from a *Homo naledi* and the left metacarpal from an *Australopithecus*

sediba. The *H. neanderthalensis* sample was found in La Ferrassie archaeological site

in Savignac-de-Miremont, France. The skeleton was discovered in 1909 and is

estimated to be 70–50,000 years old (Guérin et al., 2015). The *Homo naledi* sample

(Morphosource identifier: S2110) was recovered in 2013 from the Rising Star cave

system in South Africa and has been dated to around 250,000 years ago (Dirks et al.,

2017). The *A. sediba* sample (Morphosource identifier: S2490) was taken from the

near complete wrist and hand of an adult female [Malapa Hominin 2 (MH2)] discovered

in Malapa, South Africa (Berger et al., 2010). The latter fossils were downloaded from

Morphosource https://www.morphosource.org/, whereas the Neanderthal was

obtained from a cast housed at the Catalan Institute of Human Paleoecology and

Social Evolution (IPHES).

2.2. 3DGM

3D coordinates were collected using the software Landmark Editor 3.6 (Wiley et al.,

2005) to quantify the MC1's morphology. Eight curves comprising 20 equidistant

coordinates each were placed at pre-defined points on the MC1 (Figure 1). These

coordinates were chosen to provide a good representation of the overall shape of the

shaft of the bone. The first and last coordinates from each one of the eight curves were

treated as fixed landmarks, whereas all the rest of the coordinates (i.e., 144

coordinates) were considered as semi-landmarks. A generalized Procrustes

superimposition was performed on the coordinate data to remove differences due to

scale, translation, and rotation, thus obtaining shape variables (Bookstein, 1991). The

semi-landmarks were slid on the MC1's surface by minimizing bending energy

(Bookstein, 1997; Gunz, Mitteroecker, & Bookstein, 2005).

These obtained shape variables were then used in a principal component analysis (PCA) to quantify overall shape variation. The data set of extant hominoids was then grouped by genus and the Procrustes variance of observations in each group (i.e., the mean squared Procrustes distance of each specimen from the mean shape of the respective group) was computed as a simple measure to assess morphological disparity within each one (Klingenberg & McIntyre, 1998; Zelditch, Sheets, & Fink, 2003). Procrustes variance was applied here as way to evaluate intra-genus variation, and absolute differences in Procrustes variances were computed to test differences in morphological disparity among groups (these differences statistically evaluated through permutation). Then, a multi-group linear discriminant analysis (LDA) (also known as canonical variate analysis) was run to maximize separation between groups using the principal components (PCs) that accounted for 90% of the sample variance. Performance was calculated using the confusion matrix from which the overall classification accuracy was computed, as well as the Cohen's Kappa statistic (Püschel, Marcé-Nogué, Gladman, et al., 2020; Püschel, Marcé-Nogué, Gladman, Bobe, & Sellers, 2018). The complete dataset was resampled using a 'leave-one-subject-out' cross-validation, as a way to asses classification performance (Kuhn & Johnson, 2013). Pairwise PERMANOVA tests with Bonferroni corrections for multiple comparisons were performed to assess for shape differences between the three extant genera using the again PCs that accounted for 90% of the sample variance. Euclidean distances were selected as similarity index.

All these analyses were carried out in R 3.5.1 (R Core Team, 2019), using the 'geomorph' 3.1.2 (Adams, Collyer, & Kaliontzopoulou, 2019) and 'MASS' 7.3-51.5 packages (Venables & Ripley, 2002).

Figure 1. Illustration of the 16 landmarks (black spheres) and 144 semi-landmarks (red spheres) used to quantify MC1's body morphology.

3 Results

3.1 Principal component analysis

The PCA performed using the shape variables returned 102 PCs. The first 22 PCs accounted for ~ 90% of the total variance of the sample, hence offering a reasonable estimate of the total amount of MC1's shape variation, which were then used in the LDA and pairwise PERMANOVA tests. The first three PCs in the PCA account for ~ 57% of the total variance and display a relatively clear separation between the extant African ape genera (Fig. 2a). PC1 explains 40.8 %, PC2 10.44% and PC3 5.43% of total variance, respectively (Fig. 1a-d).

Violin plots of PC1 (Fig. 2b) show a notable difference between gorillas and humans vs. chimpanzees. Humans and gorillas exhibit the highest PC1 scores, representing a wider distal articular surface, a larger proximal articular surface, a significantly more

robust shaft. Chimpanzees show the lowest PC1 scores, representing a narrower

proximal articular surface, a smaller distal head, smaller radial and ulnar epicondyles

and a more gracile shaft. *H. neanderthalensis* falls within the human and gorilla

distributions and is distinct completely from the chimpanzees. *H. naledi* falls within the

human distribution, whilst *A. sediba* is characterized by a lower PC1 score and aligns

closer to the *Pan* distribution. None of the analyzed fossils fall within any of

interquartile ranges (IQR) (i.e., black bars in Fig. 1b-d) of any of the extant genera.

Violin plots of PC2 (Fig. 2c) shows distinct variation between the extant genera, with

a morphological continuum ranging from *Gorilla* (higher PC2 values), *Pan* (central PC2

values) and *H. sapiens* (lower PC2 scores). Interestingly, due to the presence of a

couple of outliers, the morphological variation in *Gorilla* encompasses the whole range

of observed morphological variation. The *Gorilla* sample has the highest PC2 scores,

representing an extended palmar lip, a more curved shaft and more rounded ends.

The modern human distribution shows the lowest PC2 scores, representing flatter

distal and proximal articular ends, as well as larger radial palmar condyles at the distal

end. The chimpanzee sample lies in between the gorilla and modern human samples

displaying an intermediate morphology. In a similar fashion as chimpanzees, the three

fossils are located at intermediate positions in PC2 distribution. *H. neanderthalensis*

and *H. naledi* display PC2 scores that are within the *Pan* IQR, whilst *A. sediba* has

lower values.

Violin plots of PC3 (Fig. 2d) show a similar distribution of PC scores for the three extant

genera. From a morphological perspective, higher scores are associated with more

robust morphologies displaying more marked muscular attachments (for the opponens

pollicis, first dorsal interosseous and abductor pollicis longus muscles), while lower

values correspond to more gracile MC1s. *H. naledi* and *A. sediba* show values which

are within the *Pan* or *H. sapiens* distribution, but outside their IQR and at opposite

extremes of the axis. *H. neanderthalensis* lies outside the distribution of any of the

extant genera, probably due to its particularly robust morphology and associated

marked muscular insertion areas.

Figure 2. Principal component analysis of the shape data: the a) three main axes of morphological variation are

displayed (ellipses represent 95% confidence intervals, whilst fossils are shown as red tetrahedrons); Violin plots

of the PCs scores of the analyzed sample are shown for b) PC1, c) PC2 and d) PC3 (fossil values are displayed

as red triangles). The models closest to the mean shape was warped to match the multivariate mean using the thin

plate spline method. The obtained average model was then warped to represent the variation along the three

plotted PC axes (mag = 1).

3.2 Morphological disparity

To compare the amounts of shape variation between the extant genera, we used

Procrustes variance as a way to assess intra-genus variation. The obtained results

show that three extant genera show a similar magnitude of disparity. Nevertheless,

gorillas exhibit a higher Procrustes variance as compared to modern humans and

chimpanzees (Table 1a). Gorillas are significantly different to modern humans, and

chimpanzees when comparing absolute variance differences, whilst modern human

do not significantly differ from chimpanzees (Table 1b).

Table 1. Morphological disparity results

a) Procrustes variance			
Chimpanzees	0.004840776		
Gorillas	0.006378662		
Modern humans	0.004423507		
b) Pairwise differences			
	Chimpanzees	Gorillas	Modern humans
Chimpanzees		**0.009**	0.393
Gorillas	0.001537886		**0.001**
Modern humans	0.000417269	0.001955155	

Above the diagonal: p-values (significant in bold); below the diagonal:

absolute differences.

3.3 Linear discriminant analysis

The LDA model using the first 22 PCs clearly distinguishes between the three extant

genera, displaying an outstanding performance with almost perfect classification

results after cross-validation (Accuracy: 0.98; Cohen's Kappa: 0.97; Figure 3). When

using the obtained discriminant function to classify the fossils into the extant categories

(as way of assessing morphological affinities) (Table 2), the three of them were

robustly classified into the *Homo* category (all posterior probabilities were extremely

close to 1), hence indicating that, in spite of their differences, their morphology is closer

to that of modern humans. There were significant differences between all extant

genera when analyzing 22 PCs from the PCA carried out using the shape variables

(Table 3).

Figure 3. Decision boundary plot for the first 22 PCs the PCA carried out using the shape variables.

The two variables that contributed the most to each LDA models are displayed (i.e., PC1 and PC2).

The space is colored depending on what substrate preference the (LDA model predict that region

belongs to, whereas the lines between colored areas represent the decision boundaries. Color intensity

indicates the certainty of the prediction in a particular graph area (i.e., darker colors mean a higher

probability of belonging to a certain category). Symbols surrounded by a white rim correspond to

misclassified specimens for the plotted variables, whilst fossil are in red.

Table 2. Prediction results for the fossil sample.

	LDA model		
	Posterior probabilities		
Species	Chimpanzees	Gorillas	Modern humans
Australopithecus sediba	0.09	0.00	0.91
Homo naledi	0.00	0.00	1.00
Homo neanderthalensis	0.00	0.00	1.00

Table 3. Pairwise PERMANOVA results

	F-model	Bonferroni-corrected p-value
Modern humans vs. gorillas	15.84	**0.0003**
Modern humans vs. chimpanzees	68.87	**0.0003**
Chimpanzees vs. gorillas	42.03	**0.0003**

4 Discussion

The first hypothesis was that the shape of the human MC1 would differ significantly

from that of *Pan* and *Gorilla*, due to the variation in their manipulative capabilities and

locomotive behaviors. Results from the analyses provide strong support for this

hypothesis, confirming that there is indeed significant morphological variation between

the extant great apes. Interestingly, we also found clear differences between

chimpanzees and gorillas, with gorillas closer (i.e., more similar) to humans than to

chimpanzees (PC1). The second hypothesis was that all fossil hominin species would

exhibit an MC1 morphology more similar to humans than other great apes. The results

also support this hypothesis. However, it is important to notice that even though the

three fossils are more similar to the modern humans, they also display some distinct

features, different from those which would be typically expected in modern *H. sapiens*.

5.1 Humans and great apes MC1 shape

The 3DGM data indicate that modern human MC1 morphology is significantly different

from the rest of the extant hominids, therefore allowing us to accept the first

hypothesis. The human distal head is characterized by a flatter, larger distal articular

surface and larger radial and ulnar epicondyles. The proximal base of the human MC1

is also larger and flatter in both the radioulnar and dorsovolar aspects, with less

pronounced curvature than that seen in other hominid species. These are all

morphologies that are consistent with previous 3DGM analysis of the proximal (Marchi

et al., 2017) and distal (Galletta et al. 2019) surfaces of the human MC1. The shaft,

an area that has not previously been analyzed using 3DGM, is characterized by being

significantly more robust, with a greater curvature and a larger ridge on its lateral side,

corresponding to the insertion of the opponens pollicis muscle.

The flatter and larger distal articular surface in humans has been interpreted as an

adaptation that limits dorso-palmar motion whilst preventing radioulnar motion

(Barmakian, 1992), thereby stabilizing the MC1 and facilitating forceful power and

precision grasping. In apes that distal articular surface has a more pronounced

curvature, rendering the metacarpophalangeal joint (MCPJ) less stable and unable to

sustain high loads (Galletta et al., 2019). The pronounced radial and ulnar epicondyles

found at the distal head of the human MC1 (as described by PC1) serve a similar

purpose, reducing the range of motion and stabilizing the MCPJ. These epicondyles

act as anchor points for collateral ligaments, which insert at the base of the proximal

phalanx. When the thumb is flexed these ligaments tighten and limit the radioulnar

motion of the proximal phalanx (Imaeda et al., 1992). Larger epicondyles are therefore

thought to act as stronger anchors by providing a greater area for the collateral

ligaments to attach to, helping stabilize the MCPJ during the high forces that are

experienced by the thumb during manipulation (Galletta et al., 2019). The proximal

articular surface in humans is also flatter (as described by PC2), but in contrast this is

correlated with a higher range of motion at the trapeziometacarpal joint (TMCJ), rather

than a lower one (Marzke et al., 2010). It is this combination in humans of high mobility

at the TMCJ and low mobility at the MCPJ that facilitates a high level of manual

dexterity, whilst also allowing the thumb to sustain high loads during forceful tool use.

High mobility at the TMCJ plays a key role in the pad-to-pad opposition abilities of the

human hand, in which the thumb is able to rotate and touch the apical tip of each

phalanx. In many human manipulative activities like precision grips the thumb needs

to be highly abducted, which means that the load is radially shifted on the joint surface

(Lewis, 1977; Marchi et al., 2017). The observed larger radially extended proximal

surface is therefore important because, whilst it allows for a greater radial extension,

it also helps the joint resist high levels of radial displacement by providing a greater

surface area for the abducted MC1 (Hamrick, 1996).

We hypothesize that the morphological characteristics of the human MC1 shaft

presented here, such as a significantly more robust build, are likely adaptations that

further serve to facilitate forceful tool use. Indeed, a thicker MC1 shaft would be able

to better withstand the high levels of stress placed upon the thumb by sustained power

and precision grasping (Key & Dunmore, 2015; Marzke, Wullstein, & Viegas, 1992;

Rolian, Lieberman, & Zermeno, 2011). It has been also related to a greater

development of the thenar musculature attaching into the shaft that is highly active

during hard hammer percussion and that would favor thumb opposition (Marzke, 2013;

Marzke, Toth, Schick, & Reece, 1998).

5.1 Fossil hominin MC1 shape

The general scientific consensus in recent years is that *H. neanderthalensis* had a

hand morphology and manipulative capabilities that were very similar to those of

humans, challenging the previously held beliefs that *H. neanderthalensis* lacked the

derived adaptations for advanced and precise human-like tool use (Karakostis et al.,

2017; Karakostis, Hotz, Tourloukis, & Harvati, 2018; Niewoehner, 2001, 2006; Tocheri

et al., 2008; Trinkaus & Villemeur, 1991). The results align well with this consensus,

with the *H. neanderthalensis* specimen showing several similarities with the modern

humans. The described morphology is one of a flatter (PC2) and larger (PC1) distal

articular surface, bigger epicondyles at the distal head (PC1) and a flatter proximal

articular surface (PC2). However, *H. neanderthalensis* also differs in exhibiting a

particularly robust MC1 with strongly marked muscular insertions.

Previous analysis on the thumb morphology of *Homo naledi* fossils has indicated that

it has derived characteristics compatible with forceful precision grip and human-like

manipulative abilities (Berger et al., 2015; Kivell, 2015). Such characteristics include

a well-developed crest for the opponens pollicis insertion and flat distal/proximal

articular surfaces (Kivell et al., 2015; Galletta et al. 2019). The results generally agree

with these observations and conclusions, whilst also presenting some potentially new

insights. The morphology of the *H. naledi* sample had a human-like robustness of the

shaft (PC1), suggesting that the MC1 was adapted to sustain high loads, such as those

experienced during forceful tool use. This suggests that *H. naledi* was potentially

capable of a degree of advanced manipulation, such as forceful precision and power

grasping. These findings are therefore consistent with previous functional

interpretations of *H. naledi* thumb morphology (Galletta et al. 2019). Whilst the

evidence suggests that *H. naledi* was almost certainly able to perform an certain

degree of advanced manipulation, and was likely a tool-user, it also suggests that it

had not yet showed the full repertoire of manipulative adaptations exhibited by humans

and *H. neanderthalensis* (Berger et al., 2015; Kivell et al., 2015; Galletta et al., 2019).

Previous analysis of *A. sediba* hand morphology has found that it possessed a number

of advanced *Homo*-like features, such as a longer thumb relative to shorter fingers,

that potentially indicate advanced manipulative capabilities, while retaining primitive

traits as a gracile MC1, similar to those of other australopithecines (Kivell et al., 2011).

Recent 3DGM studies that have analyzed the MC1 in particular have come to the

conclusion that if *A. sediba* was indeed utilizing tools, as some hand proportion and

trabecular evidence suggests (Kivell et al., 2011; Skinner et al., 2015), then it was

doing so in a way that differed from that of early *Homo* and modern humans (Marchi

et al., 2017; Galletta et al., 2019). This conclusion was reached due to aspects of their

MC1 morphology that were deemed inconsistent with the ability to employ forceful

precision grips, namely a gracile MC1 shaft, more curved proximal articular surface

and smaller radial and ulnar epicondyles. These morphologies suggest that the range

of motion would not have been great enough at the TMCJ to facilitate the necessary

abduction-adduction for thumb opposition and pad-pad precision grips. The results

agree with this consensus, with the *A. sediba* sample presenting a gracile shaft (PC1),

and smaller epicondyles at the distal head. These morphologies suggest that the *A.*

sediba MC1 did not have the strength or stability to withstand the forces involved with

precision grips, nor the range of motion at the TMCJ to facilitate them. Overall the

results therefore align with previous research, in the sense that they present *A. sediba*

as having a patchwork of primitive and derived characteristics, a few of which are

indicators of an ability to use tools, but most of which suggest that this ability was

incipient and certainly not comparable to the forceful precision grip abilities of humans

and *H. neanderthalensis.*

6. Conclusion

The aim of this study was to quantify the 3D morphology of the first metacarpal in

extant African hominoids, in order to facilitate a more informed functional interpretation

of fossil hominin morphology. The results are in general agreement with previous

studies on the morphology of the MC1 in extant and extinct hominids and the

inferences made by them. This study found that the human MC1 is characterized by

a distinct suite of traits, not present to the same extent in non-human great apes, that

are consistent with an ability to use forceful precision and power grips; namely flatter

proximal and distal ends, larger epicondyles at the distal head and a more robust shaft.

This morphology was also found to align very closely with that of the *H.*

neanderthalensis sample, supporting all the evidence that indicates that Neanderthals

were functionally capable of utilizing tools in the same way as modern humans.

Analysis of the *H. naledi* specimen suggested that it had a number of human-like

adaptations consistent with an ability to employ advanced manipulation and was

therefore likely able to use stone tools in a similar way to humans. The *A. sediba* fossil

presented a number of derived MC1 features that indicate a degree of dexterity, but

also several traits which were more similar to the African apes (i.e., probably primitive

traits). Overall the results obtained both aligned with and added to past functional

interpretations of hominin morphology, thereby reinforcing the validity of 3DGM as a

method of quantifying MC1 morphology and providing a deeper insight into the

function and structure of the thumb in both extant hominids and fossil hominins.

Acknowledgments

We are grateful to the following curators and institutions for the access of the ape

specimens: Emmanuel Gilissen (AfricaMuseum), Anneke H. van Heteren and Michael

Hiermeier (Zoologische Staatssammlung München), Javier Quesada (Museu de

Ciències Naturals de Barcelona), José Miguel Carretero (Universidad de Burgos). AB

was partially funded by a Becas Chile scholarship, whilst TP was funded by the

Leverhulme Trust Early Career Fellowship, ECF-2018-264. This study was funded by

the research projects AGAUR 2017 SGR 1040 and MICINN-FEDER PGC2018-

093925-B-C32.

Author contributions

TAP and AB designed the study. AB and CL generated the 3D models. JM and TAP

analyzed the data. TAP carried out the data visualizations. All authors interpreted the

data and wrote the manuscript.

ORCID

Thomas A. Püschel https://orcid.org/0000-0002-2231-2297

Ana Bucchi https://orcid.org/0000-0002-1247-230X

Carlos Lorenzo https://orcid.org/0000-0001-5706-293X

References

Adams, D. C., Collyer, M. L., & Kaliontzopoulou, A. (2019). *Geomorph: Software for geometric morphometric analysis. R package version 3.1.0.*

Alba, D. M., Moyà-Solà, S., & Köhler, M. (2003). Morphological affinities of the Australopithecus afarensis hand on the basis of manual proportions and relative thumb length. *Journal of Human Evolution*, *44*(2), 225–254. https://doi.org/10.1016/S0047-2484(02)00207-5

Almécija, S., & Alba, D. M. (2014). On manual proportions and pad-to-pad precision grasping in Australopithecus afarensis. *Journal of Human Evolution*, *73*, 88–92. https://doi.org/10.1016/j.jhevol.2014.02.006

Almécija, S., Moyà-Solà, S., & Alba, D. M. (2010). Early origin for human-like precision grasping: A comparative study of pollical distal phalanges in fossil hominins. *PLOS ONE*, *5*(7), e11727. https://doi.org/10.1371/journal.pone.0011727

Almécija, S., Smaers, J. B., & Jungers, W. L. (2015). The evolution of human and ape hand proportions. *Nature Communications*, *6*, 1–11. https://doi.org/10.1038/ncomms8717

Aubriot, J. H., & Tubiana, R. (1981). The metacarpophalangeal joint of the thumb. *The Hand*, *1*, 184–187.

Barmakian, J. T. (1992). Anatomy of the joints of the thumb. *Hand Clinics*, *8*(4), 683–691.

Berger, L. R., Hawks, J., de Ruiter, D. J., Churchill, S. E., Schmid, P., Delezene, L. K., … Zipfel, B. (2015). Homo naledi, a new species of the genus Homo from the Dinaledi Chamber, South Africa. *ELife*, *4*(September2015), 1–35. https://doi.org/10.7554/eLife.09560

Berger, L. R., Ruiter, D. J. de, Churchill, S. E., Schmid, P., Carlson, K. J., Dirks, P. H. G. M., & Kibii, J. M. (2010). Australopithecus sediba: A new species of Homo-like

australopith from South Africa. *Science*, *328*(5975), 195–204.

https://doi.org/10.1126/science.1184944

Boesch, C., & Boesch, H. (1990). Tool use and tool making in wild chimpanzees. *Folia*

Primatologica, *54*(1–2), 86–99.

Bookstein, F. L. (1991). *Morphometric Tools for Landmark Data: Geometry and Biology*.

Cambridge: Cambridge University Press.

Bookstein, F. L. (1997). Landmark methods for forms without landmarks: morphometrics of

group differences in outline shape. *Medical Image Analysis*, *1*(3), 225–243.

https://doi.org/10.1016/S1361-8415(97)85012-8

Breuer, T., Ndoundou-Hockemba, M., & Fishlock, V. (2005). First observation of tool use in

wild gorillas. *PLoS Biology*, *3*(11), e380.

Bucchi, A., Luengo, J., Fuentes, R., Arellano-Villalón, M., Lorenzo, C. (2020).

Recommendations for improving photo quality in close range photogrammetry,

exemplified in hand bones of chimpanzees and gorillas. *International Journal of*

Morphology, *38*(2), 348–355. https://doi.org/10.4067/S0717-95022020000200348

Casillas Garcia, J. A., & Alvarez, G. A. (2005). *La Actuacion Arqueologica en el "Solar de*

Caballeria" y el Convento de San Pablo de Burgos. Burgos: Ayuntamiento de

Burgos, Instituto Municipal de Cultura.

Dirks, P. H., Roberts, E. M., Hilbert-Wolf, H., Kramers, J. D., Hawks, J., Dosseto, A., …

Berger, L. R. (2017). The age of homo naledi and associated sediments in the rising

star cave, South Africa. *ELife*, *6*. https://doi.org/10.7554/eLife.24231

Feix, T., Kivell, T. L., Pouydebat, E., & Dollar, A. M. (2015). Estimating thumb-index finger

precision grip and manipulation potential in extant and fossil primates. *Journal of the*

Royal Society, Interface, *12*(106), 20150176–. https://doi.org/10.1098/rsif.2015.0176

Galletta, L., Stephens, N. B., Bardo, A., Kivell, T. L., & Marchi, D. (2019). Three-

dimensional geometric morphometric analysis of the first metacarpal distal articular

surface in humans, great apes and fossil hominins. *Journal of Human Evolution*, *132*,

119–136. https://doi.org/10.1016/j.jhevol.2019.04.008

Giacomini, G., Scaravelli, D., Herrel, A., Veneziano, A., Russo, D., Brown, R. P., & Meloro,

C. (2019). 3D photogrammetry of bat skulls: Perspectives for macro-evolutionary

analyses. *Evolutionary Biology*, 46, 249–259. https://doi.org/10.1007/s11692-019-

09478-6

Gommery, D., & Senut, B. (2006). La phalange distale du pouce d'Orrorin tugenensis

(Miocene supérieur du Kenya). *Geobios*, *39*(3), 372–384.

Green, D. J., & Gordon, A. D. (2008). Metacarpal proportions in Australopithecus africanus.

Journal of Human Evolution, *54*(5), 705–719.

https://doi.org/10.1016/j.jhevol.2007.10.007

Guérin, G., Frouin, M., Talamo, S., Aldeias, V., Bruxelles, L., Chiotti, L., … Jain, M. (2015).

A multi-method luminescence dating of the Palaeolithic sequence of La Ferrassie

based on new excavations adjacent to the La Ferrassie 1 and 2 skeletons. *Journal of*

Archaeological Science, *58*, 147–166.

Gunz, P., Mitteroecker, P., & Bookstein, F. L. (2005). Semilandmarks in three dimensions. In

D.E. Slice (Ed.), *Modern morphometrics in physical anthropology* (pp. 73–98).

Boston, MA: Springer US.

Hamrick, M. W. (1996). Articular size and curvature as determinants of carpal joint mobility

and stability in strepsirhine primates. *Journal of Morphology*, *230*(2), 113–127.

Holliday, T. W., Churchill, S. E., Carlson, K. J., DeSilva, J. M., Schmid, P., Walker, C. S., &

Berger, L. R. (2018). Body size and proportions of Australopithecus sediba.

PaleoAnthropology, (January), 406–422. https://doi.org/10.4207/PA.2018.ART118

Imaeda, T., An, K. N., & Cooney 3rd, W. P. (1992). Functional anatomy and biomechanics

of the thumb. *Hand Clinics*, *8*(1), 9–15.

Karakostis, F. A., Hotz, G., Scherf, H., Wahl, J., & Harvati, K. (2017). Occupational manual

activity is reflected on the patterns among hand entheses. *American Journal of*

Physical Anthropology, *164*(1), 30–40. https://doi.org/10.1002/ajpa.23253

Karakostis, F. A., Hotz, G., Tourloukis, V., & Harvati, K. (2018). Evidence for precision

grasping in Neandertal daily activities. *Science Advances*, *4*(9), eaat2369.

https://doi.org/10.1126/sciadv.aat2369

Key, A. J. (2016). Manual loading distribution during carrying behaviors: Implications for

the evolution of the hominin hand. *PloS One*, *11*(10), e0163801.

Key, A. J. M., & Dunmore, C. J. (2015). The evolution of the hominin thumb and the

influence exerted by the non-dominant hand during stone tool production. *Journal of*

Human Evolution, *78*, 60–69. https://doi.org/10.1016/j.jhevol.2014.08.006

Kivell, T. L., Kibii, J. M., Churchill, S. E., Schmid, P., & Berger, L. R. (2011).

Australopithecus sediba Hand Demonstrates Mosaic Evolution of Locomotor and

Manipulative Abilities. *Science*, *333*(6048), 1411–1417.

https://doi.org/10.1126/science.1202625

Kivell, T.L. (2015). Evidence in Hand: recent discoveries and the early evolution of human

manual manipulation. *Philosophical Transactions of the Royal Society*, *370*.

Klingenberg, C. P., & McIntyre, G. S. (1998). Geometric morphometrics of developmental

instability: Analyzing patterns of fluctuating asymmetry with Procrustes methods.

Evolution, *52*(5), 1363–1375. https://doi.org/10.2307/2411306

Kuhn, M., & Johnson, K. (2013). *Applied Predictive Modeling*. New York: Springer.

https://doi.org/10.1007/978-1-4614-6849-3_11

Lewis, O. J. (1977). Joint remodelling and the evolution of the human hand. *Journal of*
Anatomy.

Marchi, D., Proctor, D. J., Huston, E., Nicholas, C. L., & Fischer, F. (2017). Morphological
correlates of the first metacarpal proximal articular surface with manipulative
capabilities in apes, humans and South African early hominins. *Comptes Rendus*
Palevol, *16*(5), 645–654. https://doi.org/10.1016/j.crpv.2016.09.002

Marzke, M. W. (2013). Tool making, hand morphology and fossil hominins. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *368*(1630), 20120414.
https://doi.org/10.1098/rstb.2012.0414

Marzke, M. W., & Marzke, R. F. (2000). Evolution of the human hand: approaches to
acquiring, analysing and interpreting the anatomical evidence. *The Journal of*
Anatomy, *197*(1), 121–140.

Marzke, M. W., Tocheri, M. W., Steinberg, B., Femiani, J. D., Reece, S. P., Linscheid, R. L.,
… Marzke, R. F. (2010). Comparative 3D quantitative analyses of trapeziometacarpal
joint surface curvatures among living catarrhines and fossil hominins. *American*
Journal of Physical Anthropology, *141*(1), 38–51. https://doi.org/10.1002/ajpa.21112

Marzke, M. W., Wullstein, K. L., & Viegas, S. F. (1992). Evolution of the power ("squeeze")
grip and its morphological correlates in hominids. *American Journal of Physical*
Anthropology, *89*(3), 283–298. https://doi.org/10.1002/ajpa.1330890303

Marzke, M. W., Marchant, L. F., McGrew, W. C., & Reece, S. P. (2015). Grips and hand
movements of chimpanzees during feeding in Mahale Mountains National Park,
Tanzania. *American Journal of Physical Anthropology*, *156*(3), 317–326.

Marzke, M. W, Toth, N., Schick, K., & Reece, S. (1998). EMG Study of Hand Muscle
Recruitment During Hard Hammer Percussion Manufacture of Oldowan Tools.
American Journal of Physical Anthropology, 105(3), 315-32.

Marzke, M. W., & Wullstein, K. L. (1996). Chimpanzee and human grips: A new

classification with a focus on evolutionary morphology. *International Journal of*

Primatology, *17*(1), 117–139. https://doi.org/10.1007/BF02696162

Napier, J. R. (1955). The form and function of the carpo-metacarpal joint of the thumb.

Journal of Anatomy, *89*(3), 362–369.

Napier, J. R. (1956). The prehensile movements of the human hand. *The Journal of Bone and*

Joint Surgery. British Volume, *38*(4), 902–913.

Napier, J. R. (1960). Studies of the hands of living primates. *Proceedings of the Zoological*

Society of London, *134*(4), 647–657.

Niewoehner, W. A. (2001). Behavioral inferences from the Skhul/Qafzeh early modern

human hand remains. *Proceedings of the National Academy of Sciences*, *98*(6), 2979–

2984. https://doi.org/10.1073/pnas.041588898

Niewoehner, W. A. (2006). Neanderthal hands in their proper perspective. In J-J. Hublian, K.

Harvati & T. Harrison (Eds.), Neanderthals *revisited: New approaches and*

perspectives (pp. 157–190). Dordrecht: Springer.

Püschel, T. A., Marcé-Nogué, J., Chamberlain, A. T., Yoxall, A., & Sellers, W. I. (2020). The

biomechanical importance of the scaphoid-centrale fusion during simulated knuckle-

walking and its implications for human locomotor evolution. *Scientific Reports*, *10*(1),

1–7. https://doi.org/10.1038/s41598-020-60590-6

Püschel, T. A., Marcé-Nogué, J., Gladman, J., Patel, B. A., Almécija, S., & Sellers, W. I.

(2020). Getting its feet on the ground: Elucidating Paralouatta's semi-terrestriality

using the virtual morpho-functional toolbox. *Frontiers in Earth Science*, 8, 79.

https://doi.org/10.3389/feart.2020.00079

Püschel, T. A., Marcé-Nogué, J., Gladman, J. T., Bobe, R., & Sellers, W. I. (2018). Inferring

locomotor behaviours in Miocene New World monkeys using finite element analysis,

geometric morphometrics and machine-learning classification techniques applied to

talar morphology. *Journal of The Royal Society Interface*, *15*(146), 20180520.

https://doi.org/10.1098/rsif.2018.0520

R Core Team. (2019). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-

project.org/.

Richmond, B. G., & Strait, D. S. (2000). Evidence that humans evolved from a knuckle-

walking ancestor. *Nature*, *404*(6776), 382–385. https://doi.org/10.1038/35006045

Rightmire, G. P. (1972). Analysis Metacarpal Swartkrans. *Science*, *176*, 159–161.

Rolian, C., Lieberman, D. E., & Zermeno, J. P. (2011). Hand biomechanics during simulated

stone tool use. *Journal of Human Evolution*, *61*(1), 26–41.

https://doi.org/10.1016/j.jhevol.2011.01.008

Sanz, C., Call, J., & Morgan, D. (2009). Design complexity in termite-fishing tools of

chimpanzees (Pan troglodytes). *Biology Letters*, *5*(3), 293–296.

Sanz, C. M., & Morgan, D. B. (2007). Chimpanzee tool technology in the Goualougo

Triangle, Republic of Congo. *Journal of Human Evolution*, *52*(4), 420–433.

Skinner, M. M., Stephens, N. B., Tsegai, Z. J., Foote, A. C., Nguyen, N. H., Gross, T., …

Kivell, T. L. (2015). Human-like hand use in Australopithecus africanus. *Science*,

347(6220), 395–399. https://doi.org/10.1126/science.1261735

Susman, R. L. (1998). Hand function and tool behavior in early hominids. *Journal of Human

Evolution*, *35*(1), 23–46. https://doi.org/10.1006/jhev.1998.0220

Tocheri, M. W., Orr, C. M., Jacofsky, M. C., & Marzke, M. W. (2008). The evolutionary

history of the hominin hand since the last common ancestor of Pan and Homo.

Journal of Anatomy, *212*(4), 544–562. https://doi.org/10.1111/j.1469-

7580.2008.00865.x

Trinkaus, E., & Villemeur, I. (1991). Mechanical advantages of the Neandertal thumb in

flexion: A test of an hypothesis. *American Journal of Physical Anthropology*, *84*(3),

249–260. https://doi.org/10.1002/ajpa.1330840303

Tuttle, R. H. (1969). Quantitative and functional studies on the hands of the Anthropoidea. I.

The Hominoidea. *Journal of Morphology*, *128*(3), 309–363.

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.). New

York: Springer-Verlag.

Wiley, D. F., Amenta, N., Alcantara, D. A., Ghost, D., Kil, Y. J., Delson, E., … Hamann, B.

(2005). Evolutionary morphing. *EEE Visualization, 2005. VIS 05*.

Zelditch, M. L., Sheets, H. D., & Fink, W. L. (2003). The ontogenetic dynamics of shape

disparity. *Paleobiology*, *29*(1), 139–156.

Anterior   Posterior   Medial   Lateral