Shared Characteristics of Intrinsic Connectivity Networks Underlying Interoceptive Awareness and Empathy

Stoica, T.*[1], Depue, B.E. [1,2]

[1]University of Louisville, Department of Psychological and Brain Sciences, University of Louisville, Louisville, KY

[2]University of Louisville, Department of Anatomical Sciences and Neurobiology, University of Louisville, Louisville, KY

*Correspondence:
Teodora Stoica
teodora.stoica@lousville.edu

Keywords: Empathy, Affective, Cognitive, Interoceptive Awareness, fMRI, resting state, network connectivity, network variability

Abstract

Awareness of internal bodily sensations (interoception, IA) and its connection to complex socioemotional phenomena like empathy have been postulated, yet its neural basis remains poorly understood. The present fMRI study employs independent component analysis (ICA) to investigate whether the Cognitive or Affective facets of empathy and IA share resting state network connectivity and/or variability (SD_{BOLD}). Healthy participants viewed an abstract movie demonstrated to evoke strong connectivity in resting state brain networks (*InScapes*), and resultant connectivity and variability data was correlated with self-reported empathy and IA questionnaires. We demonstrate a bidirectional behavioral and neurobiological relationship between empathy and IA, depending on the type of empathy interrogated: Affective empathy and IA share both connectivity and variability, while Cognitive empathy and IA only share variability. Specifically, increased connectivity in the right inferior frontal operculum (rIFO) of a larger attention network was associated with increased vicarious experience but decreased awareness of inner body sensations. Furthermore, increased variability between brain regions of an interoceptive network was related to increased sensitivity to internal sensations along with discomfort alleviation arising from witnessing another's distress. Finally, increased variability between brain regions subserving a mentalizing network related to not only an improved ability to take someone's perspective, but also a better sense of mind-body interconnectedness. Overall, these findings suggest that the awareness of one's own internal body changes (IA) is related to the ability to feel and understand another's emotional state (empathy) and critically, that this relationship is not task-dependent, but is reflected in the brain's resting state neuroarchitecture. Methodologically, this work highlights the importance of utilizing network variability as a complementary window alongside functional connectivity to better understand neurological phenomena. Our results may be beneficial in aiding diagnosis in clinical populations such as autism spectrum disorder, where participants may be unable to complete tasks or questionnaires due to the severity of their socioemotional symptoms.

Words: 5161; Figures: 4; Tables: 3

Interoceptive Awareness and Empathy Share Intrinsic Network Characteristics

1 Introduction

Internal body signals relative to emotion processing has been a topic of long-standing interest (Gurney 1884; Strack, Martin, and Stepper 1988), with more recent evidence highlighting an intriguing bidirectional relationship between sensations that arise internally and emotional phenomena (Lane 2008; Craig 2009; Cameron 2001; Damasio 2005). A proposed biological basis that may clarify this interplay is interoception, namely – the afferent processing of internal bodily signals that arise from visceral organs (Craig 2009; Cameron 2001; Wiens 2005; Johnson 2001; Cacioppo et al. 2000). Indeed, neuroimaging findings corroborate a substantial overlap between the neural substrates of emotion and interoception (Critchley and Garfinkel 2017; Adolfi et al. 2017), supporting the idea that the central monitoring and representation of internal bodily signals plays a fundamental role in emotion. However, the relationship between the emotions of *another* individual and one's own body remains relatively unexplored.

Consider the ubiquitous scenario of listening to a close friend share a difficult moment. In order to accurately grasp the story from their perspective, you must map their internal sensations onto your emotional landscape, thereby creating a bridge towards proper understanding. This process hinges on two crucial socioemotional constructs: interoceptive awareness (IA) and empathy. IA represents the ability to accurately and consciously perceive changes in your internal bodily signals (Khalsa et al. 2018), while empathy refers to both vicariously experiencing and understanding the mental state of your friend (Davis 1980). Thus, empathy can be further fractioned into two interrelated facets: Affective and Cognitive empathy. Affective empathy is conceptualized as the automatic process of vicariously experiencing the emotional state of another (Baron-Cohen and Wheelwright 2004; Davis 1980), while Cognitive empathy describes the individual's ability to accurately imagine another person's perspective (Davis, 1980; Decety & Jackson, 2004; Lawrence et al., 2006). Hence, in the context of the anecdote above, genuine understanding arises when the listener is aware of their own internal bodily sensations while taking into account the speaker's feelings and mental state. However, despite their importance in harmonious socioemotional exchange, it is currently unknown which specific facets of empathy dovetail with IA.

Neuroimaging research demonstrates that the neural substrates of empathy consistently overlap with the cortical regions involved in self-experience (Iacoboni 2005; Jackson, Meltzoff, and Decety 2005; Keysers et al. 2004; Wicker et al. 2003). For instance, during the experience of various aversive taste stimuli, activation of the anterior insula (AI) and inferior frontal operculum (IFO) were observed in both the observer and experiencer (Jabbi, Swart, and Keysers 2007). Similarly, observing others' pain has been found to robustly activate the AI and anterior cingulate cortex (ACC) (Lieberman and Eisenberger 2009; Jackson, Meltzoff, and Decety 2005; Singer et al. 2004), regions associated with one's own pain. One popular interpretation of such 'shared representation' between self and other's experience (Decety and Sommerville 2003) posits that the brain represents others' experiences in terms of the experiences of the self. To wit, one's own internal state serves as a blueprint for understanding the experiences of others (Iacoboni 2009; Singer and Lamm 2009; Rizzolatti, Fogassi, and Gallese 2006). Although this view has been broadly (and often implicitly) accepted, the neurobiological details of this putative self-referential process for understanding others remain largely unclear.

This lacuna may exist due to the interacting, but only partially overlapping neural bases of empathy's two facets (Fan et al. 2011). Affective empathy primarily elicits activations from regions implicated in rapid and prioritized processing of emotion signals, including: the amygdala, hypothalamus, orbitofrontal cortex (OFC) and AI (Decety et al. 2013). By comparison, Cognitive empathy, which shares similar neural networks with perspective-taking and mentalizing (Pardini

and Nichelli 2009) additionally involves the superior temporal sulcus (STS), temporoparietal
junction (TPJ), fusiform gyrus (FG), and medial prefrontal cortex (mPFC; (Saxe and Powell
2006)). To date, only one recent meta-analysis investigated convergent areas of activation between
interoception, emotion and social processing (Adolfi et al. 2017). The results for the three domains
converged in the AI, amygdala, right inferior frontal gyrus (rIFG), basal ganglia and medial
anterior temporal lobe (mATL), ascribing particular importance to the fronto-temporo-insular
nodes (Adolfi et al. 2017). The authors conclude co-activation of these regions may result in an
evaluative association of the internal milieu, and in combination with external cues, leads to
complex social behavior (i.e. empathy) (Adolfi et al. 2017).

Although this activation-based analysis revealed a number of regions known to play a role
within a distributed socioemotional network, scant functional connectivity data exists directly
addressing how these regions communicate. Indeed, only one study investigating empathy deficits
in a patient with depersonalization disorder (body self-awareness disruption) employed graph-
theory analyses, and demonstrated impaired Affective empathy and IA network measurements in
the AI, ACC and somatosensory cortex (Sedeño et al. 2014). Although germane, the study only
supports an association between these domains during active, task-relevant network
configurations. Thus far, no evidence exists regarding this interrelation in a healthy sample during
resting state (rsfMRI), despite studies showing that individual differences in complex social
processing (i.e. empathy) are reflected in the brain's intrinsic connectivity networks (ICNs) (Tavor
et al. 2016; Smith et al. 2009; Bilevicius et al. 2018; Christov-Moore et al. 2020; Cox et al. 2012).

In addition to connectivity, ICN variability is an often-discounted neuroimaging measurement
that may offer complementary information regarding network function and organization. Recent
theories consider high variability necessary for the neural system's adaptability, efficiency and
cognitive performance (Dai et al. 2016; Garrett et al. 2010; Garrett, Kovacevic, et al. 2013;
McIntosh, Kovacevic, and Itier 2008; Vakorin et al. 2011; Vakorin, Lippé, and McIntosh 2011).
Specifically, according to the coordination dynamics theory, networks demonstrating increased
variability can flexibly shift through integrative and segregative configurations, maintaining the
neural system in balance (Tognoli and Kelso 2014). Thus far, network variability has been
predominantly used in clinical populations to investigate alteration in the organization of brain
networks (Scarapicchia et al. 2018; Zhang et al. 2020; Good et al. 2020; Kumral et al. 2020) and
in healthy populations to investigate brain maturation trajectories (Nomi et al. 2017). To the best
of knowledge however, no study has explored the intrinsic network characteristics of connectivity
and variability that putatively underlie the socioemotional constructs of empathy and IA in a
healthy sample.

Therefore, the present study employs a data-driven approach to explore resting state
connectivity and variability related to brain networks underlying Cognitive, Affective empathy
and IA. Specifically, we aim to understand whether Cognitive and/or Affective empathy as
measured by self-report questionnaires share intrinsic connectivity and/or variability with IA
during the viewing of naturalistic stimuli in healthy adults. We hypothesize based on previous
literature that 1) Affective empathy will share network connectivity with IA within amygdala, AI
and rIFO, given their involvement in the processing of emotion experienced in oneself and
vicariously for others (Singer et al. 2004), while 2) Cognitive empathy will share network
variability with IA within the rTPJ and precuneus, as these regions are posited to underlie explicit
mentalizing (Naughtin et al. 2017; Kovács et al. 2014; Hyde, Aparicio Betancourt, and Simon
2015; Bardi et al. 2017).

2 Materials and Methods

2.1 <u>Participants and Procedure</u>
26 healthy young adults (m=21.85 y.o./16 female) without a reported history of neurological
or psychiatric disorders were recruited for this study **(Table 1).** All participants were right-handed
and had normal or corrected-to-normal vision and hearing. Participants were recruited through on-
campus flyers. All participants were paid $20 for their participation. Experimental protocols were
approved by University of Louisville's Institutional Review Board prior to data collection, and
written informed consent was obtained from each participant prior to experimental sessions. The
study took part on two separate days. On the first day, participants visited the lab to be briefed on
the MRI protocol, fill out consent forms and behavioral assessments. On the second day,
participants completed the rsfMRI scan at the University of Louisville, School of Medicine.
2.2 <u>Resting State Scan: Naturalistic Paradigm</u>
The participants watched a 7 min abstract, nonsocial movie titled *Inscapes* previously
demonstrated to evoke strong connectivity in networks that resemble rest more than those
exhibited during conventional movies (Vanderwal et al. 2015). The movie features a series of
technological-looking abstract shapes (**Figure 1**). Participants were told to keep their eyes open
and relax while watching and listening to the movie. The stimulus was displayed using E-prime
on an Invivo Esys LCD TV monitor at the back of the scanner bore, which was viewed by
participants through a mirror on the head-coil. The video is freely available for download from
<u>HeadSpace Studios</u>.
2.3 <u>Behavioral Assessments</u>
2.3.1   *Empathy Questionnaire – Interpersonal Reactivity Index (IRI)*
Affective and Cognitive empathy was assessed using the Interpersonal Reactivity Index (IRI)
(Davis 1983). The IRI consists of 28 items rated on a 5-point scale with the anchors: "does not
describe me well" to "describes me very well". The items are arranged into four subscales with
seven items. Each subscale measures a distinct component of empathy: empathic concern (EC)
(feelings of compassion and concern for others); personal distress (PD) (feelings of anxiety and
discomfort that result from observing another person's negative experience); perspective taking
(PT) (the ability to adopt the perspectives of other people and see things from their point of view);
and fantasy subscale (FS) (the tendency to identify with characters in movies, books, or other
fictional situations) (Davis 1983). Affective empathy, the ability to infer an agent's feelings or
emotions, was derived from summing the EC and PD subscales. Cognitive empathy, the ability to
infer an agent's beliefs or thoughts, was derived from summing the FS and PT subscales. Total
empathy was derived from aggregating Affective and Cognitive empathy scores. All scores were
standardized by applying a z-score transformation, and later compared with the MAIA assessment
and its subscales (see below) in subsequent analyses.
2.3.2   *Multidimensional Assessment of Interoceptive Awareness (MAIA)*
The Multidimensional Assessment of Interoceptive Awareness (MAIA) is a 32-item instrument
assessing IA: "the conscious perception of sensations from inside the body that create the sense of
the physiological condition of the body, such as heart beat, respiration, satiety, and the autonomic
nervous system sensations related to emotions" (Mehling et al. 2012). Each statement is rated from
0 (never) to 5 (always) in terms of how often it applies to the participant generally in daily life.
The statements are then separated into 8 subscales: Noticing, Non-Distracting, Not-Worrying,
Attention-Regulation, Emotional Awareness, Self-Regulation, Body Listening and Trusting,
which are in turn aggregated into 5 overall scales used in the present study: Awareness of Body
Sensations (Noticing); Emotional Reaction and Attentional Response to Sensations (Not-
Distracting, Not-Worrying); Capacity to Regulate Attention (Attention Regulation), Awareness of

Mind-Body Integration (Emotional Awareness, Self-Regulation, Body Listening) and Trusting
Body Sensations (Trusting). A total Interoceptive Score (MAIA Total) was derived by summing
all the aggregate scales. All scores were standardized by applying a z-score transformation, and
later compared with the IRI and its subscales (see above) in subsequent analyses. Correlations
between behavioral measurements were conducted with the Statistical Package for Social Sciences
(Version 25.0.0; SPSS, INC.), and corrected for age, gender and multiple comparisons using false
discovery rate $p<0.05$ (FDR) (Benjamini and Hochberg 1995).
2.4 MRI Data Acquisition and Preprocessing
All structural MRI images were acquired using a Siemens 3-T Skyra MR scanner. A 20-
channel head coil was used for radiofrequency reception. Participants were given earplugs to
reduce scanner noise and were additionally given headphones to receive instructions. Foam
padding was added to limit motion if additional room remained within the head coil, and a piece
of folded tape was placed over the participant's forehead as a reminder to remain still throughout
the scan. Structural images were obtained via a T1-weighted magnetization-prepared rapid
gradient-echo sequence (MPRAGE) in 208 sagittal slices. Imaging parameters were as follows:
echo time (TE) = 2.26 ms, repetition time (TR) = 1700 ms, flip angle = 9.0°, field of view (FoV)
= 204 mm, and voxel size = 0.8 x 0.8 x 0.8 mm. Scan parameters were consistent for all imaging
sessions associated with this study. Functional blood oxygenation level-dependent (BOLD)
images were collected using multi-band acceleration factor of 3. Imaging parameters were as
follows: TE = 29 ms; TR = 2000 ms; flip angle = 62°; FoV = 250 mm; isotropic voxel size = 2.0
mm^3; 78 interleaved slices, GRAPPA on, Partial Fourier 7/8. Slices were oriented obliquely along
the AC–PC line.
All analyses were conducted using the CONN toolbox 19.c (Whitfield-Gabrieli and Nieto-
Castanon 2012) based on SPM12 (Penny et al. 2007) in the 2017 version of MATLAB. Spatial
preprocessing in the CONN toolbox included realignment, normalization and smoothing (6 mm
FWHM Gaussian filter) using SPM12 default parameter settings. Anatomical volumes were
segmented into gray matter, white matter and cerebrospinal fluid (CSF) areas, and the resulting
masks were eroded to minimize partial volume effects. Physiological and other sources of noise
were estimated and regressed out using CompCor (Behzadi et al. 2007), a method that performs
principal component analysis to estimate the physiological noise from white matter and
cerebrospinal fluid for each participant. Motion from the ART toolbox was also included as a
confound regressor. Next, a 0.01–0.10 Hz temporal band-pass filter standard for resting-state
connectivity analyses was applied to the time series (Nieto-Castanon 2015). In sum, detrending,
outlier censoring, motion regression, and CompCor correction were performed simultaneously in
a single first-level regression model, followed by band-pass filtering. These corrections yielded a
residual BOLD time course at each voxel that was used for subsequent analyses.
2.5 Neuroimaging Analysis
2.5.1 *Network Connectivity*
Group-level independent component analysis (ICA) using the CONN toolbox was conducted
to identify networks of functionally connected brain regions during resting state that may be
associated with the IRI and MAIA scores. This involved the application of the fastICA algorithm
to volumes concatenated across subject and resting state condition in order to identify independent
spatial components (ICs) and back-projection of these components to individual subjects, resulting
in maps of regression coefficients representing connectivity between the network and every voxel
in the brain (see Calhoun et al., 2001 for details). Forty ICs were identified using spatial overlap
of suprathreshold areas (Dice coefficient (Rombouts et al. 1998)), based on CONN's default

network atlas with ROIs characterizing an extended set of classical brain networks: Default Mode
Network (4 ROIs), SensoriMotor (2 ROIs), Visual (4 ROIs), Salience / Cingulo-Opercular (7
ROIs), DorsalAttention (4 ROIs), FrontoParietal / Central Executive (4 ROIs), Language (4 ROIs),
Cerebellar (2 ROIs) (all ROIs defined from CONN's ICA analyses of HCP dataset / 497 subjects)
(Nieto-Castanon 2014). We chose 40 ICs due to research suggesting ICA results are only affected
by the number of ICs when the number is smaller than the number of source signals (Ma et al.
2007), in addition to assuring coverage of a majority of signal variance. Noise components were
further identified through visual inspection by authors (TS and BD) (e.g., components largely
overlapping CSF), resulting in the exclusion of 4 out of 40 ICs from further consideration. Each
of the remaining 36 ICs was subsequently entered in multiple regressions with the IRI and MAIA
subscales, aggregate and Total scores. For each network, the resulting statistical maps were cluster
thresholded at $p \leq 0.05$, voxel thresholded at $p < 0.001$ (FDR-corrected) using Gaussian Random
Field Theory (Worsley et al. 1996), and corrected for age and gender. All coordinates reported
below refer to peak activations in anatomical MNI space.
2.5.2 *Network Variability*
In order to assess network variability and its relationship to either empathy or IA, we regressed
each IC's network variability (calculated in CONN as SD of each IC's BOLD timeseries: SD_{BOLD}
(Nieto-Castanon 2020)) with the IRI and MAIA subscales.
3   RESULTS
3.1 <u>Behavioral Results</u>
In order to establish the relationship between empathy and IA, all IRI and MAIA subscales,
aggregate and Total scores were correlated (**Table 2**).
3.1.2 Relationship between Affective Empathy and Interoceptive Awareness
Excluding same-subscale correlations, negative relationships were observed between
Affective empathy and the following MAIA subscales: Capacity to Regulate Attention ($r(26) = -$
0.83, $p < 0.01$), Trusting Body Sensations ($r(26) = -0.55$, $p < 0.01$) and MAIA Total ($r(26) = -0.64$,
$p < 0.01$). Similarly, we observed a negative relationship between the Personal Distress (PD)
subscale and Capacity to Regulate Attention ($r(26) = -0.74$, $p < 0.01$), Awareness of Mind Body
Integration ($r(26) = -0.49$, $p < 0.01$), Trusting Body Sensations ($r(26) = -0.53$, $p < 0.01$) and MAIA
Total ($r(26) = -0.66$, $p < 0.01$). Therefore, we report a negative relationship between Affective
empathy and IA (most influenced by the PD scale, since EC exhibited no significant relationship).
3.1.3 Relationship between Cognitive Empathy and Interoceptive Awareness
We observed a positive relationship between Cognitive Empathy and the Awareness of Mind
Body Integration subscale, $r(26) = 0.35$, $p = 0.06$, although it did not survive multiple comparison
correction. In addition, we observed a significant positive relationship between the IRI Perspective
Taking (PT) subscale and the MAIA Awareness of Mind Body Integration subscale, $r(26) = 0.41$,
$p < 0.01$. Therefore, we report a positive relationship between Cognitive empathy and IA (most
influenced by the PT subscale, since Fantasy exhibited no significant relationship).
Therefore, taken together, our behavioral results show a bidirectional relationship between
empathy and IA, depending on the facet of empathy interrogated and mainly driven by the
subscales of PD (Affective Empathy) and PT (Cognitive Empathy).
3.2 <u>Functional Connectivity Results</u>
3.2.1 Network Connectivity
We observed that within a network comprising right inferior frontal operculum (rIFO),
bilateral superior parietal lobule (SPL) and bilateral middle temporal gyrus (MTG), greater
connectivity in the rIFO was associated with greater overall empathy as measured by the IRI Total,

and with greater IRI Affective empathy. Conversely, lower connectivity in the rIFO was associated
with increased overall IA as measured by the MAIA Total and increased MAIA Capacity to
Regulate Attention (**Figure 2, Table 3**).
3.2.2 Network variability
Network variability analyses revealed that within a network comprising left IFO (L IFO),
Cerebellum, and rAI, IRI Personal Distress was negatively related to variability of the network
(T(22) = -1.04, *p*=0.02)), while conversely, MAIA Awareness of Body Sensations was positively
related (T(22) = 1.48, *p*=0.001)), (**Figure 3**).
Lastly, within a network comprising right Precuneus, rMTG, bilateral supramarginal gyrus
(SMG) and rIFO, IRI Perspective Taking (T(22) = 1.45, *p*=0.001) and MAIA Mind Body
Integration (T(22) = 2.52, *p*=0.01) were positively related to network variability (**Figure 4**).
4   Discussion
Interoceptive awareness and empathy are core social abilities that enable humans to create
meaningful social exchanges by accessing one's own internal state in order to understand the
experiences of others. Our resting state fMRI study employed Independent Component Analysis
(ICA) to investigate which empathy facet (Cognitive or Affective) shares network connectivity or
variability with interoceptive awareness (IA) while healthy adults viewed naturalistic stimuli. We
observed a bidirectional behavioral relationship between empathy and IA, whereby Affective
empathy and IA were negatively related, and Cognitive empathy and IA were positively related.
This bidirectional link also appeared in the neuroimaging results, such that Affective empathy and
IA were inversely related to increased connectivity within the rIFO, and also inversely related to
network variability; whereas Cognitive empathy and IA showed only a positive relationship with
network variability. Together, these results suggest a double disassociation between empathy and
IA depending on the type of empathy interrogated, which is reflected in the brain network's
intrinsic connectivity and variability patterns.
Behaviorally, we observed a negative relationship between the Personal Distress (PD) subscale
of the Affective empathy aggregate IRI scale and the total MAIA score, Capacity to Regulate
Attention and Trusting Body Sensations subscales. The Capacity to Regulate Attention subscale
pertains to various ways of controlling one's attention towards bodily sensations, as part of an
active regulatory process; while Trusting Body Sensations reflects the extent to which one views
awareness of bodily sensations as helpful for decision making (Mehling et al. 2012). The Empathic
Concern (EC) subscale of Affective empathy exhibited no significant relationship, implying that
PD is the dominant subscale of the Affective empathy aggregate scale when relating to the MAIA.
This distinction is important, considering that previous data suggests EC motivates individuals to
pay attention *to others'* emotions in order to try to comfort them, while conversely, PD drives
attention *away from others* in order to reduce the aversive effects for oneself, perhaps as a form of
emotion regulation (Zaki et al. 2014). Indeed, Decety & Jackson, 2004 proposed that PD may arise
from the failure of applying sufficient self-regulatory control over the shared emotional state. In
line with previous studies, we report an inverse relationship between PD and a an attention
regulation measure – MAIA's Capacity to Regulate Attention subscale. Together with the Trusting
Body Sensations subscale, our findings suggest the increased ability to regulate internal attention
and rely on this discrete information may be linked to a decrease in the discomfort experienced
while witnessing another's distress.
Furthermore, we found a positive relationship between the Perspective Taking (PT) subscale
of the Cognitive empathy aggregate IRI scale and the Awareness of Mind Body Integration of the
MAIA. This MAIA subscale represents the integration of several higher level cognitive processes

necessary for socially relevant goal-directed behavior (i.e. executive functions (Pribram 1973))
including: emotional awareness, self-regulation of emotions, and the ability to feel a sense of an
embodied self, that is – "a sense of the interconnectedness of mental, emotional, and physical
processes as opposed to a disembodied sense of alienation from one's body" (Mehling et al. 2012).
Thus, our results support previous findings suggesting Cognitive empathy, and in particular PT, is
related to a wide array of executive function skills such as working memory, inhibitory control
and cognitive flexibility (Yan et al. 2020; Aliakbari et al. 2013; Healey and Grossman 2018).
Taken together with the aforementioned negative relationship between PD and IA, these
behavioral results suggest a bidirectional association between empathy and IA, contingent on the
type of empathy interrogated. To wit, directing attention towards internal bodily sensations may
relieve vicarious emotional pain but flexibly employing cognitive-control skills may increase the
ability to take the perspective of another in pain.
Our connectivity results provide further support for this inverse relationship. Within a network
of brain regions previously shown to underlie attentional processing (superior parietal lobule
(SPL), medial temporal gyrus (MTG) and right inferior frontal operculum (rIFO) (Corbetta and
Shulman 2002; Wu et al. 2016; Perrett et al. 1992; Perrett, Rolls, and Caan 1982; Perrett et al.
1985), we observed that increased connectivity in the rIFO was associated with increased overall
empathy (total IRI score) and the Affective empathy aggregate scale on one hand, but reduced
overall IA (total MAIA score) and Capacity to Regulate Attention on the other hand. Previous
studies investigating both personal (Craig 2009; Johnson 2001; Damasio 2005; Critchley 2005;
Gray et al. 2007) and vicarious emotional experience (Singer et al. 2004; Jabbi, Swart, and Keysers
2007) show the consistent activation of the anterior insula (AI) and frontal operculum, and
therefore the IFO is thought of as a continuum between these two structures (Jabbi, Swart, and
Keysers 2007; Wicker et al. 2003). Because Affective empathy was driven by the PD subscale,
increased connectivity within the rIFO in the present study may relate to intensified personal
suffering from witnessing another's distress, but decreased awareness of one's own body
sensations, perhaps due to allocating attention externally (for example, away from oneself and
toward another's distress). Our findings reflect previous results showing neural overlap of
interoception and empathy in this region (Ernst et al. 2013; Adolfi et al. 2017), and extend these
findings by providing intrinsic connectivity evidence of a double dissociation between empathy
and IA.
Our network variability results offer a complementary perspective that further supports this
bidirectional relationship. We observed increased scores on the MAIA Awareness of Body
Sensations subscale and decreased scores on the IRI PD subscale was associated with increased
variability of brain regions previously shown to underlie processing and integration of visceral
information (i.e., Cerebellum, L IFO, L AI) (J. D. Schmahmann 2001; Baumann and Mattingley
2012; Schienle and Scharmüller 2013; Adamaszek et al. 2017; Bogg and Lasecki 2015; Terasawa,
Fukushima, and Umeda 2013). Despite the prevailing focus on the AI as a hub for processing body
sensations (Terasawa, Fukushima, and Umeda 2013; Kuehn et al. 2016; Pollatos, Gramann, and
Schandry 2007; Singer, Critchley, and Preuschoff 2009), additional brain regions are also
commonly implicated in interoceptive experience. For example, fMRI studies identify the
involvement of the IFO and cerebellum, reinforcing the notion that interoceptive processing (and
perhaps especially nociceptive information) may occur through multiple neural pathways (Rapps
et al. 2008; Peiffer et al. 2008; Garcia-Larrea 2012). In the same vein, observing distress in others
without actually experiencing it may rely on high-order cognitive functions to access minor
changes in physical state, as a tool to modulate negative stimulus input (Preckel, Kanske, and

Singer 2018). The implication of the cerebellum in a shared network underlying both PD and
interoceptive processing is not surprising, since the cerebellum serves as an integral node in the
distributed cortical–subcortical neural circuitry supporting an array of social cognitive operations
(Jeremy D. Schmahmann and Pandya 1995). Taken together, we suggest variability of this network
of brain regions underlying interoceptive experience may reflect increased flexibility of the
network to process internal sensations, but less adaptability to integrate vicarious emotions arising
from witnessing another's distress.
In addition, we observed a positive relationship between variability of both the IRI PT subscale
and the MAIA Awareness of Mind Body Integration subscale within a network of brain regions
previously associated with the process of mentalizing – the precuneus, rIFO, SMG and MTG (Mar
2011; Northoff et al. 2006; Schurz et al. 2014; Spreng, Mar, and Kim 2009; Vogeley et al. 2001).
Mentalizing, also referred to as Theory of Mind (TOM), signifies the ability to attribute mental
states to another individual, allowing the observer to predict intent and direct their behavior
appropriately (Frith, Morton, and Leslie 1991; Premack and Woodruff 1978) . Researchers agree
that mentalizing differs from the vicarious sharing of emotion in its psychological complexity,
combining observation, memory, knowledge, and reasoning to provide insight into the thoughts
and feelings of others (van der Heiden et al. 2013; Decety and Jackson 2004). Therefore, its
connection with MAIA's Awareness of Mind Body Integration scale is not surprising, since both
concepts require not only affective experience, but also comprehension and integration of that
particular state of mind within one's own emotional schema. We therefore suggest that increased
variability of brain regions underlying a mentalizing network may point to enhanced network
flexibility subserving not only a better ability to take another's perspective, but also an improved
sense of interconnectedness between one's own mind and body.
Lastly, our data shows an interesting convergence of empathy and IA within the IFO. Research
suggests the IFO serves as both a sensory-cognitive integration area and a control node of the
ventral attention network (Craig 2009; Corbetta and Shulman 2002), conjectured to maintain goal-
related information online until a decision is reached (Tops and De Jong 2006; Tops et al. 2010).
Moreover, recent evidence suggests a hemispheric specialization of the IFO related to
reactive/proactive goal maintenance (Tops et al. 2010). On one hand, the rIFO may facilitate
immediate somatosensory processing and attentional shifting whilst a response is ongoing
(reactive) (Tops et al. 2010; Hampshire et al. 2009; Higo et al. 2011; Nelson et al. 2010), through
its connections to rostral ACC, superior temporal gyrus (STG) and occipital cortex (Cauda et al.
2011). On the other hand, the lIFO may exert top-down control whilst preparing a response
(proactive) (Tops et al. 2010), through its connections to dorsolateral prefrontal cortex and bilateral
supplementary motor area (Cauda et al. 2011). Taking this evidence into consideration, we
speculate the positive association between connectivity within the rIFO and Affective empathy
indicates a propensity in the highly empathic individual to shift attention toward salient cues in
their environment (for example, another individual in distress). In contrast, the negative
relationship between connectivity within the rIFO and IA may indicate an inability to redirect
attention toward external salient cues, and therefore may lead to increased awareness of internal
sensations. Our network variability findings offer complementary evidence regarding the role of
the IFO in socioemotional processes. We show that increased network flexibility within an
interoceptive experience network (comprised of lIFO, L AI, cerebellum) is linked to increased
Awareness of Body Sensations as well as decreased PD. We suggest the lIFO plays a crucial part
in this network's ability to modulate attention from one's own internal sensations to discomfort
arising from witnessing another's distress, perhaps in an effort to plan an appropriate emotional

response. In the same vein, we show enhanced network flexibility within a mentalizing network
(comprised of rIFO, precuneus, SMG, MTG) is related to both better PT and increased Mind Body
Integration. These relationships may illustrate that heightened ability to determine intent and
integrate sensory information into one's own emotional schema could relate to flexibly shifting
attention towards the target of interest (either self or other). In sum, our data suggests the IFO
serves as an internal/external attention modulator and thus may play a critical role in switching
attention from one's own body sensations (IA) to another's (Affective and/or Cognitive empathy).
Our study's findings should be considered along with its limitations. The definition of network
variability has been inconsistent across previous studies, (e.g., amplitude, variance, standard
deviation, mean squared successive difference; for a review, see (Garrett, Samanez-Larkin, et al.
2013) with considerable range in the methodology used to derive them. Therefore, implementation
of network variability as a consistently used neuroimaging measure will require increased efforts
toward methodological standardization. It is also important to note that due to the nature of the
analyses used, the findings of this study do not represent causal relationships. That is, the results
represent a correlational relationship between a questionnaire-based measure of IA or empathy and
functional connectivity and/or variability. Our sample was unfortunately not large enough for a
gender-specific analysis, as evidence suggests there are differences in the capacity for empathy
between males and females (Christov-Moore et al. 2014). Future research should be conducted in
this regard. Similarly, in using an undergraduate sample, the generalizability of these findings is
limited.
In conclusion, the current research provides novel information about the relationship between
internal body sensation awareness and empathy. In contrast to previous studies which used task-
based fMRI to assess the neurobiology of these two constructs separately, we used a data-driven
resting state approach to test whether distinct empathy facets share network characteristics
(connectivity and/or variability) with IA. We demonstrate a bidirectional relationship between
empathy and IA, depending on the type of empathy investigated. Specifically, Affective empathy
and IA share network connectivity and variability, while Cognitive empathy and IA only share
variability. In regard to Affective empathy, increased vicarious emotional experience and
decreased IA were associated with increased connectivity within the rIFO of a larger attention
network; while increased IA and reduced personal discomfort arising from witnessing another's
distress were related to increased network flexibility within an internal sensation network.
Concerning Cognitive empathy, perspective-taking ability and a sense of mind-body
connectedness related to increased network flexibility between brain regions subserving a
mentalizing network. We also suggest the role of the IFO as an internal/external attention
modulator that may play a critical role in switching attention from one's own body sensations (IA)
to another's (Affective and/or Cognitive empathy). Overall, we show that the ability to feel and
understand another's emotional state is related to one's own awareness of internal body changes,
and that this relationship is not task-dependent, but is reflected in the brain's resting state
neuroarchitecture. Methodologically, this work highlights the importance of utilizing network
variability alongside functional connectivity as an important complementary route into
understanding neurological phenomena. Our results hold promise in aiding diagnosis of clinical
disorders characterized by IA and empathy deficits such as the autism spectrum disorders (ASD),
where participants may be unable to complete tasks or questionnaires due to the severity of their
socioemotional symptoms.
460
461

462
463
464
Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

Author Contributions

Teodora Stoica: Data Curation, Investigation, Project administration, Formal analysis,
Methodology, Visualization, Writing – original draft **Brendan Depue**: Resources, Supervision,
Writing – Review and editing
472
473 **Funding**
474
475 This research did not receive any specific grant from funding agencies in the public, commercial,
476 or not-for-profit sectors.
477
478 **Acknowledgements**
479
480 Many thanks to Dr. Nick Hindy for invaluable advice regarding writing.
481
482 **Data Availability Statement**

483 Raw data and behavioral variables of interest for the empirical results presented here can be found
484 on the Mendeley Data Repository at http://dx.doi.org/10.17632/fmch6sj4sw.1

References

Adamaszek, Michael, Federico D'Agata, Roberta Ferrucci, Christophe Habas, S Keulen, K C
    Kirkby, Maria Leggio, Peter Mariën, Marco Molinari, and E Moulton. 2017. "Consensus
    Paper: Cerebellum and Emotion." *The Cerebellum* 16 (2): 552–76.

Adolfi, Federico, Blas Couto, Fabian Richter, Jean Decety, Jessica Lopez, Mariano Sigman,
    Facundo Manes, and Agustín Ibáñez. 2017. "Convergence of Interoception, Emotion, and
    Social Cognition: A Twofold FMRI Meta-Analysis and Lesion Approach." *Cortex* 88
    (March): 124–42. https://doi.org/10.1016/j.cortex.2016.12.019.

Aliakbari, Mahnaz, Aliasghar Kaku Juibari, Fatemeh Amirabadi, Farhad Shaghaghi, Nushin
    Zare, and Faeze Khaleghi. 2013. "The Role of Theory of Mind and Executive Function in
    Predicting Empathy. [The Role of Theory of Mind and Executive Function in Predicting
    Empathy.]." *Advances in Cognitive Science*. Amirabadi, Fatemeh: f.amirabady@gmail.com:
    Institute for Cognitive Science Studies.

Bardi, Lara, Charlotte Desmet, Annabel Nijhof, Jan R Wiersema, and Marcel Brass. 2017.
    "Brain Activation for Spontaneous and Explicit False Belief Tasks Overlaps: New FMRI
    Evidence on Belief Processing and Violation of Expectation." *Social Cognitive and
    Affective Neuroscience* 12 (3): 391–400.

Baron-Cohen, Simon, and Sally Wheelwright. 2004. "The Empathy Quotient: An Investigation
    of Adults with Asperger Syndrome or High Functioning Autism, and Normal Sex
    Differences." *Journal of Autism and Developmental Disorders*.
    https://doi.org/10.1023/B:JADD.0000022607.19833.00.

Baumann, Oliver, and Jason B. Mattingley. 2012. "Functional Topography of Primary Emotion
    Processing in the Human Cerebellum." *NeuroImage* 61 (4): 805–11.
    https://doi.org/10.1016/j.neuroimage.2012.03.044.

Behzadi, Yashar, Khaled Restom, Joy Liau, and Thomas T. Liu. 2007. "A Component Based
    Noise Correction Method (CompCor) for BOLD and Perfusion Based FMRI." *NeuroImage*
    37 (1): 90–101. https://doi.org/10.1016/j.neuroimage.2007.04.042.

Benjamini, Y, and Y Hochberg. 1995. "Benjamini-1995.Pdf." *Journal of the Royal Statistical
    Society B*. https://doi.org/10.2307/2346101.

Bilevicius, Elena, Tiffany A. Kolesar, Stephen D. Smith, Paul D. Trapnell, and Jennifer
    Kornelsen. 2018. "Trait Emotional Empathy and Resting State Functional Connectivity in
    Default Mode, Salience, and Central Executive Networks." *Brain Sciences* 8 (7).
    https://doi.org/10.3390/brainsci8070128.

Bogg, Tim, and Leanne Lasecki. 2015. "Reliable Gains? Evidence for Substantially
    Underpowered Designs in Studies of Working Memory Training Transfer to Fluid
    Intelligence." *Frontiers in Psychology*. https://doi.org/10.3389/fpsyg.2015.00589.

Cacioppo, John T., Gary G. Berntson, John F. Sheridan, and Martha K. McClintock. 2000.
    "Multilevel Integrative Analyses of Human Behavior: Social Neuroscience and the
    Complementing Nature of Social and Biological Approaches." *Psychological Bulletin* 126
    (6): 829–43. https://doi.org/10.1037/0033-2909.126.6.829.

Calhoun, V. D., T. Adali, G. D. Pearlson, and J. J. Pekar. 2001. "A Method for Making Group
    Inferences from Functional MRI Data Using Independent Component Analysis." *Human
    Brain Mapping* 14 (3): 140–51. https://doi.org/10.1002/hbm.1048.

Cameron, O. G. 2001. "Interoception: The inside Story - A Model for Psychosomatic Processes."
    Psychosomatic Medicine 63 (5): 697–710. https://doi.org/10.1097/00006842-200109000-

00001.

Cauda, Franco, Federico D'Agata, Katiuscia Sacco, Sergio Duca, Giuliano Geminiani, and

Alessandro Vercelli. 2011. "Functional Connectivity of the Insula in the Resting Brain."

NeuroImage 55 (1): 8–23. https://doi.org/https://doi.org/10.1016/j.neuroimage.2010.11.049.

Christov-Moore, Leonardo, Nicco Reggente, Pamela K. Douglas, Jamie D. Feusner, and Marco

Iacoboni. 2020. "Predicting Empathy From Resting State Brain Connectivity: A

Multivariate Approach." *Frontiers in Integrative Neuroscience*.

https://doi.org/10.3389/fnint.2020.00003.

Christov-Moore, Leonardo, Elizabeth A. Simpson, Gino Coudé, Kristina Grigaityte, Marco

Iacoboni, and Pier Francesco Ferrari. 2014. "Empathy: Gender Effects in Brain and

Behavior." *Neuroscience and Biobehavioral Reviews* 46 (P4): 604–27.

https://doi.org/10.1016/j.neubiorev.2014.09.001.

Corbetta, Maurizio, and Gordon L. Shulman. 2002. "Control of Goal-Directed and Stimulus-

Driven Attention in the Brain." *Nature Reviews Neuroscience* 3 (3): 201–15.

https://doi.org/10.1038/nrn755.

Cox, Christine L., Lucina Q. Uddin, Adriana Di martino, F. Xavier Castellanos, Michael P.

Milham, and Clare Kelly. 2012. "The Balance between Feeling and Knowing: Affective and

Cognitive Empathy Are Reflected in the Brain's Intrinsic Functional Dynamics." *Social*

Cognitive and Affective Neuroscience 7 (6): 727–37. https://doi.org/10.1093/scan/nsr051.

Craig, A. D. 2009. "How Do You Feel - Now? The Anterior Insula and Human Awareness."

Nature Reviews Neuroscience. https://doi.org/10.1038/nrn2555.

Critchley, Hugo D., and Sarah N. Garfinkel. 2017. "Interoception and Emotion." *Current*

Opinion in Psychology 17: 7–14. https://doi.org/10.1016/j.copsyc.2017.04.020.

Critchley, Hugo D. 2005. "Neural Mechanisms of Autonomic, Affective, and Cognitive

Integration." *Journal of Comparative Neurology* 493 (1): 154–66.

https://doi.org/10.1002/cne.20749.

Dai, Rui, Zirui Huang, Huihui Tu, Luoyu Wang, Sean Tanabe, Xuchu Weng, Sheng He, and

Dongfeng Li. 2016. "Interplay between Heightened Temporal Variability of Spontaneous

Brain Activity and Task-Evoked Hyperactivation in the Blind." *Frontiers in Human*

Neuroscience 10 (December): 632. https://doi.org/10.3389/fnhum.2016.00632.

Damasio, Antonio R. 2005. *Descartes' Error : Emotion, Reason, and the Human Brain*.

Davis, Mark H. 1983. "Measuring Individual Differences in Empathy: Evidence for a

Multidimensional Approach." *Journal of Personality and Social Psychology* 44 (1): 113–

26. https://doi.org/10.1037/0022-3514.44.1.113.

Davis, Mark H. 1980. "A Multidimensional Approach to Individual Differences in Empathy."

Catalog of Selected Documents in Psychology 10: 85.

Decety, Jean, Chenyi Chen, Carla Harenski, and Kent A Kiehl. 2013. "An FMRI Study of

Affective Perspective Taking in Individuals with Psychopathy: Imagining Another in Pain

Does Not Evoke Empathy." *Frontiers in Human Neuroscience* 7 (September): 489.

https://doi.org/10.3389/fnhum.2013.00489.

Decety, Jean, and Philip L Jackson. 2004. "The Functional Architecture of Human Empathy."

Behavioral and Cognitive Neuroscience Reviews 3 (2): 71–100.

https://doi.org/10.1177/1534582304267187.

Decety, Jean, and Jessica A Sommerville. 2003. "Shared Representations between Self and

Other: A Social Cognitive Neuroscience View." *Trends in Cognitive Sciences* 7 (12): 527–

33. https://doi.org/10.1016/j.tics.2003.10.004.

Ernst, Jutta, Georg Northoff, Heinz Böker, Erich Seifritz, and Simone Grimm. 2013.
"Interoceptive Awareness Enhances Neural Activity during Empathy." *Human Brain*
Mapping 34 (7): 1615–24. https://doi.org/10.1002/hbm.22014.
Fan, Yan, Niall W. Duncan, Moritz de Greck, and Georg Northoff. 2011. "Is There a Core
Neural Network in Empathy? An FMRI Based Quantitative Meta-Analysis." *Neuroscience*
and Biobehavioral Reviews. Pergamon. https://doi.org/10.1016/j.neubiorev.2010.10.009.
Frith, Uta, John Morton, and Alan M Leslie. 1991. "The Cognitive Basis of a Biological
Disorder: Autism." *Trends in Neurosciences* 14 (10): 433–38.
Garcia-Larrea, L. 2012. "The Posterior Insular-Opercular Region and the Search of a Primary
Cortex for Pain." *Neurophysiologie Clinique* 42 (5): 299–313.
https://doi.org/10.1016/j.neucli.2012.06.001.
Garrett, Douglas D., Natasa Kovacevic, Anthony R. McIntosh, and Cheryl L. Grady. 2010.
"Blood Oxygen Level-Dependent Signal Variability Is More than Just Noise." *Journal of*
Neuroscience 30 (14): 4914–21. https://doi.org/10.1523/JNEUROSCI.5166-09.2010.
———. 2013. "The Modulation of BOLD Variability between Cognitive States Varies by Age
and Processing Speed." *Cerebral Cortex* 23 (3): 684–93.
https://doi.org/10.1093/cercor/bhs055.
Garrett, Douglas D, Gregory R Samanez-Larkin, Stuart W S MacDonald, Ulman Lindenberger,
Anthony R McIntosh, and Cheryl L Grady. 2013. "Moment-to-Moment Brain Signal
Variability: A next Frontier in Human Brain Mapping?" *Neuroscience & Biobehavioral*
Reviews 37 (4): 610–24.
Good, Tyler J, Joshua Villafuerte, Jennifer D Ryan, Cheryl L Grady, and Morgan D Barense.
2020. "Resting State BOLD Variability of the Posterior Medial Temporal Lobe Correlates
with Cognitive Performance in Older Adults with and without Risk for Cognitive Decline."
Eneuro 7 (3): ENEURO.0290-19.2020. https://doi.org/10.1523/ENEURO.0290-19.2020.
Gray, Marcus A, Neil A Harrison, Stefan Wiens, and Hugo D Critchley. 2007. "Modulation of
Emotional Appraisal by False Physiological Feedback during FMRI." *PLOS ONE* 2 (6):
e546. https://doi.org/10.1371/journal.pone.0000546.
Gurney, Edmund. 1884. *"What Is an Emotion?" Mind*. Vol. os-IX.
https://doi.org/10.1093/mind/os-IX.35.421.
Hampshire, Adam, Russell Thompson, John Duncan, and Adrian M Owen. 2009. "Selective
Tuning of the Right Inferior Frontal Gyrus during Target Detection." *Cognitive, Affective &*
Behavioral Neuroscience 9 (1): 103–12. https://doi.org/10.3758/CABN.9.1.103.
Healey, Meghan L, and Murray Grossman. 2018. "Cognitive and Affective Perspective-Taking:
Evidence for Shared and Dissociable Anatomical Substrates ." *Frontiers in Neurology* .
https://www.frontiersin.org/article/10.3389/fneur.2018.00491.
Heiden, Linda van der, Sigrid Scherpiet, Lilian Konicar, Niels Birbaumer, and Ralf Veit. 2013.
"Inter-Individual Differences in Successful Perspective Taking during Pain Perception
Mediates Emotional Responsiveness in Self and Others: An FMRI Study." *NeuroImage* 65:
387–94. https://doi.org/10.1016/j.neuroimage.2012.10.003.
Higo, Takayasu, Rogier B Mars, Erie D Boorman, Ethan R Buch, and Matthew F S Rushworth.
2011. "Distributed and Causal Influence of Frontal Operculum in Task Control."
Proceedings of the National Academy of Sciences 108 (10): 4230 LP – 4235.
https://doi.org/10.1073/pnas.1013361108.
Hyde, Daniel C, Mariana Aparicio Betancourt, and Charline E. Simon. 2015. "Human
Temporal-Parietal Junction Spontaneously Tracks Others' Beliefs: A Functional near-

Infrared Spectroscopy Study." *Human Brain Mapping* 36 (12): 4831–46.
https://doi.org/10.1002/hbm.22953.

Iacoboni, Marco. 2005. "Understanding Others: Imitation, Language, Empathy." *Perspectives on Imitation: From Cognitive Neuroscience to Social Science* 1: 77–99.

———. 2009. "Imitation, Empathy, and Mirror Neurons." *Annual Review of Psychology* 60 (1): 653–70. https://doi.org/10.1146/annurev.psych.60.110707.163604.

Jabbi, Mbemba, Marte Swart, and Christian Keysers. 2007. "Empathy for Positive and Negative Emotions in the Gustatory Cortex." *NeuroImage* 34 (4): 1744–53. https://doi.org/10.1016/j.neuroimage.2006.10.032.

Jackson, Philip L., Andrew N. Meltzoff, and Jean Decety. 2005. "How Do We Perceive the Pain of Others? A Window into the Neural Processes Involved in Empathy." *NeuroImage* 24 (3): 771–79. https://doi.org/10.1016/j.neuroimage.2004.09.006.

Johnson, Mark. 2001. *The Feeling of What Happens: Body and Emotion in the Making of Consciousness (Review)*. *The Journal of Speculative Philosophy*. Vol. 15. Houghton Mifflin Harcourt. https://doi.org/10.1353/jsp.2001.0038.

Keysers, Christian, Bruno Wicker, Valeria Gazzola, Jean Luc Anton, Leonardo Fogassi, and Vittorio Gallese. 2004. "A Touching Sight: SII/PV Activation during the Observation and Experience of Touch." *Neuron* 42 (2): 335–46. https://doi.org/10.1016/S0896-6273(04)00156-4.

Khalsa, Sahib S., Ralph Adolphs, Oliver G. Cameron, Hugo D. Critchley, Paul W. Davenport, Justin S. Feinstein, Jamie D. Feusner, et al. 2018. "Interoception and Mental Health: A Roadmap." *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* 3 (6): 501–13. https://doi.org/10.1016/j.bpsc.2017.12.004.

Kovács, Ágnes Melinda, Simone Kühn, György Gergely, Gergely Csibra, and Marcel Brass. 2014. "Are All Beliefs Equal? Implicit Belief Attributions Recruiting Core Brain Regions of Theory of Mind." *PLoS ONE* 9 (9). https://doi.org/10.1371/journal.pone.0106558.

Kuehn, Esther, Karsten Mueller, Gabriele Lohmann, and Simone Schuetz-Bosbach. 2016. "Interoceptive Awareness Changes the Posterior Insula Functional Connectivity Profile." *Brain Structure and Function* 221 (3): 1555–71. https://doi.org/10.1007/s00429-015-0989-8.

Kumral, D, F Şansal, E Cesnaite, K Mahjoory, E Al, M Gaebler, V V Nikulin, and A Villringer. 2020. "BOLD and EEG Signal Variability at Rest Differently Relate to Aging in the Human Brain." *NeuroImage* 207: 116373. https://doi.org/https://doi.org/10.1016/j.neuroimage.2019.116373.

Lane, Richard D. 2008. "Neural Substrates of Implicit and Explicit Emotional Processes: A Unifying Framework for Psychosomatic Medicine." *Psychosomatic Medicine* 70 (2). https://journals.lww.com/psychosomaticmedicine/Fulltext/2008/02000/Neural_Substrates_of_Implicit_and_Explicit.12.aspx.

Lawrence, E. J., P. Shaw, V. P. Giampietro, S. Surguladze, M. J. Brammer, and A. S. David. 2006. "The Role of 'shared Representations' in Social Perception and Empathy: An FMRI Study." *NeuroImage* 29 (4): 1173–84. https://doi.org/10.1016/j.neuroimage.2005.09.001.

Lieberman, Matthew D., and Naomi I. Eisenberger. 2009. "Neuroscience: Pains and Pleasures of Social Life." *Science* 323 (5916): 890–91. https://doi.org/10.1126/science.1170008.

Ma, Liangsuo, Binquan Wang, Xiying Chen, and Jinhu Xiong. 2007. "Detecting Functional Connectivity in the Resting Brain: A Comparison between ICA and CCA." *Magnetic Resonance Imaging* 25 (1): 47–56. https://doi.org/https://doi.org/10.1016/j.mri.2006.09.032.

Mar, Raymond A. 2011. "The Neural Bases of Social Cognition and Story Comprehension."
Annual Review of Psychology 62: 103–34.
McIntosh, Anthony Randal, Natasa Kovacevic, and Roxane J. Itier. 2008. "Increased Brain
Signal Variability Accompanies Lower Behavioral Variability in Development." *PLoS*
Computational Biology 4 (7): e1000106–e1000106.
https://doi.org/10.1371/journal.pcbi.1000106.
Mehling, Wolf E., Cynthia Price, Jennifer J. Daubenmier, Mike Acree, Elizabeth Bartmess, and
Anita Stewart. 2012. "The Multidimensional Assessment of Interoceptive Awareness
(MAIA)." Edited by Manos Tsakiris. *PLoS ONE* 7 (11): e48230.
https://doi.org/10.1371/journal.pone.0048230.
Naughtin, Claire K, Kristina Horne, Dana Schneider, Dustin Venini, Ashley York, and Paul E
Dux. 2017. "Do Implicit and Explicit Belief Processing Share Neural Substrates?" *Human*
Brain Mapping 38 (9): 4760–72.
Nelson, Steven M, Nico U F Dosenbach, Alexander L Cohen, Mark E Wheeler, Bradley L
Schlaggar, and Steven E Petersen. 2010. "Role of the Anterior Insula in Task-Level Control
and Focal Attention." *Brain Structure and Function* 214 (5): 669–80.
https://doi.org/10.1007/s00429-010-0260-2.
Nieto-Castanon, Alfonso. 2014. "NITRC: CONN : Functional Connectivity Toolbox: File
Release Notes and Changelog." 2014.
https://www.nitrc.org/frs/shownotes.php?release_id=2733.
———. 2015. "NITRC: CONN : Functional Connectivity Toolbox: RE: Conn-Preprocessing-
Band-Pass." 2015. https://www.nitrc.org/forum/message.php?msg_id=12691.
———. 2020. "NITRC: CONN : ICA Variability Calculation Method." 2020.
https://www.nitrc.org/forum/forum.php?set=custom&forum_id=1144&style=nested&max_r
ows=50.
Nomi, Jason S, Taylor S Bolt, C E Chiemeka Ezie, Lucina Q Uddin, and Aaron S Heller. 2017.
"Moment-to-Moment BOLD Signal Variability Reflects Regional Changes in Neural
Flexibility across the Lifespan." *The Journal of Neuroscience : The Official Journal of the*
Society for Neuroscience 37 (22): 5539–48. https://doi.org/10.1523/JNEUROSCI.3408-
16.2017.
Northoff, Georg, Alexander Heinzel, Moritz De Greck, Felix Bermpohl, Henrik Dobrowolny,
and Jaak Panksepp. 2006. "Self-Referential Processing in Our Brain—a Meta-Analysis of
Imaging Studies on the Self." *Neuroimage* 31 (1): 440–57.
Pardini, Matteo, and Paolo F. Nichelli. 2009. "Age-Related Decline in Mentalizing Skills across
Adult Life Span." *Experimental Aging Research* 35 (1): 98–106.
https://doi.org/10.1080/03610730802545259.
Peiffer, Claudine, Nicolas Costes, Philippe Hervé, and Luis Garcia-Larrea. 2008. "Relief of
Dyspnea Involves a Characteristic Brain Activation and a Specific Quality of Sensation."
American Journal of Respiratory and Critical Care Medicine 177 (4): 440–49.
https://doi.org/10.1164/rccm.200612-1774OC.
Penny, William, Karl Friston, John Ashburner, Stefan Kiebel, and Thomas Nichols. 2007.
Statistical Parametric Mapping: The Analysis of Functional Brain Images. *Statistical*
Parametric Mapping: The Analysis of Functional Brain Images. Elsevier Ltd.
https://doi.org/10.1016/B978-0-12-372560-8.X5000-1.
Perrett, David I, Jari K Hietanen, Michael W Oram, and Philip J Benson. 1992. "Organization
and Functions of Cells Responsive to Faces in the Temporal Cortex." *Philosophical*

Transactions of the Royal Society of London. Series B: Biological Sciences 335 (1273): 23–30.

Perrett, David I, Edmond T Rolls, and Woody Caan. 1982. "Visual Neurones Responsive to Faces in the Monkey Temporal Cortex." *Experimental Brain Research* 47 (3): 329–42.

Perrett, David I, P A J Smith, D D Potter, A J Mistlin, A S Head, Arthur David Milner, and M A Jeeves. 1985. "Visual Cells in the Temporal Cortex Sensitive to Face View and Gaze Direction." *Proceedings of the Royal Society of London. Series B. Biological Sciences* 223 (1232): 293–317.

Pollatos, Olga, Klaus Gramann, and Rainer Schandry. 2007. "Neural Systems Connecting Interoceptive Awareness and Feelings." *Human Brain Mapping* 28 (1): 9–18. https://doi.org/10.1002/hbm.20258.

Preckel, Katrin, Philipp Kanske, and Tania Singer. 2018. "On the Interaction of Social Affect and Cognition: Empathy, Compassion and Theory of Mind." *Current Opinion in Behavioral Sciences* 19: 1–6. https://doi.org/https://doi.org/10.1016/j.cobeha.2017.07.010.

Premack, David, and Guy Woodruff. 1978. "Chimpanzee Problem-Solving: A Test for Comprehension." *Science* 202 (4367): 532–35.

Pribram, Karl H. 1973. "The Primate Frontal Cortex–Executive of the Brain." In *Psychophysiology of the Frontal Lobes*, 293–314. Elsevier.

Rapps, Nora, Lukas van Oudenhove, Paul Enck, and Qasim Aziz. 2008. "Brain Imaging of Visceral Functions in Healthy Volunteers and IBS Patients." *Journal of Psychosomatic Research* 64 (6): 599–604. https://doi.org/10.1016/j.jpsychores.2008.02.018.

Rizzolatti, Giacomo, Leonardo Fogassi, and Vittorio Gallese. 2006. "Mirrors in the Mind." *Scientific American* 295 (5): 54–61.

Rombouts, Serge A.R.B., Frederik Barkhof, Frank G.C. Hoogenraad, Michiel Sprenger, and Philip Scheltens. 1998. "Within-Subject Reproducibility of Visual Activation Patterns with Functional Magnetic Resonance Imaging Using Multislice Echo Planar Imaging." *Magnetic Resonance Imaging* 16 (2): 105–13. https://doi.org/10.1016/S0730-725X(97)00253-1.

Saxe, Rebecca, and Lindsey J. Powell. 2006. "It's the Thought That Counts: Specific Brain Regions for One Component of Theory of Mind." *Psychological Science* 17 (8): 692–99. https://doi.org/10.1111/j.1467-9280.2006.01768.x.

Scarapicchia, Vanessa, Erin L Mazerolle, John D Fisk, Lesley J Ritchie, and Jodie R Gawryluk. 2018. "Resting State BOLD Variability in Alzheimer's Disease: A Marker of Cognitive Decline or Cerebrovascular Status? ." *Frontiers in Aging Neuroscience* . https://www.frontiersin.org/article/10.3389/fnagi.2018.00039.

Schienle, A., and W. Scharmüller. 2013. "Cerebellar Activity and Connectivity during the Experience of Disgust and Happiness." *Neuroscience* 246: 375–81. https://doi.org/10.1016/j.neuroscience.2013.04.048.

Schmahmann, J. D. 2001. "The Cerebrocerebellar System: Anatomic Substrates of the Cerebellar Contribution to Cognition and Emotion." *International Review of Psychiatry* 13 (4): 247–60. https://doi.org/10.1080/09540260120082092.

Schmahmann, Jeremy D., and Deepak N. Pandya. 1995. "Prefrontal Cortex Projections to the Basilar Pons in Rhesus Monkey: Implications for the Cerebellar Contribution to Higher Function." *Neuroscience Letters* 199 (3): 175–78. https://doi.org/10.1016/0304-3940(95)12056-A.

Schurz, Matthias, Joaquim Radua, Markus Aichhorn, Fabio Richlan, and Josef Perner. 2014. "Fractionating Theory of Mind: A Meta-Analysis of Functional Brain Imaging Studies."

Neuroscience & Biobehavioral Reviews 42: 9–34.

Sedeño, Lucas, Blas Couto, Margherita Melloni, Andrés Canales-Johnson, Adrián Yoris, Sandra
Baez, Sol Esteves, et al. 2014. "How Do You Feel When You Can't Feel Your Body?
Interoception, Functional Connectivity and Emotional Processing in Depersonalization-
Derealization Disorder." *PLoS ONE* 9 (6). https://doi.org/10.1371/journal.pone.0098769.

Singer, Tania, Hugo D. Critchley, and Kerstin Preuschoff. 2009. "A Common Role of Insula in
Feelings, Empathy and Uncertainty." *Trends in Cognitive Sciences* 13 (8): 334–40.
https://doi.org/10.1016/j.tics.2009.05.001.

Singer, Tania, and Claus Lamm. 2009. "The Social Neuroscience of Empathy." *Annals of the*
New York Academy of Sciences 1156 (1): 81–96. https://doi.org/10.1111/j.1749-
6632.2009.04418.x.

Singer, Tania, Ben Seymour, John O'Doherty, Holger Kaube, Raymond J. Dolan, and Chris D.
Frith. 2004. "Empathy for Pain Involves the Affective but Not Sensory Components of
Pain." *Science* 303 (5661): 1157–62. https://doi.org/10.1126/science.1093535.

Smith, Stephen M., Peter T. Fox, Karla L. Miller, David C. Glahn, P. Mickle Fox, Clare E.
Mackay, Nicola Filippini, et al. 2009. "Correspondence of the Brain's Functional
Architecture during Activation and Rest." *Proceedings of the National Academy of Sciences*
of the United States of America 106 (31): 13040–45.
https://doi.org/10.1073/pnas.0905267106.

Spreng, R Nathan, Raymond A Mar, and Alice S N Kim. 2009. "The Common Neural Basis of
Autobiographical Memory, Prospection, Navigation, Theory of Mind, and the Default
Mode: A Quantitative Meta-Analysis." *Journal of Cognitive Neuroscience* 21 (3): 489–510.

Strack, Fritz, Leonard L Martin, and Sabine Stepper. 1988. "Inhibiting and Facilitating
Conditions of the Human Smile: A Nonobtrusive Test of the Facial Feedback Hypothesis."
Journal of Personality and Social Psychology 54 (5): 768–77. https://doi.org/10.1037/0022-
3514.54.5.768.

Tavor, I., O. Parker Jones, R. B. Mars, S. M. Smith, T. E. Behrens, and S. Jbabdi. 2016. "Task-
Free MRI Predicts Individual Differences in Brain Activity during Task Performance."
Science 352 (6282): 216–20. https://doi.org/10.1126/science.aad8127.

Terasawa, Yuri, Hirokata Fukushima, and Satoshi Umeda. 2013. "How Does Interoceptive
Awareness Interact with the Subjective Experience of Emotion? An FMRI Study." *Human*
Brain Mapping 34 (3): 598–612. https://doi.org/10.1002/hbm.21458.

Tognoli, Emmanuelle, and J A Scott Kelso. 2014. "The Metastable Brain." *Neuron* 81 (1): 35–
48. https://doi.org/10.1016/j.neuron.2013.12.022.

Tops, Mattie, Maarten Boksem, Phan Luu, and Don Tucker. 2010. "Brain Substrates of
Behavioral Programs Associated with Self-Regulation." *Frontiers in Psychology* 1: 152.
https://doi.org/10.3389/fpsyg.2010.00152.

Tops, Mattie, and Ritske De Jong. 2006. "Posing for Success: Clenching a Fist Facilitates
Approach." *Psychonomic Bulletin & Review* 13 (2): 229–34.
https://doi.org/10.3758/BF03193835.

Vakorin, Vasily A, Sarah Lippé, and Anthony R McIntosh. 2011. "Variability of Brain Signals
Processed Locally Transforms into Higher Connectivity with Brain Development." *The*
Journal of Neuroscience : The Official Journal of the Society for Neuroscience 31 (17):
6405–13. https://doi.org/10.1523/JNEUROSCI.3153-10.2011.

Vakorin, Vasily A, Bratislav Mišić, Olga Krakovska, and Anthony Randal McIntosh. 2011.
"Empirical and Theoretical Aspects of Generation and Transfer of Information in a

Neuromagnetic Source Network." *Frontiers in Systems Neuroscience* 5 (November): 96.
https://doi.org/10.3389/fnsys.2011.00096.
Vanderwal, Tamara, Clare Kelly, Jeffrey Eilbott, Linda C. Mayes, and F. Xavier Castellanos.
2015. "Inscapes: A Movie Paradigm to Improve Compliance in Functional Magnetic
Resonance Imaging." *NeuroImage* 122 (November): 222–32.
https://doi.org/10.1016/j.neuroimage.2015.07.069.
Vogeley, Kai, Patrick Bussfeld, Albert Newen, Sylvie Herrmann, Francesca Happé, Peter Falkai,
Wolfgang Maier, Nadim J Shah, Gereon R Fink, and Karl Zilles. 2001. "Mind Reading:
Neural Mechanisms of Theory of Mind and Self-Perspective." *Neuroimage* 14 (1): 170–81.
Whitfield-Gabrieli, Susan, and Alfonso Nieto-Castanon. 2012. "Conn: A Functional
Connectivity Toolbox for Correlated and Anticorrelated Brain Networks." *Brain
Connectivity* 2 (3): 125–41. https://doi.org/10.1089/brain.2012.0073.
Wicker, Bruno, Christian Keysers, Jane Plailly, Jean-Pierre Royet, Vittorio Gallese, and
Giacomo Rizzolatti. 2003. "Both of Us Disgusted in My Insula: The Common Neural Basis
of Seeing and Feeling Disgust." *Neuron* 40 (3): 655–64.
Wiens, Stefan. 2005. "Interoception in Emotional Experience." *Current Opinion in Neurology*.
Lippincott Williams and Wilkins. https://doi.org/10.1097/01.wco.0000168079.92106.99.
Worsley, K. J., S. Marrett, P. Neelin, A. C. Vandal, K. J. Friston, and A. C. Evans. 1996. "A
Unified Statistical Approach for Determining Significant Signals in Images of Cerebral
Activation." *Human Brain Mapping* 4 (1): 58–73. https://doi.org/10.1002/(SICI)1097-
0193(1996)4:1<58::AID-HBM4>3.0.CO;2-O.
Wu, Yan, Jiaojian Wang, Yun Zhang, Dingchen Zheng, Jinfeng Zhang, Menglin Rong, Huawang
Wu, Yinyan Wang, Ke Zhou, and Tianzi Jiang. 2016. "The Neuroanatomical Basis for
Posterior Superior Parietal Lobule Control Lateralization of Visuospatial Attention."
Frontiers in Neuroanatomy 10 (March): 32. https://doi.org/10.3389/fnana.2016.00032.
Yan, Zhiqiang, Skylar Hong, Fuli Liu, and Yanjie Su. 2020. "A Meta-Analysis of the
Relationship between Empathy and Executive Function." *PsyCh Journal* 9 (1): 34–43.
https://doi.org/10.1002/pchj.311.
Zaki, Jamil, Jamil Zaki, Carol Dweck, and Andrew Ickes. 2014. "Empathy : A Motivated
Account Empathy : A Motivated Account Experiences." *Psychological Bulletin* 140 (6):
1608–47.
Zhang, Liwen, Xi-Nian Zuo, Kwun Kei Ng, Joanna Su Xian Chong, Hee Youn Shim, Marcus
Qin Wen Ong, Yng Miin Loke, et al. 2020. "Distinct BOLD Variability Changes in the
Default Mode and Salience Networks in Alzheimer's Disease Spectrum and Associations
with Cognitive Decline." *Scientific Reports* 10 (1): 6457. https://doi.org/10.1038/s41598-
020-63540-4.

Table 1

Sociodemographic Characteristics of Participants

	Frequency	Min	Max	%
Gender	10	Male		38.5
	16	Female		61.5
	Total: 26			
Ethnicity	1	Hispanic/Latino		3.8
	24	Not Hispanic/Latino		92.3
	1	Unknown		3.8
Race	22	White		84
	1	African American		4
	2	Asian		8
	1	More than 1 Race		4
				Mean
Age		18	31	21.8
Education (Years)		12	21	14.2

Table 2

Intercorrelations between IRI and MAIA Questionnaires and Descriptive Statistics

	1	2	3	4	5	6	7	8	9	10	11	12	Mean	STD
IRI Total (1)	-												75.79	11.12
IRI Affective Empathy (2)	0.60	-											31.84	6.3
IRI Personal Distress (3)	0.11	0.79	-										10.24	5.61
IRI Empathic Concern (4)	0.80	0.48	-0.17	-									21.6	3.85
IRI Cognitive Empathy (5)	0.83	0.05	-0.41	0.67	-								44	8.52
IRI Perspective Taking (6)	0.58	-0.15	-0.52	0.50	0.83	-							21.64	4.97
IRI Fantasy (7)	0.80	0.22	-0.18	0.61	0.84	0.39	-						22.36	5.25
MAIA Total (8)	-0.23	**-0.64***	**-0.66***	-0.08	0.16	0.31	0.04	-					23.28	4.36
MAIA Awareness of Body Sensations (9)	-0.30	-0.36	-0.19	-0.29	-0.12	-0.17	0.03	0.54	-				2.98	0.81
MAIA Emotional Reaction (10)	-0.33	-0.23	-0.21	-0.08	-0.25	-0.11	0.31	0.37	0.03	-			5.65	1.26
MAIA Capacity to Regulate Attention (11)	-0.33	**-0.83***	**-0.74***	-0.27	0.17	0.32	-0.03	0.79	0.48	0.24	-		3.05	0.94
MAIA Awareness of Mind Body Integration (12)	0.09	-0.34	**-0.49***	0.16	0.35	**0.41***	0.18	0.82	0.37	0.05	0.50	--	8.12	2.46
MAIA Trusting Body Sensations (13)	-0.25	**-0.55***	**-0.53***	-0.12	0.08	0.31	0.18	0.71	0.21	0.32	0.53	0.40	3.48	1.08

Table 2: Intercorrelations between IRI and MAIA Subscales and Descriptive Statistics (same subscale correlations not flagged for significance). * represents correlations statistically significant after multiple comparison correction. Uncorrected Mean and STD reported.

Table 3

Network Connectivity Statistics

Questionnaire Subscale	Region	Laterality	Peak Cluster (X, Y, Z)	size	p-FDR	T stat	Effect Size	90% CI
MAIA Total	IFO	R	46, 48, 2	289	0.04	-6.84	2.7	2, 3.25
Capacity to Regulate Attention	IFO	R	46, 48, 14	407	0.0003	-6.02	2.8	2, 3.25
IRI Total	IFO	R	36, 56, 6	207	0.03	5.26	2.7	1.5, 3
Affective Empathy	IFO	R	40, 48, -4	284	0.02	5.67	2.8	2, 3.5

Table 3: Connectivity Statistics of rIFO cluster within network associated with lower total IA and greater Affective empathy.

Figure 1: Still-shots from the InScapes Movie

Figure 2: Greater connectivity in the rIFO is associated with lower total IA and greater Affective empathy. Statistical maps are thresholded at FDR-corrected cluster-based $p < 0.05$ after voxel threshold at $p < 0.001$.

Figure 3: Shared network variability between IRI Personal Distress and MAIA Awareness of Body Sensations. Statistical maps are thresholded at FDR-corrected cluster-based $p < 0.05$ after voxel threshold at $p < 0.001$. *Note:* Red dots in graphs represent observed values, Blue dots fitted values.

Figure 4: Shared network variability between IRI Perspective Taking and MAIA Mind Body Integration. Statistical maps are thresholded at FDR-corrected cluster-based $p < 0.05$ after voxel threshold at $p < 0.001$. *Note:* Red dots in graphs represent observed values, Blue dots fitted values.