Title: Evidence for unique small RNA modifications in Alzheimer's disease

Xudong Zhang[1]*, Fatima Trebak[2]*, Lucas AC Souza[2], Junchao Shi[3], Tong Zhou[3],

Patrick G. Kehoe[4], Qi Chen[1#], Yumei Feng Earley[2#]

1. Division of Biomedical Sciences, School of Medicine, University of California, Riverside, Riverside, CA, USA
2. Departments of Pharmacology, Physiology & Cell Biology, Center for Molecular & Cellular Signal Transduction in the Cardiovascular System, University of Nevada, Reno, School of Medicine, Reno, NV, USA
3. Departments of Physiology & Cell Biology, University of Nevada, Reno, School of Medicine, Reno, NV, USA
4. Dementia Research Group, Translational Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

*Contributed equally to this work

#**Correspondence to:**

Yumei Feng Earley, MD, PhD, FAHA
Associate Professor of Pharmacology and Physiology & Cell Biology
Center for Molecular & Cellular Signal Transduction in the Cardiovascular System
University of Nevada, Reno, School of Medicine
1664 North Virginia Street, Mail-stop 0318
Reno, NV, 89557, USA
E-mail: yumeifeng@med.unr.edu

Qi Chen, MD, PhD
Assistant Professor
Division of Biomedical Sciences, School of Medicine
University of California, Riverside
Riverside, CA, 92521, USA
E-mail: Qi.Chen@medsch.ucr.edu

Abstract

Background: While significant advances have been made in uncovering the aetiology of Alzheimer's disease and related dementias at the genetic level, molecular events at the epigenetic level remain largely un-investigated. Emerging evidence indicates that small non-coding RNAs (sncRNAs) and their associated RNA modifications are important regulators of complex physiological and pathological processes, including aging, stress responses, and epigenetic inheritance. However, whether small RNAs and their modifications are altered in dementia is not known.

Methods: We performed LC-MS/MS–based, high-throughput small RNA modification assays on post-mortem samples of the prefrontal lobe of Alzheimer's disease (AD), vascular dementia+AD (VaD+AD), and control individuals.

Findings: We report altered small RNA modifications in the 15–25-nt fractions enriched for microRNAs, and 30–40-nt RNA fractions enriched for tRNA-derived small RNAs (tsRNAs), rRNA-derived small RNAs (rsRNAs), and YRNA-derived small RNAs (ysRNAs). Interestingly, most of these altered RNA modifications were unique to AD and were not identified in VaD+AD subjects. In addition, sequencing of small RNA in the 30–40-nt fraction from AD cortices revealed reductions in rsRNA-5S, tsRNA-Tyr, and tsRNA-Arg.

Interpretation: These data suggest that sncRNAs and their associated modifications are novel signals that may be linked to the pathogenesis and development of Alzheimer's Disease.

Funding: NIH grants (R01HL122770, R01HL091905, 1P20GM130459, R01HD092431, P50HD098593, GM103440), AHA grant (17IRG33370128), Sigmund Gestetner Foundation Fellowship to P Kehoe.

Keywords: Non-coding RNA, Alzheimer's disease, Prefrontal cortex, Epigenetics

Research in Context

Evidence before this study

Alzheimer's disease (AD) and vascular dementia (VaD) are marked by cognitive impairment and neuropathologies caused by significant neuronal death. Associated gene mutations are rare in subjects with dementia, and the aetiology of these diseases is still not completely understood. Recent emerging evidence suggests that epigenetic changes are risk factors for the development of dementia. However, studies assessing small RNA modifications—one of the features of epigenetics—in dementia are lacking.

Added value of this study

We used high-throughput liquid chromatography-tandem mass spectrometry and small RNA sequencing to profile small RNA modifications and the composition of small RNAs in post-mortem samples of the prefrontal cortex of AD and control subjects. We detected and quantified 16 types of RNA modifications and identified distinct small non-coding RNAs and modification signatures in AD subjects compared with controls.

Implications of all the available evidence

This study identified novel types and compositions of small RNA modifications in the prefrontal cortex of AD patients compared with control subjects in post-mortem samples. The cellular locations of these RNA modifications and whether they are drivers or outcomes of AD are still not known. However, results from the present study might open new possibilities for dissecting the dementia pathology.

Introduction

Alzheimer's disease (AD) and severe vascular cognitive impairment, also known as vascular dementia (VaD), are two of the most common types of dementia. They are characterized by cognitive impairment and neuropathology caused by neurotoxic forms of amyloid beta peptide, loss of synapses and neuronal function and, ultimately, significant neuronal loss. The aetiology of these forms of dementia is still not fully understood.[1]

Recent studies have highlighted epigenetic changes associated with aging or acquired through interactions with the environment as important risk factors for dementia, reflecting the fact that causative gene mutations are rare and are only present in a very small percentage of dementia patients. RNA modification is one of the most recently discovered epigenetic-mediated mechanisms for regulating physiological and pathological processes. Small RNA modifications, particularly those involving microRNAs (miRNAs) and transfer RNA-derived small RNAs (tsRNAs), have attracted considerable recent attention for their potential clinical relevance and possible use as diagnostic markers or therapeutic targets of diseases.[2-4] To date, evidence for changes in small RNA modifications in dementia is lacking. Thus, to gain insights into possible new mechanisms and identify potential biomarkers for dementia, we undertook an exploratory study to profile small RNA expression and modification status in dementia.

In this report, we profiled small RNAs modifications in post-mortem samples of human brain cortical tissue from patients with clinically and pathologically diagnosed AD, with or without VaD, as well as control individuals. To simultaneously detect and quantify 16 types of RNA modifications[5,6] in different small RNA fractions [15–25 nucleotides (nt), 30–40 nt] from AD patients and control individuals, we used high-throughput liquid chromatography-tandem mass spectrometry (LC-MS/MS). We also profiled the composition of small RNAs in AD and control

subjects using small RNA sequencing. These exploratory examinations revealed distinct sncRNA expression and RNA modification signatures between AD and controls, suggesting that these small RNAs are novel factors associated with the pathogenesis and/or progression of AD.

Materials and methods

Human subjects for RNA modification assays

A total of 34 post-mortem brain prefrontal lobe cortex samples from AD patients (n=18), VaD+AD patients (n=9), and control human subjects (n=7) were collected by post-mortem autopsy, snap-frozen, and stored in liquid nitrogen. The time between biospecimen acquisition and analysis ranged from 5 to 30 years, according to the Brain Bank database. Biospecimens were shipped on dry ice. Information from clinical and pathology reports was obtained from the brain bank database, developed as described.[7] On the basis of their clinical diagnosis, subjects were categorized into three groups: no significant abnormalities, AD, and VaD. All clinically diagnosed VaD subjects used in this study exhibited clear AD tau pathology and Aβ plaques with a neuropathological report of AD. Thus, this group was termed VaD+AD for purposes of distinguishing it from the AD group. Subjects were selected according to their confirmed clinical and pathological diagnosis as AD or normal, and were obtained from the Human Tissue Authority-licensed South West Dementia Brain Bank http://www.bristol.ac.uk/translational-health-sciences/research/neurosciences/research/dementia/swdbb/, accessed on March 24th, 2020), University of Bristol, with tissue bank ethics approval from the South West–Central Bristol Research Ethics Committee. Clinical and pathological data included patient history, diagnosis and medications; pathological reports were available for retrospective analysis. Patients' personal information has been de-identified. Data for this study was collected between 1989

and 2015. The Research Integrity Offices at the University of Nevada, Reno, and University of

Bristol have determined that this project complies with human research protection oversight by

the Institutional Review Board.

Isolation of total RNA from the human prefrontal cortex

Total RNA was isolated from 100 mg of human prefrontal tissues using the TRIzol reagent

(Thermo Fisher Scientific, Inc. Waltham, MA, USA), as described by the manufacturer. RNA

quantity was evaluated using a NanoDrop 2000 microvolume spectrophotometer (Thermo

Fisher). RNA samples were stored at -80°C until analysis. Researchers performing small RNA

sequencing and RNA modifications assays on RNA samples were blinded to diagnosis and

group-identifying information.

Small RNA Sequencing

Ten RNA samples from prefrontal lobe cortexes of control subjects and AD patients were

submitted to BGI (Cambridge, MA, USA) for sequencing of 15–40-nt small RNAs. The number

of clean reads varied from 30M to 40M across all samples. The Q20 score was greater than

99·5% for all samples. Raw sequencing data for each sample were processed and analysed using

our   newly   developed   computational   framework,   *SPORTS1·0*[8]

(https://github.com/junchaoshi/sports1.0), which is designed to optimize the annotation and

quantification of non-canonical small RNAs (e.g., tsRNAs) in small RNA sequencing data.

Sequencing adapters were removed, after which reads with lengths outside of the defined range

and those with nucleotides other than ATUCG were discarded. Clean reads were mapped

against   the   precompiled   human   small   RNA   annotation   database

(https://github.com/junchaoshi/sports1.0/blob/master/precompiled_annotation_database.md).

Small RNAs 15–25 and 30–40 nt in length were summarized separately.

Small RNA purification and analysis of RNA modifications using LC-MS/MS

Standardized ribonucleoside preparations, small RNA purification, and LC-MS/MS-based analysis of RNA modifications in RNA samples were performed as previously described.[6] Purified small RNAs (100–200 ng) from human brain samples were digested as input and loaded onto a ThermoFisher Vantage Quadrupole mass spectrometer connected to a Thermo Ultimate 3000 UHPLC system equipped with an electrospray ionization source. The MS system was operated in positive ion mode using a multiple reaction monitoring (MRM) approach. LC-MS/MS raw data were acquired using Xcalibur Workstation software and were processed using Xcalibur QuanBrowser for quantification of modified ribonucleoside concentrations. The percentage of each modified ribonucleoside was normalized to the total amount of quantified ribonucleosides containing the same nucleobase, an approach that decreases/eliminates errors caused by sample loading variation. For example, the percentage of m^5C = mole concentration (m^5C)/mole concentration $(m^5C + Cm + C + ac^4C)$. Fold-changes in RNA modifications between different groups were calculated based on the percentage of modified ribonuclosides. Brain RNAs 15–25 and 30–40 nucleotides (nt) in length were examined.

Statistical analysis

Data are presented as means ± SEM and were plotted using Prism8 software (GraphPad, La Jolla, CA, USA). Differential expression analyses were performed using the *edgeR* tool (pmid: 19910308), controlling for age and sex. RNAs with a false discovery rate < 10% were deemed to be differentially expressed. RNA modification levels between groups were compared using a linear model that controls for age and sex. Unpaired *t* tests were used for comparisons of participants' characteristics between groups. A *P*-value < 0·05 was considered statistically significant.

Results

Participant characteristics

As summarized in Table 1, patients (AD and VaD+AD) and control individuals were similar in terms of age. Almost all patients were White/European except for one patient whose race was unknown. There were slightly more males (58·8%) than females (41·2%). Brain weight was significantly lower in AD (P = 0·0099) and VaD+AD (P = 0·046) groups compared with the control group. The post-mortem autopsy delay time was similar among groups.

Altered small RNA modification profiles in the cerebral cortex of AD patients

To systemically analyse small RNA expression and modification profiles, we collected the 15–40-nt RNA fraction from the brain frontal lobe cortex for small RNA sequencing as we previously described[5,6], and then performed bioinformatic analyses using our recently developed software, *SPORTS1.0*.[8] In addition, 15–25-nt and 30–40-nt RNA fractions were collected for high-throughput examination of RNA modification by LC-MS/MS (Figure 1).[5,9] These analyses resulted in the identification and quantification of 16 types of RNA modifications.

In the 15–25-nt RNA fraction from the cortex of AD brains (Figure 2A), we found an increase in 2'-O-methylcytidine (Cm) and 7-methylguanosine (m^{7}G), and significant reductions in N^2,N^2,7-trimethylguanosine (m^{2,2,7}G) and N^2,N^2-dimethylguanosine (m^{2,2}G) compared with controls (Figure 2B–E). Interestingly, these changes were not present in the cortex of VaD+AD patients' brains, and thus were unique to the brains' of AD-only subjects. Other RNA modifications that were not significantly different among groups were also detected, as shown in Supplemental Figure S1.

In the 30–40-nt small RNA fraction from the cortex of AD brains (Figure 3A), we found significantly higher levels of Cm, 2'-O-methyluridine (Um) and 7-methylguanosine (m^7G) modifications, and reductions in 1-methylguanosine (m^1G), $m^{2,2,7}G$, and pseudouridine (psi or Ψ) modifications compared with controls (Figure 3B–G). Among 30–40-nt small RNAs, we found only one specific RNA modification, psi, that was changed in the VaD+AD brain cortex, where this modification was reduced compared with controls (Figure 3G). Again, other RNA modifications were detected in this 30–40-nt fraction that did not exhibit significant differences among groups, as shown in Supplemental Figure S2.

Altered 30–40-nt small RNA expression profiles in the cerebral cortex of AD patients

To systemically analyse small RNA expression profiles, we collected the 15–40-nt fraction from the brain frontal lobe cortex of AD patients and controls for small RNA sequencing, as we previously described[5,6], and then performed bioinformatic analyses using our recently developed software, *SPORTS1·0*[8]. Our RNA-seq analysis showed that the 15–25-nt RNA population was predominantly miRNAs (Figure 4A) in AD and control samples, whereas the expression profile of the small RNA population in the 30–40-nt fraction showed more dynamic changes. Our RNA-seq data revealed three major subtypes in the 30–40-nt fraction, namely tRNA-derived small RNAs (tsRNA), rRNA-derived small RNAs (rsRNAs) and Y RNA-derived small RNAs (ysRNAs); other un-annotated RNAs were also found (Figure 4B). Although there were trends toward an increase in tsRNAs and a reduction in ysRNAs in AD patients, these differences did not reach statistical significance. A further analysis of specific small RNA fragments in these samples revealed that the majority of the 15–25-nt fraction was miRNAs (Figure 4A). We did not find significant differences in the specific small RNA sequences in this fraction between AD and control samples (data not shown). Interestingly, in

the 30–40-nt fraction, we found a significant reduction in rRNA-5S, tsRNA-Tyr and tsRNA-

Arg fragments in AD patients compared with controls (Figure 4C–E).

Discussion

Emerging evidence shows that small RNAs, including miRNAs, tsRNAs and piwi-interacting

RNAs, harbour a diversity of RNA modifications.[10] The physiological and pathological

importance of these small, non-coding RNA modifications has only recently begun to emerge,

as highlighted by studies revealing their active involvement in stress responses, metabolism,

immunity, and epigenetic inheritance of environmentally acquired traits.[2] RNA modifications

have recently been detected in the nervous system, where they are involved in the regulation of

cortical differentiation, behaviour, and brain functions.[11,12] However, to our knowledge, there

have been no reports on the status of small RNA modifications in AD or VaD to date.

The current study showed intriguing differences in small RNA-modification signatures between

AD patients and control subjects, suggesting an association and possible contribution of such

modifications to the pathogenesis and/or progression of AD. Notably, most of the altered RNA

modifications were not observed in VaD+AD patients, with the exception of psi, as noted above,

providing molecular evidence for potential distinct signatures between these two patient groups.

In the last three decades, dementia research has demonstrated significant overlaps between AD

and VaD in terms of clinical symptoms, risk factors, and post-mortem brain autopsy findings.[13-

15] Therapeutically, drugs that enhance cholinergic activity are as effective in patients suffering

from VaD as they are in AD patients.[16] More recently, drugs that have traditionally been used

for cardiovascular diseases, especially renin-angiotensin system blockers, have shown benefits

not only in VaD patients, but also in AD patients.[17-19] However, the pathogenesis of AD, VaD,

and VaD+AD remains incompletely understood. We were surprised to find only one common

signature among RNA modifications in AD and VaD+AD patients that might contribute to

shared mechanisms of dementia pathology. Alternatively, these distinct molecular signatures

could suggest the existence of as-yet undiscovered differences in pathological processes

between these conditions, a possibility that might be harnessed for future condition-specific

treatments.

The 15–25-nt RNAs were predominantly miRNAs, whereas 30–40-nt RNAs were

predominantly tsRNAs, rsRNAs and ysRNAs, as demonstrated by small RNA sequencing. We

found changes in RNA-modification profiles in the AD brain cortex in both fractions of small

RNAs. While the importance of miRNA in the pathogenesis of AD is relatively well described[20],

the role of rsRNAs and ysRNAs in a pathophysiological context has only begun to emerge.[6,21,22]

The dynamic expression of these non-canonical small RNAs suggests the possibility of

unidentified biological functions that warrant further investigation[23,24], particularly in relation

to several risk factor genes that have been identified for AD in the last few decades.[25,26] This

situation is similar to that for tsRNAs, which have shown recently expanding functions.[4,27,28]

The novel RNA modifications that showed dynamic changes in AD patients included Cm, Um,

m^7G, m^1G, $m^{2,2,7}G$, and psi. The functions and identities of these small RNA modifications in

AD are not yet understood. We identified increases in 2'-O-methylation of cytidine (Cm),

guanosine (Gm) and uridine (Um), as well as m^7G, while observing a reduction in $m^{2,2,7}G$ in

the brain cortex of dementia patients, suggesting the potential importance of these RNA

modifications, either as compensatory responses (Cm, Gm, Um, m^7G)[29-31] or as actual causes

of pathogenesis ($m^{2,2,7}G$).[32,33]

tRNA-derived small RNAs are reported to contribute to multiple pathological processes,

including cancer, viral infection, and age-related neurodegenerative diseases.[2] In the

senescence-accelerated mouse prone 8 (SAMP8) model, which mimics age-related neuro-

disorders such as AD and Parkinson disease, tsRNAs, including tsRNA-Tyr and tsRNA-Arg,

in brain tissue are dysregulated compared with normal brain tissue. The targets of these

dysregulated tsRNAs are enriched in neurodevelopment pathways such as synapse formation.[34]

In addition, knockout of the RNA kinase, Clp1, in mice leads to progressive loss of motor

neuron function in association with the accumulation of Try-tRNA and Arg-tRNA small RNAs.

These tsRNAs sensitize motor neurons to oxidative stress-induced cell death, suggesting that

tsRNAs are involved in normal motor neuron functions and responses to oxidative stress.[35] A

similar phenomenon was reported in a human neurological disease cohort, and it was shown

that transfection of small RNAs from 5' Tyr-tRNA can protect CLP-mutant cells from oxidative

stress-induced cell death.[36] In our study, the decrease in 5'tsRNA-Tyr in AD subjects may

confer greater vulnerability to oxidative stress on neurons. Interestingly another report showed

that the expression of 5s rRNA and the oxidized level of 5S rRNA are dynamically modulated

in AD's subjects[37], findings that may be related to the biogenesis of rsRNAs and might explain

alterations of rsRNA-5s in the AD group in the current study.

Whether the changes in small RNAs and RNA modifications observed here represent drivers

or outcomes of AD remains unknown. The precise meaning of each of these changes in terms

of their specific functions awaits discovery. Future efforts to pinpoint the locations of these

modifications in each sncRNA population and identify enzymes involved in their regulation

would be invaluable[38,39] and will open new avenues for dissecting the nature of dementia

pathology.

Funding

This work was supported, in part, by grants from the National Institutes of Health (NIH/NHLBI) (R01HL122770, R01HL091905), NIH/NIGMS (1P20GM130459) and the American Heart Association National Center (17IRG33370128) to Y. Feng Earley, and from the NIH/NICHD to Q. Chen (R01HD092431) and T. Zhou (P50HD098593). Research reported in this publication used instruments from the Dr. Mick Hitchcock Proteomics Core supported by a grant from the NIH/NIGMS (GM103440) of the National Institutes of Health. University of Bristol Alumni. P Kehoe is supported by a Fellowship from the Sigmund Gestetner Foundation. The funders had no role in study design, data collection, analysis, and interpretation, as well as writing of the manuscript.

Declaration of competing interests

The authors declare no competing interests.

Data Availability

All data associated with this study are contained in the paper or Supplementary Materials. RNA sequencing data will be deposited in an online repository (Mendeley Data) upon acceptance of the manuscript.

Acknowledgements

We would like to thank Dr. David Quilici and Rebekah Woolsey from the Mick Hitchcock, Ph.D. Nevada Proteomics Center for technical support with the HPLC/MS instruments. We would especially like to thank Dr. Laura Palmer for tissue processing and facilitating access to clinical data from the South West Dementia Brain Bank at the University of Bristol, which is supported by grants from BRACE, ABBUK.

Supplemental Materials

336 Supplementary material associated with this article can be found in the online version at doi:

337 10.6084/m9.figshare.12227393

References:

1. Panza F, Lozupone M, Watling M, Imbimbo BP. Do BACE inhibitor failures in
Alzheimer patients challenge the amyloid hypothesis of the disease? *Expert review of*
neurotherapeutics 2019; **19**(7): 599-602.
2. Zhang X, Cozen AE, Liu Y, Chen Q, Lowe TM. Small RNA Modifications: Integral to
Function and Disease. *Trends Mol Med* 2016; **22**(12): 1025-34.
3. Chen Q, Yan W, Duan E. Epigenetic inheritance of acquired traits through sperm RNAs
and sperm RNA modifications. *Nature reviews Genetics* 2016; **17**(12): 733-43.
4. Schimmel P. The emerging complexity of the tRNA world: mammalian tRNAs beyond
protein synthesis. *Nature reviews Molecular cell biology* 2018; **19**(1): 45-58.
5. Chen Q, Yan M, Cao Z, et al. Sperm tsRNAs contribute to intergenerational inheritance
of an acquired metabolic disorder. *Science* 2016; **351**(6271): 397-400.
6. Zhang Y, Zhang X, Shi J, et al. Dnmt2 mediates intergenerational transmission of
paternally acquired metabolic disorders through sperm small non-coding RNAs. *Nature cell*
biology 2018; **20**(5): 535-40.
7. Hyman BT, Phelps CH, Beach TG, et al. National Institute on Aging-Alzheimer's
Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimer's*
& dementia : the journal of the Alzheimer's Association 2012; **8**(1): 1-13.
8. Shi J, Ko EA, Sanders KM, Chen Q, Zhou T. SPORTS1.0: A Tool for Annotating and
Profiling Non-coding RNAs Optimized for rRNA- and tRNA-derived Small RNAs. *Genomics*
Proteomics Bioinformatics 2018; **16**(2): 144-51.
9. Fang H, Deng M, Zhang L, et al. Role of (Pro)Renin Receptor in Albumin Overload-
Induced Nephropathy in Rats. *Am J Physiol Renal Physiol* 2018.
10. Roundtree IA, Evans ME, Pan T, He C. Dynamic RNA Modifications in Gene
Expression Regulation. *Cell* 2017; **169**(7): 1187-200.
11. Satterlee JS, Basanta-Sanchez M, Blanco S, et al. Novel RNA modifications in the
nervous system: form and function. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience 2014; **34**(46): 15170-7.
12. Jung Y, Goldman D. Role of RNA modifications in brain and behavior. *Genes, brain,*
and behavior 2018; **17**(3): e12444.
13. Ravona-Springer R, Davidson M, Noy S. Is the distinction between Alzheimer's disease
and vascular dementia possible and relevant? *Dialogues in clinical neuroscience* 2003; **5**(1): 7-
15.
14. Mirra SS, Heyman A, McKeel D, et al. The Consortium to Establish a Registry for
Alzheimer's Disease (CERAD). *Neurology* 1991; **41**(4): 479.
15. Meyer JS, Xu G, Thornby J, Chowdhury MH, Quach M. Is mild cognitive impairment
prodromal for vascular dementia like Alzheimer's disease? *Stroke* 2002; **33**(8): 1981-5.
16. Erkinjuntti T, Kurz A, Gauthier S, Bullock R, Lilienfeld S, Damaraju CV. Efficacy of
galantamine in probable vascular dementia and Alzheimer's disease combined with
cerebrovascular disease: a randomised trial. *Lancet* 2002; **359**(9314): 1283-90.
17. Zhuang S, Wang H-F, Wang X, Li J, Xing C-M. The association of renin-angiotensin
system blockade use with the risks of cognitive impairment of aging and Alzheimer's disease:
A meta-analysis. *Journal of Clinical Neuroscience* 2016; **33**: 32-8.
18. Chou C-L, Yeh H-I. The Role of the Renin-Angiotensin System in Amyloid Metabolism
of Alzheimer's Disease. *Acta Cardiologica Sinica* 2014; **30**(2): 114-8.
19. Gebre AK, Altaye BM, Atey TM, Tuem KB, Berhe DF. Targeting Renin-Angiotensin
System Against Alzheimer's Disease. *Frontiers in pharmacology* 2018; **9**: 440.
20. Ayers D, Scerri C. Non-coding RNA influences in dementia. *Noncoding RNA Res* 2018;
3(4): 188-94.

21.   Hizir Z, Bottini S, Grandjean V, Trabucchi M, Repetto E. RNY (YRNA)-derived small
RNAs regulate cell death and inflammation in monocytes/macrophages. *Cell Death Dis* 2017;
8(1): e2530.
22.   Zhang Y, Shi J, Rassoulzadegan M, Tuorto F, Chen Q. Sperm RNA code programmes
the metabolic health of offspring. *Nature reviews Endocrinology* 2019; **15**(8): 489-98.
23.   Kowalski MP, Krude T. Functional roles of non-coding Y RNAs. *The international*
journal of biochemistry & cell biology 2015; **66**: 20-9.
24.   Kabeerdoss J, Sandhya P, Danda D. Y RNA derived small RNAs in Sjogren's syndrome:
Candidate biomarkers? *International journal of rheumatic diseases* 2017; **20**(11): 1763-6.
25.   Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046 individuals
identifies 11 new susceptibility loci for Alzheimer's disease. *Nature genetics* 2013; **45**(12):
1452-8.
26.   Kunkle BW, Grenier-Boley B, Sims R, et al. Genetic meta-analysis of diagnosed
Alzheimer's disease identifies new risk loci and implicates Abeta, tau, immunity and lipid
processing. *Nature genetics* 2019; **51**(3): 414-30.
27.   Oberbauer V, Schaefer MR. tRNA-Derived Small RNAs: Biogenesis, Modification,
Function and Potential Impact on Human Disease Development. *Genes (Basel)* 2018; **9**(12).
28.   Shi J, Zhang Y, Zhou T, Chen Q. tsRNAs: The Swiss Army Knife for Translational
Regulation. *Trends Biochem Sci* 2019; **44**(3): 185-9.
29.   Ji L, Chen X. Regulation of small RNA stability: methylation and beyond. *Cell research*
2012; **22**(4): 624-36.
30.   Ren G, Xie M, Zhang S, Vinovskis C, Chen X, Yu B. Methylation protects microRNAs
from an AGO1-associated activity that uridylates 5' RNA fragments generated by AGO1
cleavage. *Proc Natl Acad Sci U S A* 2014; **111**(17): 6365-70.
31.   Lee M, Kim B, Kim VN. Emerging roles of RNA modification: m(6)A and U-tail. *Cell*
2014; **158**(5): 980-7.
32.   Will CL, Luhrmann R. Spliceosomal UsnRNP biogenesis, structure and function.
Current opinion in cell biology 2001; **13**(3): 290-301.
33.   Hausmann S, Zheng S, Costanzo M, et al. Genetic and biochemical analysis of yeast
and human cap trimethylguanosine synthase: functional overlap of 2,2,7-trimethylguanosine
caps, small nuclear ribonucleoprotein components, pre-mRNA splicing factors, and RNA decay
pathways. *J Biol Chem* 2008; **283**(46): 31706-18.
34.   Zhang S, Li H, Zheng L, Li H, Feng C, Zhang W. Identification of functional tRNA-
derived fragments in senescence-accelerated mouse prone 8 brain. *Aging (Albany NY)* 2019;
11(22): 10485-98.
35.   Hanada T, Weitzer S, Mair B, et al. CLP1 links tRNA metabolism to progressive motor-
neuron loss. *Nature* 2013; **495**(7442): 474-80.
36.   Schaffer AE, Eggens VR, Caglayan AO, et al. CLP1 founder mutation links tRNA
splicing and maturation to cerebellar development and neurodegeneration. *Cell* 2014; **157**(3):
651-63.
37.   Ding Q, Zhu H, Zhang B, Soriano A, Burns R, Markesbery WR. Increased 5S rRNA
oxidation in Alzheimer's disease. *J Alzheimers Dis* 2012; **29**(1): 201-9.
38.   Lyons SM, Fay MM, Ivanov P. The role of RNA modifications in the regulation of
tRNA cleavage. *FEBS Lett* 2018; **592**(17): 2828-44.
39.   Frye M, Harada BT, Behm M, He C. RNA modifications modulate gene expression
during development. *Science* 2018; **361**(6409): 1346-9.
434
435

436   **Table 1.** Characteristics of study subjects

	No abnormalities (CONT)	Alzheimer disease (AD)	Alzheimer disease with Vascular dementia (VaD+AD)
Demographics			
Total patients (n)	7	18	9
Sex, male/female	5/2	9/9	6/3
Ethnicity			
White /unknown	6/1	18/0	9/0
Age (years)	85·4 ± 2·6	77·3 ± 2·6	79·6 ± 2·5
Brain			
Weight (g)	1286 ± 34·3	1130 ± 33 **	1180 ± 33·4*
PH	6·03 ± 0·08	6·15 ± 0·09	5·96 ± 0·08
Post-Mortem Delay Time	41·2 ± 4·9	36·9 ± 5·4	48·5 ± 7·6

437
438   Values are means ± SE, where appropriate. *$p < 0.05$, vs. control patients; unpaired t-test.
439

Figure legends:

Figure 1. Schematic illustration of LC-MS/MS for detection of RNA modifications. Total RNAs were extracted from brain tissue using the Trizol reagent and then urea-denatured-PAGE was performed to separate different RNA fragments. Targeted RNA fragments were cut according to their size, recovered from the gel, enzymatically treated to obtain single nucleosides, and then subjected to LC/MS-MS profiling. Raw mass spectrometry data were analysed using Xcalibur software, and the relative levels of different RNAs modifications were calculated.

Figure 2. Small RNA (15–25 nt) modifications in the prefrontal lobe cortex of dementia patients. (a) Representative image showing purification of 15–25-nt small RNAs for the analysis of RNA modifications. (b–e) Increases in 2'-O-methylcytidine (Cm) and 7-methylguanosine (m^7G) modifications, and reductions in $N^2,N^2,7$-trimethylguanosine ($m^{2,2,7}G$) and N^2,N^2-dimethylguanosine ($m^{2,2}G$) modifications in the cortex of AD and VaD+AD patients compared with CONT subjects (n = 7 for CONT, n = 18 for AD, n = 9 for VaD+AD; *$P < 0\cdot05$ vs. CONT).

Figure 3. Small RNA (30–40 nt) modifications in the prefrontal lobe cortex of dementia patients. (a) Representative image showing purification of 30–40-nt small RNAs for the analysis of RNA modifications. (b–g) Increases in 2'-O-methylcytidine (Cm), 2'-O-methyluridine (Um) and 7-methylguanosine (m^7G) modifications, and reductions in 1-methylguanosine (m^1G), N^2,N^2-dimethylguanosine ($m^{2,2}G$) and pseudouridine (psi or Ψ) modifications in the cortex of AD and VaD+AD patients compared with CONT subjects (n = 7 for CONT, n = 27 for AD; * $P < 0\cdot05$ vs. CONT).

Figure 4. Reductions in 5S rRNA, tRNA-Tyr, and tRNA-Arg in prefrontal lobe cortex samples of AD patients, identified by small RNA sequencing. (a) Composition of small RNAs (15–25 nt). (b) Composition of small RNAs (34–40 nt). (c) 5s RNA, (d) tRNA-Tyr, (e) tRNA-Arg in the cortex of AD patients compared with CONT subjects (n = 4 for CONT, n = 6 for AD; *$P < 0\cdot05$ vs. CONT).

Figure 1.

Figure 2.

A

B   **Cm**

C   **m⁷G** (m^7G)

D $m^{2,2,7}G$

E $m^{2,2}G$

Figure 3.

A

B Cm

C Um

D m⁷G

E m¹G

F m²,²,⁷G

G psi

Figure 4.

A **RNA Sequencing: Percentage of small RNA from 15-25nt**

B **RNA Sequencing: Percentage of small RNA from 30-40nt**