Title: Small RNA modifications in Alzheimer's disease

Xudong Zhang[1]*, Fatima Trebak[2]*, Lucas AC Souza[2], Junchao Shi[1], Tong Zhou[3], Patrick G.

Kehoe[4], Qi Chen[1#], Yumei Feng Earley[2#]

1. Division of Biomedical Sciences, School of Medicine, University of California, Riverside,
Riverside, CA, USA
2. Departments of Pharmacology, Physiology & Cell Biology, Center for Molecular &
Cellular Signal Transduction in the Cardiovascular System, University of Nevada, Reno,
School of Medicine, Reno, NV, USA
3. Departments of Physiology & Cell Biology, University of Nevada, Reno, School of
Medicine, Reno, NV, USA
4. Dementia Research Group, Translational Health Sciences, Bristol Medical School,
University of Bristol, Bristol, UK
14

*Contributed equally to this work

#**Correspondence to:**

Yumei Feng Earley, MD, PhD, FAHA
Associate Professor of Pharmacology and Physiology & Cell Biology
Center for Molecular & Cellular Signal Transduction in the Cardiovascular System
University of Nevada, Reno, School of Medicine
1664 North Virginia Street, Mail-stop 0318
Reno, NV, 89557, USA
E-mail: yumeifeng@med.unr.edu

Qi Chen, MD, PhD
Assistant Professor
Division of Biomedical Sciences, School of Medicine
University of California, Riverside
Riverside, CA, 92521, USA
E-mail: Qi.Chen@medsch.ucr.edu
34

Abstract

Background: While significant advances have been made in uncovering the aetiology of Alzheimer's disease and related dementias at the genetic level, molecular events at the epigenetic level remain largely undefined. Emerging evidence indicates that small non-coding RNAs (sncRNAs) and their associated RNA modifications are important regulators of complex physiological and pathological processes, including aging, stress responses, and epigenetic inheritance. However, whether small RNAs and their modifications are altered in dementia is not known.

Methods: We performed LC-MS/MS–based, high-throughput assays of small RNA modifications in post-mortem samples of the prefrontal lobe cortices of Alzheimer's disease (AD) and control individuals. We noted that some of the AD patients has co-occurring vascular cognitive impairment-related pathology (VaD).

Findings: We report altered small RNA modifications in AD samples compared with normal controls. The 15–25-nucleotide (nt) RNA fraction of these samples was enriched for microRNAs, whereas the 30–40-nt RNA fraction was enriched for tRNA-derived small RNAs (tsRNAs), rRNA-derived small RNAs (rsRNAs), and YRNA-derived small RNAs (ysRNAs). Interestingly, most of these altered RNA modifications were detected both in the AD and AD with co-occurring vascular dementia subjects. In addition, sequencing of small RNA in the 30–40-nt fraction from AD cortices revealed reductions in rsRNA-5S, tsRNA-Tyr, and tsRNA-Arg.

Interpretation: These data suggest that sncRNAs and their associated modifications are novel signals that may be linked to the pathogenesis and development of Alzheimer's disease.

Funding: NIH grants (R01HL122770, R01HL091905, 1P20GM130459, R01HD092431, P50HD098593, GM103440), AHA grant (17IRG33370128), Sigmund Gestetner Foundation Fellowship to P Kehoe.

Keywords: Non-coding RNA; Alzheimer's disease; Prefrontal cortex; Epigenetics

Research in Context

Evidence before this study

Alzheimer's disease (AD) and vascular dementia (VaD) are marked by cognitive impairment

and neuropathologies caused by significant neuronal death. Associated gene mutations are

rare in subjects with dementia, and the aetiology of these diseases is still not completely

understood. Recent emerging evidence suggests that epigenetic changes are risk factors for

the development of dementia. However, studies assessing small RNA modifications—one of

the features of epigenetics—in dementia are lacking.

Added value of this study

We used high-throughput liquid chromatography-tandem mass spectrometry and small RNA

sequencing to profile small RNA modifications and the composition of small RNAs in post-

mortem samples of the prefrontal cortex of AD and control subjects. We detected and

quantified 16 types of RNA modifications and identified distinct small non-coding RNAs and

modification signatures in AD subjects compared with controls.

Implications of all the available evidence

This study identified novel types and compositions of small RNA modifications in the

prefrontal cortex of AD patients compared with control subjects in post-mortem samples. The

cellular locations of these RNA modifications and whether they are drivers or outcomes of

AD are still not known. However, results from the present study may open new possibilities

for dissecting the dementia pathology.

Introduction

Alzheimer's disease (AD) and severe vascular cognitive impairment, also known as vascular dementia (VaD), are two of the most common types of dementia. They are characterized by cognitive impairment and neuropathology caused by neurotoxic forms of amyloid beta peptide, loss of synapses and neuronal function and, ultimately, significant neuronal loss. The aetiology of these forms of dementia is still not fully understood.[1]

Recent studies have highlighted epigenetic changes associated with aging or acquired through interactions with the environment as important risk factors for dementia, reflecting the fact that causative gene mutations are rare and are only present in a very small percentage of dementia patients. RNA modification is one of the most recently discovered epigenetic-mediated mechanisms for regulating physiological and pathological processes. Small RNA modifications, particularly those involving microRNAs (miRNAs)[2,3] and transfer RNA-derived small RNAs (tsRNAs), have attracted considerable recent attention for their potential clinical relevance and possible use as diagnostic markers or therapeutic targets of diseases.[4-6] To date, evidence for changes in small RNA modifications in dementia is lacking. Thus, to gain insights into possible new mechanisms and identify potential biomarkers for dementia, we undertook an exploratory study to profile small RNA expression and modification status in dementia.

In this report, we profiled small RNA modifications in post-mortem samples of human brain cortical tissue from patients with clinically and pathologically diagnosed AD and control individuals. To simultaneously detect and quantify 16 types of RNA modifications[7,8] in different small RNA fractions [15–25 nucleotides (nt), 30–40 nt] from AD patients and control individuals, we used high-throughput liquid chromatography-tandem mass

spectrometry (LC-MS/MS). We also profiled the composition of small RNAs in AD and

control subjects using small RNA sequencing. These exploratory examinations revealed

distinct sncRNA expression and RNA modification signatures between AD and controls,

suggesting that these small RNAs are novel factors associated with the pathogenesis and/or

progression of AD.

Materials and methods

Human subjects for RNA modification assays

A total of 34 post-mortem brain prefrontal lobe cortex samples from AD patients (n = 27) and

control human subjects (n = 7) were collected by post-mortem autopsy, snap-frozen, and

stored in liquid nitrogen. The time between biospecimen acquisition and analysis ranged from

5 to 30 years, according to the Brain Bank database. Biospecimens were shipped on dry ice.

Information from clinical and pathology reports was obtained from the brain bank database,

developed as described.[9] On the basis of their clinical diagnosis, subjects were first

categorized into two groups: no significant abnormalities (control) and AD. We realized that

some AD patients had co-occurring vascular dementia. This type of patient usually has a more

rapid course of disease clinically. We thus further divided the AD patients into AD without

VaD (AD-w/o-VaD, n = 18) and AD with VaD (AD-w-VaD, n = 9) for the purpose of

analysis. Subjects in the AD-w-VaD group were clinically diagnosed with clear AD tau

pathology and Aβ plaques in post-mortem analysis by the UK Brain Bank, and a

neuropathological report of AD. Subjects were selected according to their confirmed clinical

and pathological diagnosis as AD or normal, and were obtained from the Human Tissue

Authority-licensed South West Dementia Brain Bank http://www.bristol.ac.uk/translational-

health-sciences/research/neurosciences/research/dementia/swdbb/, accessed on March 24[th],

2020). detailed information of these subject is also summarized in Supplemental Table S1.

University of Bristol, with tissue bank ethics approval from the South West–Central Bristol

Research Ethics Committee. Clinical and pathological data included patient history, diagnosis

and medications; pathological reports were available for retrospective analysis. Patients'

personal information has been anonymized. Data and tissues for this study were collected

between 1989 and 2015. The Research Integrity Offices at the University of Nevada, Reno,

and University of Bristol have determined that this project complies with human research

protection oversight by the Institutional Review Board.

Isolation of total RNA from the human prefrontal cortex

Total RNA was isolated from 100 mg of human prefrontal tissues using the TRIzol reagent

(Thermo Fisher Scientific, Inc. Waltham, MA, USA), as described by the manufacturer. RNA

quantity was evaluated using a NanoDrop 2000 microvolume spectrophotometer (Thermo

Fisher). RNA samples were stored at -80°C until analysis. Researchers performing small

RNA sequencing and RNA modifications assays on RNA samples were blinded to diagnosis

and group-identifying information.

Small RNA Sequencing

Ten RNA samples from prefrontal lobe cortexes of control subjects and AD patients were

submitted to BGI (Cambridge, MA, USA) for sequencing of 15–40-nt small RNAs. The

number of clean reads varied from 30M to 40M across all samples. The Q20 score was

greater than 99·5% for all samples. Raw sequencing data for each sample were processed and

analysed using our newly developed computational framework, *SPORTS1·0*[10]

(https://github.com/junchaoshi/sports1.0), which is designed to optimize the annotation and

quantification of non-canonical small RNAs (e.g., tsRNAs) in small RNA sequencing data.

Sequencing adapters were removed, after which reads with lengths outside of the defined

range and those with nucleotides other than ATUCG were discarded. Clean reads were

mapped against the precompiled human small RNA annotation database

(https://github.com/junchaoshi/sports1.0/blob/master/precompiled_annotation_database.md).

Small RNAs 15–25 and 30–40 nt in length were summarized separately.

Small RNA purification and analysis of RNA modifications using LC-MS/MS

Standardized ribonucleoside preparations, small RNA purification, and LC-MS/MS-based

analysis of RNA modifications in RNA samples were performed as previously described.[8]

Purified small RNAs (100–200 ng) from human brain samples were digested as input and

loaded onto a ThermoFisher Vantage Quadrupole mass spectrometer connected to a Thermo

Ultimate 3000 UHPLC system equipped with an electrospray ionization source. The MS

system was operated in positive ion mode using a multiple reaction monitoring (MRM)

approach. LC-MS/MS raw data were acquired using Xcalibur Workstation software and were

processed using Xcalibur QuanBrowser for quantification of modified ribonucleoside

concentrations. We totally performed 34 samples. However, during the analysis of the mass

spectrometry data, we exclude data points when the signal peak is not unique (e.g. two peaks,

elevated baseline), or drifting off the detection time predetermined by each standardized

ribonucleoside, which resulted in inaccurate reading (or loss of reading) of the examined

signal. The percentage of each modified ribonucleoside was normalized to the total amount of

quantified ribonucleosides containing the same nucleobase, an approach that

decreases/eliminates errors caused by sample loading variation. For example, the percentage

of m^5C = mole concentration (m^5C)/mole concentration (m^5C + Cm + C + ac^4C). Fold-

changes in RNA modifications between different groups were calculated based on the

percentage of modified ribonuclosides. Brain RNAs 15–25 and 30–40 nucleotides (nt) in

length were examined.

Statistical analysis

Data are presented as means ± SEM and were plotted using Prism8 software (GraphPad, La

Jolla, CA, USA). Differential expression analyses were performed using the *edgeR* tool (pmid:

19910308), controlling for age and sex. RNAs with a false discovery rate < 10% were deemed

to be differentially expressed. RNA modification levels among groups were compared using a

linear model that controls for age and sex. Student *t*-test or Ordinary One-way ANOVA with

post hoc Bonferroni correction was used as appropriate for comparisons among groups. A *P*-

value < 0·05 was considered statistically significant.

Results

Participant characteristics

As summarized in Tables 1 and 2, patients and control individuals were similar in terms of

age. Almost all patients were White/European except for one patient whose race was

unknown. There were slightly more males (58·8%) than females (41·2%). As expected, brain

weight was significantly lower in AD ($P = 0·0099$) compared with the controls (Table 1). This

phenomenon persisted when we subdivided the AD into (AD-w/o-VaD) ($P = 0·018$) when

compared with controls; however, there was no significant difference between AD-w-VaD

and controls for the brain weight (Table 2). No significant differences were observed for the

brain pH and post-mortem delay time among groups.

Altered small RNA modification profiles in the cerebral cortex of AD patients

To systemically analyse small RNA expression and modification profiles, we collected the

15–40-nt RNA fraction from the brain frontal lobe cortex for small RNA sequencing as we

previously described[7,8], and then performed bioinformatic analyses using our recently

developed software, *SPORTS1.0.*[10] In addition, 15–25-nt and 30–40-nt RNA fractions were

collected for high-throughput examination of RNA modification by LC-MS/MS

(Supplemental Figure S1).[7,11] These analyses resulted in the identification and quantification

of 16 types of RNA modifications.

In the 15–25-nt RNA fraction from the cortex of AD brains (Figure 1A), we found an increase

in 2'-O-methylcytidine (Cm), 7-methylguanosine (m^{7}G), 2'-O-methylguanosine (Gm), and

significant reductions in N^2,N^2,7-trimethylguanosine (m^{2,2,7}G) and N^2,N^2-dimethylguanosine

(m^{2,2}G), compared with controls (Figure 1B and 1C). Other RNA modifications that were not

significantly different among groups were also detected and are shown in Supplemental

Figure S2. Since AD Patients with co-occurring several vascular cognitive impairment-related

pathologies (e.g. VaD) tend to have a more rapid course of disease clinically.[12] We further test

whether these patients have a distinct difference at RNA modifications. To this end, we

divided AD patients into two groups, the AD-w/o-VaD and AD-w-VaD and performed

further analysis. We found that both AD-w/o-VaD group and AD-w-VaD group showed

similar trends in RNA modification changes when compare to the normal subjects, but some

RNA modifications cannot reach statistical significance when the two groups are divided

(Figure 2), possibly due to decreased sample sized in both groups.

In the 30–40-nt small RNA fraction from the cortex of AD brains (Figure 3A), we found

significantly higher levels of Cm, 2'-O-methyluridine (Um) and 7-methylguanosine (m^{7}G)

modifications, and reductions in 1-methylguanosine (m^{1}G), m^{2,2,7}G and pseudouridine (psi or

Ψ) modifications compared with controls (Figure 3B and 3C). Other RNA modifications

detected in this 30–40-nt fraction that did not exhibit significant differences among groups are

shown in Supplemental Figure S3. When further divided AD into with or without VaD

(Figure 4), we found that AD-w/o-VaD maintained significant changes in most of these RNA

modifications. One the other hand, the AD-w-VaD group maintained the same trend but did

not reach significant difference in most of the RNA modifications compared with the controls

possibly due to the small sample size; however, there was no difference between the AD-w-

VaD and the AD-w/o-VaD suggesting the similar expression profile.

Altered 30–40-nt small RNA expression profiles in the cerebral cortex of AD patients

To systemically analyse small RNA expression profiles, we collected the 15–40-nt fraction

from the brain frontal lobe cortex of AD patients and controls for small RNA sequencing, as

we previously described[7,8], and then performed bioinformatic analyses using our recently

developed software, *SPORTS1·0*[10]. Our RNA-seq analysis showed that the 15–25-nt RNA

population was predominantly miRNAs (Figure 5A) in both the AD and control samples; we

did not find significant differences in specific small RNA sequences in this fraction between

AD and control samples (data not shown). Whereas the expression profile of the small RNA

population in the 30–40-nt fraction showed more dynamic changes. (Figure 5B-E). Our RNA-

seq data revealed three major subtypes in the 30–40-nt fraction, namely tRNA-derived small

RNAs (tsRNA), rRNA-derived small RNAs (rsRNAs) and Y RNA-derived small RNAs

(ysRNAs); other un-annotated RNAs were also found (Figure 5B). Although there were

trends toward an increase in tsRNAs and a reduction in ysRNAs in AD patients, these

differences did not reach statistical significance. Interestingly, in the 30–40-nt fraction, we

found a significant reduction in rsRNA-5S, tsRNA-Tyr and tsRNA-Arg fragments in AD

patients compared with controls (Figure 5C–E).

Discussion

Emerging evidence shows that small RNAs, including miRNAs, tsRNAs and piwi-interacting

RNAs, harbour a diversity of RNA modifications.[13] The physiological and pathological

importance of these small, non-coding RNA modifications has only recently begun to emerge,

as highlighted by studies revealing their active involvement in stress responses, metabolism,

immunity, and epigenetic inheritance of environmentally acquired traits.[4,14,15] RNA

modifications have recently been detected in the nervous system, where they are involved in

the regulation of cortical differentiation, behaviour, and brain functions.[16,17] However, to our

knowledge, there have been no reports on the status of small RNA modifications in AD or

VaD to date.

The current study showed intriguing differences in small RNA-modification signatures

between AD patients and control subjects, suggesting an association and possible contribution

of such modifications to the pathogenesis and/or progression of AD. Notably, most of the

altered RNA modifications were observed in the AD with and without VaD, providing

molecular evidence for potential distinct signatures for AD patients. In the last three decades,

dementia research has demonstrated significant overlaps between AD and VaD in terms of

clinical symptoms, risk factors, and post-mortem brain autopsy findings.[18-20] Therapeutically,

drugs that enhance cholinergic activity are as effective in patients suffering from VaD as they

are in AD patients.[21] More recently, drugs that have traditionally been used for cardiovascular

diseases, especially renin-angiotensin system blockers, have shown benefits not only in VaD

patients, but also in AD patients.[22-24] However, the molecular pathogenesis of AD with co-

occurring sever vascular cognitive impairment remains incompletely understood. It is

however worth highlighting that patients with AD and co-occurring severe Vascular

Cognitive Impairment-related pathology (e.g. VaD) tend to have a more rapid course of

disease clinically because of the co-occurrence of heightened cerebrovascular disease and

hypoxia and related white matter and neuronal damage.[12,25] It was therefore prudent to

consider these as a separate group to explore whether RNA modifications might be more

prevalent and thus potentially associated with the disease. We were not surprised to find many

common signatures among RNA modifications in AD with or without VaD patients,

particularly in the 15–25-nt small RNA modifications that might contribute to shared

mechanisms of dementia pathology. Yet, the lack of apparent differences between the groups

does suggest, notwithstanding the small sample sizes, that the RNA modifications may relate

more to AD-related processes than neuropathology caused by VaD. Further studies, involving

post-mortem analysis of other cerebrovascular-related diseases and in the absence of AD or

with minimal pathology, and with larger sample size, would be useful studies to explore this

further.

The 15–25-nt RNAs were predominantly miRNAs, whereas 30–40-nt RNAs were

predominantly tsRNAs, rsRNAs and ysRNAs, as demonstrated by small RNA sequencing.

We found changes in RNA-modification profiles in the AD brain cortex in both fractions of

small RNAs. While the involvement of miRNA in the pathogenesis of AD has been

explored[2,3,26], the identifications of rsRNAs and ysRNAs in a pathophysiological context has

only begun to emerge [8,27,28] and has not been studied in AD. The dynamic expression of these

non-canonical small RNAs suggests the possibility of unidentified biological functions that

warrant further investigation[29,30], particularly in relation to several risk factor genes that have

been identified for AD in the last few decades.[31,32] This situation is similar to that for tsRNAs,

which have shown recently expanding functions.[6,33,34] The novel RNA modifications that

showed dynamic changes in AD patients included Cm, Um, m^7G, m^1G, m^3C, $m^{2,2}G$, $m^{2,2,7}G$,

and psi. The functions and identities of these small RNA modifications in AD are not yet

understood, but our data suggest that these RNA modifications in the brain cortex of dementia

patients are potentially important, either as a consequence or an actual cause of pathogenesis.

For example, it has been reported that pH, which may affect the efficiency of related RNA-

modifying enzymes and thus contribute to the RNA modification profile, is lower in the

brains of AD patients compared with normal aging controls[35,36]. Although we observed a

trend toward lower pH values in post-mortem samples from AD w/o VaD brains, this

difference did not reach statistical significance.

tRNA-derived small RNAs are reported to contribute to multiple pathological processes,

including cancer, viral infection, and age-related neurodegenerative diseases.[4] In the

senescence-accelerated mouse prone 8 (SAMP8) model, which mimics age-related neuro-

disorders such as AD and Parkinson disease, tsRNAs, including tsRNA-Tyr and tsRNA-Arg,

in brain tissue are dysregulated compared with normal brain tissue. The targets of these

dysregulated tsRNAs are enriched in neurodevelopment pathways such as synapse

formation.[37] In addition, knockout of the RNA kinase, CLP1, in mice leads to progressive loss

of motor neuron function in association with the accumulation of tsRNAs derived from Try-

tRNA and Arg-tRNA .[38,39] These tsRNAs sensitize motor neurons to oxidative stress-induced

cell death, suggesting that tsRNAs are involved in normal motor neuron functions and

responses to oxidative stress.[38] A similar phenomenon was reported in a human neurological

disease cohort, and it was shown that transfection of small RNAs derived from 5' Tyr-tRNA

can protect CLP-mutant cells from oxidative stress-induced cell death.[39] In our study, the

decrease in 5'tsRNA-Tyr in AD subjects may confer greater vulnerability to oxidative stress

on neurons. Interestingly another report showed that the expression of 5S rRNA and levels of

oxidized 5S rRNA are dynamically modulated in AD's subjects[40], findings that may be

related to the biogenesis of rsRNAs and could explain alterations of rsRNA-5S in the AD

group in the current study.

Whether the changes in small RNAs and RNA modifications observed here represent drivers

or outcomes of AD remains unknown. In our study, the sample number from the control and

patient groups is relatively small, which prevent deeper analyses to consider other factors

such as ages, NFT pathology stage, and PMI. One limitation of the current study is the

prolonged PMI in some of the brain samples as we have no access to adequate number of

samples with shorter PMI (e.g. < 24 hours PMI). Another potential limitation of our study is

the prolonged storage time in some of the tissues, as long-term storage might affect the RNA

modification results. To examine this possibility. We performed principle components

analysis (PCA), which converts the observations of possibly correlated variables (RNA

modifications) into a set of values of linearly uncorrelated variables. This analysis of the 27

AD samples ranging from 1985 to 2015 does not support this possibility that storage time

affects RNA modification as showed in Supplemental Figure S4. This result support the

conclusion that the detected changes in RNA modifications indeed represent intrinsic

difference in the AD samples and the storage time is not a major contributing factor on the

result.

In summary, the current pioneer study based on limited sample numbers have revealed that

RNA modifications in the sncRNAs are associated with AD, although the precise meaning of

each of these changes in terms of their specific functions awaits discovery. Future efforts to

pinpoint the locations of these modifications in each sncRNA population and identify

enzymes involved in their regulation would be invaluable[41,42] and will open new avenues for

dissecting the nature of dementia pathology.

Funding

This work was supported, in part, by grants from the National Institutes of Health (NIH/NHLBI) (R01HL122770, R01HL091905), NIH/NIGMS (1P20GM130459) and the American Heart Association National Center (17IRG33370128) to Y. Feng Earley, and from the NIH/NICHD to Q. Chen (R01HD092431) and T. Zhou (P50HD098593). Research reported in this publication used instruments from the Dr. Mick Hitchcock Proteomics Core supported by a grant from the NIH/NIGMS (GM103440) of the National Institutes of Health, University of Bristol Alumni. P Kehoe is supported by a Fellowship from the Sigmund Gestetner Foundation. The funders had no role in study design, data collection, analysis and interpretation, or writing of the manuscript.

Declaration of competing interests

The authors declare no competing interests.

Data availability

Small RNA-seq data that support the findings of this study have been deposited in the Gene Expression Omnibus (GEO) under accession code GSE153284. All other data supporting the findings of this study are available from the corresponding author on reasonable request.

Acknowledgements

386 We would like to thank Dr. David Quilici and Rebekah Woolsey from the Mick Hitchcock,

387 Ph.D. Nevada Proteomics Center for technical support with HPLC/MS instruments. We

388 would especially like to thank Dr. Laura Palmer for tissue processing and facilitating access

389 to clinical data from the South West Dementia Brain Bank at the University of Bristol, which

390 is supported by grants from BRACE, ABBUK.

References:

1.   Panza F, Lozupone M, Watling M, Imbimbo BP. Do BACE inhibitor failures in Alzheimer patients challenge the amyloid hypothesis of the disease? *Expert review of neurotherapeutics* 2019; **19**(7): 599-602.

2.   Hébert SS, Wang W-X, Zhu Q, Nelson PT. A study of small RNAs from cerebral neocortex of pathology-verified Alzheimer's disease, dementia with lewy bodies, hippocampal sclerosis, frontotemporal lobar dementia, and non-demented human controls. *Journal of Alzheimer's disease : JAD* 2013; **35**(2): 335-48.

3.   Lugli G, Cohen AM, Bennett DA, et al. Plasma Exosomal miRNAs in Persons with and without Alzheimer Disease: Altered Expression and Prospects for Biomarkers. *PloS one* 2015; **10**(10): e0139233-e.

4.   Zhang X, Cozen AE, Liu Y, Chen Q, Lowe TM. Small RNA Modifications: Integral to Function and Disease. *Trends Mol Med* 2016; **22**(12): 1025-34.

5.   Chen Q, Yan W, Duan E. Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nature reviews Genetics* 2016; **17**(12): 733-43.

6.   Schimmel P. The emerging complexity of the tRNA world: mammalian tRNAs beyond protein synthesis. *Nature reviews Molecular cell biology* 2018; **19**(1): 45-58.

7.   Chen Q, Yan M, Cao Z, et al. Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 2016; **351**(6271): 397-400.

8.   Zhang Y, Zhang X, Shi J, et al. Dnmt2 mediates intergenerational transmission of paternally acquired metabolic disorders through sperm small non-coding RNAs. *Nature cell biology* 2018; **20**(5): 535-40.

9.   Hyman BT, Phelps CH, Beach TG, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 2012; **8**(1): 1-13.

10.   Shi J, Ko EA, Sanders KM, Chen Q, Zhou T. SPORTS1.0: A Tool for Annotating and Profiling Non-coding RNAs Optimized for rRNA- and tRNA-derived Small RNAs. *Genomics Proteomics Bioinformatics* 2018; **16**(2): 144-51.

11.   Fang H, Deng M, Zhang L, et al. Role of (Pro)Renin Receptor in Albumin Overload-Induced Nephropathy in Rats. *Am J Physiol Renal Physiol* 2018.

12.   Iadecola C. The pathobiology of vascular dementia. *Neuron* 2013; **80**(4): 844-66.

13.   Roundtree IA, Evans ME, Pan T, He C. Dynamic RNA Modifications in Gene Expression Regulation. *Cell* 2017; **169**(7): 1187-200.

14.   Lovell MA, Soman S, Bradley MA. Oxidatively modified nucleic acids in preclinical Alzheimer's disease (PCAD) brain. *Mech Ageing Dev* 2011; **132**(8-9): 443-8.

15.   Pietrzak M, Rempala G, Nelson PT, Zheng J-J, Hetman M. Epigenetic silencing of nucleolar rRNA genes in Alzheimer's disease. *PloS one* 2011; **6**(7): e22585-e.

16.   Satterlee JS, Basanta-Sanchez M, Blanco S, et al. Novel RNA modifications in the nervous system: form and function. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2014; **34**(46): 15170-7.

17.   Jung Y, Goldman D. Role of RNA modifications in brain and behavior. *Genes, brain, and behavior* 2018; **17**(3): e12444.

18.   Ravona-Springer R, Davidson M, Noy S. Is the distinction between Alzheimer's disease and vascular dementia possible and relevant? *Dialogues in clinical neuroscience* 2003; **5**(1): 7-15.

19.   Mirra SS, Heyman A, McKeel D, et al. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). *Neurology* 1991; **41**(4): 479.

20.   Meyer JS, Xu G, Thornby J, Chowdhury MH, Quach M. Is mild cognitive impairment prodromal for vascular dementia like Alzheimer's disease? *Stroke* 2002; **33**(8): 1981-5.

21.   Erkinjuntti T, Kurz A, Gauthier S, Bullock R, Lilienfeld S, Damaraju CV. Efficacy of
galantamine in probable vascular dementia and Alzheimer's disease combined with
cerebrovascular disease: a randomised trial. *Lancet* 2002; **359**(9314): 1283-90.
22.   Zhuang S, Wang H-F, Wang X, Li J, Xing C-M. The association of renin-angiotensin
system blockade use with the risks of cognitive impairment of aging and Alzheimer's disease:
A meta-analysis. *Journal of Clinical Neuroscience* 2016; **33**: 32-8.
23.   Chou C-L, Yeh H-I. The Role of the Renin-Angiotensin System in Amyloid
Metabolism of Alzheimer's Disease. *Acta Cardiologica Sinica* 2014; **30**(2): 114-8.
24.   Gebre AK, Altaye BM, Atey TM, Tuem KB, Berhe DF. Targeting Renin-Angiotensin
System Against Alzheimer's Disease. *Frontiers in pharmacology* 2018; **9**: 440.
25.   Vinters HV, Zarow C, Borys E, et al. Review: Vascular dementia: clinicopathologic
and genetic considerations. *Neuropathol Appl Neurobiol* 2018; **44**(3): 247-66.
26.   Ayers D, Scerri C. Non-coding RNA influences in dementia. *Noncoding RNA Res*
2018; **3**(4): 188-94.
27.   Hizir Z, Bottini S, Grandjean V, Trabucchi M, Repetto E. RNY (YRNA)-derived
small RNAs regulate cell death and inflammation in monocytes/macrophages. *Cell Death Dis*
2017; **8**(1): e2530.
28.   Zhang Y, Shi J, Rassoulzadegan M, Tuorto F, Chen Q. Sperm RNA code programmes
the metabolic health of offspring. *Nature reviews Endocrinology* 2019; **15**(8): 489-98.
29.   Kowalski MP, Krude T. Functional roles of non-coding Y RNAs. *The international*
journal of biochemistry & cell biology 2015; **66**: 20-9.
30.   Kabeerdoss J, Sandhya P, Danda D. Y RNA derived small RNAs in Sjogren's
syndrome: Candidate biomarkers? *International journal of rheumatic diseases* 2017; **20**(11):
1763-6.
31.   Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046
individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nature genetics* 2013;
45(12): 1452-8.
32.   Kunkle BW, Grenier-Boley B, Sims R, et al. Genetic meta-analysis of diagnosed
Alzheimer's disease identifies new risk loci and implicates Abeta, tau, immunity and lipid
processing. *Nature genetics* 2019; **51**(3): 414-30.
33.   Oberbauer V, Schaefer MR. tRNA-Derived Small RNAs: Biogenesis, Modification,
Function and Potential Impact on Human Disease Development. *Genes (Basel)* 2018; **9**(12).
34.   Shi J, Zhang Y, Zhou T, Chen Q. tsRNAs: The Swiss Army Knife for Translational
Regulation. *Trends Biochem Sci* 2019; **44**(3): 185-9.
35.   Lyros E, Ragoschke-Schumm A, Kostopoulos P, et al. Normal brain aging and
Alzheimer's disease are associated with lower cerebral pH: an in vivo histidine 1H-MR
spectroscopy study. *Neurobiology of Aging* 2020; **87**: 60-9.
36.   Ponto L, Schultz S, Jacob M, Menda Y, Wemmie J, Magnotta V. Brain pH and
Alzheimer's pathology. *Journal of Nuclear Medicine* 2014; **55**(supplement 1): 192.
37.   Zhang S, Li H, Zheng L, Li H, Feng C, Zhang W. Identification of functional tRNA-
derived fragments in senescence-accelerated mouse prone 8 brain. *Aging (Albany NY)* 2019;
11(22): 10485-98.
38.   Hanada T, Weitzer S, Mair B, et al. CLP1 links tRNA metabolism to progressive
motor-neuron loss. *Nature* 2013; **495**(7442): 474-80.
39.   Schaffer AE, Eggens VR, Caglayan AO, et al. CLP1 founder mutation links tRNA
splicing and maturation to cerebellar development and neurodegeneration. *Cell* 2014; **157**(3):
651-63.
40.   Ding Q, Zhu H, Zhang B, Soriano A, Burns R, Markesbery WR. Increased 5S rRNA
oxidation in Alzheimer's disease. *J Alzheimers Dis* 2012; **29**(1): 201-9.

41.   Lyons SM, Fay MM, Ivanov P. The role of RNA modifications in the regulation of
tRNA cleavage. *FEBS Lett* 2018; **592**(17): 2828-44.
42.   Frye M, Harada BT, Behm M, He C. RNA modifications modulate gene expression
during development. *Science* 2018; **361**(6409): 1346-9.
493
494

Figure legends:

Figure 1. Small RNA modifications in the 15–25-nt fraction from samples of the prefrontal lobe cortex of dementia patients. (**A**) Representative image showing purification of 15–25-nt small RNAs for the analysis of RNA modifications. (**B**) Increases in 2'-O-methylcytidine (Cm) 7-methylguanosine (m^7G), 2'-O-methylguanosine (Gm), and 3-Methylcytidine(m^3C) modifications. (**C**) Reductions in N^2,N^2,7-trimethylguanosine ($m^{2,2,7}G$) and N^2,N^2-dimethylguanosine ($m^{2,2}G$) modifications in the cortex of AD compared with CONT subjects (each dot in the figure represents value from one independent sample); $*P < 0.05;**P <0.01$ vs. CONT by Student t-test.

Figure 2. Small RNA modifications in the 15–25-nt fraction from samples of the prefrontal lobe cortex of dementia patients with and without co-occurring vascular dementia patients. (**A**) Increases in 2'-O-methylcytidine (Cm) 7-methylguanosine (m^7G), 2'-O-methylguanosine (Gm), and 3-Methylcytidine(m^3C) modifications. (**B**) Reductions in N^2,N^2,7-trimethylguanosine ($m^{2,2,7}G$) and N^2,N^2-dimethylguanosine ($m^{2,2}G$) modifications in the cortex of AD-w/o-VaD, AD-w-VaD, and CONT subjects (each dot in the figure represents value from one independent sample); $*P < 0.05;**P <0.01$ vs. CONT by Ordinary One-way ANOVA with post hoc Bonferroni correction.

Figure 3. Small RNA modifications in the 30–40-nt fraction from samples of the prefrontal lobe cortex of dementia patients. (**A**) Representative image showing purification of 30–40-nt small RNAs for the analysis of RNA modifications. (**B**) Increases in 2'-O-methylcytidine (Cm), 2'-O-methyluridine (Um) and 7-methylguanosine (m^7G) modifications, and (**C**) reductions in 1-methylguanosine (m^1G), N^2,N^2-dimethylguanosine ($m^{2,2}G$) and pseudouridine (psi or Ψ) modifications in the cortex of AD compared with CONT subjects (each dot in the figure represents value from one independent sample); $*P < 0.05$ vs. CONT by Student t-test.

Figure 4. Small RNA modifications in the 30–40-nt fraction from samples of the prefrontal lobe cortex of dementia patients with and without co-occurring vascular dementia patients. Changes in (**A**) 2'-O-methylcytidine (Cm), 2'-O-methyluridine (Um) and 7-methylguanosine (m^7G) modifications, and (**B**) 1-methylguanosine (m^1G), N^2,N^2-dimethylguanosine ($m^{2,2}G$) and pseudouridine (psi or Ψ) modifications in the cortex of AD-w/o-VaD, AD-w-VaD, and CONT subjects (each dot in the figure represents value from one independent sample); $*P < 0.05$ vs. CONT by Ordinary One-way ANOVA with post hoc Bonferroni correction.

Figure 5. Reductions in rsRNA-5S, tsRNA-Tyr, and tsRNA-Arg in prefrontal lobe cortex samples of AD patients, identified by small RNA sequencing. (**A**) Composition of the 15–25 nt fraction of small RNAs. (**B**) Composition of the 34–40 nt fraction of small RNAs. (**C**) rsRNA-5S, (**D**) tsRNA-Tyr, (**E**) tsRNA-Arg in the cortex of AD patients compared with CONT subjects (n = 4 for CONT, n = 6 for AD in panel A-E); $*P < 0.05$ vs. CONT by Student t-test.

Figure 1

Figure 2.

Figure 3

Figure 4.

Figure 5.

A   **RNA Sequencing: Percentage of small RNA from 15-25nt**

B   **RNA Sequencing: Percentage of small RNA from 30-40nt**